Fatal neuroinvasion of SARS-CoV-2 in K18-hACE2 mice is partially dependent on hACE2 expression

Mariano Carossino[6], Paige Montanaro[1*], Aoife O'Connell[1*], Devin Kenney[1*], Hans Gertje[1], Kyle A. Grosz [1], Susanna A. Kurnick[1], Markus Bosmann[1,4,7], Mohsan Saeed[1,5] Udeni B. R. Balasuriya[6], Florian Douam[1,3$#], and Nicholas A. Crossland[1-2$#]

[1]National Emerging Infectious Diseases Laboratories (NEIDL), Boston University, Boston, MA 02118, USA

[2]Department of Pathology and Laboratory Medicine, Boston University School of Medicine, Boston, MA 02118, USA

[3]Department of Microbiology, Boston University School of Medicine, Boston, MA 02118, USA

[4]Department of Medicine, Pulmonary Center, Boston University School of Medicine, Boston, MA 02118, USA

[5]Department of Biochemistry, Boston University, Boston, MA 02118, USA

[6]Louisiana Animal Disease Diagnostic Laboratory (LADDL) and Department of Pathobiological Sciences, School of Veterinary Medicine, Louisiana State University, Baton Rouge, LA, USA

[7] Center for Thrombosis and Hemostasis, University Medical Center of the Johannes Gutenberg-University, Mainz, Germany

*, $ These authors contributed equally to the work.

Co-corresponding author's contact information: Florian Douam, PhD, fdouam@bu.edu, (617)-358-9174; Nicholas Crossland, DVM, ncrossla@bu.edu, (617)-358-9285.

Running Title: Spatiotemporal analysis of SARS-CoV-2 and *hACE2* in K18-hACE2 mice

Keywords: Translational animal model, comparative pathology, immunohistochemistry, *in situ* hybridization, viral pathogenesis

SUMMARY

COVID-19 is a respiratory disease caused by SARS-CoV-2, a betacoronavirus. Here,

we show that in a widely used transgenic mouse model of COVID-19, lethality is

invariably associated with viral neuroinvasion and the ensuing neuronal disease, while

lung inflammation remains moderate.

ABSTRACT

Animal models recapitulating the distinctive features of severe COVID-19 are critical to

enhance our understanding of SARS-CoV-2 pathogenesis. Transgenic mice expressing

human angiotensin-converting enzyme 2 (hACE2) under the cytokeratin 18 promoter

(K18-hACE2) represent a lethal model of SARS-CoV-2 infection. However, the cause(s)

and mechanisms of lethality in this mouse model remain unclear. Here, we evaluated

the spatiotemporal dynamics of SARS-CoV-2 infection for up to 14 days post-infection.

Despite infection and moderate inflammation in the lungs, lethality was invariably

associated with viral neuroinvasion and neuronal damage (including spinal motor

neurons). Neuroinvasion occurred following virus transport through the olfactory

neuroepithelium in a manner that was only partially dependent on hACE2. Interestingly,

SARS-CoV-2 tropism was overall neither widespread among nor restricted to only

ACE2-expressing cells. Although our work incites caution in the utility of the K18-hACE2

model to study global aspects of SARS-CoV-2 pathogenesis, it underscores this model

as a unique platform for exploring the mechanisms of SARS-CoV-2 neuropathogenesis.

INTRODUCTION

The world is experiencing the devastating effects of the Coronavirus Disease 2019 (COVID-19) pandemic, a highly contagious viral respiratory disease caused by the newly emerged betacoronavirus, Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2) (Andersen et al., 2020; Coronaviridae Study Group of the International Committee on Taxonomy of, 2020; Wang et al., 2020a). The initial index case was reported at a seafood market in Wuhan, Hubei Province, China in late 2019 (Andersen et al., 2020). While still under investigation, it has been postulated that the progenitor of SARS-CoV-2 may have originated from horseshoe bats (*Rhinolophus affinis*) or Malayan pangolins (*Manis javanica*) that, following spill over into humans, acquired the genomic features leading to adaptation and human-to-human transmission (Andersen et al., 2020). SARS-CoV-2 has a high transmissibility rate and, to date, it has infected nearly 84.5 million people with over 1.8 million fatalities (Johns Hopkins University & Medicine, 2020). COVID-19 causes respiratory disease of variable severity, ranging from mild to severe, with the development of acute respiratory distress syndrome (ARDS) requiring intensive care and mechanical ventilation (Goyal et al., 2020; Tenforde et al., 2020; Wang et al., 2020a; Wang et al., 2020b). Numerous comorbidities including hypertension, obesity and diabetes, among others, are affiliated with an increased risk of developing severe COVID-19 (Goyal et al., 2020; Simonnet et al., 2020; Tartof et al., 2020; Team, 2020; Tenforde et al., 2020). Furthermore, a proportion of infected patients develop poorly understood neurological signs and/or symptoms mostly associated with the loss of smell and taste (anosmia and ageusia), headache, dizziness, encephalopathy (delirium), and ischemic injury (stroke), among other uncommon symptoms (DosSantos et al., 2020; Eliezer et al., 2020; Ellul et al., 2020;

Goyal et al., 2020; Lee et al., 2020; Liu et al., 2020; Solomon et al., 2020; Walker et al.,

2020; Wang et al., 2020b). Recently, SARS-CoV-2 RNA and antigen has been reported

in the brain of COVID-19 patients and the olfactory mucosa postulated as a port of entry

(Meinhardt et al., 2020). COVID-19 has severely challenged health care systems

around the globe, with the urgent need for medical countermeasures including the

development of efficacious vaccines and therapeutics.

Animal models permissive to SARS-CoV-2 that could serve as suitable models to

understand the pathogenesis of COVID-19 and as preclinical models for the evaluation

of vaccine and therapeutic targets are critically needed (Johansen et al., 2020;

McNamara et al., 2020; Munoz-Fontela et al., 2020). While various animal models

(mice, hamsters, non-human primates, ferrets, minks, dogs, and cats) have been

evaluated to date (Gaudreault et al., 2020; Imai et al., 2020; Meekins et al., 2020;

Munoz-Fontela et al., 2020; Rockx et al., 2020; Shi et al., 2020; Shuai et al., 2020; Sia

et al., 2020; Winkler et al., 2020), none faithfully recapitulates all of the pathological

features of COVID-19. The main limitation in the development of suitable murine models

of COVID-19 is related to the virus entry mechanism: SARS-CoV-2 binds to target cells

via interaction between the viral spike protein (S) and the host angiotensin-converting

enzyme 2 (ACE2), considered to be the major host entry receptor (Hoffmann et al.,

2020). The low binding affinity between the S protein and murine ACE2 (mACE2)

renders conventional mouse strains naturally resistant to infection, posing a challenge in

the development of murine models of COVID-19 (Conceicao et al., 2020; Damas et al.,

2020; Dinnon et al., 2020). These difficulties have been circumvented by the

development of transgenic murine models that express human ACE2 (hACE2) under

different promoters including cytomegalovirus (CMV), hepatocyte nuclear factor-

3/forkhead homologue 4 (HFH4), and cytokeratin 18 (K18) (Jiang et al., 2020; McCray

et al., 2007; Rathnasinghe et al., 2020; Winkler et al., 2020; Zheng et al., 2020). The

transgenic murine model expressing hACE2 under a K18 promoter (namely K18-

hACE2) was developed by McCray et al in 2007 to study SARS-CoV (McCray et al.,

2007), which shares the same host receptor as SARS-CoV-2 (Li et al., 2003).

SARS-CoV-2 infection of K18-hACE2 mice results in up to 100% lethality similar to what

has been reported for SARS-CoV (McCray et al., 2007; Winkler et al., 2020; Zheng et

al., 2020). Lethality was reported to be associated with lung inflammation and impaired

respiratory function, suggesting that this model can recapitulate features of the

respiratory disease observed in severe cases of COVID-19 (Rathnasinghe et al., 2020;

Winkler et al., 2020; Zheng et al., 2020). While several studies have reported SARS-

CoV-2 neuroinvasion and neurological signs of disease in infected K18-hACE2 mice

(Golden et al., 2020; Zheng et al., 2020), the contribution of CNS disease in the

lethality of the model and the mechanism by which neuroinvasion occurs, remain

unclear.

Under K18 regulation, the expression of hACE2 is reported to be limited mainly to

airway epithelial cells and enterocytes lining the colonic mucosa, to a lower degree

within kidney, liver, spleen and small intestine, and a minor level of expression in the

brain (McCray et al., 2007). However, the cellular distribution of ACE2, and particularly

hACE2, in tissues of K18-hACE2 mice remains largely undetermined. We hypothesized

that the nature, severity, and outcome of disease in K18-hACE2 mouse model is not

solely dictated by the expression and tissue distribution of hACE2 and that increased

lethality in this model is related to neuroinvasion. To investigate this hypothesis, we

undertook a comprehensive spatiotemporal analysis of histologic changes, cellular

distribution and abundance of viral antigen and RNA along with detailed analysis of the

distribution of hACE2 and its correlation with SARS-CoV-2 tropism.

Although SARS-CoV-2 antigen and RNA could be detected in ACE2-expressing cells

such as olfactory neuroepithelium (ONE) and alveolar type 2 (AT2) cells, we found that

SARS-CoV-2 primarily infected neurons and alveolar type 1 (AT1) cells lacking ACE2

expression. The lethality of the K18-hACE2 model was entirely due to viral

neuroinvasion, which occurred through the olfactory neuroepithelium. This process was

accompanied by clinically detectable neurological manifestations, and was partially

driven by ACE2-independent mechanisms. Altogether, this study expands the current

knowledge on the K18-hACE2 murine model to study COVID-19. Our findings will help

refine its utilization for providing a relevant understanding of the molecular mechanisms

driving neuropathogenesis and pulmonary pathology.

RESULTS

SARS-CoV-2 is invariably fatal in infected K18-hACE2 mice with evidence of

neuroinvasion. K18-hACE2 mice inoculated intranasally with SARS-CoV-2 (1x10^6

plaque-forming units [PFU]; n=47 [n=25 male and n=12 female) began losing weight as

early as 2-3 days post-infection (dpi) irrespective of sex, with maximum weight loss

occurring at 7 dpi (17.7% $\pm$ 7.8% in male mice, 21.8% $\pm$ 3.1% in female, and combined

18.9% $\pm$ 6.9%; Figure 1A). Trends in weight loss were associated with an increase in

clinical scores and a decline in core body temperature, both of which became most pronounced near the time of death (Figure 1B-C). SARS-CoV-2-infected K18-hACE2 mice exhibited neurological signs by 6-7 dpi, characterized by tremors, proprioceptive defects, abnormal gait and imbalance. The majority of the animals were either euthanized at this time or were found dead in their cage (93%; 28/30 [Figure 1D]). At the time of death (7-10 dpi), the median clinical score was 3 (interquartile range = 2) and the mean body temperature 30.6 ± 3.7 °C. Two male mice survived to the end of the 14-day observation period. One of these animals had a maximum clinical score of 3/5 at 10 dpi and lost a maximum of 6% body weight, while the other displayed no clinical signs or appreciable weight loss over the observation period. Furthermore, neither of the two male survivors were hypothermic at any point in time, a feature that was consistently observed in animals that succumbed to disease.

When compared with C57BL/6J mice, which were concomitantly infected with SARS-CoV-2, the K18-hACE2 mice had several dark red, slightly depressed foci of variable size (2-3 mm), scattered throughout the lung lobes at 2 and 4 dpi. By 7 dpi, these foci grew into large coalescing, dark red depressions encompassing around 30% of the pulmonary parenchyma, leading to the abnormal pulmonary buoyancy (i.e. tissues sank when placed in formalin). Additionally, the mice euthanized due to clinical deterioration frequently had a markedly dilated urinary bladder filled with concentrated urine. The mice euthanized at 14 dpi (n=2) had sporadic dark red pinpoint foci in the pulmonary parenchyma.

Lethality was associated with significant viral load in the lung and brain of the K18-hACE2 mice (Figure 1E), as previously reported (Golden et al., 2020; Rathnasinghe et

al., 2020; Winkler et al., 2020; Zheng et al., 2020). In the lung, viral RNA was detectable

at the earliest experimental timepoint (2 dpi), with a mean of 6.42×10^7 RNA copies/mg,

peaked at 4 dpi (mean of 5.8×10^8 RNA copies/mg) and remained high at 7 dpi (mean of

3.71×10^7 RNA copies/mg). In contrast, viral RNA in the brain was low at 2 dpi (mean of

2.46×10^5 RNA copies/mg) but dramatically increased at 7 dpi (4-log increase)

representing the highest viral RNA load during the course of the study, with a mean of

1.27×10^9 RNA copies/mg. A small amount of viral RNA was detected in the serum

($<10^6$ RNA copies/ml). However, incubation of SARS-CoV-2 permissive cell lines (i.e.

Vero E6 cells) with serum samples did not result in any detectable productive infection

in vitro, confirming that K18-hACE2 are not viremic during infection and that small

amounts of naked viral RNA might be circulating in the bloodstream following release

from infected tissues (data not shown). Altogether, these data illustrate that lethality was

associated with increasing viral load in the brain until time of death, in contrast to lung

viral load.

Transient SARS-CoV-2 infection in the nasal cavity of K18-hACE2 mice. We next

performed detailed histologic analysis of various tissues to uncover the mechanism of

lethality in K18-hACE2 mice. For this, we first focused on the spatial and temporal

dynamics of SARS-CoV-2 infection in the upper respiratory tract and analyzed the

anterior/rostral nasal cavity and olfactory neuroepithelium for disease-associated

lesions. To do so, we performed a thorough sequential histologic analysis combined

with immunohistochemistry (IHC) and RNAscope® *in situ* hybridization (ISH) in order to

determine the cellular localization and abundance of SARS-CoV-2 antigen and RNA,

using an anti-spike monoclonal antibody and an S-specific RNA probe, respectively.

At 2 dpi, the anterior/rostral nasal cavity was characterized by mild, multifocal

neutrophilic inflammation (rhinitis) in all infected K18-hACE2 mice. In the rostral and

intermediate turbinates, histologic changes were characterized by segmental

degeneration and necrosis of the lining transitional and respiratory epithelium, with cell

rounding, loss of cilia and sloughing, along with scattered intracytoplasmic SARS-CoV-2

antigen and RNA (Figure 2). Nasal passages were partially filled with small amounts of

cellular debris, degenerate neutrophils and small numbers of erythrocytes. The lamina

propria underlying affected areas was infiltrated by low to mild numbers of neutrophils

and fewer lymphocytes (Figure 2). SARS-CoV-2 antigen and RNA were detected within

the epithelium lining the mucosa of rostral and intermediate turbinates. At 4 dpi,

epithelial degeneration and necrosis in the rostral and intermediate turbinates were no

longer observed, and histologic changes were consistent with mild, multifocal

lymphocytic rhinitis with rare clusters of few neutrophils within the lamina propria (Figure

2). Nasal passages were clear of exudate. The abundance of SARS-CoV-2 antigen and

RNA decreased, and was mostly restricted to rare positive cells in the respiratory

epithelium (Figure 2, Table 1 and Table S1). By 7 dpi, the nasal passages (including the

ONE) were histologically within normal limits and no SARS-CoV-2 antigen or RNA were

detectable in the rostral and intermediate turbinates (Figure S1).

The ONE was histologically unremarkable at all timepoints despite the variable amounts

of viral antigen and RNA detected during the course of the study. At 2 dpi, multifocal

clusters of cells within the ONE contained abundant SARS-CoV-2 antigen and RNA

(Figure 2, Table 1 and Table S1), becoming gradually more rare at 4 and 7 dpi. No

SARS-CoV-2 antigen or RNA were detected in the ONE by 14 dpi (Table 1, Table S1,

and Figure S1).

Moderate interstitial pneumonia in K18-hACE2 mice following SARS-CoV-2

infection. To evaluate the effects of SARS-CoV-2 infection in the lower respiratory

tract of K18-hACE2 mice, we performed similar analysis to that described above. In the

lower respiratory tract, histologic alterations in the pulmonary parenchyma mainly

involved the alveoli, interstitium and vasculature. Pathologic alterations in the lungs

were overall characterized by variable degree of interstitial pneumonia and variable

abundance of SARS-CoV-2 antigen and RNA, which were strictly dependent on the

experimental timepoint.

At 2 dpi, mild perivascular and peribronchiolar inflammation, consisting primarily of

lymphocytes and lesser numbers of histiocytes, and occasional perivascular edema

were evident (Figure 3). Pulmonary vessels were frequently reactive and lined by a

plump endothelium with large numbers of marginating mononuclear leukocytes and few

neutrophils (Figure 3). Multifocal to coalescing areas of the pulmonary parenchyma,

mostly associated with a bronchopulmonary segment, were composed of alveolar septa

mildly expanded by lymphocytes, histiocytes and fewer neutrophils along with scattered

air spaces containing low numbers of neutrophils and mild numbers of histiocytes and

lymphocytes (Figure 3). SARS-CoV-2 antigen and RNA were multifocally detected

within the pulmonary parenchyma, and localized in alveolar type (AT) 1 and fewer AT2

pneumocytes predominantly within and adjacent to areas of interstitial pneumonia as

demonstrated by singleplex and multiplex IHC, and RNAscope® ISH (Figure 3 and

Figure S2).

At 4 dpi, peak in viral antigen and RNA abundance was reached (correlating with the

highest viral RNA load as determined by RT-qPCR) along with a higher number of

infiltrating lymphocytes (Figure 3 and Figure S2). SARS-CoV-2 cellular tropism did not

differ from that described at 2 dpi.

At 7 dpi, there was an increase in the severity of the interstitial pneumonia, which

affected up to ~30% of the parenchyma (Figure 3). Multifocal bronchopulmonary

segments were composed of alveolar septa expanded by numerous mononuclear cells

and fewer neutrophils that extend into the alveolar spaces, occasional septal necrotic

debris, and mild proliferation of AT2 cells (Figure 3). In some animals, the affected

alveoli were multifocally denuded to completely disrupted by necrosis, and filled with

necrotic cell debris and alveolar edema. As for earlier timepoints, mild perivascular and

peribronchiolar cuffing of lymphocytes and histiocytes was a common feature. While

bronchioles were mostly unaffected, sporadically they were partially lined by an

attenuated epithelium and contain rare, individualized mononuclear cells within their

lumina. SARS-CoV-2 antigen and RNA were still abundant (Figure 3, Table 1, Figure

S2, and Table S1), albeit to a lower extent than at 4 dpi suggesting progressive

resolution of viral infection by the host consistent with the RT-qPCR data (Figure 1E).

Interestingly, viral antigen was mostly distributed in AT1 and AT2 cells within

histologically normal areas of lung adjacent to areas of intense inflammation, while

areas of subacute interstitial pneumonia and mild AT2 hyperplasia were characterized

by scant to absent SARS-CoV-2 S immunoreactivity (Figure 3, Table 1 and Table S1).

Finally, multifocal areas of similar interstitial pneumonia with sporadic aggregates of

interstitial lymphocytes and more prominent AT2 hyperplasia were identified in the two

male survivors at 14 dpi (Figure 3). No SARS-CoV-2 antigen or RNA was detected at

this time point (Figure 3 and Figure S2).

Of note, no evidence of SARS-CoV-2 infection in bronchiolar epithelium and pulmonary

vasculature was seen at any time during the course of the study (Figure 3, Table 1,

Figure S2, and Table S1). Similarly, hyaline membranes, vascular thrombosis, and

syncytial cells were not observed at any time point across all animals; which strikingly

contrasts with the human patient (Martines et al., 2020) and non-human primate studies

(Aid et al., 2020; Blair et al., 2020). Hemorrhage and pulmonary edema were only rarely

observed and of minimal-to-moderate severity. In one animal (7 dpi), there was

localized flooding of bronchioles by degenerate neutrophils and cellular debris mixed

with birefringent foreign material consistent with aspiration pneumonia, a reported rare

complication in K18 hACE2 mice infected with SARS-CoV-1 that is ultimately attributed

to pharyngeal and laryngeal dysfunction secondary to central nervous system (CNS)

disease (McCray et al., 2007).

Altogether, our data shows evidence of a significant but moderate interstitial

pneumonia in infected K18-hACE2 mice. The histopathological features and the

moderate extent of the pneumonia observed in the lung of K18-hACE2 mice therefore

contrast with those observed in severe cases of COVID-19 in humans, and suggest that

the lethality observed in this model might be independent of lung inflammatory

mechanisms.

Effective control of SARS-CoV-2 infection in the lower respiratory tract is associated with T-cell and macrophage infiltration. Subsequently, we aimed to further investigate SARS-CoV-2 tropism in the lower respiratory tract of K18-hACE2 mice, as well as the dynamics of the host response in this tissue upon infection. We first performed multiplex IHC to probe the localization of SARS-CoV-2 antigen in AT1 pneumocytes (cell maker: receptor for advanced glycation end-products [RAGE]), AT2 pneumocytes (cell marker: surfactant protein C [SPC]), and endothelial cells (cell marker: CD31). We found that SARS-CoV-2 showed preferential tropism for RAGE+ AT1 pneumocytes, as well as scattered SPC+ AT2 pneumocytes, but not for CD31+ endothelial cells (Figure 4A).

Next, we optimized and applied a 5-plex IHC to characterize the inflammatory response in the lungs of K18-hACE2 mice infected with SARS-CoV-2. At 2 dpi, the affected areas of the pulmonary parenchyma were characterized by abundant viral antigen accompanied by an early host response predominantly mediated by Iba-1+ macrophages and low numbers of CD3+CD8- T lymphocytes (presumably CD4+ T lymphocytes, Figure 4C). At 7 dpi, the parenchyma was infiltrated by CD3+CD8+ and CD3+CD8- T lymphocytes and abundant Iba-1+ macrophages with a reduction in the number of SARS-CoV-2-infected cells, the latter mostly restricted to areas of the parenchyma with less intense inflammatory response (Figure 4D). At 14 dpi, the inflammatory response was characterized by multifocal interstitial aggregates of CD19+ B lymphocytes, abundant Iba-1+ macrophages, moderate numbers of CD3+CD8- T lymphocytes, (scattered throughout the interstitium as well as associated with CD19+ B lymphocyte aggregates) and a reduction in the number of infiltrating CD3+CD8+ T

lymphocytes with undetectable SARS-CoV-2 antigen (Figure 4E). Altogether, our data

showed that control of SARS-CoV-2 in the lung of K18-hACE2 mice is associated with

moderate interstitial pneumonia characterized by a strong local lymphocytic and

histiocytic host response. Effective immune control and moderate inflammation once

again suggest that lethality observed in K18-hACE2 mice is attributable to lung-

independent disease.

Severe SARS-CoV-2 neuroinvasion and neurological damage in K18-hACE2 mice.

Pursuing our hypothesis that the lethality of the K18-hACE2 mice is associated with

brain neuroinvasion, we analyzed sagittal sections of the brain including representation

of the cribriform plate and olfactory bulb at different timepoints post-infection (2, 4, 6-7

and 14 dpi) for histologic changes and viral antigen distribution (Figures 5 and 6).

Histologic alterations in the brain were first observed as early as 6 dpi but mostly at 7

dpi, and consisted of mild, multifocal neuronal spongiosis primarily throughout the

cerebral cortical layers of the somatomotor, somatosensory and visual areas (Figure 5).

Additionally, multifocal blood vessels were delimited by delicate cuffs of few

lymphocytes with mild numbers of reactive glial cells in the adjacent neuroparenchyma.

At 7 dpi, histologic alterations worsened in severity and became more widespread with

involvement of the olfactory bulb, cerebral cortex (most predominantly somatosensory

and somatomotor areas), hippocampus (mainly CA1 region), brainstem (thalamus),

midbrain and the dentate nucleus ventral to the cerebellum (Figure 5). At the level of the

olfactory bulb, the glomerular, external plexiform, and mitral cell layers were

characterized by moderate to marked neuropil vacuolization (spongiosis) with

occasional vacuoles containing intralesional cell debris. Elsewhere, the grey and white

matter within affected areas of the neuroparenchyma were extensively and predominantly characterized by marked spongiosis with frequent clear vacuoles containing intralesional cell debris, and multifocal shrunken, angular, hypereosinophilic and pyknotic neuronal bodies with loss of Nissl substance/chromatolysis (neuronal degeneration and necrosis, Figure 5) and occasionally delimited by multiple glial cells (satellitosis). There were multifocal, delicate perivascular cuffs composed of lymphocytes and the adjacent neuroparenchyma had an increase in the number of reactive glial cells (gliosis). Notably, the cerebellum (cortical layers and associated white matter of the cerebellar folia) was spared of histologic changes.

Neuronal changes correlated with abundant neuronal immunoreactivity for SARS-CoV-2 S antigen and viral RNA, with exclusive localization within the perikaryon and neuronal processes (Figure 6). SARS-CoV-2 antigen and RNA had a widespread distribution throughout the brain in roughly 85% (11/13) of infected K18-hACE2 mice at 7 dpi, including neuronal bodies within the cerebral cortex, CA1, CA2 and CA3 regions of the hippocampus, anterior olfactory nucleus, caudoputamen, nucleus accumbens, thalamic nuclei including hypothalamus, midbrain, pons and medulla oblongata nuclei (Figure 6). Few vestibulocochlear nerve fascicles showed immunoreactivity for viral antigen; while no viral S antigen or RNA was detected in areas spared of histological changes including the cerebellar cortex and white matter, optic nerve and retina, and the spiral ganglion of the inner ear (albeit the eye and inner ear were not present in the majority of sections examined). SARS-CoV-2 S antigen and RNA preceded histological findings with rare viral antigen and RNA detected as early as 4 dpi in mitral and inner nuclear

neurons of the olfactory bulb, as well as small clusters of neurons within the anterior olfactory nucleus and orbital area of the cerebral cortex (Figure 6).

Multiplex IHC for co-detection of SARS-CoV-2 antigen, Iba-1+ and GFAP+ glial cells was performed at 7 dpi, timepoint at which widespread viral antigen and RNA along with histological lesions were observed. Multiple IHC revealed that the gliosis in the brain of infected mice was attributed to abundant GFAP+ astrocytes as well as Iba-1+ microglial cells, which did not colocalize with SARS-CoV-2 S protein expression (Figure 7). Iba-1+ microglial cells with broad cytoplasmic ramifications were seen to be tightly associated with SARS-CoV-2+ neuronal bodies, suggestive of an active host response against infection (Figure 7).

We evaluated the extent of the CNS involvement in infected K18-hACE2 mice at 7 dpi by additionally examining the cervicothoracic and lumbosacral segments of the spinal cord. In 9/11 infected K18-hACE2 mice that showed widespread viral antigen in the brain, the spinal cord also had mild-to-moderate viral antigen that predominated within the cervicothoracic segments (Figure S3, Table 1 and Table S1). Neuronal changes were similar to those noted in affected areas of the brain; however, gliosis and perivascular cuffing were sparse.

Finally, Luxol Fast Blue was utilized to assess myelin loss (demyelination) following SARS-CoV-2 invasion in the brain and spinal cord at 7 dpi. No evidence of demyelination was noted.

Taken together, our data show that SARS-CoV-2 infection of K18-hACE2 results in severe neuronal invasion of the CNS, likely via retrograde transport through the

olfactory bulb originating from axonal processes traversing the ONE. Viral neuroinvasion resulted in extensive cytopathic effect in neurons, comprising not only the brain but also the spinal cord.

ACE2 expression and distribution does not fully reflect SARS-CoV-2 tissue tropism in K18-hACE2 mice. To further explore the mechanism driving lethal SARS-CoV-2 infection in K18-hACE2 mice, we first investigated the tissue and cellular distribution of the ACE2 receptor in both C57BL/6J and K18-hACE2 mice by IHC using a cross-reactive anti-ACE2 antibody (cross-reactive to hACE2 and mACE2) (Table S2). In the lower respiratory tract (lungs), ACE2 was ubiquitously expressed along the apical membrane of bronchiolar epithelium and, less commonly, in rare and scattered AT2 pneumocytes (Figure 8). No ACE2 expression was found in AT1 pneumocytes. Strikingly, no differences in the distribution and abundance of ACE2 expression were identified between uninfected C57BL/6J, sham-inoculated K18-hACE2, and terminal (7 dpi) K18-hACE2 mice inoculated with SARS-CoV-2 (Figure 8).

We therefore aimed at analyzing expression and distribution of *hACE2* mRNA using RNAscope® ISH. Although no expression of *hACE2* mRNA was detected in the lungs of non-transgenic C57BL/6J mice (Figure 9), expression of *hACE2* mRNA was detectable but low in the lungs of K18-hACE2 mice, and mostly involved bronchiolar epithelial cells with sporadic expression within few pneumocytes (Figure 9). In summary, these findings indicate that *hACE2* expression in the lungs of K18-hACE2 is low, and that expression of *hACE2* is not the sole host factor determinant of susceptibility to SARS-CoV-2. This is clearly exemplified by the following: 1) certain cell types that, while expressing

hACE2, were non-permissive to SARS-CoV-2 infection throughout the experiment (i.e.

bronchiolar epithelial cells); and 2) the near diffuse infection of AT1 cells by 4 dpi

despite sparse to absent expression of *hACE2* in these cells. These observations

support evidence for an ACE2-independent viral entry mechanism playing a major role

in the pulmonary dissemination of K18-hACE2 mice.

In contrast to the lung, increased ACE2 expression was clearly evident in the nasal

cavity of K18-hACE2 mice compared to C57BL/6J mice. We assessed ACE2

expression on the rostral nasal epithelium, respiratory epithelium at the level of the

intermediate turbinates (Figure 8) as well as in the ONE and olfactory bulb (Figure 8).

Unlike C57BL/6J mice, in which ACE2 was undetectable within the nasal cavity, ACE2

was diffusely expressed within the apical membrane of transitional and respiratory

epithelium, and segmentally within the apical surface of the ONE in both sham-

inoculated and SARS-CoV-2-infected K18-hACE mice (Figure 8). While minimal to rare

expression of *hACE2* mRNA was identified in the neurons of the mitral layer of the

olfactory bulb, olfactory neuroepithelium and respiratory epithelium of rostral turbinates,

estimation of its abundance and distribution could not be fully assessed since the

decalcification procedure had a significant impact in the quality of cellular mRNA as

demonstrated by the low detection of the housekeeping mRNA, *Ppib* (data not shown).

In the brain of both C57BL/6J and K18-hACE2 mice, ACE2 immunoreactivity was

observed in the vascular endothelium lining blood vessels (Figure 8), as well as

ependymal and choroid plexus epithelium. In contrast, distribution of *hACE2* mRNA

expression involved clusters of neurons within the cerebral cortex, hippocampus,

midbrain, brainstem and Purkinje cells from the cerebellum, with no expression noted in

non-transgenic C57BL/6J mice (Figure 10 and Figure S4). There was no expression of

hACE2 mRNA in vascular endothelial cells. Taken together, our data show a

discrepancy between ACE2 protein and RNA expression and distribution within the

CNS. This is partly attributable to the fact that the ACE2 antibody we utilized cross

reacts with both hACE2 and mACE2, while the *ACE2* probe employed was human

specific. The absence of *hACE2* hybridization with simultaneous ACE2

immunoreactivity in the capillary endothelium supports the notion that ACE2 expression

in these cells is of murine origin. The absence of ACE2 immunoreactivity in neurons is

suggestive of a potential restriction in the translation (or post-translation) of the ACE2

receptor in these cells. This, in addition to the fact that Purkinje cells of the cerebellum

are non-permissive to SARS-CoV-2 infection in spite of the expression of *hACE2*

mRNA, also suggests that ACE2 is likely not the sole host factor associated with

neuroinvasion and that other ACE2-independent entry mechanisms contribute to

neuroinvasion and spread by SARS-CoV-2 in this murine model.

Absence of infection and histologic lesions in extrapulmonary and extraneural

tissues despite ACE2 expression. Other tissues examined included heart, kidney,

stomach, duodenum, jejunum, ileum, cecum, and colon. All of these were histologically

within normal limits (data not shown). No SARS-CoV-2 S antigen was detected in any of

these tissues at any time point (data not shown). ACE2 distribution was evaluated in

sections of the heart, stomach, small intestine and colon. While ACE2 expression was

limited to the capillary vascular endothelium in the heart and stomach, intense

expression was noted in the non-glandular mucosa of the stomach and apical surface of

enterocytes lining the small intestinal mucosa. Colonic enterocytes rarely expressed

ACE2 (Figure S5).

DISCUSSION

The K18-hACE2 transgenic mouse model has become a widespread laboratory animal

model suitable for studying SARS-CoV-2 pathogenesis as well as medical

countermeasures against COVID-19 (Johansen et al., 2020). The suitability of this

model relies on the common host entry receptor shared between SARS-CoV-1 and

SARS-CoV-2 (Hoffmann et al., 2020; Li et al., 2003), and transgenic mice expressing

hACE2 under the K18 promoter develop lethal clinical disease associated with

pulmonary pathology and neuroinvasion, with high viral titers (Golden et al., 2020;

McCray et al., 2007; Moreau et al., 2020; Oladunni et al., 2020; Rathnasinghe et al.,

2020; Winkler et al., 2020; Yinda et al., 2020; Zheng et al., 2020). In contrast, other

murine models of SARS-CoV-2 (e.g. adenovirus-transduced hACE2 mice and

transgenic mice expressing hACE2 under the HFH4 promoter) develop only mild

disease with limited and short-lived viral replication and pulmonary pathology, and low

to no lethality (Jiang et al., 2020; Rathnasinghe et al., 2020). While the K18-hACE2

murine model has been critical in shedding light on mechanisms of lung injury and

dysfunction, it fails to faithfully recapitulate several key histologic features of severe and

lethal cases of COVID-19 in humans, such as diffuse alveolar damage (DAD) with

hyaline membrane formation and multi-organ failure associated with hypercoagulability

and widespread microthrombi formation (Maiese et al., 2020; Martines et al., 2020).

In order to better understand the pathogenesis of SARS-CoV-2, well-characterized animal models are critically needed (Munoz-Fontela et al., 2020). Even though the K18-hACE2 murine model is currently under extensive use, several aspects associated with the temporospatial dynamics of SARS-CoV-2 infection remain poorly characterized, including the expression and cellular distribution of hACE2. We attempted to further characterize pathological aspects related to viral pathogenesis in this unique murine model, and hypothesized that the temporospatial distribution of SARS-CoV-2 and pathological outcomes following infection in the K18-hACE2 murine model is partially but not solely associated with hACE2 and that increased lethality in this model is related to neuroinvasion. The study presented herein provides additional novel information regarding the temporal and spatial aspects of SARS-CoV-2 infection in the K18-hACE2 mouse model with emphasis on pathological outcomes as well as a thorough and methodical characterization of ACE2 expression in this transgenic mouse model, which contributes to our understanding of this critical model used for preclinical evaluation of vaccines and antiviral therapeutics. The results presented herein not only demonstrate that lethality of this murine model is associated with neuroinvasion and subsequent neuronal cytopathic effect, but that SARS-CoV-2 tropism is not solely restricted to ACE2-expressing cells in K18-hACE2 mice. Thus, the neuropathogenic potential of SARS-CoV-2 is dependent on other host factors.

In this study, we have utilized a large cohort of K18-hACE2 mice (n=50) in order to sequentially evaluate SARS-CoV-2 tropism and pathological alterations, spatial and temporal analysis of host factors including inflammatory response and ACE2/*hACE2* expression, and survival curve analysis for a period of 14 dpi. Similarly to previous

studies, which only observed the outcome of SARS-CoV-2 for up to 7 dpi (Golden et al., 2020; Moreau et al., 2020; Oladunni et al., 2020; Rathnasinghe et al., 2020; Winkler et al., 2020), infected K18-hACE2 mice initially developed significant clinical disease with marked weight loss and increased respiratory effort associated with the development of interstitial pneumonia and high viral load as determined via IHC, ISH and RT-qPCR; an unquestionable feature of this model. Survival curve analysis clearly demonstrated that lethality in infected mice only occurs after 6 dpi and in the vast majority of mice (96.3%), coincided with the initiation of neurologic signs and/or symptoms, neuronal cytopathic effect, and abundance of viral S antigen and RNA in the CNS. These observations are a clear indication of the rapid and fatal neuroinvasive nature of this model. Our study also demonstrates that SARS-CoV-2 has a tropism for motor neurons within the spinal cord (predominantly within the cervicothoracic segments), which was only observed in mice with concurrent brain involvement suggesting a descending infection. This finding helps rationalize the neurologic signs observed with this model including decreased mobility/responsiveness and decreased urine voiding that was evident at necropsy, reflective of severe urinary bladder dilation and accumulation of concentrated urine. Given the spinal cord involvement, the latter is likely attributed to altered spinal reflexes and/or decreased intervention of the detrusor muscle, which is required for normal micturation. An additional striking clinical feature in infected K18-hACE2 mice at 7 dpi was hypothermia (mean body temperature of 30.6 °C), which is likely a consequence of hypothalamic and motor neuron dysfunction associated with SARS-CoV-2 neurotropism and could serve as a clinical indicator of CNS involvement. Our results unequivocally demonstrate that neuroinvasion represents the driving component of fatality in this

animal model compared to others such as Syrian hamsters, which display more severe

pulmonary disease and infection of the ONE, but no evidence of neuroinvasion (Bryche

et al., 2020). Furthermore, these animals invariably recover within 14 days following

intranasal infection with SARS-CoV-2 (Bryche et al., 2020; Imai et al., 2020; Osterrieder

et al., 2020; Rosenke et al., 2020; Sia et al., 2020). Very few infected K18-hACE2 mice

(2/30) from our cohort survived until the end of the study (14 dpi) and, while ameliorated

pulmonary lesions were noticed in these, they did not exhibit involvement of the CNS

further supporting the observation that mice suffering from SARS-CoV-2 neuroinvasion

invariably succumb at earlier time points post-infection. Both of these survivor mice

developed pulmonary interstitial aggregates of B lymphocytes that were distinct from

earlier time points, which could be suggestive of the development of protective adaptive

immunity. The normal histologic appearance of the CNS in surviving mice and lack of

any residual neurologic signs supports the notion that animals can fully recover from

infection, albeit we acknowledge that limited neuroinvasion in these animals could not

be disproven. Furthermore, we acknowledge that extensive neurobehavior testing,

which is beyond the expertise of the authors, would be required to rule out any long-

term sequelae in the rare instance of survivors. Overall, these findings are of

importance to researchers with a particular interest in studying SARS-CoV-2-associated

neuropathogenesis, as premature euthanasia due to other clinical features (i.e., weight

loss, ruffled fur, and/or respiratory distress) have the potential to precede CNS disease

and such terminal endpoints, if elected, may preclude evaluation of the effects of SARS-

CoV-2 in the CNS. Instead decreased responsiveness/mobility and tremors are

interpreted to reflect better clinical findings supportive of neuroinvasive disease.

To date, the precise mechanism(s) enabling neuroinvasion in the K18-hACE2 model is

poorly understood (Bryche et al., 2020; DosSantos et al., 2020; Ellul et al., 2020; Liu et

al., 2020; Solomon et al., 2020). Here, we determined that K18-hACE2 transgenic mice

show a significant upregulation in the expression of ACE2 in the nasal cavity

(particularly in the neuroepithelium) compared to wild-type C57BL/6J mice, in which

ACE2 expression is undetectable by IHC. This difference between K18-hACE2 and

C57BL/6J mice is clearly attributed to the expression of the *hACE2* transgene, and is a

key feature to the neuropathogenesis of this model. Interestingly, temporal analysis of

SARS-CoV-2 S antigen and RNA in the ONE of transgenic mice preceded and/or

occurred simultaneously with infection of neurons within the glomerular and mitral layers

of the olfactory bulb, indicative of possible viral retrograde transport through axonal

processes traversing the cribriform plate and allowing viral spread within cortical

neurons and beyond. Expression of *hACE2* within neurons in the CNS is overall low and

does not correlate with our immunohistochemical findings, where ACE2 protein was

restricted to capillary endothelium, ependymal and choroid epithelium with sparing of

neurons and their processes. These findings suggest the ACE2 expression in these

anatomical compartments could be attributed to *mACE2* or indicative of a post-

transcriptional event that could be limiting neuronal expression of *hACE2*. These along

with the fact that *hACE2* mRNA is not abundantly and equally expressed among

different neuronal populations and that Purkinje cells in the cerebellum express *hACE2*

mRNA, but are not permissive to SARS-CoV-2 infection, suggest that entry of SARS-

CoV-2 into neurons is likely mediated by other host receptors independent of ACE2.

Infection of brain organoids has been shown to be inhibited by the use of anti-ACE2

antibodies (Song et al., 2020). However, brain organoids do not recapitulate the

complexity of the entire CNS, and retrograde transport of viral particles into the CNS

can hardly be modeled *in vitro*. Altogether, this suggests that while ACE2 is assuredly

an important mediator of CNS neuroinvasion, *in vitro* platforms to study mechanisms of

SARS-CoV-2 neuroinvasion are limiting, and require the use of more complex

experimental systems. Recently, neuropilin-1, a transmembrane glycoprotein serving as

cell surface receptor for semaphorins and other ligands, was shown to serve as an

alternative receptor for SARS-CoV-2 mediating entry into the neuroepithelium and

neurons (Cantuti-Castelvetri et al., 2020). Even though we analyzed the expression of

neuropilin-1 in this study (data not shown), we observed ubiquitous expression in the

nasal passages, brain, kidneys, liver and lungs, precluding any definitive conclusions in

support or against these claims (Cantuti-Castelvetri et al., 2020).

Anosmia and ageusia (loss of smell and taste, respectively) represent the earliest and

most common but transient neurologic symptoms in people with COVID-19, being

reported in ≥ 50% of cases (Eliezer et al., 2020; Ellul et al., 2020; Walker et al., 2020).

Hyposmia or anosmia has also been clearly characterized in K18-hACE2 mice,

occurring between 2-3 dpi, which was characterized through a series of unique

behavioral tests requiring a normal sense of smell (Zheng et al., 2020). Other neurologic

manifestations of COVID-19 are associated with acute cerebrovascular disease, with

cohort studies reporting strokes in 2–6% of hospitalized patients (Ellul et al., 2020;

Wang et al., 2020b). Long-term neurologic sequelae associated with COVID-19 or its

effect on neurodegenerative diseases remain unclear (Wang et al., 2020b). Very little is

known about the pathogenesis of these neurologic manifestations and whether they are directly or indirectly associated with SARS-CoV-2. ACE2 expression has been described in humans both in health and with chronic rhinosinusitis, with expression noted in sustentacular cells of the ONE, but not within immature and mature olfactory neurons (Chen et al., 2020). This observation led the authors to suggest that anosmia in COVID-19 is likely attributable to an indirect effect of SARS-CoV-2 infection. However, recent studies evaluated brain and nasal autopsies from patients who died of COVID-19, in which SARS-CoV-2 antigen and RNA was detected in cells of neural origin within the ONE and cortical neurons occasionally associated with locally ischemic regions (Meinhardt et al., 2020; Song et al., 2020). These studies provide conclusive evidence that the K18-hACE2 mice serves as a model with translational significance, even though ischemic lesions have not been reported including those in our study, further justifying its use to dissect the intricate mechanisms involved in SARS-CoV-2-mediated CNS invasion. Even though SARS-CoV-2 infects sustentacular cells within the neuroepithelium of Syrian hamsters (Bryche et al., 2020), the K18-hACE2 and a CRISPR/Cas9 hACE2 knock-in mouse model (Sun et al., 2020) are the only models that develop neuroinvasion with wild-type virus and, thus, will be particularly useful for studying SARS-CoV-2 neuropathogenesis, particularly the mechanisms of viral trafficking of into the CNS through the neuroepithelium.

Another important observation of the K18-hACE2 model is that SARS-CoV-2 tropism extensively involves infection of ACE2 and *hACE2* negative cells, including certain population of neurons and the vast majority of AT1 pneumocytes. Similarly, sole expression of *hACE2* in some cell types (i.e., bronchiolar epithelial cells) clearly does

not render these cells susceptible to SARS-CoV-2 even following intranasal exposure,

and underscore that other undetermined host factors are most likely required to allow

viral entry. Therefore, this model is extremely relevant for investigating the role of

ACE2-independent entry.

In conclusion, this study provides a comprehensive spatiotemporal analysis of SARS-

CoV-2 infection in the K18-hACE2 transgenic murine model along with an analysis of

the contribution of ACE2 in the permissiveness of the model. Our work provides

evidence that SARS-CoV-2 exhibits a marked neurotropism, and that this process likely

occurs through mechanisms that are in part hACE2-independent. Lethal CNS invasion,

combined with the absence of several of the pulmonary hallmarks associated with

severe human COVID-19, therefore calls for attentive caution when utilizing the K18-

hACE2 mouse model to investigate certain aspects of SARS-CoV-2 pathogenesis. The

protective ability of anti-viral therapies and T-cell based vaccines against lethal

challenge in this model might indeed be then underestimated due to the increased

susceptibility to CNS invasion. Regardless, the K18-hACE2 mouse model represents a

promising model for understanding the mechanisms governing SARS-CoV-2

neuroinvasion, ACE2-independent virus entry, and evaluating potent and fast-acting

prophylactic countermeasures.

MATERIALS AND METHODS

Biosafety. All aspects of this study were approved by the Institutional Biosafety

Committee and the office of Environmental Health and Safety at Boston University prior

to study initiation. Work with SARS-CoV-2 was performed in a biosafety level-3

laboratory by personnel equipped with powered air-purifying respirators.

Cells and viruses. African green monkey kidney Vero E6 cells (ATCC® CRL-1586™,

American Type Culture Collection, Manassas, VA) were maintained in Dulbecco's

minimum essential medium (DMEM; Gibco, Carlsbad, CA [#11995-065]) containing

10% fetal bovine serum (FBS, ThermoFisher Scientific, Waltham, MA), 1X non-essential

amino acids (ThermoFisher Scientific), penicillin and streptomycin (100 U/ml and 100

µg/ml), and 0.25 µg/ml of amphotericin B (Gibco®, Carlsbad, CA), and incubated at 37

°C and 5% CO_2 in a humidified incubator.

SARS-CoV-2 isolate stock preparation and titration. All replication-competent

SARS-CoV-2 experiments were performed in a biosafety level 3 laboratory (BSL-3) at

the Boston University' National Emerging Infectious Diseases Laboratories. 2019-

nCoV/USA-WA1/2020 isolate (NCBI accession number: MN985325) of SARS-CoV-2

was obtained from the Centers for Disease Control and Prevention (Atlanta, GA) and

BEI Resources (Manassas, VA). To generate the passage 1 (P1) virus stock, Vero E6

cells, pre-seeded the day before at a density of 10 million cells, were infected in T175

flasks with the master stock, diluted in 10 ml final volume of Opti-MEM (ThermoFisher

Scientific). Following virus adsorption to the cells at 37 ºC for 1 h, 15 ml DMEM

containing 10% FBS and 1X penicillin/streptomycin was added to the flask. The next

day, media was removed, the cell monolayer was rinsed with 1X phosphate buffered

saline (PBS) pH 7.5 (ThermoFisher Scientific) and 25 ml of fresh DMEM containing 2%

FBS was added. Two days later, when the cytopathic effect of the virus was clearly

visible, culture medium was collected, filtered through a 0.2 µm filter, and stored at -80

ºC. Our P2 working stock of the virus was prepared by infecting Vero E6 cells with the

P1 stock, at a multiplicity of infection (MOI) of 0.1. Cell culture media was harvested at 2

and 3 dpi, and after the last harvest, ultracentrifuged (Beckman Coulter Optima L-100k;

SW32 Ti rotor) for 2 h at 25,000 rpm (80,000 X g) over a 20% sucrose cushion (Sigma-

Aldrich, St. Louis, MO). Following centrifugation, the media and sucrose were discarded

and pellets were left to dry for 5 minutes at room temperature. Pellets were then

resuspended overnight at 4 ºC in 500 µl of 1X PBS. The next day, concentrated virions

were aliquoted at stored at -80 ºC.

The titer of our viral stock was determined by plaque assay. Vero E6 cells were seeded

into a 12-well plate at a density of 2.5 x 10^5 cells per well and infected the next day with

serial 10-fold dilutions of the virus stock for 1 h at 37 ºC. Following virus adsorption, 1 ml

of overlay media, consisting of 2X DMEM supplemented with 4% FBS and mixed at a

1:1 ratio with 1.2% Avicel (DuPont; RC-581), was added in each well. Three days later,

the overlay medium was removed, the cell monolayer was washed with 1X PBS and

fixed for 30 minutes at room temperature with 4% paraformaldehyde (Sigma-Aldrich).

Fixed cells were then washed with 1X PBS and stained for 1h at room temperature with

0.1% crystal violet (Sigma-Aldrich) prepared in 10% ethanol/water. After rinsing with tap

water, the number of plaques were counted and the virus titer was calculated. The titer

of our P2 virus stock was 4 x 10^8 plaque forming units (PFU)/ml.

Mice. Mice were maintained in a facility accredited by the Association for the

Assessment and Accreditation of Laboratory Animal Care (AAALAC). All protocols were

approved by the Boston University Institutional Animal Care and Use Committee

(PROTO202000020). Heterozygous K18-hACE2 C57BL/6J mice of both sexes (strain:

2B6.Cg-Tg(K18-ACE2)2Prlmn/J) were obtained from the Jackson Laboratory (Jax, Bar

Harbor, ME). Animals were group-housed by sex in Tecniplast green line individually

ventilated cages (Tecniplast, Buguggiate, Italy). Mice were maintained on a 12:12 light

cycle at 30-70% humidity and provided ad-libitum water and standard chow diets

(LabDiet, St. Louis, MO).

Intranasal inoculation with SARS-CoV-2. At 4 months of age, K18-hACE2 mice of

both sexes were intranasally inoculated with 1 x 10^6 PFU of SARS-CoV-2 in 50 µl of

sterile 1X PBS (n=47 [n=25 male and n=12 female), or sham inoculated with 50 µl of

sterile 1X PBS (n=3; female). Inoculations were performed under 1-3% isoflurane

anesthesia. Animals were either enrolled in a 14-day survival curve study (n=33).

Animals were only euthanized early if they reached euthanasia criteria, or if the

experiment included predetermined euthanasia timepoints for tissue sampling (2, 4, or 7

dpi; n=15).

Clinical monitoring. Animals included in the 14-day survival curve study were

intraperitoneally implanted with an RFID temperature-monitoring microchip (Unified

Information Devices, Lake Villa, IL, USA) 48-72 hours prior to inoculation. An IACUC-

approved clinical scoring system was utilized to monitor disease progression and

establish humane endpoints (Table 2). Categories evaluated included body weight,

general appearance, responsiveness, respiration, and neurological signs for a

maximum score of 5. Animals were considered moribund and humanely euthanized in

the event of the following: a score of 4 or greater for 2 consecutive observation periods,

weight loss greater than or equal to 20%, severe respiratory distress, or lack of

responsiveness. Clinical signs and body temperature were recorded once per day for

the duration of the study.

Tissue processing and viral RNA isolation. Tissues were collected from mice and

stored in 600 µl of RNA*later* (Sigma-Aldrich; # R0901500ML) and stored at -80 °C. For

processing, 20 – 30 mg of tissue were placed into a 2 ml tube with 600 µl of RLT buffer

with 1% β-mercaptoethanol and a 5 mm stainless steel bead (Qiagen, Valencia, CA;

#69989). Tissues were then dissociated using a Qiagen TissueLyser II (Qiagen) with

the following cycle parameters: 20 cycles/s for 2 min, 1 min wait, 20 cycles/s for 2 min.

Samples were centrifuged at 17,000 X g (13,000 rpm) for 10 minutes and supernatant

was transferred to a new 1.5 ml tube. Viral RNA isolation was performed using a Qiagen

RNeasy Plus Mini Kit (Qiagen; #74134), according to the manufacturer's instructions,

with an additional on-column DNase treatment (Qiagen; #79256). RNA was finally

eluted in 30 µl of RNase/DNase-free water and stored at -80 °C until used.

RNA isolation from serum. Total viral RNA was isolated from serum using a Zymo

Research Corporation Quick-RNA^{TM} Viral Kit (Zymo Research, Tustin, CA; #R1040)

according to the manufacturer's instructions. RNA was eluted in 15 µl of RNase/DNase-

free water and stored at -80 °C until used.

SARS-CoV-2 E-specific reverse transcription quantitative polymerase chain

reaction (RT-qPCR). Viral RNA was quantitated using single-step RT-quantitative real-

time PCR (Quanta qScript One-Step RT-qPCR Kit, QuantaBio, Beverly, MA; VWR;

#76047-082) with primers and TaqMan® probes targeting the SARS-CoV-2 E gene as

previously described (Corman et al., 2020). Briefly, a 20 µl reaction mixture containing

10 µl of Quanta qScript™ XLT One-Step RT-qPCR ToughMix, 0.5 µM Primer

E_Sarbeco_F1 (ACAGGTACGTTAATAGTTAATAGCGT), 0.5 µM Primer

E_Sarbeco_R2 (ATATTGCAGCAGTACGCACACA), 0.25 µM Probe E_Sarbeco_P1

(FAM-ACACTAGCCATCCTTACTGCGCTTCG-BHQ1), and 2 µl of template RNA was

prepared. RT-qPCR was performed using an Applied Biosystems QuantStudio 3

(ThermoFisher Scientific) and the following cycling conditions: reverse transcription for

10 minutes at 55 °C, an activation step at 94 °C for 3 min followed by 45 cycles of

denaturation at 94 °C for 15 seconds and combined annealing/extension at 58 °C for 30

seconds. Ct values were determined using QuantStudio™ Design and Analysis software

V1.5.1 (ThermoFisher Scientific). For absolute quantitation of viral RNA, a 389 bp

fragment from the SARS-CoV-2 E gene was cloned onto pIDTBlue plasmid under an

SP6 promoter using NEB PCR cloning kit (New England Biosciences, Ipswich, MA).

The cloned fragment was then *in vitro* transcribed (mMessage mMachine SP6

transcription kit; ThermoFisher) to generate an RT-qPCR standard.

Serum infectivity assay. Vero E6 cells were plated in a 24-well plate at a density of

50,000 cells per well. To assess for infectious particles, 20 µl of serum was added onto

cells with 250 µl of DMEM supplemented with 2% FBS and 1% penicillin and

streptomycin and incubated for 6 hours at 37 °C. After incubation, serum was removed,

cells were washed twice with 1 ml of 1X PBS, and 1 ml of DMEM supplemented with

2% FBS 1% penicillin and streptomycin was added to each well. Media was collected

24- and 96-hours post infection for RT-qPCR analysis.

Histology. Animals were anesthetized with 1-3% isoflurane and euthanized with an

intraperitoneal overdose of ketamine and xylazine before harvest and fixation of tissues.

Lungs were insufflated with ~1.5mL of 1% low melting point agarose (Sigma-Aldrich)

diluted in 1X PBS using a 24-gauge catheter placed into the trachea. The skull cap was

removed and the animal decapitated and immersed in fixative. Additional tissues

harvested included the heart, kidneys, and representative sections of the

gastrointestinal tract, which included the duodenum, jejunum, ileum, cecum, and colon.

Tissues were inactivated in 10% neutral buffered formalin at a 20:1 fixative to tissue

ratio for a minimum of 72 hours before removal from BSL-3 in accordance with an

approved institutional standard operating procedure. Following fixation, the whole head

was decalcified in Immunocal™ Decalcifier (StatLab, McKinney, TX) for 7 days before

performing a mid-sagittal section dividing the two hemispheres into even sections.

Tissues were subsequently processed and embedded in paraffin following standard

histological procedures. Five-micron sections were obtained and stained with

hematoxylin and eosin or Luxol Fast Blue (myelin stain).

Immunohistochemistry and RNAscope[®] *in situ* hybridization.

Immunohistochemistry (IHC) was performed using a Ventana BenchMark Discovery

Ultra autostainer (Roche Diagnostics, Indianapolis, IN) using tyramide signaling

amplification (TSA) technology. Specific IHC assay details including antibodies, antigen

retrieval, sequence of multiplex assays, and incubation periods are found in Table S2.

SARS-CoV-2 S was semiquantitatively scored as follows: 0, no viral antigen observed;

1, up to 5% positive cells per 400X field examined; 2, 5-25% positive cells per 400X

field examined; and 3, up to 50% positive cells per 400X field examined.

For SARS-CoV-2 RNAscope[®] ISH, an anti-sense probe targeting the spike (S;

nucleotide sequence: 21,563-25,384) of SARS-CoV-2, USA-WA1/2020 isolate

(GenBank accession number MN985325.1) was used as previously described

(Carossino et al., 2020; Gaudreault et al., 2020). The RNAscope[®] ISH assay was

performed using the RNAscope 2.5 LSx Reagent Kit (Advanced Cell Diagnostics,

Newark, CA) on the automated BOND RXm platform (Leica Biosystems, Buffalo Grove,

IL) as described previously (Meekins et al., 2020). Briefly, four-micron sections of

formalin-fixed paraffin-embedded tissue were subjected to automated baking and

deparaffinization followed by heat-induced epitope retrieval (HIER) using a ready-to-use

EDTA-based solution (pH 9.0; Leica Biosystems) at 100 °C for 15 min. Subsequently,

tissue sections were treated with a ready-to-use protease (RNAscope® 2.5 LSx

Protease) for 15 min at 40 °C followed by a ready-to-use hydrogen peroxide solution for

10 min at room temperature. Slides were then incubated with the ready-to-use probe

mixture for 2 h at 40 °C, and the signal amplified using a specific set of amplifiers

(AMP1 through AMP6 as recommended by the manufacturer). The signal was detected

using a Fast-Red solution for 10 minutes at room temperature. Slides were

counterstained with a ready-to-use hematoxylin for 5 min, followed by five washes with

1X BOND Wash Solution (Leica Biosystems) for bluing. Slides were finally rinsed in

deionized water, dried in a 60 °C oven for 30 min, and mounted with Ecomount[®]

(Biocare, Concord, CA, USA). A SARS-CoV-2-infected Vero E6 cell pellet was used as

a positive assay control. For all assays, an uninfected mouse was used as a negative

control.

For *hACE2* mRNA RNAscope[®] ISH, an anti-sense probe targeting *hACE2* (GenBank

accession number NM_021804.3; Cat. No. 848038) with no cross-reactivity to murine

Ace2 was used in a similar manner as described above with the exception that AMP5

and AMP6 were incubated for 45 min and 30 min, respectively. Murine *peptidylprolyl*

isomerase B (Ppib) mRNA was used as a housekeeping gene to determine RNA quality

and a Vero E6 cell pellet was used as a positive assay control.

Multispectral microscopy. Fluorescently labeled slides were imaged using a Mantra

2.0^TM Quantitative Pathology Workstation (Akoya Biosciences, Marlborough, MA). To

maximize signal-to-noise ratios, images were spectrally unmixed using a synthetic

library specific for the Opal fluorophores used for each assay and for 4′,6-diamidino-2-

phenylindole (DAPI). Furthermore, an unstained section was used to create an

autofluorescence signature that was subsequently removed from whole-slide images

using InForm software version 2.4.8 (Akoya Biosciences).

Statistical analysis. Descriptive statistics and graphics as well as Kaplan-Meier

(survival) curves and statistical tests were performed using GraphPad Prism statistical

analysis software (GraphPad, San Diego, CA). Clinical parameters were analyzed using

a Student *t*-test. Significance levels were set at p-value<0.05 in all cases.

FUNDING

We thank the Evans Center for Interdisciplinary Biomedical Research at Boston

University School of Medicine for their support of the Affinity Research Collaborative on

'Respiratory Viruses: A Focus on COVID-19'. This work utilized a Ventana Discovery

Ultra autostainer that was purchased with funding from a National Institutes of Health

SIG grant (S10-OD026983). This work was also supported by a Boston University Start-

up fund, and a Peter Paul Career Development Professorship (to F.D.). This study was

also partially supported by start-up funds provided by the School of Veterinary Medicine,

Louisiana State University to Dr. Udeni Balasuriya (PG002165) and by pilot funding to

Dr. Markus Bosmann from the National Institutes of Health grant 1UL1TR001430.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest.

AUTHOR CONTRIBUTIONS: F. Douam, N. Crossland, M. Carossino, U. Balasuriya

and M. Bosmann designed the study; F. Douam, N Crossland, M. Saeed, M. Carossino,

P. Montanaro, A. O'Connell, D. Kenney, H. Gertje, K Grosz, and S Kurnick performed

the experiments; F. Douam, N. Crossland, M. Carossino, P. Montanaro, A. O'Connell,

and D. Kenney performed data analysis; and M. Carossino, N. Crossland, and F.

Douam wrote the manuscript.

ACKNOWLEDGEMENTS

Drs. Crossland and Carossino would like to thank our pathology mentors Drs. Fabio Del

Piero and Ingeborg M. Langohr for helping instill our passion for pathology and for

introducing us to each other. We are hopeful these efforts will represent the early days

of a fruitful and long-lasting collaboration. We acknowledge the Histology and

Immunohistochemistry sections at the Louisiana Animal Disease Diagnostic Laboratory

for their technical assistance. The following reagent was deposited by the Centers for

Disease Control and Prevention and obtained through BEI Resources, NIAID, NIH:

SARS-Related Coronavirus 2, Isolate USA-WA1/2020, NR-52281.

825
REFERENCES

Aid, M., K. Busman-Sahay, S.J. Vidal, Z. Maliga, S. Bondoc, C. Starke, M. Terry, C.A. Jacobson, L. Wrijil, S.
Ducat, O.R. Brook, A.D. Miller, M. Porto, K.L. Pellegrini, M. Pino, T.N. Hoang, A. Chandrashekar,
S. Patel, K. Stephenson, S.E. Bosinger, H. Andersen, M.G. Lewis, J.L. Hecht, P.K. Sorger, A.J.
Martinot, J.D. Estes, and D.H. Barouch. 2020. Vascular Disease and Thrombosis in SARS-CoV-2-
Infected Rhesus Macaques. *Cell* 183:1354-1366 e1313.
Andersen, K.G., A. Rambaut, W.I. Lipkin, E.C. Holmes, and R.F. Garry. 2020. The proximal origin of SARS-
CoV-2. *Nat Med* 26:450-452.
Blair, R.V., M. Vaccari, L.A. Doyle-Meyers, C.J. Roy, K. Russell-Lodrigue, M. Fahlberg, C.J. Monjure, B.
Beddingfield, K.S. Plante, J.A. Plante, S.C. Weaver, X. Qin, C.C. Midkiff, G. Lehmicke, N. Golden, B.
Threeton, T. Penney, C. Allers, M.B. Barnes, M. Pattison, P.K. Datta, N.J. Maness, A. Birnbaum, T.
Fischer, R.P. Bohm, and J. Rappaport. 2020. Acute Respiratory Distress in Aged, SARS-CoV-2-
Infected African Green Monkeys but Not Rhesus Macaques. *Am J Pathol*
Bryche, B., A. St Albin, S. Murri, S. Lacote, C. Pulido, M. Ar Gouilh, S. Lesellier, A. Servat, M. Wasniewski,
E. Picard-Meyer, E. Monchatre-Leroy, R. Volmer, O. Rampin, R. Le Goffic, P. Marianneau, and N.
Meunier. 2020. Massive transient damage of the olfactory epithelium associated with infection
of sustentacular cells by SARS-CoV-2 in golden Syrian hamsters. *Brain Behav Immun* 89:579-586.
Cantuti-Castelvetri, L., R. Ojha, L.D. Pedro, M. Djannatian, J. Franz, S. Kuivanen, F. van der Meer, K. Kallio,
T. Kaya, M. Anastasina, T. Smura, L. Levanov, L. Szirovicza, A. Tobi, H. Kallio-Kokko, P. Osterlund,
M. Joensuu, F.A. Meunier, S.J. Butcher, M.S. Winkler, B. Mollenhauer, A. Helenius, O. Gokce, T.
Teesalu, J. Hepojoki, O. Vapalahti, C. Stadelmann, G. Balistreri, and M. Simons. 2020. Neuropilin-
1 facilitates SARS-CoV-2 cell entry and infectivity. *Science* 370:856-860.
Carossino, M., H.S. Ip, J.A. Richt, K. Shultz, K. Harper, A.T. Loynachan, F. Del Piero, and U.B.R. Balasuriya.
2020. Detection of SARS-CoV-2 by RNAscope((R)) in situ hybridization and
immunohistochemistry techniques. *Arch Virol* 165:2373-2377.
Chen, M., W. Shen, N.R. Rowan, H. Kulaga, A. Hillel, M. Ramanathan, Jr., and A.P. Lane. 2020. Elevated
ACE-2 expression in the olfactory neuroepithelium: implications for anosmia and upper
respiratory SARS-CoV-2 entry and replication. *Eur Respir J* 56:
Conceicao, C., N. Thakur, S. Human, J.T. Kelly, L. Logan, D. Bialy, S. Bhat, P. Stevenson-Leggett, A.K.
Zagrajek, P. Hollinghurst, M. Varga, C. Tsirigoti, M. Tully, C. Chiu, K. Moffat, A.P. Silesian, J.A.
Hammond, H.J. Maier, E. Bickerton, H. Shelton, I. Dietrich, S.C. Graham, and D. Bailey. 2020. The
SARS-CoV-2 Spike protein has a broad tropism for mammalian ACE2 proteins. *PLoS Biol*
18:e3001016.
Corman, V.M., O. Landt, M. Kaiser, R. Molenkamp, A. Meijer, D.K. Chu, T. Bleicker, S. Brunink, J.
Schneider, M.L. Schmidt, D.G. Mulders, B.L. Haagmans, B. van der Veer, S. van den Brink, L.
Wijsman, G. Goderski, J.L. Romette, J. Ellis, M. Zambon, M. Peiris, H. Goossens, C. Reusken, M.P.
Koopmans, and C. Drosten. 2020. Detection of 2019 novel coronavirus (2019-nCoV) by real-time
RT-PCR. *Euro Surveill* 25:
Coronaviridae Study Group of the International Committee on Taxonomy of, V. 2020. The species Severe
acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-
2. *Nat Microbiol* 5:536-544.
Damas, J., G.M. Hughes, K.C. Keough, C.A. Painter, N.S. Persky, M. Corbo, M. Hiller, K.P. Koepfli, A.R.
Pfenning, H. Zhao, D.P. Genereux, R. Swofford, K.S. Pollard, O.A. Ryder, M.T. Nweeia, K.
Lindblad-Toh, E.C. Teeling, E.K. Karlsson, and H.A. Lewin. 2020. Broad host range of SARS-CoV-2
predicted by comparative and structural analysis of ACE2 in vertebrates. *Proc Natl Acad Sci U S A*
117:22311-22322.
Dinnon, K.H., 3rd, S.R. Leist, A. Schafer, C.E. Edwards, D.R. Martinez, S.A. Montgomery, A. West, B.L.
Yount, Jr., Y.J. Hou, L.E. Adams, K.L. Gully, A.J. Brown, E. Huang, M.D. Bryant, I.C. Choong, J.S.
Glenn, L.E. Gralinski, T.P. Sheahan, and R.S. Baric. 2020. A mouse-adapted model of SARS-CoV-2

to test COVID-19 countermeasures. *Nature* In press (https://doi.org/10.1038/s41586-020-2708-
8):
DosSantos, M.F., S. Devalle, V. Aran, D. Capra, N.R. Roque, J.d.M. Coelho-Aguiar, T.C.L.d.S.e. Spohr, J.G.
Subilhaga, C.M. Pereira, I. D'Andrea Meira, P. Niemeyer Soares Filho, and V. Moura-Neto. 2020.
Neuromechanisms of SARS-CoV-2: A Review. *Frontiers in Neuroanatomy* 14:
Eliezer, M., A.L. Hamel, E. Houdart, P. Herman, J. Housset, C. Jourdaine, C. Eloit, B. Verillaud, and C.
Hautefort. 2020. Loss of smell in patients with COVID-19: MRI data reveal a transient edema of
the olfactory clefts. *Neurology* 95:e3145-e3152.
Ellul, M.A., L. Benjamin, B. Singh, S. Lant, B.D. Michael, A. Easton, R. Kneen, S. Defres, J. Sejvar, and T.
Solomon. 2020. Neurological associations of COVID-19. *Lancet Neurol* 19:767-783.
Gaudreault, N., J. Trujillo, M. Carossino, D. Meekins, D. Madden, V. Balaraman, J. Henningson, I.
Morozov, D. Bold, T. Kwong, G. Roman-Sosa, B. Artiaga, S. Indran, K. Cool, A. Garcia-Sastre, W.
Ma, W. Wilson, U. Balasuriya, and J. Richt. 2020. SARS-CoV-2 infection, disease and transmission
in domestic cats. *Emerg Microbes Inf* In press
(https://doi.org/10.1080/22221751.2020.1833687):
Golden, J.W., C.R. Cline, X. Zeng, A.R. Garrison, B.D. Carey, E.M. Mucker, L.E. White, J.D. Shamblin, R.L.
Brocato, J. Liu, A.M. Babka, H.B. Rauch, J.M. Smith, B.S. Hollidge, C. Fitzpatrick, C.V. Badger, and
J.W. Hooper. 2020. Human angiotensin-converting enzyme 2 transgenic mice infected with
SARS-CoV-2 develop severe and fatal respiratory disease. *JCI Insight* 5:
Goyal, P., J.J. Choi, L.C. Pinheiro, E.J. Schenck, R. Chen, A. Jabri, M.J. Satlin, T.R. Campion, Jr., M. Nahid,
J.B. Ringel, K.L. Hoffman, M.N. Alshak, H.A. Li, G.T. Wehmeyer, M. Rajan, E. Reshetnyak, N.
Hupert, E.M. Horn, F.J. Martinez, R.M. Gulick, and M.M. Safford. 2020. Clinical Characteristics of
Covid-19 in New York City. *N Engl J Med* 382:2372-2374.
Hoffmann, M., H. Kleine-Weber, S. Schroeder, N. Kruger, T. Herrler, S. Erichsen, T.S. Schiergens, G.
Herrler, N.H. Wu, A. Nitsche, M.A. Muller, C. Drosten, and S. Pohlmann. 2020. SARS-CoV-2 Cell
Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor.
Cell 181:271-280 e278.
Imai, M., K. Iwatsuki-Horimoto, M. Hatta, S. Loeber, P.J. Halfmann, N. Nakajima, T. Watanabe, M. Ujie, K.
Takahashi, M. Ito, S. Yamada, S. Fan, S. Chiba, M. Kuroda, L. Guan, K. Takada, T. Armbrust, A.
Balogh, Y. Furusawa, M. Okuda, H. Ueki, A. Yasuhara, Y. Sakai-Tagawa, T.J.S. Lopes, M. Kiso, S.
Yamayoshi, N. Kinoshita, N. Ohmagari, S.I. Hattori, M. Takeda, H. Mitsuya, F. Krammer, T. Suzuki,
and Y. Kawaoka. 2020. Syrian hamsters as a small animal model for SARS-CoV-2 infection and
countermeasure development. *Proc Natl Acad Sci U S A* 117:16587-16595.
Jiang, R.D., M.Q. Liu, Y. Chen, C. Shan, Y.W. Zhou, X.R. Shen, Q. Li, L. Zhang, Y. Zhu, H.R. Si, Q. Wang, J.
Min, X. Wang, W. Zhang, B. Li, H.J. Zhang, R.S. Baric, P. Zhou, X.L. Yang, and Z.L. Shi. 2020.
Pathogenesis of SARS-CoV-2 in Transgenic Mice Expressing Human Angiotensin-Converting
Enzyme 2. *Cell* 182:50-58 e58.
Johansen, M.D., A. Irving, X. Montagutelli, M.D. Tate, I. Rudloff, M.F. Nold, N.G. Hansbro, R.Y. Kim, C.
Donovan, G. Liu, A. Faiz, K.R. Short, J.G. Lyons, G.W. McCaughan, M.D. Gorrell, A. Cole, C.
Moreno, D. Couteur, D. Hesselson, J. Triccas, G.G. Neely, J.R. Gamble, S.J. Simpson, B.M.
Saunders, B.G. Oliver, W.J. Britton, P.A. Wark, C.A. Nold-Petry, and P.M. Hansbro. 2020. Animal
and translational models of SARS-CoV-2 infection and COVID-19. *Mucosal Immunol* 13:877-891.
Johns Hopkins University & Medicine. 2020. Coronavirus Resource Center. In.
Lee, M.H., D.P. Perl, G. Nair, W. Li, D. Maric, H. Murray, S.J. Dodd, A.P. Koretsky, J.A. Watts, V. Cheung, E.
Masliah, I. Horkayne-Szakaly, R. Jones, M.N. Stram, J. Moncur, M. Hefti, R.D. Folkerth, and A.
Nath. 2020. Microvascular Injury in the Brains of Patients with Covid-19. *N Engl J Med*

Li, W., M.J. Moore, N. Vasilieva, J. Sui, S.K. Wong, M.A. Berne, M. Somasundaran, J.L. Sullivan, K.
Luzuriaga, T.C. Greenough, H. Choe, and M. Farzan. 2003. Angiotensin-converting enzyme 2 is a
functional receptor for the SARS coronavirus. *Nature* 426:450-454.
Liu, J.M., B.H. Tan, S. Wu, Y. Gui, J.L. Suo, and Y.C. Li. 2020. Evidence of central nervous system infection
and neuroinvasive routes, as well as neurological involvement, in the lethality of SARS-CoV-2
infection. *J Med Virol*
Maiese, A., A.C. Manetti, R. La Russa, M. Di Paolo, E. Turillazzi, P. Frati, and V. Fineschi. 2020. Autopsy
findings in COVID-19-related deaths: a literature review. *Forensic Sci Med Pathol*
Martines, R.B., J.M. Ritter, E. Matkovic, J. Gary, B.C. Bollweg, H. Bullock, C.S. Goldsmith, L. Silva-Flannery,
J.N. Seixas, S. Reagan-Steiner, T. Uyeki, A. Denison, J. Bhatnagar, W.J. Shieh, S.R. Zaki, and C.-
P.W. Group. 2020. Pathology and Pathogenesis of SARS-CoV-2 Associated with Fatal Coronavirus
Disease, United States. *Emerg Infect Dis* 26:2005-2015.
McCray, P.B., Jr., L. Pewe, C. Wohlford-Lenane, M. Hickey, L. Manzel, L. Shi, J. Netland, H.P. Jia, C. Halabi,
C.D. Sigmund, D.K. Meyerholz, P. Kirby, D.C. Look, and S. Perlman. 2007. Lethal infection of K18-
hACE2 mice infected with severe acute respiratory syndrome coronavirus. *J Virol* 81:813-821.
McNamara, T., J.A. Richt, and L. Glickman. 2020. A Critical Needs Assessment for Research in Companion
Animals and Livestock Following the Pandemic of COVID-19 in Humans. *Vector Borne Zoonotic*
Dis
Meekins, D.A., I. Morozov, J.D. Trujillo, N.N. Gaudreault, D. Bold, M. Carossino, B.L. Artiaga, S.V. Indran,
T. Kwon, V. Balaraman, D.W. Madden, H. Feldmann, J. Henningson, W. Ma, U.B.R. Balasuriya,
and J.A. Richt. 2020. Susceptibility of swine cells and domestic pigs to SARS-CoV-2. *Emerg*
Microbes Inf In press (https://doi.org/10.1080/22221751.2020.1831405):
Meinhardt, J., J. Radke, C. Dittmayer, J. Franz, C. Thomas, R. Mothes, M. Laue, J. Schneider, S. Brunink, S.
Greuel, M. Lehmann, O. Hassan, T. Aschman, E. Schumann, R.L. Chua, C. Conrad, R. Eils, W.
Stenzel, M. Windgassen, L. Rossler, H.H. Goebel, H.R. Gelderblom, H. Martin, A. Nitsche, W.J.
Schulz-Schaeffer, S. Hakroush, M.S. Winkler, B. Tampe, F. Scheibe, P. Kortvelyessy, D. Reinhold,
B. Siegmund, A.A. Kuhl, S. Elezkurtaj, D. Horst, L. Oesterhelweg, M. Tsokos, B. Ingold-Heppner, C.
Stadelmann, C. Drosten, V.M. Corman, H. Radbruch, and F.L. Heppner. 2020. Olfactory
transmucosal SARS-CoV-2 invasion as a port of central nervous system entry in individuals with
COVID-19. *Nat Neurosci*
Moreau, G.B., S.L. Burgess, J.M. Sturek, A.N. Donlan, W.A. Petri, and B.J. Mann. 2020. Evaluation of K18-
hACE2 Mice as a Model of SARS-CoV-2 Infection. *Am J Trop Med Hyg* 103:1215-1219.
Munoz-Fontela, C., W.E. Dowling, S.G.P. Funnell, P.S. Gsell, A.X. Riveros-Balta, R.A. Albrecht, H.
Andersen, R.S. Baric, M.W. Carroll, M. Cavaleri, C. Qin, I. Crozier, K. Dallmeier, L. de Waal, E. de
Wit, L. Delang, E. Dohm, W.P. Duprex, D. Falzarano, C.L. Finch, M.B. Frieman, B.S. Graham, L.E.
Gralinski, K. Guilfoyle, B.L. Haagmans, G.A. Hamilton, A.L. Hartman, S. Herfst, S.J.F. Kaptein, W.B.
Klimstra, I. Knezevic, P.R. Krause, J.H. Kuhn, R. Le Grand, M.G. Lewis, W.C. Liu, P. Maisonnasse,
A.K. McElroy, V. Munster, N. Oreshkova, A.L. Rasmussen, J. Rocha-Pereira, B. Rockx, E.
Rodriguez, T.F. Rogers, F.J. Salguero, M. Schotsaert, K.J. Stittelaar, H.J. Thibaut, C.T. Tseng, J.
Vergara-Alert, M. Beer, T. Brasel, J.F.W. Chan, A. Garcia-Sastre, J. Neyts, S. Perlman, D.S. Reed,
J.A. Richt, C.J. Roy, J. Segales, S.S. Vasan, A.M. Henao-Restrepo, and D.H. Barouch. 2020. Animal
models for COVID-19. *Nature* 586:509-515.
Oladunni, F.S., J.-G. Park, P.P. Tamayo, O. Gonzalez, A. Akhter, A. Allué-Guardia, A. Olmo-Fontánez, S.
Gautam, A. Garcia-Vilanova, C. Ye, K. Chiem, C. Headley, V. Dwivedi, L.M. Parodi, K.J. Alfson,
H.M. Staples, A. Schami, J.I. Garcia, A. Whigham, R.N. Platt, M. Gazi, J. Martinez, C. Chuba, S.
Earley, O.H. Rodriguez, S.D. Mdaki, K.N. Kavelish, R. Escalona, C.R.A. Hallam, C. Christie, J.L.
Patterson, T.J.C. Anderson, R. Carrion, E.J. Dick, S. Hall-Ursone, L.S. Schlesinger, D. Kaushal, L.D.
Giavedoni, X. Alvarez, J. Turner, L. Martinez-Sobrido, and J.B. Torrelles. 2020. Lethality of SARS-

CoV-2 infection in K18 human angiotensin converting enzyme 2 transgenic mice. *bioRxiv*
preprint (https://doi.org/10.1101/2020.07.18.210179):
Osterrieder, N., L.D. Bertzbach, K. Dietert, A. Abdelgawad, D. Vladimirova, D. Kunec, D. Hoffmann, M.
Beer, A.D. Gruber, and J. Trimpert. 2020. Age-Dependent Progression of SARS-CoV-2 Infection in
Syrian Hamsters. *Viruses* 12:
Rathnasinghe, R., S. Strohmeier, F. Amanat, V.L. Gillespie, F. Krammer, A. Garcia-Sastre, L. Coughlan, M.
Schotsaert, and M.B. Uccellini. 2020. Comparison of transgenic and adenovirus hACE2 mouse
models for SARS-CoV-2 infection. *Emerg Microbes Infect* 9:2433-2445.
Rockx, B., T. Kuiken, S. Herfst, T. Bestebroer, M.M. Lamers, B.B. Oude Munnink, D. de Meulder, G. van
Amerongen, J. van den Brand, N.M.A. Okba, D. Schipper, P. van Run, L. Leijten, R. Sikkema, E.
Verschoor, B. Verstrepen, W. Bogers, J. Langermans, C. Drosten, M. Fentener van Vlissingen, R.
Fouchier, R. de Swart, M. Koopmans, and B.L. Haagmans. 2020. Comparative pathogenesis of
COVID-19, MERS, and SARS in a nonhuman primate model. *Science* 368:1012-1015.
Rosenke, K., K. Meade-White, M. Letko, C. Clancy, F. Hansen, Y. Liu, A. Okumura, T.L. Tang-Huau, R. Li, G.
Saturday, F. Feldmann, D. Scott, Z. Wang, V. Munster, M.A. Jarvis, and H. Feldmann. 2020.
Defining the Syrian hamster as a highly susceptible preclinical model for SARS-CoV-2 infection.
bioRxiv
Shi, J., Z. Wen, G. Zhong, H. Yang, C. Wang, B. Huang, R. Liu, X. He, L. Shuai, Z. Sun, Y. Zhao, P. Liu, L.
Liang, P. Cui, J. Wang, X. Zhang, Y. Guan, W. Tan, G. Wu, H. Chen, and Z. Bu. 2020. Susceptibility
of ferrets, cats, dogs, and other domesticated animals to SARS-coronavirus 2. *Science* 368:1016-
1020.
Shuai, L., G. Zhong, Q. Yuan, Z. Wen, C. Wang, X. He, R. Liu, J. Wang, Q. Zhao, Y. Liu, N. Huo, J. Deng, J.
Bai, H. Wu, Y. Guan, J. Shi, K. Tian, N. Xia, H. Chen, and Z. Bu. 2020. Replication, pathogenicity,
and transmission of SARS-CoV-2 in minks. *National Science Review*
Sia, S.F., L.M. Yan, A.W.H. Chin, K. Fung, K.T. Choy, A.Y.L. Wong, P. Kaewpreedee, R. Perera, L.L.M. Poon,
J.M. Nicholls, M. Peiris, and H.L. Yen. 2020. Pathogenesis and transmission of SARS-CoV-2 in
golden hamsters. *Nature* 583:834-838.
Simonnet, A., M. Chetboun, J. Poissy, V. Raverdy, J. Noulette, A. Duhamel, J. Labreuche, D. Mathieu, F.
Pattou, M. Jourdain, Licorn, C. the Lille, and g. Obesity study. 2020. High Prevalence of Obesity in
Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2) Requiring Invasive Mechanical
Ventilation. *Obesity (Silver Spring)* 28:1195-1199.
Solomon, I.H., E. Normandin, S. Bhattacharyya, S.S. Mukerji, K. Keller, A.S. Ali, G. Adams, J.L. Hornick, R.F.
Padera, and P. Sabeti. 2020. Neuropathological Features of Covid-19. *New England Journal of*
Medicine
Song, E., C. Zhang, B. Israelow, A. Lu-Culligan, A.V. Prado, S. Skriabine, P. Lu, O.E. Weizman, F. Liu, Y. Dai,
K. Szigeti-Buck, Y. Yasumoto, G. Wang, C. Castaldi, J. Heltke, E. Ng, J. Wheeler, M.M. Alfajaro, E.
Levavasseur, B. Fontes, N.G. Ravindra, D. Van Dijk, S. Mane, M. Gunel, A. Ring, S.A. Jaffar Kazmi,
K. Zhang, C.B. Wilen, T.L. Horvath, I. Plu, S. Haik, J.L. Thomas, A. Louvi, S.F. Farhadian, A. Huttner,
D. Seilhean, N. Renier, K. Bilguvar, and A. Iwasaki. 2020. Neuroinvasion of SARS-CoV-2 in human
and mouse brain. *bioRxiv*
Sun, S.H., Q. Chen, H.J. Gu, G. Yang, Y.X. Wang, X.Y. Huang, S.S. Liu, N.N. Zhang, X.F. Li, R. Xiong, Y. Guo,
Y.Q. Deng, W.J. Huang, Q. Liu, Q.M. Liu, Y.L. Shen, Y. Zhou, X. Yang, T.Y. Zhao, C.F. Fan, Y.S. Zhou,
C.F. Qin, and Y.C. Wang. 2020. A Mouse Model of SARS-CoV-2 Infection and Pathogenesis. *Cell*
Host Microbe 28:124-133 e124.
Tartof, S.Y., L. Qian, V. Hong, R. Wei, R.F. Nadjafi, H. Fischer, Z. Li, S.F. Shaw, S.L. Caparosa, C.L. Nau, T.
Saxena, G.K. Rieg, B.K. Ackerson, A.L. Sharp, J. Skarbinski, T.K. Naik, and S.B. Murali. 2020.
Obesity and Mortality Among Patients Diagnosed With COVID-19: Results From an Integrated
Health Care Organization. *Ann Intern Med* In press (https://doi.org/10.7326/M20-3742):

Team, C.C.-R. 2020. Preliminary Estimates of the Prevalence of Selected Underlying Health Conditions
Among Patients with Coronavirus Disease 2019 - United States, February 12-March 28, 2020.
MMWR Morb Mortal Wkly Rep 69:382-386.
Tenforde, M.W., S.S. Kim, C.J. Lindsell, E. Billig Rose, N.I. Shapiro, D.C. Files, K.W. Gibbs, H.L. Erickson, J.S.
Steingrub, H.A. Smithline, M.N. Gong, M.S. Aboodi, M.C. Exline, D.J. Henning, J.G. Wilson, A.
Khan, N. Qadir, S.M. Brown, I.D. Peltan, T.W. Rice, D.N. Hager, A.A. Ginde, W.B. Stubblefield,
M.M. Patel, W.H. Self, L.R. Feldstein, I.V.Y.N. Investigators, C.C.-R. Team, and I.V.Y.N.
Investigators. 2020. Symptom Duration and Risk Factors for Delayed Return to Usual Health
Among Outpatients with COVID-19 in a Multistate Health Care Systems Network - United States,
March-June 2020. *MMWR Morb Mortal Wkly Rep* 69:993-998.
Walker, A., G. Pottinger, A. Scott, and C. Hopkins. 2020. Anosmia and loss of smell in the era of covid-19.
BMJ 370:m2808.
Wang, C., P.W. Horby, F.G. Hayden, and G.F. Gao. 2020a. A novel coronavirus outbreak of global health
concern. *Lancet* 395:470-473.
Wang, F., R.M. Kream, and G.B. Stefano. 2020b. Long-Term Respiratory and Neurological Sequelae of
COVID-19. *Med Sci Monit* 26:e928996.
Winkler, E.S., A.L. Bailey, N.M. Kafai, S. Nair, B.T. McCune, J. Yu, J.M. Fox, R.E. Chen, J.T. Earnest, S.P.
Keeler, J.H. Ritter, L.I. Kang, S. Dort, A. Robichaud, R. Head, M.J. Holtzman, and M.S. Diamond.
2020. SARS-CoV-2 infection in the lungs of human ACE2 transgenic mice causes severe
inflammation, immune cell infiltration, and compromised respiratory function. *Nature* In press
(https://doi.org/10.1038/s41590-020-0778-2):
Yinda, C.K., J.R. Port, T. Bushmaker, I.O. Owusu, V.A. Avanzato, R.J. Fischer, J.E. Schulz, M.G. Holbrook,
M.J. Hebner, R. Rosenke, T. Thomas, A. Marzi, S.M. Best, E. de Wit, C. Shaia, N. van Doremalen,
and V.J. Munster. 2020. K18-hACE2 mice develop respiratory disease resembling severe COVID-
19. *bioRxiv* preprint (10.1101/2020.08.11.246314):
Zheng, J., L.R. Wong, K. Li, A.K. Verma, M.E. Ortiz, C. Wohlford-Lenane, M.R. Leidinger, C.M. Knudson,
D.K. Meyerholz, P.B. McCray, Jr., and S. Perlman. 2020. COVID-19 treatments and pathogenesis
including anosmia in K18-hACE2 mice. *Nature*

1045

FIGURE LEGENDS

FIGURE LEGENDS

Figure 1. SARS-CoV-2 caused lethal disease in K18-hACE2 mice (n=33) inoculated

intranasally with 1×10^6 plaque forming units (PFU). Body weight (A), clinical signs (B),

temperature (C), and mortality (D) were monitored daily. Viral loads (genome copy

numbers/mg or ml) were monitored in the brain, lungs (E) and serum (F) throughout the

study. Mean genome copy numbers are depicted. The limit of detection (LOD) is shown

with a dashed line.

Figure 2. Temporal analysis of SARS-CoV-2 infection in the nasal cavity of K18-hACE2 mice at 2 (A-D) and 4 (E-H) days following intranasal inoculation. Histological changes and viral antigen (brown) and RNA (red) distribution and abundance were assessed. At 2 dpi, suppurative rhinitis in the rostral and intermediate turbinates (A, arrow) correlated with abundant intraepithelial SARS-CoV-2 antigen (C) and RNA (C, inset). Abundant viral antigen and RNA was detected in the olfactory neuroepithelium (ONE, D and inset) in the absence of histologic lesions (B). At 4 dpi, only sporadically infected cells were noted in the epithelium lining the turbinates and ONE (G and H, arrow and insets) in the absence of histologic lesions (E and F). H&E and Fast Red (viral RNA), 200X total magnification. Bar = 100 µm.

Figure 3. Temporal analysis of SARS-CoV-2 infection in the lungs of K18-hACE2 mice at 2 (D-F), 4 (G-I), 7 (J-L) and 14 (M-O) days following intranasal inoculation. Histological changes and viral RNA (red) distribution and abundance were assessed. Mild to moderate interstitial pneumonia was evident starting at 2 days post-infection (dpi) with frequently reactive blood vessels (E, arrow) and a peak in viral RNA at 4 days post-infection (I). At 7 and 14 days post-infection, there was evidence of alveolar type 2 (AT2) cell hyperplasia (K and N, arrows), with viral RNA restricted to areas spared of inflammation (L). The pneumonia was significantly ameliorated by 14 dpi (M) with no viral RNA (O). H&E and Fast Red (viral RNA), 50X (A, D, G, J, and M; bar = 500 µm) and 200X (B, C, E, F, H, I, K, L, M and N; bar = 100 µm) total magnification.

Figure 4. SARS-CoV-2 tropism and temporal immunoprofiling of the pulmonary host inflammatory response following SARS-CoV-2 intranasal inoculation in K18-hACE2. (A) SARS-CoV-2 (yellow) showed tropism for RAGE$^+$ alveolar type 1 (AT1, magenta) and

scattered SPC$^+$ alveolar type 2 (AT2, red) pneumocytes (A1 and A2, arrowheads and arrows, respectively) but not for CD31$^+$ endothelial cells (green). SARS-CoV-2 (orange) was abundant and localized within AT1 and AT2 pneumocytes, with a progressive increase in the number of CD3+CD8- T cells (blue, presumed to be CD4+ T cells), Iba-1+ histiocytes (magenta), and CD8+ T cells (red) throughout the course of the experiment (2 days post-infection [dpi, C], 7 dpi [D] and 14 dpi [E]). SARS-CoV-2 S antigen was no longer detectable by 14 dpi, and numerous aggregates of CD19+ B cells (green) were noted (E). Multiplex fluorescent IHC, 200X total magnification. Bar = 50 µm

Figure 5. Temporal neuronal damage in K18-hACE2 mice following intranasal inoculation with SARS-CoV-2. No histologic changes were noted in the cerebrum or olfactory bulb until 6-7 days post-infection (dpi, A-D). SARS-CoV-2 antigen (brown) was evident as early as 4 dpi (F, arrow). At 6-7 dpi, mild (G, arrowheads) to marked (G, inset) spongiosis with neuronal degeneration and necrosis involving multiple areas within the cerebral cortex and elsewhere were noted. Similar changes were evident in the olfactory bulb, with occasional perivascular cuffs/gliosis (H, arrowhead) and abundant viral antigen (I, arrows). No histologic alterations or viral antigen was detected in survivor mice at 14 dpi (J-L). H&E and DAB (viral antigen), 100X (A, B, D, E, G, H, J, K; bar = 200 µm) and 200X (C, F, I, L; bar = 100 µm) total magnification.

Figure 6. Invasion of SARS-CoV-2 into the central nervous system. Sagittal sections of the head were analyzed for viral antigen and RNA distribution. SARS-CoV-2 infected neurons within the mitral layer of the olfactory bulb (1, arrow) as well as small clusters of neuronal bodies within the cerebral cortex (2, SARS-CoV-2 RNA in inset) as early as

4 days post-infection. At 7 days post-infection, SARS-CoV-2 antigen was widespread

along the mitral layer of the olfactory bulb (1) and throughout the central nervous

system (2, SARS-CoV-2 RNA in inset) with exception of the cerebellum. EPL, external

plexiform layer; GCL, granular cell layer; GL, glomerular layer; ML, mitral layer. DAB

(viral antigen) and Fast Red (viral RNA). 7.5X (bar = 2.5 mm) and 200X (bar = 100 μm)

total magnification.

Figure 7. Analysis of SARS-CoV-2 tropism within the central nervous system at 7 days

post-infection using multiplex immunohistochemistry. Abundant SARS-CoV-2 antigen

(green) localized within cortical (A), hippocampal (B) and brainstem (C) neurons with

abundant neighboring Iba-1+ microglia (orange) and clusters of GFAP+ astrocytes

(gliosis, magenta). 40X total magnification. Bar = 50 μm.

Figure 8. Distribution of ACE2 in lungs, nasal cavity, brain and olfactory bulb of wild-

type C57BL/6J and transgenic (mock and SARS-CoV-2-infected) K18-hACE2 mice was

analyzed via immunohistochemistry using a cross-reactive anti-ACE2 antibody. In the

lungs (A-C), ACE2 expression (brown) was mostly restricted to the apical membrane of

bronchiolar epithelial cells with scattered positive pneumocytes (inset arrows). Nasal

(rostral/intermediate turbinates [R/I]) and olfactory epithelium (ONE) were devoid of

ACE2 in C57BL/6J mice (D) but expression was enhanced in K18-hACE2 mice with

intense apical expression (E and F). ACE2 expression within the brain (G-I) and

olfactory bulb (J-L) was restricted to capillary endothelium with no neuronal expression.

DAB, 200X total magnification. Bar = 100 μm.

Figure 9. Expression and distribution of *hACE2* mRNA in the lungs of C57BL/6J and

K18-hACE2 transgenic mice via RNAscope® ISH. While no expression of *hACE2* was

noted in the lungs of wild-type C57BL/6J mice (A), *hACE2* was expressed in the

bronchiolar epithelium (arrowheads) and sporadic pneumocytes (arrows) in transgenic

K18-hACE2 mice (B and C), which correlated with immunohistochemical findings. Fast

Red, 400X total magnification. Bar = 50 µm.

Figure 10. Expression and distribution of *hACE2* mRNA in the brain of K18-hACE2

transgenic mice via RNAscope® ISH. *hACE2* was expressed in clusters of neurons

within the cerebral cortex (A), hippocampus (B) as well as in other locations including

Purkinje cells of the cerebellum. Fast Red, 400X total magnification. Bar = 50 µm.

SUPPLEMENTARY FIGURES

Figure S1. Histologic and viral antigen distribution in the nasal cavity of K18-hACE2

mice following SARS-CoV-2 infection. No histologic lesions or antigen were detected at

7 and 14 days post-infection (dpi). 200X total magnification. Bar = 100 µm.

Figure S2. Localization and abundance of SARS-CoV-2 RNA (red) and antigen (brown)

in the lungs of infected K18-hACE2 mice as demonstrated by anti-Spike RNAscope® *in

situ* hybridization and immunohistochemistry. 2 (A and B), 4 (C and D), 7 (E and F) and

14 (G and H) days post-infection (dpi). 50X (bar = 500 µm) and 100X (bar = 100 µm)

total magnification.

Figure S3. Histological and immunohistochemical findings in the cervicothoracic spinal

cord of SARS-CoV-2-infected K18-hACE2 mice at 7 days post-infection. Histologically

(A), Multifocal neuronal bodies within the grey matter are shrunken, angular and

hyperchromatic (neuronal degeneration and necrosis), and the neuroparenchyma has

multiple clear spaces filled with small amounts of debris (spongiosis) with a slight

increase in the number of glial cells (gliosis). H&E, 100X total magnification. (B)

Abundant SARS-CoV-2 spike protein localized within the perikaryon and processes of

motor neurons within the spinal cord. DAB, 200X total magnification. Bar = 100 μm.

Figure S4. Expression of *hACE2* mRNA (red) in the brain of C57BL/6J and K18-hACE2

mice determined by RNAscope® ISH. While wild-type mice exhibit no expression (A),

hACE2 is expressed in clusters of neurons at multiple locations in K18-hACE2 mice (B).

Fast Red, 400X total magnification. Bar = 50 μm.

Figure S5. Expression of ACE2 (red) in the enterocytes lining the small intestine of

C57BL/6J and K18-hACE2 mice. Immunohistochemistry was performed using a cross-

reactive anti-ACE2 antibody. In the stomach, expression was intense in the non-

glandular mucosa and capillaries of the glandular mucosa (A-C, arrows). In the colon,

scattered enterocytes express ACE2 (G-I, arrows). Fast Red, 100X (A-C; bar = 200 μm)

and 200X (D-I; bar = 100 μm) total magnification.

1158

1159

1160

Table 1. SARS-CoV-2 viral antigen abundance in tissues derived from SARS-CoV-2-infected K18-hACE2 mice. Median scores are represented along with ranges between brackets when applicable.

DPI	AT1/AT2	Bronchioles	Rostral turbinates	Intermediate turbinates	ONE	Olf. bulb	Brain	Spinal cord (CT)	Spinal cord (LS)	GI*	Kidneys
Mock	0	0	0	0	0	0	0	0	0	0	0
2	2 (1-2)	0	1 (0-2)	2 (1-2)	1 (1-2)	0	0	0	0	0	0
4	2 (1-3)	0	0 (0-1)	0 (0-1)	1 (0-1)	0 (0-1)	0 (0-1)	0	0	0	0
6-8	2 (1-3)	0	0	0	1 (0-1)	1 (0-2)	3 (0-3)	1 (0-2)	0 (0-1)	0	0
14	0	0	0	0	0	0	0	0	0	0	0

0, no SARS-CoV-2 antigen observed; 1, 0 to 5% of cells within a high magnification (400X) field are positive for viral antigen; 2, 5 to 25% of cells within a high magnification (400X) field are positive for viral antigen; 3, >25 to <50% of cells within a high magnification (400X) field are positive for viral antigen. NA, not available. AT1, alveolar type 1 pneumocytes; AT2, alveolar type 2 pneumocytes; ONE, olfactory neuroepithelium; CT, cervicothoracic segment; LS, lumbosacral segment; GI, gastrointestinal tract.

*Sections examined included stomach, small intestine (duodenum, jejunum and ileum) and large intestine (cecum and colon).

1170   **Table 2.** Clinical scoring system used for clinical monitoring of SARS-CoV-2-infected K18-hACE2 mice.

Category	Score = Criteria
Body weight	1 = 10-19% loss
Respiration	1 = rapid, shallow, increased effort
Appearance	1 = ruffled fur, hunched posture
Responsiveness	1 = low to moderate unresponsiveness
Neurologic signs	1 = tremors

1171

1172

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

C57BL/6J - Mock   K18-hACE2 - Mock   K18-hACE2 – 7 dpi

Lung

Nasal cavity — R/I ONE   R/I ONE   R/I ONE

Cerebrum

Olf. bulb

Figure 9

Figure 10

Cerebral cortex

A

Hippocampus

B