Inhibitors of ERp44, PDIA1, and AGR2 induce disulfide-mediated

oligomerization of Death Receptors 4 and 5 and cancer cell death

[1]Mary E. Law, [2]Elham Yaaghubi, [2]Amanda F. Ghilardi, [1]Bradley J. Davis, [2]Renan B. Ferreira,

[3]Jin Koh, [3,4]Sixue Chen, [2]Sadie F. DePeter, [2]Christopher M. Schilson, [5]Chi-Wu Chiang, [6,7]Coy

D. Heldermon, [8]Peter Nørgaard, [2,7]Ronald K. Castellano[#], and [1,7]Brian K. Law[#]

Affiliations

[1]Department of Pharmacology & Therapeutics, University of Florida, Gainesville, FL 32610

USA.

[2]Department of Chemistry, University of Florida, Gainesville, FL 32611 USA.

[3]Proteomics and Mass Spectrometry Facility, Interdisciplinary Center for Biotechnology

Research, University of Florida, Gainesville, FL 32610 USA.

[4]Department of Biology, Genetics Institute, University of Florida, Gainesville, FL 32610 USA

[5]Institute of Molecular Medicine, College of Medicine and Center

for Infectious Disease and Signaling Research, National Cheng

Kung University, Tainan, Taiwan.

[6]Department of Medicine, University of Florida, Gainesville, FL 32610 USA.

[7]UF-Health Cancer Center, University of Florida, Gainesville, FL 32610 USA.

[8]Department of Pathology, Copenhagen University Hospital Herlev, DK-2730 Herlev,

Denmark.

[#]Co-Contributing Corresponding Authors

Addresses of Corresponding Authors

Brian K. Law

1200 Newell Drive

Room R5-210 Academic Research Building

University of Florida

Gainesville, FL 32610

Email: bklaw@ufl.edu Phone: (352) 273-9423

Ronald K. Castellano

Department of Chemistry

University of Florida

P.O. Box 117200

Gainesville, FL 32611

Email: castellano@chem.ufl.edu Phone: (352) 392-2752

Conflict of Interest

The Authors have no conflicts of interest to declare.

Running Title: Molecular Targets of DDAs

Abstract

Breast cancer mortality remains unacceptably high, indicating a need for safer and more

effective therapeutic agents. Disulfide bond Disrupting Agents (DDAs) were previously

identified as a novel class of anticancer compounds that selectively kill cancers that

overexpress the Epidermal Growth Factor Receptor (EGFR) or its family member HER2. DDAs

kill EGFR+ and HER2+ cancer cells via the parallel downregulation of EGFR, HER2, and

HER3 and activation/oligomerization of Death Receptors 4 and 5 (DR4/5). However, the

mechanisms by which DDAs mediate these effects are unknown. Affinity purification analyses

employing biotinylated-DDAs reveal that the Protein Disulfide Isomerase (PDI) family members

AGR2, PDIA1, and ERp44 are DDA target proteins. Further analyses demonstrate that

shRNA-mediated knockdown of AGR2 and ERp44, or expression of ERp44 mutants, enhance

basal and DDA-induced DR5 oligomerization. DDA treatment of breast cancer cells disrupts

PDIA1 and ERp44 mixed disulfide bonds with their client proteins. Together, the results herein

reveal DDAs as the first small molecule, active site inhibitors of AGR2 and ERp44, and

demonstrate roles for AGR2 and ERp44 in regulating the activity, stability, and localization of

DR4 and DR5, and activation of Caspase 8.

Key Words: Apoptosis, Endoplasmic Reticulum Stress, Cancer, Disulfide bond Disrupting

Agents, DR5, DR4, EGFR, AGR2, ERp44, PDIA1, ERAP1

Introduction

In spite of significant improvements in early detection, and new molecularly targeted therapeutics, breast cancer remains the second largest cancer killer of women in America after lung cancer, and is the most frequently diagnosed cancer in American women [1]. Much of the lethality of breast cancer derives from our inability to successfully treat patients with metastatic and/or drug-resistant malignancies. New drugs are needed that bypass resistance to currently used regimens by targeting novel cancer cell-specific vulnerabilities. A class of agents termed Disulfide bond Disrupting Agents (DDAs) was previously identified and shown to kill breast cancer cells in association with downregulation of the HER-family proteins EGFR, HER2, and HER3, and decreased activating phosphorylation of the Akt kinase [2]. Subsequent work showed that DDAs induce endoplasmic reticulum stress (ERS) [3] and alter the pattern of disulfide bonding of EGFR and the TRAIL receptors, Death Receptor 4 (DR4) and Death Receptor 5 (DR5) [4]. DDA-induced cell death is mediated by activation of the Caspase 8-Caspase 3 extrinsic apoptotic pathway.

Little is known regarding how the disulfide bonding of HER-family and Death Receptor-family proteins are chaperoned within the endoplasmic reticulum, and how the disulfide bonding patterns of these proteins control their stability, intracellular localization, and downstream signaling. TNF-Related Apoptosis Inducing Ligand (TRAIL) has been considered a promising cancer therapeutic agent due to its selectivity for killing cancer cells, while leaving normal tissues unharmed [5-8]. However, TRAIL analogs and DR4/5 agonistic antibodies have yet to be clinically approved for anticancer therapy. This is due in part to pharmacokinetic issues related to these protein drugs themselves, and in part to the ability of cancer cells to acquire resistance through a variety of mechanisms, including downregulation of DR4 and

DR5 (reviewed in [9]). Since DDAs upregulate DR5 and activate DR4 and DR5 in a ligand-

independent manner [4, 10], DDAs may overcome the resistance mechanisms that suppress

the efficacy of other activators of the TRAIL/Death Receptor pathway. A recent report showed

that the extracellular domain of DR5 prevents oligomerization and downstream activation of

Caspase 8 in the absence of ligand engagement, and that excision of the DR5 extracellular

domain permits full activation of DR5 in a ligand-independent manner [11]. It was previously

proposed that DDAs activate DR5 in a ligand-independent manner by permitting inter-

molecular disulfide bond formation at the expense of intramolecular disulfide bonding, resulting

in DR5 oligomerization and activation [4, 10]. However, the precise mechanisms by which this

happens are unknown. Similarly, DDA treatment causes disulfide bond-dependent EGFR

oligomerization followed by degradation [2, 4], but how DDAs perturb EGFR disulfide bonding

is unknown. The Protein Disulfide Isomerase (PDI) family member AGR2 is essential for EGFR

folding and surface localization [12], and likewise facilitates expression of Mucins [13, 14].

The PDI family contains 21 members (reviewed in [15]). Canonical PDIs are considered

promising targets for anticancer therapeutics [16] and significant attention has focused on the

roles of canonical PDIs in mediating disulfide bond oxidation, reduction, or exchange via their

active site thioredoxin-like CXXC motifs. However, other PDI family members are less well

studied. The PDIs AGR2 and AGR3 contain a non-canonical CXXS active site motif. Based on

biochemical PDI mutagenesis studies [17] it is predicted that AGR2/3 will exhibit disulfide

isomerase activity, but not oxidase or reductase activity. Consistent with this, AGR2 forms

stable mixed disulfide bonds with its client proteins [12, 13]. Another PDI family member,

ERp44, also contains an active site CXXS motif, and has multiple functions within the

endoplasmic reticulum (ER). ERp44 plays a critical role in the disulfide-mediated

oligomerization of Adiponectin [18, 19] and Immunoglobulin M [20, 21]. Additionally, ERp44

anchors client/partner proteins such as ERO1α, Prx4, FGE/SUMF1, and ER Aminopeptidase 1

(ERAP1) [22-25] that do not possess C-terminal -KDEL ER retention sequences to prevent

their secretion (reviewed in [26]).

HER-family proteins are the only sub-family of receptor tyrosine kinases that contain

two cysteine-rich repeats in their extracellular domain [27]. DR4 and DR5 also harbor

numerous disulfide bonds in their extracellular domains [28]. Based on these observations, we

previously hypothesized that HER-family proteins and DR4/5 may be particularly sensitive to

agents that impede native disulfide bond formation. The mechanisms by which DDAs cause

downregulation of EGFR, HER2, and HER3, and activate DR4/5 are unknown. Here we show

using novel biotinylated DDA derivatives that DDAs covalently modify ERp44, PDIA1 and

AGR2 both in vitro and in intact cells, and that DDA treatment of cells disrupts PDIA1 and

ERp44 disulfide bonds with their client proteins, and disulfide-mediated AGR2 dimerization.

Together, the results indicate that DDAs activate DR4/5 by inhibiting the activity of the PDI

family members ERp44, PDIA1, and AGR2. DDAs are to our knowledge the first active site

inhibitors of ERp44 and AGR2.

Results

Characterization of novel DDAs with enhanced potency: Previously described DDAs, including

RBF3, DTDO, and tcyDTDO (Fig. 1A), were demonstrated to suppress tumor growth through

induction of apoptotic cancer cell death [4, 29]. As shown in our previous studies, modifications

of the parent cyclic DDA, DTDO, resulted in a more potent DDA, tcyDTDO, with improved

pharmacodynamic and pharmacokinetic properties [4, 10, 29]. With tcyDTDO in hand, we then

explored the effect of substitutions on its cyclohexane ring to determine if its potency and drug-

like properties could be further improved. Our efforts led to the design and synthesis of novel

DDAs including dimethoxy-tcyDTDO (dMtcyDTDO) and difluoro-tcyDTDO (dFtcyDTDO) (Fig.

1B). Well appreciated, fluorine's size and high electronegativity lead to strong C–F bonds that

resist metabolic degradation making fluorine substitution an effective way to increase drug

stability. For instance, replacing hydrogen with fluorine significantly alters the rate and extent of

Cytochrome P450 metabolism [30-33]. Both dMtcyDTDO and dFtcyDTDO exhibit

approximately two-fold greater potency than tcyDTDO in cell viability assays (Fig. 1C).

DMtcyDTDO also exhibited two-fold greater potency than tcyDTDO in DNA and protein

synthesis assays (Fig. S1A). DDAs induce apoptotic cell death associated with ERS, disulfide

bond-mediated oligomerization of Death Receptors 4 and 5 (DR4/5) and EGFR, and

upregulation of DR5 [2-4, 29]. Therefore, we examined the effect of the novel DDAs on

disulfide bonding of these proteins by immunoblot analysis under non-reducing conditions ([4]

and Supplemental Information). When applied to cancer cells at the same concentration,

dMtcyDTDO and dFtcyDTDO induced more DR4/5 oligomerization, higher DR5 levels, and

greater ERS (Fig. 1D, left panel). Induction of ERS by dFtcyDTDO and dMtcyDTDO, as

indicated by GRP78 elevation, and disulfide-mediated oligomerization of DR4 and EGFR, were

detected at 625 nM and oligomerization of DR5 was observed at 1.25 µM of either dFtcyDTDO

or dMtcyDTDO (Fig. 1D, right panel). Additional immunoblot analyses performed under

reducing conditions revealed that dMtcyDTDO is more potent than tcyDTDO in the induction of

HER3 downregulation, DR5 upregulation, and ERS as indicated by elevated CHOP and

XBP1s (Fig. S1B). Time-course and concentration-response analyses of these endpoints

showed that dMtcyDTDO induced these responses as early as four hours after treatment (Fig.

S1C).

Quantitation of dMtcyDTDO effects on the levels and state of oligomerization of DR4

and DR5 revealed that dMtcyDTDO increases DR4 oligomerization without affecting overall

DR4 levels, while dMtcyDTDO increases the levels of both oligomeric and monomeric forms of

DR5 (Fig. 1E). Tumor studies using the HER2+ BT474 xenograft model [2] were performed to

examine if dMtcyDTDO and dFtcyDTDO induce rapid breast cancer cell death in mice bearing

palpable breast tumors. Tumors from animals treated with the vehicle for four consecutive

days did not show evidence of tumor necrosis (Fig. 1F). In contrast, tumors regressed in

mammary fat pads from dFtcyDTDO treated animals. Mammary glands from dMtcyDTDO

treated animals exhibited tumors that had undergone coagulation necrosis, consistent with

rapid and uniform death of the cancer cells.

AGR2 as a DDA binding protein: To understand how DDAs kill breast tumors, identification of

their direct protein targets is essential. The development of molecular probes proved crucial in

identifying DDA biological targets and linking them to the observed DDA responses. For this

purpose, we used biotinylated-DDA based affinity pulldown techniques.

To generate DDA probes, we sought a biologically active cyclic DDA with a functional

group that enables installation of a biotin tag. This was achieved by fusing a Boc-protected

pyrrolidine ring at the 4,5-positions of the parent DTDO compound to generate ($\pm$)-

BocPyrDTDO (Fig. 2A), which is biologically active and showed a potency similar to DTDO

(Fig. S1D). The synthesis of ($\pm$)-BocPyrDTDO was accomplished by cleaving the acetyl groups

of dithioacetate ($\pm$)-**3** followed by oxidation to furnish the 1,2-dithiane 1,1-dioxide ring (Fig. 2A).

Intermediate ($\pm$)-**3** can be obtained through a few functional group interconversion steps from

diester ($\pm$)-**1** which itself could be obtained according to literature procedures. The amino

group on the pyrrolidine ring, after Boc group deprotection, enables installation of the linker

and biotin tag as described in full detail in the Supplemental Information. Biotinylated DDA

analogs, including Biotin-PyrDTDO (BPD) and Biotin-GlyPyrDTDO (BGPD), were generated in

this fashion and used as affinity tags to identify DDA bound proteins (Fig. 2A).

The protein disulfide isomerase (PDI) family member AGR2 was previously implicated

in disulfide bonding of EGFR [12], and DDAs alter EGFR disulfide bonding ([4] and Fig. 1D,

right panel). We hypothesized that AGR2 might be a target of DDAs and examined if

biotinylated DDAs bind covalently with recombinant AGR2. Biotin-GlyPyrDTDO labeled AGR2

in a concentration-dependent manner (Fig. 2B). Under these conditions, Biotin-GlyPyrDTDO

labeled monomeric, but not dimeric AGR2. We further hypothesized that in addition to EGFR,

DR4 or DR5 may be AGR2 client proteins. Stable knockdown of AGR2 in breast cancer cells

increased disulfide-mediated DR5 oligomerization in the absence of DDA treatment (Fig. 2C).

AGR2 forms mixed disulfide bonds to its client proteins with Cys81 of its thioredoxin-like repeat

[13], and Cys81 is the only Cys residue present in AGR2. Therefore, we examined if mutating

Cys81 to Ser prevented DDA labeling. The C81S AGR2 mutant was not labeled by Biotin-

GlyPyrDTDO (Fig. 2D). AGR3 is the PDI paralog most closely related to AGR2 and it was also

labeled by Biotin-GlyPyrDTDO. Addition of a molar excess of dMtcyDTDO to the labeling

reactions partially prevented Biotin-GlyPyrDTDO binding to AGR2 and AGR3 (Fig. 2D),

suggesting competition for binding to the same Cys residue(s). A comparison of the ability of

dMtcyDTDO and tcyDTDO to block Biotin-GlyPyrDTDO binding to AGR2 indicated that

dMtcyDTDO was a more effective competitor, indicating that competition in this binding assay

reflects DDA potency in cancer cells (Fig. 2E, S1E, S1F). AGR2 is expressed at low levels in

MDA-MB-468 cells, and disulfide-mediated AGR2 complexes with client proteins were not

detected in either the presence or absence of dMtcyDTDO treatment (Fig. 2F). Enforced AGR2

expression revealed formation of dimeric AGR2 that was not observed with the C81S AGR2

active site mutant. Treatment of the cells with dMtcyDTDO blocked dimer formation by wild

type AGR2, consistent with blockade of C81S by dMtcyDTDO binding.

Identification of PDIA1 and ERp44 as DDA target proteins: A panel of three breast cancer cell

lines was treated with Biotin-PyrDTDO to label DDA target proteins in intact cells. In addition to

affinity purification of endogenous biotinylated proteins, such as carboxylases, which use biotin

as a cofactor, two biotinylated proteins were observed exclusively in the context of cell labeling

with Biotin-PyrDTDO in multiple experiments and termed DDA target proteins 1 and 2

(DDAT1/2). DDAT1/2 migrated at 60 kDa and 44 kDa, respectively, and were observed in

three different breast cancer lines in crude cell extracts and in samples purified using

Streptavidin-agarose (Fig. 3A). The Streptavidin-agarose purification was scaled up to

generate DDAT1/2 in sufficient amounts for identification by mass spectrometry, and aliquots

of this preparation and the corresponding crude extracts were analyzed by blotting with

Streptavidin-Alkaline Phosphatase, and by silver staining (Fig. 3B). DDAT1 and DDAT2 from a

Coomassie stained gel were excised and analyzed by trypsin digestion followed by mass

spectrometry as described in Materials and Methods. DDAT1 and DDAT2 were identified as

the PDI-family members PDIA1 and ERp44, respectively (Fig. S2A, S2B). In support of these

assignments, ERp44, PDIA1, and AGR2 were isolated from the luminal A T47D and luminal B

BT474 breast cancer cell lines treated with Biotin-PyrDTDO using Streptavidin-Agarose (Fig.

3C, left panel). In additional Biotin-PyrDTDO pulldown experiments, ERp44 and PDIA1 were

detected, but the PDIs ERp57 and ERp5 were not (Fig. 3D). Endogenous proteins are

biotinylated via amide linkages. Thus, DDAT1 and DDAT2 were distinguished from

endogenous biotinylated proteins by disruption of Biotin-DDA/target protein linkages with the

reducing agent 2-Mercaptoethanol (Fig. S2C). Immunoprecipitation with an AGR2 antibody

showed that ERp44 and PDIA1 co-purify with AGR2 from cells irrespective of treatment with

Biotin-PyrDTDO (Fig. 3C, left panel). ERp44 immunoprecipitates contained PDIA1, but AGR2

co-purification was not detected. Biotin-PyrDTDO/Streptavidin-Agarose also affinity purified

ERp44 and PDIA1 from the MDA-MB-468 Triple-Negative breast cancer cell line (Fig. 3C, right

panel). In vitro binding studies showed that Biotin-PyrDTDO labeled ERp44 and PDIA1, and in

both cases, biotinylation was blocked by pre-incubation with a molar excess of dMtcyDTDO

(Fig. 3E). If ERp44 and PDIA1 are cellular DDA targets, then DDA treatment may alter the

pattern of ERp44 and PDIA1 disulfide bonding with their client proteins. Consistent with this,

treatment of breast cancer cells with dMtcyDTDO reduced disulfide bonding of ERp44 and

PDIA1 with multiple proteins (Fig. 3F, red arrows) with minimal effect on monomeric levels of

ERp44 and PDIA1. PDIA1 is a validated target for anticancer drugs [16]. However, ERp44 has

not been investigated extensively as a therapeutic target for cancer, and small molecule

ERp44 inhibitors have not been reported. Small molecule active site AGR2 inhibitors have not

been reported, but under the conditions used here, AGR2 binding to client proteins was not

detected. Therefore we focused most of our efforts on examining the consequences of ERp44

inhibition by DDAs.

DDAs block the ER retention function of ERp44: ERp44 carries out two important roles by

disulfide bonding with client proteins and cycling between the Golgi and the ER as modeled in

Fig. 4A. First, ERp44 retrotranslocates inappropriately oligomerized proteins from the Golgi to

ER for additional rounds of disulfide bond formation and shuffling. Second, ERp44 forms

disulfide bonds with select partner proteins to retain them in the ER/Golgi, preventing their

secretion. To determine if DDAs block ERp44 retention of partners by binding to the ERp44

active site, MDA-MB-468 cells were treated with dMtcyDTDO for 24 h and the levels of the

ERp44 partner protein ERAP1 were examined. DMtcyDTDO treatment decreased cellular

ERAP1 levels (Fig. 4B). Quantitation of dMtcyDTDO effects on ERAP1 levels and dMtcyDTDO

binding to its client proteins across multiple experiments showed that dMtcyDTDO decreased

ERAP1 levels and ERp44 binding to client proteins in a statistically significant manner (Fig. 4C,

D). ERAP1 migrates primarily as a monomer even under nonreducing conditions, indicating

that ERAP1 disulfide bonding with ERp44 is transient. Enforced expression of wild type ERp44

(CXXS) resulted in the formation of a higher molecular mass band recognized by ERAP1

antibodies (Fig. 4E, red arrow). This band was not observed when the C58S ERp44 active site

mutant (SXXS) was expressed, but was observed when the S61C mutant (CXXC) was

expressed. The ERAP1-ERp44 disulfide-bonded dimer was not observed when analyses were

performed under reducing conditions (not shown) and was reduced if the cells were treated

with the DDA dFtcyDTDO for 24h (Fig. 4E).

DDA-induced secretion of ERAP1 was examined by treating MDA-MB-468 breast

cancer cells or A431 skin cancer cells with dMtcyDTDO for 24 h and examining ERAP1 levels

in cell extracts and in the culture medium. ERAP1 levels in the medium dramatically increased

upon dMtcyDTDO treatment (Fig. 4F, upper panels), and dMtcyDTDO induced ERAP1

secretion in a statistically significant manner (Fig. 4F, lower panels).

A role for ERp44 in the control of Death Receptor and EGFR disulfide bonding: Immunoblot

analysis of a panel of breast cancer cell lines revealed that PDIA1 levels were similar among

the lines (Fig. 5A). In contrast, ERp44, AGR2, and AGR3 levels varied among the lines, as did

the oxidoreductase ERO1, which plays a key role in producing oxidizing equivalents for

disulfide bond formation [16, 34-36]. A BT474 breast cancer cell line stably overexpressing

ERp44 was generated to examine the effect of elevated ERp44 levels on DR4/5 disulfide

bonding and DDA responses. ERp44 overexpression reduced DDA induction of high molecular

mass DR4 oligomers, and to a lesser extent high molecular mass DR5 oligomers (Fig. 5B, left

panel, red arrows). In both cases, the overall pattern was a shift from high to lower molecular

mass disulfide-bonded species. These oligomeric protein forms were not observed when the

analyses were performed under reducing conditions (Fig. 5B, right panel). Next, ERp44

knockdown MDA-MB-468 cells were generated to examine the effect of ERp44 depletion on

DR4/5 disulfide bonding. Although we were only able to achieve modest ERp44 knockdown, in

multiple independent experiments ERp44 knockdown consistently increased high molecular

mass oligomers of DR4, DR5, and EGFR in the absence of DDA treatment (Fig. 5C, red

arrows; expanded view in Fig. 5D, red asterisks; Fig. S2D). ERp44 knockdown augmented

PDIA1 oligomerization and increased dMtcyDTDO induction of Caspase 8 cleavage/activation

(CC8). Together, the results of the AGR2 and ERp44 knockdown experiments are consistent

with roles for AGR2 and ERp44 in facilitating native disulfide bond formation of DR4 and DR5,

and for DDAs in blocking these AGR2 and ERp44 functions. The observation that ERp44

knockdown increases PDIA1 binding to clients (Fig. 5C, 5D) and that ERp44 and PDIA1 form

complexes (Fig. 3C) may indicate that PDIA1 and ERp44 chaperone native disulfide bond

formation in a coordinated manner.

We next examined if mutations within the thioredoxin-like repeat of ERp44 alter its

disulfide bonding in the presence or absence of DDA treatment. Overexpressed, wild type

ERp44 exhibited a high level of disulfide bond-mediated oligomer formation with client proteins

(Fig. 6A). Both endogenous and overexpressed ERp44 monomers migrated as a doublet,

which may result from differential O-linked glycosylation [37]. The C58S ERp44 mutant

(SXXS), which lacks a Cys residue in its thioredoxin-like repeat, showed lower binding to client

proteins at similar expression levels. The S61C ERp44 mutant (CXXC) exhibited slightly

increased oligomerization compared with wild type ERp44.

We previously showed that DDAs alter the electrophoretic mobility of EGFR and CUB

Domain-Containing Protein 1 (CDCP1), block the formation of complexes between these two

proteins, and alter their tyrosine phosphorylation patterns [38]. Overexpression of both ERp44

point mutants was associated with increased oligomerization of EGFR and CDCP1, but did not

alter the ratio between the full length (FL) and cleaved (Cl.) forms of CDCP1. Analysis of the

samples from Fig. 4E for DR4/5 oligomerization revealed that the C58S ERp44 mutant induced

the strongest potentiation of DR5 upregulation by dFtcyDTDO (Fig. 6B). Expression of this

mutant was also associated with partial oligomerization of EGFR in vehicle treated cells that

correlated with increased expression of the ERS marker GRP78 (Fig. 6B). Overall, these

observations indicate that ERp44^{C58S}, and to a lesser extent, ERp44^{S61C}, interfere with disulfide

bond formation of ERp44 clients, resulting in higher baseline ERS and potentiating DDA

effects on DR4 and DR5. However, these observations do not explain why cancer cells that

overexpress EGFR or HER2 have heightened DDA sensitivity [2, 3].

T47D breast cancer cells express low levels of EGFR, HER2, and HER3 and were

previously used as a model system to study the effects of EGFR overexpression on DDA

responses [2, 3, 39]. Examination of ERp44 and PDIA1 disulfide bonding in control and EGFR

overexpressing T47D cells showed that EGFR overexpression resulted in recruitment of

ERp44 and PDIA1 into disulfide-bonded protein complexes (Fig. 6C, left panels). Quantitation

of the percentages of oligomeric ERp44 and PDIA1 in three experiments employing

independently derived pairs of vector control and EGFR overexpressing T47D cell lines

showed that EGFR overexpression increased ERp44 and PDIA1 oligomerization (Fig. 6C, right

panel). Oligomeric, disulfide-bonded forms of ERp44 and PDIA1 are expected to be

inaccessible for binding to newly produced client proteins, while the monomeric forms of

ERp44 and PDIA1 are accessible to facilitate native disulfide bonding of their client proteins.

Thus, EGFR overexpression may increase DDA sensitivity by reducing the pools of monomeric

ERp44 and PDIA1 that DDAs must inhibit in order to block native disulfide bond formation of

ERp44 and PDIA1 client proteins.

In addition to chaperoning DR4 and DR5 disulfide bonding, we hypothesized that

ERp44, PDIA1, or AGR2 disulfide bonding with DR4 or DR5 may sequester DR4 and DR5 in

the ER or Golgi, preventing their trafficking to the cell surface. If so, blocking active site ERp44,

PDIA1, and AGR2 Cys residues, DDA treatment may permit DR4 and DR5 trafficking to the

plasma membrane. Cell labeling using membrane impermeable, protein-reactive probes is

useful for confirming the presence of receptors at the cell surface. Surface biotinylation studies

performed on cells after treatment with dMtcyDTDO showed increased cell surface levels of

DR4 and DR5, irrespective of whether labeling was performed with amine- or thiol-reactive

cell-impermeable biotinylation probes (Fig. 6D). This increased accumulation of DR4/5 at the

cell surface may explain previous findings showing that DDAs synergize with the DR4/5 ligand

TRAIL to induce cancer cell apoptosis [4].

DDAs potentiate non-covalent binding of DR5 to ERp44: Data in Fig. 5 and 6 showing that

ERp44 knockdown and overexpression of ERp44 mutants alter DR5 levels or oligomerization

suggested that DR5 may be an ERp44 client protein. We reasoned that if DR5 is an ERp44

client, the two proteins may bind to each other. ERp44 co-immunoprecipitation experiments

showed that ERp44 associates with DR5 and to a lesser extent DR4 in MDA-MB-468 cells,

and these interactions were potentiated by DDA treatment (Fig. 7A; Fig. S2E). Unexpectedly,

these were non-covalent ERp44/DR5 complexes because although the analyses were

performed under non-reducing conditions, each protein migrated as a monomer. ERp44 co-

immunoprecipitation studies were next carried out in cells ectopically expressing ERp44 or the

C58S or S61C ERp44 mutants. DR5 immunoprecipitated with wild type ERp44 and the S61C

ERp44 mutant, but not the C58S ERp44 mutant (Fig. 7B, left panel). DDA treatment increased

DR5 and decreased PDIA1 association with ERp44, which may be partly explained by

changes in PDIA1 and ERp44 expression levels (Fig. 7B, right panel). Since DDAs increase

DR5 levels, which could account for the increased DR5 binding to ERp44, we used the tet-ON

system to drive transient high-level DR4 or DR5 expression in the absence of DDA treatment.

Doxycycline-induced DR4 expression increased DR4 binding to ERp44 and binding was

further enhanced by DDA treatment (Fig. 7C). Doxycycline-induced DR5 expression caused

detectable DR5 binding to ERp44, which was strongly enhanced by DDA treatment. The

results in Fig. 7A-C demonstrate that DR5 forms a non-covalent complex with ERp44 and that

formation of this complex is strongly enhanced by DDA treatment. To determine if ERp44 in

dMtcyDTDO-treated cells associates with the disulfide bond-rich extracellular domain of DR5,

ERp44 immunoprecipitation experiments were performed in cells that inducibly express wild

type DR5 or the Δ81-178 DR5 deletion construct lacking part of the disulfide-rich region. The

results indicate that while wild type DR5 bound ERp44 in the context of dMtcyDTDO treatment,

ERp44 binding to DR5[Δ81-178] was barely detectable (Fig. 7D). This result suggests that

DDA binding to the ERp44 active site Cys residue not only inactivates its catalytic function, but

increases its association with the disulfide-rich region of DR5. These catalytically null

ERp44(DDA)-DR4/5 complexes may sequester DR4/5, preventing catalytically active forms of

ERp44, PDIA1, or AGR2 from chaperoning native DR5 disulfide bond formation.

The results presented thus far indicate that DDAs alter the disulfide bonding status of

DR4 and DR5, which correlates with inhibition of ERp44 and PDIA1 binding to their client

proteins. However, it is unclear if genetic perturbation of DR5 disulfide bonding mimics the

effects of DDAs on DR5, which include upregulation of DR5 protein levels and increased pro-

apoptotic signaling. This issue was addressed by inducibly expressing DR5 mutants in which a

Cys residue was mutated to Ser to disrupt one of the seven extracellular DR5 disulfide bonds.

The C119S mutant was constructed to disrupt the center/fourth disulfide bond, and the C160S

mutant was constructed to disrupt the seventh, most C-terminal DR5 disulfide bond. As

observed previously using this doxycycline-inducible expression system [39], induction of wild

type DR5 expression weakly increased DR5 expression (Fig. 7E). However,

dMtcyDTDO/doxycycline co-treatment further increased DR5 expression, demonstrating DR5

stabilization by DDA treatment. The C119S and C160S DR5 mutants were highly expressed in

both the monomeric and oligomeric forms, and this was associated with Caspase 8

cleavage/activation. DMtcyDTDO treatment modestly increased Caspase 8 cleavage, but did

not further increase expression of the C119S or C160S DR5 mutants. Expression of the DR5

mutants did not upregulate the ERS marker GRP78, indicating that the mutation-driven

upregulation of DR5 was not a result of ER stress induction. These results indicate that

disruption of individual DR5 disulfide bonds is sufficient to promote high-level DR5 protein

expression and pro-apoptotic signaling that is not further enhanced by DDA treatment.

Discussion

Models for DDA mechanisms of action and compound selectivity: The findings presented here

suggest that DDAs bind to the PDI-family proteins ERp44, PDIA1, AGR2, and AGR3 and block

their ability to chaperone the disulfide bonding of subsets of client proteins. The resulting

alteration of disulfide bonding causes upregulation of DR5, oligomerization of DR4 and DR5,

and accumulation of DR4 and DR5 at the cell surface. Oligomerization of DR4 and DR5

causes Caspase 8 activation, which in turn activates Caspase 3 to mediate apoptosis.

Caspase activation plays a key role in downregulating EGFR and HER2, and decreasing Akt

phosphorylation because these DDA responses are largely ablated by a pan-Caspase inhibitor

[4]. To our knowledge, DDAs are the first small molecule agents that block the active site Cys

residues of ERp44 and AGR2/3. Further, DDAs are the first experimental anticancer agents

that trigger DR4 and DR5-mediated Caspase 8 activation by inducing disulfide bond-mediated

DR4/5 clustering in the absence of their ligand, TRAIL. A recent report showed that the

extracellular domain of DR5 acts to prevent its oligomerization in the absence of TRAIL [11].

DDAs may overcome this autoinhibition by altering the conformations of the DR4/5

extracellular domains or favoring intermolecular disulfide bonding between DR4/5 extracellular

domains over intramolecular disulfide bonding. Consistent with these interpretations,

mutational disruption of individual DR5 disulfide bonds is sufficient to trigger DR5 upregulation,

oligomerization, and downstream activation of Caspase 8 (Fig. 7E).

Previous reports showed that overexpression of EGFR amplifies multiple DDA responses,

including ERS induction, DR5 upregulation, HER1-3 downregulation, and reduction of Akt

phosphorylation [2-4]. Discovery of ERp44 and PDIA1 as DDA target proteins led to the

observation that EGFR overexpression reduced the fractions of ERp44 and PDIA1 in the

monomeric forms and increased the relative abundance of the oligomeric forms. Since ERp44

and PDIA1 oligomerization with client proteins involves their active site Cys residues, only

monomeric ERp44 and PDIA1 are accessible for facilitating native disulfide bonding of client

proteins. Because DDAs covalently modify monomeric ERp44 and PDIA1 (Fig. 3), these

observations may explain the molecular basis for how EGFR overexpression sensitizes cells to

DDAs. Given the importance of EGFR as a potential oncogenic driver in a number of cancers,

including glioblastoma [40, 41] and Triple-Negative Breast Cancer [42-44], DDAs may have

utility against these malignancies.

Because of the relative structural simplicity of DDA molecules, the mechanisms

responsible for their target protein selectivity are of interest. PDIA1 has been investigated as a

target for anticancer therapeutics, and PDIA1 inhibitors have established anticancer efficacy in

preclinical models [45, 46]. AGR2 contributes to tumor progression by facilitating EGFR

disulfide bonding and surface presentation [47]. However, AGR2 may also be secreted and

promote tumor angiogenesis [48, 49], and act on cell surface receptors [50]. Monoclonal

antibodies designed to block extracellular AGR2 functions are under development [50, 51], but

agents that ablate the role of AGR2 in the folding of client proteins are not available.

The canonical thioredoxin-like catalytic site sequence CXXC is featured in PDIA1 and

several other PDI enzymes (reviewed in [15]). Interestingly, AGR2, AGR3, and ERp44 share a

non-canonical CXXS thioredoxin-like repeat sequence. A report employing both yeast genetics

and biochemical enzyme assays showed that the essential function of the PDI enzyme for

yeast cell survival is catalysis of disulfide exchange or "scramblase" activity and that this could

be carried out by the CXXS mutant of PDI (PDI^{CXXS}) [17]. Enzyme assays showed that PDI^{CXXS}

was able to reactivate inappropriately disulfide bonded RNAse A via its scramblase activity. In

contrast, PDI^{CXXS} was unable to catalyze disulfide bond oxidation or reduction like the wild type

PDI^{CXXC} enzyme.

The C-terminal Cys residue in the CXXC motif of PDIs accelerates substrate release.

Consequently, CXXS substrate-trapping mutants of PDIs have been used to isolate mixed

disulfide bonded dimers with their client/substrate proteins [52, 53]. That the CXXS motif is

associated with slow client protein off-rate is consistent with the established roles of ERp44 in

retaining specific client/partner proteins such as ERO1 [22] and ERAP1 [25] in the ER, and in

retrotranslocation of inappropriately disulfide bonded Adiponectin [54, 55] and IgM [21, 56]

from the Golgi to the ER (see model in Fig. 4A). Thus, the CXXS motifs of AGR2, AGR3, and

ERp44 bound to DDAs may exhibit a slow DDA off-rate for the same reason, and could explain

stable DDA binding to these enzymes. Our current working model for DDA binding to AGR2,

AGR3, and ERp44 is shown in Fig. 7F. To wit, binding of Biotin-PyrDTDO is initiated by

nucleophilic attack of the thiosulfonate group of Biotin-PyrDTDO by the thiolate nucleophile of

the Cys residue of the CXXS motif. This results in reversible ring opening. We hypothesize that

the rate of ring closure and DDA compound excision is reduced by stabilization of the open

form of the DDA by solvation of the sulfinate group through interactions with functional groups

in or near the PDI active site. The reason for stable DDA binding to PDIAI, but not ERp57 or

ERp5 is unclear since each protein contains two canonical CGHC repeats. Further work is

required to determine which PDIA1 Cys residue binds DDAs and to identify features in the

local environment that stabilize DDA binding.

It is clear from the experiments with intact cancer cells that DDAs alter the disulfide

bonding of DR4, DR5, and EGFR and change the pattern of disulfide bonding between ERp44

and PDIA1 and their client proteins. It is difficult to compare these DDA actions with those of

published PDI inhibitors since most of these studies did not examine perturbation of protein

disulfide bonding in cancer cells, but rather examined compound activity in the Insulin

reduction/aggregation assay. While the Insulin reduction assay is valuable for optimizing the

potency of PDI inhibitors, it may not be salient in instances where disulfide isomerase activity,

ER retention, or Golgi-ER retrotranslocation are more important than disulfide oxidase or

reductase activity.

The presence of 21 different PDIs in humans and the diverse phenotypes of the

corresponding knockout mice (e.g., [25, 57-59]) suggests that individual PDIs chaperone the

disulfide bonding, ER retention, or Golgi-ER retrotranslocation of different subsets of client

proteins. For instance, ERp44 is unique in its ability to retain the aminopeptidase ERAP1 in the

ER/Golgi and this results in ERAP1 secretion in ERp44 knockout mice or ERp44-deficient cell

lines [25]. A biological consequence of this is that ERp44 deletion is associated with

hypotension in mice since secreted ERAP1 destabilizes Angiotensin II, leading to reduced

blood pressure. It will be of interest to determine if DDAs exhibit antihypertensive activity.

ERAP1 functions as a "molecular ruler" to process peptides for loading onto MHC I for

presentation to T cells [60-62]. ERAP1 inhibitors or ERAP1 knockdown changes the spectrum

of the MHC I-associated peptides that constitute the "immunopeptidome." Since DDAs cause

ERAP1 secretion, it will be important to determine if DDAs alter the immunopeptidome, and if

so, how this impacts anticancer immunity.

In summary, the results presented here establish perturbation of disulfide bonding by

inhibition of subsets of PDIs as a new strategy for activating DR5 and DR4 in cancer cells

independently of their engagement by agonistic antibodies or TRAIL. Further, DDAs

upregulate DR5 and increase DR4 and DR5 localization to the cell surface. Consequently,

DDAs may overcome some of the shortcomings associated with TRAIL analogs and agonistic

antibodies specific for DR4 and DR5. Additional research is needed to elucidate the

mechanisms by which disrupting individual DR5 disulfide bonds increases its expression

levels, disulfide-mediated oligomerization, and activation of Caspase 8. In future studies, it will

be important to determine if C-S point mutants of DR4 mimic the DR5 mutants and drive pro-

apoptotic signaling. It will also be interesting to examine whether activated point mutants of

DR4 and DR5 exhibit the same cancer-selective killing as their ligand, TRAIL.

Materials and Methods

In vivo tumor studies and histochemical analysis

Breast tumor formation was initiated in NOD-SCID-gamma (NSG) mice obtained from Jackson

Laboratories (Bar Harbor, ME USA) by injecting 1×10^6 cancer cells into the #4 mammary

fat pads as described previously [4]. Animals bearing HER2+ BT474 tumors [2] were treated

with the vehicle or 20 mg/kg dMtcyDTDO or dFtcyDTDO by intraperitoneal injection for four

consecutive days. Tissue samples fixed in 4% paraformaldehyde/Phosphate-Buffered Saline

(PBS) and paraffin-embedded were sectioned and stained with hematoxylin and eosin (H&E)

by the University of Florida Molecular Pathology Core (https://molecular.pathology.ufl.edu/).

Streptavidin-Agarose Pulldowns

Cells were treated with Biotin-PyrDTDO with or without the pan-Caspase inhibitor Q-VD-OPH

to prevent apoptosis (MedChemExpress, Monmouth Junction, NJ USA) for 16 h in 10% FBS–

DMEM, followed by the addition of 25 mM N-ethylmaleimide (NEM) (Thermo Scientific,

Rockford, IL USA) for 20 min at 37^oC. Cells were subsequently scraped directly into the media,

washed twice with ice-cold PBS, and pelleted by centrifugation. Cells were extracted in buffer

(20 mM HEPES (pH 7.4), 1 mM EDTA, 1mM EGTA, 5% Glycerol, 1 nM Microcystin, 1 mM

Na_3VO_4, and 40 mM $Na_2H_2P_2O_7$) containing 1% Triton X-100. Samples were sonicated,

followed by centrifugation for 20 min at 4^oC. The supernatant was pre-cleared with Sepharose

CL-6B (Amersham Biosciences AP, Uppsala, Sweden) for 2h at 4^oC. Pre-cleared samples

were then incubated with Streptavidin (SA)-Agarose (Thermo Scientific, Rockford IL USA) and

mixed overnight at 4^oC. The SA-Agarose was washed three times with extraction buffer

containing 0.1% Triton X-100, and eluted by boiling in 2X-SDS Sample Buffer containing 5 mM

biotin. Samples were analyzed by immunoblot, silver stain, or probing proteins transferred to

nitrocellulose membranes with Streptavidin-conjugated Alkaline Phosphatase (Rockland

Immunochemicals, Inc., Limerick, PA USA).

Disulfide Bond-mediated Oligomerization

To evaluate protein disulfide bond-mediated oligomerization, cells were treated with DDAs for

24 h in 10% FBS–DMEM. Cells were subsequently scraped into the media and pelleted by

centrifugation. Cell pellets were extracted directly into boiling 2X-SDS-Sample Buffer

containing 100 mM NEM and boiled for 10 min. Samples were sonicated, followed by

centrifugation for 20 minutes at room temperature, and subsequently analyzed by SDS-

PAGE/immunoblot.

Surface Biotinylation

Cell surface proteins were labeled by incubating cells with 1.6 mM Sulfo-NHS-SS-Biotin

(Thermo Fisher Scientific, Waltham, MA USA) or 1.6 mM Biotin-dPEG$_3$-Maleimide

(MilliporeSigma, St. Louis, MO USA) in PBS, pH 8.0 for 30 minutes at 4 °C. The cells were

subsequently washed three times with PBS, and proteins were extracted in 1% Triton X-100

extraction buffer in the absence of reducing agents. Cell extracts were then mixed with

Streptavidin-agarose beads (Thermo Fisher Scientific, Waltham, MA USA) for 2.5 h at 4^{o}C.

Following centrifugation, the supernatant containing non-biotinylated proteins (flow-through)

was retained and boiled in 2X-SDS-Sample Buffer containing 1% 2-mercaptoethanol.

Streptavidin-agarose beads bound to biotinylated proteins were washed four times with 0.1%

Triton X-100 extraction buffer, and eluted by boiling in 2X-SDS-Sample Buffer containing 1%

2-mercaptoethanol. Samples were then analyzed by immunoblot or blotting with Streptavidin-

conjugated Alkaline Phosphatase.

His$_6$-AGR2 TALON Purification

Competent BL21(DE3) cells were transformed with either the AGR2/pET-45b(+) or AGR2

C81S/pET-45b(+) vectors. Protein expression was induced by the addition of 50 μM Isopropyl

β-D-1-thiogalactopyranoside (Fisher Scientific, Fair Lawn, NJ USA) to bacterial cell cultures for

2.5 hours with shaking at 37^{o}C. Bacterial cells were subsequently pelleted and dissolved in

Mouse Tonicity Phosphate Buffered Saline (MTPBS) containing 1% Triton X-100 and

sonicated for 20 seconds on ice, followed by centrifugation for 10 min at 4^{o}C to pellet cell

debris. The supernatant was transferred to a new tube, followed by the addition of TALON

Resin (Clontech, Mountain View, CA USA) and mixed for 30 min at 4^{o}C. The TALON Resin

was pelleted by centrifugation and washed twice with MTPBS and twice with MTPBS

containing 30 mM Imidazole. Protein bound to the TALON Resin was eluted with 400 mM

Imidazole and precipitated with 60% saturated ammonium sulfate. The precipitated protein

was pelleted by centrifugation, dissolved in MTPBS containing 1 mM DTT and 1 mM EDTA,

and subsequently dialyzed against a solution of MTPBS containing 10 μM DTT and 30%

Glycerol.

DDA-AGR2 Binding Experiments

Purified His$_6$-AGR2 or His$_6$-AGR2[C81S], as prepared above, or His$_6$-AGR2 (PRO-580) or

His$_6$-AGR3 (PRO-242) obtained from ProSpec (New Brunswick NJ USA) were incubated with

or without varying concentrations of the unlabeled DDA competitors, tcyDTDO or dMtcyDTDO,

and 30 μM Biotin-GlyPyrDTDO or Biotin-PyrDTDO in buffer (20 mM HEPES (pH 7.4), 1 mM

EDTA, 1mM EGTA, 5% Glycerol, 1 nM Microcystin, 1 mM Na_3VO_4, and 40 mM $Na_2H_2P_2O_7$)

containing 1% Triton-X 100 for 30 min at 37^oC. Preliminary studies showed that Biotin-

GlyPyrDTDO or Biotin-PyrDTDO exhibited indistinguishable labeling of proteins in intact cells

and purified proteins in vitro, and are referred to collectively as "Biotin-DDA" and the two

probes were used interchangeably. Reactions were stopped by boiling the samples in 2X-

SDS-Sample Buffer containing 100 mM NEM for 10 min. Biotin-DDA labeling of protein was

verified by blotting with Streptavidin-Alkaline Phosphatase. ERp44 (PRO-547) and PDIA1

(ENZ-262) were obtained from ProSpec and used in experiments examining the binding of

these proteins to Biotin-PyrDTDO.

Quantitative Analysis of Immunoblot Results

Immunoblots were scanned, converted to grayscale and inverted using Adobe Photoshop

(Berkeley, CA). Bands were quantified using ImageJ (NIH, Bethesda, MD). Expression levels

were normalized using Actin as a loading control.

Chemical Synthesis of DDAs

Full experimental and characterization details associated with the synthesis of (±)-

BocPyrDTDO, Biotin-GlyPyrDTDO, and Biotin-PyrDTDO are provided in the Supplemental

Information. Detailed discussion of the syntheses and analytical characterization of

dMtcyDTDO and dFtcyDTDO will be published elsewhere.

Detailed Methods in Supplemental Information

At least three biological replicates of each of the studies presented were performed. Please see Supplemental Information for in-depth descriptions of approaches relating to cell culture, preparation of cell extracts, immunoblot analysis, construction of retroviral and lentiviral vectors and their use to generate stable cell lines, DNA and protein synthesis assays, cell viability assays, sample preparation for analysis by tandem mass spectrometry, and statistical methods..

Acknowledgements

These studies were supported in part by grants from the Florida Breast Cancer Foundation (BL

and RC), the Ocala Royal Dames for Cancer Research (BL), and the Office of the Assistant

Secretary of Defense for Health Affairs through the Breast Cancer Research Program under

Award Nos. W81XWH-15-1-0199 (BL) and W81XWH-15-1-0200 (RC), and NIH/NCI grant

CA252400 (BL). RF is grateful to the University of Florida for a Graduate School Fellowship.

EY thanks the Dr. Howard and Brenda Sheridan Fund in Chemistry for a 2020 Howard and

Brenda Sheridan Summer Fellowship. AG is grateful to the Tarrant Organic Chemistry Fund

for a 2021 Tarrant Summer Graduate Research Fellowship. Mass spectrometric data on DDA

protein targets were obtained by the Proteomics and Mass Spectrometry Facility at the

Interdisciplinary Center for Biotechnology Research. Mass spectrometric data on DDA

compound structures and DDA adducts were obtained by the UF Department of Chemistry

Mass Spectrometry Research and Education Center supported, in part, by the National

Institutes of Health (NIH S10 OD021758-01A1).

Conflicts of Interest

None.

References

608 1 DeSantis CE, Ma J, Gaudet MM, Newman LA, Miller KD, Goding Sauer A *et al*. Breast
609   cancer statistics, 2019. *CA Cancer J Clin* 2019; 69: 438-451.
610

611 2 Ferreira RB, Law ME, Jahn SC, Davis BJ, Heldermon CD, Reinhard M *et al*. Novel
612   agents that downregulate EGFR, HER2, and HER3 in parallel. *Oncotarget* 2015; 6:
613   10445-10459.
614

615 3 Ferreira RB, Wang M, Law ME, Davis BJ, Bartley AN, Higgins PJ *et al*. Disulfide bond
616   disrupting agents activate the unfolded protein response in EGFR- and HER2-positive
617   breast tumor cells. *Oncotarget* 2017; 8: 28971-28989.
618

619 4 Wang M, Law ME, Davis BJ, Yaaghubi E, Ghilardi AF, Ferreira RB *et al*. Disulfide bond-
620   disrupting agents activate the tumor necrosis family-related apoptosis-inducing
621   ligand/death receptor 5 pathway. *Cell Death Discovery* 2019; 5: 153.
622

623 5 Wiley SR, Schooley K, Smolak PJ, Din WS, Huang CP, Nicholl JK *et al*. Identification
624   and characterization of a new member of the TNF family that induces apoptosis.
625   *Immunity* 1995; 3: 673-682.
626

627 6 Rieger J, Naumann U, Glaser T, Ashkenazi A, Weller M. APO2 ligand: a novel lethal
628   weapon against malignant glioma? *FEBS Lett* 1998; 427: 124-128.
629

630 7 Zamai L, Ahmad M, Bennett IM, Azzoni L, Alnemri ES, Perussia B. Natural killer (NK)
631   cell-mediated cytotoxicity: differential use of TRAIL and Fas ligand by immature and
632   mature primary human NK cells. *J Exp Med* 1998; 188: 2375-2380.
633

634 8 Ashkenazi A, Pai RC, Fong S, Leung S, Lawrence DA, Marsters SA *et al*. Safety and
635   antitumor activity of recombinant soluble Apo2 ligand. *J Clin Invest* 1999; 104: 155-162.
636

637 9 Thapa B, Kc R, Uludag H. TRAIL therapy and prospective developments for cancer
638   treatment. *J Control Release* 2020; 326: 335-349.
639

640 10 Wang M, Law ME, Davis BJ, Yaaghubi E, Ghilardi AF, Ferreira RB *et al*. Mechanistic
641   Elucidation of the Antitumor Properties of a Novel Death Receptor 5 Activator. *bioRxiv*
642   2019: 700906.
643

644 11 Pan L, Fu TM, Zhao W, Zhao L, Chen W, Qiu C *et al*. Higher-Order Clustering of the
645   Transmembrane Anchor of DR5 Drives Signaling. *Cell* 2019; 176: 1477-1489 e1414.
646

647 12 Dong A, Wodziak D, Lowe AW. Epidermal growth factor receptor (EGFR) signaling
648   requires a specific endoplasmic reticulum thioredoxin for the post-translational control of
649   receptor presentation to the cell surface. *J Biol Chem* 2015; 290: 8016-8027.
650

651 13 Park SW, Zhen G, Verhaeghe C, Nakagami Y, Nguyenvu LT, Barczak AJ *et al*. The
652   protein disulfide isomerase AGR2 is essential for production of intestinal mucus. *Proc*
653   *Natl Acad Sci U S A* 2009; 106: 6950-6955.
654
655 14 Schroeder BW, Verhaeghe C, Park SW, Nguyenvu LT, Huang X, Zhen G *et al*. AGR2 is
656   induced in asthma and promotes allergen-induced mucin overproduction. *Am J Respir*
657   *Cell Mol Biol* 2012; 47: 178-185.
658
659 15 Galligan JJ, Petersen DR. The human protein disulfide isomerase gene family. *Hum*
660   *Genomics* 2012; 6: 6.
661
662 16 Shergalis AG, Hu S, Bankhead A, 3rd, Neamati N. Role of the ERO1-PDI interaction in
663   oxidative protein folding and disease. *Pharmacol Ther* 2020; 210: 107525.
664
665 17 Laboissiere MC, Sturley SL, Raines RT. The essential function of protein-disulfide
666   isomerase is to unscramble non-native disulfide bonds. *J Biol Chem* 1995; 270: 28006-
667   28009.
668
669 18 Hampe L, Radjainia M, Xu C, Harris PW, Bashiri G, Goldstone DC *et al*. Regulation and
670   Quality Control of Adiponectin Assembly by Endoplasmic Reticulum Chaperone ERp44.
671   *J Biol Chem* 2015; 290: 18111-18123.
672
673 19 Long Q, Lei T, Feng B, Yin C, Jin D, Wu Y *et al*. Peroxisome proliferator-activated
674   receptor-gamma increases adiponectin secretion via transcriptional repression of
675   endoplasmic reticulum chaperone protein ERp44. *Endocrinology* 2010; 151: 3195-3203.
676
677 20 Cortini M, Sitia R. From antibodies to adiponectin: role of ERp44 in sizing and timing
678   protein secretion. *Diabetes Obes Metab* 2010; 12 Suppl 2: 39-47.
679
680 21 Cortini M, Sitia R. ERp44 and ERGIC-53 synergize in coupling efficiency and fidelity of
681   IgM polymerization and secretion. *Traffic* 2010; 11: 651-659.
682
683 22 Otsu M, Bertoli G, Fagioli C, Guerini-Rocco E, Nerini-Molteni S, Ruffato E *et al*.
684   Dynamic retention of Ero1alpha and Ero1beta in the endoplasmic reticulum by
685   interactions with PDI and ERp44. *Antioxid Redox Signal* 2006; 8: 274-282.
686
687 23 Yang K, Li DF, Wang X, Liang J, Sitia R, Wang CC *et al*. Crystal Structure of the
688   ERp44-Peroxiredoxin 4 Complex Reveals the Molecular Mechanisms of Thiol-Mediated
689   Protein Retention. *Structure* 2016; 24: 1755-1765.
690
691 24 Fraldi A, Zito E, Annunziata F, Lombardi A, Cozzolino M, Monti M *et al*. Multistep,
692   sequential control of the trafficking and function of the multiple sulfatase deficiency gene
693   product, SUMF1 by PDI, ERGIC-53 and ERp44. *Hum Mol Genet* 2008; 17: 2610-2621.
694
695 25 Hisatsune C, Ebisui E, Usui M, Ogawa N, Suzuki A, Mataga N *et al*. ERp44 Exerts
696   Redox-Dependent Control of Blood Pressure at the ER. *Mol Cell* 2015; 58: 1015-1027.

698   26   Tempio T, Anelli T. The pivotal role of ERp44 in patrolling protein secretion. *J Cell Sci*
699   2020; 133.
700

701   27   Lemmon MA, Schlessinger J. Cell signaling by receptor tyrosine kinases. *Cell* 2010;
702   141: 1117-1134.
703

704   28   Cha SS, Sung BJ, Kim YA, Song YL, Kim HJ, Kim S *et al.* Crystal structure of TRAIL-
705   DR5 complex identifies a critical role of the unique frame insertion in conferring
706   recognition specificity. *J Biol Chem* 2000; 275: 31171-31177.
707

708   29   Wang MX, Ferreira RB, Law ME, Davis BJ, Yaaghubi E, Ghilardi AF *et al.* A novel
709   proteotoxic combination therapy for EGFR+ and HER2+cancers. *Oncogene* 2019; 38:
710   4264-4282.
711

712   30   Bondi A. Van Der Waals Volumes + Radii. *Journal of Physical Chemistry* 1964; 68: 441-
713   +.
714

715   31   Gillis EP, Eastman KJ, Hill MD, Donnelly DJ, Meanwell NA. Applications of Fluorine in
716   Medicinal Chemistry. *Journal of Medicinal Chemistry* 2015; 58: 8315-8359.
717

718   32   Wang J, Sanchez-Rosello M, Acena JL, del Pozo C, Sorochinsky AE, Fustero S *et al.*
719   Fluorine in Pharmaceutical Industry: Fluorine-Containing Drugs Introduced to the
720   Market in the Last Decade (2001-2011). *Chemical Reviews* 2014; 114: 2432-2506.
721

722   33   Park BK, Kitteringham NR, O'Neill PM. Metabolism of fluorine-containing drugs. *Annual*
723   *Review of Pharmacology and Toxicology* 2001; 41: 443-470.
724

725   34   Frand AR, Kaiser CA. The ERO1 gene of yeast is required for oxidation of protein
726   dithiols in the endoplasmic reticulum. *Mol Cell* 1998; 1: 161-170.
727

728   35   Pollard MG, Travers KJ, Weissman JS. Ero1p: a novel and ubiquitous protein with an
729   essential role in oxidative protein folding in the endoplasmic reticulum. *Mol Cell* 1998; 1:
730   171-182.
731

732   36   Bertoli G, Simmen T, Anelli T, Molteni SN, Fesce R, Sitia R. Two conserved cysteine
733   triads in human Ero1alpha cooperate for efficient disulfide bond formation in the
734   endoplasmic reticulum. *J Biol Chem* 2004; 279: 30047-30052.
735

736   37   Sannino S, Anelli T, Cortini M, Masui S, Degano M, Fagioli C *et al.* Progressive quality
737   control of secretory proteins in the early secretory compartment by ERp44. *J Cell Sci*
738   2014; 127: 4260-4269.
739

740   38   Law ME, Ferreira RB, Davis BJ, Higgins PJ, Kim JS, Castellano RK *et al.* CUB domain-
741   containing protein 1 and the epidermal growth factor receptor cooperate to induce cell
742   detachment. *Breast Cancer Res* 2016; 18: 80.

Wang M, Law ME, Davis BJ, Yaaghubi E, Ghilardi AF, Ferreira RB *et al.* Disulfide bond-disrupting agents activate the tumor necrosis family-related apoptosis-inducing ligand/death receptor 5 pathway. *Cell Death Discov* 2019; 5: 153.

Stockhausen MT, Broholm H, Villingshoj M, Kirchhoff M, Gerdes T, Kristoffersen K *et al.* Maintenance of EGFR and EGFRvIII expressions in an in vivo and in vitro model of human glioblastoma multiforme. *Exp Cell Res* 2011; 317: 1513-1526.

Cho J, Pastorino S, Zeng Q, Xu X, Johnson W, Vandenberg S *et al.* Glioblastoma-derived epidermal growth factor receptor carboxyl-terminal deletion mutants are transforming and are sensitive to EGFR-directed therapies. *Cancer Res* 2011; 71: 7587-7596.

Liu D, He J, Yuan Z, Wang S, Peng R, Shi Y *et al.* EGFR expression correlates with decreased disease-free survival in triple-negative breast cancer: a retrospective analysis based on a tissue microarray. *Med Oncol* 2012; 29: 401-405.

de Ruijter TC, Veeck J, de Hoon JP, van Engeland M, Tjan-Heijnen VC. Characteristics of triple-negative breast cancer. *J Cancer Res Clin Oncol* 2011; 137: 183-192.

Toyama T, Yamashita H, Kondo N, Okuda K, Takahashi S, Sasaki H *et al.* Frequently increased epidermal growth factor receptor (EGFR) copy numbers and decreased BRCA1 mRNA expression in Japanese triple-negative breast cancers. *BMC Cancer* 2008; 8: 309.

Stojak M, Milczarek M, Kurpinska A, Suraj-Prazmowska J, Kaczara P, Wojnar-Lason K *et al.* Protein Disulphide Isomerase A1 Is Involved in the Regulation of Breast Cancer Cell Adhesion and Transmigration via Lung Microvascular Endothelial Cells. *Cancers (Basel)* 2020; 12.

Xu S, Liu Y, Yang K, Wang H, Shergalis A, Kyani A *et al.* Inhibition of protein disulfide isomerase in glioblastoma causes marked downregulation of DNA repair and DNA damage response genes. *Theranostics* 2019; 9: 2282-2298.

Gupta A, Dong A, Lowe AW. AGR2 gene function requires a unique endoplasmic reticulum localization motif. *J Biol Chem* 2012; 287: 4773-4782.

Guo H, Zhu Q, Yu X, Merugu SB, Mangukiya HB, Smith N *et al.* Tumor-secreted anterior gradient-2 binds to VEGF and FGF2 and enhances their activities by promoting their homodimerization. *Oncogene* 2017; 36: 5098-5109.

Jia M, Guo Y, Zhu D, Zhang N, Li L, Jiang J *et al.* Pro-metastatic activity of AGR2 interrupts angiogenesis target bevacizumab efficiency via direct interaction with VEGFA and activation of NF-kappaB pathway. *Biochim Biophys Acta Mol Basis Dis* 2018; 1864: 1622-1633.

790   50   Arumugam T, Deng D, Bover L, Wang H, Logsdon CD, Ramachandran V. New Blocking
791   Antibodies against Novel AGR2-C4.4A Pathway Reduce Growth and Metastasis of
792   Pancreatic Tumors and Increase Survival in Mice. *Mol Cancer Ther* 2015; 14: 941-951.
793

794   51   Liu AY, Kanan AD, Radon TP, Shah S, Weeks ME, Foster JM *et al*. AGR2, a unique
795   tumor-associated antigen, is a promising candidate for antibody targeting. *Oncotarget*
796   2019; 10: 4276-4289.
797

798   52   Eriksson O, Stopa J, Furie B. Identification of PDI Substrates by Mechanism-Based
799   Kinetic Trapping. *Methods Mol Biol* 2019; 1967: 165-182.
800

801   53   Stopa JD, Baker KM, Grover SP, Flaumenhaft R, Furie B. Kinetic-based trapping by
802   intervening sequence variants of the active sites of protein-disulfide isomerase identifies
803   platelet protein substrates. *J Biol Chem* 2017; 292: 9063-9074.
804

805   54   Wang ZV, Schraw TD, Kim JY, Khan T, Rajala MW, Follenzi A *et al*. Secretion of the
806   adipocyte-specific secretory protein adiponectin critically depends on thiol-mediated
807   protein retention. *Mol Cell Biol* 2007; 27: 3716-3731.
808

809   55   Wolf G. New insights into thiol-mediated regulation of adiponectin secretion. *Nutr Rev*
810   2008; 66: 642-645.
811

812   56   Anelli T, Ceppi S, Bergamelli L, Cortini M, Masciarelli S, Valetti C *et al*. Sequential steps
813   and checkpoints in the early exocytic compartment during secretory IgM biogenesis.
814   *EMBO J* 2007; 26: 4177-4188.
815

816   57   Jang I, Pottekat A, Poothong J, Yong J, Lagunas-Acosta J, Charbono A *et al*.
817   PDIA1/P4HB is required for efficient proinsulin maturation and ss cell health in response
818   to diet induced obesity. *Elife* 2019; 8.
819

820   58   Torres M, Medinas DB, Matamala JM, Woehlbier U, Cornejo VH, Solda T *et al*. The
821   Protein-disulfide Isomerase ERp57 Regulates the Steady-state Levels of the Prion
822   Protein. *J Biol Chem* 2015; 290: 23631-23645.
823

824   59   Nyirimigabo E, Jin M, Yang Z, Wang J, Zhai K, Mao Y *et al*. The role of ERp44 in
825   glucose and lipid metabolism. *Arch Biochem Biophys* 2019; 671: 175-184.
826

827   60   Sui L, Gandhi A, Guo HC. Crystal structure of a polypeptide's C-terminus in complex
828   with the regulatory domain of ER aminopeptidase 1. *Mol Immunol* 2016; 80: 41-49.
829

830   61   Chang SC, Momburg F, Bhutani N, Goldberg AL. The ER aminopeptidase, ERAP1,
831   trims precursors to lengths of MHC class I peptides by a "molecular ruler" mechanism.
832   *Proc Natl Acad Sci U S A* 2005; 102: 17107-17112.
833

62   Koumantou D, Barnea E, Martin-Esteban A, Maben Z, Papakyriakou A, Mpakali A *et al*.
Editing the immunopeptidome of melanoma cells using a potent inhibitor of endoplasmic
reticulum aminopeptidase 1 (ERAP1). *Cancer Immunol Immunother* 2019; 68: 1245-
1261.

Figure Legends

Figure 1 – Characterization of new, more potent, orally available DDAs that induce rapid tumor

regression.

A. Representative first and second generation DDAs (RBF3, DTDO, and tcyDTDO) and

illustration of interconversion between the linear and cyclic forms of the DDA pharmacophore.

B. Structures of new tcyDTDO derivatives dMtcyDTDO and dFtcyDTDO.

C. MDA-MB-468 cells were treated for 24 h with the indicated concentrations of DDAs and cell

viability (MTT) assays were performed. Three biological replicates were performed with similar

results.

D. MDA-MB-468 cells were treated for 24 h as indicated and subjected to immunoblot analysis

after non-reducing SDS-PAGE. O and M represent disulfide bonded oligomeric and

monomeric protein forms, respectively. (N=4 for each treatment group with similar within-group

results).

E. Levels and degree of oligomerization of DR4 (left panel) and DR5 (right panel) were

determined by densitometry analysis of bands across multiple experiments and subjected to

statistical analysis. Values plotted represent averages and error bars represent S.E. P-values

for pairwise comparisons were determined using Student's t-test.

F. Mice bearing BT474 cell xenograft tumors were treated once daily for five days with vehicle,

20 mg/kg dFtcyDTDO, or 20 mg/kg dMtcyDTDO. Two hours after the fifth treatment, mammary

fat pads/tumors were excised and tumor sections were stained with H&E. T denotes tumor

tissue, FP indicates mammary fat pads, and CN represents coagulation necrosis.

Figure 2 - AGR2 and AGR3 are DDA target proteins.

A. Abbreviated synthetic approach and chemical structures of the biotinylated DDA analogs

Biotin-PyrDTDO and Biotin-GlyPyrDTDO.

B. Purified, recombinant AGR2 was reacted with Biotin-GlyPyrDTDO for 1 h at 37°C, resolved

by non-reducing SDS-PAGE, transferred to membranes, and probed with Streptavidin-Alkaline

Phosphatase, and by anti-AGR2 immunoblot.

C. Vector control or AGR2 knockdown MDA-MB-468 cells were treated for 24 h as indicated

and analyzed by non-reducing (left panels) or reducing (right panels) immunoblot. Red arrow

indicates oligomerized forms of DR5 that are present at higher levels in the AGR2 knockdown

cells. O and M represent disulfide-bonded oligomeric and monomeric protein forms,

respectively.

D. Biotin-GlyPyrDTDO binding assays performed as in Fig. 2B, employing the C81S AGR2

mutant and AGR3 in addition to AGR2. In the indicated samples, the proteins were pre-

incubated with 600 μM dMtcyDTDO.

E. AGR2 binding assay performed as in Fig. 2B to examine the relative abilities of pre-

incubation of AGR2 with dMtcyDTDO or tcyDTDO to block subsequent binding of Biotin-

GlyPyrDTDO.

F. The indicated MDA-MB-468 stable cell lines were treated for 24 h with 5 μM dMtcyDTDO

and subjected to analysis by non-reducing SDS-PAGE/immunoblot. O, D, and M represent the

oligomeric, dimeric, and monomeric forms of AGR2 detected. P43/p41 and p18 denote

different Caspase 8 cleavage products.

Figure 3 - ERp44 and PDIA1 are additional DDA targets.

A. Breast cancer cell lines were treated for 16 h with vehicle, 100 μM Biotin-PyrDTDO, or 100

μM Biotin-PyrDTDO + 10 μM Q-VD-OPH for 16 h. Biotinylated proteins were isolated using

Streptavidin-coated beads and the crude extracts and Streptavidin pulldowns were analyzed

by blotting with Streptavidin-Alkaline Phosphatase. Red arrows indicate bands, designated

DDAT1 and DDAT2, present exclusively after Biotin-PyrDTDO treatment.

B. Biotin-PyrDTDO/Streptavidin pulldowns were performed in BT474 and MDA-MB-468 cells.

Purified material was analyzed by Streptavidin-Alkaline Phosphatase detection and silver stain.

C, left panel. The indicated breast cancer lines were treated with vehicle or 100 μM Biotin-

PyrDTDO for 16 h and extracts were subjected to Streptavidin-agarose pulldowns, or anti-

AGR2 or anti-ERp44 immunoprecipitations. The purified material was analyzed by immunoblot

with the indicated antibodies. C, right panel. BT474 and MDA-MB-468 breast cancer lines were

treated with vehicle or 100 μM Biotin-PyrDTDO for 16 h and extracts were subjected to

Streptavidin-agarose pulldowns. The affinity-purified samples were analyzed by immunoblot.

D. MDA-MB-468 cells were treated for 24 h with Biotin-PyrDTDO. Cell extracts were subjected

to Streptavidin-agarose pulldowns and the beads were washed and eluted by incubation with 1

M 2-mercaptoethanol (Elution 1) followed by elution with SDS-sample buffer containing 5 mM

biotin (Elution 2). The eluents and the flow-throughs were analyzed by immunoblot (left panel)

or probed with Streptavidin-conjugated with Alkaline Phosphatase (right panel).

E. Binding assays employing recombinant, purified ERp44 and PDIA1. Proteins were labeled

with Biotin-PyrDTDO after pretreatment with vehicle or dMtcyDTDO. Reactions were analyzed

by non-reducing SDS-PAGE, followed by detection with Streptavidin-Alkaline Phosphatase.

F. MDA-MB-468 cells were treated for 24 h with vehicle (control) or 5 μM dMtcyDTDO. Cell

extracts were resolved by non-reducing SDS-PAGE and analyzed by immunoblot with the

indicated antibodies. O and M represent disulfide bonded oligomeric and monomeric protein

forms, respectively (CC8, Cleaved Caspase 8). Red arrows denote bands stained with ERp44

or PDIA1 antibodies that change in intensity with dMtcyDTDO treatment.

Figure 4 – DDAs block ER retention of ERAP1 by ERp44

A. Model for the role of ERp44 in retrotranslocation of client proteins from the Golgi to the ER

and retention of partner proteins within the ER.

B. Immunoblot analysis of the levels of ERAP1 and MHC I in MDA-MB-468 cells after

treatment with dMtcyDTDO for 24 h.

C. Quantitative analysis of cellular ERAP1 levels in control and dMtyDTDO treated MDA-MB-

468 cells across four experiments.

D. Quantitative analysis of cellular ERp44 binding to client proteins in control and dMtyDTDO

treated MDA-MB-468 cells across four experiments.

E. The indicated stable MDA-MB-468 cell lines were treated for 24 h with dFtcyDTDO and

subjected to non-reducing SDS-PAGE followed by immunoblot. O and M represent oligomeric

and monomeric protein isoforms.

F. MDA-MB-468 (left panels) or A431 cells (right panels) were treated for 24 h with 5 μM

dMtcyDTDO and cell extracts and conditioned culture media were subjected to immunoblot

analysis as indicated. ERAP1 levels in the cell extracts and culture medium were subjected to

statistical analysis (lower panels, N=3).

Figure 5 - Evidence that disulfide bonding of DR4 and DR5 are regulated by ERp44.

A. Immunoblot analysis of the expression levels of EGFR, ERO1-α, AGR2, AGR3, PDIA1, and

ERp44 across a panel of breast cancer cell lines. Actin serves as a loading control.

B. BT474 cells stably expressing ERp44 or transduced with the vector control were treated for

24 h with vehicle, 2.5 μM, or 5 μM dMtcyDTDO. Cell extracts were resolved by SDS-PAGE

under non-reducing (left panel) or reducing (right panel) conditions and analyzed by

immunoblot. O and M are disulfide bonded oligomeric and monomeric protein forms,

respectively. Red arrows highlight dMtcyDTDO-induced changes in DR4 and DR5

oligomerization.

C. Vector control or ERp44 knockdown MDA-MB-468 cell lines were treated for 24 h with

vehicle, 2.5 μM, or 5 μM dMtcyDTDO. Cell extracts were separated by non-reducing SDS-

PAGE and analyzed by immunoblot (CC8, Cleaved Caspase 8). O and M are disulfide bonded

oligomeric and monomeric protein bands, respectively.

D. Expanded portions of the blots from panel C showing high molecular mass bands at the

stacking gel/resolving gel interface (SGI). Asterisks denote bands at the SGI that are more

prominent in the ERp44 knockdown cells in the absence of dMtcyDTDO treatment.

Figure 6 – Expression of ERp44 thioredoxin-like repeat mutants alters DDA responses.

A. Non-reducing immunoblot analysis of vector control MDA-MB-468 cells, or cells

overexpressing wild type ERp44, ERp44[C58S], or ERp44[S61C]. O and M are disulfide

bonded oligomeric and monomeric protein bands, respectively. M (FL) and M (Cl.) represent

the monomeric full-length and cleaved forms of CDCP1, respectively.

B. Non-reducing immunoblot analysis of the samples from Fig. 4E with the indicated

antibodies. O and M are disulfide bonded oligomeric and monomeric protein bands,

respectively.

C, left panel. Extracts from vector control or EGFR overexpressing T47D cells were analyzed

by non-reducing SDS-PAGE followed by immunoblot. O and M are oligomeric and monomeric

protein bands, respectively. Two replicate experiments are shown involving independent

isolation of the T47D vector control and EGFR overexpressing lines. C, right panel.

Quantitation of the percentage of PDIA1 and ERp44 present in oligomeric or monomeric forms

as assessed by immunoblot and densitometry analysis. Graphic representation of the results

of three experiments using independent isolates of T47D/Vector and T47D/EGFR lines

examining levels of monomeric and oligomeric PDIA1 and ERp44. Results are shown as the

mean ± standard error.

D. MDA-MB-468 cells were treated for 24 h with vehicle or 2.5 μM dMtcyDTDO. The cells were

then subjected to cell surface protein labeling using membrane-impermeable amine- or thiol-

reactive biotinylation probes as described in Supplemental Information. Surface proteins were

collected using Streptavidin-coated beads and analyzed along with the flow-though (internal,

non-biotinylated proteins) by immunoblot.

Figure 7 – Scission of DR5 disulfide bonds triggers high-level DR5 expression,

oligomerization, and Caspase 8 activation.

A. MDA-MB-468 cells were treated for 24 h with 10 μM dMtcyDTDO or vehicle and cell

extracts were immunoprecipitated with ERp44 antibody and analyzed by immunoblot under

non-reducing conditions.

1001

1002 B. MDA-MB-468 cells ectopically expressing wild type ERp44 or the C58S or S61C ERp44

1003 mutants were treated with 10 μM dMtcyDTDO or vehicle. Left panels, cell extracts were

1004 immunoprecipitated with ERp44 antibody and analyzed by immunoblot. Right panels, crude

1005 extracts were analyzed for protein expression levels and oligomerization by immunoblot. O and

1006 M represent disulfide bonded oligomeric and monomeric protein forms.

1007

1008 C. MDA-MB-468 expressing doxycycline-inducible DR4 and DR5 were treated with 1 μg/ml

1009 doxycycline, 10 μM dMtcyDTDO, or doxycycline + dMtcyDTDO for 24 h followed by

1010 immunoprecipitation of ERp44 and immunoblot analysis of the immunoprecipitates and crude

1011 lysates. O and M represent disulfide bonded oligomeric and monomeric protein forms.

1012

1013 D. The indicated MDA-MB-468 cell lines stably expressing wild type DR5 or the Δ81-178

1014 mutant lacking part of the disulfide-rich extracellular domain were treated for 24 h as indicated

1015 and subjected to immunoprecipitation with ERp44 antibodies. The immunoprecipitates and

1016 crude cell lysates were analyzed by non-reducing SDS-PAGE/immunoblot. O and M represent

1017 disulfide bonded oligomeric and monomeric protein forms.

1018

1019   E. The indicated stable MDA-MB-468 cell lines were treated for 24 h as indicated and

1020   subjected to immunoblot analysis after resolving proteins by non-reducing SDS-PAGE. O and

1021   M represent disulfide-bonded oligomeric and monomeric protein forms.

1022

1023   F. Model for how the biotin-labelled DDA biotin-PyrDTDO reacts covalently with the PDI active

1024   site thiolate. Covalent DDA binding is associated with ring-opening and exposure of a

1025   nucleophilic sulfinate group, which is capable of attacking the DDA-PDI disulfide bond,

1026   resulting in ring closure and DDA excision. It is hypothesized that solvation of the DDA

1027   sulfinate group by the PDI slows the rate of DDA self-excision.

1028

1029

Fig. 1, Law et al.

Fig. 2, Law et al.

Fig. 3, Law et al.

Fig. 4, Law et al.

Fig. 5, Law et al.

Fig. 6, Law et al.

Fig. 7, Law et al.