1   # Inhibitors of ERp44, PDIA1, and AGR2 induce disulfide-mediated

2   # oligomerization of Death Receptors 4 and 5 and cancer cell death

4   [1]Mary E. Law, [2]Elham Yaaghubi, [2]Amanda F. Ghilardi, [1]Bradley J. Davis, [2]Renan B. Ferreira,

5   [3]Jin Koh, [3,4]Sixue Chen, [2]Sadie F. DePeter, [2]Christopher M. Schilson, [5]Chi-Wu Chiang, [6,7]Coy

6   D. Heldermon, [8]Peter Nørgaard, [2,7]Ronald K. Castellano[#], and [1,7]Brian K. Law[#]

Affiliations

[1]Department of Pharmacology & Therapeutics, University of Florida, Gainesville, FL 32610

USA.

[2]Department of Chemistry, University of Florida, Gainesville, FL 32611 USA.

[3]Proteomics and Mass Spectrometry Facility, Interdisciplinary Center for Biotechnology

Research, University of Florida, Gainesville, FL 32610 USA.

[4]Department of Biology, Genetics Institute, University of Florida, Gainesville, FL 32610 USA

[5]Institute of Molecular Medicine, College of Medicine and Center

for Infectious Disease and Signaling Research, National Cheng

Kung University, Tainan, Taiwan.

[6]Department of Medicine, University of Florida, Gainesville, FL 32610 USA.

[7]UF-Health Cancer Center, University of Florida, Gainesville, FL 32610 USA.

[8]Department of Pathology, Copenhagen University Hospital Herlev, DK-2730 Herlev,

Denmark.

[#]Co-Contributing Corresponding Authors

Addresses of Corresponding Authors

Brian K. Law

1200 Newell Drive

Room R5-210 Academic Research Building

University of Florida

P.O. Box 100267

Gainesville, FL 32610

Email: bklaw@ufl.edu Phone: (352) 273-9423

Ronald K. Castellano

Department of Chemistry

University of Florida

P.O. Box 117200

Gainesville, FL 32611

Email: castellano@chem.ufl.edu Phone: (352) 392-2752

Conflict of Interest

The Authors have no conflicts of interest to declare.

Running Title: Molecular Targets of DDAs

Abstract

Breast cancer mortality remains unacceptably high, indicating a need for safer and more

effective therapeutic agents. Disulfide bond Disrupting Agents (DDAs) were previously

identified as a novel class of anticancer compounds that selectively kill cancers that

overexpress the Epidermal Growth Factor Receptor (EGFR) or its family member HER2. DDAs

kill EGFR+ and HER2+ cancer cells via the parallel downregulation of EGFR, HER2, and

HER3 and activation/oligomerization of Death Receptors 4 and 5 (DR4/5). However, the

mechanisms by which DDAs mediate these effects are unknown. Affinity purification analyses

employing biotinylated-DDAs reveal that the Protein Disulfide Isomerase (PDI) family members

AGR2, PDIA1, and ERp44 are DDA target proteins. Further analyses demonstrate that

shRNA-mediated knockdown of AGR2 and ERp44, or expression of ERp44 mutants, alter

basal and DDA-induced DR5 oligomerization. DDA treatment of breast cancer cells disrupts

PDIA1 and ERp44 mixed disulfide bonds with their client proteins. Together, these results

reveal DDAs as the first small molecule, active site inhibitors of AGR2 and ERp44, and

demonstrate roles for AGR2 and ERp44 in regulating the activity, stability, and localization of

DR4 and DR5, and activation of Caspase 8.

Key Words: Apoptosis, Endoplasmic Reticulum Stress, Cancer, Disulfide bond Disrupting

Agents, DR5, DR4, EGFR, AGR2, ERp44, PDIA1, ERAP1

Introduction

In spite of significant improvements in early detection, and new molecularly targeted therapeutics, breast cancer remains the second largest cancer killer of women in America after lung cancer, and is the most frequently diagnosed cancer in American women [1]. Much of the lethality of breast cancer derives from our inability to successfully treat patients with metastatic and/or drug-resistant malignancies. New drugs are needed that bypass resistance to currently used regimens by targeting cancer cell-specific vulnerabilities. A class of agents termed Disulfide bond Disrupting Agents (DDAs) was previously identified and shown to kill breast cancer cells in association with downregulation of the HER-family proteins EGFR, HER2, and HER3, and decreased activating phosphorylation of the Akt kinase [2]. Subsequent work showed that DDAs induce endoplasmic reticulum stress (ERS) [3] and alter the pattern of disulfide bonding of EGFR and the TRAIL receptors, Death Receptor 4 (DR4) and Death Receptor 5 (DR5) [4]. DDA-induced cell death is mediated by activation of the Caspase 8-Caspase 3 extrinsic apoptotic cascade.

Little is known regarding how the disulfide bonding of HER-family and Death Receptor-family proteins are chaperoned within the endoplasmic reticulum, and how the disulfide bonding patterns of these proteins control their stability, intracellular localization, and downstream signaling. TNF-Related Apoptosis Inducing Ligand (TRAIL) has been considered a promising cancer therapeutic agent due to its selectivity for killing cancer cells, while leaving normal tissues unharmed [5-8]. However, TRAIL analogs and DR4/5 agonistic antibodies have yet to be clinically approved for anticancer therapy. This is due in part to pharmacokinetic issues related to these protein drugs themselves, and in part to the ability of cancer cells to acquire resistance through a variety of mechanisms, including downregulation of DR4 and

DR5 (reviewed in [9]). Since DDAs upregulate DR5 and activate DR4 and DR5 in a ligand-

independent manner [4, 10], DDAs may overcome the resistance mechanisms that suppress

the efficacy of other activators of the TRAIL/Death Receptor pathway. A recent report showed

that the extracellular domain of DR5 prevents oligomerization and downstream activation of

Caspase 8 in the absence of ligand engagement, and that excision of the DR5 extracellular

domain permits full activation of DR5 in a ligand-independent manner [11]. It was previously

proposed that DDAs activate DR5 in a ligand-independent manner by permitting inter-

molecular disulfide bond formation at the expense of intramolecular disulfide bonding, resulting

in DR5 oligomerization and activation [4, 10]. However, the precise mechanisms by which this

happens are unknown. Similarly, DDA treatment causes disulfide bond-dependent EGFR

oligomerization followed by degradation [2, 4], but how DDAs perturb EGFR disulfide bonding

is unknown. The Protein Disulfide Isomerase (PDI) family member AGR2 is essential for EGFR

folding and surface localization [12], and likewise facilitates expression of Mucins [13, 14].

The PDI family contains 21 members (reviewed in [15]). Canonical PDIs are considered

promising targets for anticancer therapeutics [16] and significant attention has focused on the

roles of canonical PDIs in mediating disulfide bond formation via their active site thioredoxin-

like CXXC motifs. However, other PDI family members are less well studied. The PDIs AGR2

and AGR3 contain a non-canonical CXXS active site motif. Based on biochemical PDI

mutagenesis studies [17] it is predicted that AGR2/3 will exhibit disulfide isomerase activity, but

not oxidase or reductase activity. Consistent with this, AGR2 forms stable mixed disulfide

bonds with its client proteins [12, 13]. Another PDI family member, ERp44, also contains an

active site CXXS motif, and has multiple functions within the endoplasmic reticulum (ER).

ERp44 plays a critical role in the disulfide-mediated oligomerization of Adiponectin [18, 19]

and Immunoglobulin M [20, 21]. Additionally, ERp44 anchors client/partner proteins such as

ERO1α, Prx4, FGE/SUMF1, and ER Aminopeptidase 1 (ERAP1) [22-25] that do not possess

C-terminal -KDEL ER retention sequences to prevent their secretion (reviewed in [26]).

HER-family proteins are the only sub-family of receptor tyrosine kinases that contain

two cysteine-rich repeats in their extracellular domain [27]. DR4 and DR5 also harbor

numerous disulfide bonds in their extracellular domains [28]. Based on these observations, we

previously hypothesized that HER-family proteins and DR4/5 may be particularly sensitive to

agents that impede native disulfide bond formation. The mechanisms by which DDAs cause

downregulation of EGFR, HER2, and HER3, and activate DR4/5 are unknown. Here we show

using novel biotinylated DDA derivatives that DDAs covalently modify ERp44, PDIA1, and

AGR2 both in vitro and in intact cells, and that DDA treatment of cells disrupts PDIA1 and

ERp44 disulfide bonds with their client proteins, and disulfide-mediated AGR2 dimerization.

Together, the results indicate that DDAs activate DR4/5 by inhibiting the activity of the PDI

family members ERp44, PDIA1, and AGR2. DDAs are the first agents to our knowledge that

function in part by modifying the active site Cys residues of ERp44 and AGR2.

Results

Characterization of novel DDAs with enhanced potency: Previously described DDAs, including RBF3, DTDO, and tcyDTDO (Fig. 1A), were demonstrated to suppress tumor growth through induction of apoptotic cancer cell death [4, 29]. As shown in our previous studies, modifications of the parent cyclic DDA, DTDO, resulted in a more potent DDA, tcyDTDO, with improved pharmacodynamic and pharmacokinetic properties [4, 10, 29]. With tcyDTDO in hand, we then explored the effect of substitutions on its cyclohexane ring to determine if its potency and drug-like properties could be further improved. Our efforts led to the design and synthesis of novel DDAs including dimethoxy-tcyDTDO (dMtcyDTDO) and difluoro-tcyDTDO (dFtcyDTDO) (Fig. 1B). Well appreciated, fluorine's size and high electronegativity lead to strong C–F bonds that resist metabolic degradation making fluorine substitution an effective way to increase drug stability. For instance, replacing hydrogen with fluorine significantly alters the rate and extent of Cytochrome P450 metabolism [30-33]. Both dMtcyDTDO and dFtcyDTDO exhibit approximately two-fold greater potency than tcyDTDO in cell viability assays (Fig. 1C). DMtcyDTDO also exhibited two-fold greater potency than tcyDTDO in DNA and protein synthesis assays (Fig. S1A). DDAs were previously shown to induce apoptotic cell death associated with ERS, disulfide bond-mediated oligomerization of Death Receptors 4 and 5 (DR4/5) and EGFR, and upregulation of DR5 [2-4, 29]. Therefore, we examined the effect of the novel DDAs on disulfide bonding of these proteins by immunoblot analysis under non-reducing conditions ([4] and Supplemental Information). When applied to cancer cells at the same concentration, dMtcyDTDO and dFtcyDTDO induced more DR4/5 oligomerization, higher DR5 levels, and greater ERS (Fig. 1D, left panel). Induction of ERS by dFtcyDTDO and dMtcyDTDO, was indicated by GRP78 elevation, and disulfide-mediated oligomerization of

DR4 and EGFR, and oligomerization of DR5 was observed at 1.25 μM of each dFtcyDTDO or

dMtcyDTDO (Fig. 1D, right panel). Additional immunoblot analyses performed under reducing

conditions revealed that dMtcyDTDO is more potent than tcyDTDO in the induction of HER3

downregulation, DR5 upregulation, and ERS as indicated by elevated CHOP and XBP1s (Fig.

S1B). Time-course and concentration-response analyses of these endpoints showed that

dMtcyDTDO induced these responses as early as four hours after treatment (Fig. S1C).

Quantitation of dMtcyDTDO effects on the levels and state of oligomerization of DR4

and DR5 revealed that dMtcyDTDO increases DR4 oligomerization without affecting overall

DR4 levels, while dMtcyDTDO increases the levels of both oligomeric and monomeric forms of

DR5 (Fig. 1E). Tumor studies using the HER2+ BT474 xenograft model [2] were performed to

examine if dMtcyDTDO and dFtcyDTDO induce rapid breast cancer cell death in mice bearing

palpable breast tumors. Tumors from animals treated with the vehicle for four consecutive

days did not show evidence of tumor necrosis (Fig. 1F). In contrast, tumors were not observed

in mammary fat pads from dFtcyDTDO treated animals. Mammary glands from dMtcyDTDO

treated animals exhibited tumors that had undergone coagulation necrosis, consistent with

rapid and uniform death of the cancer cells.

AGR2 as a DDA binding protein: To understand how DDAs kill breast tumors, identification of

their direct protein targets is essential. The development of molecular probes proved crucial in

identifying DDA biological targets and linking them to the observed DDA responses. For this

purpose, we used activity-based protein profiling and standard affinity pulldown techniques

with biotin as the affinity label.

To generate DDA probes, we sought a biologically active cyclic DDA with a functional

group that enables installation of a biotin tag. This was achieved by fusing a Boc-protected

pyrrolidine ring at the 4,5-positions of the parent DTDO compound to generate (±)-

BocPyrDTDO (Fig. 2A), which is biologically active and showed a potency similar to DTDO

(Fig. S1D). The synthesis of (±)-BocPyrDTDO was accomplished by cleaving the acetyl groups

of dithioacetate (±)-**3** followed by oxidation to furnish the 1,2-dithiane 1,1-dioxide ring (Fig. 2A).

Intermediate (±)-**3** can be obtained through a few functional group interconversion steps from

diester (±)-**1** which itself could be obtained according to literature procedures. The amino

group on the pyrrolidine ring, after Boc group deprotection, enables installation of the linker

and biotin tag as described in full detail in the Supplemental Information. Biotinylated DDA

analogs, including Biotin-PyrDTDO (BPD) and Biotin-GlyPyrDTDO (BGPD), were generated in

this fashion and used as affinity tags to identify DDA bound proteins (Fig. 2A).

The protein disulfide isomerase (PDI) family member AGR2 was previously implicated

in disulfide bonding of EGFR [12], and DDAs alter EGFR disulfide bonding ([4] and Fig. 1D,

right panel). We hypothesized that AGR2 might be a target of DDAs and examined if

biotinylated DDAs bind covalently with recombinant AGR2. Biotin-GlyPyrDTDO labeled AGR2

in a concentration-dependent manner (Fig. 2B). Under these conditions, Biotin-GlyPyrDTDO

labeled monomeric, but not disulfide bonded dimeric AGR2. We further hypothesized that in

addition to EGFR, DR4 or DR5 may be AGR2 client proteins. Stable knockdown of AGR2 in

breast cancer cells increased disulfide-mediated DR5 oligomerization in the absence of DDA

treatment (Fig. 2C). AGR2 forms mixed disulfide bonds to its client proteins with Cys81 in its

thioredoxin-like repeat [13], and Cys81 is the only Cys residue present in AGR2. Therefore, we

examined if mutating Cys81 to Ser prevented DDA labeling. The C81S AGR2 mutant was not

labeled by Biotin-GlyPyrDTDO (Fig. 2D). AGR3 is the PDI paralog most closely related to

AGR2 and it was also labeled by Biotin-GlyPyrDTDO. Addition of a molar excess of

dMtcyDTDO to the labeling reactions partially prevented Biotin-GlyPyrDTDO binding to AGR2

and AGR3 (Fig. 2D), suggesting competition for binding to the same Cys residue(s). A

comparison of the ability of dMtcyDTDO and tcyDTDO to block Biotin-GlyPyrDTDO binding to

AGR2 indicated that dMtcyDTDO was a more effective competitor, indicating that competition

in this binding assay reflects DDA potency (Fig. 2E, S1E, S1F). AGR2 is expressed at low

levels in MDA-MB-468 cells, and disulfide-bonded AGR2 complexes with client proteins were

not detected in either the presence or absence of dMtcyDTDO treatment (Fig. 2F). Enforced

AGR2 expression revealed formation of dimeric AGR2 that was not observed with the C81S

AGR2 active site mutant. Treatment of the cells with dMtcyDTDO blocked dimer formation by

wild type AGR2, consistent with blockade of C81S by dMtcyDTDO binding.

Identification of PDIA1 and ERp44 as DDA target proteins: A panel of three breast cancer cell

lines was treated with Biotin-PyrDTDO to label DDA target proteins in intact cells. In addition to

affinity purification of endogenous biotinylated proteins, such as carboxylases, which use biotin

as a cofactor, two biotinylated proteins were observed exclusively in the context of cell labeling

with Biotin-PyrDTDO in multiple experiments and termed DDA target proteins 1 and 2

(DDAT1/2). DDAT1/2 migrated at 60 kDa and 44 kDa, respectively, and were observed in

three different breast cancer lines in crude cell extracts and in samples purified using

Streptavidin-agarose (Fig. 3A). The Streptavidin-agarose purification was scaled up to

generate DDAT1/2 in sufficient amounts for identification by mass spectrometry, and aliquots

of this preparation and the corresponding crude extracts were analyzed by blotting with

Streptavidin-Alkaline Phosphatase, and by silver staining (Fig. 3B). DDAT1 and DDAT2 from a

Coomassie stained gel were excised and analyzed by trypsin digestion followed by mass

spectrometry as described in Materials and Methods. DDAT1 and DDAT2 were identified as

the PDI-family members PDIA1 and ERp44, respectively (Fig. S2A, S2B). In support of these

assignments, ERp44, PDIA1, and AGR2 were isolated from the luminal A T47D and luminal B

BT474 breast cancer cell lines treated with Biotin-PyrDTDO using Streptavidin-Agarose (Fig.

3C, left panel). In additional Biotin-PyrDTDO pulldown experiments, ERp44 and PDIA1 were

detected, but the PDIs ERp57 and ERp5 were not (Fig. 3D). Endogenous proteins are

biotinylated via amide linkages. Thus, DDAT1 and DDAT2 were distinguished from

endogenous biotinylated proteins by disruption of Biotin-DDA/target protein linkages with the

reducing agent 2-Mercaptoethanol (Fig. S2C). Immunoprecipitation with an AGR2 antibody

showed that ERp44 and PDIA1 co-purify with AGR2 from cells irrespective of treatment with

Biotin-PyrDTDO (Fig. 3C, left panel). ERp44 immunoprecipitates contained PDIA1, but AGR2

co-purification was not detected. Biotin-PyrDTDO/Streptavidin-Agarose also affinity purified

ERp44 and PDIA1 from the MDA-MB-468 Triple-Negative breast cancer cell line (Fig. 3C, right

panel). In vitro binding studies showed that Biotin-PyrDTDO labeled ERp44 and PDIA1, and in

both cases, biotinylation was blocked by pre-incubation with a molar excess of dMtcyDTDO

(Fig. 3E). If ERp44 and PDIA1 are cellular DDA targets, then DDA treatment may alter the

pattern of ERp44 and PDIA1 disulfide bonding with their client proteins. Consistent with this,

treatment of breast cancer cells with dMtcyDTDO reduced disulfide bonding of ERp44 and

PDIA1 with multiple proteins (Fig. 3F, red arrows) with minimal effect on monomeric levels of

ERp44 and PDIA1. PDIA1 is a validated target for anticancer drugs [16]. However, ERp44 has

not been investigated extensively as a therapeutic target for cancer, and small molecule

ERp44 inhibitors have not been reported. Small molecule active site AGR2 inhibitors have not

been reported, but under the conditions used here, AGR2 binding to client proteins was not

detected. Therefore we focused most of our efforts on examining the consequences of ERp44

inhibition by DDAs.

DDAs block the ER retention function of ERp44: ERp44 carries out two important functions by

disulfide bonding with client proteins and cycling between the Golgi and the ER as modeled in

Fig. 4A. First, ERp44 retrotranslocates inappropriately oligomerized proteins from the Golgi to

ER for additional rounds of disulfide bond formation and shuffling. Second, ERp44 disulfide

bonds with select partner proteins to retain them in the ER/Golgi, preventing their secretion. To

determine if DDAs block ERp44 retention of partners by binding to the ERp44 active site,

MDA-MB-468 cells were treated with dMtcyDTDO for 24 h and the levels of the ERp44 partner

ERAP1 were examined. DMtcyDTDO treatment decreased cellular ERAP1 levels (Fig. 4B).

Quantitation of dMtcyDTDO effects on ERAP1 levels and dMtcyDTDO binding to its client

proteins across multiple experiments showed that dMtcyDTDO decreased ERAP1 levels and

ERp44 binding to client proteins in a statistically significant manner (Fig. 4C, D). ERAP1

migrates primarily as a monomer even under nonreducing conditions, indicating that ERAP1

disulfide bonding with ERp44 is transient. Enforced expression of wild type ERp44 (CXXS)

resulted in the formation of a higher molecular mass band recognized by ERAP1 antibodies

(Fig. 4E). This band was not observed when the C58S ERp44 active site mutant (SXXS) was

expressed, but was observed when the S61C mutant (CXXC) was expressed. The ERAP1-

ERp44 disulfide-bonded dimer was not observed when analyses were performed under

reducing conditions (not shown) and was decreased if the cells were treated with the DDA

dFtcyDTDO for 24h (Fig. 4E).

DDA-induced secretion of ERAP1 was examined by treating MDA-MB-468 breast

cancer cells or A431 skin cancer cells with dMtcyDTDO for 24 h and examining ERAP1 levels

in cell extracts and in the culture medium. ERAP1 was only observed in the medium upon

dMtcyDTDO treatment (Fig. 4F, upper panels), and dMtcyDTDO induced ERAP1 secretion in a

statistically significant manner (Fig. 4F, lower panels).

A role for ERp44 in the control of Death Receptor and EGFR disulfide bonding: Immunoblot

analysis of a panel of breast cancer cell lines revealed that PDIA1 levels were similar among

the lines (Fig. 5A). In contrast, ERp44, AGR2, and AGR3 levels varied among the lines, as did

the oxidoreductase ERO1, which plays a key role in producing oxidizing equivalents for

disulfide bond formation [16, 34-36]. A BT474 breast cancer cell line stably overexpressing

ERp44 was generated to examine the effect of elevated ERp44 levels on DR4/5 disulfide

bonding and DDA responses. ERp44 overexpression reduced DDA induction of high molecular

mass DR4 oligomers, and to a lesser extent high molecular mass DR5 oligomers (Fig. 5B, left

panel, red arrows). In both cases, the overall pattern was a shift from high to lower molecular

mass disulfide-bonded species. These oligomeric protein forms were not observed when the

analyses were performed under reducing conditions (Fig. 5B, right panel). Next, ERp44

knockdown MDA-MB-468 cells were generated and the effect on DR4/5 disulfide bonding was

examined. Although we were only able to achieve modest ERp44 knockdown in multiple

independent experiments, ERp44 knockdown consistently increased high molecular mass

oligomers of DR4, DR5, and EGFR in the absence of DDA treatment (Fig. 5C, red arrows;

expanded view in Fig. 5D, red asterisks; Fig. S2D). ERp44 knockdown augmented PDIA1

oligomerization and increased dMtcyDTDO induction of Caspase 8 cleavage/activation (CC8).

Together, the results of the AGR2 and ERp44 knockdown experiments are consistent with

roles for AGR2 and ERp44 in facilitating native disulfide bond formation of DR4 and DR5, and

for DDAs in blocking these AGR2 and ERp44 functions. The observation that ERp44

knockdown increases PDIA1 binding to clients may indicate that PDIA1 partially compensates

for decreased ERp44 function.

294   We next examined if mutations within the thioredoxin-like repeat of ERp44 alter DR5

levels and disulfide bonding, and other DDA responses. Overexpressed wild type ERp44

exhibited a high level of disulfide bond-mediated oligomer formation with client proteins (Fig.

6A). Both endogenous and overexpressed ERp44 monomers migrated as a doublet, which

may result from differential O-linked glycosylation [37]. The C58S ERp44 mutant (SXXS),

which lacks a Cys residue in its thioredoxin-like repeat, showed lower binding to client proteins,

although the levels of the monomer were similar to that of the wild type protein. The S61C

ERp44 mutant (CXXC) exhibited slightly increased oligomerization compared with wild type

ERp44. We previously showed that DDAs alter the electrophoretic mobility of EGFR and CUB

Domain-Containing Protein 1 (CDCP1), block the formation of complexes between these two

proteins, and alter their tyrosine phosphorylation patterns [38]. Overexpression of both ERp44

point mutants was associated with increased oligomerization of EGFR and CDCP1, but did not

alter the ratio between the full length (FL) and cleaved (Cl.) forms of CDCP1. Overexpressed

ERp44 exhibited disulfide bonding to client proteins and these interactions were partially

blocked by dFtcyDTDO treatment (Fig. 6B, e.g., bands marked by asterisks). As observed in

Fig. 6A, the C58S mutant showed decreased binding to client proteins. Interestingly, the S61C

ERp44 mutant bound client proteins similarly to the wild type protein, but was less sensitive to

disruption of binding to its clients by dFtcyDTDO. ERp44 overexpression had little effect on

DR4 levels with or without dFtcyDTDO treatment. In contrast, overexpression of either the

C58S or S61C ERp44 mutants modestly increased steady-state DR5 expression and

potentiated DR5 upregulation by dFtcyDTDO. Overexpression of ERp44 and its mutants

increased PDIA1 levels and binding to its client proteins. Consistent with the C58S ERp44

mutant inducing the strongest potentiation of DR5 upregulation by dFtcyDTDO, expression of

this mutant was also associated with increased expression of the ERS marker GRP78 (Fig.

6B). Overall, these observations indicate that ERp44^{C58S}, and to a lesser extent, ERp44^{S61C},

interfere with disulfide bond formation of ERp44 clients, resulting in higher baseline ERS and

potentiating DDA effects on DR4 and DR5. However, these observations do not explain why

cancer cells that overexpress EGFR or HER2 have heightened DDA sensitivity [2, 3]. T47D

breast cancer cells express low levels of EGFR, HER2, and HER3 and were previously used

as a model system to study the effects of EGFR overexpression on DDA responses [2, 3, 39].

Examination of ERp44 and PDIA1 disulfide bonding in control and EGFR overexpressing T47D

cells showed that EGFR overexpression resulted in recruitment of ERp44 and PDIA1, and to a

lesser extent DR4/5, into disulfide-bonded protein complexes (Fig. 6C). Under these

conditions, levels of monomeric ERp44 and PDIA1 are decreased. Oligomeric, disulfide-

bonded forms of ERp44 and PDIA1 are expected to be inaccessible for binding to newly

produced client proteins, while the monomeric forms of ERp44 and PDIA1 are accessible to

facilitate native disulfide bonding of their client proteins. Thus, EGFR may increase DDA

sensitivity by reducing the pools of monomeric ERp44 and PDIA1 that DDAs must inhibit in

order to block native disulfide bond formation of ERp44 and PDIA1 client proteins.

ERp44 mediates quality control of Adiponectin [18] and Immunoglobulin M [21]

maturation by shuttling inappropriately disulfide bonded protein forms from the Golgi back into

the ER, while permitting secretion of correctly disulfide bonded proteins. We hypothesized that

ERp44, PDIA1, or AGR2 disulfide bonding with DR4 or DR5 may sequester DR4 and DR5 in

the ER or Golgi, preventing their trafficking to the cell surface. Therefore, we hypothesized that

by blocking active site ERp44, PDIA1, and AGR2 Cys residues, DDA treatment may permit

DR4 and DR5 trafficking to the cell surface. Cell labeling using membrane impermeable,

protein-reactive probes is useful for confirming the presence of receptors at the cell surface.

Surface biotinylation studies performed on cells after treatment with dMtcyDTDO showed

increased cell surface levels of DR4 and DR5, irrespective of whether labeling was performed

with amine- or thiol-reactive cell-impermeable biotinylation probes (Fig. 6D). This increased

accumulation of DR4/5 at the cell surface may explain previous findings showing that DDAs

synergize with the DR4/5 ligand TRAIL to induce cancer cell apoptosis [4].

DDAs potentiate non-covalent binding of DR5 to ERp44: Data in Fig. 5 and 6 showing that

ERp44 knockdown and overexpression of ERp44 mutants alter DR5 levels or oligomerization

suggested that DR5 may be an ERp44 client protein. We reasoned that if DR5 is an ERp44

client, the two proteins may bind to each other. ERp44 co-immunoprecipitation experiments

showed that ERp44 associates with DR5 and to a lesser extent DR4 in MDA-MB-468 cells,

and these interactions were potentiated by DDA treatment (Fig. 7A; Fig. S2E). These were

non-covalent ERp44/DR5 complexes because although the analyses were performed under

non-reducing conditions, each protein migrated as a monomer. ERp44 co-immunoprecipitation

studies were next carried out in cells ectopically expressing ERp44 or the C58S or S61C

ERp44 mutants. DR5 immunoprecipitated with wild type ERp44 and the S61C ERp44 mutant,

but not the C58S ERp44 mutant (Fig. 7B, left panel). DDA treatment increased DR5 and

decreased PDIA1 association with ERp44, which may be partly explained by changes in

PDIA1 and ERp44 expression levels (Fig. 7B, right panel). Since DDAs increase DR5 levels,

which could account for the increased DR5 binding to ERp44, we used the tet-ON system to

drive high-level DR4 or DR5 expression. Doxycycline-induced DR4 expression increased DR4

binding to ERp44 and binding was further enhanced by DDA treatment (Fig. 7C). Doxycycline-

induced DR5 expression caused detectable DR5 binding to ERp44, which was strongly

enhanced by DDA treatment. The results in Fig. 7A-C demonstrate that DR5 forms a non-

covalent complex with ERp44 and that formation of this complex is strongly enhanced by DDA

treatment. To determine if ERp44 in dMtcyDTDO-treated cells associates with the disulfide

bond-rich extracellular domain of DR5, ERp44 immunoprecipitation experiments were

performed in cells that inducibly express wild type DR5 or the Δ81-178 DR5 deletion construct

lacking part of the disulfide-rich region. The results indicate that while wild type DR5 bound

ERp44 in the context of dMtcyDTDO treatment, ERp44 binding to DR5[Δ81-178] was barely

detectable (Fig. 7D). This result may indicate that DDA binding to the ERp44 active site Cys

residue not only inactivates its catalytic function, but increases its association with the

disulfide-rich region of DR5. This may, in turn, prevent catalytically active forms of ERp44,

PDIA1, or AGR2 from chaperoning native DR5 disulfide bond formation.

The results presented thus far indicate that DDAs alter the disulfide bonding status of

DR4 and DR5, and that this is associated with inhibition of ERp44 and PDIA1 binding to their

client proteins. However, it is unclear if perturbation of DR5 disulfide bonding mimics the

effects of DDAs, which include upregulation of DR5 protein levels and increased pro-apoptotic

signaling. This issue was addressed by inducibly expressing DR5 mutants in which a Cys

residue was mutated to Ser to disrupt one of the seven DR5 disulfide bonds. The C119S

mutant was constructed to disrupt the center/fourth disulfide bond, and the C160S mutant was

constructed to disrupt the seventh, most C-terminal DR5 disulfide bond. Induction of ectopic

wild type DR5 expression with low concentrations of doxycycline (e.g. 100 ng/ml) weakly

increased DR5 expression (Fig. 7E). However, dMtcyDTDO/doxycycline co-treatment further

increased DR5 expression, demonstrating DR5 stabilization by DDAs. The C119S and C160S

DR5 mutants were highly expressed in both the monomeric and oligomeric forms, and this was

associated with Caspase 8 cleavage/activation. DMtcyDTDO treatment modestly increased

Caspase 8 cleavage, but did not further increase expression of the C119S or C160S DR5

mutants. This suggests that disruption of individual DR5 disulfide bonds may be sufficient to

promote high-level DR5 protein expression, oligomerization, and pro-apoptotic signaling.

Discussion

Models for DDA mechanisms of action and compound selectivity: The findings presented here

suggest that DDAs bind to the PDI-family proteins ERp44, PDIA1, AGR2, and AGR3 and block

their ability to chaperone the disulfide bonding of subsets of client proteins. The resulting

alteration of disulfide bonding causes upregulation of DR5, oligomerization of DR4 and DR5,

and accumulation of DR4 and DR5 at the cell surface. Oligomerization of DR4 and DR5

causes Caspase 8 activation, which in turn activates Caspase 3 to execute apoptosis.

Caspase activation plays a key role in downregulating EGFR and HER2, and decreasing Akt

phosphorylation because these DDA responses are largely ablated by a pan-Caspase inhibitor

[4]. To our knowledge, DDAs are the first small molecule agents that block the active site Cys

residues of ERp44 and AGR2/3. Further, DDAs are the first experimental anticancer agents

that trigger DR4 and DR5-mediated Caspase 8 activation by inducing disulfide bond-mediated

DR4/5 clustering in the absence of their ligand, TRAIL. A recent report showed that the

extracellular domain of DR5 acts to prevent its oligomerization in the absence of TRAIL [11].

DDAs may overcome this autoinhibition by altering the conformations of the DR4/5

extracellular domains or favoring intermolecular disulfide bonding between DR4/5 extracellular

domains over intramolecular disulfide bonding. Consistent with these interpretations,

mutational disruption of individual DR5 disulfide bonds is sufficient to trigger its upregulation,

oligomerization, and downstream activation of Caspase 8 (Fig. 7E).

Previous reports showed that overexpression of EGFR amplifies multiple DDA responses,

including ERS induction, DR5 upregulation, HER1-3 downregulation, and reduction of Akt

phosphorylation [2-4]. Discovery of ERp44 and PDIA1 as DDA target proteins led to the

observation that EGFR overexpression reduced the levels of "free", monomeric ERp44 and

PDIA1 accessible for facilitating native disulfide bonding of client proteins. Since DDAs

covalently modify monomeric ERp44 and PDIA1 (Fig. 3), these observations explain the

molecular basis for the sensitivity of EGFR-overexpressing cells to DDAs. Given the

importance of EGFR as a potential oncogenic driver in a number of cancers, including

glioblastoma [40, 41] and Triple-Negative Breast Cancer [42-44], DDAs may have utility

against these malignancies.

Because of the relative structural simplicity of DDA molecules, the mechanisms

responsible for their target protein selectivity are of interest. PDIA1 has been investigated as a

target for anticancer therapeutics, and PDIA1 inhibitors have established anticancer efficacy in

preclinical models [45, 46]. AGR2 contributes to tumor progression by facilitating EGFR

disulfide bonding and surface presentation [47]. However, AGR2 may also be secreted and

promote tumor angiogenesis [48, 49] and act on cell surface receptors [50]. Monoclonal

antibodies designed to block extracellular AGR2 functions are under development [50, 51], but

agents that ablate the essential role of AGR2 in the folding of client proteins are not available.

The canonical thioredoxin-like catalytic site motif CXXC is featured in PDIA1 and

several other PDI enzymes (reviewed in [15]). Interestingly, AGR2, AGR3, and ERp44 share a

non-canonical CXXS thioredoxin-like repeat sequence. A report employing both yeast genetics

and biochemical enzyme assays showed that the essential function of the PDI enzyme for

yeast cell survival is catalysis of disulfide exchange or "scramblase" activity and that this could

be carried out by the CXXS mutant of PDI (PDI^{CXXS}) [17]. Enzyme assays showed that PDI^{CXXS}

could reactivate inappropriately disulfide bonded RNAse A via its scramblase activity. In

contrast, PDI^{CXXS} was unable to catalyze disulfide bond oxidation or reduction like the wild type

PDI^{CXXC} enzyme.

The C-terminal Cys residue in the CXXC motif of PDIs accelerates substrate release.

Consequently, CXXS substrate-trapping mutants of PDIs have been used to isolate mixed

disulfide bonded dimers with their client/substrate proteins [52, 53]. That the CXXS motif is

associated with slow client protein off-rate is consistent with the established roles of ERp44 in

retaining specific client/partner proteins such as ERO1 [22] and ERAP1 [25] in the ER, and in

retrotranslocation of inappropriately disulfide bonded Adiponectin [54, 55] and IgM [21, 56]

from the Golgi to the ER (see Fig. 4A). Thus, the CXXS motifs of AGR2, AGR3, and ERp44

bound to DDAs may exhibit a slow DDA off-rate for the same reason, and explain in part the

selectivity of DDAs for these enzymes. Our current working model for DDA binding to AGR2,

AGR3, and ERp44 is shown in Fig. 7F, where binding of Biotin-PyrDTDO is initiated by

nucleophilic attack of the thiosulfonate group of Biotin-PyrDTDO by the thiolate nucleophile of

the Cys residue of the CXXS motif. This results in reversible ring opening. We hypothesize that

the rate of ring closure and DDA compound excision is decreased by stabilization of the open

form of the DDA by interaction of the sulfinate group with functional groups in or near the PDI

active site. The reason for stable DDA binding to PDIAI, but not ERp57 or ERp5 is unclear

since each protein contains two canonical CGHC repeats. Further work is required to

determine which PDIA1 Cys binds DDAs and to identify structural features in the local

environment that stabilize DDA binding.

It is clear from the experiments with intact cancer cells that DDAs alter the disulfide

bonding of DR4, DR5, and EGFR and change the pattern of disulfide bonding between ERp44

and PDIA1 and their client proteins. It is difficult to compare these DDA actions with those of

published PDI inhibitors since most of these studies did not examine perturbation of protein

disulfide bonding in cancer cells, but rather examined compound activity in the Insulin

reduction/aggregation assay. While the Insulin reduction assay is valuable for optimizing the

potency of PDI inhibitors, it may not be salient in instances where disulfide isomerase activity,

ER retention, or Golgi-ER retrotranslocation are more important than disulfide oxidase or

reductase activity.

The presence of 21 different PDIs in humans and the diverse phenotypes of the

corresponding knockout mice (e.g., [25, 57-59]) suggest that individual PDIs chaperone the

disulfide bonding, ER retention, or Golgi-ER retrotranslocation of different subsets of client

proteins. For instance, ERp44 is unique in its ability to retain the aminopeptidase ERAP1 in the

ER/Golgi and this results in ERAP1 secretion in ERp44-deficient mice or cell lines [25]. A

biological consequence of this is that ERp44 deletion is associated with hypotension in mice

since secreted ERAP1 destabilizes Angiotensin II, leading to reduced blood pressure. It will be

of interest to determine if DDAs exhibit antihypertensive activity.

ERAP1 also functions as a "molecular ruler" to process peptides for loading onto MHC I

for presentation to T cells [60-62]. ERAP1 inhibitors or ERAP1 knockdown changes the

spectrum of the MHC I-associated peptides that constitute the "immunopeptidome." Since

DDAs cause ERAP1 secretion, it will be important to determine if DDAs alter the

immunopeptidome, and if so, how this alters anticancer immunity.

In summary, the results presented here establish perturbation of disulfide bonding by

inhibition of subsets of PDIs as a new strategy for activating DR5 and DR4 in cancer cells

independently of their engagement by agonistic antibodies or TRAIL. Further, DDAs

upregulate DR5 and increase DR4 and DR5 localization to the cell surface. Consequently,

DDAs may overcome many of the problems associated with TRAIL analogs and agonistic

antibodies specific for DR4 and DR5. Additional research is needed to elucidate the

mechanisms by which disrupting individual DR5 disulfide bonds increases its expression

levels, disulfide-mediated oligomerization, and activation of Caspase 8. In future studies, it will

be important to determine if C-S point mutants of DR4 mimic the corresponding DR5 mutants

and drive pro-apoptotic signaling. It will also be interesting to examine whether activated point

mutants of DR4 and DR5 exhibit the same selectivity for killing cancer cells in preference to

normal cells as their ligand, TRAIL.

Materials and Methods

In vivo tumor studies and histochemical analysis

Breast tumor formation was initiated in NOD-SCID-gamma (NSG) mice obtained from Jackson

Laboratories (Bar Harbor, ME USA) by injecting 1×10^6 cancer cells into the #4 mammary fat

pads as described previously [4]. Animals bearing HER2+ BT474 tumors [2] were treated with

the vehicle or 20 mg/kg dMtcyDTDO or dFtcyDTDO by intraperitoneal injection for five

consecutive days. Tissue samples fixed in 4% paraformaldehyde/Phosphate-Buffered Saline

(PBS) and paraffin-embedded were sectioned and stained with hematoxylin and eosin (H&E)

by the University of Florida Molecular Pathology Core (https://molecular.pathology.ufl.edu/).

Streptavidin-Agarose Pulldowns

Cells were treated with Biotin-PyrDTDO with or without the pan-Caspase inhibitor Q-VD-OPH

to prevent apoptosis (MedChemExpress, Monmouth Junction, NJ USA) for 16 h in 10% FBS–

DMEM, followed by the addition of 25 mM *N*-ethylmaleimide (NEM) (Thermo Scientific,

Rockford, IL USA) for 20 min at 37°C. Cells were subsequently scraped directly into the media,

washed twice with ice-cold PBS, and pelleted by centrifugation. Cells were extracted in buffer

(20 mM HEPES (pH 7.4), 1 mM EDTA, 1mM EGTA, 5% Glycerol, 1 nM Microcystin, 1 mM

Na_3VO_4, and 40 mM $Na_2H_2P_2O_7$) containing 1% Triton X-100. Samples were sonicated,

followed by centrifugation for 20 min at 4°C. The supernatant was pre-cleared with Sepharose

CL-6B (Amersham Biosciences AP, Uppsala, Sweden) for 2h at 4°C. Pre-cleared samples

were then incubated with Streptavidin (SA)-Agarose (Thermo Scientific, Rockford IL USA) and

mixed overnight at 4°C. The SA-Agarose was washed three times with extraction buffer

containing 0.1% Triton X-100, and eluted by boiling in 2X-SDS Sample Buffer containing 5 mM

biotin. Samples were analyzed by immunoblot, silver stain, or probing with Streptavidin-

conjugated Alkaline Phosphatase.

Disulfide Bond-mediated Oligomerization

To evaluate protein disulfide bond-mediated oligomerization, cells were treated with DDAs for

24 h in 10% FBS–DMEM. Cells were subsequently scraped into the media and pelleted by

centrifugation. Cell pellets were extracted directly into boiling 2X-SDS-Sample Buffer

containing 100 mM NEM and boiled for 10 min. Samples were sonicated, followed by

centrifugation for 20 minutes at room temperature, and subsequently analyzed by SDS-

PAGE/immunoblot.

Surface Biotinylation

Cell surface proteins were labeled by incubating cells with 1.6 mM Sulfo-NHS-SS-Biotin

(Thermo Fisher Scientific, Waltham, MA USA) or 1.6 mM Biotin-dPEG$_3$-Maleimide

(MilliporeSigma, St. Louis, MO USA) in PBS, pH 8.0 for 30 minutes at 4 °C. The cells were

subsequently washed three times with PBS, and proteins were extracted in 1% Triton X-100

extraction buffer in the absence of reducing agents. Cell extracts were then mixed with

Streptavidin-agarose beads (Thermo Fisher Scientific, Waltham, MA USA) for 2.5 h at 4°C.

Following centrifugation, the supernatant containing non-biotinylated proteins (flow-through)

was retained and boiled in 2X-SDS-Sample Buffer containing 1% 2-mercaptoethanol.

Streptavidin-agarose beads bound to biotinylated proteins were washed four times with 0.1%

Triton X-100 extraction buffer, and eluted by boiling in 2X-SDS-Sample Buffer containing 1%

2-mercaptoethanol. Samples were then analyzed by immunoblot or blotting with Streptavidin-

conjugated Alkaline Phosphatase.

His$_6$-AGR2 TALON Purification

Competent BL21(DE3) cells were transformed with either the AGR2/pET-45b(+) or AGR2

C81S/pET-45b(+) vectors. Protein expression was induced by the addition of 50 μM Isopropyl

β-D-1-thiogalactopyranoside (Fisher Scientific, Fair Lawn, NJ USA) to bacterial cell cultures for

2.5 hours and shaking at 37°C. Bacterial cells were subsequently pelleted and dissolved in

Mouse Tonicity Phosphate Buffered Saline (MTPBS) containing 1% Triton X-100 and

sonicated for 20 seconds on ice, followed by centrifugation for 10 min at 4°C to pellet cell

debris. The supernatant was transferred to a new tube, followed by the addition of TALON

Resin (Clontech, Mountain View, CA USA) and mixed for 30 min at 4°C. The TALON Resin

was pelleted by centrifugation and washed twice with MTPBS and twice with MTPBS

containing 30 mM Imidazole. Protein bound to the TALON Resin was eluted with 400 mM

Imidazole and precipitated with 60% saturated ammonium sulfate. The precipitated protein

was pelleted by centrifugation, dissolved in MTPBS containing 1 mM DTT and 1 mM EDTA,

and subsequently dialyzed against a solution of MTPBS containing 10 μM DTT and 30%

Glycerol.

DDA-AGR2 Binding Experiments

Purified His$_6$-AGR2 or His$_6$-AGR2[C81S], as prepared above, or His$_6$-AGR2 (PRO-580) or

His$_6$-AGR3 (PRO-242) obtained from ProSpec (New Brunswick NJ USA) were incubated with

or without varying concentrations of the unlabeled DDA competitors, tcyDTDO or dMtcyDTDO,

and 30 μM Biotin-GlyPyrDTDO or Biotin-PyrDTDO in buffer (20 mM HEPES (pH 7.4), 1 mM

EDTA, 1mM EGTA, 5% Glycerol, 1 nM Microcystin, 1 mM Na_3VO_4, and 40 mM $Na_2H_2P_2O_7$)

containing 1% Triton-X 100 for 30 min at 37°C. Preliminary studies showed that Biotin-

GlyPyrDTDO or Biotin-PyrDTDO exhibited indistinguishable labeling of proteins in intact cells

and purified proteins in vitro. Therefore, the two probes were used interchangeably and are

referred to collectively as "Biotin-DDA." Reactions were stopped by boiling the samples in 2X-

SDS-Sample Buffer containing 100 mM NEM for 10 min. Biotin-DDA labeling of protein was

verified by blotting with Streptavidin-Alkaline Phosphatase (Rockland Immunochemicals, Inc.,

Limerick, PA USA). ERp44 (PRO-547) and PDIA1 (ENZ-262) were obtained from ProSpec and

used in experiments examining the binding of these proteins to Biotin-PyrDTDO.

Quantitative Analysis of Immunoblot Results

Immunoblots were scanned, converted to grayscale and inverted using Adobe Photoshop

(Berkeley, CA). Bands were quantified using ImageJ (NIH, Bethesda, MD). Expression levels

were normalized using Actin as a loading control.

Chemical Synthesis of DDAs

Full experimental and characterization details associated with the synthesis of (±)-

BocPyrDTDO, Biotin-GlyPyrDTDO, and Biotin-PyrDTDO are provided in the Supplemental

Information. Detailed discussion of the syntheses and analytical characterization of

dMtcyDTDO and dFtcyDTDO will be published elsewhere.

Detailed Methods in Supplemental Information

At least three biological replicates of each of the studies presented were performed. In-depth

descriptions of approaches relating to cell culture, preparation of cell extracts, immunoblot

analysis, construction of retroviral and lentiviral vectors and their use to generate stable cell

lines, DNA and protein synthesis assays, cell viability assays, sample preparation for analysis

by tandem mass spectrometry, and statistical methods are presented in Supplemental

Information.

Acknowledgements

These studies were supported in part by grants from the Florida Breast Cancer Foundation (BL

and RC), the Ocala Royal Dames for Cancer Research (BL), and the Office of the Assistant

Secretary of Defense for Health Affairs through the Breast Cancer Research Program under

Award Nos. W81XWH-15-1-0199 (BL) and W81XWH-15-1-0200 (RC), and NIH/NCI grant

CA252400 (BL). RF is grateful to the University of Florida for a Graduate School Fellowship.

EY thanks the Dr. Howard and Brenda Sheridan Fund in Chemistry for a 2020 Howard and

Brenda Sheridan Summer Fellowship. AG is grateful to the Tarrant Organic Chemistry Fund

for a 2021 Tarrant Summer Graduate Research Fellowship. Mass spectrometric data on DDA

protein targets were obtained by the UF Proteomics and Mass Spectrometry Facility at the

Interdisciplinary Center for Biotechnology Research. Mass spectrometric data on DDA

compound structures and DDA adducts were obtained by the UF Department of Chemistry

Mass Spectrometry Research and Education Center supported, in part, by the National

Institutes of Health (NIH S10 OD021758-01A1).

Conflicts of Interest

None.

References

607   1   DeSantis CE, Ma J, Gaudet MM, Newman LA, Miller KD, Goding Sauer A *et al*. Breast
608   cancer statistics, 2019. *CA Cancer J Clin* 2019; 69: 438-451.
609

610   2   Ferreira RB, Law ME, Jahn SC, Davis BJ, Heldermon CD, Reinhard M *et al*. Novel
611   agents that downregulate EGFR, HER2, and HER3 in parallel. *Oncotarget* 2015; 6:
612   10445-10459.
613

614   3   Ferreira RB, Wang M, Law ME, Davis BJ, Bartley AN, Higgins PJ *et al*. Disulfide bond
615   disrupting agents activate the unfolded protein response in EGFR- and HER2-positive
616   breast tumor cells. *Oncotarget* 2017; 8: 28971-28989.
617

618   4   Wang M, Law ME, Davis BJ, Yaaghubi E, Ghilardi AF, Ferreira RB *et al*. Disulfide bond-
619   disrupting agents activate the tumor necrosis family-related apoptosis-inducing
620   ligand/death receptor 5 pathway. *Cell Death Discovery* 2019; 5: 153.
621

622   5   Wiley SR, Schooley K, Smolak PJ, Din WS, Huang CP, Nicholl JK *et al*. Identification
623   and characterization of a new member of the TNF family that induces apoptosis.
624   *Immunity* 1995; 3: 673-682.
625

626   6   Rieger J, Naumann U, Glaser T, Ashkenazi A, Weller M. APO2 ligand: a novel lethal
627   weapon against malignant glioma? *FEBS Lett* 1998; 427: 124-128.
628

629   7   Zamai L, Ahmad M, Bennett IM, Azzoni L, Alnemri ES, Perussia B. Natural killer (NK)
630   cell-mediated cytotoxicity: differential use of TRAIL and Fas ligand by immature and
631   mature primary human NK cells. *J Exp Med* 1998; 188: 2375-2380.
632

633   8   Ashkenazi A, Pai RC, Fong S, Leung S, Lawrence DA, Marsters SA *et al*. Safety and
634   antitumor activity of recombinant soluble Apo2 ligand. *J Clin Invest* 1999; 104: 155-162.
635

636   9   Thapa B, Kc R, Uludag H. TRAIL therapy and prospective developments for cancer
637   treatment. *J Control Release* 2020; 326: 335-349.
638

639   10 Wang M, Law ME, Davis BJ, Yaaghubi E, Ghilardi AF, Ferreira RB *et al*. Mechanistic
640   Elucidation of the Antitumor Properties of a Novel Death Receptor 5 Activator. *bioRxiv*
641   2019: 700906.
642

643   11 Pan L, Fu TM, Zhao W, Zhao L, Chen W, Qiu C *et al*. Higher-Order Clustering of the
644   Transmembrane Anchor of DR5 Drives Signaling. *Cell* 2019; 176: 1477-1489 e1414.
645

646   12 Dong A, Wodziak D, Lowe AW. Epidermal growth factor receptor (EGFR) signaling
647   requires a specific endoplasmic reticulum thioredoxin for the post-translational control of
648   receptor presentation to the cell surface. *J Biol Chem* 2015; 290: 8016-8027.
649

650   13   Park SW, Zhen G, Verhaeghe C, Nakagami Y, Nguyenvu LT, Barczak AJ *et al*. The
651   protein disulfide isomerase AGR2 is essential for production of intestinal mucus. *Proc*
652   *Natl Acad Sci U S A* 2009; 106: 6950-6955.

654   14   Schroeder BW, Verhaeghe C, Park SW, Nguyenvu LT, Huang X, Zhen G *et al*. AGR2 is
655   induced in asthma and promotes allergen-induced mucin overproduction. *Am J Respir*
656   *Cell Mol Biol* 2012; 47: 178-185.

658   15   Galligan JJ, Petersen DR. The human protein disulfide isomerase gene family. *Hum*
659   *Genomics* 2012; 6: 6.

661   16   Shergalis AG, Hu S, Bankhead A, 3rd, Neamati N. Role of the ERO1-PDI interaction in
662   oxidative protein folding and disease. *Pharmacol Ther* 2020; 210: 107525.

664   17   Laboissiere MC, Sturley SL, Raines RT. The essential function of protein-disulfide
665   isomerase is to unscramble non-native disulfide bonds. *J Biol Chem* 1995; 270: 28006-
666   28009.

668   18   Hampe L, Radjainia M, Xu C, Harris PW, Bashiri G, Goldstone DC *et al*. Regulation and
669   Quality Control of Adiponectin Assembly by Endoplasmic Reticulum Chaperone ERp44.
670   *J Biol Chem* 2015; 290: 18111-18123.

672   19   Long Q, Lei T, Feng B, Yin C, Jin D, Wu Y *et al*. Peroxisome proliferator-activated
673   receptor-gamma increases adiponectin secretion via transcriptional repression of
674   endoplasmic reticulum chaperone protein ERp44. *Endocrinology* 2010; 151: 3195-3203.

676   20   Cortini M, Sitia R. From antibodies to adiponectin: role of ERp44 in sizing and timing
677   protein secretion. *Diabetes Obes Metab* 2010; 12 Suppl 2: 39-47.

679   21   Cortini M, Sitia R. ERp44 and ERGIC-53 synergize in coupling efficiency and fidelity of
680   IgM polymerization and secretion. *Traffic* 2010; 11: 651-659.

682   22   Otsu M, Bertoli G, Fagioli C, Guerini-Rocco E, Nerini-Molteni S, Ruffato E *et al*.
683   Dynamic retention of Ero1alpha and Ero1beta in the endoplasmic reticulum by
684   interactions with PDI and ERp44. *Antioxid Redox Signal* 2006; 8: 274-282.

686   23   Yang K, Li DF, Wang X, Liang J, Sitia R, Wang CC *et al*. Crystal Structure of the
687   ERp44-Peroxiredoxin 4 Complex Reveals the Molecular Mechanisms of Thiol-Mediated
688   Protein Retention. *Structure* 2016; 24: 1755-1765.

690   24   Fraldi A, Zito E, Annunziata F, Lombardi A, Cozzolino M, Monti M *et al*. Multistep,
691   sequential control of the trafficking and function of the multiple sulfatase deficiency gene
692   product, SUMF1 by PDI, ERGIC-53 and ERp44. *Hum Mol Genet* 2008; 17: 2610-2621.

694   25   Hisatsune C, Ebisui E, Usui M, Ogawa N, Suzuki A, Mataga N *et al*. ERp44 Exerts
695   Redox-Dependent Control of Blood Pressure at the ER. *Mol Cell* 2015; 58: 1015-1027.

26   Tempio T, Anelli T. The pivotal role of ERp44 in patrolling protein secretion. *J Cell Sci*
2020; 133.

27   Lemmon MA, Schlessinger J. Cell signaling by receptor tyrosine kinases. *Cell* 2010;
141: 1117-1134.

28   Cha SS, Sung BJ, Kim YA, Song YL, Kim HJ, Kim S *et al*. Crystal structure of TRAIL-
DR5 complex identifies a critical role of the unique frame insertion in conferring
recognition specificity. *J Biol Chem* 2000; 275: 31171-31177.

29   Wang MX, Ferreira RB, Law ME, Davis BJ, Yaaghubi E, Ghilardi AF *et al*. A novel
proteotoxic combination therapy for EGFR+ and HER2+cancers. *Oncogene* 2019; 38:
4264-4282.

30   Bondi A. Van Der Waals Volumes + Radii. *Journal of Physical Chemistry* 1964; 68: 441-
+.

31   Gillis EP, Eastman KJ, Hill MD, Donnelly DJ, Meanwell NA. Applications of Fluorine in
Medicinal Chemistry. *Journal of Medicinal Chemistry* 2015; 58: 8315-8359.

32   Wang J, Sanchez-Rosello M, Acena JL, del Pozo C, Sorochinsky AE, Fustero S *et al*.
Fluorine in Pharmaceutical Industry: Fluorine-Containing Drugs Introduced to the
Market in the Last Decade (2001-2011). *Chemical Reviews* 2014; 114: 2432-2506.

33   Park BK, Kitteringham NR, O'Neill PM. Metabolism of fluorine-containing drugs. *Annual*
Review of Pharmacology and Toxicology 2001; 41: 443-470.

34   Frand AR, Kaiser CA. The ERO1 gene of yeast is required for oxidation of protein
dithiols in the endoplasmic reticulum. *Mol Cell* 1998; 1: 161-170.

35   Pollard MG, Travers KJ, Weissman JS. Ero1p: a novel and ubiquitous protein with an
essential role in oxidative protein folding in the endoplasmic reticulum. *Mol Cell* 1998; 1:
171-182.

36   Bertoli G, Simmen T, Anelli T, Molteni SN, Fesce R, Sitia R. Two conserved cysteine
triads in human Ero1alpha cooperate for efficient disulfide bond formation in the
endoplasmic reticulum. *J Biol Chem* 2004; 279: 30047-30052.

37   Sannino S, Anelli T, Cortini M, Masui S, Degano M, Fagioli C *et al*. Progressive quality
control of secretory proteins in the early secretory compartment by ERp44. *J Cell Sci*
2014; 127: 4260-4269.

38   Law ME, Ferreira RB, Davis BJ, Higgins PJ, Kim JS, Castellano RK *et al*. CUB domain-
containing protein 1 and the epidermal growth factor receptor cooperate to induce cell
detachment. *Breast Cancer Res* 2016; 18: 80.

39   Wang M, Law ME, Davis BJ, Yaaghubi E, Ghilardi AF, Ferreira RB *et al*. Disulfide bond-
disrupting agents activate the tumor necrosis family-related apoptosis-inducing
ligand/death receptor 5 pathway. *Cell Death Discov* 2019; 5: 153.
746

747   40   Stockhausen MT, Broholm H, Villingshoj M, Kirchhoff M, Gerdes T, Kristoffersen K *et al*.
748   Maintenance of EGFR and EGFRvIII expressions in an in vivo and in vitro model of
749   human glioblastoma multiforme. *Exp Cell Res* 2011; 317: 1513-1526.
750

751   41   Cho J, Pastorino S, Zeng Q, Xu X, Johnson W, Vandenberg S *et al*. Glioblastoma-
752   derived epidermal growth factor receptor carboxyl-terminal deletion mutants are
753   transforming and are sensitive to EGFR-directed therapies. *Cancer Res* 2011; 71: 7587-
754   7596.
755

756   42   Liu D, He J, Yuan Z, Wang S, Peng R, Shi Y *et al*. EGFR expression correlates with
757   decreased disease-free survival in triple-negative breast cancer: a retrospective
758   analysis based on a tissue microarray. *Med Oncol* 2012; 29: 401-405.
759

760   43   de Ruijter TC, Veeck J, de Hoon JP, van Engeland M, Tjan-Heijnen VC. Characteristics
761   of triple-negative breast cancer. *J Cancer Res Clin Oncol* 2011; 137: 183-192.
762

763   44   Toyama T, Yamashita H, Kondo N, Okuda K, Takahashi S, Sasaki H *et al*. Frequently
764   increased epidermal growth factor receptor (EGFR) copy numbers and decreased
765   BRCA1 mRNA expression in Japanese triple-negative breast cancers. *BMC Cancer*
766   2008; 8: 309.
767

768   45   Stojak M, Milczarek M, Kurpinska A, Suraj-Prazmowska J, Kaczara P, Wojnar-Lason K
769   *et al*. Protein Disulphide Isomerase A1 Is Involved in the Regulation of Breast Cancer
770   Cell Adhesion and Transmigration via Lung Microvascular Endothelial Cells. *Cancers*
771   *(Basel)* 2020; 12.
772

773   46   Xu S, Liu Y, Yang K, Wang H, Shergalis A, Kyani A *et al*. Inhibition of protein disulfide
774   isomerase in glioblastoma causes marked downregulation of DNA repair and DNA
775   damage response genes. *Theranostics* 2019; 9: 2282-2298.
776

777   47   Gupta A, Dong A, Lowe AW. AGR2 gene function requires a unique endoplasmic
778   reticulum localization motif. *J Biol Chem* 2012; 287: 4773-4782.
779

780   48   Guo H, Zhu Q, Yu X, Merugu SB, Mangukiya HB, Smith N *et al*. Tumor-secreted
781   anterior gradient-2 binds to VEGF and FGF2 and enhances their activities by promoting
782   their homodimerization. *Oncogene* 2017; 36: 5098-5109.
783

49   Jia M, Guo Y, Zhu D, Zhang N, Li L, Jiang J *et al*. Pro-metastatic activity of AGR2
interrupts angiogenesis target bevacizumab efficiency via direct interaction with VEGFA
and activation of NF-kappaB pathway. *Biochim Biophys Acta Mol Basis Dis* 2018; 1864:
1622-1633.

50   Arumugam T, Deng D, Bover L, Wang H, Logsdon CD, Ramachandran V. New Blocking
Antibodies against Novel AGR2-C4.4A Pathway Reduce Growth and Metastasis of
Pancreatic Tumors and Increase Survival in Mice. *Mol Cancer Ther* 2015; 14: 941-951.

51   Liu AY, Kanan AD, Radon TP, Shah S, Weeks ME, Foster JM *et al*. AGR2, a unique
tumor-associated antigen, is a promising candidate for antibody targeting. *Oncotarget*
2019; 10: 4276-4289.

52   Eriksson O, Stopa J, Furie B. Identification of PDI Substrates by Mechanism-Based
Kinetic Trapping. *Methods Mol Biol* 2019; 1967: 165-182.

53   Stopa JD, Baker KM, Grover SP, Flaumenhaft R, Furie B. Kinetic-based trapping by
intervening sequence variants of the active sites of protein-disulfide isomerase identifies
platelet protein substrates. *J Biol Chem* 2017; 292: 9063-9074.

54   Wang ZV, Schraw TD, Kim JY, Khan T, Rajala MW, Follenzi A *et al*. Secretion of the
adipocyte-specific secretory protein adiponectin critically depends on thiol-mediated
protein retention. *Mol Cell Biol* 2007; 27: 3716-3731.

55   Wolf G. New insights into thiol-mediated regulation of adiponectin secretion. *Nutr Rev*
2008; 66: 642-645.

56   Anelli T, Ceppi S, Bergamelli L, Cortini M, Masciarelli S, Valetti C *et al*. Sequential steps
and checkpoints in the early exocytic compartment during secretory IgM biogenesis.
EMBO J 2007; 26: 4177-4188.

57   Jang I, Pottekat A, Poothong J, Yong J, Lagunas-Acosta J, Charbono A *et al*.
PDIA1/P4HB is required for efficient proinsulin maturation and ss cell health in response
to diet induced obesity. *Elife* 2019; 8.

58   Torres M, Medinas DB, Matamala JM, Woehlbier U, Cornejo VH, Solda T *et al*. The
Protein-disulfide Isomerase ERp57 Regulates the Steady-state Levels of the Prion
Protein. *J Biol Chem* 2015; 290: 23631-23645.

59   Nyirimigabo E, Jin M, Yang Z, Wang J, Zhai K, Mao Y *et al*. The role of ERp44 in
glucose and lipid metabolism. *Arch Biochem Biophys* 2019; 671: 175-184.

60   Sui L, Gandhi A, Guo HC. Crystal structure of a polypeptide's C-terminus in complex
with the regulatory domain of ER aminopeptidase 1. *Mol Immunol* 2016; 80: 41-49.

61   Chang SC, Momburg F, Bhutani N, Goldberg AL. The ER aminopeptidase, ERAP1,
trims precursors to lengths of MHC class I peptides by a "molecular ruler" mechanism.
Proc Natl Acad Sci U S A 2005; 102: 17107-17112.

62   Koumantou D, Barnea E, Martin-Esteban A, Maben Z, Papakyriakou A, Mpakali A *et al*.
Editing the immunopeptidome of melanoma cells using a potent inhibitor of endoplasmic
reticulum aminopeptidase 1 (ERAP1). *Cancer Immunol Immunother* 2019; 68: 1245-
1261.

Figure Legends

Figure 1 – Characterization of new, more potent DDAs that induce rapid tumor regression.

A. Representative first and second generation DDAs (RBF3, DTDO, and tcyDTDO) and

illustration of interconversion between the linear and cyclic forms of the DDA pharmacophore.

B. Structures of new tcyDTDO derivatives dMtcyDTDO and dFtcyDTDO.

C. MDA-MB-468 cells were treated for 24 h with the indicated concentrations of DDAs and cell

viability (MTT) assays were performed. Three biological replicates were performed with similar

results.

D. MDA-MB-468 cells were treated for 24 h as indicated and subjected to immunoblot analysis

after non-reducing SDS-PAGE. O and M represent disulfide bonded oligomeric and

monomeric protein forms, respectively.

E. Levels and degree of oligomerization of DR4 (left panel) and DR5 (right panel) were

determined by densitometry analysis of bands across multiple experiments and subjected to

statistical analysis. Values plotted represent averages and error bars represent S.E. P-values

for pairwise comparisons were determined using Student's t-test. (N=4 for each treatment

group with similar within-group results).

F. Mice bearing BT474 cell xenograft tumors were treated once daily for five days with vehicle,

20 mg/kg dFtcyDTDO, or 20 mg/kg dMtcyDTDO. Two hours after the final treatment,

mammary fat pads/tumors were excised and tumor sections were stained with H&E. T denotes

tumor tissue, FP indicates mammary fat pads, and CN represents coagulation necrosis.

Figure 2 - AGR2 and AGR3 are DDA target proteins.

A. Abbreviated synthetic approach and chemical structures of the biotinylated DDA analogs

Biotin-PyrDTDO and Biotin-GlyPyrDTDO.

B. Purified, recombinant AGR2 was reacted with Biotin-GlyPyrDTDO for 1 h at 37°C, resolved

by non-reducing SDS-PAGE, transferred to membranes, and probed with Streptavidin-Alkaline

Phosphatase, and by anti-AGR2 immunoblot.

C. Vector control or AGR2 knockdown MDA-MB-468 cells were treated for 24 h as indicated

and analyzed by non-reducing (left panels) or reducing (right panels) immunoblot (CC8,

Cleaved Caspase 8). Red arrow indicates oligomerized forms of DR5 that are present at

higher levels in the AGR2 knockdown cells. O and M represent disulfide bonded oligomeric

and monomeric protein forms, respectively.

D. Biotin-GlyPyrDTDO binding assays performed as in Fig. 2B, employing the C81S AGR2

mutant and AGR3 in addition to AGR2. In the indicated samples, the proteins were pre-

incubated with 600 μM dMtcyDTDO to determine if this prevented subsequent Biotin-

GlyPyrDTDO binding.

E. AGR2 binding assay performed as in Fig. 2B to examine the relative abilities of pre-

incubation of AGR2 with dMtcyDTDO or tcyDTDO to block subsequent binding of Biotin-

GlyPyrDTDO.

F. The indicated MDA-MB-468 stable cell lines were treated for 24 h with 5 μM dMtcyDTDO

and subjected to analysis by non-reducing SDS-PAGE/immunoblot. O, D, and M represent the

Oligomeric, Dimeric, and Monomeric forms of AGR2 detected. P43/p41 and p18 denote

different Caspase 8 cleavage products.

Figure 3 - ERp44 and PDIA1 are additional DDA targets.

A. Breast cancer cell lines were treated for 16 h with vehicle, 100 μM Biotin-PyrDTDO, or 100

μM Biotin-PyrDTDO + 10 μM Q-VD-OPH for 16 h. Biotinylated proteins were isolated using

Streptavidin-coated beads and the crude extracts and Streptavidin pulldowns were analyzed

by blotting with Streptavidin-Alkaline Phosphatase. Red arrows indicate bands, designated

DDAT1 and DDAT2, present exclusively after Biotin-PyrDTDO treatment.

B. Biotin-PyrDTDO/Streptavidin pulldowns were performed in BT474 and MDA-MB-468 cells.

Purified material was analyzed by Streptavidin-Alkaline Phosphatase detection and silver stain.

C, left panel. The indicated breast cancer lines were treated with vehicle or 100 μM Biotin-

PyrDTDO for 16 h and extracts were subjected to Streptavidin-agarose pulldowns, or anti-

AGR2 or anti-ERp44 immunoprecipitations. The purified material was analyzed by immunoblot

with the indicated antibodies. C, right panel. BT474 and MDA-MB-468 breast cancer lines were

treated with vehicle or 100 μM Biotin-PyrDTDO for 16 h and extracts were subjected to

Streptavidin-agarose pulldowns. The affinity-purified samples were analyzed by immunoblot.

D. MDA-MB-468 cells were treated for 24 h with Biotin-PyrDTDO. Cell extracts were subjected

to Streptavidin-agarose pulldowns and the beads were washed and eluted by incubation with 1

M 2-mercaptoethanol (Elution 1) followed by elution with SDS-sample buffer containing 5 mM

biotin (Elution 2). The eluents and the flow-throughs were analyzed by immunoblot (left panel)

and probed with Streptavidin-conjugated with Alkaline Phosphatase (right panel).

E. Binding assays employing recombinant, purified ERp44 and PDIA1. Proteins were labeled

with Biotin-PyrDTDO after pretreatment with vehicle or dMtcyDTDO. Reactions were analyzed

by non-reducing SDS-PAGE followed by detection with Streptavidin-Alkaline Phosphatase.

F. MDA-MB-468 cells were treated or 24 h with vehicle (control) or 5 μM dMtcyDTDO. Cell

extracts were resolved by non-reducing SDS-PAGE and analyzed by immunoblot with the

indicated antibodies. O and M represent disulfide bonded oligomeric and monomeric protein

forms, respectively (CC8, Cleaved Caspase 8). Red arrows denote bands stained with ERp44

or PDIA1 antibodies that change in intensity with dMtcyDTDO treatment.

Figure 4 – DDAs block ER retention of ERAP1 by ERp44

A. Model for the role of ERp44 in retrotranslocation of client proteins from the Golgi to the ER

and retention of partner proteins within the ER.

B. Immunoblot analysis of the levels of ERAP1 and MHC I in MDA-MB-468 cells after

treatment with dMtcyDTDO for 24 h.

C. Quantitative analysis of cellular ERAP1 levels in control and dMtyDTDO treated MDA-MB-

468 cells across multiple experiments (N=4).

D. Quantitative analysis of cellular ERp44 binding to client proteins in control and dMtyDTDO

treated MDA-MB-468 cells across multiple experiments (N=4).

E. The indicated stable MDA-MB-468 cell lines were treated for 24 h with dFtcyDTDO and

subjected to non-reducing SDS-PAGE followed by immunoblot. O and M represent Oligomeric

and Monomeric protein isoforms.

F. MDA-MB-468 (left panels) or A431 cells (right panels) were treated for 24 h with 5 μM

dMtcyDTDO and cell extracts and conditioned culture media were subjected to immunoblot

analysis as indicated (upper panels). ERAP1 levels in the cell extracts and culture medium

were subjected to statistical analysis (lower panels, N=3).

Figure 5 - Evidence that disulfide bonding of DR4 and DR5 are regulated by ERp44.

A. Immunoblot analysis of the expression levels of EGFR, ERO1-α, AGR2, AGR3, PDIA1, and

ERp44 across a panel of breast cancer cell lines. Actin serves as a loading control.

B. BT474 cells stably expressing ERp44 or transduced with the vector control were treated for

24 h with vehicle, 2.5 μM, or 5 μM dMtcyDTDO. Cell extracts were resolved by SDS-PAGE

under non-reducing (left panel) or reducing (right panel) conditions and analyzed by

immunoblot. O and M are disulfide bonded oligomeric and monomeric protein forms,

respectively.

C. Vector control or ERp44 knockdown MDA-MB-468 cell lines were treated for 24 h with

vehicle, 2.5 μM, or 5 μM dMtcyDTDO. Cell extracts were resolved by non-reducing SDS-PAGE

and analyzed by immunoblot (CC8, Cleaved Caspase 8). O and M are disulfide bonded

oligomeric and monomeric protein bands, respectively.

D. Expanded portions of the blots from panel C showing high molecular mass bands at the

stacking gel/resolving gel interface (SGI). Asterisks denote bands at the SGI that are more

prominent in the ERp44 knockdown cells.

Figure 6 – Expression of ERp44 thioredoxin-like repeat mutants alters DDA responses.

A. Non-reducing immunoblot analysis of vector control MDA-MB-468 cells, or cells

overexpressing wild type ERp44, ERp44[C58S], or ERp44[S61C]. O and M are disulfide

bonded oligomeric and monomeric protein bands, respectively. M (FL) and M (Cl.) represent

the monomeric full-length and cleaved forms of CDCP1, respectively.

B. Non-reducing immunoblot analysis of the indicated cell lines treated with vehicle or 7.5 μM

dFtcyDTDO for 24 h. O and M are disulfide bonded oligomeric and monomeric protein bands,

respectively.

C. Extracts from vector control or EGFR overexpressing T47D cells were analyzed by non-

reducing SDS-PAGE followed by immunoblot. O and M are disulfide bonded oligomeric and

monomeric protein bands, respectively.

D. MDA-MB-468 cells were treated for 24 h with vehicle or 2.5 μM dMtcyDTDO. The cells were

then subjected to cell surface protein labeling using membrane-impermeable amine- or thiol-

reactive biotinylation probes as described in Supplemental Information. Surface proteins were

collected using Streptavidin-coated beads and analyzed along with the flow-though (internal,

non-biotinylated proteins) by immunoblot.

Figure 7 – Scission of DR5 disulfide bonds triggers high-level DR5 expression,

oligomerization, and Caspase 8 activation.

A. MDA-MB-468 cells were treated for 24 h with 10 μM dMtcyDTDO or vehicle and cell

extracts were immunoprecipitated with ERp44 antibody and analyzed by immunoblot under

non-reducing conditions.

B. MDA-MB-468 cells ectopically expressing wild type ERp44 or the C58S or S61C mutants of

ERp44 were treated for 24 h with 10 μM dMtcyDTDO or vehicle. Left panels, cell extracts were

immunoprecipitated with ERp44 antibody and analyzed by immunoblot. Right panels, crude

extracts were analyzed for protein expression levels by immunoblot.

C. MDA-MB-468 cells expressing doxycycline-inducible DR4 and DR5 were treated with 1

μg/ml doxycycline, 10 μM dMtcyDTDO, or doxycycline + dMtcyDTDO for 24 h followed by

immunoprecipitation of ERp44 and immunoblot analysis of the immunoprecipitates and crude

lysates.

1004

1005   D. The indicated MDA-MB-468 cell lines stably expressing wild type DR5 or the Δ81-178

1006   mutant lacking part of the disulfide-rich extracellular domain were treated for 24 h as indicated

1007   and subjected to immunoprecipitation with ERp44 antibodies. The immunoprecipitates and

1008   crude cell lysates were analyzed by non-reducing SDS-PAGE/immunoblot. O and M represent

1009   disulfide bonded oligomeric and monomeric protein forms.

1010

1011   E. The indicated stable MDA-MB-468 cell lines were treated for 24 h with 5 μM dMtcyDTDO

1012   and/or 100 ng/ml doxycycline as indicated and subjected to immunoblot analysis after

1013   resolving proteins by non-reducing SDS-PAGE. O and M represent disulfide bonded

1014   oligomeric and monomeric protein forms.

1015

1016   F. Model for how the biotin-labelled DDA biotin-PyrDTDO reacts in a reversible covalent

1017   fashion with the PDI active site thiolate. Covalent DDA binding is associated with ring-opening

1018   and exposure of a nucleophilic sulfinate group; it is hypothesized that favorable interaction of

1019   the DDA sulfinate group by the PDI decreases the rate of DDA self-excision.

Fig. 1, Law et al.

Fig. 2, Law et al.

Fig. 3, Law et al.

Fig. 4, Law et al.

Fig. 5, Law et al.

Fig. 6, Law et al.

Fig. 7, Law et al.