Epidemiological and evolutionary consequences of CRISPR-Cas reactivity

Hélène Chabas[a,*], Viktor Müller[b], Sebastian Bonhoeffer[a,1], Roland R. Regoes[a,1]

[a]*Institute for Integrative Biology, ETH Zürich, Switzerland*
[b]*Institute of Biology, Eötvös Loránd University, Budapest, Hungary*

Abstract

Adaptive immune systems face a control challenge: they should react with enough strength to clear an infection while avoiding to harm their organism. CRISPR-Cas systems are adaptive immune systems of prokaryotes that defend against fast evolving viruses. Here, we explore the CRISPR-Cas control challenge and look how its reactivity, i.e. its probability to acquire a new resistance, impacts the epidemiological outcome of a phage outbreak and the prokaryote's fitness. We show that in the absence of phage evolution, phage extinction is driven by the probability to acquire at least one resistance. However, when phage evolution is fast, phage extinction is driven by an epidemiological critical threshold: any reactivity below this critical threshold leads to phage survival whereas any reactivity above it leads to phage extinction. We also show that in the absence of autoimmunity, high levels of reactivity evolve. However, when CRISPR-Cas systems are prone to autoimmune reactions, intermediate levels of reactivity are evolutionarily optimal. These results help explaining why natural CRISPR-Cas systems do not show high levels of reactivity.

Keywords: phage-bacteria interactions, CRISPR-Cas, evolutionary epidemiology, microbial evolution

Introduction

The pressure viruses exert on their host has resulted in the evolution of various anti-viral immune defences, usually classified as innate or adaptive immunity. Adaptive immunity refers to systems that can acquire new specific targets of immunity during the first encounter with a parasite and memorise this resistance. To be efficient, these systems face a control challenge: they must lessen damage caused by the virus by limiting the spread of an infection and while avoiding dangerous autoimmune reactions (Bergstrom and Antia, 2006). This challenge has resulted in a tight regulation of their reactivity and any misregulation is harmful: lower reactivity results in a predisposition to severe infections whereas higher reactivity results in a propensity for autoimmunity.

The concept of adaptive immunity originated during the study of mammalian immune systems. However, it was later discovered that many others organisms carry some forms of adaptive immunity

[*]Corresponding author
[1]These authors contributed equally to this work.

(Müller et al., 2018). In prokaryotes, adaptive immunity is represented by CRISPR-Cas systems (Clustered Regularly Interspaced Short Palindromic Repeats – CRISPR ASsociated) (Barrangou et al., 2007). These systems are composed of two loci: a CRISPR locus, that can be seen as a heritable library of small sequences derived from encountered viruses and a Cas locus that codes for all the proteins required for the system to work (Koonin et al., 2017). There are six known types of CRISPR-Cas systems that differ in their Cas genes composition, which can result in differences in their molecular mechanism (Koonin et al., 2017). However, the most common mechanism is as follows: when a cell is infected by a prokaryotic virus (phage), the invader can be detected by some Cas proteins and a sequence of approximatively 30-60 bp of its DNA (the protospacer) is integrated into the chromosomal CRISPR locus (where it is called a spacer): this is the acquisition step. Then, the CRISPR locus is transcribed and the spacer RNA is used as a guide to target the invader DNA and, upon matching, to trigger its degradation (expression and interference steps) (Barrangou et al., 2007; Hampton et al., 2020). As this mechanism relies on Watson-Crick pairing, viruses can escape by mutating their targeted protospacer (Deveau et al., 2008; Semenova et al., 2011; Sun et al., 2013).

Phages can evolve rapidly and, because for most CRISPR-Cas systems a single mutation is all that is needed to escape immunity, the escape from a spacer is usually fast (Deveau et al., 2008; Semenova et al., 2011; Sun et al., 2013). Recently, it was discovered that CRISPR-Cas systems prevent this through the generation of a diversity of spacers (van Houte et al., 2016; Chabas et al., 2018). This diversity of spacers arises during the acquisition step: phage genomes contain hundreds or thousands of potential protospacers and the choice among them is stochastic. As a consequence, each acquisition event leads to the acquisition of a different spacer in each prokaryotic cell, which at the population level, generates a diversity of spacers (Childs et al., 2012; Sun et al., 2013; van Houte et al., 2016; Paez-Espino et al., 2013; Heler et al., 2019). This diversity acts as an epidemiological shield against the spread of newly evolved escape mutants because when an escape mutant evolve against a given spacer, all hosts carrying another spacer still degrade this mutant, preventing its spread (Chabas et al., 2018). Overall, if diversity is high enough, this results in rapid phage extinction (van Houte et al., 2016).

The efficiency of CRISPR-Cas immunity (i.e. its ability to rapidly eradicate the phage population) therefore relies on its ability to generate diversity. The generation of diversity derives from the probability per infection that a cell acquires a random spacer (later called CRISPR-Cas reactivity) (Heler et al., 2017). However, CRISPR-Cas systems are also prone to autoimmunity i.e. to the acquisition of a spacer that targets a chromosomal sequence, which triggers its degradation and this is thought to be often lethal (Stern et al., 2010; Jiang et al., 2013; Vercoe et al., 2013; Gomaa et al., 2014; Wimmer and Beisel, 2020). Importantly, the level of autoimmunity of a CRISPR-Cas system is related to its reactivity (Levy et al., 2015; Workman et al., 2020). Therefore, it is likely that CRISPR-Cas systems face a similar control challenge as vertebrate adaptive immune systems: too

much reactivity increases autoimmunity but low levels of reactivity decrease the efficiency of the immune response (Weissman et al., 2020).

This control challenge is evidenced by three observations. First, CRISPR-Cas reactivity is low: one cell in a million for example for the *Streptococcus thermophilus* most active CRISPR-Cas system (Hynes et al., 2017). Second, scientists can easily generate CRISPR-Cas systems with higher reactivity but these are not the forms found naturally (Heler et al., 2017; Levy et al., 2015; Workman et al., 2020). Third, even if the study of CRISPR-Cas regulation is in its early stages, it is already clear that the reactivity of the system is tightly regulated (Patterson et al., 2017).

A previous theoretical work attempted to explore CRISPR-Cas control challenge (Bradde et al., 2019). They found that autoimmunity decreases the optimal reactivity and that the optimal reactivity is governed by the interactions of the initial size of the bacteria and phages populations. However, in their work, they do not account that i) phages can infect bacteria carrying a spacer, which results in phage degradation (Barrangou et al., 2007), ii) spacers can be escaped by phage mutation (Deveau et al., 2008; Semenova et al., 2011; Sun et al., 2013), iii) CRISPR-Cas generates a diversity of resistance which protects the population against phage evolution (Childs et al., 2012; van Houte et al., 2016; Chabas et al., 2018). Consequently, it is unclear how a change in CRISPR-Cas reactivity changes the epidemiological outcome of a phage infection and the fitness of the prokaryote carrying it. Does an increase in CRISPR-Cas reactivity always improve its efficiency? Does a low reactivity always decrease bacterial fitness in the presence of phage? What is the impact of phage evolution on CRISPR-Cas efficiency and optimal reactivity? How does autoimmunity affect the optimal reactivity?

Answering these questions is experimentally challenging as it would require to finely modify CRISPR-Cas reactivity and phage mutation rates without altering other biological parameters. We therefore chose mathematical modeling to explore the consequences of CRISPR-Cas reactivity on the epidemiological outcome and on the bacterial fitness. Specifically, we developed a stochastic model of coupled population dynamics and genetics that simulates the early dynamics of a virulent phage outbreak in a population carrying naive CRISPR-Cas immunity. Our model accounts i) the ability of CRISPR-Cas to generate spacer diversity by the stochastic acquisition of spacers, ii) for phage escape of a given spacer through mutation and iii) for phage clearance resulting from the infection of cells carrying non-escaped spacers. In the absence of autoimmunity, we found that when phage evolution is slow, increasing CRISPR-Cas reactivity is beneficial as it increases the probability of generating at least one spacer that can target a phage. However, when phage evolution is fast, phage population survival is governed by an epidemiological critical threshold: below a certain reactivity, the phage always remains in the system, whereas above this critical threshold, it is always driven to extinction. This is due to a non-linear relationship between CRISPR-Cas reactivity and the diversity of spacers that is generated. We also show that strains with higher reactivities

have higher fitness. In the presence of autoimmunity, epidemiological outcomes are not modified, except for very high reactivities that impair bacterial survival. However, when strains with different reactivities compete, intermediate reactivities are selected in the presence of phages whereas the lowest reactivity is selected for in their absence. Finally, we show that lowering propensity for autoimmunity (for example by the presence of a self/non-self discrimination mechanism) decreases its cost and therefore selects for higher levels of reactivity. We then discuss the implications of these findings for the evolutionary ecology of CRISPR-Cas and phages.

Results

To study the relationship between CRISPR-Cas reactivity, spacer diversity and the outcome of a phage infection, we developed a stochastic epidemiological model (see Methods for a full presentation of the model and Figure 1 for a graphic representation). Briefly, naive CRISPR-Cas bacteria are infected by virulent wildtype phage WT. Infection can lead to one of three outcomes with different probabilities: 1) the phage kills the bacteria and produces WT progeny, 2) the phage kills the cell and produces WT and a randomly-chosen mutated phage, 3) CRISPR-Cas kills the infecting phage by acquiring a randomly-selected spacer. WT phages can also enter a cell carrying a spacer and this results in phage death, without consequences for the cell. Escape phages can also infect a cell with a spacer and in this case, the outcome depends on the identity of the spacer and of the escape mutation. If the mutation escapes this specific spacer, the infection kills the cell and produces escape virions; if not, the phage is killed by the CRISPR-Cas system without consequences for the cell. These processes have been implemented stochastically and we can therefore follow the dynamics of each genotype, including its appearance and extinction (see Figure 2 for representative simulations).

The diversity of spacers protects against phage epidemics

To evaluate the validity of our model, we checked whether it predicts outcomes that have been experimentally observed. First, we tested if the model predicts that initial spacer diversity protects the bacterial population against phage outbreaks. Indeed, it was shown experimentally that spacer diversity protects the bacterial population by preventing the spread of escape phages (van Houte et al., 2016). To do so, we ran 100 simulations with an initial bacterial population composed of 50% of naive CRISPR-Cas bacteria and 50% of resistant cells with various levels of diversity. This population was infected by 10^5 WT phages and we looked at the probability of phage extinction at the end of the simulations. We find (Figure S1) that increasing the initial diversity of spacers in the host population increases the probability of phage extinction, which fits experimental results obtained by van Houte et al. (2016).

Higher reactivities lead to higher diversity of spacers

Second, we wanted to confirm that higher reactivities lead to higher levels of spacers diversity. Indeed, Heler et al. found that a mutant with a higher spacer acquisition generates more diversity

Figure 1: Important transitions in the model

Figure 2: Example of time courses the model produces when $6 * 10^6$ S bacteria are infected by 10^5 WT phages ($\alpha = 5.12 * 10^{-6}$, $\mu = 10^{-6}$). In panel A, B and C, the phage drives the bacteria to extinction whereas in panel D, E and F the phage is driven to extinction by the CRISPR-Cas system. Panels A and D show the number of bacteria that are sensitive (black) and resistant (blue). Panels B and E show the number of phages that are WT (black) and escape mutants (purple). Panels C and F show the strains diversity of the bacterial population through time, each colour representing a strain.

(Heler et al., 2017). We ran 100 simulations of an outbreak due to an infection by 10^5 WT phages that cannot evolve in a naive bacterial population. We look at the diversity of the bacterial population at the very beginning of the outbreak (at the time when sensitive bacteria are driven extinct). As expected, we observe that higher reactivities lead to higher levels of spacers diversity (Figure S2, panel A, black line). Notably, spacers diversity is not linearly linked to the probability of acquisition. Instead, three phases can be observed: at very low reactivity, the level of spacers diversity does not increase; at intermediate reactivity, the diversity increases rapidly and at high reactivities, the system has reached the maximal number of potential spacers and saturates. Therefore, if any increases of spacers diversity are selected for, only increases in the probability of acquisition that result in a transition from very low levels to intermediate levels of acquisition or an increase in the intermediate range can lead to an increase in bacterial spacers diversity.

Phage survival depends on CRISPR-Cas reactivity and phage evolution

To understand the relationship between CRISPR-Cas reactivity and phage extinction, we ran 100 simulations for various reactivities. For each, we calculated the probability of phage extinction by looking if at least one of the phage genotype remains at the end of the simulation. We first looked at the outcomes of outbreaks that are caused by phages that cannot escape CRISPR-Cas ($\mu = 0$). We observe that for low reactivities, rises in reactivities increase the probability of phage extinction (Figure 3, panel A, black line). When the probability of acquisition reaches a certain value (around 10^{-6}), the probability of phage extinction saturates to 1. Therefore, up to a certain value, evolving higher reactivities enhances the efficacy of CRISPR-Cas.

However in living systems, phages can escape CRISPR-Cas immunity by mutations. How does phage evolution change the outcome of an infection? We repeated the previous analysis for various probabilities of phage evolution. We find that when phages evolve slowly ($\mu = 10^{-8}$ and $\mu = 10^{-7}$), the situation does not massively differ from the case without evolution (Figure 3, panel A, purple and green lines). However, when phage evolution is fast ($\mu = 10^{-6}$), we observe that the outcome of the outbreak is determined by an epidemiological critical threshold: below a probability of acquisition around 10^{-5}, the probability of phage extinction is equal to zero, whereas above this critical threshold, phages are always driven to extinction by CRISPR-Cas immunity (Figure 3, panel A, red line). It is notable that phage evolution increases the value of the minimal CRISPR-Cas reactivity that assures phage extinction: frequent phage evolution puts a high pressure on the CRISPR-Cas system.

In addition to purging viruses from the host population, immune systems also prevent the spread of a parasite. To see whether increased CRISPR-Cas reactivities decrease the spread of phages, we looked at the size of phage outbreaks for 100 simulations. We found that increasing CRISPR-Cas reactivity decreases the size of the outbreak (Figure S3). Importantly, increasing the probability of spacer acquisition when it is too small to lead the phage to extinction, still results in a decrease in the

Figure 3: Influence of CRISPR-Cas reactivity on the probability of phage extinction in the absence of autoimmunity. A) Probability of phage survival when infecting bacteria with various reactivities. The different colours corresponds to different levels of phage evolution (μ): black, no evolution ($\mu = 0$), purple $\mu = 10^{-8}$; green $\mu = 10^{-7}$ and red $\mu = 10^{-6}$. B) Probability for bacteria with various reactivities to generate at least one single resistant cells. The different colours corresponds to different levels of phage evolution (μ): in black, no evolution ($\mu = 0$), in purple $\mu = 10^{-8}$; in green $\mu = 10^{-7}$ and in red $\mu = 10^{-6}$. The grey line corresponds to the probability of phage extinction in the absence of phage evolution $\mu = 0$. C) Relationship between CRISPR reactivity (x-axis), initial spacer diversity (y-axis) and phage extinction (color) when phage evolution is high ($\mu = 10^{-6}$).

size of the outbreak. As a consequence it seems beneficial for bacteria to evolve higher reactivities, even if these levels do not increase the probability of phage extinction.

When phage evolution is rare, the probability of generating one single spacer drives the epidemiological outcome

How can we explain these results? First, we tried to understand how in the absence of phage evolution, the increase in CRISPR-Cas reactivity results in a higher probability of phage extinction. We reasoned that if CRISPR-Cas reactivity is too low, there may be a non-zero probability that no bacteria acquires a spacer: then, the phage would spread in the population until all the bacteria have been killed. To check for this, we looked at the number of simulations in which the proportion of resistant strains is positive at the beginning of the outbreak i.e. at the time the sensitive cells reach extinction. We find that indeed increasing CRISPR-Cas reactivity increases the probability of generating at least one resistant genotype (Figure 3, panel B). Critically, the relationship between CRISPR-Cas reactivity and the probability of phage extinction is identical as the one between CRISPR-Cas reactivity and the probability of generating at least one single resistant strain. This means that the probability of generating at least one resistant strain explains the impact of CRISPR-Cas reactivity on the probability of phage extinction.

When phage evolution is frequent, the probability to generate enough diversity explains the epidemiological outcome

When phage evolution is frequent, the probability of generating one spacer does not recapitulate the epidemiological outcome. We wondered how can the epidemiological critical threshold be explained. We reasoned that spacer diversity is related to phage extinction and to CRISPR-Cas reactivity and that it is likely that phage are driven extinct when initial spacer diversity is high. Because spacer acquisition and phage evolutionary emergence are stochastic processes, each value of reactivity would then be associated with a probability to generate a certain diversity and each diversity in turn has a certain probability to drive phage to extinction. This should result in simulations with low initial diversity having a low probability of phage extinction. Indeed, when we look at the initial spacer diversity depending on CRISPR-Cas reactivity and at the epidemiological outcome, we observe that simulations with low initial spacer diversity tends to result in phage survival, whereas simulations with high initial spacer diversity are likely to result in phage extinction (Figure 3, panel C). Importantly, there seems to be a diversity critical threshold: above this value, the phage is very likely to go extinct whereas below this diversity phage extinction is unlikely.

In the absence of autoimmunity, evolving higher reactivities is always beneficial

So far, we have looked at the impact on the epidemiological outcome of various CRISPR-Cas reactivities. What are the fitness consequences for the bacteria to evolve a higher CRISPR-Cas reactivity? To determine this, we simulated competition experiments. Two sensitive strains, that differ only by their reactivity, are mixed together and we infect this population with the phage

Figure 4: Fitness of a bacterial strain with various reactivity when in competition with a strain with a reactivity $\alpha = 10^{-5}$.

A) In the absence of autoimmunity, B) In the presence of autoimmunity (Propensity = 40). Colours corresponds to phage evolution: in black, no evolution ($\mu = 0$) and in red high phage evolution ($\mu = 10^{-6}$) Each point corresponds to one simulation. Only simulations in which bacteria survive the infection have been plotted.

P_{WT}. At the end of the simulation, we calculate the proportion of each strain (sensitive + resistant cells) and we deduce the relative fitness. We observe that it is always beneficial for a strain to evolve higher reactivities (Figure 4, panel A). However, this is not what is observed in nature, when CRISPR-Cas reactivity is usually low (Hynes et al., 2017). In addition, scientists can easily generate CRISPR-Cas systems with higher reactivities (Heler et al., 2017; Levy et al., 2015; Workman et al., 2020), so their absence is not the result of evolutionary constraints.

Autoimmunity selects for intermediate levels of reactivity

Natural CRISPR-Cas systems are prone to autoimmunity (Stern et al., 2010; Jiang et al., 2013; Vercoe et al., 2013; Gomaa et al., 2014), i.e. they can acquire spacers derived from the prokaryote chromosome. To explore the consequences of this autoimmunity, we added autoimmunity as an additional cause of bacterial death (see Methods for details).

Using this extended model, we looked at the probability of phage extinction for various CRISPR-Cas reactivities. We found that the overall outcome of an outbreak is not modified by autoimmunity, except for very high reactivities, for which the bacteria are driven to extinction (Figure S4). Then we wanted to understand the evolutionary consequences of autoimmunity on bacteria: to do so, we simulate competition experiments between a WT strain with a reactivity of 10^{-5} and a mutant strain with various levels of reactivities. At the beginning of the simulation, both strains have equal representation. We observe that in the presence of autoimmunity, intermediate levels of reactivity are selected for (Figure 4, panel B). To get better insights into the cost of autoimmunity, we ran competition in the absence of phages (Figure S5). We observe that it is always beneficial to have a lower reactivity, i.e. that autoimmunity always causes a detectable cost. In addition, the cost on

Figure 5: High levels of propensity selects for lower optimal reactivity.
A WT strain with a reactivity at 10^{-5} competes with a mutant strain with various reactivity in the presence of phage that can escape spacers ($\mu = 10^{-6}$). For each propensity (0.4, 4, 40, 400 and 4000), we took the median fitness of the bacteria and reported as the optimal reactivity the reactivity associated with the highest fitness.

fitness increases when autoimmunity increases: as a consequence, very high reactivity are massively selected against, whereas low and intermediate levels of reactivity are slightly deleterious.

CRISPR-Cas propensity for autoimmunity determines the optimal reactivity

We reasoned that various CRISPR-Cas systems may not have the same propensity for autoimmunity and consequently the cost of autoimmunity can vary independently of the reactivity. To explore the consequences for bacterial fitness of various propensities, we ran 100 competition simulations for various levels of propensity. We observe that the higher the propensity, the lower the optimal reactivity (Figure 5). Interestingly, we observe that high levels of propensity decreases the optimal reactivity at the level of the minimal reactivity needed to assures phage extinction. We wondered then if high levels of propensity can impair the efficiency of CRISPR-Cas immunity. To test for this, we ran 100 simulations of the infection of a bacterial population with high propensity to autoimmunity ($Propensity = 4000$) and look at phage survival. We found that indeed high levels of propensity makes really high levels of reactivity unbearable, however, the effect is not strong enough to prevent phage extinction for intermediate levels of reactivity (Figure S6).

Discussion

We studied the control challenge faced by the prokaryotic adaptive immune system, CRISPR-Cas. We find that the outcome of a phage outbreak is governed by CRISPR-Cas reactivity: 1) when resistance escape is impossible, rises in CRISPR-Cas reactivity increases the probability of phage extinction and this is governed by the probability for CRISPR-Cas to generate at least one single resistant genotype; 2) when the infected phage has a high rate of resistance escape, phage extinction is controlled by an epidemiological critical threshold: any value below a certain level of reactivity leads to phage persistence whereas any reactivity above leads to phage extinction. Importantly, this

critical threshold is above the minimal reactivity that assures phage extinction in the absence of phage evolution. We also find that in the absence of autoimmunity, evolving higher reactivities is always beneficial. However, CRISPR-Cas susceptibility for autoimmunity results in the selection for intermediate reactivities: indeed, during an outbreak, high reactivities levels results in high fitness costs that negate the benefits of CRISPR-Cas. Finally, we show that the optimal reactivity level depends on the propensity of the system for autoimmunity.

In this work, we simulate outbreaks of two types of phages: phages that have a high rate of escaping spacers and phages that cannot escape spacers. This can correspond to two types of CRISPR-Cas immunity. Indeed, even if all types of CRISPR-Cas rely on Watson-Crick pairing to guide the CRISPR complex, their sensitivity to phage mutations varies greatly. For type I and type II systems, a single mutation in the protospacer adjacent motif or in the seed sequence is all that is needed to completely escape the spacer and this usually happens rapidly (Deveau et al., 2008; Chabas et al., 2018, 2019): therefore the model with various phage probability of evolution can be used to study these CRISPR-Cas systems. On the other hand, the interference mechanism of type III CRISPR-Cas systems is resilient to phage mutation and phage escape is extremely rare (Manica et al., 2013; Pyenson et al., 2017): the model in the absence of phage evolution can therefore be used to study interactions between phages and type III CRISPR-Cas systems. Our model predicts important differences in epidemiological outcomes for type I/II and type III CRISPR-Cas systems. For type I/II CRISPR-Cas systems, the control of phage evolution is mediated by spacer diversity and there is a reactivity critical threshold whereas for type III CRISPR-Cas systems, spacers diversity does not impact phage extinction and the epidemiological outcome is driven by the probability to generate at least one resistant genotype. Because the minimal reactivity that assures to acquire at least one spacer is lower than the reactivity critical threshold, this would suggest that type III CRISPR-Cas systems evolve lower reactivities than type I/II CRISPR-Cas systems. This lower reactivity for type III would indeed advantage the cell as it would minimise the cost of autoimmunity.

Experimental data to support these predictions are lacking. Even if the importance of spacers diversity for the efficiency of type I/II CRISPR-Cas systems has been documented both experimentally and theoretically (van Houte et al., 2016; Chabas et al., 2018; Payne et al., 2018), the relationship between phage extinction probability and CRISPR-Cas reactivity has, as far as we know, not been explored. In addition, to the best of our knowledge, there has been no experimental evolution studies between phages and naive type III CRISPR-Cas system. This is most likely due to the lack of type III CRISPR-Cas systems showing naive spacer acquisition in laboratory conditions and this makes the discovery of naive acquisition by the type III CRISPR system of *Thermus thermophilus* very promising (Artamonova et al., 2020).

Considering the importance of CRISPR-Cas reactivity for the epidemiological outcome and for the fitness of the prokaryote carrying the system, we could expect CRISPR-Cas reactivity to be under tight control and its evolution to be under strong selection pressure. Indeed, studies of the regulation of CRISPR-Cas show that CRISPR-Cas expression is finely regulated (Patterson et al., 2017). For example, it was shown that in some bacteria, CRISPR-Cas reactivity is upregulated by quorum sensing and this has been explained by the higher risk of phage outbreaks when cell density is high (Høyland-Kroghsbo et al., 2013; Patterson et al., 2016; Høyland-Kroghsbo et al., 2017). This finding makes sense in the light of our predictions as in the absence of phage infection, autoimmunity makes the system always costly and therefore CRISPR-Cas reactivity should be maintained to a minimum when the risk of infection is low. On the phage side, we show that lower reactivities increase phage survival. Consequently, it is probably beneficial for phages to decrease CRISPR-Cas reactivity. Many phages carry anti-CRISPR proteins, i.e. small proteins that inhibit some CRISPR-Cas systems (Borges et al., 2017). To the best of our knowledge, most of them inhibit interference (Wiegand et al., 2020), a handful inhibit both acquisition and interference (Vorontsova et al., 2015) and none inhibit only acquisition: our model suggests that families of anti-CRISPR that decrease CRISPR-Cas reactivity may exist. Finally, one should not forget that an observed reactivity is also influenced by the ecological conditions: for example, it has been proposed that infections at lower temperature boosts CRISPR-Cas reactivity by slowing down the intra-host phage kinetics and therefore letting more time for the system to react (Høyland-Kroghsbo et al., 2018). This calls for caution when measuring and studying of CRISPR-Cas reactivity as the value measured might be the result of the system itself, its regulation, the infective phage and ecological conditions and may have limited prediction power if any of this changes.

It is striking that our model predicts an epidemiological critical threshold that is around 10^{-5}, ten times higher than the reactivity of *S. thermophilus*' most active CRISPR-Cas system (Hynes et al., 2017). Importantly, when infected by a virulent phage, *S. thermophilus*, defending exclusively with its CRISPR-Cas systems, survives and lead the phage to extinction in the long term (Paez-Espino et al., 2013; Common et al., 2019). There are two potential hypotheses that may explain this discrepancy between theoretical predictions and experimental data. First, the epidemiological outcome may depend on the value of the parameters. While we carefully parameterized our model using all available estimates from the experimental literature, some key parameters have, to our knowledge, not been estimated yet. Most notably, there is no estimate of phage infectivity.

To test if our particular choice for the infectivity is the reason for the discrepancy, we ran the model with various infectivities. Varying the value of the infectivity only marginally affected the results (see supplementary figure S7). Therefore, we are confident that this is unlikely to explain this discrepency. Therefore, we are confident that this is unlikely to explain this discrepancy. Second, we have been making too simplifying assumptions. Importantly, contrary to our model, CRISPR systems can acquire multiple spacers against a single phage and this is likely important for phage

extinction. In addition, many CRISPR-Cas systems display some form of priming, which boosts the acquisition of a second spacer, either because the cell already carries a partially matching spacer (type I) or because it already possesses an efficient spacer (type II) (Datsenko et al., 2012; Fineran et al., 2014; Nussenzweig et al., 2019). The impact of these on the relationship between CRISPR-Cas reactivity, the epidemiological outcome, autoimmunity and prokaryote fitness remain to be explored and it is possible that this would change the epidemiological outcome for a given reactivity and the value of the optimal reactivity.

Acknowledgements

The authors thank Judith Bouman and Peter Ashcroft for their help in coding and Thomas Aubier and Mircea Sofonea for scientific discussions. This work was supported by an ETH Zurich Postdoctoral Fellowship.

Authors contributions

Conceptualization: HC; Methodology: HC, VM, RRR; Investigation: HC; Writing - Original Draft: HC; Writing - Review & Editing: HC, VM, SB, RRR; Visualization: HC; Supervision: SB, RRR; Project Administration: SB; Funding Acquisition: HC.

Declaration of interests

The authors declare no competing interests.

References

Artamonova, D., Karneyeva, K., Medvedeva, S., Klimuk, E., Kolesnik, M., Yasinskaya, A., Samolygo, A., Severinov, K., 2020. Spacer acquisition by type iii crispr–cas system during bacteriophage infection of thermus thermophilus. Nucleic Acids Research.

Barrangou, R., Fremaux, C., Deveau, H., Richards, M., Boyaval, P., Moineau, S., Romero, D. A., Horvath, P., 2007. Crispr provides acquired resistance against viruses in prokaryotes. Science 315 (5819), 1709–1712.

Bergstrom, C. T., Antia, R., 2006. How do adaptive immune systems control pathogens while avoiding autoimmunity? Trends in ecology & evolution 21 (1), 22–28.

Borges, A. L., Davidson, A. R., Bondy-Denomy, J., 2017. The discovery, mechanisms, and evolutionary impact of anti-crisprs.

Bradde, S., Mora, T., Walczak, A. M., 2019. Cost and benefits of clustered regularly interspaced short palindromic repeats spacer acquisition. Philosophical Transactions of the Royal Society B 374 (1772), 20180095.

Chabas, H., Lion, S., Nicot, A., Meaden, S., van Houte, S., Moineau, S., Wahl, L. M., Westra, E. R., Gandon, S., 2018. Evolutionary emergence of infectious diseases in heterogeneous host populations. PLoS biology 16 (9), e2006738.

Chabas, H., Nicot, A., Meaden, S., Westra, E. R., Tremblay, D. M., Pradier, L., Lion, S., Moineau, S., Gandon, S., 2019. Variability in the durability of crispr-cas immunity. Philosophical Transactions of the Royal Society B 374 (1772), 20180097.

Childs, L. M., Held, N. L., Young, M. J., Whitaker, R. J., Weitz, J. S., 2012. Multiscale model of crispr-induced coevolutionary dynamics: diversification at the interface of lamarck and darwin. Evolution: International Journal of Organic Evolution 66 (7), 2015–2029.

Common, J., Morley, D., Westra, E. R., van Houte, S., 2019. Crispr-cas immunity leads to a coevolutionary arms race between streptococcus thermophilus and lytic phage. Philosophical Transactions of the Royal Society B 374 (1772), 20180098.

Datsenko, K. A., Pougach, K., Tikhonov, A., Wanner, B. L., Severinov, K., Semenova, E., 2012. Molecular memory of prior infections activates the crispr/cas adaptive bacterial immunity system. Nature communications 3 (1), 1–7.

Deveau, H., Barrangou, R., Garneau, J. E., Labonté, J., Fremaux, C., Boyaval, P., Romero, D. A., Horvath, P., Moineau, S., 2008. Phage response to crispr-encoded resistance in streptococcus thermophilus. Journal of bacteriology 190 (4), 1390–1400.

Fineran, P. C., Gerritzen, M. J., Suárez-Diez, M., Künne, T., Boekhorst, J., van Hijum, S. A., Staals, R. H., Brouns, S. J., 2014. Degenerate target sites mediate rapid primed crispr adaptation. Proceedings of the National Academy of Sciences 111 (16), E1629–E1638.

Gomaa, A. A., Klumpe, H. E., Luo, M. L., Selle, K., Barrangou, R., Beisel, C. L., 2014. Programmable removal of bacterial strains by use of genome-targeting crispr-cas systems. MBio 5 (1).

Hampton, H. G., Watson, B. N., Fineran, P. C., 2020. The arms race between bacteria and their phage foes. Nature 577 (7790), 327–336.

Heler, R., Wright, A. V., Vucelja, M., Bikard, D., Doudna, J. A., Marraffini, L. A., 2017. Mutations in cas9 enhance the rate of acquisition of viral spacer sequences during the crispr-cas immune response. Molecular cell 65 (1), 168–175.

Heler, R., Wright, A. V., Vucelja, M., Doudna, J. A., Marraffini, L. A., 2019. Spacer acquisition rates determine the immunological diversity of the type ii crispr-cas immune response. Cell host & microbe 25 (2), 242–249.

Høyland-Kroghsbo, N. M., Mærkedahl, R. B., Svenningsen, S. L., 2013. A quorum-sensing-induced bacteriophage defense mechanism. MBio 4 (1).

Høyland-Kroghsbo, N. M., Muñoz, K. A., Bassler, B. L., 2018. Temperature, by controlling growth rate, regulates crispr-cas activity in pseudomonas aeruginosa. MBio 9 (6).

Høyland-Kroghsbo, N. M., Paczkowski, J., Mukherjee, S., Broniewski, J., Westra, E., Bondy-Denomy, J., Bassler, B. L., 2017. Quorum sensing controls the pseudomonas aeruginosa crispr-cas adaptive immune system. Proceedings of the National Academy of Sciences 114 (1), 131–135.

Hynes, A. P., Lemay, M.-L., Trudel, L., Deveau, H., Frenette, M., Tremblay, D. M., Moineau, S., 2017. Detecting natural adaptation of the streptococcus thermophilus crispr-cas systems in research and classroom settings. nature protocols 12 (3), 547–565.

Jiang, W., Bikard, D., Cox, D., Zhang, F., Marraffini, L. A., 2013. Rna-guided editing of bacterial genomes using crispr-cas systems. Nature biotechnology 31 (3), 233–239.

Johnson, P., 2019. adaptivetau: Tau-Leaping Stochastic Simulation. R package version 2.2-3.
URL https://CRAN.R-project.org/package=adaptivetau

Koonin, E. V., Makarova, K. S., Zhang, F., 2017. Diversity, classification and evolution of crispr-cas systems. Current opinion in microbiology 37, 67–78.

Levy, A., Goren, M. G., Yosef, I., Auster, O., Manor, M., Amitai, G., Edgar, R., Qimron, U., Sorek, R., 2015. Crispr adaptation biases explain preference for acquisition of foreign dna. Nature 520 (7548), 505–510.

Manica, A., Zebec, Z., Steinkellner, J., Schleper, C., 2013. Unexpectedly broad target recognition of the crispr-mediated virus defence system in the archaeon sulfolobus solfataricus. Nucleic acids research 41 (22), 10509–10517.

Modell, J. W., Jiang, W., Marraffini, L. A., 2017. Crispr–cas systems exploit viral dna injection to establish and maintain adaptive immunity. Nature 544 (7648), 101–104.

Müller, V., De Boer, R. J., Bonhoeffer, S., Szathmáry, E., 2018. An evolutionary perspective on the systems of adaptive immunity. Biological Reviews 93 (1), 505–528.

Nussenzweig, P. M., McGinn, J., Marraffini, L. A., 2019. Cas9 cleavage of viral genomes primes the acquisition of new immunological memories. Cell host & microbe 26 (4), 515–526.

Paez-Espino, D., Morovic, W., Sun, C. L., Thomas, B. C., Ueda, K.-i., Stahl, B., Barrangou, R., Banfield, J. F., 2013. Strong bias in the bacterial crispr elements that confer immunity to phage. Nature communications 4 (1), 1–7.

Patterson, A. G., Jackson, S. A., Taylor, C., Evans, G. B., Salmond, G. P., Przybilski, R., Staals, R. H., Fineran, P. C., 2016. Quorum sensing controls adaptive immunity through the regulation of multiple crispr-cas systems. Molecular cell 64 (6), 1102–1108.

Patterson, A. G., Yevstigneyeva, M. S., Fineran, P. C., 2017. Regulation of crispr–cas adaptive immune systems. Current opinion in microbiology 37, 1–7.

Payne, P., Geyrhofer, L., Barton, N. H., Bollback, J. P., 2018. Crispr-based herd immunity can limit phage epidemics in bacterial populations. Elife 7, e32035.

Pyenson, N. C., Gayvert, K., Varble, A., Elemento, O., Marraffini, L. A., 2017. Broad targeting specificity during bacterial type iii crispr-cas immunity constrains viral escape. Cell host & microbe 22 (3), 343–353.

R Core Team, 2018. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.
URL https://www.R-project.org/

Semenova, E., Jore, M. M., Datsenko, K. A., Semenova, A., Westra, E. R., Wanner, B., Van Der Oost, J., Brouns, S. J., Severinov, K., 2011. Interference by clustered regularly interspaced short palindromic repeat (crispr) rna is governed by a seed sequence. Proceedings of the National Academy of Sciences 108 (25), 10098–10103.

Stern, A., Keren, L., Wurtzel, O., Amitai, G., Sorek, R., 2010. Self-targeting by crispr: gene regulation or autoimmunity? Trends in genetics 26 (8), 335–340.

Sun, C. L., Barrangou, R., Thomas, B. C., Horvath, P., Fremaux, C., Banfield, J. F., 2013. Phage mutations in response to crispr diversification in a bacterial population. Environmental microbiology 15 (2), 463–470.

van Houte, S., Ekroth, A. K., Broniewski, J. M., Chabas, H., Ashby, B., Bondy-Denomy, J., Gandon, S., Boots, M., Paterson, S., Buckling, A., et al., 2016. The diversity-generating benefits of a prokaryotic adaptive immune system. Nature 532 (7599), 385–388.

Vercoe, R. B., Chang, J. T., Dy, R. L., Taylor, C., Gristwood, T., Clulow, J. S., Richter, C., Przybilski, R., Pitman, A. R., Fineran, P. C., 2013. Cytotoxic chromosomal targeting by crispr/cas systems can reshape bacterial genomes and expel or remodel pathogenicity islands. PLoS Genet 9 (4), e1003454.

Vorontsova, D., Datsenko, K. A., Medvedeva, S., Bondy-Denomy, J., Savitskaya, E. E., Pougach, K., Logacheva, M., Wiedenheft, B., Davidson, A. R., Severinov, K., et al., 2015. Foreign dna acquisition by the if crispr–cas system requires all components of the interference machinery. Nucleic acids research 43 (22), 10848–10860.

Weissman, J. L., Stoltzfus, A., Westra, E. R., Johnson, P. L., 2020. Avoidance of self during crispr immunization. Trends in Microbiology.

Wiegand, T., Karambelkar, S., Bondy-Denomy, J., Wiedenheft, B., 2020. Structures and strategies of anti-crispr-mediated immune suppression. Annual Review of Microbiology 74.

Wimmer, F., Beisel, C. L., 2020. Crispr-cas systems and the paradox of self-targeting spacers. Frontiers in Microbiology 10, 3078.

Workman, R. E., Pammi, T., Nguyen, B. T., Graeff, L. W., Smith, E., Sebald, S. M., Stoltzfus, M. J., Modell, J. W., 2020. A natural single-guide rna repurposes cas9 to autoregulate crispr-cas expression. bioRxiv.

Methods

Resource availability

Lead Contact

Further inquiries and requests should be addressed to Hélène Chabas (helene.chabas@ens-lyon.org).

Materials Availability

This study did not generate new unique reagents.

Data and Code Availability

All code will be made available on github at the time of publication.

Method details

Model definition

In our mathematical model, we consider a bacterial population composed of a sensitive strain S carrying a naive CRISPR-Cas system, i. e. with no pre-exisiting spacers. Upon infection by phage P_{WT}, S can acquire a spacer with a probability α and evolve resistance against P_{WT}. We assume that the phage genome possesses n_s protospacers and that the CRISPR-Cas system randomly acquires one of them with equal probability. As a result, the infection of S will lead to the evolution of a diversified population composed of a subset of resistant strains R_i, all carrying a single different spacer (=resistance) against phage WT. All these strains follow a logistic growth with a growth rate g and a total carrying capacity K.

However, the infection of S by P_{WT} can also be successful because the CRISPR-Cas system fails to stop the infection and the phage reproduces on S. A successful infection can either lead to the amplification of P_{WT} and produce b_{WT} progeny phages, or result in the escape of the phage from one spacer. Specifically, each protospacer has a probability μ to mutate and escape the spacer that targets it. In this case, we assume that each phage genotype produces half of their respective progeny: $\frac{b_{WT}}{2}$ P_{WT} and $\frac{b_m}{2}$ P_i). Each of the escape phage P_i can successfully infect the

Variables	Biological interpretation	
S	Population size of sensitive bacteria (no spacer)	
R_i	Population size of resistant bacteria carrying spacer i	
T	Sum of all bacterial population sizes	
P	Population size of WT phage (no escape mutation)	
P_i	Population size of escape phage (escape mutation in protospacer i)	

Parameters	Biological interpretation	Value
α	CRISPR-Cas reactivity: probability to acquire 1 spacer	variable
μ	Mutation probability of a protospacer	variable
g	Bacterial growth rate per hour	$1.3/h$
K	Bacterial carrying capacity	10^{11}
n_s	Number of phage protospacers	100
β	Infectivity rate constant	$10^{-6}/h$
b_{WT}	Burst size of the WT phage (P)	100
b_m	Burst size of the escape phages (P_i)	70
$Propensity$	CRISPR-Cas propensity to autoimmunity	variable

Table 1: Summary of the variables and parameters used in the model.

corresponding resistant bacteria R_i in addition to the sensitive strain S. If they infect another resistant strain R_j, they are degraded by the CRISPR-Cas system with no consequences for the bacteria.

The interaction between bacterial and phage populations described above can be summarized by the following differential equations:

$$\frac{dS}{dt} = \underbrace{g_s S}_{\text{Growth}} - \underbrace{g_s S \frac{T}{K}}_{\text{Death}} - \underbrace{(1-\alpha)(1-n_s\mu)\beta SP}_{\text{Predation by WT phages}} - \underbrace{\beta S \sum_{j=1}^{n_s} P_i}_{\text{Predation by escape phages}} - \underbrace{\alpha\beta SP}_{\text{Spacer acquisition}} - \underbrace{(1-\alpha)n_s\mu\beta SP}_{\text{Mutation of phage}}$$

(1)

$$\frac{dR_i}{dt} = \underbrace{g_r R_i}_{\text{Growth}} - \underbrace{g_r R_i \frac{T}{K}}_{\text{Death}} + \underbrace{\frac{\alpha}{n_s}\beta SP}_{\text{Spacer acquisition}} - \underbrace{\beta P_i R_i}_{\text{Predation by escape phages}}$$

(2)

$$\frac{dP}{dt} = \underbrace{b_{\text{wt}}(1-\alpha)(1-n_s\mu)\beta SP}_{\text{Successful infection of S}} + \underbrace{\frac{b_{\text{wt}}}{2}(1-\alpha)n_s\mu\beta SP}_{\text{Production of WT when phage escape}} - \underbrace{\beta P \sum_{j=1}^{n_s} R_j}_{\text{Failed infections of R bacteria}} - \underbrace{\alpha\beta SP}_{\substack{\text{Failed infections when} \\ \text{S acquires a spacer}}}$$

(3)

$$\frac{dP_i}{dt} = \underbrace{b_m \beta P_i (S+R_i)}_{\text{Amplification on S and } R_i \text{ hosts}} + \underbrace{\frac{b_{\text{m}}}{2}(1-\alpha)\mu\beta SP}_{\text{Mutation of WT Phage}} - \underbrace{\beta P_i \left(\left(\sum_{j=1}^{n_s} R_j \right) - R_i \right)}_{\text{Failed infections on resistant bacteria}}$$

(4)

We implemented the dynamics described by Equations 1–4 stochastically. A list of all model variables and parameters and their definition can be found in table 1. Figure 1 illustrates the

different processes we considered in our model.

We are aware that our model does not take into considerations some complexities of CRISPR-Cas biology, such as priming (Fineran et al., 2014), acquisition of multiple spacers (Barrangou et al., 2007), heterogeneity in the probability of choosing a spacer (Heler et al., 2019; Modell et al., 2017), or multiple phage mutations.

Overall, our model is similar (though simpler) of the model developed by Childs et al. (2012).

Mathematical modelling of autoimmunity

To test the impact of autoimmunity on bacterial fitness and CRISPR-Cas immunity, we added autoimmunity as an additional cause of bacterial death. To the best of our knowledge, the precise relationship between CRISPR-Cas reactivity and CRISPR-Cas autoimmunity is unknown. However, we know that an increase in reactivity results in higher autoimmunity (Levy et al., 2015; Workman et al., 2020). In addition, there is limited evidence that an increase in reactivity is proportional to autoimmunity (Levy et al., 2015). We therefore modelled the rate of death due to autoimmunity as being proportional to CRISPR-Cas reactivity and to the number of bacteria. In addition, the cost of autoimmunity can be modulated by a parameter representing the propensity of CRISPR-Cas for autoimmunity, independently of its reactivity: biologically, this can be understood as the propensity for the bacterial chromosome to be targeted by CRISPR-Cas, for example because of the density of the bacterial chromosome in potential protospacers or in sequences inhibiting acquisition (like Chi sites (Levy et al., 2015)) or because the bacterial CRISPR-Cas system possesses an efficient self/non-self distinction mechanism.

The rate of death due to autoimmunity is therefore implemented as the following term that is subtracted from Equation 1 and 2:

$$Autoimmunity = \alpha \times \text{Propensity} \times B, \text{ with B being S or } R_i$$

Bacterial spacer diversity

As a quantitative measure of the bacterial spacer diversity, we calculated the Simpson index from the frequency of each bacterial genotype p_i:

$$^2D = \frac{1}{\sum_{i=1}^{n_s} p_i^2} \tag{5}$$

Fitness calculation

We calculate relative fitness of two competing bacterial strains from their initial and final frequencies, p_i and p_f using the following formula:

$$\text{Relative Fitness} = \frac{p_f(1-p_i)}{p_i(1-p_f)}, \text{ which with } p_i = 0.5 \tag{6}$$

$$= \frac{p_f}{(1-p_f)} \tag{7}$$

Model implementation

All simulations and analyses were implemented and conducted in the language for statistical computing R 3.5.1 (R Core Team, 2018). To implement the model (Equation 1–4) stochastically, we used the Gillespie algorithm in the package adaptivetau (Johnson, 2019).