Myoglobin primary structure reveals multiple convergent transitions to semi-aquatic life in the world's smallest mammalian divers

Kai He,[*,1,2,3] Triston G. Eastman,[1] Hannah Czolacz,[4] Shuhao Li,[1] Akio Shinohara,[5] Shin-ichiro Kawada,[6] Mark S. Springer,[7] Michael Berenbrink,[*,†,4] and Kevin L. Campbell[*,†,1]

[1]Department of Biological Sciences, University of Manitoba, Winnipeg, MN, Canada
[2]State Key Laboratory of Genetic Resources and Evolution, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming, Yunnan, China
[3]School of Basic Medical Sciences, Southern Medical University, Guangzhou, Guangdong, China
[4]Department of Evolution, Ecology and Behaviour, University of Liverpool, Liverpool, United Kingdom
[5]Department of Bio-resources, Division of Biotechnology, Frontier Science Research Center, University of Miyazaki, Miyazaki, Japan
[6]Department of Zoology, Division of Vertebrates, National Museum of Nature and Science, Tokyo, Japan
[7]Department of Evolution, Ecology, and Organismal Biology, University of California, Riverside, CA, USA

*Corresponding authors: Email: hekai2018@smu.edu.cn, Michael.Berenbrink@liverpool.ac.uk, or Kevin.Campbell@umanitoba.ca
†These authors contributed equally to this work

Abstract

Identifying the phylogenomic underpinnings of specialized phenotypes that fueled transitions into new adaptive zones is central to evolutionary biology but is often confounded by a fragmentary fossil record, morphological convergence, and unresolved phylogenetic relationships. The speciose mammalian order Eulipotyphla (e.g., moles, shrews, hedgehogs, solenodons) combines an unusual diversity of semi-aquatic, semi-fossorial, and fossorial forms that arose from terrestrial forbearers, yet the ecomorphological pathways leading to these lifestyles have been disputed for a century and more calling for novel approaches. Here we resolve previously intractable eulipotyphlan intra-family relationships and establish the net surface charge of the oxygen-storing muscle protein myoglobin—readily determined from its primary structure—as a molecular signature to trace ancient lifestyle transitions based on protein sequence alone. Our analyses confidently resolve fossorial habits having evolved twice in talpid moles and a reveal five independent origins of a semi-aquatic lifestyle in the order housing the world's smallest endothermic divers.

Introduction

A fundamental challenge of evolutionary biology is to understand the phylogenomic foundations of biochemical and physiological specializations that have enabled organisms to proliferate into new adaptive zones. While species of most mammalian orders generally occupy a single niche, members of a few mammalian lineages have radiated into a diverse range of environmental habitats. The order Eulipotyphla is one such assemblage, in that terrestrial forms repeatedly evolved into high-altitude, scansorial (climbing), fossorial (subterranean), semi-fossorial, and semi-aquatic niches (Burgin and He 2018; He, et al. 2017).

The fourth largest among the 27 mammalian orders after Rodentia, Chiroptera, and Cetartiodactyla, the eulipotyphlan clade consists of 527 recognized species of small insectivorous mammals distributed among four extant families (Burgin, et al. 2018): Erinaceidae (hedgehogs, gymnures, and moonrats), Soricidae (shrews), Talpidae (moles, shrew moles, shrew-like moles, and desmans), and Solenodontidae (solenodons). A fifth family (Nesophontidae, Caribbean 'island-shrews') only became extinct in the past 500 years following the arrival of Europeans in the West Indies (MacPhee, et al. 1999). Fossil evidence supports terrestrial habits as ancestral for Eulipotyphla, with shrew-like moles, erinaceids, and the majority of shrews

retaining morphological characteristics (e.g., slender legs and a long, pointed mouse-like snout;

fig. 1) effective for hunting and avoiding predation on land (Churchfield 1990; Nowak 1999).

Semi-aquatic moles and shrews possess a stream-lined body and limb adaptations for underwater

locomotion (Nowak 1999; Burgin and He 2018), with desmans having an outward appearance

more similar to that of semi-aquatic rodents (e.g., muskrat) than to that of any other

eulipotyphlan mammal (fig. 1). By contrast, fossorial moles have evolved powerful forelimbs

that are held out to the sides of their tube-shaped bodies and have large, spade-like hands facing

posteriorly and oriented vertically allowing them to be effective burrowers (Gorman and Stone

1990). Finally, semi-fossorial shrews, solenodons, and shrew moles exhibit specializations in-

between that of terrestrial and fossorial lineages, often retaining pointed mouse-like snouts yet

possessing forelimbs that are oriented downwards and backwards to facilitate digging (Gorman

and Stone 1990; Nowak 1999).

While the lifestyle of extant eulipotyphlan mammals is easily predicted based on external

morphological features alone (e.g., Woodman and Gaffney 2014; Sansalone, et al. 2016;

Sansalone, et al. 2019), definitive assignments of transitional evolutionary stages have remained

elusive due to an incomplete, fragmentary fossil record (Hickman 1984; Sánchez-Villagra, et al.

2006; Piras, et al. 2012; Hooker 2016). Additionally, because morphological specializations

coincide with lifestyle across the different clades of eulipotyphlan mammals, it has been difficult

to discern whether fossorial and semi-aquatic specializations found in Eurasian and North

American moles arose in a common ancestor or are due to convergent evolution (Schwermann

and Thompson 2015; He, et al. 2017). As a result, long-standing questions persist regarding the

evolutionary pathways leading to the diverse ecomorphological variation seen in eulipotyphlan

mammals.

For example, examinations of shoulder anatomy (Campbell 1939) and skeletal musculature

(Whidden 1999) indicated that the shoulder and humeroclavicular joint morphology of fossorial

talpid species aligned more closely with semi-aquatic locomotion, suggesting that the ancestor of

Talpinae moles was semi-aquatic, which we here term the 'aquatic mole' hypothesis.

Conversely, Reed (1951) analyzed anatomical features from four talpid genera and proposed that

semi-fossorial and fossorial forms evolved from a terrestrial ancestor and that the semi-aquatic

lifestyle of desmans was secondary to a semi-fossorial phase, a view also shared by Grand et al.

(1998). Skeletal and fossil evidence also suggested that a period of strictly fossorial life preceded

the invasion of semi-aquatic habits by the ancestors of star-nosed moles (*Condylura cristata*)

(Hickman 1984; Grand, et al. 1998; Sanchez-Villagra, et al. 2006; Piras, et al. 2012; Hooker

2016), though Sansalone et al. (2016) proposed that the semi-aquatic lifestyle of this genus

instead represents an autapomorphy from a semi-fossorial (shrew mole-like) ancestor.

Adaptive morphology and behavioural/physiological specializations for diving have also

been well documented for members of the family Soricidae (Hutterer 1985; Catania, et al. 2008;

Burgin and He 2018), though uncertainties regarding the evolution of semi-aquatic habits in this

clade are also pervasive. For example, semi-aquatic members of four genera, *Sorex* (North

American water shrews), *Neomys* (European water shrews), *Chimarrogale* (Asian water shrews),

and *Nectogale* (Elegant water shrews) show a graded series of progressively greater

morphological adaptations for underwater foraging (Hutterer 1985), yet phylogenetic analyses

strongly refute a common ancestry of aquatic exploitation (He et al. 2010). Ancestral-state

reconstructions and the placement of *Chimarrogale* + *Nectogale* as sister-taxa instead provided

support for semi-aquatic habits having evolved three times (He et al. 2010), as opposed to four

convergent invasions (Churchfield 1990). However, based on associated fauna/flora and

paleoecological indicators of one of the oldest fossil nectogaline genera (*Asoriculus*), it has been

hypothesized that a shared Late Miocene ancestor of *Neomys*, *Chimarrogale*, and *Nectogale* may

have been adapted to humid environments with permanent open water (Rofes and Cuenca-

Bescos 2006), thus allowing the possibility that semi-aquatic habits evolved only twice in

shrews. Taken together, potential morphological convergence and an incomplete fossil record

have precluded reliable conclusions to be made regarding the evolutionary pathways underlying

the origins of the diverse phenotypes of eulipotyphlan mammals.

The monomeric oxygen-binding protein myoglobin, which plays an essential role in O_2

storage and facilitates O_2 diffusion within the heart and skeletal muscle of most vertebrates,

provides an intriguing, objective molecular target to address this question. Indeed, the O_2 storage

function of myoglobin has long been known to contribute integrally to the mammalian dive

response, with maximum active submergence times of birds and mammals being strongly

correlated to muscle myoglobin concentrations (Noren and Williams 2000; Lestyk, et al. 2009;

Ponganis, et al. 2011). Links between the molecular evolution of this molecule and aquatic life

had already been suspected for a half century or more (Scholander 1962; Romero-Herrera, et al.

1973), leading to the speculation that "there might be functional reasons, perhaps associated with

diving" underlying the presence of parallel amino acid replacements in the myoglobins of seals

and cetaceans (Romero-Herrera, et al. 1978). Mirceta et al. (2013) extended this work to reveal

that maximal muscle myoglobin concentration was mechanistically linked to myoglobin net

surface charge (Z_{Mb}) in mammals via adaptive changes in primary structure, with convergent

increases in Z_{Mb} found in members of all eight lineages with an extended aquatic/semi-aquatic

evolutionary history. This trait presumably represents an adaptive response to combat the general

propensity for proteins to precipitate at high concentration, thereby allowing for advantageous

elevations of this muscle O_2 store without deleterious self-aggregation (Mirceta, et al. 2013). An

increase in Z_{Mb} may also reduce the tendency of the myoglobin peptide (apomyoglobin) to

partially unfold prior to heme insertion, thus limiting hydrophobic interactions and making the

formation of an aggregate to be energetically unfavourable (Samuel, et al. 2015; Isogai, et al.

2018). It should be stressed that increases in myoglobin concentration afforded by an elevated

Z_{Mb} presumably represent an evolutionary trade off because there is less room for contractile

units (sarcomeres) and organelles (e.g., mitochondria) in muscle tissue as more energy and space

is allocated to myoglobin. Therefore, an increased Z_{Mb} only offers an advantage to species

experiencing hypoxic episodes as for example occurs during breath-hold dives when access to

environmental air is impeded. Importantly, the finding that Z_{Mb} and maximal myoglobin

concentrations are only slightly elevated within mammals that live at high elevations or have

fossorial lifestyles (Mirceta, et al. 2013) suggests that this locus may be useful to discern past

semi-aquatic versus high-altitude/fossorial evolutionary histories within eulipotyphlan mammals.

Support for the above contention is provided by the molecular modelling and phylogenetic

reconstruction of Z_{Mb} in six eulipotyphlan species (Mirceta, et al. 2013), which revealed

convergent elevations in Z_{Mb} in both semi-aquatic taxa included in the dataset—the American

water shrew (*Sorex palustris*) and the star-nosed mole. While myoglobin concentration data is

sparse for eulipotyphlan mammals, available data indicates that myoglobin levels are also higher

in these two species relative to non-aquatic forms (McIntyre, et al. 2002; Gusztak 2008).

Notably, Z_{Mb} was reconstructed to secondarily decline in the branch containing the single strictly

fossorial talpid species (Mirceta, et al. 2013), thus providing tentative support for the 'aquatic

mole' hypothesis. The sparse taxon sampling in the cited study (two moles, three shrews, and a

hedgehog), however, did not account for the broad phylogenetic and ecomorphological diversity

within these families. Additionally, current phylogenetic hypotheses for Eulipotyphla lack

definitive resolution below the family level (He, et al. 2010; He, et al. 2017) thereby precluding

reliable ancestral reconstructions. To overcome these shortcomings, we used a capture

hybridization approach to target coding sequences of myoglobin together with 25 tree-of-life

genes from 61 eulipotyphlan DNA libraries (44 moles, 11 shrews, 5 hedgehogs, and 1

solenodon) that included representatives from all seven recognized semi-aquatic genera within

this order. We then tested whether Z_{Mb} is elevated within members of all living genera of semi-

aquatic Eulipotyphla, and if so, whether there is a significant correlation between this lifestyle

and Z_{Mb}. Having shown that these important conditions are met, we then traced Z_{Mb} across

eulipotyphlan evolutionary history, thereby allowing us to determine when and how many times

semi-aquatic specializations for increased dive durations evolved in both shrews and moles, and

evaluate the merit of the 'aquatic mole' hypothesis versus alternative evolutionary scenarios of

talpid lifestyle evolution.

Results

Phylogenetic relationships within Eulipotyphla

To obtain mammalian tree-of-life genes (Meredith, et al. 2011) for phylogenetic estimation,

we conducted in-solution probe-hybridization for segments of 25 single-copy genes from 61

eulipotyphlan DNA libraries (supplementary table S1). Twenty-three of these loci were

efficiently captured, with usable sequence obtained from 51 to 61 libraries per locus, equivalent

to a ~95% (1330/1403) success rate (supplementary table S2). By contrast, probes for two gene

segments only successfully hybridized to 21 and 15 of the 61 libraries, respectively, and were

subsequently not included in the phylogenetic analyses. Specifically, we failed to capture the

prepronociceptin (PNOC) gene for any shrew and many of the talpid species (supplementary

table S2). Probes for the interphotoreceptor retinoid binding protein (IRBP) also failed to retrieve

any sequence from the solenodon and 37 of the 38 non-uropsiline talpid specimens. Notably, the

putative 234 bp IRBP fragment recovered from True's shrew mole (*Dymecodon pilirostris*)

contained gaps and premature stop codons, suggesting this gene was inactivated in the common

ancestor of the subfamily Talpinae. Similarly, the Chinese mole shrew (*Anourosorex squamipes*)

IRBP sequence contained premature stop codons and was presumed to be non-functional. After

incorporating orthologous tree-of-life sequence data from ten additional eulipotyphlan specimens

and five outgroup species downloaded from GenBank (see Methods for details), our 76 specimen

dataset (71 eulipotyphlans) resulted in a final alignment of 39,414 bp.

To estimate evolutionary relationships, we used maximum likelihood (RAxML) and

Bayesian (BEAST) approaches on the concatenated alignment, and coalescent-based species tree

methods (ASTRAL-III, *BEAST) on the 23 individual gene trees. These analyses resulted in

highly congruent topologies (fig. 2 and supplementary figs. S1, S2, S3), with family level

relationships corresponding to those obtained in recent eulipotyphlan (Brace, et al. 2016;

Springer, et al. 2018) and mammal wide studies (Meredith, et al. 2011; Esselstyn, et al. 2017).

Higher-level relationships within Soricidae and Erinaceidae also corresponded closely with

previous studies on these families (He, et al. 2010; He, et al. 2012). Principal among these is the

non-sister relationship of the Nectogalini water shrew clades *Neomys* and *Chimarrogale* +

Nectogale, as the latter was strongly supported as sister to terrestrial *Episoriculus* by RAxML

(bootstrap support [ML-BS] = 97; supplementary fig. S1), BEAST (posterior probability [PP] =

1.0; fig. 2), ASTRAL-III (coalescent bootstrap support based on gene-wise resampling [C-BS] =

76; supplementary fig. S2), and *BEAST (posterior probability [C-PP] = 0.99; supplementary

fig. S3). By contrast, previously intractable relationships within Talpidae are now resolved, with

the interrelationships among fossorial and semi-aquatic clades consistently recovered (fig. 2 and

supplementary figs. S1, S2, S3). Specifically, desmans are placed sister to a clade containing

shrew moles, star-nosed moles, and Eurasian fossorial moles (BS = 100, PP = 1.0, C-BS = 87, C-

PP = 0.97), with Condylurini and Talpini recovered as sister lineages (BS = 72, PP = 1.0, C-BS

=52, C-PP = 0.93). North American fossorial Scalopini moles are also supported as sister to all

other Talpinae moles with high support scores (BS = 97, PP = 1.0, C-BS = 87, C-PP = 0.97).

Accordingly, a sister group relationship of fully fossorial Scalopini and Talpini moles was

statistically rejected by the Shimodaira–Hasegawa (SH) test ($p < 0.01$, supplementary table S3).

The only minor incongruences among phylogenies was the position of the tribe Soricini and of

the Taiwanese brown-toothed shrew (*Episoriculus fumidus*) within Nectogalini in the ASTRAL

tree (fig. 2 and supplementary figs. S1, S2, S3).

Myoglobin primary structure

Complete myoglobin coding sequences (465 base pairs including initiation and stop codons)

were obtained for 55 eulipotyphlan species (38 moles, 14 shrews, 2 erinacids, and 1 solenodon)

using capture-hybridization, transcriptome sequencing, genome mining, and PCR approaches.

Conceptual translations revealed the expected 153 amino-acid peptides in most cases. However,

the translated myoglobin proteins of the Chinese mole shrew, both members of the genus

Scaptonyx, and the two desman genera are only 152 amino acids in length. In every case the

shorter myoglobin sequences are the result of 3 bp deletions in exon 3 that corresponded to

residue position 121 (supplementary table S4) and shorten the loop between helices G and H

from 6 to 5 residues. These deletions were confirmed for each lineage using at least two of the

three sequencing approaches noted above. To our knowledge, a similar deletion was previously

only known for three bird of prey species (Enoki et al. 2008), but we have detected it also in the

predicted myoglobin sequences of the draft genomes of several burrowing species of the order

Rodentia (data not shown), notably including the fully fossorial Transcaucasian and Northern

mole voles (*Ellobius lutescens* and *E. talpinus*, respectively; Mulugeta, et al. 2016).

Myoglobin homology modelling

Homology modelling of myoglobin structure was conducted for one extant species from

each of the five diving lineages together with that predicted for the last common eulipotyphlan

ancestor based on ancestral sequence reconstruction (see next section). This analysis confirmed

that all charge-changing substitutions were located at the solvent-exposed surface of the protein

(supplementary table S4). These comparisons further suggested that deletion of position 121 in

the loop between helices G and H in the myoglobin of the Russian desman has minimal effect on

the tertiary structure of the protein (supplementary fig. S5). In addition to their effect on Z_{Mb},

some of the charge-increasing substitutions were associated with a complex re-arrangement of

the network of salt bridges in the tertiary structure of the protein. Thus, for example, during the

evolution of the Russian desman, arguably the most aquatic of extant Eulipotyphla, the two

neighbouring substitutions Gln²⁶→Arg²⁶ (charge-increasing) and Glu²⁷→Asp²⁷ (charge-neutral)

in the B-helix allow for the formation of a new intra-helical salt bridge between Arg²⁶ and Asp²⁷,

while at the same time breaking the inter-helical salt bridge between Glu²⁷ and Lys¹¹⁸ in the B-

and G-helices, respectively (fig. 3). However, charge-decreasing Asn³⁵→Asp³⁵ followed by

charge-increasing Gln¹¹³→Lys¹¹³ subsequently again tethered the B- and G-helices to each other

by the salt bridge Asp³⁵- Lys¹¹³. Further, the removal of a negative charge by the substitution

Asp⁴⁴→Ala⁴⁴ destroyed a salt bridge between Asp⁴⁴ and Lys⁴⁷ in the CD-corner of the protein,

thereby potentially affecting the flexibility of the loop between the C- and D-helices (fig. 3).

However, a detailed assessment of any changes in the folding stability of the proteins that are

associated with the identified charge-changing substitutions, let alone with any charge neutral

replacements (see, e.g., Isogai, et al. 2018), is difficult and beyond the scope of this study, not

least because of the potentially opposing effects of salt bridge formation and the associated

desolvation of charges for the folding stability of proteins (for discussion see Bosshard, et al.

2004).

Electrophoretic mobility and ancestral reconstruction of Z_{Mb}

Without exception, the modelled Z_{Mb} values of extant semi-aquatic taxa (2.07 to 3.07) were

substantially higher than those of terrestrial Eulipotyphla (-0.46 to 0.63), with fossorial species

generally exhibiting intermediate Z_{Mb} values (typically 1.07; fig. 4A and supplementary fig. S6).

To assess the reliability of our Z_{Mb} determinations, we measured the electrophoretic mobility of

the myoglobin band of muscle extracts from two semi-aquatic, two strictly fossorial, and one

terrestrial eulipotyphlan species (fig. 4B). The close correspondence between the two variables

validates Z_{Mb} as a molecular marker for inferring present and past semi-aquatic habits in

Eulipotyphla.

Our myoglobin nucleotide and amino acid gene trees retrieved few well-supported

phylogenetic relationships (especially at deeper nodes), and no compelling evidence of an

evolutionary history that deviated from the species trees (supplementary figs. S8A, B), which

would potentially result in erroneous ancestral reconstructions (Hahn and Nakhleh, 2016). We

thus conducted a maximum likelihood ancestral amino acid sequence reconstruction using the

species tree in fig. 1 as the phylogenetic backbone and the best fitting model of the Dayhoff

amino acid substitution matrix with a gamma distribution of rate variation among sites. This

analysis yielded highly supported ancestral amino acid identities across all 153 residues and 58

internal nodes of the species tree (supplementary fig. S7A), which was presumably due to our

dense taxon sampling and the relatively highly conserved primary structure of mammalian

myoglobins. Of the 8,874 (= 58 × 153) reconstructed ancestral sites on the species tree, 8,799

(99.15%) had maximal probabilities of reconstructed amino acid identities of $p > 0.95$ under the

given phylogeny and amino acid substitution model. In 50 cases (0.55%), alternatively

reconstructed amino acids with $p > 0.05$ were of the same charge and thus did not affect the

calculated Z_{Mb} values. Only in 25 cases (0.28%), one or more alternatively reconstructed amino

acids with $p > 0.05$ carried a different charge from the most probable amino acid at that site.

However, even in those cases this usually had only a minimal effect on Z_{Mb} (± 0.11; 3 sites), or

the summed probabilities of alternative charge states were comparatively small ($p \leq 0.21$; 23

sites) such that we regard both the results of the ancestral sequence reconstruction and of the

resulting values of Z_{Mb} as robust. This is supported by generally congruent results obtained by

codon-based ancestral sequence reconstructions (supplementary fig. S7B), which primarily

deviated from the above empirical amino acid matrix-based method at some deeper nodes, where

a higher number of negatively charged amino acid residues (and thus anomalously low values for

Z_{Mb}; e.g., -1.93) were reconstructed relative to those obtained for extant mammalian species

(Mirceta, et al. 2013; this study). Consequently, the following discussion largely focuses on the

Z_{Mb} reconstruction using the empirical Dayhoff substitution matrix (supplementary fig. S7A),

which moreover—in contrast to the codon-based analysis (supplementary fig. S7B)—includes

the effects of purifying selection acting on replacements with dissimilar amino acid properties.

Ancestral Z_{Mb} estimates arising from the best fitting amino-acid based reconstruction model

indicated that the most recent common ancestor of Eulipotyphla displayed a Z_{Mb} of 0.18. Within

Soricidae, distinct increases in Z_{Mb} were found on the branches leading to the three semi-aquatic

clades (fig. 5). Specifically, the Z_{Mb} of the American water shrew (*Sorex palustris*) branch

increased from -0.46 to 2.48 and was characterized by three integral (+1) charge increasing

residue replacements (Asp^{122}→Asn^{122}, Ser^{132}→Lys^{132}, and Glu^{136}→Gln^{136}; supplementary fig.

S4A). The other three genera of water shrews reside in the tribe Nectogalini, the stem branch of

which evolved a single charge increasing substitution (Glu^{27}→Ala^{27}) and thus had a Z_{Mb} of 1.07

(fig. 5). The European water shrew branch (*Neomys fodiens*) subsequently acquired two charge

increasing substitutions (Asn^{12}→Lys^{12}, Asp^{53}→Ala^{53}; Z_{Mb}=3.07; supplementary fig. S4B), while

the common ancestor of *Nectogale + Chimarrogale* evolved a separate charge increasing

replacement (Asp^{44}→Ser^{44}; Z_{Mb}=2.07; supplementary fig. S4C). By contrast, members of the

terrestrial genus *Episoriculus*, which are nested between the semi-aquatic nectogaline lineages,

exhibited secondary reductions in Z_{Mb} towards neutrality via different residue substitutions in

each case (supplementary fig. S7A). Importantly, similar results were obtained when the above

amino-acid based reconstructions were re-run on an alternative topology (supplementary fig. S3)

that supported the monophyly of *Episoriculus* (supplementary fig. S7C).

Within the Talpidae, Z_{Mb} exhibited an increase from an ancestral value of 0.07 to 1.07

(Ser^{132}→Lys^{132}) in the stem Talpinae branch (fig. 5), with the probabilities of the identities of

reconstructed amino acids at this node under the given tree and substitution model being 1.00 for

all 153 sites (supplementary fig. S7A). With few exceptions, these values remained highly

conserved in the clades containing either semi-fossorial or fossorial taxa (fig. 5 and

supplementary figs. S7A, B). By contrast, Z_{Mb} increased to >2 in members of both semi-aquatic

lineages—star-nosed moles and the desmans (fig. 3). For the star-nosed mole branch this entailed

two charge increasing substitutions (Asp^{53}→Gly^{53}, Gln^{128}→Lys^{128}) and one charge decreasing

substitution (Gly^{129}→Glu^{129}; supplementary figs. S4D and S7A). The ancestral branch of the

desmans also acquired two charge increasing replacements (Gln^{26}→Arg^{26}, Asp^{44}→Ala^{44})

together with one charge decreasing (Asn^{35}→Asp^{35}) replacement, with Z_{Mb} further being

elevated in the Russian desman branch via the acquisition of an additional charge increasing

substitution (Gln^{113}→Lys^{113}; Z_{Mb}=3.07; figs. 3, 5). Notably, while the same residue positions are

occasionally recruited in the charge altering replacements of semi-aquatic taxa, the derived

residues are different in all cases (supplementary table S5).

To evaluate the above reconstructions of semi-aquatic lifestyles based on Z_{Mb}, we coded each

species as semi-aquatic or non-aquatic and estimated ancestral lifestyles using both maximum

parsimony and threshold models. Although both of these latter analyses suggested that semi-

aquatic lifestyles evolved five times independently, this result was not strongly supported by the

threshold model (supplementary fig. S9A). For example, the posterior probabilities of the most

recent common ancestor of *Desmana + Galemys* and *Chimarrogale + Nectogale* being semi-

aquatic was only 0.85 and 0.75, respectively, while the posterior probability for the most recent

common ancestor of Nectogalini being semi-aquatic was 0.40. To account for the alternative

placement of *E. fumidus*, we repeated this analysis using the results of the *BEAST species tree

(supplementary fig. S3). The maximum parsimony reconstruction yielded two equi-parsimonious

ancestral reconstructions of semi-aquatic lifestyle in nectogaline shrews, encompassing a single

origin at the base of nectogaline shrews with a secondary loss at the base of the *Episoriculus*

clade (supplementary fig. S9B). By contrast, the threshold model only weakly supported two

independent origins of a semi-aquatic lifestyle in *Neomys* and *Nectogale + Chimarrogale*.

Lifestyle correlation analyses

As a final test regarding the reliability of using Z_{Mb} to predict ancient semi-aquatic lifestyles, we first assigned species as semi-aquatic or non-aquatic (see fig. 2 and supplementary table S1 for lifestyle assignments), and used a threshBayes analysis to estimate covariances between Z_{Mb} and a semi-aquatic lifestyle. This analysis revealed a strong correlation between Z_{Mb} and aquatic adaptation (correlation coefficient r = 0.78, 95% highest posterior density [HPD] = 0.48-0.93; supplementary fig. S10A). Conversely, threshBayes analyses did not support a correlation between Z_{Mb} and adaptations for digging, or between Z_{Mb} and a fully fossorial habit, when terrestrial and semi-aquatic eulipotyphlan species were included (supplementary figs. S10B, C). When applying threshBayes to subsets of habits that included only terrestrial and fully fossorial species, or terrestrial and burrowing species, weak correlations between Z_{Mb} and fossoriality (r = 0.55) and digging habits (r = 0.32) were revealed, but not significantly supported (i.e., 95%HPD overlaps with 0; supplementary figs. S10D, E). Importantly, Z_{Mb} comparisons between semi-aquatic and terrestrial, and between semi-aquatic and fossorial habits (supplementary figs. S10F, G) were both significant.

Discussion

The phylogenetic estimates constructed from our comprehensive tree-of-life gene set provide a robust framework to interpret ecomorphological evolution within Eulipotyphla. The close correspondence of our concatenation and coalescent phylogenetic topologies not only support key findings of previous studies—i.e., the monophyly of shrew moles, the non-monophyly of nectogaline water shrews (Whidden 2000; He, et al. 2010; He, et al. 2017)—but finally puts to rest the long hypothesized monophyletic origin of the fully fossorial tribes Talpini and Scalopini, which have been routinely grouped together based on morphological data (see, e.g., Whidden 2000; Piras, et al. 2012; Schwermann and Thompson 2015; Hooker 2016; Sansalone, et al. 2019). Results of the present study thus provide compelling evidence that the extreme anatomical specializations for subterranean life evolved convergently in these two tribes. The molecular phylogenetic position of the amphibious desmans and semi-aquatic/fossorial star-nosed mole are also finally resolved (fig. 2), with the latter placed sister to Talpini, a relationship not supported by morphological-based hypotheses (see, e.g., Whidden 2000; Motokawa 2004; Sanchez-Villagra, et al. 2006).

Previous studies have failed to reach consensus on the lifestyle evolution of Eulipotyphla.

Here we show that ancestral sequence reconstruction of myoglobin primary structure and Z_{Mb}

modelling, with their well established mechanistic and biophysical underpinnings, outperform

discrete, two character-state lifestyle reconstructions based on maximum parsimony and

threshold approaches. These attributes, together with demonstrated links between Z_{Mb} and

maximal Mb concentration, and hence muscle oxygen storage capacity (Mirceta, et al. 2013;

Berenbrink 2020), provide strong support that this quantitative metric is well suited to resolve

long-standing questions regarding lifestyle evolution within Eulipotyphla. For example, despite

being less specialized for aquatic life than marine mammals, the strong positive correlation

between Z_{Mb} and semi-aquatic specialization indicates that Z_{Mb} is a powerful marker to identify

secondary aquatic transitions in even the world's smallest mammalian divers. Although a strong

correlation was not recovered between Z_{Mb} and strictly fossorial habitation, Z_{Mb} is highly

conserved in fossorial Scalopini and Talpini, with 23 of 25 species exhibiting a value of 1.07.

This conservation is presumably driven by selective pressures to maintain moderately elevated

tissue myoglobin levels to help foster burst burrowing activities in their hypoxic underground

environment. By contrast, the Z_{Mb} of terrestrial species in our dataset were consistently close to

neutrality, though many lineages exhibited clear signals of Z_{Mb} fluctuation over time

(supplementary figs. S7A, B, C). For example, the shrew gymnure (*Neotetracus sinensis*) branch

evolved nine charge altering residue substitutions since its split from European hedgehogs

(*Erinaceus europaeus*), including the charge inversion $Asp^{126} \rightarrow Lys^{126}$ that increases Z_{Mb} by +2.

Similarly, the Taiwanese brown-toothed shrew branch (*Episoriculus fumidus*) has fourteen

charge altering residue substitutions, including one negative-to-positive ($Glu^{109} \rightarrow Lys^{109}$) and two

positive-to-negative charge inversions ($Lys^{102} \rightarrow Glu^{102}$ and $Lys^{132} \rightarrow Glu^{132}$). These observations

are consistent with a stochastic evolutionary process operating under purifying selection. Charge

fluctuation is also apparent on the Hispaniolan solenodon (*Solenodon paradoxus*) branch, with

the moderately elevated Z_{Mb} value (1.85) in line with their terrestrial/burrowing lifestyle (Nowak,

1999). Myoglobin sequence data from additional extant/extinct members of this family (e.g.,

Cuban solenodons) together with that from the recently extinct terrestrial/fossorial Antillean

family Nesophontidae (MacPhee, et al. 1999)—which is placed sister to solenodons (Brace et al.

2016)—may help resolve the life history evolution of this poorly understood insectivore clade.

Similar, albeit less pronounced charge fluctuations are evident in several fossorial mole branches

(as well as the stem *Condylura* and desman branches), suggesting that the single fossorial Z_{Mb}

outlier (hairy-tailed mole, *Parascalops breweri*; $Z_{Mb} = 2.07$) may also represent stochastic

variation that has not yet been selected against and that presumably will return to an ecologically

normalized value over evolutionary time.

Our results indicate that charge increases in Z_{Mb} to >2 is essential for members of

eulipotyphlan mammals to successfully exploit a semi-aquatic lifestyle. Because Z_{Mb} of the most

recent common ancestor of Talpinae was confidently estimated to be below this value (fig. 5),

our results do not support the "aquatic mole" hypothesis which posits that a semi-aquatic stage

predated the invasion of fossorial habits in talpid moles (Campbell 1939; Whidden 1999). This

conclusion is supported by the results of our lifestyle reconstructions, which uniformly revealed

that stem Talpinae had a low probability of semi-aquatic habits. These findings are instead

consistent with the interpretation that fossorial forms evolved directly from terrestrial/semi-

fossorial ancestors, without passing through a semi-aquatic phase (Reed 1951; Hickman 1984;

Grand, et al. 1998). The interpretation of a semi-fossorial ancestry for early non-uropsiline

(Talpinae) moles is further supported by our finding of a presumed inactivation/deletion of IRBP

on this branch (supplementary Table S2). This eye-specific locus has been shown to be

inactivated/lost in numerous mammalian lineages that inhabit subterranean and other dim-light

niches (Emerling and Springer 2014), and also appears to be non-functional in solendons and the

Chinese mole shrew (this study). The Z_{Mb} charge increase (to 2.07) in the stem desman branch,

paired with a lack of pronounced forelimb specializations for digging (He, et al. 2017) and a

basal placement among non-scalopine Talpinae (fig. 2), also chimes with Reed's (1951)

suggestion that the semi-aquatic lifestyle of this clade was secondary to a semi-fossorial phase.

By contrast, semi-aquatic/fossorial star-nosed mole exhibits prominent morphological

adaptations for burrowing and is placed sister to fossorial Talpini, consistent with Grand et al.'s

(1998) hypothesis that this lineage passed through a specialized fossorial stage prior to invasion

of the semi-aquatic niche. However, a semi-fossorial—as opposed to a fully fossorial—ancestry

for *Condylura* (Sansalone, et al. 2016) cannot be excluded based on the pattern of Z_{Mb} evolution

on this branch.

The charge elevation to 1.07 in stem Nectogalini shrews is enticing, as it temporally

corresponds to the early Miocene fossil *Asoriculus*, which was theorized to have inhabited wet

environments though was unlikely to have been an efficient semi-aquatic predator (Rofes and

Cuenca-Bescos 2006). The single charge increasing substitution that evolved at this stage

($Asp^{27} \rightarrow Ala^{27}$) is retained in the three semi-aquatic nectogaline genera and presumably

facilitated the adaptation of early *Neomys* and *Chimarrogale + Nectogale* for aquatic food

resources starting some 15 and 10 million years ago, respectively. The presence of separate

charge increasing replacements within the evolutionary branches of each of the three

Episoriculus species also opens the possibility of additional semi-aquatic 'experiments' in this

genus. Regardless, independent reductions in Z_{Mb} to neutrality or below for each of the latter

extant species, which today inhabit damp areas in vegetated environments (Nowak 1999), is

consistent with convergent reversions of *Episoriculus* lineages to a predominantly terrestrial

foraging habit.

Our results provide strong evidence that the exploitation of semi-aquatic habits by extant

shrews and talpids occurred at least five times, and was accompanied by convergently evolved

charge-increasing substitutions at different surface sites on the myoglobin protein in each case

(figs. 3 and 5 and supplementary figs. S4, S7). This finding provides additional support for our

contention that adaptive increases in Z_{Mb} underlie the invasion of (semi-)aquatic niches by

mammals, presumably by allowing for higher skeletal muscle myoglobin concentration (Mirceta,

et al. 2013; Berenbrink 2020). This elevated Z_{Mb} presumably underlies the elevated O_2 reserves

in muscle of star-nosed moles and American water shrews compared to non-aquatic relatives

(McIntyre, et al. 2002; Gusztak 2008), and likely contributes to the extended dive times and

remarkable underwater foraging efficiency of these species (Catania, et al. 2008). The increase in

Z_{Mb} must be particularly important for semi-aquatic soricine shrews due to allometic

considerations that have resulted in extremely high muscle mitochondrial contents (which may

comprise up to 45% of the cell volume; Weibel 1985) and mass-specific tissue O_2 requirements

that may be >100-fold higher than those of large-bodied marine mammals (Butler 1998; Gusztak,

et al. 2005). It is notable that the highest Z_{Mb} values were found for Russian desmans (3.07) and

European water shrews (3.07), consistent with the exceptional diving abilities of these species

(Vogel 1998; Ivlev 2010), and it is predicted that these two species will also possess the highest

muscle myoglobin concentrations among Eulipotyphla.

Adaptive evolution of similar phenotypic and physiological features occurring in distantly

related lineages are not uncommon in mammals (Madsen, et al. 2001). For example, adaptive

radiations in Afrotheria and Laurasiatheria resulted in striking morphological convergence of

species occupying semi-aquatic (otter shrews vs. desmans) and subterranean (golden moles vs. true moles) habitats (Madsen, et al. 2001; Springer, et al. 2004). A unifying pattern underlying these and most other large-scale mammalian radiations over the past 200 million years is that they all involved ecological and locomotory diversification from ancestral lineages of small insectivores (Grossnickle, et al. 2019). The extensive radiation of small terrestrial Eulipotyphla into different adaptive zones, including four independent origins of venom systems in shrews and solenodons (Casewell, et al. 2019) and multiple independent invasions of shrews and moles to semi-aquatic, semi-fossorial, and subterranean environments that occurred on shallow timescales of only a few million years, further demonstrates the high intrinsic evolutionary potential of this Bauplan. Morphological, physiological, and even behavioral convergence have previously been identified within semi-aquatic eulipotyphlan species. For example, *D. moschatus*, *C. cristata*, and *S. palustris* can all detect prey scent while under water via the rapid exhalation and inhalation of air bubbles (Catania 2006; Catania, et al., 2008; Ivlev, et al. 2013), with the latter two species also being characterized by an elevated proton-buffering capacity in muscle (McIntyre, et al. 2002; Gusztak 2008). The results presented here add to this list of convergences, and indicate that semi-aquatic eulipotyphlans have evolved similar Z_{Mb} (and presumably elevated myoglobin concentration) phenotypes via the same selection pressure acting on different sites of the protein and by dissimilar combinations of amino acid substitutions (i.e., differential gains and losses of cationic and anionic residues, respectively). In other words, molecular adaptation of myoglobin towards life in a semi-aquatic environment is predictable at the protein level but underpinned by unpredictable genotypic evolution. As such, the phylogenomic analysis of myoglobin loci from tissue samples is not only able to provide insights into the lifestyles of rare and recently extinct mammalian species (e.g., museum specimens and subfossil material from the obscure Caribbean nesophontids), but also offers a useful tool to infer past semi-aquatic transitions based on myoglobin primary structure alone.

Materials and Methods

Eulipotyphlan taxon sampling and tree-of-life sequence data collection

Our taxon sampling of eulipotyphlan mammals included 44 talpids, 11 shrews, five erinaceids, and one solenodon (61 specimens encompassing 60 species). Note that this sampling incorporates talpid specimens from five putative 'cryptic lineages' (denoted by 'sp.', 'sp. 1', or

'sp. 2' in the figures and tables); for the purpose of this study, each of these genetically distinct

lineages are considered independent species. The tissue samples were from various resources

(supplementary table S1), with most tissue samples provided by co-authors from China, Japan,

Canada, and the USA. Voucher specimens collected by co-authors were deposited in the

Kunming Institute of Zoology (KIZ, China), the National Museum of Nature and Science

(NMNS, Japan) or kept in personal collections (A.S. and S.I.K.). Additional tissue samples were

obtained with permission from the National Museum of Natural History (USNM, USA), the

Burke Museum of Natural History and Culture (NWBM, USA), the Field Museum of Natural

History (FMNH, USA), and the New Mexico Museum of Natural History (NMMNH, USA).

For each specimen, we used a capture hybridization approach (Mason, et al. 2011; Horn

2012) to enrich segments of 25 mammalian tree-of-life genes (Meredith, et al. 2011) for

phylogenetic analyses. We first downloaded tree-of-life sequences from three eulipotyphlan

whole genome sequences available in GenBank (*Erinaceus europaeus*, *Sorex araneus*,

Condylura cristata), together with 60 bp of 5'- and 3'- flanking sequence for each target. We

then aligned each gene segment using MAFFT (Katoh and Standley 2013). The resulting

alignments were used to design 120 mer RNA probes (baits) that overlapped by 90 bp (4x tiling),

and collapsed any replicates with up to six mismatches (95% similarity) for each segment. For

example, if the 120-bp gene fragments from all species were 95% similar with each other, only

one probe was designed for this region, otherwise two or more probes were designed to cover the

heterogeneity. The probes were synthesized by Arbor Biosciences (Ann Arbor, MI, USA). As a

first step in DNA library construction we extracted total DNA from each specimen using a

Qiagen DNeasy Blood & Tissue Kit (Qiagen, Canada). The quality and quantity of each DNA

sample was measured using a Nanodrop 2000. We then sheared the total DNA into smaller

fragments using NEBNext dsDNA Fragmentase (New England Biolabs, Canada), and used this

as template to construct DNA libraries using a NEBNext Fast DNA Library Prep Set for Ion

Torrent kit (New England Biolabs, Canada). Each sample library contained a unique barcode

adapter (NEXTflex DNA Barcodes for Ion Torrent, BIOO Scientific, USA). We selected

libraries within the size range of 450-500 bp using a 2% E-gel on an E-Gel Electrophoresis

System (Invitrogen, Canada), and re-amplified the size-selected libraries using a NEBNext High-

Fidelity 2X PCR Master Mix (New England Biolabs, Canada). Finally, we purified the libraries

using Serapure magnetic beads, and measured DNA concentrations using a Qubit 2 Fluorometer
(Thermo Fisher Scientific, Canada).

We pooled up to four DNA libraries of similar quality and concentrations before
hybridization to avoid biased target captures (e.g., baits being used up by one sample).
Approximately 500 ng (100 ng to 1000 ng) pooled DNA library was used for each hybridization.
We conducted in-solution hybridization using a myBaits® custom target capture kit (Arbor
Biosciences, Ann Arbor, MI, USA) following the MYbaits user manual v3.0. The enriched
libraries were re-amplified and purified as above. We thereafter measured the DNA
concentration using a Qubit flourometer and pooled the enriched libraries for sequencing. The
libraries were sequenced using either v318 chips on an Ion Torrent Personal Genome Machine
(PGM) or an Ion PI Chip v3 via an Ion Proton Machine.

Sequence assembly
Ion Torrent sequencing technology is characterized by higher error rates than Illumina
(Jünemann, et al. 2013), and Ion Torrent platforms produce single-end (rather than pair-end)
reads. We therefore conducted comprehensive data cleaning and reconciliation procedures, and
selected software which could handle single-end sequencing data. The raw data were
automatically demultiplexed, trimmed, and converted to FASTQ format on the Torrent Suite
v4.0.2 (Thermo Fisher Scientific, Canada) after sequencing. Briefly, we trimmed contaminant
(adapters and barcodes) sequences with AlienTrimmer (Criscuolo and Brisse 2013) using
conservative parameters (-k 15 -m 5 -l 15 -q 0 -p 0). To remove poor quality data, we used the
DynamicTrim function of the software SolexaQA++ v3.1 (Cox, et al. 2010) to trim sequences
dynamically and crop the longest contiguous segment for each read. We set the probability value
to 0.01 (i.e., one base call error every 100 nucleotides) in this analysis. We removed duplicated
and near-duplicated reads for each sample as implemented in ParDRe using all default
parameters (Gonzalez-Dominguez and Schmidt 2016). Finally, we conducted data correction
using Karect, a multiple sequence alignment-based approach (Allam, et al. 2015), because this
software handles substitution, insertion, and deletion errors. The output files of Karect were used
for sequence assembly.

We de novo assembled the raw sequences for each sample using Abyss v2.0 (Simpson, et al.
2009), MRIA v4.0 (Chevreux, et al. 2004), and SPAdes v3.10 (Bankevich, et al. 2012), all of

which were designed for short read sequencing data. Abyss is able to use a paired de Bruijn

graph instead of a standard de Bruijn graph by specifying a k-mer size (K) and a k-mer pair span

(k). We set the K and k to 17 and 33, respectively, and set the maximum number of branches of a

bubble to 5 in our analyses. MIRA is based on a Smith-Waterman algorithm. We ran MIRA

using specific parameters including bases_per_hash = 31 and minimum_read_length = 35. The

SPAdes assembler is also based on a de Bruijn graph, and we set only one k-mer value of 33 for

analyses. It is known that merging different draft assemblies (i.e., reconciliation) could improve

the assembly quality (Zimin, et al. 2008). We therefore conducted reconciliation using Geneious

R11 (https://www.geneious.com). We concatenated the assembled draft contigs generated in

three assemblers into a list. We removed contigs shorter than 120 bps, and used the BBMap

dedupe function to remove duplicate contigs. We conducted assemblies using the Geneious

assembler to group draft contigs with a minimum overlap identity of 96% to a new contig.

Finally, all the new contigs and the leftover draft contigs were grouped into a contig list for

subsequent analyses.

Tree-of-life data analysis

We additionally extracted myoglobin coding sequences and the 25 mammalian tree-of-life

gene segments from seven publicly available eulipotyphlan genomes on GenBank using

PHYLUCE (Faircloth 2015) and Geneious R11: Indochinese shrew (*Crocidura indochinensis*),

gracile shrew-like mole (*Uropsilus nivatus*), Eastern mole (*Scalopus aquaticus*), Hispaniolan

solenodon (*Solenodon paradoxus*), European hedgehog (*Erinaceus europaeus*), common shrew

(*Sorex araneus*), and the star-nosed mole (*Condylura cristata*). Corresponding sequences from

five outgroup taxa were also mined: guinea pig (*Cavia porcellus*), horse (*Equus caballus*), cat

(*Felis catus*), pig (*Sus scrofa*) and bat (*Pteropus alecto*). PHYLUCE was originally developed

for ultra-conserved elements (UCE). We followed the "harvesting UCE loci from genomes"

protocol, but used the tree-of-life reference genes as probes instead of the original UCE probe

sets. We extracted genomic regions which were at least 75% similar to the tree-of-life reference

sequences. We also mapped the genomes to the tree-of-life references using Geneious with a

minimum overlap identity of 75%. These two packages were generally equally efficient at

capturing target genes, though in a few cases only one successfully captured the target genes

from the genome.

We used the above eulipotyphlan myoglobin and tree-of-life gene sequences to generate consensus sequences for the 61 specimens employed for the hybridization capture experiments (supplementary table S2). Briefly, the GenBank sequences were used as reference scaffolds to individually map the Ion Torrent generated reads of each sample using the Geneious "Map to Reference" function, and allowing for a mismatch of 35% per contig. This package conducts iterative mapping and outperforms many other algorithms by higher mapping rates and better consensus accuracy (Kearse, et al. 2012). Approximately 1-4 contigs from each sample were mapped to each gene reference. For the TTN gene segment, whose reference sequence was 4452 bp in length, as many as 10 contigs from each sample could be mapped to the reference. In addition, sequences of 19 nuclear gene segments obtained from 21 eulipotyphlan samples collected as part of previous studies (He, et al. 2014; He, et al. 2017), were also used for assemblies as above (and included in the final assemblies). Three shrew species (*Episoriculus umbrinus*, *Episoriculus fumidus*, and *Sorex bedfordiae*) for which we obtained myoglobin coding sequences via transcriptome sequencing (see below) were not included in our hybridization capture experiments. We thus downloaded the available tree-of-life genes from these species (*APOB*, *BRCA1*, and *RAG2*) on GenBank and included them in our analysis.

The resulting 25 tree-of-life gene segments were aligned separately using MAFFT. We then removed sequences shorter than 247 bp, and estimated gene trees using FastTree v2.1.5 to check for potential non-homologs for each gene. We also checked each alignment by eye, and removed ambiguous regions from the alignment. As segments of two genes (*IRBP*, *PNOC*) failed to hybridize to the majority of the DNA libraries, they were removed from subsequent analysis. The final 23 gene segment alignment (39,414 bp) included sequences from 76 samples (supplementary fig. S11). Because ASTRAL assumes no intra-locus recombination, we also tested recombination events using both RDP and GENECONV methods to examine each gene as implemented in RDP v5.5 (Martin, et al. 2015). When the program detected a signal of recombination, we checked UPGMA trees estimated using the two non-overlapping fragments and also examined the original alignment to see whether the signal was likely due to recombination or other evolutionary processes. We did not observe strong evidence of cross-species recombination in gene alignments (data not shown).

We estimated evolutionary relationships using several different approaches. We first constructed a summary-coalescent tree whereby we simultaneously conducted rapid bootstrap

analyses and searched for the best scoring maximum likelihood tree using RAXML v8.2
(Stamatakis 2014) for each gene alignment, allowing the program to determine the number of
bootstraps (-#autoMRE). We used *Cavia porcellus* as the root of the tree employing GTR+γ
during both ML searches and bootstrapping phases, and disabled the BFGS searching algorithm
for optimizing branch lengths and GTR parameters (--no-bfgs). We followed Simmons and
Kessenich's (2020) recommendation to remove dubiously supported clades and increase accuracy
of tree estimation. We used RAxML to estimate SH-like aLRT support values for the best-
scoring gene trees (-f j; supplementary table S7). Then we collapsed branches whose SH-like
aLRT support values equal zero using Newick utilities and Tree Graph 2 (Junier and Zdobnov
2010; Stöver and Müller 2010). Then we used the collapsed trees to estimate the coalescent
species tree using ASTRAL III v5.15.0 (Zhang, et al. 2018). Because gene-wise bootstrap could
provide more conservative support than site-wise bootstrap analyses (Simmons, et al. 2019), we
conducted gene-wise bootstrapping (--gene-only) instead of the typical site-wise bootstrapping.
We also allowed the program to explore a larger search space by adding extra bipartitions to the
search space (--extraLevel 2).

In addition to the summary coalescent analyses, we also constructed a concatenation-based
tree using the same dataset. We partitioned the alignment by gene, searched for the best-scoring
tree and conducted rapid bootstrapping under the GTR+γ model using RAxML as described
above. We used seven calibrations (supplementary text S1) and BEAST v2.5 (Bouckaert, et al.
2014) to estimate divergence times for all 76 samples as well as for the 60 species (55
eulipotyphlans and 5 outgroup species) for which complete myoglobin coding sequences were
obtained. For this analysis, we first partitioned the alignment by gene and used the bModelTest
package of BEAST2 to estimate the most appropriate substitution model for each gene
(Bouckaert and Drummond 2017). We used a relaxed clock model with lognormal distribution
for estimating the branch lengths, a birth-death model for the prior of the tree, and ran the
analysis for 100 million generations. We used *Cavia porcellus* as the outgroup to Laurasiatheria,
and also fixed the relationships of the other four outgroup species, because a biased sampling
toward the ingroup (i.e., Eulipotyphla) may lead to an inaccurate estimation of outgroup
relationships (Springer, et al. 2018). Secondly, we used the BModelAnalyzer package of
BEAST2 to determine the best models for each gene based on the results of the bModelTest

(supplementary table S6; Barido-Sottani, et al. 2018). We fixed the models of evolution based on

the results of BModelAnalyzer and re-ran BEAST using the same parameters described above.

Finally, we employed a multispecies coalescent model (*BEAST; Heled and Drummond

2010) as implemented in BEAST v2.5. We grouped the samples by species. We used linear and

constant root as the prior for the population model. Substitution model, tree model, and

calibrations were set as above. All gene trees and species trees are given in supplementary table

S8.

To examine whether alternative evolutionary hypotheses of life histories (e.g., a single

origination of fully fossorial lifestyle within Talpidae) could be statistically rejected, we

performed Shimodaira–Hasegawa (SH) tests. For these analyses, we constrained the

monophyletic relationships of: i) fully fossorial Talpini and Scalopini moles, ii) semi-aquatic

desmans and the star-nosed mole, and iii) semi-aquatic nectogaline shrew genera (*Chimarrogale*,

Nectogale, and *Neomys*), one at a time and estimated the maximum likelihood concatenation

trees using RAxML as described above (supplementary table S3). Then we computed the log

likelihood between the best scoring maximum likelihood tree and the constrained alternative

phylogenies as implemented in RAxML (-f H).

Myoglobin collection and analysis

We used four strategies to obtain myoglobin coding sequences (supplementary table S1). We

first extracted available eulipotyphlan myoglobin mRNA and gene sequences from GenBank.

The three coding exons were individually used both as templates for capture hybridization probe

design (see above) and to map the hybridization contigs/generate consensus sequences for each

exon. The 5'- and 3'- ends of introns were confirmed based on the GT-AG splice site rule. Of the

61 samples that we used for hybridization capture, complete coding sequence was obtained for

27 samples, partial myoglobin sequences were obtained for 30 samples, and no sequence

obtained for 4 samples (supplementary table S1).

To cross-validate the results of our hybridization experiments, fill sequencing gaps, and

extend our taxon sampling, we also PCR amplified and Sanger sequenced whole myoglobin

exons from existing DNA samples and/or employed transcriptome sequencing on additional

eulipotyphlan specimens. For the latter, we collected heart and lung samples from five shrews,

one shrew-like mole, and one gymnure (supplementary table S1). Tissues were preserved in

RNA*later* (Qiagen, China), and stored at -80C. Total RNA was extracted using a RNeasy Mini

kit (Qiagen, China), and mRNA subsequently enriched using immobilized oligo(dT). mRNA was

sheared and reverse transcribed to DNA. The cDNA libraries were purified and re-amplified

using PCR for de novo sequencing using a HiSeq X™ Ten Sequencing System. Approximately 6

Gb data were obtained for each sample. Experiments and sequencing were conducted by

BioMarker Co. (Beijing, China). We used FastQC v0.11.5 (Andrews 2010) to access sequence

quality, and trimmed adapter sequences using Trimmomatic v0.39 (Bolger, et al. 2014). We

conducted *de no* assembly using Trinity v2.4 with default parameters (Grabherr, et al. 2011).

Finally, primers for PCR were designed for conserved exon flanking regions from available

eulipotyphlan genomes, hybridization capture, and mRNA sequences, and were used for both

PCR and Sanger sequencing (supplementary table S9). These procedures resulted in complete

coding sequences being obtained for 55 eulipotyphlan species (supplementary table S4). We

estimated the myoglobin gene trees utilizing both nucleotide and amino acid sequences as

implemented in RAxML. We used Dayhoff+γ model (Dayhoff, et al. 1978) for the amino acid

gene tree estimation using the same settings described above.

Ancestral sequence reconstruction and homology modelling

We estimated ancestral myoglobin sequences for each node of a 60 species phylogeny that

utilized both DNA sequences and amino acids. Prior to analysis, the start (methionine) and stop

codons were removed from the alignment. As in our previous study (Mirceta, et al. 2013), we

performed maximum likelihood ancestral amino acid sequence reconstruction as implemented in

MEGA (Kumar, et al. 2018) using the Dayhoff+γ model that was obtained as the best-fitting

substitution model using the model test facility in MEGA-X. Prior to conducting the codon-

based analysis, we removed codons corresponding to residue position 121 (which was absent for

5 of the 60 species; supplementary table S4). We then used the PAML package CodeML (Yang

2007) as implemented in EasyCodeML (Gao, et al. 2019), and compared codon substitution

models (site models) including M0, M1a, M2a, M3, M7, M8 and M8a using likelihood-ratio

tests. We relied on the model with the highest likelihood (M8a). Because PAML does not take

account of insertion/deletion events (indels), and instead treats gaps as missing data, the ancestral

states of the gapped codon position 121 was reconstructed separately using a likelihood-based

mixture model as implemented in FastML (Ashkenazy, et al. 2012).

To assess the three-dimensional location and any secondary or tertiary structural implications

of amino acid replacements or insertions/deletions, we used the fully automated homology

modelling facilities of the SWISS-MODEL server (Waterhouse, et al. 2018) to build protein

structural models from the reconstructed ancestral primary structure of myoglobin in the last

common eulipotyphlan ancestor and from the primary structure of the sequenced myoglobins of

one species of each of the five semiaquatic lineages. Implications of the gapped position 121 on

the tertiary structure of myoglobin in the Russian desman compared to the last eulipotyphlan

ancestor were visualised in PyMol (The PyMOL Molecular Graphics System, Version 2.1.1,

Schrödinger, LLC).

Calculation of myoglobin net surface charge and electrophoretic mobility

We calculated Z_{Mb} as the sum of the charge of all ionizable groups in myoglobin at pH = 6.5

by modelling Mb primary structures onto the tertiary structure and using published, conserved,

site-specific ionization constants (McLellan 1984; Mirceta, et al. 2013). The reliability of

modelled Z_{Mb} values was assessed by determining the electrophoretic mobility of native

myoglobin bands at the same pH in muscle extracts of representative eulipotyphlan species and

the grey seal, *Halichoerus grypus*, as an example of a marine mammal. Approximately 0.2 g of

skeletal or cardiac muscle tissue from selected species, freed from any obvious fat or connective

tissue remnants and rinsed with homogenisation buffer to move any remaining blood, was

homogenized in 5 volumes of ice-cold 0.2 M MES buffer [2-(N-morpholino)ethanesulfonic acid]

adjusted to pH 6.5, using an Ultra-turrax T25 homogeniser for 10 s at first 9,500 rpm and then 3

times at 13,500 rpm, leaving samples to cool down between steps for 1 min on ice to avoid heat

denaturation of proteins. The homogenized muscle extracts were then centrifuged at 10,500 g (20

min at 4°C) and the supernatants stored at -80°C until further use. Electrophoretic mobility of

thawed muscle extracts was assessed in 9% polyacrylamide gels containing 0.3 M MES buffer

pH 6.5, using a Bio-Rad Mini-PROTEAN II gel system with 0.2 M MES pH 6.5 as the running

buffer at 100 V and room temperature for a minimum of 3 hours. Native myoglobin bands were

identified by their distinct red-brown colour before general protein staining with EZBlue™

(G104, Sigma-Aldrich). Electrophoretic mobility was assessed on digital gel images and

expressed as distance travelled relative to the grey seal myoglobin, which was used as a standard

of a marine mammal myoglobin with high net surface charge (Mirceta, et al. 2013). The

correlation between measured relative electrophoretic mobility and modelled Z_{Mb} values was

assessed using Phylogenetic Generalized Least Squares (PGLS) analysis using the CAPER

package (Orme, et al. 2013) as implemented in R v3.6.1 and the tree from the BEAST analysis in

Fig. 2. Because of low sample size, the parameter lambda was not estimated from the data but

fixed at a value of 1.0.

Lifestyle correlation analysis and ancestral lifestyle reconstruction

We analyzed the relationship between lifestyles and Z_{Mb} based on a threshold model

(Felsenstein 2012) using the phytools function threshBayes (Revell 2012) as implemented in R

v3.6. The threshold model hypothesizes that each lifestyle is determined by an underlying,

unobserved continuous trait (i.e., liability). We first categorized the 55 eulipotyphlan species for

which Z_{Mb} was calculated as either semi-aquatic (including the semi-aquatic/fossorial star-nosed

mole in this category) or non-aquatic based on the habits described in Burgin and He (2018). We

ran Markov chain Monte Carlo (MCMC) for 10^7 generations, sampling every 500 generations,

and discarded the first 20% generations as burn-in. We plotted the posterior sample for the

correlation to examine whether analyses reached a stationary state. We also estimated the

correlation between Z_{Mb} and full fossoriality (i.e., fossorial species versus non-fossorial species),

as well as that between Z_{Mb} and 'digging' (a category that included both fossorial and semi-

fossorial species) habits using the same approach. Finally, we also created subsets of our dataset

to enable comparisons between only two ecomorphotypes, with the following four threshBayes

analyses conducted: terrestrial Z_{Mb} versus semi-aquatic Z_{Mb}, terrestrial Z_{Mb} versus fully fossorial

Z_{Mb}, terrestrial Z_{Mb} versus semi-fossorial/fossorial Z_{Mb}, fully fossorial Z_{Mb} versus semi-aquatic

Z_{Mb}.

We estimated the ancestral lifestyle using a maximum parsimony and a threshold model

based on the 76 species time-calibrated concatenated gene tree (fig. 2) and the *BEAST

coalescent species tree (fig. S3). We categorized the species into non-aquatic or semi-aquatic as

above. We used the R package castor to reconstruct ancestral lifestyles using maximum

parsimony (Louca and Doebeli 2017), treating the transition cost between non-aquatic and semi-

aquatic equally. We then used the R package phytools to estimate ancestral states using a

threshold model (Revell 2014). We ran MCMC for 1 million generations, sampling every 1000

generations, and discarded the first 20% generations as burn-in. We performed the threshold

analyses using either a Brownian motion or lambda model for estimating the liability, and

compared the results based on deviance information criterion (DIC). We selected the result of the

lambda model because it outperformed the Brownian motion model (ΔDIC =99, data not

shown).

Acknowledgements

We thank curators and staff from the Smithsonian Institute, National Museum of Natural History

(K.M. Helgen), Field Museum (L. Heaney), Burke Museum (B. Sharon) and New Mexico

Museum of Natural History (B. Oh), National Museum of Natural Science (Y. Chen), Kunming

Institute of Zoology (S. Li and X-L. Jiang) for approving our proposal for the use of tissue

samples, or conducting destructive sampling on museum skin specimens. We are grateful to K.

Wareing and S. Mirceta, University of Liverpool for muscle samples of the European mole and

hedgehog, and for help with native PAGE of muscle extracts, respectively. We thank M. Docker,

University of Manitoba, for use of the Ion torrent PGM, and Z.-L. Ding, Kunming Institute of

Zoology, for performing Ion Torrent Proton sequencing. We appreciate Ailaoshan Station for

Subtropical Forest Ecosystem Studies, Xishuangbanna Tropical Botanical Garden, Chinese

Academy of Sciences for allowing us to conduct fieldwork. We also thank Y-H. Sun and S-Y.

Liu for supporting our fieldwork in Gansu and Sichuan provinces, and S-L. Yuan and L-K. Lin

for assistance with fieldwork in Taiwan. Finally, we are grateful to Umi Matsushita for painting

eulipotyphlan species for our figures. This work was supported by the National Natural Science

Foundation of China (31970389, 31301869 to K. H.), the National Science Foundation (NSF

DEB-1457735 to M.S.S), and by the University of Manitoba Research Grants Program (41342),

National Sciences and Engineering Research Council of Canada Discovery (RGPIN/238838-

2011; RGPIN/6562-2016) and Discovery Accelerator Supplement (RGPIN/412336-2011) grants

to K.L.C.

Data Availability Statement

The newly obtained myoglobin sequences were deposited to GenBank under accession numbers

MW456061 to MW456069 and MW473727- MW473769, and sequence alignments per gene

were deposited to Dryad Digital Repository at https://doi.org/10.5061/dryad.brv15dv7q.

Figures

Fig. 1. Museum specimen photos illustrating the four major ecomorphotypes within the order Eulipotyphla. Representative terrestrial (shrew-like mole, *Uropsilus soricipes*; bottom right), semi-aquatic (Russian desman, *Desmana moschata*; right centre), strictly fossorial (Eastern mole, *Scalopus aquaticus*; left), and semi-fossorial (Chinese long-tailed mole, *Scaptonyx fusicaudus*; top centre) talpid mole species are given. Photo by Kai He.

Fig. 2. Time calibrated Bayesian phylogenetic tree of Eulipotyphla based on a concatenated alignment of 23 nuclear genes (outgroups not shown). Branch lengths represent median ages. Node bars indicate the 95% confidence interval [CI] for each clade age. Unless specified, all relationships are highly supported. Relationships weakly supported in concatenation Bayesian and maximum likelihood (PP <0.97 and/or BS <80: #) as well as *BEAST and ASTRAL coalescent analyses (C-PP <0.97 and/or C-BS <80: *) are indicated. Note that an alternative position was recovered for Soricini in the ASTRAL tree (supplementary fig. S2), while both coalescent analyses (ASTRAL, *BEAST) favored *Episoriculus* monophyly (supplementary figs. S2, S3). Coloured bars at the tips of the tree denote terrestrial (green), semi-aquatic (blue), semi-fossorial (beige) and fossorial (brown) lifestyles in extant species.

Fig. 3. Three-dimensional structural models of myoglobin in (A) the last common ancestor of Eulipotyphla and (B) the semi-aquatic Russian desman (*Desmana moschata*) obtained by homology modelling using the SWISS-MODEL server (Waterhouse, et al. 2018). The structure of the last common ancestor of the group was modelled based on results of an amino acid sequence reconstruction (see text for details). Ancestral (left) and derived (right) states of charge-changing amino acid replacements are circled and indicated with positional number and one-letter amino acid code. Blue and red color indicate amino acids with positively (H, His; K, Lys; R, Arg) and negatively charged amino acid side chains (D, Asp; E, Glu), respectively. White double arrows indicate surface amino acid side chains involved in salt bridges that are affected by charge-changing substitutions. Text boxes indicate the reconstructed temporal order (top to bottom) of charge decreasing and charge increasing amino acid substitutions (red and blue font, respectively) in the *Desmana* lineage in one letter code from ancestral (left) to derived (right)

separated by positional number. Note that charge neutral substitutions (e.g., G35N), are not given

in the text boxes.

Fig. 4. (A) Violin plot showing the distribution (y-axis) and probability density (x-axis) of

modelled myoglobin net surface charge, Z_{Mb}, among living species (black dots) of the four

prevalent eulipotyphlan ecomorphotypes. (B) Correlation between Z_{Mb} and electrophoretic

mobility of native myoglobin from five eulipotyphlan insectivores; data from the grey seal

(*Halichoerus grypus*) is added for comparison. Z_{Mb} was calculated as the sum of the charge of all

ionisable groups at pH 6.5 by modelling myoglobin primary structures onto the tertiary structure

and using published, conserved, site-specific ionization constants (McLellan 1984; Mirceta, et al.

2013). Electrophoretic mobility was assessed relative to the mobility of grey seal myoglobin

using native polyacrylamide gel electrophoresis of heart or skeletal muscle protein extracts of the

indicated species. Green, orange, brown, and blue areas (A) or symbols and fonts (B) indicate

terrestrial, semi-fossorial, fossorial, and semi-aquatic/aquatic species, respectively. Phylogenetic

Generalized Least Squares analysis in panel (B) revealed a highly significant positive correlation

($R^2 = 0.897$, $p < 0.005$) between the two parameters (solid line, $y = 0.1488x + 0.3075$).

Fig. 5. Evolutionary reconstruction of myoglobin net surface charge Z_{Mb} in 55 eulipotyphlan

insectivores mapped onto the time calibrated phylogeny of Fig. 2. Ancestral Z_{Mb} was modelled

from primary structures as in Fig. 3 and after maximum likelihood ancestral sequence

reconstruction. Major charge increasing (blue font) and charge decreasing (red font) amino acid

substitutions, from ancestral to derived and separated by positional number, inferred for the

immediate ancestry of semi-aquatic species (blue font) are indicated in textboxes alongside the

respective branches. Grey and white background shading indicates geologic epochs. See

supplementary Fig. S7A for a complete account of charge-changing substitutions, reconstructed

Z_{Mb} values, and outgroup information. Paintings of representative species by Umi Matsushita.

Supplementary files

Fig. S1. The full RAxML species tree constructed from 71 eulipotyphlan specimens. Bootstrap

supports are given next to internal nodes.

Fig. S2. The full ASTRAL-III coalescent species tree constructed from 71 eulipotyphlan

specimens. Bootstrap supports are given next to internal nodes.

Fig. S3. The full *BEAST coalescent species tree constructed from 71 eulipotyphlan specimens.

Posterior probabilities are given next to internal nodes.

Fig. S4. Location of charge-changing amino acid substitutions in the oxygen-storing protein

myoglobin of four semi-aquatic species of moles and shrews in the mammalian insectivore order

Eulipotyphla. Three-dimensional structures were obtained by homology modelling using the

SWISS-MODEL server (Waterhouse, et al. 2018) and the amino acid sequences of (A) *Sorex*

palustris, (B) *Neomys fodiens*, (C) *Nectogale elegans*, and (D) *Condylura cristata*. The three-

dimensional myoglobin structure of the last common ancestor of the group was also modelled

based on results of an amino acid sequence reconstruction (see text for details). Ancestral (left)

and derived (right) states of charge-changing amino acid replacements are circled and indicated

with positional number and one-letter amino acid code. Blue and red color indicate amino acids

with positively (H, His; K, Lys; R, Arg) and negatively charged amino acid side chains (D, Asp;

E, Glu). White double arrows indicate salt bridges that are affected by charge-changing

substitutions. Image views between panels A-D have been rotated to maximally visualise

lineage-specific replacements.

Fig. S5: Structural model of myoglobin in (A) the last common ancestor of Eulipotyphla and (B)

the semi-aquatic Russian desman (*Desmana moschata*). Structures were obtained by homology

modelling using the SWISS-MODEL server (Waterhouse, et al. 2018) and the primary structures

of myoglobin obtained by conceptual translation of the here determined nucleotide sequence

(Russian desman) or by ancestral amino acid sequence reconstruction (see text for details).

Structures were visualised in PyMOL version 2.1.1. Note that the gap at position 121 in the GH-

loop of the tertiary structure (circled in red) of the Russian desman in (B) appears to exert

negligible effect on the tertiary structure of the protein.

Fig. S6. Time-calibrated tree of 55 eulipotyphlan species for which complete myoglobin coding

sequences were determined (left). Horizontal bars on the right indicate the calculated Z_{Mb} for

each species, which are color coded according to species lifestyle.

Fig. S7. Maximum likelihood ancestral (A) amino acid-based and (B) codon-based sequence

reconstruction and net surface charge (Z_{Mb}) calculation of eulipotyphlan myoglobin mapped onto

the time calibrated phylogeny of fig. 2. Results of a separate amino-acid based reconstruction

based on the *BEAST species tree reconstruction (supplementary fig. S3) is provided in (C) with

Episoriculus fumidus as the species with an alternative position compared to *A* and *B* in red font.

Only charge-changing substitutions are shown, with blue and red font in text boxes indicating

charge increasing and charge decreasing substitutions, respectively. Absolute charge-changing

substitutions >1.0 are underlined, whereas those <0.10 are shown in brackets. Yellow

highlighting indicates amino acids reconstructed with p < 0.95 for which less likely but

differentially charged amino acids have been reconstructed with p > 0.05 (see text for details).

Reconstructed net charges are shown at nodes, though have been omitted in some cases if they

were identical to values on the preceding node, for clarity. Terminal species considered semi-

aquatic are indicated by blue font, with Z_{Mb} values >+2.0 also given in blue font.

Fig. S8. Comparisons of the Bayesian concatenation species tree estimated using the tree-of-life

genes with myoglobin RAxML gene trees estimated using (A) nucleotide and (B) amino-acid

sequences. Bootstrap supports are given next to internal nodes on the myoglobin gene trees. Only

Bootstrap supports higher than 70 are shown.

Fig. S9. Ancestral reconstruction of semi-aquatic lifestyles (blue filled circles) within

Eulipotyphla based on both the (A) RAxML concatenation gene tree and (B) the *BEAST

species tree. For each tree, reconstructions were constructed using either a maximum parsimony

(left) or a threshold (right) model. Note that *Episoriculus fumidus* (red font in *B*) was supported

at different positions on the two trees.

Fig. S10. Posterior density distribution of the correlation coefficient (r) between (A) semi-

aquatic, (B) fully fossorial, and (C) digging (to include fossorial and semi-fossorial lifestyles)

and Z_{Mb} as estimated using the threshBayes function. Figures D-G show the results based on the

analyses of subsets of samples that included only: (D) terrestrial and semi-aquatic species, (E)

terrestrial and fully fossorial species, (F) terrestrial and digging species, or (G) semi-aquatic and

fully fossorial species. The solid red line indicates the grand mean, the green dashed lines

indicate the mean of the 80% confidence intervals, and the grey dashed lines indicate the mean

of the 95% confidence intervals.

Fig. S11. A heatmap showing tree-of-life genes included for each of the 76 samples used for

phylogenetic analyses. The ultrametric tree is the BEAST concatenation gene tree and a blank

block indicates the gene is missing in the final dataset.

Table S1. Sample information of specimens used in this study.

Table S2. Hybridization capture results of tree-of-life gene segments from 61 eulipotyphlan

DNA libraries. Numbers in each column represent total base pairs captured; NA: no data.

Table S3. Result of likelihood-based Shimodaira–Hasegawa test to compare the best scoring

RAxML concatenated gene tree and alternative evolutionary hypotheses.

Table S4. Myoglobin amino acid alignment used for modeling myoglobin net surface charge

(Z_{Mb}) and ancestral sequence reconstructions. Myoglobin helices A to H are highlighted in

yellow, with amino acid positions and helical notations indicated above and below the graphic,

respectively. Internal amino acid residue positions are shaded in light grey, while deleted

residues are indicated by a dash mark. Strongly anionic residues (D [Asp] and E [Glu]) are

shaded in red, with strongly (K [Lys] and R [Arg]) and weakly (H [His]) cationic residues

shaded in dark and light green, respectively.

Table S5. Charge increasing (blue font) and decreasing (red font) residue substitutions

reconstructed for semi-aquatic eulipotyphlan branches.

Table S6. Evolutionary models estimated using bModelTest in BEAST, and used for BEAST

and *BEAST analyses.

Table S7. RAxML best scoring gene trees used for ASTRAL-III coalescent analysis before and

after collapsing 0% Shimodaira–Hasegawa (SH) scores in Newick format.

Table S8. The best scoring concatenation species trees estimated using BEAST and RAxML,

and the best species coalescence trees estimated using ASTRAL-III and *BEAST, in Newick

format.

Table S9. Primers used to amplify and sequence the protein coding exons of myoglobin.

Text S1. Calibrations used for estimating divergence times in the BEAST analyses.

References

Allam A, Kalnis P, Solovyev V. 2015. Karect: accurate correction of substitution, insertion and

deletion errors for next-generation sequencing data. *Bioinformatics* 31:3421–3428.

Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. In:

Babraham Bioinformatics, Babraham Institute, Cambridge, United Kingdom.

Ashkenazy H, Penn O, Doron-Faigenboim A, Cohen O, Cannarozzi G, Zomer O, Pupko T. 2012.

FastML: a web server for probabilistic reconstruction of ancestral sequences. *Nucleic Acids Res.*

40:W580–W584.

Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko

SI, Pham S, Prjibelski AD. 2012. SPAdes: a new genome assembly algorithm and its

applications to single-cell sequencing. *J Comput Biol.* 19:455–477.

Barido-Sottani J, Boskova V, Plessis LD, Kuhnert D, Magnus C, Mitov V, Muller NF, PecErska

J, Rasmussen DA, Zhang C, et al. 2018. Taming the BEAST-A community teaching material

resource for BEAST 2. *Syst Biol.* 67:170–174.

Berenbrink M. 2020. The role of myoglobin in the evolution of mammalian diving capacity –

The August Krogh principle applied in molecular and evolutionary physiology. *Comp Biochem*

Physiol A Mol Integr Physiol. 252:110843.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence

data. *Bioinformatics* 30:2114–2120.

Bosshard HR, Marti DN, Jelesarov I. 2004. Protein stabilization by salt bridges: concepts,

experimental approaches and clarification of some misunderstandings. *J Mol Recognit.* 17:1–16.

Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu C-H, Xie D, Suchard MA, Rambaut A,

Drummond AJ. 2014. BEAST 2: a software platform for Bayesian evolutionary analysis. *PLoS*

Comput Biol. 10:e1003537.

Bouckaert RR, Drummond AJ. 2017. bModelTest: Bayesian phylogenetic site model averaging

and model comparison. *BMC Evol Biol.* 17:42.

Brace S, Thomas JA, Dalen L, Burger J, MacPhee RDE, Barnes I, Turvey ST. 2016.

Evolutionary history of the Nesophontidae, the last unplaced recent mammal family. *Mol Biol*

Evol. 33:3095–3103.

Burgin C, He K. 2018. Family Soricidae (Shrews). In: Wilson DE, Mittermeier RA, editors.

Handbook of The Mammals of the World. Vol. 8. Insectivores, Sloths and Colugos. Barcelona:

Lynx Edicions. p. 332–551.

Burgin CJ, Colella JP, Kahn PL, Upham NS. 2018. How many species of mammals are there? *J*

Mammal. 99:1–14.

Butler PM. 1998. Fossil history of shrews in Africa. In: Wojcik JM, Wolsan M, editors.

Evolution of Shrews. Bialowieza: Mammal Research Institute. p. 121–132.

Casewell NR, Petras D, Card DC, Suranse V, Mychajliw AM, Richards D, Koludarov I,

Albulescu LO, Slagboom J, Hempel BF, Ngum NM. 2019. Solenodon genome reveals

convergent evolution of venom in eulipotyphlan mammals. *Proc Nat Acad Sci U S A.*

116:25745–25755.

Campbell B. 1939. The shoulder anatomy of the moles. A study in phylogeny and adaptation.

Am J Anat. 64:1–39.

Catania KC. 2006. Olfaction - Underwater 'sniffing' by semi-aquatic mammals. *Nature*

444:1024–1025.

Catania KC, Hare JF, Campbell KL. 2008. Water shrews detect movement, shape, and smell to

find prey underwater. *Proc Nat Acad Sci U S A.* 105:571–576.

1001

1002   Chevreux B, Pfisterer T, Drescher B, Driesel AJ, Muller WEG, Wetter T, Suhai S. 2004. Using

1003   the miraEST assembler for reliable and automated mRNA transcript assembly and SNP detection

1004   in sequenced ESTs. *Genome Res.* 14:1147-1159.

1005

1006   Churchfield S. 1990. The Natural History of Shrews: Cornell University Press.

1007

1008   Cox MP, Peterson DA, Biggs PJ. 2010. SolexaQA: At-a-glance quality assessment of Illumina

1009   second-generation sequencing data. *BMC Bioinformatics* 11:485.

1010

1011   Criscuolo A, Brisse S. 2013. AlienTrimmer: a tool to quickly and accurately trim off multiple

1012   short contaminant sequences from high-throughput sequencing reads. *Genomics* 102:500–506.

1013

1014   Dayhoff M, Schwartz R, Orcutt B. 1978. A model of evolutionary change in proteins. In. Atlas

1015   of protein sequence and structure: National Biomedical Research Foundation Silver Spring MD.

1016   p. 345–352.

1017

1018   Emerling CA, Springer MS. 2014. Eyes underground: regression of visual protein networks in

1019   subterranean mammals. *Mol Phylogenet Evol.* 78:260–270.

1020

1021   Enoki Y, Ohga Y, Ishidate H, Morimoto T. 2008. Primary structure of myoglobins from 31

1022   species of birds. *Comp Biochem Physiol B Biochem Mol Biol.* 149:11–21.

1023

1024   Esselstyn JA, Oliveros CH, Swanson MT, Faircloth BC. 2017. Investigating difficult nodes in

1025   the placental mammal tree with expanded taxon sampling and thousands of ultraconserved

1026   elements. *Genome Biol Evol.* 9:2308–2321.

1027

1028   Faircloth BC. 2015. PHYLUCE is a software package for the analysis of conserved genomic

1029   loci. *Bioinformatics* 32:786–788.

1030

1031   Felsenstein J. 2012. A comparative method for both discrete and continuous characters using the

1032   threshold model. *Am Nat.* 179:145–156.

1033

1034   Gao F, Chen C, Arab DA, Du Z, He Y, Ho SYW. 2019. EasyCodeML: A visual tool for analysis

1035   of selection using CodeML. *Ecol Evol.* 9:3891–3898.

1036

1037   Gonzalez-Dominguez J, Schmidt B. 2016. ParDRe: faster parallel duplicated reads removal tool

1038   for sequencing studies. *Bioinformatics* 32:1562–1564.

1039

1040   Grabherr M, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L,

1041   Raychowdhury R, Zeng QJNB. 2011. Full-length transcriptome assembly from RNA-Seq data

1042   without a reference genome. *Nat Protoc.* 29:644-652.

1043

1044   Gorman ML, Stone RD. 1990. The Natural History of Moles: Cornell University Press.

1045

Grand T, Gould E, Montali R. 1998. Structure of the proboscis and rays of the star-nosed mole, *Condylura cristata*. *J Mammal.* 79:492–501.

1048

Grossnickle DM, Smith SM, Wilson GP. 2019. Untangling the multiple ecological radiations of early mammals. *Trends Ecol Evol.* 34:936-949.

1051

Gusztak RW. 2008. Dive performance and aquatic thermoregulation of the world's smallest mammalian diver, the American water shrew (*Sorex palustris*). M.Sc. thesis. University of Manitoba.

1055

Gusztak RW, MacArthur RA, Campbell KL. 2005. Bioenergetics and thermal physiology of American water shrews (*Sorex palustris*). *J Comp Physiol B.* 175:87–95.

1058

Hahn MW, Nakhleh L. 2016. Irrational exuberance for resolved species trees. *Evolution* 70:7-17.

He K, Chen J-H, Gould GC, Yamaguchi N, Ai H-S, Wang Y-X, Zhang Y-P, Jiang X-L. 2012. An estimation of Erinaceidae phylogeny: A combined analysis approach. *PLoS ONE* 7:e39304.

1062

He K, Li YJ, Brandley MC, Lin LK, Wang YX, Zhang YP, Jiang XL. 2010. A multi-locus phylogeny of Nectogalini shrews and influences of the paleoclimate on speciation and evolution. *Mol Phylogenet Evol.* 56:734–746.

1066

He K, Shinohara A, Helgen KM, Springer MS, Jiang X-L, Campbell KL. 2017. Talpid mole phylogeny unites shrew moles and illuminates overlooked cryptic species diversity. *Mol Biol Evol.* 34:78–87.

1070

He K, Shinohara A, Jiang X-L, Campbell KL. 2014. Multilocus phylogeny of talpine moles (Talpini, Talpidae, Eulipotyphla) and its implications for systematics. *Mol Phylogenet Evol.* 70:513–521.

1074

1075   Heled J, Drummond AJ. 2010. Bayesian inference of species trees from multilocus data. *Mol*

1076   *Biol Evol.* 27:570–580.

1077   Hickman GC. 1984. Swimming ability of talpid moles, with particular reference to the semi-

1078   aquatic *Condylura cristata*. *Mammalia* 48:505–514.

1079

1080   Hooker JJ. 2016. Skeletal adaptations and phylogeny of the oldest mole Eotalpa (Talpidae,

1081   Lipotyphla, Mammalia) from the UK Eocene: the beginning of fossoriality in moles.

1082   *Palaeontology* 59:195–216.

1083

1084   Horn S. 2012. Target enrichment via DNA hybridization capture. In: Shapiro B, Hofreiter M,

1085   editors. Ancient DNA: Springer. p. 177–188.

1086

1087   Hutterer R. 1985. Anatomical adaptations of shrews. *Mamm Rev.* 15:43–55.

1088

1089   Isogai Y, Imamura H, Nakae S, Sumi T, Takahashi K-i, Nakagawa T, Tsuneshige A, Shirai T.

1090   2018. Tracing whale myoglobin evolution by resurrecting ancient proteins. *Sci Rep.* 8:16883.

1091

Ivlev YF, Kuznetsov A, Rutovskaya M. 2010. Preliminary data on the swimming kinematics of

the Russian desman (*Desmana moschata* L.). *Dokl Biol Sci.* 431:144–148.

Ivlev YF, Rutovskaya MV, Luchkina OS. (Ivlev2013 co-authors). 2013. The use of olfaction by

the Russian desman (*Desmana moschata* L.) during underwater swimming. *Dokl Biol Sci.*

452:280–283.

Junemann S, Sedlazeck FJ, Prior K, Albersmeier A, John U, Kalinowski J, Mellmann A,

Goesmann A, von Haeseler A, Stoye J, Harmsen D. 2013. Updating benchtop sequencing

performance comparison. *Nat Biotechnol.* 31:294–296.

1100

1101   Junier T, Zdobnov EM. 2010. The Newick utilities: high-throughput phylogenetic tree

1102   processing in the UNIX shell. *Bioinformatics* 26:1669–1670.

1103

1104   Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:

1105   improvements in performance and usability. *Mol Biol Evol.* 30:772–780.

1106

1107   Kearse M, Sturrock S, Meintjes P. 2012. The Geneious 6.0. 3 read mapper. Auckland, New

1108   Zealand: Biomatters, Ltd.

1109

1110   Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: molecular evolutionary

1111   genetics analysis across computing platforms. *Mol Biol Evol.* 35:1547–1549.

1112

1113   Lestyk KC, Folkow LP, Blix AS, Hammill MO, Burns JM. 2009. Development of myoglobin

1114   concentration and acid buffering capacity in harp (*Pagophilus groenlandicus*) and hooded

1115   (*Cystophora cristata*) seals from birth to maturity. *J Comp Physiol B* 179:985–996.

1116

1117   Louca S, Doebeli M. 2017. Efficient comparative phylogenetics on large trees. *Bioinformatics*

1118   34:1053–1055.

1119

1120   Madsen O, Scally M, Douady CJ, Kao DJ, DeBry RW, Adkins R, Amrine HM, Stanhope MJ, de

1121   Jong WW, Springer MS. 2001. Parallel adaptive radiations in two major clades of placental

1122   mammals. *Nature* 409:610–614.

1123

1124   Martin DP, Murrell B, Golden M, Khoosal A, Muhire B. 2015. RDP4: Detection and analysis of

1125   recombination patterns in virus genomes. *Virus Evol* 1:vev003.

1126

1127   MacPhee RDE, Flemming C, Lunde DP. 1999. "Last occurrence" of the Antillean insectivoran

1128   Nesophontes: new radiometric dates and their interpretation. *Am Mus Novit.* 3261:1–19.

1129

1130   Mason VC, Li G, Helgen KM, Murphy WJ. 2011. Efficient cross-species capture hybridization

1131   and next-generation sequencing of mitochondrial genomes from noninvasively sampled museum

1132   specimens. *Genome Res.* 21:1695–1704.

1133

1134   McIntyre IW, Campbell KL, MacArthur RA. 2002. Body oxygen stores, aerobic dive limits and

1135   diving behaviour of the star-nosed mole (*Condylura cristata*) and comparisons with non-aquatic

1136   talpids. *J Exp Biol.* 205:45–54.

1137

1138   McLellan T. 1984. Molecular charge and electrophoretic mobility in cetacean myoglobins of

1139   known sequence. *Biochem Genet.* 22:181–200.

1140

1141   Meredith RW, Janecka JE, Gatesy J, Ryder OA, Fisher CA, Teeling EC, Goodbla A, Eizirik E,

1142   Simao TL, Stadler T, et al. 2011. Impacts of the Cretaceous terrestrial revolution and KPg

1143   extinction on mammal diversification. *Science* 334:521–524.

1144

1145   Mirceta S, Signore AV, Burns JM, Cossins AR, Campbell KL, Berenbrink M. 2013. Evolution

1146   of mammalian diving capacity traced by myoglobin net surface charge. *Science* 340:1234192.

1147

1148   Mittermeier RA, Wilson DE, Rylands AB. 2018. Handbook of the Mammals of the World:

1149   Insectivores, Sloths and Colugos. Barcelona Lynx Edicions.

1150

1151   Motokawa M. 2004. Phylogenetic relationships within the family Talpidae (Mammalia:

1152   Insectivora). *J Zool (Lond.)* 263:147–157.

1153

1154   Mulugeta E, Wassenaar E, Sleddens-Linkels E, van Ijcken WFJ, Heard E, Grootegoed JA, Just

1155   W, Gribnau J, Baarends WM. 2016. Genomes of *Ellobius* species provide insight into the

1156   evolutionary dynamics of mammalian sex chromosomes. *Genome Res.* 26:1202–1210.

1157

1158   Noren SR, Williams TM. 2000. Body size and skeletal muscle myoglobin of cetaceans:

1159   adaptations for maximizing dive duration. *Comp Biochem Physiol A Mol Integr Physiol.*

1160   126:181–191.

1161

1162   Nowak RM. 1999. Walker's Mammals of the World. Baltimore: Johns Hopkins University Press.

1163

1164 Orme D, Freckleton R, Thomas G, Petzoldt T, 2013. The caper package: comparative analysis of

1165 phylogenetics and evolution in R. *R package version* 5:1–36.

1166

1167 Piras P, Sansalone G, Teresi L, Kotsakis T, Colangelo P, Loy A. 2012. Testing convergent and

1168 parallel adaptations in talpids humeral mechanical performance by means of geometric

1169 morphometrics and finite element analysis. *J Morphol.* 273:696–711.

1170

1171 Ponganis PJ, Meir JU, Williams CL. 2011. In pursuit of Irving and Scholander: a review of

1172 oxygen store management in seals and penguins. *J Exp Biol.* 214:3325–3339.

1173

1174 Reed CA. 1951. Locomotion and appendicular anatomy in three soricoid insectivores. *Am Midl*

1175 *Nat.* 45:513–671.

1176

1177 Revell LJ. 2014. Ancestral character estimation under the threshold model from quantitative

1178 genetics. *Evolution* 68:743–759.

1179 Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and other

1180 things). *Methods Ecol Evol.* 3:217–223.

1181

1182 Rofes J, Cuenca-Bescos G. 2006. First evidence of the Soricidae (Mammalia) *Asoriculus*

1183 *gibberodon* (Petényi, 1864) in the Pleistocene of north Iberia. *Riv Ital Paleontol Stratigr.*

1184 112:301–315.

1185

1186 Romero-Herrera A, Lehmann H, Joysey K, Friday A. 1973. Molecular evolution of myoglobin

1187 and the fossil record: a phylogenetic synthesis. *Nature* 246:389–395.

1188

1189 Romero-Herrera A, Lehmann H, Joysey K, Friday A. 1978. On the evolution of myoglobin.

1190 *Philos Trans R Soc Lond B Biol Sci.* 283:61–163.

1191

1192 Samuel P, Phillips G, Olson J. 2015. A cell-free investigation of the relationship between

1193 myoglobin expression, globin stability and heme affinity. *FASEB J.* 29:882–888.

1194

1195   Sanchez-Villagra MR, Horovitz I, Motokawa M. 2006. A comprehensive morphological analysis

1196   of talpid moles (Mammalia) phylogenetic relationships. *Cladistics* 22:5–88.

1197

1198   Sansalone G, Kotsakis T, Piras P. 2016. *Condylura* (Mammalia, Talpidae) reloaded: New

1199   insights about the fossil representatives of the genus. *Palaeontol Electron.* 19.3.54A

1200

1201   Sansalone G, Colangelo P, Loy A, Raia P, Wroe S, Piras P. 2019. Impact of transition to a

1202   subterranean lifestyle on morphological disparity and integration in talpid moles (Mammalia,

1203   Talpidae). *BMC Evol Biol.* 19:179.

1204

1205   Scholander P. 1962. Physiological adaptation to diving in animals and man. *Harvey Lect.* 57:93–

1206   110.

1207

1208   Schwermann AH, Thompson RS. 2015. Extraordinarily preserved talpids (Mammalia,

1209   Lipotyphla) and the evolution of fossoriality. *J Vert Paleontol.* 35:e934828.

1210

1211   Simmons MP, Kessenich J. 2020. Divergence and support among slightly suboptimal likelihood

1212   gene trees. *Cladistics* 36:322–340.

1213

1214   Simmons MP, Sloan DB, Springer MS, Gatesy J. 2019. Gene-wise resampling outperforms site-

1215   wise resampling in phylogenetic coalescence analyses. *Mol Phylogenet Evol.* 131:80–92.

1216

1217   Simpson JT, Wong K, Jackman SD, Schein JE, Jones SJM, Birol I. 2009. ABySS: A parallel

1218   assembler for short read sequence data. *Genome Res.* 19:1117–1123.

1219   Springer MS, Murphy WJ, Roca AL. 2018. Appropriate fossil calibrations and tree constraints

1220   uphold the Mesozoic divergence of solenodons from other extant mammals. *Mol Phylogent Evol.*

1221   121:158–165.

1222

1223   Springer MS, Stanhope MJ, Madsen O, de Jong WW. 2004. Molecules consolidate the placental

1224   mammal tree. *Trend Ecol Evol.* 19:430–438.

1225

1226   Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of

1227   large phylogenies. *Bioinformatics* 30:1312–1313.

1228

1229   Stöver BC, Müller KF. 2010. TreeGraph 2: Combining and visualizing evidence from different

1230   phylogenetic analyses. *BMC Bioinformatics* 11:7.

1231

1232   Vogel P, Bodmer C, Spreng M, Aeschimann J. 2007. Diving capacity and foraging behaviour of

1233   the water shrew. In: Gorman ML, Dunstone N, editors. Behaviour and Ecology of Riparian

1234   Mammals. New York: Cambridge University Press Cambridge, United Kingdom. p. 31–48.

1235

1236   Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, Heer FT, de Beer

1237   TAP, Rempfer C, Bordoli L, Lepore R, Schwede T. 2018. SWISS-MODEL: homology

1238   modelling of protein structures and complexes. *Nucleic Acids Res.* 46:W296–W303.

1239

1240   Whidden HP. 1999. The evolution of locomotor specializations in moles. *Am Zool.* 39: 135A.

1241

1242   Whidden HP. 2000. Comparative myology of moles and the phylogeny of the Talpidae

1243   (Mammalia, Lipotyphla). *Am Mus Novit.* 3294:1–53.

1244

1245   Woodman N, Gaffney SA. 2014. Can they dig it? Functional morphology and semifossoriality

1246   among small-eared shrews, genus *Cryptotis* (Mammalia, Soricidae). *J Morphol.* 275:745–759.

1247

1248   Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol.* 24:1586–

1249   1591.

1250

1251   Zhang C, Rabiee M, Sayyari E, Mirarab S. 2018. ASTRAL-III: polynomial time species tree

1252   reconstruction from partially resolved gene trees. *BMC Bioinformatics* 19:153.

1253

1254   Zimin AV, Smith DR, Sutton G, Yorke JA. 2008. Assembly reconciliation. *Bioinformatics*

1255   24:42–45.

20 mm

A) Last common ancestor of Eulipotyphla

B) *Desmana moschata*

H48N
S132K

Q26R
N35D
D44A

Q113K

Z_{Mb}

+3
+2
+1
±0
−1

Solenodon paradoxus
Chimarrogale styani
Chimarrogale platycephala
Nectogale elegans
Episoriculus macrurus
Episoriculus umbrinus
Episoriculus fumidus
Neomys fodiens
Anourosorex squamipes
Sorex bedfordiae
Sorex araneus
Sorex palustris
Blarinella quadraticauda
Blarina brevicauda
Crocidura indochinensis
Neotetracus sinensis
Erinaceus europaeus
Uropsilus gracilis
Uropsilus atronates
Uropsilus soricipes
Uropsilus sp. 1
Uropsilus nivatus
Scapanus orarius
Scapanus latimanus
Scapanus townsendii
Scapanus aquaticus
Scalopus aquaticus
Parascalops breweri
Parascalops oweni
Galemys pyrenaicus
Desmana moschata
Urotrichus talpoides
Dymecodon pilirostris
Neurotrichus gibbsii
Scaptonyx fusicaudu
Scaptonyx sp.
Condylura cristata
Talpa europaea
Talpa caeca
Talpa romana
Mogera robusta
Mogera wogura
Mogera etigo
Mogera tokudae
Mogera latouchei
Mogera kanoana
Mogera insularis
Oreoscaptor mizura
Scaptochirus moschatus
Parascaptor sp. 1
Euroscaptor longirostris
Euroscaptor grandis
Euroscaptor klossi
Euroscaptor malayana
Euroscaptor sp. 1
Euroscaptor sp. 2

Q113K
N12K
D53A
D44S
D122N
S132K
E136Q
E27A
E27L
Q26R
N35D
D44A
D53G
Q128K
G129E
S132K
D53G
S132K

Pleistocene
Pliocene
Mio-cene
Oligocene
Eocene
Paleocene
Late Cretaceous

Time before present (Ma)
80