Article Type: Letter

Article Title: **Thermal tolerance and preference are both consistent with the clinal**

distribution of house fly proto-Y chromosomes

Pablo J Delclos[†,1], Kiran Adhikari[†,1], Jessica E Cambric[1], Anna G Matuk[1], Rebecca I Presley[1],

Jessica Tran[1], Richard P Meisel[1]

[†]Co-first author/equal contribution

Affiliation: [1]Department of Biology & Biochemistry, University of Houston, Houston, TX, USA

Running Title: Genotypic control of concordant thermal traits

Keywords: Thermoregulation, polygenic sex determination, sex chromosomes, *Musca domestica*,

genotype-by-environment effects

Abstract

Selection pressures can vary within localized areas and across massive geographical scales. Temperature is one of the best studied ecologically variable abiotic factors that can affect selection pressures across multiple spatial scales. Organisms rely on physiological (thermal tolerance) and behavioral (thermal preference) mechanisms to thermoregulate in response to environmental temperature. In addition, spatial heterogeneity in temperatures can select for local adaptation in thermal tolerance, thermal preference, or both. However, the concordance between thermal tolerance and preference across genotypes and sexes within species and across populations is greatly understudied. The house fly, *Musca domestica*, is a well-suited system to examine how genotype and environment interact to affect thermal tolerance and preference. Across multiple continents, house fly males from higher latitudes tend to carry the male-determining gene on the Y chromosome, whereas those from lower latitudes usually have the male-determiner on the third chromosome. We tested whether these two male-determining chromosomes differentially affect thermal tolerance and preference as predicted by their geographical distributions. We identify effects of genotype and developmental temperature on male thermal tolerance and preference that are concordant with the natural distributions of the chromosomes, suggesting that temperature variation across the species range contributes to the maintenance of the polymorphism. In contrast, female thermal preference is bimodal and largely independent of congener male genotypes. These sexually dimorphic thermal preferences suggest that temperature-dependent mating dynamics within populations could further affect the distribution of the two chromosomes. Together, the differences in thermal tolerance and

preference across sexes and male genotypes suggest that different selection pressures may affect

the frequencies of the male-determining chromosomes across different spatial scales.

Impact Statement

Genetic variation within species can be maintained by environmental factors that vary

across the species' range, creating clinal distributions of alleles responsible for ecologically

important traits. Some of the best examples of clinal distributions come from

temperature-dependent phenotypes, such as thermal tolerance and preference. Although genotype

and developmental temperature strongly affect physiological and behavioral traits in ectotherms,

the correlation between these traits across genotypes and sexes within species is greatly

understudied. We show that two different male-determining chromosomes found in natural

populations of house fly affect both thermal tolerance and preference in a way that is concordant

with their clinal distributions across latitudes. This provides strong evidence that temperature

variation across the species range contributes to the maintenance of the polymorphism.

Furthermore, we find evidence that thermal preference is sexually dimorphic, suggesting that

temperature-dependent mating dynamics could further affect the distribution of genetic variation

in this system. Therefore, at a macro-geographical scale, the differences in thermal tolerance and

preference across male genotypes likely contributes to the maintenance of the cline. Within

populations, differences in thermal preference likely affect sexual selection dynamics, which may

further affect the frequencies of the chromosomes.

Introduction

Ecological variation across a species' range can select for local adaptation within populations, which can contribute to the maintenance of genetic variation by favoring different alleles across the range (Levene 1953; Felsenstein 1976; Hedrick et al. 1976; Kawecki and Ebert 2004). In addition, heterogeneous selection pressures that are distributed as a gradual continuum from one end of the species' range to another can create a cline of genetic variation responsible for phenotypes under selection (Slatkin 1973; Endler 1977). Some of the best examples of latitudinal clines come from temperature-dependent phenotypes (e.g., body size, developmental rate, and thermal tolerance) that have been well-documented in flies (Partridge et al. 1994; Eanes 1999; Robinson and Partridge 2001; Hoffmann et al. 2002). Moreover, heterogeneous selection pressures across a cline may affect males and females differently (Connallon 2015; Connallon et al. 2019), although the empirical evidence for such variation in sex-specific selection across geographic ranges is mixed (Delcourt et al. 2009; Delph et al. 2011; Allen et al. 2017; Lasne et al. 2018).

Thermal adaptation within populations and across a species range can occur via selection on physiological, anatomical, or behavioral traits. For example, north-south gradients in heat and cold tolerance have been observed in *Drosophila* (Hoffmann et al. 2002), suggesting physiological adaptation to thermal environments. In addition, ectotherms, such as flies, rely on behavioral mechanisms of thermoregulation by avoiding suboptimal temperatures in search of more optimal ones (Dillon et al. 2009), and thermal preference may be correlated with optimal thermal performance (Dawson 1975; Angilletta et al. 2002).

Concordance across genotypes between different thermal traits could reinforce the

response to selection, whereas negative correlations could constrain adaptation (Etterson and

Shaw 2001). However, it is not clear if physiological and behavioral thermal traits are genetically

correlated within a species, between sexes, or across populations (Dawson 1975; Angilletta et al.

2002; Gilbert and Miles 2017). For example, experiments in *Drosophila subobscura* identified

individual chromosomes that affected thermal tolerance or temperature preference, but no single

chromosome affected both physiological and behavioral phenotypes (Dolgova et al. 2010; Rego

et al. 2010; Castañeda et al. 2019). Furthermore, temperature-dependent traits can affect

assortative mating and male reproductive success (Dolgin et al. 2006; Keller and Seehausen

2012), suggesting inter-sexual differences in thermoregulation could affect genetic variation

within populations via sexual selection. These sex-specific selection pressures could also

contribute to the maintenance of genetic variation via inter-sexual conflict or context-dependent

selection (Kotiaho et al. 2001; Rostant et al. 2015; Meisel 2018). Despite the importance of

inter-sexual differences, previous work did not test for differences in the genetic correlation of

thermal traits between males and females.

We used a sex chromosome polymorphism in house fly, *Musca domestica*, to investigate

the concordance of thermal tolerance and preference across clinally distributed male genotypes.

House fly has a polygenic sex determination system, with a male-determining gene most

commonly found on either the third chromosome (III^M) or what was historically refered to as the

Y chromosome (Y^M) (Hamm et al. 2014; Sharma et al. 2017). Both III^M and Y^M are very young

proto-Y chromosomes that are minimally differentiated from their homologous proto-X

chromosomes (Meisel et al. 2017; Son et al. 2019; Son and Meisel 2020). Y^M and III^M are

distributed along latitudinal clines on multiple continents in the Northern Hemisphere (Tomita

and Wada 1989; Hamm et al. 2005; Kozielska et al. 2008). Y^M is most frequently found at

northern latitudes, and III^M is more common at southern latitudes (Figure 1A). This distribution

suggests that the Y^M chromosome confers higher fitness in colder climates, and, conversely, III^M

confers higher fitness in hotter climates. Therefore, variation in temperature across the species

range may create heterogeneous selection pressures that maintain the proto-Y chromosome cline

in house fly. Consistent with this hypothesis, seasonality in temperature is the best predictor of

the frequencies of the proto-Y chromosomes across natural populations (Feldmeyer et al. 2008).

We tested the hypothesis that the Y^M chromosome confers cold-adaptive phenotypes and

III^M confers heat-adaptive phenotypes in house fly males, which would be consistent with their

latitudinal distributions (Tomita and Wada 1989; Hamm et al. 2005; Feldmeyer et al. 2008;

Kozielska et al. 2008). To those ends, we first evaluated if males carrying the III^M chromosome

(hereafter III^M males) have greater tolerance to extreme heat and if males carrying the Y^M

chromosome (Y^M males) have greater cold tolerance. Second, we tested if III^M males prefer

warmer temperatures than Y^M males, and if males and females differ in their thermal preference.

We performed all experiments using flies raised at multiple developmental temperatures because

thermal acclimation strongly affects temperature-dependent phenotypes in flies and other

ectotherms (Krstevska and Hoffmann 1994; Dillon et al. 2009). Together, we evaluated if thermal

preference and tolerance are aligned for sex-linked genetic variants, tested if this alignment could

maintain polygenic sex determination, and then discuss how these temperature-dependent

phenotypes could affect the access of males to female mates.

Materials and methods

Fly strains and rearing

We performed our experiments using five nearly isogenic house fly strains, three with III^M

males and two with Y^M males (Supplementary Methods). All five strains have a common genetic

background from an inbred III^M strain that was produced from a mixture of flies collected across

the United States (Scott et al. 1996; Hamm et al. 2005). Each of the three III^M strains carries a

different III^M chromosome from a separate wild-derived line, and, likewise, the two Y^M strains

carry different Y^M chromosomes.

We reared each strain at 18°C, 22°C, and 29°C for two generations in order to evaluate

how thermal acclimation affects thermal tolerance (Chown and Terblanche 2006) and thermal

preference (Krstevska and Hoffmann 1994; Dillon et al. 2009). Flies from each developmental

temperature were assayed at equivalent physiological ages estimated by accumulated degree days

(Barnard and Geden 1993; Wang et al. 2018). For our heat and cold tolerance assays, we used

flies 22–50 total degree days after eclosion. For thermal preference assays, we used flies 96–115

total degree days after eclosion. Additional details and calculations are provided in the

Supplementary Methods.

Thermal tolerance

We measured heat and cold tolerance in individual male and female flies. To measure heat

tolerance, lightly anaesthetized individual flies were transferred to a 1.5 ml centrifuge tube that

was sealed with fabric. We placed the 1.5 ml tube in a heat block set to 53°C. This temperature

was selected because it is the lowest at which heat tolerance could be measured in a reasonable

period of time. The time at which a fly fell to the bottom of the tube and could not make its way

back to the top of the tube was considered the knockdown time. To measure cold tolerance,

lightly anaesthetized flies were transferred to a fabric-sealed 20 ml glass vial individually, and the

vials were placed in a 4°C refrigerator with a transparent door. Knockdown occured when a fly

fell on its back to the bottom of the vial. We gently tapped the assay vial every 2–3 minutes to

ensure flies were active.

For both heat and cold tolerance assays, we performed an analysis of variance (ANOVA)

using the lmer() function in the lme4 (v1.1) R package (Bates et al. 2015) to model the effect of

genotype (G: Y^M vs III^M), developmental temperature (T: 18°C or 29°C), and their interaction on

knockdown time (K):

$$K \sim G + T + G{\times}T + B + S,$$

with batch (B) and strain (S) treated as random effects. We also constructed another model

excluding the interaction term:

$$K \sim G + T + B + S.$$

We then used a drop in deviance test to compare the fit of the models with and without the

interaction term using the anova() function in R. We also compared heat and cold tolerance

between males raised at 22°C and 29°C, using the same approaches as described above.

Additional details are provided in the Supplementary Methods.

 Thermal preference

We measured thermal preference as the position of individual flies along a 17–37°C

thermal gradient (Figure S1), following a slightly modified version of previous protocols

(Anderson et al. 2013; Lynch et al. 2018). For each individual fly, we report mean thermal

preference (T_{pref}) as the average position during a 10 minute assay window (measured once per

minute). We also report thermal breadth, $T_{breadth}$ (Carrascal et al. 2016), as the coefficient of

variation of individual-level T_{pref} during the assay window. $T_{breadth}$ provides an estimate of how

individuals utilize thermal space within their environment (Slatyer et al. 2013). Choosier

individuals show a lower $T_{breadth}$ value and, thus, would be expected to occupy a narrower range of

temperatures within a given thermal habitat.

To determine the effects of developmental temperature (18°C, 22°C, and 29°C), genotype

(Y^M and III^M), and their interaction on mean T_{pref} across sexes, we created a mixed-effects model

using the lme4 package (v1.1) in R (Bates et al. 2015). Developmental temperature, genotype,

and their interaction were included as fixed effects, and strain, batch, and lane in the thermal

gradient (L) were included as random effects:

169                              $$T_{pref} \sim G + T + G{\times}T + B + S + L.$$

We did the same for $T_{breadth}$. We then determined whether groups significantly differed in T_{pref} or

$T_{breadth}$ using Tukey contrasts with the multcomp package (v1.4) in R (Hothorn et al. 2008).

Within developmental temperature treatments, we used Bayesian information criterion (BIC)

scores from the mclust (v5.4.5) package in R (Scrucca et al. 2016) to determine whether the

distribution of individual measures of T_{pref} within a group are best explained by one or multiple

normal distributions.

Results

Thermal tolerance depends on developmental temperature and male genotype

We measured extreme heat (53°C) and cold (4°C) tolerance as a readout of differences in

physiological thermal adaptation between Y^M and III^M house fly males. We observed the expected

effect of acclimation on both heat and cold tolerance (Chown and Terblanche 2006): flies raised

at 18°C tolerate cold longer than the flies raised at 29°C, and flies raised at 29°C tolerate heat

longer than flies raised at 18°C (Figure 1). We also find that Y^M males are more cold tolerant, and

III^M males are more heat tolerant, consistent with the latitudinal distributions of Y^M and III^M

males in nature (Tomita and Wada 1989; Hamm et al. 2005; Feldmeyer et al. 2008; Kozielska et

al. 2008). However, the effect of genotype on thermal tolerance depends on acclimation.

Specifically, a linear model with an interaction between genotype (Y^M or III^M) and developmental

temperature fits the cold tolerance data significantly better than a model without the interaction

term ($\chi^2_1 = 19.3$, $p = 1.1$ x 10^{-5}). This provides evidence for a G×T effect on cold tolerance—Y^M

males are more cold tolerant than III^M males, but only if they are raised at 18°C (Figure 1B).

There is also a significant G×T interaction affecting heat tolerance ($\chi^2_1 = 4.71$, $p = 0.030$

comparing models with and without the interaction term): III^M males are more heat tolerant than

Y^M males, but only if raised at 29°C (Figure 1D).

Figure 1 - Thermal tolerance in males and females. (**A**) Map of the eastern United States, showing the cline of Y^M (more common in the north) and III^M (more common in the south). (**B–E**) Graphs show the effect of developmental temperature on knockdown time at either 4°C (cold tolerance) or 53°C (heat tolerance) for Y^M (turquoise) and III^M (salmon) male flies. Proto-Y chromosomes for females reflect whether males from the strain carry the Y^M or III^M chromosome. Mean knockdown time is plotted for each combination of genotype and temperature. Error bars represent standard error.

We next attempted to identify a threshold temperature for the genotype-specific benefits of acclimation by comparing heat and cold tolerance of flies raised at 22°C and 29°C (instead of 18°C and 29°C). We did not observe a significant effect of the interaction between developmental temperature and male genotype on extreme cold tolerance ($\chi^2_1 = 0.947$, $p = 0.331$ comparing models with and without an interaction term) (Figure S2). We therefore hypothesize that there is a

threshold temperature between 18°C and 22°C, below which Y^M males experience a greater

benefit of cold acclimation than III^M males. In contrast, there is a significant interaction between

genotype and developmental temperature on heat tolerance when comparing males raised at 22°C

and 29°C ($\chi^2_1 = 11.02$, $p = 9.0$ x 10^{-4} comparing models with and without the interaction term)

(Figure S2). Therefore, the threshold for a genotype-specific benefit from heat acclimation lies

between 22°C and 29°C.

We do not expect any difference in heat or cold tolerance across females from our

different strains because all females have the same genotype, regardless of the male genotype in

the strain. Indeed, a model with an interaction between developmental temperature and male

genotype does not fit the female cold tolerance data better than a model without the interaction

term ($\chi^2_1 = 1.46$, $p = 0.23$) (Figure 1C). There is a significant effect of developmental temperature

on cold tolerance in females ($\chi^2_1 = 43.5$, $p = 4.3$ x 10^{-11} comparing a model with and without

developmental temperature), demonstrating that females benefit from cold acclimation regardless

of male genotype (Figure 1C). Surprisingly, there is a significant interaction between male

genotype and developmental temperature on heat tolerance in females ($\chi^2_1 = 10.4$, $p = 0.0013$

comparing a model with and without the interaction term). In general, females raised at warmer

temperatures are more heat tolerant (Figure 1E). However, the interaction of male genotype and

developmental temperature is in the opposite direction from what would be expected based on the

latitudinal distribution of Y^M and III^M: females from strains with Y^M males that are raised at 29°C

are more heat tolerant than females from III^M strains raised at 29°C (Figure 1E). We thus

conclude that the heat and cold tolerance differences between Y^M and III^M males are specific to

males and/or the proto-Y chromosomes (i.e., not genetic background) because we do not observe

the same heat or cold tolerance differences in females from those strains (who do not carry the

proto-Y chromosomes).

Figure 2 - Thermal preference (T_{pref}) of (**A**) male and (**B**) female house flies according to

genotype (III^M = salmon points and line, Y^M = turquoise points and line) and developmental

temperature. Each point depicts the mean thermal preference for an individual fly, with lines and

error bars denoting means within groups and standard errors of the mean, respectively. Significant

differences between groups are denoted by letters, with differing letters highlighting significantly

different mean thermal preferences (Tukey's *post hoc* test, $p < 0.05$).

Thermal preference depends on developmental temperature and male genotype

We next tested if genotype and developmental temperature affect thermal preference

(T_{pref}). First, we find that T_{pref} is inversely proportional to developmental temperature (Figure 2),

with house flies that develop at a warmer temperature preferring cooler temperatures (and *vice*

versa), regardless of sex (male: $F_{2, 742.7} = 138.4$, $p < 1.0 \times 10^{-5}$; female: $F_{2, 245.3} = 37.1$, $p = 1.19 \times$

10^{-4}; Figure 2). This is consistent with how developmental acclimation affects T_{pref} in *Drosophila*

(Dillon et al. 2009).

We also find that male proto-Y chromosome genotype (Y^M vs III^M) affects T_{pref} ($F_{1, 756.2}=$

44.5, $p < 1.0$ x 10^{-5}). There is also a significant interaction effect between developmental

temperature and genotype on T_{pref} in males ($F_{2, 756.3} = 8.47$, $p = 2.31$ x 10^{-4}, Figure 2A). Male T_{pref}

is similar across genotypes when they develop at either 18°C or 29°C. However, when reared at

22°C, III^M males prefer warmer temperatures than Y^M males (Tukey's *post hoc* test, $p < 0.001$).

This is consistent with III^M males being more common at lower latitudes (where average

temperatures are warmer), and Y^M males more common at higher latitudes (Tomita and Wada

1989; Hamm et al. 2005; Feldmeyer et al. 2008; Kozielska et al. 2008). We do not expect

differences in T_{pref} in females across strains because all females have the same genotype. Indeed,

the genotype of males in a strain (Y^M vs III^M) and the interaction between male genotype and

female developmental temperature showed no significant effect on T_{pref} in females (ANOVA, all p

> 0.1 in Figure 2B).

Thermal breadth depends on sex and thermal preference

We used thermal breadth ($T_{breadth}$) as a measure of the specificity of T_{pref}. Male $T_{breadth}$ was

not significantly affected by either developmental temperature, genotype, or the interaction

between genotype and developmental temperature (ANOVA, all $p > 0.1$, Figure S3A). In contrast,

developmental temperature ($F_{2, 236.9} = 16.5$, $p < 1.0$ x 10^{-5}), as well as the interaction between

developmental temperature and male genotype ($F_{2, 243.6} = 5.35$, $p = 0.005$), had significant effects

on $T_{breadth}$ in females. However, the significant interaction is of small effect, as females from

strains with differing male genotypes do not significantly differ in $T_{breadth}$ within any

developmental temperature treatment (Figure S3B).

Figure 3 - Thermal breadth ($T_{breadth}$) depends on male genotype and sex. (**A**) Distribution of

individual-level mean thermal preferences (T_{pref}) of III^M males, Y^M males, and pooled females that

developed at 22°C. Y-axis represents relative density of data points and is analogous to frequency

of data points for a given T_{pref} value. (**B**) $T_{breadth}$ of individuals raised at 22°C according to group

(F^C = cold-preferring females, F^W = warm-preferring females). Boxplots denote median values

and lower- and upper- quartiles. Asterisks denote significant differences in $T_{breadth}$ between groups

(*****: Tukey's *post hoc* test, $p < 0.01$).

The effect of developmental temperature on $T_{breadth}$ in females is driven by increased

variance in T_{pref} when females develop at 22°C. The increased variance in female T_{pref} can be

explained by a mixture of two normal distributions (Figure 3A, see Table S1 for statistics). This

bimodal distribution is not a result of differences across strains because the same pattern was

observed among females separately analyzed based on male genotype (Figure S4). In

comparison, a single normal distribution best fit Y^M male T_{pref} when developed at 22°C, and two

normal distributions best explained the III^M male T_{pref} when developed at 22°C. Upon inspection,

however, the two distributions representing III^M male T_{pref} likely correspond to the tail (mean of

28.7°C and large variance of 10.4°C) and peak (mean of 32.6°C and small variance of 0.4°C) of a

single skewed distribution, which we are unable to detect using the mclust package we used to fit

distributions to our data.

We used our model-based clustering analysis of T_{pref} to classify individuals that developed

at 22°C into one of four groups: Y^M males (lower T_{pref}), III^M males (higher T_{pref}), females with

cooler T_{pref} (F^C females, 59.3% of females tested), and females with warmer T_{pref} (F^W females,

40.7% of females tested). The mean T_{pref} of F^C females (26.90°C) is nearly equal to the mean T_{pref}

of Y^M males (26.87°C; Figure 3A). Similarly, the mean T_{pref} of F^W females (32.2°C) is near the

mode of the T_{pref} of III^M males (32.0–32.5°C; Figure 3A).

We further find that T_{pref} is predictive of $T_{breadth}$ for flies that develop at 22°C. We

considered flies from our four T_{pref} groups (Y^M males, III^M males, F^C females, and F^W females),

and we found a significant effect of group on $T_{breadth}$ ($F_{3, 32.9} = 9.40$, $p = 1.24$ x 10^{-4}). Specifically,

F^C females have significantly greater $T_{breadth}$ than all other groups (Tukey's *post hoc* test, all $p <$

1.0 x 10^{-5}, Figure 3B). Therefore, if we consider $T_{breadth}$ as a measure of the strength of T_{pref}, adult

house flies can be summarized by one of three phenotypes related to thermal behavior when

developed at 22°C: a relatively strong preference for warm temperatures (III^M males and F^W

females, which have high T_{pref} and low $T_{breadth}$), a strong preference for cooler temperatures (Y^M

males, with low T_{pref} and low $T_{breadth}$), and a relatively weak preference for cooler temperatures

(F^C females, with low T_{pref} and high $T_{breadth}$).

Discussion

We tested if thermal tolerance and preference depend on sex and male genotype in house

fly. We find that males carrying the Y^M chromosome (which is common in the northern end of the

species' range) are more cold tolerant and prefer colder temperatures. Conversely, males carrying

the III^M chromosome (which is common in the southern end of the species' range) are more heat

tolerant and prefer warmer temperatures. Our results are therefore consistent with the general

trend that temperate populations are typically more cold-tolerant than (sub-) tropical ones (Gibert

and Huey 2001; Hoffmann et al. 2002). Moreover, these differences in thermal tolerance and

preference depend on developmental temperature, and they are not observed in congener females

from the same strains (who do not carry the Y^M or III^M chromosome). However, females exhibit a

bimodal T_{pref}, with females from each of the two subgroups overlapping with one of the male

genotypes.

Thermal tolerance and preference depend on developmental temperature, genotype, and sex

Our results demonstrate, to the best of our knowledge, the first documented example of

concordant temperature preference, cold tolerance, and heat tolerance across genotypes within a

species. We find that Y^M males both have greater cold tolerance and prefer colder temperatures,

whereas III^M males have greater heat tolerance and prefer warmer temperatures (Figures 1 and 2),

consistent with their latitudinal distributions (Tomita and Wada 1989; Hamm et al. 2005;

Feldmeyer et al. 2008; Kozielska et al. 2008). Previous work has identified concordant T_{pref} and

heat tolerance differences across species (Qu et al. 2011), or found no clear relationship between

thermal tolerance and preference across genotypes within species (Yang et al. 2008; Rego et al.

2010; Castañeda et al. 2019).

We observed strong effects of developmental temperature on both thermal tolerance and

preference that depend on both genotype and sex. Acclimation effects on heat and cold tolerance

(Figure 1) are well-documented for ectotherms, including flies and other insects (Bowler and

Terblanche 2008). An inverse relationship between developmental temperature and thermal

preference has also been observed in other flies (Dillon et al. 2009; Castañeda et al. 2013).

Behaviorally navigating towards compensatory temperatures could serve as a means of mitigating

the costs of thermally suboptimal development (i.e., too hot or too cold). The observed

relationships between thermal tolerance and developmental temperatures are unlikely to be the

result of natural selection during acclimation within our experiment for two reasons. First, there is

unlikely to be sufficient genetic variation in these inbred strains for selection to generate these

results within 2 generations. Second, prior attempts at selecting for thermal tolerance in house

flies resulted in negligible differences in tolerance across developmental temperatures (Geden et

al. 2019). We therefore conclude that the differences in thermal tolerance (and preference)

between Y^M and III^M males have evolved across the natural populations from which we sampled

the Y^M and III^M chromosomes.

There are important methodological implications for our observation that variation in

thermal preference across genotypes depends on developmental temperature. We only observe

warmer (colder) thermal preferences in III^M (Y^M) males when developed at 22°C; thermal

preference did not differ between male genotypes when raised at more extreme (18°C, 29°C)

temperatures (Figure 2A). Previous studies attempting to estimate genetic variance in thermal

preference within or among populations of *Drosophila* have had mixed results. While some

studies identified genetic variance among populations within species (Good 1993; Castañeda et

al. 2013), others did not detect substantial variance within (Krstevska and Hoffmann 1994) or

among species (MacLean et al. 2019). Our results show that the phenotypic presentation of

genetic variation for thermal preference can depend on the environmental conditions experienced,

which could explain why this variance was not detected in other experiments.

We identified a female-specific plasticity for thermal preference that does not map to

genotype. In females, we found that neither thermal tolerance nor thermal preference differ

predictably between strains where males carry different proto-Y chromosomes (Figures 1C, E and

2B). However, there is a bimodal thermal preference for females that develop at 22°C (Figure

3A), regardless of congener male genotype. In addition, females that had colder T_{pref} when

developed at 22°C also had a larger $T_{breadth}$ (Figure 3B). This suggests that, in nature, females with

colder temperature preferences may occupy a wider range of temperatures than females with

warmer temperature preferences. Because all females in our experiment are expected to have the

same genotype, we hypothesize that these differences in T_{pref} and $T_{breadth}$ are conferred by a plastic

response to some yet to be characterized factor (e.g., microclimates within larval rearing

containers). Alternatively, this plasticity could have a stochastic origin that is intrinsic to the

development of thermal preference (Honegger and de Bivort 2018; Jensen 2018).

The correlation between thermal preference and thermal breadth at 22°C is

female-specific: Y^M and III^M males have similar $T_{breadth}$ values when raised at 22°C despite their

differences in T_{pref}. Although general sex differences in thermal tolerance (Hoffmann et al. 2005)

and thermal preference (Krstevska and Hoffmann 1994) have been documented, this is the first

study, to our knowledge, to identify sex differences in the relationship between thermal

preference and thermal breadth. Our results suggest that male and female house flies exhibit

different thermoregulatory behavioral patterns which may further be influenced by genotype.

Directly identifying a sex-by-genotype-by-environment interaction is beyond the scope of this

study because sex and genotype are confounded in our experimental design (the females in our

experiment have a different genotype from either male, characterized by a lack of either the III^M

or Y^M chromosome). Nonetheless, the house fly is a tractable system for directly testing for

sex-specific genotype-by-environment interactions on thermoregulation. For example, future

work could test for sex-specific effects of Y^M and III^M by measuring phenotypes in females

carrying a proto-Y chromosome along with the epistatic female-determining Md-tra^D allele

(Hediger et al. 2010; Hamm et al. 2014).

Environmental heterogeneity and the maintenance of polygenic sex determination

Sex determination pathways rapidly diverge across species, driving evolutionary turnover

of sex chromosomes (Bull 1983; Beukeboom and Perrin 2014). Polygenic sex determination

systems, in which more than one master sex determining locus segregate independently on

different chromosomes, have been observed in multiple animal species (Moore and Roberts

2013). Most population genetic models that attempt to explain the stable maintenance of

polygenic sex determination focus on sexually antagonstic effects of sex determining loci or

linked alleles on sex chromosomes (Rice 1986; van Doorn and Kirkpatrick 2007; Meisel et al.

2016). Less attention has been given to ecological factors that can maintain polygenic sex

determination (cf. Bateman and Anholt 2017).

Our results demonstrate how spatially variable ecological factors can maintain polygenic

sex determination. Specifically, thermal tolerance and preference phenotypes conferred by the Y^M

and III^M chromosomes (Figures 1 and 2) are consistent with the clinal and temperature-dependent

distributions of the Y^M and III^M chromosomes (Tomita and Wada 1989; Hamm et al. 2005;

Feldmeyer et al. 2008; Kozielska et al. 2008). Previous experiments identified multiple fitness

advantages conferred by the III^M chromosome over Y^M at warmer temperatures, including an

increase in frequency of III^M over generations in a laboratory population (Hamm et al. 2009).

However, these fitness differences can only explain the invasion or fixation of the III^M

chromosome, not the maintenance of the polymorphism. In contrast, differences in thermal

tolerance and preference could maintain proto-Y chromosome polymorphism across the species'

range, similar to how selection maintains other clinal variation (Slatkin 1973; Endler 1977).

The house fly system reveals how temperature variation can contribute to the maintenance

of polygenic sex determination independently of selection on the sex-determination pathway

itself. Temperature is an important contributor to the evolution of sex determination pathways in

vertebrates (Bull and Vogt 1979; Holleley et al. 2015). However, the effects of the house fly

proto-Y chromosomes on thermal tolerance and preference likely act independently of the sex

determination pathway because there are not differences in the expression of sex determination

genes across house fly male genotypes raised at different temperatures in a way that is consistent

with their clinal distribution (Adhikari et al. n.d.). This suggests that the effects of the Y^M and III^M

chromosomes on thermal phenotypes is a result of alleles on proto-Y chromosomes that are

genetically linked to the male-determining locus, as opposed to the male-determiner itself.

Therefore, our results highlight how temperature can be important for the evolution of sex

determination independently of temperature-dependent activity of the sex determination pathway.

Future theoretical work should consider the effect of spatially heterogeneous selection pressures

on the maintenance of polygenic sex determination, similar to how temporal heterogeneity can

create fluctuating selection pressures that maintain polygenic sex determination (Bateman and

Anholt 2017).

Selection on thermal phenotypes may depend on geographical scale

Our results suggest that selection on thermal traits differs between macro-geographic

species ranges and at a micro-geographical scale within populations. Thermal tolerance and

preference in male house flies depend on proto-Y chromosomes genotype in a way that is

consistent with the latitudinal distribution of the Y^M and III^M chromosomes (Figures 1 and 2).

This suggests that, at the macro-geographic scale, selection is operating on male physiology and

behavior to create or maintain the clinal distribution of Y^M and III^M (Figure 4A). In comparison,

at an intermediate developmental temperature (22°C), female thermal preference is bimodal for a

reason that we have yet to determine (Figure 3A). This raises the possibility that within

populations near the center of the cline, where Y^M and III^M both segregate (e.g., Hamm and Scott

2008; Meisel et al. 2016), sexual selection may favor males that can preferentially obtain access

to the two different female phenotypes.

III^M males may disproportionately benefit from differences in T_{pref} and $T_{breadth}$ between

males and females. F^C females that prefer colder temperatures have greater $T_{breadth}$ than warm

preferring F^W females and both male genotypes (Figure 3B), suggesting that F^C females occupy a

wider range of thermal habitats. Thus, III^M males may gain an advantage by having greater access

to F^W females, as well as occasional access to F^C females, in contrast to Y^M males who would

only be likely to encounter F^C females (Figure 4B). This raises the possibility that differences in

thermal preference across genotypes and sexes could affect the dynamics of sexual selection.

Figure 4 - Selection on the III^M and Y^M chromosomes likely differs across geographic scales. (**A**)

At the macro-geographical scale, selection for thermal tolerance and/or thermal preference results

in the clinal distribution of the Y^M (turquoise) and III^M (salmon) chromosomes. (**B**) At

intermediate developmental temperatures, male genotypes (Y^M vs III^M) differ in thermal

preference, which may create asymmetrical mating opportunities because of variation in female

thermal preference and breadth (F^C vs F^W). The asymmetry of the overlap of males and females at

the intermediate developmental temperature could affect sexual selection in populations where

435   Y^M and III^M both segregate.

Acknowledgements

Tin Nguyen, Chantalle Vincent, and Mandy Cao assisted with experiments. This material is based upon work supported by the National Science Foundation under Grant No. DEB-1845686. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.

Author Contributions

PD, KA, and RM conceived and designed the study. KA and JC collected, and KA analyzed, all thermal tolerance data. PD, RP, JT, and AM collected, and PD analyzed, all thermal preference data. PD, KA, and RM wrote the manuscript, and all authors reviewed the manuscript prior to submission.

Data Accessibility

All data files used for analyses described in this manuscript have been deposited in Dryad (doi:10.5061/dryad.n2z34tmvs). Raw video and image files from thermal preference assays are available from the authors upon request.

References

Adhikari, K., J. H. Son, A. H. Rensink, J. Jaweria, D. Bopp, L. W. Beukeboom, and R. P. Meisel. n.d. Temperature-dependent effects of house fly proto-Y chromosomes on gene expression act independently of the sex determination pathway.

Allen, S. L., R. Bonduriansky, C. M. Sgro, and S. F. Chenoweth. 2017. Sex-biased transcriptome divergence along a latitudinal gradient. Mol. Ecol. 26:1256–1272.

Anderson, R. D., S. Blanford, and M. B. Thomas. 2013. House flies delay fungal infection by fevering: at a cost. Ecol. Entomol. 38:1–10.

Angilletta, M. J., T. Hill, and M. A. Robson. 2002. Is physiological performance optimized by thermoregulatory behavior?: a case study of the eastern fence lizard, *Sceloporus undulatus*. J. Therm. Biol. 27:199–204.

Barnard, D. R., and C. J. Geden. 1993. Influence of larval density and temperature in poultry manure on development of the house fly (Diptera: Muscidae). Environ. Entomol. 22:971–977.

Bateman, A. W., and B. R. Anholt. 2017. Maintenance of polygenic sex determination in a fluctuating environment: an individual-based model. J. Evol. Biol. 30:915–925.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting linear mixed-effects models using lme4. J. Stat. Soft. 67:1–48.

Beukeboom, L. W., and N. Perrin. 2014. The Evolution of Sex Determination. Oxford University Press, New York, NY.

Bowler, K., and J. S. Terblanche. 2008. Insect thermal tolerance: what is the role of ontogeny, ageing and senescence? Biol. Rev. Camb. Philos. Soc. 83:339–355.

Bull, J. J. 1983. Evolution of Sex Determining Mechanisms. Benjamin/Cummings, Menlo Park, CA.

Bull, J. J., and R. C. Vogt. 1979. Temperature-dependent sex determination in turtles. Science

206:1186–1188.

Carrascal, L. M., S. Villén-Pérez, and D. Palomino. 2016. Preferred temperature and thermal breadth of

birds wintering in peninsular Spain: the limited effect of temperature on species distribution. PeerJ

4:e2156.

Castañeda, L. E., J. Balanyà, E. L. Rezende, and M. Santos. 2013. Vanishing chromosomal inversion

clines in *Drosophila subobscura* from Chile: is behavioral thermoregulation to blame? Am. Nat.

182:249–259.

Castañeda, L. E., V. Romero-Soriano, A. Mesas, D. A. Roff, and M. Santos. 2019. Evolutionary potential

of thermal preference and heat tolerance in *Drosophila subobscura*. J. Evol. Biol. 32:818–824.

Chown, S. L., and J. S. Terblanche. 2006. Physiological diversity in insects: ecological and evolutionary

contexts. Adv. Insect Physiol. 33:50–152.

Connallon, T. 2015. The geography of sex-specific selection, local adaptation, and sexual dimorphism.

Evolution 69:2333–2344.

Connallon, T., S. Sharma, and C. Olito. 2019. Evolutionary consequences of sex-specific selection in

variable environments: four simple models reveal diverse evolutionary outcomes. Am. Nat.

193:93–105.

Dawson, W. R. 1975. On the physiological significance of the preferred body temperatures of reptiles.

Persp. of Biophys. Ecol. 443–473.

Delcourt, M., M. W. Blows, and H. D. Rundle. 2009. Sexually antagonistic genetic variance for fitness in

an ancestral and a novel environment. Proc. Biol. Sci. 276:2009–2014.

Delph, L. F., J. Andicoechea, J. C. Steven, C. R. Herlihy, S. V. Scarpino, and D. L. Bell. 2011.

Environment-dependent intralocus sexual conflict in a dioecious plant. New Phytol. 192:542–552.

Dillon, M. E., G. Wang, P. A. Garrity, and R. B. Huey. 2009. Review: Thermal preference in *Drosophila*.

J. Therm. Biol. 34:109–119.

Dolgin, E. S., M. C. Whitlock, and A. F. Agrawal. 2006. Male *Drosophila melanogaster* have higher

mating success when adapted to their thermal environment. J. Evol. Biol. 19:1894–1900.

Dolgova, O., C. Rego, G. Calabria, J. Balanyà, M. Pascual, E. L. Rezende, and M. Santos. 2010. Genetic

constraints for thermal coadaptation in *Drosophila subobscura*. BMC Evol. Biol. 10:363.

Eanes, W. F. 1999. Analysis of selection on enzyme polymorphisms. Annu. Rev. Ecol. Syst. 30:301–326.

Annual Reviews 4139 El Camino Way, P.O. Box 10139, Palo Alto, CA 94303-0139, USA.

Endler, J. A. 1977. Geographic Variation, Speciation, and Clines. Princeton University Press.

Etterson, J. R., and R. G. Shaw. 2001. Constraint to adaptive evolution in response to global warming.

Science 294:151–154.

Feldmeyer, B., M. Kozielska, B. Kuijper, F. J. Weissing, L. W. Beukeboom, and I. Pen. 2008. Climatic

variation and the geographical distribution of sex-determining mechanisms in the housefly. Evol.

Ecol. Res. 10:797–809.

Felsenstein, J. 1976. The theoretical population genetics of variable selection and migration. Annu. Rev.

Genet. 10:253–280.

Geden, C. J., H. Biale, E. Chiel, and D. M. Johnson. 2019. Effect of fluctuating high temperatures on

house flies (Diptera: Muscidae) and their principal parasitoids (*Muscidifurax spp.* and *Spalangia spp.*

[Hymenoptera: Pteromalidae]) from the United States. J. Med. Entomol. 56:1650–1660.

Gibert, P., and R. B. Huey. 2001. Chill-coma temperature in *Drosophila*: effects of developmental

temperature, latitude, and phylogeny. Physiol. Biochem. Zool. 74:429–434.

Gilbert, A. L., and D. B. Miles. 2017. Natural selection on thermal preference, critical thermal maxima

and locomotor performance. Proc. Biol. Sci. 284.

Good, D. S. 1993. Evolution of behaviours in *Drosophila melanogaster* in high temperatures: Genetic and

environmental effects. J. Ins. Physiol. 39:537–544.

Hamm, R. L., J.-R. Gao, G. G.-H. Lin, and J. G. Scott. 2009. Selective advantage for IIIM males over YM

males in cage competition, mating competition, and pupal emergence in *Musca domestica L.*

(Diptera: Muscidae). Environ. Entomol. 38:499–504.

Hamm, R. L., R. P. Meisel, and J. G. Scott. 2014. The evolving puzzle of autosomal versus Y-linked male

determination in *Musca domestica*. G3 5:371–384.

Hamm, R. L., and J. G. Scott. 2008. Changes in the frequency of YM versus IIIM in the housefly, *Musca

domestica* L., under field and laboratory conditions. Genet. Res. 90:493–498.

Hamm, R. L., T. Shono, and J. G. Scott. 2005. A cline in frequency of autosomal males is not associated

with insecticide resistance in house fly (Diptera: Muscidae). J. Econ. Entomol. 98:171–176.

Hediger, M., C. Henggeler, N. Meier, R. Perez, G. Saccone, and D. Bopp. 2010. Molecular

characterization of the key switch *F* provides a basis for understanding the rapid divergence of the

sex-determining pathway in the housefly. Genetics 184:155–170.

Hedrick, P. W., M. E. Ginevan, and E. P. Ewing. 1976. Genetic polymorphism in heterogeneous

environments. Annu. Rev. Ecol. Syst. 7:1–32.

Hoffmann, A. A., A. Anderson, and R. Hallas. 2002. Opposing clines for high and low temperature

resistance in *Drosophila melanogaster*. Ecol. Letters 5:614–618.

Holleley, C. E., D. O'Meally, S. D. Sarre, J. A. Marshall Graves, T. Ezaz, K. Matsubara, B. Azad, X.

Zhang, and A. Georges. 2015. Sex reversal triggers the rapid transition from genetic to

temperature-dependent sex. Nature 523:79–82.

Honegger, K., and B. de Bivort. 2018. Stochasticity, individuality and behavior. Curr. Biol. 28:R8–R12.

Hothorn, T., F. Bretz, and P. Westfall. 2008. Simultaneous inference in general parametric models. Biom. J

50:346–363.

Jensen, G. 2018. Behavioral Stochasticity. Pp. 1–5 *in* J. Vonk and T. Shackelford, eds. Encyclopedia of

Animal Cognition and Behavior. Springer International, Cham, Switzerland.

Kawecki, T. J., and D. Ebert. 2004. Conceptual issues in local adaptation. Ecol. Letters 7:1225–1241.

Keller, I., and O. Seehausen. 2012. Thermal adaptation and ecological speciation. Mol. Ecol. 21:782–799.

Kotiaho, J. S., L. W. Simmons, and J. L. Tomkins. 2001. Towards a resolution of the lek paradox. Nature

410:684–686.

Kozielska, M., B. Feldmeyer, I. Pen, F. J. Weissing, and L. W. Beukeboom. 2008. Are autosomal

sex-determining factors of the housefly (*Musca domestica*) spreading north? Genet. Res. 90:157–165.

Krstevska, B., and A. A. Hoffmann. 1994. The effects of acclimation and rearing conditions on the

response of tropical and temperate populations of *Drosophila melanogaster* and *D. simulans* to a

temperature gradient (Diptera: Drosophilidae). J. Ins. Behav. 7:279–288.

Lasne, C., S. B. Hangartner, T. Connallon, and C. M. Sgrò. 2018. Cross-sex genetic correlations and the

evolution of sex-specific local adaptation: Insights from classical trait clines in *Drosophila*

melanogaster. Evolution 72:1317–1327.

Levene, H. 1953. Genetic equilibrium when more than one ecological niche is available. Am. Nat.

87:331–333. The University of Chicago Press.

Lynch, K. E., T. E. White, and D. J. Kemp. 2018. The effect of captive breeding upon adult thermal

preference in the Queensland fruit fly (*Bactrocera tryoni*). J. Therm. Biol. 78:290–297.

MacLean, H. J., J. Overgaard, T. N. Kristensen, C. Lyster, L. Hessner, E. Olsvig, and J. G. Sørensen. 2019.

Temperature preference across life stages and acclimation temperatures investigated in four species of

Drosophila. J. Therm. Biol. 86:102428.

Meisel, R. P. 2018. Sexual conflict and the maintenance of genetic variation in natural populations. Mol.

Ecol. 27:3569–3571.

Meisel, R. P., T. Davey, J. H. Son, A. C. Gerry, T. Shono, and J. G. Scott. 2016. Is multifactorial sex

determination in the house fly, *Musca domestica* (L.), stable over time? J. Hered. 107:615–625.

Meisel, R. P., C. A. Gonzales, and H. Luu. 2017. The house fly Y chromosome is young and minimally

differentiated from its ancient X chromosome partner. Genome Res. 27:1417–1426.

Moore, E. C., and R. B. Roberts. 2013. Polygenic sex determination. Curr. Biol. 23:R510–2.

Partridge, L., B. Barrie, K. Fowler, and V. French. 1994. Evolution and development of body size and cell

size in *Drosophila melanogaster* in response to temperature. Evolution 48:1269–1276.

Qu, Y., H. Li, J. Gao, X. Xu, and X. Ji. 2011. Thermal preference, thermal tolerance and the thermal

dependence of digestive performance in two *Phrynocephalus* lizards (Agamidae), with a review of

species studied. Curr. Zool. 57:684–700.

Rego, C., J. Balanyà, I. Fragata, M. Matos, E. L. Rezende, and M. Santos. 2010. Clinal patterns of

chromosomal inversion polymorphisms in *Drosophila subobscura* are partly associated with thermal

preferences and heat stress resistance. Evolution 64:385–397.

Rice, W. R. 1986. On the instability of polygenic sex determination: the effect of sex-specific selection.

Evolution 40:633–639.

Robinson, S. J. W., and L. Partridge. 2001. Temperature and clinal variation in larval growth efficiency in

Drosophila melanogaster. J. Evol. Biol. 14:14–21.

Rostant, W. G., C. Kay, N. Wedell, and D. J. Hosken. 2015. Sexual conflict maintains variation at an

insecticide resistance locus. BMC Biol. 13:34.

Scott, J. G., P. Sridhar, and N. Liu. 1996. Adult specific expression and induction of cytochrome P450lpr

in house flies. Arch. Insect Biochem. Physiol. 31:313–323.

Scrucca, L., M. Fop, T. B. Murphy, and A. E. Raftery. 2016. mclust 5: clustering, classification and density

estimation using Gaussian finite mixture models. R J. 8:289–317.

Sharma, A., S. D. Heinze, Y. Wu, T. Kohlbrenner, I. Morilla, C. Brunner, E. A. Wimmer, L. van de Zande,

M. D. Robinson, L. W. Beukeboom, and D. Bopp. 2017. Male sex in houseflies is determined by

Mdmd, a paralog of the generic splice factor gene. Science 356:642–645.

Slatkin, M. 1973. Gene flow and selection in a cline. Genetics 75:733–756.

Slatyer, R. A., M. Hirst, and J. P. Sexton. 2013. Niche breadth predicts geographical range size: a general

ecological pattern. Ecol. Letters 16:1104–1114.

Son, J. H., T. Kohlbrenner, S. Heinze, L. W. Beukeboom, D. Bopp, and R. P. Meisel. 2019. Minimal

effects of proto-Y chromosomes on house fly gene expression in spite of evidence that selection

maintains stable polygenic sex determination. Genetics 213:313–327.

Son, J. H., and R. P. Meisel. 2020. Gene-level, but not chromosome-wide, divergence between a very

young house fly proto-Y chromosome and its homologous proto-X chromosome. Mol. Biol. Evol.,

doi: 10.1093/molbev/msaa250.

Tomita, T., and Y. Wada. 1989. Multifactorial sex determination in natural populations of the housefly

(*Musca domestica*) in Japan. The Japan. J. of Gen. 64:373–382.

van Doorn, G. S., and M. Kirkpatrick. 2007. Turnover of sex chromosomes induced by sexual conflict.

Nature 449:909–912.

Wang, Y., L. Yang, Y. Zhang, L. Tao, and J. Wang. 2018. Development of *Musca domestica* at constant

temperatures and the first case report of its application for estimating the minimum postmortem

interval. Forensic Sci. Int. 285:172–180.

Yang, J., Y.-Y. Sun, H. An, and X. Ji. 2008. Northern grass lizards (*Takydromus septentrionalis*) from

different populations do not differ in thermal preference and thermal tolerance when acclimated under

identical thermal conditions. J. Comp. Physiol. B 178:343–349.