Article Type: Letter

Article Title: **Thermal tolerance and preference are both consistent with the clinal**

distribution of house fly proto-Y chromosomes

Pablo J Delclos[†,1], Kiran Adhikari[†,1], Oluwatomi Hassan[1], Jessica E Cambric[1], Anna G Matuk[1],

Rebecca I Presley[1], Jessica Tran[1], Vyshnika Sriskantharajah[1,2], Richard P Meisel[1]

[†]Co-first author/equal contribution

Affiliation: [1]Department of Biology & Biochemistry, University of Houston, Houston, TX, USA

Current affiliations:

[2]University of Texas Health Science Center School of Biomedical Informatics

Running Title: Genotypic control of concordant thermal traits

Keywords: Thermoregulation, polygenic sex determination, sex chromosomes, *Musca domestica*,

genotype-by-environment effects

Abstract

Selection pressures can vary within localized areas and across massive geographical

scales. Temperature is one of the best studied ecologically variable abiotic factors that can affect

selection pressures across multiple spatial scales. Organisms rely on physiological (thermal

tolerance) and behavioral (thermal preference) mechanisms to thermoregulate in response to

environmental temperature. In addition, spatial heterogeneity in temperatures can select for local

adaptation in thermal tolerance, thermal preference, or both. However, the concordance between

thermal tolerance and preference across genotypes and sexes within species and across

populations is greatly understudied. The house fly, *Musca domestica*, is a well-suited system to

examine how genotype and environment interact to affect thermal tolerance and preference.

Across multiple continents, house fly males from higher latitudes tend to carry the male-

determining gene on the Y chromosome, whereas those from lower latitudes usually have the

male-determiner on the third chromosome. We tested whether these two male-determining

chromosomes differentially affect thermal tolerance and preference as predicted by their

geographical distributions. We identify effects of genotype and developmental temperature on

male thermal tolerance and preference that are concordant with the natural distributions of the

chromosomes, suggesting that temperature variation across the species range contributes to the

maintenance of the polymorphism. In contrast, female thermal preference is bimodal and largely

independent of congener male genotypes. These sexually dimorphic thermal preferences suggest

that temperature-dependent mating dynamics within populations could further affect the

distribution of the two chromosomes. Together, the differences in thermal tolerance and

preference across sexes and male genotypes suggest that different selection pressures may affect

the frequencies of the male-determining chromosomes across different spatial scales.

Impact Statement

Genetic variation within species can be maintained by environmental factors that vary across the species' range, creating clinal distributions of alleles responsible for ecologically important traits. Some of the best examples of clinal distributions come from temperature-dependent phenotypes, such as thermal tolerance and preference. Although genotype and developmental temperature strongly affect physiological and behavioral traits in ectotherms, the correlation between these traits across genotypes and sexes within species is greatly understudied. We show that two different male-determining chromosomes found in natural populations of house flies affect both thermal tolerance and preference in a way that is concordant with their clinal distributions across latitudes. This provides strong evidence that temperature variation across the species range contributes to the maintenance of the polymorphism. Furthermore, we find evidence that thermal preference is sexually dimorphic, suggesting that temperature-dependent mating dynamics could further affect the distribution of genetic variation in this system. Therefore, at a macro-geographical scale, the differences in thermal tolerance and preference across male genotypes likely contributes to the maintenance of the cline. Within populations, differences in thermal preference likely affect sexual selection dynamics, which may further affect the frequencies of the chromosomes.

Introduction

Ecological variation across a species' range can select for local adaptation within populations, which can contribute to the maintenance of genetic variation by favoring different alleles across the range (Levene, 1953; Felsenstein, 1976; Hedrick *et al.*, 1976; Kawecki & Ebert, 2004). In addition, heterogeneous selection pressures that are distributed as a gradual continuum from one end of the species' range to another can create a cline of genetic variation responsible for phenotypes under selection (Slatkin, 1973; Endler, 1977). Some of the best examples of latitudinal clines come from temperature-dependent phenotypes (e.g., body size, developmental rate, and thermal tolerance) that have been well-documented in flies (Partridge *et al.*, 1994; Eanes, 1999; Robinson & Partridge, 2001; Hoffmann *et al.*, 2002). Moreover, heterogeneous selection pressures across a cline may affect males and females differently (Connallon, 2015; Connallon *et al.*, 2019), although the empirical evidence for such variation in sex-specific selection across geographic ranges is mixed (Delcourt *et al.*, 2009; Delph *et al.*, 2011; Allen *et al.*, 2017; Lasne *et al.*, 2018).

Thermal adaptation within populations and across a species range can occur via selection on physiological, anatomical, or behavioral traits. For example, north-south gradients in heat and cold tolerance have been observed in *Drosophila* (Hoffmann *et al.*, 2002), suggesting physiological adaptation to thermal environments. In addition, ectotherms, such as flies, rely on behavioral mechanisms of thermoregulation by avoiding suboptimal temperatures in search of more optimal ones (Dillon *et al.*, 2009; Kearney *et al.*, 2009), and thermal preference may be correlated with optimal thermal performance (Dawson, 1975; Angilletta *et al.*, 2002).

Concordance across genotypes between different thermal traits could reinforce the response to selection, whereas negative correlations could constrain adaptation (Etterson & Shaw,

2001). However, it is not clear if physiological and behavioral thermal traits are genetically

correlated within a species, between sexes, or across populations (Dawson, 1975; Angilletta *et*

al., 2002; Gilbert & Miles, 2017). For example, experiments in *Drosophila subobscura* identified

individual chromosomes that affected thermal tolerance or temperature preference, but no single

chromosome affected both physiological and behavioral phenotypes (Dolgova *et al.*, 2010; Rego

et al., 2010; Castañeda *et al.*, 2019). Furthermore, temperature-dependent traits can affect

assortative mating and male reproductive success (Dolgin *et al.*, 2006; Keller & Seehausen,

2012), suggesting inter-sexual differences in thermoregulation could affect genetic variation

within populations via sexual selection. These sex-specific selection pressures could also

contribute to the maintenance of genetic variation via inter-sexual conflict or context-dependent

selection (Kotiaho *et al.*, 2001; Rostant *et al.*, 2015; Meisel, 2018). Despite the importance of

inter-sexual differences, previous work did not test for differences in the genetic correlation of

thermal traits between males and females.

We used a sex chromosome polymorphism in the house fly, *Musca domestica*, to

investigate the concordance of thermal tolerance and preference across clinally distributed male

genotypes. House fly has a polygenic sex determination system, in which a male-determining

gene has been mapped to all six chromosomes, some males can carry multiple male-determining

chromosomes, and a female-determining allele segregates on one chromosome (McDonald *et al.*,

1978; Inoue & Hiroyoshi, 1986; Dübendorfer *et al.*, 2002; Hediger *et al.*, 2010). The *M.*

domestica male determiner (*Mdmd*) gene is most commonly found on either the third

chromosome (III^M) or what was historically referred to as the Y chromosome (Y^M) (Hamm *et al.*,

2014; Sharma *et al.*, 2017). Both III^M and Y^M are very young proto-Y chromosomes that are

minimally differentiated from their homologous proto-X chromosomes (Meisel *et al.*, 2017; Son

et al., 2019; Son & Meisel, 2020). Y^M and III^M are distributed along latitudinal clines on multiple

continents in the Northern Hemisphere (Tomita & Wada, 1989; Hamm *et al.*, 2005; Kozielska *et*

al., 2008). Y^M is most frequently found at northern latitudes, and III^M is more common at

southern latitudes (Figure 1A). This distribution suggests that the Y^M chromosome confers higher

fitness in colder climates, and, conversely, III^M confers higher fitness in hotter climates.

Therefore, variation in temperature across the species range may create heterogeneous selection

pressures that maintain the proto-Y chromosome cline in house fly. Consistent with this

hypothesis, seasonality in temperature is the best predictor of the frequencies of the proto-Y

chromosomes across natural populations (Feldmeyer *et al.*, 2008).

109   We tested the hypothesis that the Y^M chromosome confers cold-adaptive phenotypes and

III^M confers heat-adaptive phenotypes in house fly males, which would be consistent with their

latitudinal distributions (Tomita & Wada, 1989; Hamm *et al.*, 2005; Feldmeyer *et al.*, 2008;

Kozielska *et al.*, 2008). To those ends, we first evaluated if males carrying the III^M chromosome

(hereafter III^M males) have greater tolerance to extreme heat and if males carrying the Y^M

chromosome (Y^M males) have greater cold tolerance. Second, we tested if III^M males prefer

warmer temperatures than Y^M males, and if males and females differ in their thermal preference.

We performed all experiments using flies raised at multiple developmental temperatures because

thermal acclimation strongly affects temperature-dependent phenotypes in flies and other

ectotherms (Krstevska & Hoffmann, 1994; Dillon *et al.*, 2009). Together, we evaluated if thermal

preference and tolerance are aligned for sex-linked genetic variants, tested if this alignment is

consistent with the geographic distribution of the proto-Y chromosomes, and then discuss how

these temperature-dependent phenotypes could affect the access of males to female mates.

Materials and methods

Fly strains and rearing

We performed our experiments using five nearly isogenic house fly strains, three with III^M males and two with Y^M males (Supplementary Methods). All five strains have a common genetic background from an inbred III^M strain that was produced from a mixture of flies collected across the United States (Scott *et al.*, 1996; Hamm *et al.*, 2005). Each of the three III^M strains carries a different III^M chromosome from a separate wild-derived line, and, likewise, the two Y^M strains carry different Y^M chromosomes. Each strain is fixed for its proto-Y chromosome (either III^M or Y^M), and no other sex determiners, such as the female-determining $Md\text{-}tra^D$ allele (Hediger *et al.*, 2010), segregate within these strains.

We reared each strain at 18°C, 22°C, and 29°C for two generations in order to evaluate how thermal acclimation affects thermal tolerance (Chown & Terblanche, 2006) and thermal preference (Krstevska & Hoffmann, 1994; Dillon *et al.*, 2009). Flies from each developmental temperature were assayed at equivalent physiological ages estimated by accumulated degree days (Barnard & Geden, 1993; Wang *et al.*, 2018). For our heat and cold tolerance assays, we used flies 22–50 total degree days after eclosion. For thermal preference assays, we used flies 96–115 total degree days after eclosion. Additional details and calculations are provided in the Supplementary Methods.

Thermal tolerance

We measured heat and cold tolerance in individual male and female house flies. To measure heat tolerance, lightly anaesthetized individual flies were transferred to a 1.5 ml centrifuge tube that was sealed with fabric. We placed the 1.5 ml tube in a heat block set to 53°C.

This temperature was selected because it is the lowest at which heat tolerance could be measured

in a reasonable period of time. The time at which a fly fell to the bottom of the tube and could not

make its way back to the top was considered the knockdown time. To measure cold tolerance,

lightly anaesthetized flies were transferred to a fabric-sealed 20 ml glass vial individually, and the

vials were placed in a 4°C refrigerator with a transparent door. Knockdown occured when a fly

fell on its back to the bottom of the vial. We gently tapped the assay vial every 2–3 minutes to

ensure flies were active.

For both heat and cold tolerance assays, we performed an analysis of variance (ANOVA)

using the lmer() function in the lme4 (v1.1) R package (Bates *et al.*, 2015) to model the effect of

genotype (G: Y^M vs III^M), developmental temperature (T: 18°C or 29°C), and their interaction on

knockdown time (K):

$$K \sim G + T + G{\times}T + B + S,$$

with experimental batch (B) and strain (S) treated as random effects. We also constructed another

model excluding the interaction term:

$$K \sim G + T + B + S.$$

We then used a drop in deviance test to compare the fit of the models with and without the

interaction term using the anova() function in R. We also compared heat and cold tolerance

between males raised at 22°C and 29°C, using the same approaches as described above. As the

thermal tolerance comparisons between flies raised at 18°C and 29°C and between flies raised at

22°C and 29°C were conducted in separate experimental batches, we analyzed each comparison

separately. Additional details are provided in the Supplementary Methods.

Thermal preference

We measured thermal preference as the position of individual flies along a 17–37°C

thermal gradient (Figure S1), following a slightly modified version of previous protocols

(Anderson *et al.*, 2013; Lynch *et al.*, 2018). For each individual fly, we report mean thermal

preference (T_{pref}) as the average position during a 10 minute assay window (measured once per

minute). We also report thermal breadth, $T_{breadth}$ (Carrascal *et al.*, 2016), as the coefficient of

variation of individual-level T_{pref} during the assay window. $T_{breadth}$ provides an estimate of how

individuals utilize thermal space within their environment (Slatyer *et al.*, 2013). Choosier

individuals show a lower $T_{breadth}$ value and, thus, would be expected to occupy a narrower range

of temperatures within a given thermal habitat.

To determine the effects of developmental temperature (18°C, 22°C, and 29°C), genotype

(Y^M and III^M), and their interaction on mean T_{pref} across sexes, we created a mixed-effects model

using the lme4 package (v1.1) in R (Bates *et al.*, 2015). Developmental temperature, genotype,

and their interaction were included as fixed effects, and strain, batch, and lane in the thermal

gradient (L) were included as random effects:

180                        $$T_{pref} \sim G + T + G{\times}T + B + S + L.$$

We did the same for $T_{breadth}$. We then determined whether groups significantly differed in T_{pref} or

$T_{breadth}$ using Tukey contrasts with the multcomp package (v1.4) in R (Hothorn *et al.*, 2008).

Within developmental temperature treatments, we used Bayesian information criterion (BIC)

scores from the mclust (v5.4.5) package in R (Scrucca *et al.*, 2016) to determine whether the

distribution of individual measures of T_{pref} within a group are best explained by one or multiple

normal distributions.

Results

Thermal tolerance depends on developmental temperature and male genotype

We measured extreme heat (53°C) and cold (4°C) tolerance as a readout of differences in

physiological thermal adaptation between Y^M and III^M house fly males. We observed the

expected effect of acclimation on both heat and cold tolerance (Chown & Terblanche, 2006): flies

raised at 18°C tolerate cold longer than the flies raised at 29°C, and flies raised at 29°C tolerate

heat longer than flies raised at 18°C (Figure 1). We also find that Y^M males are more cold

tolerant, and III^M males are more heat tolerant, consistent with the latitudinal distributions of Y^M

and III^M males in nature (Tomita & Wada, 1989; Hamm *et al.*, 2005; Feldmeyer *et al.*, 2008;

Kozielska *et al.*, 2008). However, the effect of genotype on thermal tolerance depends on

acclimation temperature. Specifically, a linear model with an interaction between genotype (Y^M

or III^M) and developmental temperature fits the cold tolerance data significantly better than a

model without the interaction term ($\chi^2_1 = 19.3$, $p = 1.1 \times 10^{-5}$). This provides evidence for a G×T

effect on cold tolerance—Y^M males are more cold tolerant than III^M males, but only if they are

raised at 18°C (Figure 1B). There is also a significant G×T interaction affecting heat tolerance

($\chi^2_1 = 4.71$, $p = 0.030$ comparing models with and without the interaction term): III^M males are

more heat tolerant than Y^M males, but only if raised at 29°C (Figure 1D).

Figure 1 - Thermal tolerance in males and females. (**A**) Map of the eastern United States,

showing the cline of Y^M (more common in the north) and III^M (more common in the south). (**B–**

E) Graphs show the effect of developmental temperature on knockdown time at either 4°C (cold

tolerance) or 53°C (heat tolerance) for Y^M (turquoise) and III^M (salmon) male flies. Proto-Y

chromosome labels for females reflect whether males from the strain carry the Y^M or III^M

chromosome. Mean knockdown time is plotted for each combination of genotype and

temperature. Error bars represent standard error.

We next attempted to identify a threshold temperature for the genotype-specific benefits

of acclimation by comparing heat and cold tolerance of flies raised at 22°C and 29°C (instead of

18°C and 29°C). We did not observe a significant effect of the interaction between developmental

temperature and male genotype on extreme cold tolerance ($\chi^2_1 = 0.947$, $p = 0.331$ comparing

models with and without an interaction term) (Figure S2). We therefore hypothesize that there is

a threshold temperature between 18°C and 22°C, below which Y^M males experience a greater

benefit of cold acclimation than III^M males. In contrast, there is a significant interaction between

genotype and developmental temperature on heat tolerance when comparing males raised at 22°C

and 29°C ($\chi^2_1 = 11.02$, $p = 9.0 \times 10^{-4}$ comparing models with and without the interaction term)

(Figure S2). Therefore, the threshold for a genotype-specific benefit from heat acclimation lies

between 22°C and 29°C.

We do not expect any difference in heat or cold tolerance across females from our

different strains because all females have the same genotype, regardless of the male genotype in

the strain. Indeed, a model with an interaction between developmental temperature and male

genotype does not fit the female cold tolerance data better than a model without the interaction

term ($\chi^2_1 = 1.46$, $p = 0.23$) (Figure 1C). There is a significant effect of developmental temperature

on cold tolerance in females ($\chi^2_1 = 43.5$, $p = 4.3 \times 10^{-11}$ comparing a model with and without

developmental temperature), demonstrating that females benefit from cold acclimation regardless

of male genotype (Figure 1C). Surprisingly, there is a significant interaction between male

genotype and developmental temperature on heat tolerance in females ($\chi^2_1 = 10.4$, $p = 0.0013$

comparing a model with and without the interaction term). In general, females raised at warmer

temperatures are more heat tolerant (Figure 1E). However, the interaction of male genotype and

developmental temperature is in the opposite direction from what would be expected based on the

latitudinal distribution of Y^M and III^M: females from strains with Y^M males that are raised at 29°C

are more heat tolerant than females from III^M strains raised at 29°C (Figure 1E). We thus

conclude that the heat and cold tolerance differences between Y^M and III^M males are specific to

males and/or the proto-Y chromosomes (i.e., not genetic background) because we do not observe

the same heat or cold tolerance differences in females from those strains (who do not carry the

proto-Y chromosomes).

Figure 2 - Thermal preference (T_{pref}) of (**A**) male and (**B**) female house flies according to male

genotype (III^M = salmon points and line, Y^M = turquoise points and line) and developmental

temperature. Each point depicts the mean thermal preference for an individual fly, with lines and

error bars denoting means within groups and standard errors of the mean, respectively.

Significant differences between groups are denoted by letters, with differing letters highlighting

significantly different mean thermal preferences within each graph (Tukey's *post hoc* test, *p* <

0.05).

Thermal preference depends on developmental temperature and male genotype

We next tested if genotype and developmental temperature affect thermal preference

(T_{pref}). First, we find that T_{pref} is inversely proportional to developmental temperature (Figure 2),

with house flies that develop at a warmer temperature preferring cooler temperatures (and *vice*

versa), regardless of sex (male: $F_{2, 742.7} = 138.4$, *p* < 1.0 x 10^{-5}; female: $F_{2, 245.3} = 37.1$, *p* = 1.19 x

10^{-4}; Figure 2). This is consistent with how developmental acclimation affects T_{pref} in *Drosophila*

(Dillon *et al.*, 2009).

We also find that male proto-Y chromosome genotype (Y^M vs III^M) affects T_{pref} ($F_{1, 756.2}$=

44.5, *p* < 1.0 x 10^{-5}). There is also a significant interaction effect between developmental

temperature and genotype on T_{pref} in males ($F_{2, 756.3} = 8.47$, $p = 2.31$ x 10^{-4}, Figure 2A). Male T_{pref}

is similar across genotypes when they develop at either 18°C or 29°C. However, when reared at

22°C, III^M males prefer warmer temperatures than Y^M males (Tukey's *post hoc* test, $p < 0.001$).

This is consistent with III^M males being more common at lower latitudes (where average

temperatures are warmer), and Y^M males more common at higher latitudes (Tomita & Wada,

1989; Hamm *et al.*, 2005; Feldmeyer *et al.*, 2008; Kozielska *et al.*, 2008). We do not expect

differences in T_{pref} in females across strains because all females have the same genotype. Indeed,

the genotype of males in a strain (Y^M vs III^M) and the interaction between male genotype and

female developmental temperature showed no significant effect on T_{pref} in females (ANOVA, all

$p > 0.1$ in Figure 2B). We assayed more males than females in our thermal preference

experiments, and so we repeated our analysis by down-sampling the data to have equal numbers

of individuals across treatments. The down-sampled data give equivalent results to the full data

set (Supplementary Results).

Thermal breadth depends on sex and thermal preference

We used thermal breadth ($T_{breadth}$) as a measure of the specificity of T_{pref}. Male $T_{breadth}$ was

not significantly affected by either developmental temperature, genotype, or the interaction

between genotype and developmental temperature (ANOVA, all $p > 0.1$, Figure S3A). In

contrast, developmental temperature ($F_{2, 236.9} = 16.5$, $p < 1.0$ x 10^{-5}), as well as the interaction

between developmental temperature and male genotype ($F_{2, 243.6} = 5.35$, $p = 0.005$), had

significant effects on $T_{breadth}$ in females. However, the significant interaction is of small effect, as

females from strains with differing male genotypes do not significantly differ in $T_{breadth}$ within

any developmental temperature treatment (Figure S3B).

Figure 3 - Thermal breadth ($T_{breadth}$) depends on male genotype and sex. (**A**) Distribution of individual-level mean thermal preferences (T_{pref}) of III^M males, Y^M males, and pooled females that developed at 22°C. Y-axis represents relative density of data points and is analogous to frequency of data points for a given T_{pref} value. (**B**) $T_{breadth}$ of individuals raised at 22°C according to group (F^C = cold-preferring females, F^W = warm-preferring females). Boxplots denote median values and lower- and upper- quartiles. Asterisks denote significant differences in $T_{breadth}$ between groups (***: Tukey's *post hoc* test, $p < 0.01$).

The effect of developmental temperature on $T_{breadth}$ in females is driven by increased variance in T_{pref} when females develop at 22°C. The increased variance in female T_{pref} can be explained by a mixture of two normal distributions (Figure 3A, see Table S1 for statistics). This bimodal distribution is not a result of differences across strains because the same pattern was observed among females separately analyzed based on male genotype (Figure S4). In comparison, a single normal distribution best fit Y^M male T_{pref} when developed at 22°C, and two normal distributions best explained the III^M male T_{pref} when developed at 22°C. Upon inspection, however, the two distributions representing III^M male T_{pref} likely correspond to the tail (mean of

28.7°C and large variance of 10.4°C) and peak (mean of 32.6°C and small variance of 0.4°C) of a

single skewed distribution, which we are unable to detect using the mclust package we used to fit

distributions to our data.

We used our model-based clustering analysis of T_{pref} to classify individuals that developed

at 22°C into one of four groups: Y^M males (lower T_{pref}), III^M males (higher T_{pref}), females with

cooler T_{pref} (F^C females, 59.3% of females tested), and females with warmer T_{pref} (F^W females,

40.7% of females tested). The mean T_{pref} of F^C females (26.90°C) is nearly equal to the mean T_{pref}

of Y^M males (26.87°C; Figure 3A). Similarly, the mean T_{pref} of F^W females (32.2°C) is near the

mode of the T_{pref} of III^M males (32.0–32.5°C; Figure 3A).

We further find that T_{pref} is predictive of $T_{breadth}$ for flies that develop at 22°C. We

considered flies from our four T_{pref} groups (Y^M males, III^M males, F^C females, and F^W females),

and we found a significant effect of group on $T_{breadth}$ ($F_{3, 32.9} = 9.40$, $p = 1.24$ x 10^{-4}). Specifically,

F^C females have significantly greater $T_{breadth}$ than all other groups (Tukey's *post hoc* test, all $p <$

1.0 x 10^{-5}, Figure 3B). Therefore, if we consider $T_{breadth}$ as a measure of the strength of T_{pref}, adult

house flies can be summarized by one of three phenotypes related to thermal behavior when

developed at 22°C: a relatively strong preference for warm temperatures (III^M males and F^W

females, which have high T_{pref} and low $T_{breadth}$), a strong preference for cooler temperatures (Y^M

males, with low T_{pref} and low $T_{breadth}$), and a relatively weak preference for cooler temperatures

(F^C females, with low T_{pref} and high $T_{breadth}$). Down-sampling the data gives similar results as the

full data set (Supplementary Material).

Discussion

We tested if thermal tolerance and preference depend on sex and male genotype in house

flies. We find that males carrying the Y^M chromosome (which is common in the northern end of

the species' range) are more cold tolerant and prefer colder temperatures. Conversely, males

carrying the III^M chromosome (which is common in the southern end of the species' range) are

more heat tolerant and prefer warmer temperatures. Our results are therefore consistent with the

general trend that temperate populations are typically more cold-tolerant than (sub-) tropical ones

(Gibert & Huey, 2001; Hoffmann *et al.*, 2002). The differences in thermal preference are

consistent with the idea that behavioral thermoregulation can weaken selection for thermal

tolerance, as predicted by the "Bogert Effect" (Huey *et al.*, 2003; Huey & Pascual, 2009;

Castañeda *et al.*, 2013). However, the fact that thermal preference and tolerance are both

predicted by male genotype provides evidence that these traits are responsive to selection,

suggesting any Bogert effects are not sufficient to overwhelm thermal adaptation. These

differences in thermal tolerance and preference in males depend on developmental temperature,

and they are not observed in congener females from the same strains (who do not carry the Y^M or

III^M chromosome). However, females exhibit a bimodal T_{pref}, with females from each of the two

subgroups overlapping with one of the male genotypes.

Thermal tolerance and preference depend on developmental temperature, genotype, and sex

Our results demonstrate, to the best of our knowledge, the first documented example of

concordant temperature preference, cold tolerance, and heat tolerance across genotypes within a

species. We find that Y^M males both have greater cold tolerance and prefer colder temperatures,

whereas III^M males have greater heat tolerance and prefer warmer temperatures (Figures 1 and 2),

consistent with their latitudinal distributions (Tomita & Wada, 1989; Hamm *et al.*, 2005;

Feldmeyer *et al.*, 2008; Kozielska *et al.*, 2008). Previous work has identified concordant T_{pref} and

heat tolerance differences across species (Qu *et al.*, 2011), or found no clear relationship between

thermal tolerance and preference across genotypes within species (Yang *et al.*, 2008; Rego *et al.*,

2010; Castañeda *et al.*, 2019). Body size is also predicted to vary with thermal traits (Leiva *et al.*,

2019). In our study, we did not measure insect body size. While we did not observe any obvious

differences between strains, it is possible that some of the genotypic effect on thermal tolerance

or preference we observed is due to (temperature-dependent) morphological differences between

Y^M and III^M males. Future studies should directly test this hypothesis.

We observed strong effects of developmental temperature on both thermal tolerance and

preference that depend on both genotype and sex. Acclimation effects on heat and cold tolerance

(Figure 1) are well-documented for ectotherms, including flies and other insects (Bowler &

Terblanche, 2008). An inverse relationship between developmental temperature and thermal

preference has also been observed in other flies (Dillon *et al.*, 2009; Castañeda *et al.*, 2013).

Behaviorally navigating towards compensatory temperatures could serve as a means of mitigating

the costs of thermally suboptimal development (i.e., too hot or too cold). The observed

relationships between thermal tolerance and developmental temperatures are likely to be caused

by acclimation and unlikely to be the result of natural selection within our experiment for two

reasons. First, there is unlikely to be sufficient genetic variation in these inbred strains for

selection to generate these results within 2 generations. Second, prior attempts at selecting for

thermal tolerance in house flies resulted in negligible differences in tolerance across

developmental temperatures (Geden *et al.*, 2019) However, it is worth noting that the males used

by Geden et al. (2019) were likely all III^M based on their geographic origin. Had the experimental

population consisted of both III^M and Y^M males, a response to tolerance may have been detected.

We conclude that the differences in thermal tolerance (and preference) between Y^M and III^M

males have evolved across the natural populations from which we sampled the Y^M and III^M

chromosomes.

There are important methodological implications for our observation that variation in

thermal preference across genotypes depends on developmental temperature. We only observe

warmer (colder) thermal preferences in III^M (Y^M) males when developed at 22°C; thermal

preference did not differ between male genotypes when raised at more extreme (18°C, 29°C)

temperatures (Figure 2A). Previous studies attempting to estimate genetic variance in thermal

preference within or among populations of *Drosophila* have had mixed results. While some

studies identified genetic variance among populations within species (Good, 1993; Castañeda *et*

al., 2013), others did not detect substantial variance within (Krstevska & Hoffmann, 1994) or

among species (MacLean *et al.*, 2019). Our results show that the phenotypic presentation of

genetic variation for thermal preference can depend on the environmental conditions experienced,

which could explain why this variance was not detected in other experiments. In addition, while

the genetic mechanisms that regulate thermal tolerance in other systems have been extensively

studied (Svetec *et al.*, 2011; Königer & Grath, 2018; Königer *et al.*, 2019), it is possible that

some of the molecular pathways involved will only be revealed through experiments conducted

across developmental temperatures.

We identify multiple differences between males and females in their thermal tolerance

and preferences. The strain differences we observed are primarily limited to males, which is

expected because the males differ in genotypes (Y^M and III^M) but females are isogenic (Meisel *et*

al., 2015). However, there is a difference in heat tolerance between females from strains with Y^M

males and females from strains with III^M males (Figure 1). While we can rule out certain

genotypic explanations for this difference (i.e., all females are isogenic and do not carry *Md-*

tra^D), we do not yet have a mechanistic explanation on why females show the opposite

developmental heat tolerance from males. Nevertheless, the difference in heat tolerance observed

between females from different strains is in the opposite direction as between Y^M and III^M males

from those strains. This helps us to conclude that differences between Y^M and III^M males are

indeed a result of different proto-Y chromosomes rather than their genetic backgrounds. In other

words, the difference in heat tolerance between females is an exception that proves the rule with

respect to the effects of proto-Y chromosomes on male thermal tolerance and preference.

We identified a female-specific plasticity for thermal preference that does not map to male

genotype. In females, we found that neither thermal tolerance nor thermal preference differ

predictably between strains where males carry different proto-Y chromosomes (Figures 1C, E and

2B). However, there is a bimodal thermal preference for females that develop at 22°C (Figure

3A), regardless of congener male genotype. In addition, females that had colder T_{pref} when

developed at 22°C also had a larger $T_{breadth}$ (Figure 3B). In small ectotherms with little thermal

inertia, measures of movement along a thermal gradient (such as $T_{breadth}$) are predicted to be

positively correlated with environmental temperature (Anderson *et al.*, 2007). However, we

observe the opposite relationship between mean environmental temperature (T_{pref}) and $T_{breadth}$ in

females (Figure 3), suggesting that the difference in $T_{breadth}$ cannot be explained by thermal

inertia. Our results suggest that, in nature, females with colder temperature preferences may

occupy a wider range of temperatures than females with warmer temperature preferences.

Because all females in our experiment are expected to have the same genotype, we hypothesize

that these differences in T_{pref} and $T_{breadth}$ are conferred by a plastic response to some yet to be

characterized factor (e.g., microclimates within larval rearing containers). Alternatively, this

plasticity could have a stochastic origin that is intrinsic to the development of thermal preference

(Honegger & de Bivort, 2018; Jensen, 2018).

The correlation between thermal preference and thermal breadth at 22°C is female-

specific: Y^M and III^M males have similar $T_{breadth}$ values when raised at 22°C despite their

differences in T_{pref}. Although general sex differences in thermal tolerance (Hoffmann et al. 2005)

and thermal preference (Krstevska and Hoffmann 1994) have been documented, this is the first

study, to our knowledge, to identify sex differences in the relationship between thermal

preference and thermal breadth. Our results suggest that male and female house flies exhibit

different thermoregulatory behavioral patterns which may further be influenced by genotype.

Directly identifying a sex-by-genotype-by-environment interaction is beyond the scope of this

study because sex and genotype are confounded in our experimental design (the females in our

experiment have a different genotype from either male, characterized by a lack of either the III^M

or Y^M chromosome). Nonetheless, the house fly is a tractable system for directly testing for sex-

specific genotype-by-environment interactions on thermoregulation. For example, future work

could test for sex-specific effects of Y^M and III^M by measuring phenotypes in females carrying a

proto-Y chromosome along with the epistatic female-determining *Md-tra^D* allele (Hediger *et al.*,

2010; Hamm *et al.*, 2014).

Environmental heterogeneity and the maintenance of polygenic sex determination

Sex determination pathways rapidly diverge across species, driving evolutionary turnover

of sex chromosomes (Bull, 1983; Beukeboom & Perrin, 2014). Polygenic sex determination

systems, in which more than one master sex determining locus segregate independently on

different chromosomes, have been observed in multiple animal species (Moore & Roberts, 2013).

Most population genetic models that attempt to explain the stable maintenance of polygenic sex

determination focus on sexually antagonstic effects of sex determining loci or linked alleles on

sex chromosomes (Rice, 1986; van Doorn & Kirkpatrick, 2007; Kozielska *et al.*, 2010; Meisel *et*

al., 2016). Less attention has been given to ecological factors that can maintain polygenic sex

determination (Pen *et al.*, 2010; cf. Bateman & Anholt, 2017).

Our results demonstrate how spatially variable ecological factors can maintain polygenic

sex determination. Specifically, thermal tolerance and preference phenotypes conferred by the Y^M

and III^M chromosomes (Figures 1 and 2) are consistent with the clinal and temperature-dependent

distributions of the Y^M and III^M chromosomes (Tomita & Wada, 1989; Hamm *et al.*, 2005;

Feldmeyer *et al.*, 2008; Kozielska *et al.*, 2008). Previous experiments identified multiple fitness

advantages conferred by the III^M chromosome over Y^M at warmer temperatures, including an

increase in frequency of III^M over generations in a laboratory population (Hamm *et al.*, 2009).

However, these fitness differences can only explain the invasion or fixation of the III^M

chromosome, not the maintenance of the polymorphism. In contrast, differences in thermal

tolerance and preference could maintain proto-Y chromosome polymorphism across the species'

range, similar to how selection maintains other clinal variation (Slatkin, 1973; Endler, 1977).

The house fly system reveals how temperature variation can contribute to the maintenance

of polygenic sex determination independently of selection on the sex-determination pathway

itself. Temperature is an important contributor to the evolution of sex determination pathways in

vertebrates (Bull & Vogt, 1979; Holleley *et al.*, 2015). However, the effects of the house fly

proto-Y chromosomes on thermal tolerance and preference likely act independently of the sex

determination pathway because there are not differences in the expression of sex determination

genes across house fly male genotypes raised at different temperatures in a way that is consistent

with their clinal distribution (Adhikari *et al.*, n.d.). This suggests that the effects of the Y^M and

III^M chromosomes on thermal phenotypes is a result of alleles on proto-Y chromosomes that are

genetically linked to the male-determining locus, as opposed to the male-determiner itself.

Therefore, our results highlight how temperature can be important for the evolution of sex

determination independently of temperature-dependent activity of the sex determination pathway.

Future theoretical work should consider the effect of spatially heterogeneous selection pressures

on the maintenance of polygenic sex determination, similar to how temporal heterogeneity can

create fluctuating selection pressures that maintain polygenic sex determination (Bateman &

Anholt, 2017).

Selection on thermal phenotypes may depend on geographical scale

Our results suggest that selection on thermal traits differs between macro-geographic

species ranges and at a micro-geographical scale within populations. Similar differences in

selection pressures according to geographic scale have been documented before in other species

(Richter-Boix *et al.*, 2010; De Block *et al.*, 2013; Tüzün *et al.*, 2017). Thermal tolerance and

preference in male house flies depend on proto-Y chromosomes genotype in a way that is

consistent with the latitudinal distribution of the Y^M and III^M chromosomes (Figures 1 and 2).

This suggests that, at the macro-geographic scale, selection is operating on male physiology and

behavior to create or maintain the clinal distribution of Y^M and III^M (Figure 4A). It is also worth

noting that our study focuses on only two male genotypes (III^M and Y^M). While these are the most

prevalent genotypes in the eastern United States, other genotypes exist (including males with

multiple proto-Y chromosomes, and females with proto-Y and proto-W chromosomes) and are

common in other populations (Franco *et al.*, 1982; Feldmeyer *et al.*, 2008; Hamm & Scott, 2009;

Hamm *et al.*, 2014). Future studies should characterize thermal tolerance and preference of these

other genotypes in order to determine whether their geographical distribution is similarly

explained by thermal biology.

At an intermediate developmental temperature (22°C), female thermal preference is

bimodal for a reason that we have yet to determine (Figure 3A). This raises the possibility that

within populations near the center of the cline (i.e., at a micro-geographic scale), where Y^M and

III^M both segregate (e.g., Hamm & Scott, 2008; Meisel *et al.*, 2016), sexual selection may favor

males that can preferentially obtain access to the two different female phenotypes. While

differences in thermal preference probably did not evolve in response to sexual selection, these

differences do likely have important consequences on the reproductive success of III^M and Y^M

males where they co-occur. III^M males may disproportionately benefit from differences in T_{pref}

and $T_{breadth}$ between males and females. F^C females that prefer colder temperatures have greater

$T_{breadth}$ than warm preferring F^W females and both male genotypes (Figure 3B), suggesting that F^C

females occupy a wider range of thermal habitats. Thus, III^M males may gain an advantage by

having greater access to F^W females, as well as occasional access to F^C females, in contrast to Y^M

males who would only be likely to encounter F^C females (Figure 4B). This raises the possibility

that differences in thermal preference across genotypes and sexes could affect the dynamics of

sexual selection.

Figure 4 - Selection on the III^M and Y^M chromosomes likely differs across geographic scales. (**A**)

At the macro-geographical scale, selection for thermal tolerance and/or thermal preference results

in the clinal distribution of the Y^M (turquoise) and III^M (salmon) chromosomes. (**B**) At

intermediate developmental temperatures, male genotypes (Y^M vs III^M) differ in thermal

preference, which may create asymmetrical mating opportunities because of variation in female

thermal preference and breadth (F^C vs F^W). The asymmetry of the overlap of males and females at

the intermediate developmental temperature could affect sexual selection in populations where

Y^M and III^M both segregate.

Acknowledgements

Tin Nguyen, Chantalle Vincent, and Mandy Cao assisted with experiments. This material

is based upon work supported by the National Science Foundation under Grant No. DEB-

1845686. Any opinions, findings, and conclusions or recommendations expressed in this material

are those of the authors and do not necessarily reflect the views of the National Science

Foundation.

Author Contributions

PD, KA, and RM conceived and designed the study. KA, OH, VS, and JC collected, and

KA analyzed, all thermal tolerance data. PD, RP, JT, and AM collected, and PD analyzed, all

thermal preference data. PD, KA, and RM wrote the manuscript, and all authors reviewed the

manuscript prior to submission.

Data Accessibility

All data files used for analyses described in this manuscript have been deposited in Dryad

(doi:10.5061/dryad.n2z34tmvs). Raw video and image files from thermal preference assays are

available from the authors upon request.

References

Adhikari, K., Son, J.H., Rensink, A.H., Jaweria, J., Bopp, D., Beukeboom, L.W., *et al.* (n.d.) Temperature-

dependent effects of house fly proto-Y chromosomes on gene expression act independently of the sex

determination pathway.

Allen, S.L., Bonduriansky, R., Sgro, C.M. & Chenoweth, S.F. (2017) Sex-biased transcriptome divergence

along a latitudinal gradient. *Molecular ecology*, **26**, 1256–1272.

Anderson, J.L., Albergotti, L., Proulx, S., Peden, C., Huey, R.B. & Phillips, P.C. (2007) Thermal

preference of *Caenorhabditis elegans*: a null model and empirical tests. *The Journal of experimental*

biology, **210**, 3107–3116.

Anderson, R.D., Blanford, S. & Thomas, M.B. (2013) House flies delay fungal infection by fevering: at a

cost. *Ecol. Entomol.*, **38**, 1–10.

Angilletta, M.J., Hill, T. & Robson, M.A. (2002) Is physiological performance optimized by

thermoregulatory behavior?: a case study of the eastern fence lizard, *Sceloporus undulatus*. *J. Therm.*

Biol., **27**, 199–204.

Barnard, D.R. & Geden, C.J. (1993) Influence of larval density and temperature in poultry manure on

development of the house fly (Diptera: Muscidae). *Environmental entomology*, **22**, 971–977.

Bateman, A.W. & Anholt, B.R. (2017) Maintenance of polygenic sex determination in a fluctuating

environment: an individual-based model. *Journal of evolutionary biology*, **30**, 915–925.

Bates, D., Mächler, M., Bolker, B. & Walker, S. (2015) Fitting linear mixed-effects models using lme4. *J.*

Stat. Soft., **67**, 1–48.

Beukeboom, L.W. & Perrin, N. (2014) *The Evolution of Sex Determination*. Oxford University Press, New

York, NY.

Bowler, K. & Terblanche, J.S. (2008) Insect thermal tolerance: what is the role of ontogeny, ageing and

senescence? *Biological reviews of the Cambridge Philosophical Society*, **83**, 339–355.

Bull, J.J. (1983) *Evolution of Sex Determining Mechanisms*. Benjamin/Cummings, Menlo Park, CA.

Bull, J.J. & Vogt, R.C. (1979) Temperature-dependent sex determination in turtles. *Science*, **206**, 1186–

1188.

Carrascal, L.M., Villén-Pérez, S. & Palomino, D. (2016) Preferred temperature and thermal breadth of

birds wintering in peninsular Spain: the limited effect of temperature on species distribution. *PeerJ*, **4**,

e2156.

Castañeda, L.E., Balanyà, J., Rezende, E.L. & Santos, M. (2013) Vanishing chromosomal inversion clines

in *Drosophila subobscura* from Chile: is behavioral thermoregulation to blame? *The American naturalist*,

182, 249–259.

Castañeda, L.E., Romero-Soriano, V., Mesas, A., Roff, D.A. & Santos, M. (2019) Evolutionary potential

of thermal preference and heat tolerance in *Drosophila subobscura*. *Journal of evolutionary biology*, **32**,

818–824.

Chown, S.L. & Terblanche, J.S. (2006) Physiological diversity in insects: ecological and evolutionary

contexts. *Adv. Insect Physiol.*, **33**, 50–152.

Connallon, T. (2015) The geography of sex-specific selection, local adaptation, and sexual dimorphism.

Evolution; international journal of organic evolution, **69**, 2333–2344.

Connallon, T., Sharma, S. & Olito, C. (2019) Evolutionary consequences of sex-specific selection in

variable environments: four simple models reveal diverse evolutionary outcomes. *The American*

naturalist, **193**, 93–105.

Dawson, W.R. (1975) On the physiological significance of the preferred body temperatures of reptiles.

Persp. of Biophys. Ecol., 443–473.

De Block, M., Pauwels, K., Van Den Broeck, M., De Meester, L. & Stoks, R. (2013) Local genetic

adaptation generates latitude-specific effects of warming on predator-prey interactions. *Global change*

biology, **19**, 689–696.

Delcourt, M., Blows, M.W. & Rundle, H.D. (2009) Sexually antagonistic genetic variance for fitness in an

ancestral and a novel environment. *Proceedings. Biological sciences / The Royal Society*, **276**, 2009–2014.

Delph, L.F., Andicoechea, J., Steven, J.C., Herlihy, C.R., Scarpino, S.V. & Bell, D.L. (2011)

Environment-dependent intralocus sexual conflict in a dioecious plant. *The New phytologist*, **192**, 542–

552.

Dillon, M.E., Wang, G., Garrity, P.A. & Huey, R.B. (2009) Review: Thermal preference in *Drosophila*.

Journal of thermal biology, **34**, 109–119.

Dolgin, E.S., Whitlock, M.C. & Agrawal, A.F. (2006) Male *Drosophila melanogaster* have higher mating

success when adapted to their thermal environment. *Journal of evolutionary biology*, **19**, 1894–1900.

Dolgova, O., Rego, C., Calabria, G., Balanyà, J., Pascual, M., Rezende, E.L., *et al.* (2010) Genetic

constraints for thermal coadaptation in *Drosophila subobscura*. *BMC evolutionary biology*, **10**, 363.

Doorn, G.S. van & Kirkpatrick, M. (2007) Turnover of sex chromosomes induced by sexual conflict.

Nature, **449**, 909–912.

Dübendorfer, A., Hediger, M., Burghardt, G. & Bopp, D. (2002) *Musca domestica*, a window on the

evolution of sex-determining mechanisms in insects. *The International journal of developmental biology*,

46, 75–79.

Eanes, W.F. (1999) Analysis of selection on enzyme polymorphisms. *Annu. Rev. Ecol. Syst.*, **30**, 301–326.

Endler, J.A. (1977) *Geographic Variation, Speciation, and Clines*. Princeton University Press.

Etterson, J.R. & Shaw, R.G. (2001) Constraint to adaptive evolution in response to global warming.

Science, **294**, 151–154.

Feldmeyer, B., Kozielska, M., Kuijper, B., Weissing, F.J., Beukeboom, L.W. & Pen, I. (2008) Climatic

variation and the geographical distribution of sex-determining mechanisms in the housefly. *Evolutionary*

ecology research, **10**, 797–809.

Felsenstein, J. (1976) The theoretical population genetics of variable selection and migration. *Annual*

review of genetics, **10**, 253–280.

Franco, M.G., Rubini, P.G. & Vecchi, M. (1982) Sex-determinants and their distribution in various

597 populations of *Musca domestica* L. of Western Europe. *Genetical research*, **40**, 279–293.

598 Geden, C.J., Biale, H., Chiel, E. & Johnson, D.M. (2019) Effect of fluctuating high temperatures on house

599 flies (Diptera: Muscidae) and their principal parasitoids (*Muscidifurax spp.* and *Spalangia spp.*

600 [Hymenoptera: Pteromalidae]) from the United States. *Journal of medical entomology*, **56**, 1650–1660.

601 Gibert, P. & Huey, R.B. (2001) Chill-coma temperature in *Drosophila*: effects of developmental

602 temperature, latitude, and phylogeny. *Physiological and biochemical zoology: PBZ*, **74**, 429–434.

603 Gilbert, A.L. & Miles, D.B. (2017) Natural selection on thermal preference, critical thermal maxima and

604 locomotor performance. *Proceedings. Biological sciences / The Royal Society*, **284**.

605 Good, D.S. (1993) Evolution of behaviours in *Drosophila melanogaster* in high temperatures: Genetic and

606 environmental effects. *J. Ins. Physiol.*, **39**, 537–544.

607 Hamm, R.L., Gao, J.-R., Lin, G.G.-H. & Scott, J.G. (2009) Selective advantage for IIIM males over YM

608 males in cage competition, mating competition, and pupal emergence in *Musca domestica L.* (Diptera:

609 Muscidae). *Environmental entomology*, **38**, 499–504.

610 Hamm, R.L., Meisel, R.P. & Scott, J.G. (2014) The evolving puzzle of autosomal versus Y-linked male

611 determination in *Musca domestica*. *G3* , **5**, 371–384.

612 Hamm, R.L. & Scott, J.G. (2008) Changes in the frequency of YM versus IIIM in the housefly, *Musca*

613 *domestica* L., under field and laboratory conditions. *Genetics research*, **90**, 493–498.

614 Hamm, R.L. & Scott, J.G. (2009) A high frequency of male determining factors in male *Musca domestica*

615 (Diptera: Muscidae) from Ipswich, Australia. *Journal of medical entomology*, **46**, 169–172.

616 Hamm, R.L., Shono, T. & Scott, J.G. (2005) A cline in frequency of autosomal males is not associated

617 with insecticide resistance in house fly (Diptera: Muscidae). *Journal of economic entomology*, **98**, 171–

618 176.

619 Hediger, M., Henggeler, C., Meier, N., Perez, R., Saccone, G. & Bopp, D. (2010) Molecular

620 characterization of the key switch *F* provides a basis for understanding the rapid divergence of the sex-

621 determining pathway in the housefly. *Genetics*, **184**, 155–170.

Hedrick, P.W., Ginevan, M.E. & Ewing, E.P. (1976) Genetic polymorphism in heterogeneous

environments. *Annual review of ecology and systematics*, **7**, 1–32.

Hoffmann, A.A., Anderson, A. & Hallas, R. (2002) Opposing clines for high and low temperature

resistance in *Drosophila melanogaster*. *Ecol. Letters*, **5**, 614–618.

Holleley, C.E., O'Meally, D., Sarre, S.D., Marshall Graves, J.A., Ezaz, T., Matsubara, K., *et al.* (2015)

Sex reversal triggers the rapid transition from genetic to temperature-dependent sex. *Nature*, **523**, 79–82.

Honegger, K. & Bivort, B. de. (2018) Stochasticity, individuality and behavior. *Current biology: CB*, **28**,

R8–R12.

Hothorn, T., Bretz, F. & Westfall, P. (2008) Simultaneous inference in general parametric models. *Biom.*

J, **50**, 346–363.

Huey, R.B., Hertz, P.E. & Sinervo, B. (2003) Behavioral drive versus behavioral inertia in evolution: a

null model approach. *The American naturalist*, **161**, 357–366.

Huey, R.B. & Pascual, M. (2009) Partial thermoregulatory compensation by a rapidly evolving invasive

species along a latitudinal cline. *Ecology*, **90**, 1715–1720.

Inoue, H. & Hiroyoshi, T. (1986) A maternal-effect sex-transformation mutant of the housefly, *Musca*

domestica L. *Genetics*, **112**, 469–482.

Jensen, G. (2018) Behavioral Stochasticity. In *Encyclopedia of Animal Cognition and Behavior* (ed. by

Vonk, J. & Shackelford, T.). Springer International, Cham, Switzerland, pp. 1–5.

Kawecki, T.J. & Ebert, D. (2004) Conceptual issues in local adaptation. *Ecology letters*, **7**, 1225–1241.

Kearney, M., Shine, R. & Porter, W.P. (2009) The potential for behavioral thermoregulation to buffer

"cold-blooded" animals against climate warming. *Proceedings of the National Academy of Sciences of the*

United States of America, **106**, 3835–3840.

Keller, I. & Seehausen, O. (2012) Thermal adaptation and ecological speciation. *Molecular ecology*, **21**,

782–799.

Königer, A., Arif, S. & Grath, S. (2019) Three quantitative trait loci explain more than 60% of variation

for chill coma recovery time in a natural population of *Drosophila ananassae*. *G3*

Genes|Genomes|Genetics.

Königer, A. & Grath, S. (2018) Transcriptome analysis reveals candidate genes for cold tolerance in

Drosophila ananassae. *Genes*, **9**.

Kotiaho, J.S., Simmons, L.W. & Tomkins, J.L. (2001) Towards a resolution of the lek paradox. *Nature*,

410, 684–686.

Kozielska, M., Feldmeyer, B., Pen, I., Weissing, F.J. & Beukeboom, L.W. (2008) Are autosomal sex-

determining factors of the housefly (*Musca domestica*) spreading north? *Genetics research*, **90**, 157–165.

Kozielska, M., Weissing, F.J., Beukeboom, L.W. & Pen, I. (2010) Segregation distortion and the evolution

of sex-determining mechanisms. *Heredity*, **104**, 100–112.

Krstevska, B. & Hoffmann, A.A. (1994) The effects of acclimation and rearing conditions on the response

of tropical and temperate populations of *Drosophila melanogaster* and *D. simulans* to a temperature

gradient (Diptera: Drosophilidae). *J. Ins. Behav.*, **7**, 279–288.

Lasne, C., Hangartner, S.B., Connallon, T. & Sgrò, C.M. (2018) Cross-sex genetic correlations and the

evolution of sex-specific local adaptation: Insights from classical trait clines in *Drosophila melanogaster*.

Evolution; international journal of organic evolution, **72**, 1317–1327.

Leiva, F.P., Calosi, P. & Verberk, W.C.E.P. (2019) Scaling of thermal tolerance with body mass and

genome size in ectotherms: a comparison between water- and air-breathers. *Philosophical transactions of*

the Royal Society of London. Series B, Biological sciences, **374**, 20190035.

Levene, H. (1953) Genetic equilibrium when more than one ecological niche is available. *The American*

naturalist, **87**, 331–333.

Lynch, K.E., White, T.E. & Kemp, D.J. (2018) The effect of captive breeding upon adult thermal

preference in the Queensland fruit fly (*Bactrocera tryoni*). *Journal of thermal biology*, **78**, 290–297.

MacLean, H.J., Overgaard, J., Kristensen, T.N., Lyster, C., Hessner, L., Olsvig, E., *et al.* (2019)

Temperature preference across life stages and acclimation temperatures investigated in four species of

Drosophila. *Journal of thermal biology*, **86**, 102428.

McDonald, I. c., Evenson, P., Nickel, C.A. & Johnson, O.A. (1978) House fly genetics: isolation of a

female determining factor on chromosome 4. *Annals of the Entomological Society of America*, **71**, 692–

694.

Meisel, R.P. (2018) Sexual conflict and the maintenance of genetic variation in natural populations.

Molecular ecology, **27**, 3569–3571.

Meisel, R.P., Davey, T., Son, J.H., Gerry, A.C., Shono, T. & Scott, J.G. (2016) Is multifactorial sex

determination in the house fly, *Musca domestica* (L.), stable over time? *The Journal of heredity*, **107**, 615–

625.

Meisel, R.P., Gonzales, C.A. & Luu, H. (2017) The house fly Y chromosome is young and minimally

differentiated from its ancient X chromosome partner. *Genome research*, **27**, 1417–1426.

Meisel, R.P., Scott, J.G. & Clark, A.G. (2015) Transcriptome differences between alternative sex

determining genotypes in the house fly, *Musca domestica*. *Gen. Biol. and Evol.*, **7**, 2051–2061.

Moore, E.C. & Roberts, R.B. (2013) Polygenic sex determination. *Current biology: CB*, **23**, R510–2.

Partridge, L., Barrie, B., Fowler, K. & French, V. (1994) Evolution and development of body size and cell

size in *Drosophila melanogaster* in response to temperature. *Evolution; international journal of organic

evolution*, **48**, 1269–1276.

Pen, I., Uller, T., Feldmeyer, B., Harts, A., While, G.M. & Wapstra, E. (2010) Climate-driven population

divergence in sex-determining systems. *Nature*, **468**, 436–438.

Qu, Y., Li, H., Gao, J., Xu, X. & Ji, X. (2011) Thermal preference, thermal tolerance and the thermal

dependence of digestive performance in two *Phrynocephalus* lizards (Agamidae), with a review of species

studied. *Curr. Zool.*, **57**, 684–700.

Rego, C., Balanyà, J., Fragata, I., Matos, M., Rezende, E.L. & Santos, M. (2010) Clinal patterns of

chromosomal inversion polymorphisms in *Drosophila subobscura* are partly associated with thermal

preferences and heat stress resistance. *Evolution; international journal of organic evolution*, **64**, 385–397.

Rice, W.R. (1986) On the instability of polygenic sex determination: the effect of sex-specific selection.

Evolution; international journal of organic evolution, **40**, 633–639.

Richter-Boix, A., Teplitsky, C., Rogell, B. & Laurila, A. (2010) Local selection modifies phenotypic

divergence among *Rana temporaria* populations in the presence of gene flow. *Molecular ecology*, **19**,

716–731.

Robinson, S.J.W. & Partridge, L. (2001) Temperature and clinal variation in larval growth efficiency in

Drosophila melanogaster. *Journal of evolutionary biology*, **14**, 14–21.

Rostant, W.G., Kay, C., Wedell, N. & Hosken, D.J. (2015) Sexual conflict maintains variation at an

insecticide resistance locus. *BMC biology*, **13**, 34.

Scott, J.G., Sridhar, P. & Liu, N. (1996) Adult specific expression and induction of cytochrome P450lpr in

house flies. *Archives of insect biochemistry and physiology*, **31**, 313–323.

Scrucca, L., Fop, M., Murphy, T.B. & Raftery, A.E. (2016) mclust 5: clustering, classification and density

estimation using Gaussian finite mixture models. *The R journal*, **8**, 289–317.

Sharma, A., Heinze, S.D., Wu, Y., Kohlbrenner, T., Morilla, I., Brunner, C., *et al.* (2017) Male sex in

houseflies is determined by *Mdmd*, a paralog of the generic splice factor gene. *Science*, **356**, 642–645.

Slatkin, M. (1973) Gene flow and selection in a cline. *Genetics*, **75**, 733–756.

Slatyer, R.A., Hirst, M. & Sexton, J.P. (2013) Niche breadth predicts geographical range size: a general

ecological pattern. *Ecol. Letters*, **16**, 1104–1114.

Son, J.H., Kohlbrenner, T., Heinze, S., Beukeboom, L.W., Bopp, D. & Meisel, R.P. (2019) Minimal

effects of proto-Y chromosomes on house fly gene expression in spite of evidence that selection maintains

stable polygenic sex determination. *Genetics*, **213**, 313–327.

Son, J.H. & Meisel, R.P. (2020) Gene-level, but not chromosome-wide, divergence between a very young

house fly proto-Y chromosome and its homologous proto-X chromosome. *Molecular biology and*

evolution.

Svetec, N., Werzner, A., Wilches, R., Pavlidis, P., Alvarez-Castro, J.M., Broman, K.W., *et al.* (2011)

Identification of X-linked quantitative trait loci affecting cold tolerance in *Drosophila melanogaster* and

fine mapping by selective sweep analysis. *Molecular ecology*, **20**, 530–544.

Tomita, T. & Wada, Y. (1989) Multifactorial sex determination in natural populations of the housefly

(*Musca domestica*) in Japan. *The Japan. J. of Gen.*, **64**, 373–382.

Tüzün, N., Op de Beeck, L., Brans, K.I., Janssens, L. & Stoks, R. (2017) Microgeographic differentiation

in thermal performance curves between rural and urban populations of an aquatic insect. *Evolutionary*

applications, **10**, 1067–1075.

Wang, Y., Yang, L., Zhang, Y., Tao, L. & Wang, J. (2018) Development of *Musca domestica* at constant

temperatures and the first case report of its application for estimating the minimum postmortem interval.

Forensic science international, **285**, 172–180.

Yang, J., Sun, Y.-Y., An, H. & Ji, X. (2008) Northern grass lizards (*Takydromus septentrionalis*) from

different populations do not differ in thermal preference and thermal tolerance when acclimated under

identical thermal conditions. *J. Comp. Physiol. B*, **178**, 343–349.