1   # Genetic analysis of the contribution of the proprotein convertases

2   to the *in vivo* regulation of FGF23

4   Running title: Proprotein convertases in the regulation of FGF23

6   Omar Al Rifai[1,2], Delia Susan-Resiga[3], Rachid Essalmani[3], John W.M.

7   Creemers[4], Nabil G. Seidah[3] and Mathieu Ferron[1,2,5,6]#

8
[1] Unité de recherche en physiologie moléculaire, Institut de Recherches Cliniques de Montréal, Montréal,
Québec H2W 1R7, Canada
[2] Programme de biologie moléculaire, Université de Montréal, Québec H3T 3J7, Canada
[3] Unité de recherche en biochimie neuroendocrinienne, Institut de Recherches Cliniques de Montréal,
Montréal, Québec H2W 1R7, Canada
[4] Department of Human Genetics, KU Leuven, Herestraat 49, B-3000 Leuven, Belgium
[5] Département de Médecine, Université de Montréal, Québec H3T 3J7, Canada
[6] Division of Experimental Medicine, McGill University, Montréal, Québec H3A 1A3, Canada
17

Omar.alrifai@ircm.qc.ca; Delia.Susan-Resiga@ircm.qc.ca; Rachid.Essalmani@ircm.qc.ca;

john.creemers@kuleuven.be; Nabil.Seidah@ircm.qc.ca; mathieu.ferron@ircm.qc.ca

Correspondence to:

Mathieu Ferron, PhD
Institut de Recherches Cliniques de Montréal
110 Ave. des Pins O.
Montréal, QC,
H2W 1R7, Canada
Phone: 514-987-5754
Fax: 514-987-5649
Email: mathieu.ferron@ircm.qc.ca
30

Author contributions: MF, DSR and OAR conceived and planned the experiments. OAR

and DSR carried out the experiments. OAR and MF analyzed the data. RE, JWMC and

NGS provided critical mouse lines. OAR and MF wrote the manuscript with input from all

authors.

Keywords

FGF23, endoproteolysis, proprotein convertases, Furin, PC5, PACE4, phosphate

ABSTRACT

Fibroblast growth factor 23 (FGF23) is a hormone secreted from terminally differentiated osteoblasts and by osteocytes that inhibits phosphate reabsorption by kidney proximal tubules. The full-length (i.e., intact) protein mediates FGF23 endocrine functions, while endoproteolytic cleavage at a consensus cleavage sequence for the proprotein convertases (PCs) inactivates it. Two PCs, furin and PC5, were shown to cleave FGF23 *in vitro* at RHTR$_{179}\downarrow$, but those involved in FGF23 processing *in vivo* are not defined. Here, we assessed FGF23 processing and phosphate metabolism in mice lacking either or both furin and PC5 in the osteoblastic lineage and/or in hematopoietic cells, and in mice lacking the paired basic amino acid-cleaving enzyme 4 (PACE4) in all tissues. Our results showed that furin inactivation in osteoblasts and osteocytes results in a 25% increase in circulating intact FGF23, without any significant impact on serum phosphate levels, whether mice are maintained on a normal or a low phosphate diet. Under conditions of iron deficiency, FGF23 processing is impaired in mice lacking furin in osteoblasts and osteocytes, while it is normally processed in control mice. In contrast, FGF23 is normally processed following erythropoietin or IL-1β injections in mice lacking furin or both furin and PC5, and in PACE4-deficient mice. Altogether, these studies suggest that furin is only partially responsible for FGF23 cleavage under certain conditions *in vivo*. The processing of FGF23 may therefore involve the redundant action of multiple PCs or of other peptidases in osteoblasts, osteocytes and hematopoietic cells.

INTRODUCTION

Fibroblast growth factor 23 (FGF23) is a bone-derived hormone (i.e., osteokine) secreted by terminally differentiated cells of the osteoblast lineage, including osteoblasts and osteocytes. It plays a fundamental role in phosphate and vitamin D metabolism in particular through a direct action on the kidney [1,2]. In this tissue, its signaling is mediated through interaction with the FGF receptor 1 (FGFR1) and its co-receptor α-klotho [3]. FGFR1/α-klotho signaling in the kidney proximal tubule suppresses the transcription of *Slc34a1* and *Slc34a3*, the genes encoding for the sodium phosphate co-transporters NaPi2A and NaPi2C, resulting in increased urinary phosphate excretion and decreased serum phosphate levels. In addition, FGF23 reduces the 1,25-dihydroxyvitamin D_3 (calcitriol) level by decreasing expression of *Cyp27b1* which encodes the 25-hydroxyvitamin D-1 alpha hydroxylase and increasing expression of *Cyp24a1* which encodes the 1,25-dihydroxyvitamin D_3 24-hydroxylase [4-7]. These endocrine functions are mediated through full-length "intact" FGF23 (amino acids [aa] 25-251). Endoproteolytic cleavage of FGF23 at the basic motif $\underline{R}_{176}HTR_{179}\downarrow SA$, a consensus P1 and P4 arginine (Arg) cleavage site for proprotein convertases (PCs), inactivates FGF23 by releasing N-terminal (aa 25-179) and C-terminal (aa 180-251) fragments.

The circulating level of intact FGF23 is tightly regulated in part through various post-translational modifications of amino acids surrounding the PC cleavage site. The *O*-glycosylation of threonine 178 (T_{178}) by the *N*-acetylgalactosaminyltransferase 3 (GalNAc-T3) inhibits FGF23 processing [8-10]. FGF23 *O*-glycosylation is in turn reduced when the serine 180 (S_{180}) is phosphorylated by Fam20c, a kinase residing in the Golgi

[11]. Impaired FGF23 glycosylation caused by a defective GalNAc-T3 or mutation in FGF23

glycosylation sites cause an increase in FGF23 processing and subsequent decrease in

circulating intact FGF23 levels. These molecular defects lead to familial tumoral calcinosis

(FTC), a human disease characterized by hyperphosphatemia and heterotopic

calcification [8-10]. On the other hand, mutations in the FGF23 cleavage site and in the X-

linked phosphate-regulating neutral endopeptidase (PHEX) respectively cause

autosomal dominant hypophosphatemic rickets (ADHR) and X-linked hypophosphatemic

rickets (XLH). Patients with ADHR and XLH develop hypophosphatemia and

osteomalacia due to the increased level of active intact FGF23 [12,13]. Iron deficiency,

erythropoietin treatment or interleukin-1 beta (IL-1β) were shown to stimulate FGF23

production by osteocytes and bone marrow hematopoietic cells [14-19]. However, this

increase in FGF23 production only results in the accumulation of intact FGF23 levels and

hypophosphatemia in ADHR patients or in mice with an ADHR-like mutation in *Fgf23*. In

wildtype animals or healthy subjects, the same stimuli result in increased circulating levels

of cleaved FGF23, revealing the importance of endoproteolysis by PC for the regulation

of this hormone.

The proprotein convertases furin and PC5 were shown to cleave FGF23 *in vitro* [20], while

inactivation of *FURIN* in the human osteosarcomas U2-OS cell line impaired the cleavage

of ectopically expressed human FGF23 [11]. Nevertheless, despite the implication of FGF23

cleavage in several human diseases, the identity of the enzyme(s) responsible for FGF23

endoproteolysis *in vivo* remains undetermined. In the present study we assessed the

requirement of furin and PC5 for FGF23 processing and phosphate metabolism *in vivo*.

We showed that furin inactivation in the osteoblastic lineage results in a modest increase

in intact FGF23 under normal or low phosphate diet, while it impairs FGF23 cleavage in

conditions of iron deficiency. We also show that furin, PC5 and PACE4, another PC

expressed in osteoblasts, which can cleave FGF23 *in vitro*, are dispensable for the

cleavage and inactivation of FGF23 *in vivo* following treatment with erythropoietin or IL-

1β. Together, our data suggest that furin is involved in FGF23 cleavage *in vivo*, but that

multiple other PC(s) are also implicated in this process.

RESULTS

Furin inactivation in osteoblasts and osteocytes partially impairs FGF23 cleavage, but decreases phosphate excretion

To investigate the role of furin in the regulation of FGF23 processing *in vivo*, we generated mice in which the *Furin* gene was specifically inactivated in osteoblasts and osteocytes (i.e., *Furin*osb*-/-*) by breeding *Furin*fl/fl* mice with *hOCN-Cre* transgenic animals which express the Cre recombinase under the control of human osteocalcin promoter [21]. We previously reported that the *Furin* gene was efficiently and specifically inactivated in osteoblasts and bones in these mice [22]. When fed a normal chow diet containing ~0.7% inorganic phosphate (Pi), 4-month-old *Furin*osb*-/-* female mice displayed a significant 25% rise in circulating intact FGF23 (Figure 1A). This moderate increase did not significantly alter the expression of *Slc34a1*, *Slc34a3*, *Cyp27b1* or *Cyp24a1* (Figure 1B). Accordingly, these mice maintained normal serum phosphate levels and bone histology did not reveal presence of osteomalacia (Figure 1C and D).

Serum phosphate level is a potent regulator of FGF23 and previous studies have shown that in mice fed a low-phosphate synthetic diet the serum level of intact FGF23 is reduced, while the processed fraction is increased, presumably due to the action of one or several PCs [23-25]. We therefore next assessed if severe hypophosphatemia could be induced in *Furin*osb*-/-* mice under low phosphate dietary intake. Feeding control and *Furin*osb*-/-* mice normal phosphate diet (i.e., 0.6% Pi) or low-phosphate diet (0.02% Pi) for one week did not affect body weight or kidney weight (Figure 1E and F). As expected, serum

Figure 1: Increased FGF23 level and reduced phosphate excretion in *Furin^osb^-/-* mice. FGF23 and phosphate levels in in 4-month-old *Furin^flox/flox^* (n=6) and *Furin^osb^-/-* (n=6) mice fed on normal chow diet **(A-D)**. Intact FGF23 levels in plasma **(A)**. FGF23 target genes expression in kidney **(B)**. Serum phosphate level **(C)**. Von Kossa / van Gieson staining on vertebrae sections **(D)**. FGF23 and phosphate parameters in 6-months-old *Furin^flox/flox^* (n=6) and *Furin^osb^-/-* (n=6-7) mice fed a normal phosphate diet (0.6%) or a low phosphate diet (0.02%) for one week **(E-N)**. Body weight **(E)**. Kidney weight normalize to body weight **(F)**. Serum phosphate levels **(G)**. Plasma level of C-terminal FGF23 **(H)**, intact FGF23 **(I)**, percentage (%) of intact over C-terminal FGF23 **(J)** and cleaved FGF23 levels **(K)** calculated as (C-terminal FGF23) - (intact FGF23). Expression level in kidney of the sodium phosphate cotransporter *Slc34a1* **(L)** and *Slc34a3* **(M)**. **(N)** Urine phosphate measurements normalized to creatinine. Results represent the mean ± SEM. *$P < 0.05$, **$P < 0.01$, and ***$P < 0.001$, by 2-way ANOVA with Bonferroni's multiple comparisons test or by Student's t-test (A, C, G, H, K, and M) #: $P<0.05$; # #: $P<0.01$, # # #: $P<0.001$ &: $0.05<P<0.1$.

phosphate level decreased significantly in both control and *Furin^osb^-/-* mice fed the low

phosphate diet (Figure 1G). Under the same conditions, *Furin^osb^-/-* mice tend to have a

lower serum phosphate level compared to control mice, although this did not reach

statistical significance. *Furin^osb^-/-* mice on normal diet had >25% increase in intact and C-

terminal (total) FGF23, while maintaining normal intact/C-terminal FGF23 ratio compared

to control littermates (Figure 1H-J). Under low phosphate diet circulating intact FGF23

was reduced in control mice but remains significantly higher in *Furin^osb^-/-* mice (Figure 1I).

Yet, the ratio of intact over C-terminal FGF23 was decreased in both control and *Furin^osb^-*

/- mice under low phosphate diet (Figure 1J). Moreover, the amount of cleaved FGF23

(i.e., C-terminal FGF23 minus intact FGF23) was increased regardless of the genotype

on low phosphate diet, suggesting efficient cleavage of FGF23 even in absence of furin

(Figure 1K). Surprisingly, when fed these synthetic diets, *Furin^osb^-/-* mice displayed an

increased expression in the kidney of the sodium phosphate transporter genes *Slc34a1*

and *Slc34a3* compared to control group, independently of the level of phosphate in the

diet (Figure 1L and M). In addition, *Furin^osb^-/-* mice on normal phosphate diet showed a

decrease in urinary phosphate compared to control littermate (Figure 1N). Challenging

the mice for one week on the low phosphate diet reduced urinary phosphate in control

mice as expected, but not in *Furin^osb^-/-* mice, as these mice maintained the low urinary

phosphate observed on normal phosphate diet (Figure 1N). Together these results show

that under physiological conditions or in conditions of decreased phosphate intake,

inactivation of *Furin* in the osteoblast lineage modestly impacted FGF23 processing. They

also suggest that furin may influence phosphate excretion through FGF23-independent

mechanisms.

Impact of iron deficiency on FGF23 processing and phosphate metabolism in

absence of furin in osteoblasts and osteocytes

It was previously reported that mice carrying an ADHR mutation (R176Q, i.e., loss of P4

Arg in the motif $\underline{\mathbf{R}}_{176}HT\underline{\mathbf{R}}_{179}\downarrow SA$) in the *Fgf23* gene did not develop hypophosphatemia

unless they were fed a low-iron diet [15]. Therefore, iron-deficiency was induced in control

and *Furinᵒˢᵇ-/-* mice by feeding them a low-iron diet in combination with repeated tail

bleedings (i.e., ~70 µl every other week) for 14 weeks, before assessing the impact on

FGF23 and phosphate metabolism. Mice of both genotypes maintained normal growth

curves in these conditions as compared to mice fed the control chow diet (Figure 2A).

Figure 2: Impaired FGF23 processing in *Furinᵒˢᵇ-/-* mice following iron restriction. *Furin^{flox/flox}* (n=3-5)
and *Furinᵒˢᵇ-/-* (n=7-8) 21-day-old mice were fed normal chow diet (ND) or a low iron diet (LI) for 14 weeks.
(A) Body weight growth curve presented as percentage of initial body weight. **(B-C)** Expression of iron-
regulated genes in liver. Hepcidin (*Hamp*) expression **(B)**. Transferrin receptor (*Tfrc*) expression **(C). (D)**
Fgf23 expression in long bone. **(E-H)** Plasma level of C-terminal FGF23 **(E)**, intact FGF23 **(F)**, percentage
(%) of intact over C-terminal FGF23 **(G)** and cleaved FGF23 levels **(H)**, calculated as (C-terminal FGF23)
- (intact FGF23). **(I)** Serum phosphate levels. Results represent the mean ± SEM. *$P < 0.05$, **$P < 0.01$,
and ***$P < 0.001$, by 2-way ANOVA with Bonferroni's multiple comparisons test or by Student's t-test (D
and E) #: $P<0.05$; # #: $P<0.01$, # # #: $P<0.001$ &: $0.05<P<0.1$.
192
However, challenged mice developed iron deficiency as reflected by the decrease in

hepcidin (*Hamp*) expression and the increase in transferrin receptor (*Tfrc*) expression in

liver (Figure 2B-C). As expected from a previous study [15], iron deficient control and

Furin^osb^-/- mice displayed an increase in *Fgf23* gene expression in bone (Figure 2D).

However, iron-deficiency resulted in increased C-terminal FGF23 only in control mice,

although both genotypes showed an increase in intact FGF23 (Figure 2E-F). FGF23

processing appears to be totally impaired in *Furin^osb^-/-* mice under conditions of iron

deficiency as these mice maintained a 1:1 ratio (i.e., 100%) of intact over C-terminal

FGF23 and very low level of cleaved FGF23 (Figure 2G-H). Yet, despite this near

complete inhibition in FGF23 processing, *Furin^osb^-/-* mice displayed a paradoxical

increase in serum phosphate level compared to the control littermates under iron

deficiency (Figure 2I). Altogether these results suggest that furin is responsible for FGF23

processing under iron deficiency. However, under these conditions the absence of furin

in osteoblasts and osteocytes also results in a compensatory increase in serum

phosphate levels in an FGF23-independent manner.

Normal FGF23 processing in absence of furin in osteoblasts and osteocytes

following rhEPO and IL-1β injections

Anemia and inflammation may contribute to the onset of ADHR [16-19,26]. Erythropoietin and

interleukin 1 beta (IL-1β), which are increased in condition of anemia and inflammation

respectively, are two potent inducers of *Fgf23* at the transcriptional level [16-19,26,27]. In wild

type mice, most (i.e., >80%) of the circulating FGF23 is in its cleaved form following

recombinant human erythropoietin (rhEPO) or IL-1β injections, suggesting the action of

one or multiple PC in these particular conditions [16,17,19]. We therefore next tested the

impact of furin deficiency in osteoblasts and osteocytes on FGF23 processing 6 hours

after a single injection of rhEPO or IL-1β.

Figure 3: FGF23 is normally processed following erythropoietin and IL-1β injection in *Furin^osb^-/-*
mice. Genes expression in long bone **(A, B and E)** and bone marrow **(C, D and F)** of 2-month-old C57BL/6J
mice 6 hours following the injection of vehicle (n=4), 300 U/kg (n=4) or 3000 U/kg (n=4) of recombinant
human erythropoietin (rhEPO). Erythropoietin receptor (*Epor*) **(A and C)**, erythroferrone (*Erfe*) **(B and D)**
and FGF23 (*Fgf23*) **(C and F)**. **(G-J)** Plasma FGF23 measurements in 4-month-old *Furin^flox/flox^* (n=3) and
Furin^osb^-/- (n=3) mice 6 hours following the injection of vehicle, 300 U/kg or 3000 U/kg of rhEPO. C-terminal
FGF23 **(G)**, intact FGF23 **(H),** percentage (%) of intact over C-terminal FGF23 **(I)** and cleaved FGF23 levels
(J), calculated as (C-terminal FGF23) - (intact FGF23). **(K-N)** Plasma FGF23 measurements in 4-month-
old *Furin^flox/flox^* (n=4) and *Furin^osb^-/-* (n=4) mice before (T0) and 6 hours following the injection of 50 ng/g IL-
1β (T=6hrs). C-terminal FGF23 **(K)**, intact FGF23 **(L),** percentage (%) of intact over C-terminal FGF23 **(M)**
and cleaved FGF23 levels **(N)**. Results represent the mean ± SEM. *$P < 0.05$, **$P < 0.01$, and ***$P < 0.001$,
by 1-way ANOVA **(A-F)** or 2-way ANOVA **(G-N)** with Bonferroni's multiple comparisons test, or by Student's
t-test **(A and E)** #: $P<0.05$; # #: $P<0.01$, # # #: $P<0.001$ &: $0.05<P<0.1$.

Because of variation in the doses of rhEPO used in the literature to induce FGF23 in

rodents [17-19,26,27], we first confirmed that two relatively low doses of rhEPO (i.e., 300U/kg

and 3000U/kg) were sufficient to induce in bones and bone marrow of wild type C57BL/6J

mice the expression of two known EPO target genes, erythropoietin receptor (*Epor*) and

erythroferrone (*Erfe*) (Figure 3A-D). These two doses also robustly induced the

expression of *Fgf23* in both tissues in the same animals (Figure 3E-F). We next assessed

the effect of the same treatment in control and *Furin^{osb}-/-* mice. Circulating level of total

(C-terminal) FGF23 was increased in a dose dependent manner by rhEPO, while intact

FGF23 was modestly changed regardless of the genotype (Figure 3G-H). The ratio of

intact over C-terminal FGF23 level was robustly reduced by the same treatment (Figure

3I) and cleaved FGF23 was increased more than 10-fold by rhEPO in both control and

Furin^{osb}-/- mice (Figure 3J), suggesting that in this context, FGF23 is normally processed

even in absence of furin in osteoblasts and osteocytes.

In separate experiments, control and *Furin^{osb}-/-* mice were administered a single dose of

IL-1β (50ng/g), a treatment which was previously shown to induce *Fgf23* expression in

bone and to raise total FGF23 circulating level within a few hours [16]. Six hours after the

injection of IL-1β, the level of total (C-terminal) FGF23 increased by more than 30-fold,

while intact FGF23 increased by less than 10-fold, in both control and *Furin^{osb}-/-* mice

(Figure 3K-L). Following the IL-1β treatment and regardless of the mouse genotype, the

ratio of intact over C-terminal FGF23 level decreased from 60% to less than 10% (Figure

3M) and the plasma concentration of cleaved FGF23 increased more than 10 times

(Figure 3N). Together, these data show that FGF23 is efficiently processed in absence of

furin in osteoblasts and osteocytes following rhEPO or IL-1β injections, likely to the

contribution of other processing enzymes.

FGF23 is processed in mice lacking furin in osteoblasts, osteocytes and

hematopoietic cells following rhEPO injections

The data presented above suggest that under normal conditions furin is partially

responsible for FGF23 cleavage, but that in the context of iron deficiency, osteocytes-

derived furin may be a major contributor to the processing of FGF23. In contrast, FGF23

was normally processed following rhEPO or IL-1β injection in *Furin*^{osb}*-/-* mice. Osteoblasts

and osteocytes are the main source of circulating FGF23 in normal conditions [26,28], but

bone marrow hematopoietic cells contribute to approximately 40% of circulating FGF23

following rhEPO treatment [26]. Therefore, in *Furin*^{osb}*-/-* mice, bone marrow hematopoietic

cells could be a significant source of cleaved FGF23 following rhEPO injections, thereby

masking the impact of furin inactivation in osteoblasts and osteocytes. To test this

possibility, we generated mice lacking furin in all hematopoietic cells by breeding

Furin$^{flox/flox}$ mice with *Vav1-Cre* to generate *Furin*$^{flox/flox}$*; Vav1-Cre* mice (*Furin*^{BM}*-/-*). *Vav1-*

Cre transgenic animals expresses the Cre recombinase under the control of the *Vav1*

promoter, allowing the efficient deletion of furin in fetal and adult hematopoietic stem cells

(HSC) [29] and hence in all hematopoietic cells. To address any potential redundancy

between osteocytes and hematopoietic cells in the cleavage of FGF23, we also generated

mice lacking *Furin* in both cell types, i.e., *Furin*$^{flox/flox}$*;* hOCN-Cre*;* Vav1-Cre mice

(*Furin*$^{osb;BM}$*-/-*). Furin deficiency in T cells impairs the function of regulatory and effector T

cells, causing at around 6 month of age severe inflammatory bowel disease, weight loss,

ruffled hair and hunched appearance [30]. Confirming efficient inactivation of *Furin* in

hematopoietic cells, these defects were also observed in our *Furin^BM-/-* mice between 4

and 6 months of age (data not shown). Hence, in the following experiments all mice were

8 weeks of age, a time point at which the *Furin^BM-/-* mice are still healthy.

Figure 4: *Furin* inactivation in osteoblasts, osteocytes and hematopoietic cells did not impair FGF23
processing following erythropoietin injection. Plasma FGF23 measurements in 8-week-old *Furin^flox/flox*
(n=7), *Furin^osb-/-* (n=5), *Furin^osb-/-* (n=5) and *Furin^osb;BM-/-* (n=6) mice 6 hours following injection of vehicle
or 3000 U/kg of recombinant human erythropoietin (rhEPO): C-terminal FGF23 **(A)**, intact FGF23 **(B)**,
percentage (%) of intact over C-terminal FGF23 **(C)** and cleaved FGF23 levels **(D)**, calculated as (C-
terminal FGF23) - (intact FGF23). Results represent the mean ± SEM. *$P < 0.05$ and ***$P < 0.001$, by 2-
way ANOVA with Bonferroni's multiple comparisons test.
289

A single injection of rhEPO (3000U/kg) increased total (C-terminal) FGF23 by 20-30 fold

and intact FGF23 by about 2 fold in control (*Furin^flox/flox*), *Furin^osb-/-*, *Furin^BM-/-* and

Furin^osb;BM-/- mice (Figure 4A-B). The same treatment also reduced the ratio of intact over

C-terminal FGF23 from 40% to <10%, while cleaved FGF23 level increased to the same

extent (>10 fold) in all genotypes (Figure 4C-D). Together these results show that FGF23

is still processed in absence of furin in osteoblasts/osteocytes and bone marrow

hematopoietic cells following rhEPO injections, suggesting that additional PCs may

redundantly cleave FGF23 in these cells *in vivo*.

298
Figure 5: Multiple proprotein
convertases can process FGF23
in mammalian cells. (A) Western
blot analysis of the secretion media
(SM) from CHO-K1 cells transfected
with V5 tagged human FGF23
(hFGF23-V5) and cultured in the
absence (-) or presence (+) of PC
inhibitors, decanoyl-RVKR-CMK
(RVKR) (50 μM) or hexa-D-arginine
(D6R) (20 μM). **(B)** Western blot
analysis of the secretion media (SM)
from CHO furin-deficient FD11 cells
co-transfected with V5 tagged
human FGF23 (hFGF23-V5) and
proprotein convertase Furin, PC5A,
PC5B, PACE4 or PC7. Intact and
processed C-terminal (CT) FGF23
forms are detected with V5 antibody.

Several PCs can process FGF23 in cell culture

Our data suggests that in some conditions (e.g., low phosphate diet, rhEPO injection or

IL-1β injection), more than one PC may be involved in FGF23 cleavage. In order to identify

which additional PC(s) might be involved in FGF23 processing, we tested FGF23

processing in cell lines expressing or not furin. In Chinese hamster ovary (CHO-K1) cells,

which endogenously express furin [31], ectopically transfected human FGF23 is secreted

and partially processed (Figure 5A). In the same cells, cleavage of FGF23 was inhibited

by decanoyl-RVKR-CMK (RVKR) a pan-PC cell-permeable inhibitor, but not by hexa-D-

arginine (D6R), a cell-surface PC inhibitor (Figure 5A), suggesting that cleavage of

FGF23 occurs before it is secreted in an intracellular secretory compartment, likely the

trans-Golgi network [32]. In addition, FGF23 was not cleaved in CHO-FD11 cells, which do

not endogenously express furin [31], and co-transfection with a furin expressing vector was

sufficient to restore efficient cleavage (Figure 5B, first two lanes). We next tested the

capacity of other PCs, i.e., PC5A, PC5B, PACE4 and PC7 [33], to process FGF23 in the

same cells. Interestingly in this assay, PC5A was almost as efficient as furin in processing

FGF23. Moreover, PC5B, PACE4 and PCSK7 were also found to process FGF23

although to a lower extent than furin and PC5A (Figure 5B). Altogether, these results

indicate that in addition to furin, several other PCs have the capacity to process FGF23

in a mammalian cell line.

Genetic ablation of PC5 and furin in osteoblasts and osteocytes does not impair

FGF23 processing

Our data shows that PC5A and PC5B, which are both encoded by *Pcsk5* [33], can cleave

FGF23 in cell culture, and PC5A is expressed in differentiated osteoblasts and in

osteocytes [20,22]. We therefore next generated mice in which *Pcsk5* was inactivated

specifically in osteoblasts and osteocytes by breeding *Pcsk5*[fl/fl] mice [34] with hOCN-Cre

transgenic animals. When fed a normal chow diet, control (*Pcsk5*[fl/fl]) and *Pcsk5*[osb]-/- mice

displayed similar circulating level of intact FGF23 and serum phosphate (Figure 6A-B).

Von Kossa/van Gieson staining on non-decalcified bone sections did not reveal any sign

of osteomalacia in these mice (Figure 6C). These analyses suggest that PC5 in the

osteoblast lineage is dispensable for the regulation of FGF23 and phosphate levels *in*

vivo.

To test if PC5 and furin redundantly cleave FGF23 *in vivo*, we next generated mice lacking

both furin and PC5 specifically in osteoblasts and osteocytes (i.e., *Furin^flox/flox; Pcsk5^flox/flox;*

hOCN-Cre or *Furin;Pcsk5^osb-/-* mice). Similar to the single inactivation of furin in the same

cells, osteoblast and osteocytes-specific inactivation of furin and PC5 resulted in a

modest increase in circulating intact FGF23 when mice were fed normal or low phosphate

diet (Figure 6D). A single injection of rhEPO (300U/kg) increased total (C-terminal) FGF23

by almost 10-fold and intact FGF23 by 25% in both control and *Furin;Pcsk5^osb-/-* mice

(Figure 6E-F). Following the same treatment, intact over C-terminal ratio decreased by

approximately 80%, while cleaved FGF23 increased more than 10-fold, regardless of the

Figure 6: Genetic ablation of PC5 alone or in combination with Furin does not impair FGF23
processing *in vivo*. Plasma intact FGF23 **(A)** and serum phosphate levels **(B)** in 4-month-old *Pcsk5^flox/flox*
(n=6) and *Pcsk5^osb-/-* (n=7) mice fed a normal chow diet. **(C)** Von Kossa / van Gieson staining of vertebrae
sections of *Pcsk5^flox/flox* and *Pcsk5^osb-/-*. **(D)** Plasma intact FGF23 in 6-month-old *Furin;Pcsk5^flox/flox* (n=7-8)
and *Furin;Pcsk5^osb-/-* (n=8) fed a normal phosphate diet (0.6%) or a low phosphate diet (0.02%) for one
week. **(E and H)** Plasma FGF23 measurements in 4-month-old *Furin;Pcsk5^flox/flox* (n=4) and *Furin;Pcsk5^osb-*
/- (n=4) 6 hours following an injection of vehicle and 300 U/kg of recombinant human erythropoietin (rhEPO):
C-terminal FGF23 **(E)**, intact FGF23 **(F)**, percentage (%) of intact over C-terminal FGF23 **(G)** and cleaved
FGF23 levels **(H)**, calculated as (C-terminal FGF23) - (intact FGF23). Results represent the mean ± SEM.
*$P < 0.05$ and ***$P < 0.001$, by 2-way ANOVA with Bonferroni's multiple comparisons test, or by Student's
t-test **(A and B)**.

mice genotype (Figure 6G-H). Altogether, these results show that FGF23 is still efficiently

processed *in vivo* even when both furin and PC5 were inactivated in osteoblasts and

osteocytes.

Figure 7: PACE4 inactivation in mice does not impair FGF23 processing. Plasma FGF23 measurements in 4-month-old *Pcsk6 +/+* (n=4) and *Pcsk6 -/-* (n=4) male mice before (T0) and 6 hours following the injection of 50 ng/g of IL-1β: C-terminal FGF23 **(A)**, intact FGF23 **(B)**, percentage (%) of intact over C-terminal FGF23 **(C)**, and cleaved FGF23 levels **(D)**, calculated as (C-terminal FGF23) - (intact FGF23). Results represent the mean ± SEM. *$P < 0.05$ and ***$P < 0.001$, by 2-way ANOVA with Bonferroni's multiple comparisons test.

Whole body PACE4 deletion does not impair FGF23 processing under physiological conditions or following IL-1β injections

Combined inactivation of furin and PC5 failed to inhibit FGF23 processing in mice,

suggesting the potential involvement of at least one other enzyme in the regulation of

FGF23 *in vivo*. PACE4 can cleave FGF23 *in vitro* (Figure 5B) and is strongly expressed

in mouse osteoblasts [22]. We therefore next assessed FGF23 processing in PACE4

deficient mice (*Pcsk6-/-*) under physiological conditions and in response to IL-1β. When

fed a standard chow diet, *Pcsk6-/-* mice showed normal total (C-terminal) and intact

FGF23 plasma level compared to wild type littermates. However, 6h following a single IL-

1β (50ng/g) injection, these mice displayed a significant decrease in total (C-terminal)

FGF23 while intact FGF23 was not significantly changed (Figure 7A and 7B). The ratio of

intact over C-terminal FGF23 was equally decreased following IL-1β treatment in wild

type and *Pcsk6-/-* mice. Processed FGF23 was robustly increased by the same treatment

regardless of the genotype, although PACE4 deficient mice did displayed a significant

30% reduction compared to wild type mice (Figure 7C-D). Altogether, these results show

that PACE4 may partially contribute to FGF23 expression or secretion *in vivo* in response

to IL-1β. Yet, FGF23 is still processed in absence of PACE4, suggesting the redundant

action of other PCs.

DISCUSSION

More than 20 years ago, mutations in a putative proprotein convertase (PC) target sequence within human FGF23 were found to cause ADHR in humans. Yet, the identity of the specific PC(s) involved in FGF23 cleavage *in vivo* remained elusive. Taking advantage of floxed alleles of *Furin* and *Psck5*, we generated mice lacking furin and/or PC5 specifically in the osteoblasts and osteocytes lineage and assessed the role of these two enzymes in regulating FGF23 processing *in vivo*. We provide evidence suggesting that furin, but not PC5, partially regulates FGF23 processing *in vivo* under normal conditions. We also exposed the mice to different challenges previously shown to regulate FGF23 expression and its processing and show that FGF23 cleavage is largely mediated by furin under conditions of iron deficiency. The requirement of furin for FGF23 processing appears to be context or cell-type dependent, since FGF23 is still properly cleaved following rhEPO or IL-1β injection when furin is inactivated alone or in combination with PC5. Our results therefore suggest that additional peptidases may redundantly cleave FGF23 *in vivo*.

When mice were fed standard chow diet, the modest increase in circulating intact FGF23 in *Furin*^{osb}*-/-* mice was not sufficient to cause hypophosphatemia and osteomalacia. This result is consistent with the observation that ADHR patients show variable age of onset and penetrance of the disease [35]. Since iron deficiency was shown to positively correlate with FGF23 circulating levels in ADHR patient and to induce hypophosphatemia and osteomalacia in mice carrying an AHDR mutation [14,15], we induced iron deficiency in

control and furin deficient mice. Under these conditions, furin deficient mice did not

develop hypophosphatemia and in fact displayed higher serum phosphate level despite

a near complete impairment of FGF23 cleavage. These observations suggest that

although furin may be important for the cleavage of FGF23 in osteocytes in the context

of iron deficiency, it may also be involved in the regulation of phosphate metabolism

through FGF23-independent mechanism(s).

The results obtained in the context of iron deficiency raised the question whether under

normal conditions FGF23 is redundantly cleaved by another enzyme that is

downregulated in osteocytes following iron deficiency. To clarify whether FGF23 cleavage

by furin is context specific, we injected controls and furin deficient mice with rhEPO or IL-

1β. These two circulating factors are produced in response to anemia and inflammation

respectively [16,36], and were shown to induce FGF23 secretion in bone and bone marrow

cells in rodents [16-19,26,27]. Carboplatin-mediated ablation of bone marrow hematopoietic

cells in mice suggest that osteoblasts and osteocytes may contribute to ~60% of the

increase in circulating FGF23 in response to rhEPO injections [26]. Herein, despite a strong

increase in circulating total (C-terminal) FGF23 level following rhEPO or IL-1β injection,

Furin^{osb}-/- mice did not show a concurrent increase in intact FGF23 suggesting cleavage

by one or several other PC(s) in osteocytes, or that cleaved FGF23 originates mainly from

bone marrow hematopoietic cells in this context. To address this second possibility, we

generated mice deficient in furin in osteoblast-derived cells and/or in all hematopoietic

cells. Surprisingly, our data shows that in these two cellular compartments, furin is

dispensable for FGF23 cleavage following the injection of rhEPO. These results provide

additional credence to the notion that furin is not the only peptidase involved in FGF23

processing *in vivo*.

We previously showed that PC5 (*Pcsk5*), PACE4 (*Pcsk6*) and PC7 (*Pcsk7*) are expressed

in differentiated osteoblasts [22]. These PCs also target the minimal consensus cleavage

site (i.e., RXXR) recognized by furin [37], suggesting that these PCs may also cleave

FGF23. We show herein that a mammalian cell line, furin, but also PC5, PACE4 and PC7

can process FGF23. However, inactivation of *Pcsk5* specifically in osteoblasts and

osteocytes did not impact intact FGF23 level and serum phosphate concentrations. In

addition, mice deficient in both furin and PC5 in the osteoblast lineage are comparable to

furin deficient mice in terms of FGF23 circulating level under physiological conditions and

following rhEPO injection, suggesting that PC5 does not compensate for furin absence *in*

vivo. Finally, even though PACE4 is highly expressed in differentiated osteoblasts [22] and

cleaves FGF23 in mammalian cells, *Pcsk6* inactivation in mice only reduced FGF23

production following IL-1β injection and did not impact its cleavage.

Altogether, these findings provide genetic evidence that furin in osteoblasts and

osteocytes is only partially responsible for FGF23 processing *in vivo*. Our data also

showed that FGF23 cleavage is differentially regulated depending on the physiological

context. Under conditions of iron deficiency furin is responsible for this process, while

following single injection of rhEPO or IL-1β, additional proteases may be involved in

FGF23 processing. Alternately, the increase in cleaved FGF23 following rhEPO and IL-

1β injection could originate from non-osteoblastic cells such as bone marrow

hematopoietic cells and involve multiple proteases including or not furin. Our study also

excluded PC5 and PACE4 as major FGF23 processing enzymes *in vivo.* In conclusion,

additional studies are required to determine precisely which enzyme(s) are involved in

FGF23 cleavage *in vivo* and how furin regulates phosphate metabolism independently of

FGF23.

MATERIALS AND METHODS

Mice and diets

Furin^flox/flox and *Pcsk5^flox/flox* mice were generated by introducing loxP site flanking exon 2 of furin gene and exon 1 of Pcsk5 gene as described previously [38,39]. To generate conditional knockout of furin (*Furin^osb-/-*) or PC5 (*Pcsk5^osb-/-*) in osteoblasts and osteocytes, *Furin^flox/flox* and *Pcsk5^flox/flox* were crossed with OCN-Cre transgenic mice [22], which express the Cre-recombinase under the control of human osteocalcin promoter [21]. *Furin;Pcsk5^osb-/-* mice are deficient in both *Furin* and *Pcsk5* in osteoblasts/osteocytes and were generated by breeding *Furin^osb-/-* and *Pcsk5^flox/flox* mice. *Furin^BM-/-* and *Furin^osb;BM-/-* mice were generated by breeding *Furin^osb-/-* mice with *Furin^flox/+;Vav-iCre* mice. The Vav-iCre expresses the Cre recombinase under the control of the *Vav* promoter, allowing the efficient deletion of furin in fetal and adult hematopoietic stem cells (HSC) [29]. *Pcsk6-/-* mice were provided by Dr. Robert Day (Université de Sherbrooke), these mice were generated as described previously [40]. All mice were backcrossed at least 10 time with the C57BL/6J genetic background, they were maintained at the IRCM pathogen free animal facility on 12h light/dark cycle and fed a normal chow diet. In all experiments, unless otherwise indicated, female mice were used.

In some indicated experiments mice were fed synthetic diet to induce iron or phosphate deficiency. To induce iron deficiency, after weaning at 21 days old, mice were fed either with normal chow diet or an iron deficient diet (Research diet, AIN-76A modified rodent diet, D09070102i) for 14 weeks in combination with bleeding equivalent to one blood

capillary (70 μl) every two weeks. To induce phosphate deficiency, mice were fed for 5

days a normal phosphate diet (Envigo custom diet TD. 98243, 0.6% phosphorus), then

divided to two groups. The first group was maintained for one more week on 0.6% PI diet

while the second group was fed a phosphate deficient diet (Envigo custom diet, TD.

86128, 0.02% phosphorus). In other experiments, mice fed a normal chow diet were

injected with vehicle (saline or PBS), 300U/kg or 3000U/kg of recombinant human

erythropoietin (EPREX, Janssen-Cilag) or 50ng/g IL-1β (STEMCELL Technologies).

Serum biochemistry

Intact and C-terminal FGF23 measurements were performed on EDTA plasma collected

from mice under fed conditions using ELISA assay specific for mouse intact or C-terminal

FGF23 (Quidel, cat#: 60-6800 and cat#: 60-6300). Since C-terminal FGF23 ELISA

detects both intact and processed FGF23, it reflects the total concentration of FGF23 in

the circulation. We also calculated the level of cleaved FGF23 by subtracting intact from

C-terminal values. At end of experiments, mice were anesthetized, and serum was

collected by heart puncture. Serum phosphate level was determined using a phosphorus

assay kit (Sekisui Diagnostic, cat#:117-30). For urine phosphate measurements, urine

was collected in the morning for 3 consecutive days and phosphate level was normalized

to creatinine measured using creatinine assay (Quidel, cat#: 8009).

RNA extraction

Liver, kidney and bone marrow RNA was extracted using the standard protocol previously

described by Chomczynski *et al* [41], while bone RNA was extracted using Trizol reagent

(Thermo Fisher). cDNA was generated by RNA reverse transcription using M-MLV

Reverse Transcriptase (Thermo Fisher). Quantitative real-time PCR (qPCR) was

performed using specific primers (Key Resources Table Appendix) and SYBR Green

qPCR Master Mix (BiMake). Expression levels were normalized to *Actb* expression levels.

Von Kossa/ Van Gieson staining

Mouse skeletons were collected and fixed in 10% formalin for 24 hours and in 70%

ethanol for at least an additional 24 hours. Lumbar vertebrae were collected and

dehydrated by gradual increase of ethanol percentage. Vertebrae were then embedded

in methyl methacrylate resin as described previously [42]. Vertebrae were sectioned at 7-

μm sections and stained with Von kossa/Van Gieson and images were analyzed using

the Osteomeasure analysis system (Osteometrics, Atlanta, GA).

FGF23 processing in cell culture

The cDNAs coding for human furin, PC7, and PACE4, and mouse PC5A and PC5B were

cloned into the pIRES2-EGFP vector [43], while the cDNA of hFGF23 was cloned in

pIRES2-EGFP-V5. CHO-K1 and furin-deficient CHO-FD11 cells were cultured at 37 °C

and 5% CO_2 in Ham's F-12 medium supplemented with 10% (v/v) FBS [44] and transfected

with 3 μg of plasmid following the standard protocol for Lipofectamine 2000 (Thermo

Fisher). For the PCs inhibition, CHO-K1 cells were transfected with cDNAs encoding for

hFGF23-V5 and PCs in the ratio of 5:1 and 24 hours later pretreated for 5 hours with Dec-

RVKR-CMK (50 μM; Tocris) or D6R (20 μM; Calbiochem) followed by 22 hours of

treatment in serum free media. Cells supernatant was then collected in the presence of

1x complete protease inhibitor cocktail (Roche) and FGF23 processing was analyzed by

SDS-PAGE (15% Tris-glycine) followed by western blot using V5 antibody (Thermo

Fisher).

Statistics

Statistical analyses were performed using GraphPad Prism software version 7.03.

Results are shown as the mean ± SEM. Unpaired, 2-tailed Student's *t* test was used for

single measurements or comparison between two groups, while 1-way or 2-way ANOVA

followed by Bonferroni's post-test were used for comparison of more than 2 groups.

A *P* value of less than 0.05 was considered statistically significant. All experiments were

performed at least on 3 independent animals. Before starting each *in vivo* experiment, the

mice were randomized into experimental groups with similar average body weight.

ACKNOWLEDGEMENTS

We thank Dr. R. Day for providing *Pcsk6$^{-/-}$* mice, Dr. T. Clemens for sharing the hOC-Cre mice and Dr. T. Möröy for the iVav-Cre strain. This work was supported by funding from the Canada Research Chair program (M.F.) and the Natural Sciences and Engineering Research Council of Canada (RGPIN-2016-05213, to MF) and a CIHR Foundation grant (# 148363 to NGS). OAR received scholarships from IRCM and FRQS.

CONFLICTS OF INTERESTS

O.A.R., D.S.R., R.E., J.W.M.C., N.G.S., and M.F. have nothing to declare.

REFERENCES

Meyer, R. A., Jr., Meyer, M. H. & Gray, R. W. Parabiosis suggests a humoral factor is involved in X-linked hypophosphatemia in mice. *J Bone Miner Res* **4**, 493-500, doi:10.1002/jbmr.5650040407 (1989).

Miyauchi, A., Fukase, M., Tsutsumi, M. & Fujita, T. Hemangiopericytoma-induced osteomalacia: tumor transplantation in nude mice causes hypophosphatemia and tumor extracts inhibit renal 25-hydroxyvitamin D 1-hydroxylase activity. *J Clin Endocrinol Metab* **67**, 46-53, doi:10.1210/jcem-67-1-46 (1988).

Urakawa, I. *et al.* Klotho converts canonical FGF receptor into a specific receptor for FGF23. *Nature* **444**, 770-774, doi:10.1038/nature05315 (2006).

Larsson, T. *et al.* Transgenic mice expressing fibroblast growth factor 23 under the control of the alpha1(I) collagen promoter exhibit growth retardation, osteomalacia, and disturbed phosphate homeostasis. *Endocrinology* **145**, 3087-3094, doi:10.1210/en.2003-1768 (2004).

Shimada, T. *et al.* FGF-23 is a potent regulator of vitamin D metabolism and phosphate homeostasis. *J Bone Miner Res* **19**, 429-435, doi:10.1359/JBMR.0301264 (2004).

Shimada, T. *et al.* Targeted ablation of Fgf23 demonstrates an essential physiological role of FGF23 in phosphate and vitamin D metabolism. *J Clin Invest* **113**, 561-568, doi:10.1172/JCI19081 (2004).

Shimada, T. *et al.* FGF-23 transgenic mice demonstrate hypophosphatemic rickets with reduced expression of sodium phosphate cotransporter type IIa. *Biochem Biophys Res Commun* **314**, 409-414, doi:10.1016/j.bbrc.2003.12.102 (2004).

Benet-Pages, A., Orlik, P., Strom, T. M. & Lorenz-Depiereux, B. An FGF23 missense mutation causes familial tumoral calcinosis with hyperphosphatemia. *Hum Mol Genet* **14**, 385-390, doi:10.1093/hmg/ddi034 (2005).

Ichikawa, S. *et al.* Genetic rescue of glycosylation-deficient Fgf23 in the Galnt3 knockout mouse. *Endocrinology* **155**, 3891-3898, doi:10.1210/en.2014-1199 (2014).

Topaz, O. *et al.* Mutations in GALNT3, encoding a protein involved in O-linked glycosylation, cause familial tumoral calcinosis. *Nat Genet* **36**, 579-581, doi:10.1038/ng1358 (2004).

Tagliabracci, V. S. *et al.* Dynamic regulation of FGF23 by Fam20C phosphorylation, GalNAc-T3 glycosylation, and furin proteolysis. *Proc Natl Acad Sci U S A* **111**, 5520-5525, doi:10.1073/pnas.1402218111 (2014).

Consortium, A. Autosomal dominant hypophosphataemic rickets is associated with mutations in FGF23. *Nat Genet* **26**, 345-348, doi:10.1038/81664 (2000).

Nesbitt, T., Coffman, T. M., Griffiths, R. & Drezner, M. K. Crosstransplantation of kidneys in normal and Hyp mice. Evidence that the Hyp mouse phenotype is

unrelated to an intrinsic renal defect. *J Clin Invest* **89**, 1453-1459,
doi:10.1172/JCI115735 (1992).

14   Imel, E. A. *et al.* Iron modifies plasma FGF23 differently in autosomal dominant
hypophosphatemic rickets and healthy humans. *J Clin Endocrinol Metab* **96**, 3541-
3549, doi:10.1210/jc.2011-1239 (2011).

15   Farrow, E. G. *et al.* Iron deficiency drives an autosomal dominant
hypophosphatemic rickets (ADHR) phenotype in fibroblast growth factor-23
(Fgf23) knock-in mice. *Proc Natl Acad Sci U S A* **108**, E1146-1155,
doi:10.1073/pnas.1110905108 (2011).

16   David, V. *et al.* Inflammation and functional iron deficiency regulate fibroblast
growth factor 23 production. *Kidney Int* **89**, 135-146, doi:10.1038/ki.2015.290
(2016).

17   Hanudel, M. R. *et al.* Effects of erythropoietin on fibroblast growth factor 23 in mice
and humans. *Nephrol Dial Transplant,* doi:10.1093/ndt/gfy189 (2018).

18   Flamme, I., Ellinghaus, P., Urrego, D. & Kruger, T. FGF23 expression in rodents
is directly induced via erythropoietin after inhibition of hypoxia inducible factor
proline hydroxylase. *PLoS One* **12**, e0186979, doi:10.1371/journal.pone.0186979
(2017).

19   Daryadel, A. *et al.* Erythropoietin stimulates fibroblast growth factor 23 (FGF23) in
mice and men. *Pflugers Arch* **470**, 1569-1582, doi:10.1007/s00424-018-2171-7
(2018).

20   Yamamoto, H. *et al.* Posttranslational processing of FGF23 in osteocytes during
the osteoblast to osteocyte transition. *Bone* **84**, 120-130,
doi:10.1016/j.bone.2015.12.055 (2016).

21   Zhang, M. *et al.* Osteoblast-specific knockout of the insulin-like growth factor (IGF)
receptor gene reveals an essential role of IGF signaling in bone matrix
mineralization. *J Biol Chem* **277**, 44005-44012 (2002).

22   Al Rifai, O. *et al.* Proprotein convertase furin regulates osteocalcin and bone
endocrine function. *J Clin Invest* **127**, 4104-4117, doi:10.1172/JCI93437 (2017).

23   Bon, N. *et al.* Phosphate-dependent FGF23 secretion is modulated by
PiT2/Slc20a2. *Mol Metab* **11**, 197-204, doi:10.1016/j.molmet.2018.02.007 (2018).

24   Yu, X., Sabbagh, Y., Davis, S. I., Demay, M. B. & White, K. E. Genetic dissection
of phosphate- and vitamin D-mediated regulation of circulating Fgf23
concentrations. *Bone* **36**, 971-977, doi:10.1016/j.bone.2005.03.002 (2005).

25   Perwad, F. *et al.* Dietary and serum phosphorus regulate fibroblast growth factor
23 expression and 1,25-dihydroxyvitamin D metabolism in mice. *Endocrinology*
146, 5358-5364, doi:10.1210/en.2005-0777 (2005).

26   Clinkenbeard, E. L. *et al.* Erythropoietin stimulates murine and human fibroblast
growth factor-23, revealing novel roles for bone and bone marrow. *Haematologica*
102, e427-e430, doi:10.3324/haematol.2017.167882 (2017).

27   Toro, L. *et al.* Erythropoietin induces bone marrow and plasma fibroblast growth
factor 23 during acute kidney injury. *Kidney Int* **93**, 1131-1141,
doi:10.1016/j.kint.2017.11.018 (2018).

28   Clinkenbeard, E. L. *et al.* Conditional Deletion of Murine Fgf23: Interruption of the
Normal Skeletal Responses to Phosphate Challenge and Rescue of Genetic
Hypophosphatemia. *J Bone Miner Res* **31**, 1247-1257, doi:10.1002/jbmr.2792
(2016).

29   de Boer, J. *et al.* Transgenic mice with hematopoietic and lymphoid specific
expression of Cre. *European journal of immunology* **33**, 314-325,
doi:10.1002/immu.200310005 (2003).

30   Pesu, M. *et al.* T-cell-expressed proprotein convertase furin is essential for
maintenance of peripheral immune tolerance. *Nature* **455**, 246-250 (2008).

31   Susan-Resiga, D. *et al.* Furin is the major processing enzyme of the cardiac-
specific growth factor bone morphogenetic protein 10. *J Biol Chem* **286**, 22785-
22794, doi:10.1074/jbc.M111.233577 (2011).

32   Susan-Resiga, D. *et al.* Furin Is the Major Processing Enzyme of the Cardiac-
specific Growth Factor Bone Morphogenetic Protein 10. *J Biol. Chem.* **286**, 22785-
22794 (2011).

33   Seidah, N. G. & Prat, A. The biology and therapeutic targeting of the proprotein
convertases. *Nat Rev Drug Discov.* **11**, 367-383 (2012).

34   Essalmani, R. *et al.* In vivo functions of the proprotein convertase PC5/6 during
mouse development: Gdf11 is a likely substrate. *Proc. Natl. Acad. Sci. U. S. A.*
105, 5750-5755 (2008).

35   Imel, E. A., Hui, S. L. & Econs, M. J. FGF23 concentrations vary with disease
status in autosomal dominant hypophosphatemic rickets. *J Bone Miner Res* **22**,
520-526, doi:10.1359/jbmr.070107 (2007).

36   Jelkmann, W. Regulation of erythropoietin production. *J Physiol* **589**, 1251-1258,
doi:10.1113/jphysiol.2010.195057 (2011).

37   Seidah, N. G. & Prat, A. The biology and therapeutic targeting of the proprotein
convertases. *Nature reviews. Drug discovery* **11**, 367-383 (2012).

38   Essalmani, R. *et al.* In vivo functions of the proprotein convertase PC5/6 during
mouse development: Gdf11 is a likely substrate. *Proceedings of the National*
Academy of Sciences of the United States of America **105**, 5750-5755,
doi:10.1073/pnas.0709428105 (2008).

39   Roebroek, A. J. *et al.* Limited redundancy of the proprotein convertase furin in
mouse liver. *J Biol Chem* **279**, 53442-53450, doi:10.1074/jbc.M407152200 (2004).

40   Constam, D. B. & Robertson, E. J. SPC4/PACE4 regulates a TGFbeta signaling
network during axis formation. *Genes & development* **14**, 1146-1155 (2000).

41   Chomczynski, P. & Sacchi, N. The single-step method of RNA isolation by acid
guanidinium thiocyanate-phenol-chloroform extraction: twenty-something years
on. *Nature protocols* **1**, 581-585, doi:10.1038/nprot.2006.83 (2006).

42   Chappard, D., Palle, S., Alexandre, C., Vico, L. & Riffat, G. Bone embedding in
pure methyl methacrylate at low temperature preserves enzyme activities. *Acta*
Histochem **81**, 183-190 (1987).

43   Bergeron, E. *et al.* Implication of proprotein convertases in the processing and
spread of severe acute respiratory syndrome coronavirus. *Biochem Biophys Res*
Commun **326**, 554-563, doi:10.1016/j.bbrc.2004.11.063 (2005).

44   Gordon, V. M., Rehemtulla, A. & Leppla, S. H. A role for PACE4 in the proteolytic
activation of anthrax toxin protective antigen. *Infection and immunity* **65**, 3370-
3375, doi:10.1128/IAI.65.8.3370-3375.1997 (1997).

 FIGURES LEGENDS

Figure 1: Increased FGF23 level and reduced phosphate excretion in *Furin^osb^-/-*

mice. FGF23 and phosphate levels in in 4-month-old *Furin^flox/flox^* (n=6) and *Furin^osb^-/-*

(n=6) mice fed on normal chow diet **(A-D)**. Intact FGF23 levels in plasma **(A)**. FGF23

target genes expression in kidney **(B)**. Serum phosphate level **(C)**. Von Kossa / van

Gieson staining on vertebrae sections **(D)**. FGF23 and phosphate parameters in 6-

months-old *Furin^flox/flox^* (n=6) and *Furin^osb^-/-* (n=6-7) mice fed a normal phosphate diet

(0.6%) or a low phosphate diet (0.02%) for one week **(E-N)**. Body weight **(E)**. Kidney

weight normalize to body weight **(F)**. Serum phosphate levels **(G)**. Plasma level of C-

terminal FGF23 **(H)**, intact FGF23 **(I)**, percentage (%) of intact over C-terminal FGF23 **(J)**

and cleaved FGF23 levels **(K)** calculated as (C-terminal FGF23) - (intact FGF23).

Expression level in kidney of the sodium phosphate cotransporter *Slc34a1* **(L)** and

Slc34a3 **(M). (N)** Urine phosphate measurements normalized to creatinine. Results

represent the mean ± SEM. *$P < 0.05$, **$P < 0.01$, and ***$P < 0.001$, by 2-way ANOVA

with Bonferroni's multiple comparisons test or by Student's t-test (A, C, G, H, K, and M)

#: $P<0.05$; # #: $P<0.01$, # # #: $P<0.001$ &: $0.05<P<0.1$.

Figure 2: Impaired FGF23 processing in *Furin^osb^-/-* mice following iron restriction.

*Furin^flox/flox^* (n=3-5) and *Furin^osb^-/-* (n=7-8) 21-day-old mice were fed normal chow diet (ND)

or a low iron diet (LI) for 14 weeks. **(A)** Body weight growth curve presented as percentage

of initial body weight. **(B-C)** Expression of iron-regulated genes in liver. Hepcidin (*Hamp*)

expression **(B)**. Transferrin receptor (*Tfrc*) expression **(C)**. **(D)** *Fgf23* expression in long

bone. **(E-H)** Plasma level of C-terminal FGF23 **(E)**, intact FGF23 **(F)**, percentage (%) of

intact over C-terminal FGF23 **(G)** and cleaved FGF23 levels **(H)**, calculated as (C-terminal

FGF23) - (intact FGF23). **(I)** Serum phosphate levels. Results represent the mean ± SEM.

*$P < 0.05$, **$P < 0.01$, and ***$P < 0.001$, by 2-way ANOVA with Bonferroni's multiple

comparisons test or by Student's t-test (D and E) #: *$P<0.05$; # #: *$P<0.01$, # # #: *$P<0.001$

&: $0.05<P<0.1$.

Figure 3: FGF23 is normally processed following erythropoietin and IL-1β injection

in *Furin^{osb}-/- * mice. Genes expression in long bone **(A, B and E)** and bone marrow **(C, D**

and F) of 2-month-old C57BL/6J mice 6 hours following the injection of vehicle (n=4), 300

U/kg (n=4) or 3000 U/kg (n=4) of recombinant human erythropoietin (rhEPO).

Erythropoietin receptor (*Epor*) **(A and C)**, erythroferrone (*Erfe*) **(B and D)** and FGF23

(*Fgf23*) **(C and F)**. **(G-J)** Plasma FGF23 measurements in 4-month-old *Furin$^{flox/flox}$* (n=3)

and *Furin^{osb}-/- * (n=3) mice 6 hours following the injection of vehicle, 300 U/kg or 3000 U/kg

of rhEPO. C-terminal FGF23 **(G)**, intact FGF23 **(H),** percentage (%) of intact over C-

terminal FGF23 **(I)** and cleaved FGF23 levels **(J)**, calculated as (C-terminal FGF23) -

(intact FGF23). **(K-N)** Plasma FGF23 measurements in 4-month-old *Furin$^{flox/flox}$* (n=4) and

*Furin^{osb}-/- * (n=4) mice before (T0) and 6 hours following the injection of 50 ng/g IL-1β

(T=6hrs). C-terminal FGF23 **(K)**, intact FGF23 **(L),** percentage (%) of intact over C-

terminal FGF23 **(M)** and cleaved FGF23 levels **(N)**. Results represent the mean ± SEM.

*$P < 0.05$, **$P < 0.01$, and ***$P < 0.001$, by 1-way ANOVA **(A-F)** or 2-way ANOVA **(G-N)**

with Bonferroni's multiple comparisons test, or by Student's t-test **(A and E)** #: *$P<0.05$; #

#: *$P<0.01$, # # #: *$P<0.001$ &: $0.05<P<0.1$.

Figure 4: *Furin* inactivation in osteoblasts, osteocytes and hematopoietic cells did

not impair FGF23 processing following erythropoietin injection. Plasma FGF23

measurements in 8-week-old *Furin$^{flox/flox}$* (n=7), *Furin^{osb}-/-* (n=5), *Furin^{osb}-/-* (n=5) and

Furin$^{osb;BM}$-/- (n=6) mice 6 hours following injection of vehicle or 3000 U/kg of recombinant

human erythropoietin (rhEPO): C-terminal FGF23 **(A)**, intact FGF23 **(B)**, percentage (%)

of intact over C-terminal FGF23 **(C)** and cleaved FGF23 levels **(D)**, calculated as (C-

terminal FGF23) - (intact FGF23). Results represent the mean ± SEM. *$P < 0.05$ and

***$P < 0.001$, by 2-way ANOVA with Bonferroni's multiple comparisons test.

Figure 5: Multiple proprotein convertases can process FGF23 in mammalian cells.

(A) Western blot analysis of the secretion media (SM) from CHO-K1 cells transfected with

V5 tagged human FGF23 (hFGF23-V5) and cultured in the absence (-) or presence (+)

of PC inhibitors, decanoyl-RVKR-CMK (RVKR) (50 μM) or hexa-D-arginine (D6R) (20

μM). **(B)** Western blot analysis of the secretion media (SM) from CHO furin-deficient FD11

cells co-transfected with V5 tagged human FGF23 (hFGF23-V5) and proprotein

convertase Furin, PC5A, PC5B, PACE4 or PC7. Intact and processed C-terminal (CT)

FGF23 forms are detected with V5 antibody.

Figure 6: Genetic ablation of PC5 alone or in combination with Furin does not

impair FGF23 processing *in vivo*. Plasma intact FGF23 **(A)** and serum phosphate

levels **(B)** in 4-month-old *Pcsk5$^{flox/flox}$* (n=6) and *Pcsk5^{osb}-/-* (n=7) mice fed a normal chow

diet. **(C)** Von Kossa / van Gieson staining of vertebrae sections of *Pcsk5$^{flox/flox}$* and

Pcsk5^{osb}-/-. **(D)** Plasma intact FGF23 in 6-month-old *Furin;Pcsk5$^{flox/flox}$* (n=7-8) and

783   *Furin;Pcsk5^{osb}-/-* (n=8) fed a normal phosphate diet (0.6%) or a low phosphate diet

784   (0.02%) for one week. **(E and H)** Plasma FGF23 measurements in 4-month-old

785   *Furin;Pcsk5$^{flox/flox}$* (n=4) and *Furin;Pcsk5^{osb}-/-* (n=4) 6 hours following an injection of

786   vehicle and 300 U/kg of recombinant human erythropoietin (rhEPO): C-terminal FGF23

787   **(E)**, intact FGF23 **(F)**, percentage (%) of intact over C-terminal FGF23 **(G)** and cleaved

788   FGF23 levels **(H)**, calculated as (C-terminal FGF23) - (intact FGF23). Results represent

789   the mean ± SEM. *$P < 0.05$ and ***$P < 0.001$, by 2-way ANOVA with Bonferroni's multiple

790   comparisons test, or by Student's t-test **(A and B)**.

792   **Figure 7: PACE4 inactivation in mice does not impair FGF23 processing.** Plasma

793   FGF23 measurements in 4-month-old *Pcsk6 +/+* (n=4) and *Pcsk6 -/-* (n=4) male mice

794   before (T0) and 6 hours following the injection of 50 ng/g of IL-1β: C-terminal FGF23 **(A)**,

795   intact FGF23 **(B)**, percentage (%) of intact over C-terminal FGF23 **(C)**, and cleaved

796   FGF23 levels **(D)**, calculated as (C-terminal FGF23) - (intact FGF23). Results represent

797   the mean ± SEM. *$P < 0.05$ and ***$P < 0.001$, by 2-way ANOVA with Bonferroni's multiple

798   comparisons test.

Key Resources Table Appendix: Primers used				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
Sequence-based reagent	Actin beta- For	This paper	QPCR primer (amplify *Actb*, M. musculus)	GACCTCTAT GCCAACACAGT
Sequence-based reagent	Actin beta- Rev	This paper	QPCR primer (amplify *Actb*, M. musculus)	AGTACTTGC GCTCAGGAGGA
Sequence-based reagent	Slc34a1-Fw	This paper	QPCR primer (amplify *Slc34a1*, M. musculus)	GCATCCTACTGTGGTAC CCG
Sequence-based reagent	Slc34a1-Rv	This paper	QPCR primer (amplify *Slc34a1*, M. musculus)	CAGCAAACCAGCGGTA CTTG
Sequence-based reagent	Slc34a3-Fw	This paper	QPCR primer (amplify *Slc34a3*, M. musculus)	CCTTTGGACTTTCCCTG GCA
Sequence-based reagent	Slc34a3-Rv	This paper	QPCR primer (amplify *Slc34a3*, M. musculus)	GATGGTCGGTGTTGTT GCAG
Sequence-based reagent	Hamp-Fw	This paper	QPCR primer (amplify *Hamp*, M. musculus)	AGAGCTGCAGCCTTTG CAC
Sequence-based reagent	Hamp-Rv	This paper	QPCR primer (amplify *Hamp*, M. musculus)	GAGGTCAGGATGTGGC TCTA
Sequence-based reagent	Tfrc-Fw	This paper	QPCR primer (amplify *Tfrc*, M. musculus)	TAAATTCCCCGTTGTTG AGG
Sequence-based reagent	Tfrc-Rv	This paper	QPCR primer (amplify *Tfrc*, M. musculus)	CAGGACAGCTTCCTTC CATT
Sequence-based reagent	Cyp27b1-Fw	This paper	QPCR primer (amplify *Cyp27b1*, M. musculus)	GAAGCTGCGATGAGGA ACCA
Sequence-based reagent	Cyp27b1-Rv	This paper	QPCR primer (amplify *Cyp27b1*, M. musculus)	TTCCCCACTATGGACTG GACA
Sequence-based reagent	Cyp24a1-Fw	This paper	QPCR primer (amplify *Cyp24a1*, M. musculus)	CCCAGCGGCTAGAGAT CAAA
Sequence-based reagent	Cyp24a1-Rv	This paper	QPCR primer (amplify *Cyp24a1*, M. musculus)	TTTCTTTTGGAAGGCGC TGC

Sequence-based reagent	Fgf23-Fw	This paper	QPCR primer (amplify *Fgf23*, M. musculus)	CCGCTGCTGCATTTCTACAC
Sequence-based reagent	Fgf23-Rv	This paper	QPCR primer (amplify *Fgf23*, M. musculus)	CTTGAGCACGTTCAGTGGGT
Sequence-based reagent	Epor-Fw	This paper	QPCR primer (amplify *Epor*, M. musculus)	GGACACAAAGGGTGGAGGTC
Sequence-based reagent	Epor-Rv	This paper	QPCR primer (amplify *Epor*, M. musculus)	GCTCGAACAGCGAAGGTGTA
Sequence-based reagent	Erfe-Fw	This paper	QPCR primer (amplify *Erfe*, M. musculus)	GTTCCAGCTGTTGCTGAAAGG
Sequence-based reagent	Erfe-Rv	This paper	QPCR primer (amplify *Erfe*, M. musculus)	TACCCGAGGCTGGTGTAGTG