Protein Phosphatase 1 regulates atypical chromosome segregation and cell polarity during mitotic and meiotic division in *Plasmodium* sexual stages

Mohammad Zeeshan[1], Rajan Pandey[1,$], Amit Kumar Subudhi[2,$], David J. P. Ferguson[3,4,$], Gursimran Kaur[1], Ravish Rashpa[5], Raushan Nugmanova[2], Declan Brady[1], Andrew R. Bottrill[6], Sue Vaughan[4], Mathieu Brochet[5], Mathieu Bollen[7], Arnab Pain[2,8], Anthony A. Holder[9], David S. Guttery[1,10], Rita Tewari[1,*]

[1]*School of Life Sciences, University of Nottingham, Nottingham, UK*

[2]*Pathogen Genomics Group, BESE Division, King Abdullah University of Science and Technology (KAUST), Thuwal, Kingdom of Saudi Arabia*

[3]*Nuffield Department of Clinical Laboratory Sciences, University of Oxford, John Radcliffe Hospital, Oxford, UK*

[4]*Department of Biological and Medical Sciences, Faculty of Health and Life Science, Oxford Brookes University, Gipsy Lane, Oxford, UK*

[5]*Department of Microbiology and Molecular Medicine, Faculty of Medicine, University of Geneva, Geneva, Switzerland*

[6]School of Life Sciences, Gibbet Hill Campus, University of Warwick, Coventry CV4 7AL, UK

[7]*Laboratory of Biosignaling and Therapeutics, KU Leuven Department of Cellular and Molecular Medicine, University of Leuven, B-3000 Leuven, Belgium.*

[8]*Research Center for Zoonosis Control, Global Institution for Collaborative Research and Education (GI-CoRE); Hokkaido University, Sapporo, Japan*

[9]*Malaria Parasitology Laboratory, The Francis Crick Institute, London, UK*

[10]*Leicester Cancer Research Centre, University of Leicester, Leicester, UK*

[$]These authors contributed equally

*For correspondence

Rita Tewari: rita.tewari@nottingham.ac.uk

Short title: Role of PP1 during mitosis and meiosis in *Plasmodium*

Abstract

PP1 is a conserved eukaryotic serine/threonine phosphatase that regulates many aspects of mitosis and meiosis, often working in concert with other phosphatases, such as CDC14 and CDC25. The proliferative stages of the parasite life cycle include sexual development within the mosquito vector, with male gamete formation characterized by an atypical rapid mitosis, consisting of three rounds of DNA synthesis, successive spindle formation with clustered kinetochores, and a meiotic stage during zygote to ookinete development following fertilization. It is unclear how PP1 is involved in these unusual processes. Using real-time live-cell and ultrastructural imaging, conditional gene knockdown, RNA-seq and proteomic approaches, we show that *Plasmodium* PP1 is involved in both chromosome segregation during mitotic exit, and establishment of cell polarity during zygote development in the mosquito midgut, suggesting that small molecule inhibitors of PP1 should be explored for blocking parasite transmission.

Introduction

Cell cycle progression involves sequential and highly ordered DNA replication and chromosome segregation in eukaryotes[1, 2], which is tightly controlled and coordinated by reversible protein phosphorylation catalysed by protein kinases (PKs) and protein phosphatases (PPs)[3]. Numerous studies have highlighted the importance of the phosphoprotein phosphatase (PPP) family in regulating mitosis in model eukaryotic organisms, often working in conjunction with CDC25 and CDC14 phosphatases, as key regulators of mitotic entry and exit, respectively[4, 5, 6, 7].

Protein Phosphatase 1 (PP1) is a member of the PPP family and is expressed in all eukaryotic cells. It plays a key role in the progression of mitosis through dephosphorylation of a large variety of proteins, including mitotic kinases (such as cyclin-dependent kinase 1 (CDK1)[8, 9, 10, 11], and regulators of chromosome segregation[12]. Chromosome segregation is orchestrated at the kinetochore, a protein complex that assembles on the centromeres, located at the constriction point of sister chromatids to facilitate and monitor attachment of the sister chromatids to spindle microtubules[13, 14]. Correct attachment of the kinetochore to spindle microtubules is regulated by the KMN (KNL1, MIS12 and NDC80) protein network[15], which in mammalian cells integrates the activities of at least five protein kinases (including MPS1, Aurora B, BUB1, PLK1, and CDK1) and two protein phosphatases (PP1 and PP2A-B56). The KMN network mediates kinetochore-microtubule attachments and scaffolds the spindle assembly checkpoint (SAC) to prevent chromosome segregation until all sister chromatids are properly connected to the spindle[16]. The orchestration of reversible protein phosphorylation is crucial to control the spatial-temporal progression of the cell cycle and PP1 has a key role in this

process, in particular during mitotic exit. Throughout mitosis, PP1 is inhibited by the

cyclin-dependent kinase (CDK1)/cyclin B complex and Inhibitor 1[17, 18]; however,

during mitotic exit concomitant destruction of cyclin B and reduced activity of CDK1

through dephosphorylation by CDC14 results in subsequent reactivation of PP1 via

autophosphorylation and completion of mitotic exit[10, 17, 19, 20, 21].

In *Plasmodium*, the causative organism of malaria, there is a single PP1 orthologue,

which is expressed throughout the parasite's complex life-cycle[22] and located in both

nucleus and cytoplasm[23]. Previous studies determined that *P. falciparum* PP1

functionally complements the *Saccharomyces cerevisiae* glc7 (PP1) homologue[24],

and subsequent phylogenetic analyses revealed that homologues of the

phosphatases CDC14 and CDC25 are absent from *Plasmodium*[23]. Genetic screens

and inhibitor studies have shown that PP1 is essential for asexual blood stage

development[23, 25, 26, 27] (also reviewed[28]), in particular for parasite egress from the

host erythrocyte[29]. Most studies in *Plasmodium* have been focused on these asexual

blood stages of parasite proliferation, and very little is known about the importance of

PP1 for transmission stages within the mosquito vector. Recent phosphoproteomic

studies and chemical genetics analysis have identified a number of potential

Plasmodium falciparum (Pf) PP1 substrates modified during egress, including a

HECT E3 protein-ubiquitin ligase and GCα, a guanylyl cyclase with a phospholipid

transporter domain[29]. Other biochemical studies in *Plasmodium* have identified

numerous PP1-interacting partners (PIPs) that are structurally conserved and

regulate PP1 activity, including LRR1 (a human SDS22 orthologue), Inhibitor 2 and

Inhibitor 3. Other PIPs including eif2β, and GEXP15 have also been identified[9, 28, 30,

[31]. However, little is known regarding how PP1 is involved in regulating mitosis and

meiosis in the absence of CDC14 and CDC25.

The *Plasmodium* life-cycle is characterised by several mitotic stages and a single

meiotic stage (reviewed in[32, 33, 34]). However, in this organism kinetochore dynamics

and chromosome segregation are poorly understood, especially in mitosis during

male gametogenesis. This is a very rapid process with three rounds of spindle

formation and genome replication from 1N to 8N within 12 minutes. In addition, little

is known about the first stage of meiosis that occurs during the zygote to ookinete

transition. We have followed recently the spatio-temporal dynamics of NDC80

throughout mitosis in schizogony, sporogony and male gametogenesis, and during

meiosis in ookinete development[35], using approaches that offer the opportunity to

study the key molecular players in these crucial stages of the life cycle. Although

PP1 has been partially characterised and shown to have an essential role during

asexual blood stage development in the vertebrate host[23], the role and importance of

PP1 during sexual stage development in the mosquito is completely unknown.

Here, using the *Plasmodium berghei* (Pb) mouse model of malaria, we determined

the importance of PP1 during the sexual stages within the mosquito vector. PP1

expression and location were studied using the endogenous, GFP-tagged protein

and co-localisation with the kinetochore marker, NDC80, to follow progression

through chromosome segregation during male gamete formation and zygote

differentiation. Using a conditional gene knockdown approach we examined how

PP1 orchestrates atypical mitosis and meiosis, and investigated the ultrastructural

consequences of PP1 gene knockdown for cell morphology, nuclear pole

multiplication and flagella formation during male gamete formation. RNA-Seq

analysis was used to determine the consequence of PP1 gene knockdown on global

transcription, which disclosed a marked differential expression of genes involved in

reversible phosphorylation, motor activity and the regulation of cell polarity.

Proteomics studies identified motor protein kinesins as interacting partners of PP1 in

the gametocyte. The knockdown of PP1 gene expression blocks parasite

transmission by the mosquito, showing that this protein has a crucial function in

Plasmodium sexual development during both mitosis in male gamete formation and

meiosis during zygote to ookinete differentiation.

Results

PP1-GFP has a diffuse subcellular location during asexual blood stage

schizogony and forms discrete foci during endomitosis.

To examine the spatio-temporal expression of *Plasmodium* PP1 in real-time during

cell division, we generated a transgenic *P. berghei* line expressing endogenous PP1

with a C-terminal GFP tag. An in-frame *gfp* coding sequence was inserted at the 3'

end of the endogenous *pp1* locus using single crossover homologous recombination

(**Fig S1A**), and successful insertion was confirmed by diagnostic PCR (**Fig S1B**).

Western blot analysis of schizont protein extracts using an anti-GFP antibody

revealed a major 62-kDa band, the expected size for PP1-GFP protein, compared to

the 29 kDa WT-GFP (**Fig S1C**).

During asexual blood stage proliferation, schizogony is characterised by multiple

rounds of asynchronous nuclear division without chromosome condensation or

cytokinesis. Nuclear division is a closed mitotic process without dissolution-

reformation of the nuclear envelope and with the spindle-pole body

(SPB)/microtubule-organising centre (MTOC) embedded within the nuclear

membrane[36]. It results in a multinucleated coenocyte termed a schizont, which is

resolved by cytokinesis at the end of schizogony into individual merozoites. Live cell

imaging of *P. berghei* asexual blood stages revealed a diffuse cytoplasmic and

nuclear distribution of PP1-GFP, together with a distinct single focus of concentrated

PP1-GFP in the nucleus of early trophozoite stage (**Fig 1A**), representing the early

S-phase of the cell cycle when DNA synthesis starts. As schizogony proceeds, the

diffuse distribution of PP1-GFP remained; however, in early schizonts each cell

displayed two distinct PP1-GFP foci in close association with the stained nuclear

DNA. These pairs of PP1-GFP foci became clearer during the middle and late

schizont stages but following merozoite maturation and egress the intensity of the

foci diminished (**Fig 1A**).

To study further the location of the PP1-GFP foci throughout mitosis, we generated

by genetic cross parasite lines expressing either PP1-GFP and the kinetochore

marker NDC80-mCherry[35] or PP1-GFP and the inner membrane complex (IMC)-

associated myosin A (MyoA)-mCherry. Live cell imaging of these lines revealed co-

localisation of PP1-GFP and NDC80-mCherry foci close to the DNA of the nucleus

through blood stage development, and especially during late schizogony and

segmentation (**Fig 1B**); whereas the PP1-GFP foci at the outer periphery of the cell

showed a partial co-location with MyoA-mCherry, and only during mid- to late

schizogony (**Fig 1C**). Thus, in schizonts there is a concentration of PP1-GFP at the

kinetochore and at a transient peripheral location during specific stages of

schizogony, as well as the diffuse distribution throughout the cytoplasm (**Fig.1**).

PP1-GFP is enriched on kinetochores during chromosome segregation associated with putative mitotic exit in male gametogony

To study PP1 expression and location through the three rounds of DNA replication and chromosome segregation prior to nuclear division during male gametogony, we examined PP1-GFP by live-cell imaging over a 15-minute period following male gametocyte activation. Following activation with xanthurenic acid and decreased temperature, male gametocytes undergo three rounds of DNA replication and mitotic division followed by chromosome condensation and exflagellation, resulting in eight gametes[37]. Before activation (0 min), PP1-GFP was detected with a diffuse location throughout gametocytes (**Fig 2A**). At one minute post-activation, PP1-GFP accumulated at one end of the nucleus at a single focal point (**Fig 2A**). After 2-3 min two distinct foci were observed on one side of the nucleus, concurrent with the first round of chromosome replication/segregation (**Fig 2A**). Subsequently, four and eight PP1-GFP foci were observed at 6 to 8 min and 8 to 12 min post-activation, respectively, corresponding to the second and third rounds of chromosome replication/segregation. These discrete PP1-GFP foci dispersed prior to karyokinesis and exflagellation of the mature male gamete 15 min post-activation, leaving residual protein remaining diffusely distributed throughout the remnant gametocyte and flagellum (**Fig 2A**).

To investigate further the location of PP1-GFP during spindle formation and chromosome segregation, the parasite line expressing both PP1-GFP and NDC80-mCherry was examined by live cell imaging to establish the spatio-temporal relationship of the two proteins. We found that the discrete PP1-GFP foci colocalized

with NDC80-mCherry at different stages of male gametogony, including when up to

eight kinetochores were visible **(Fig 2B)**, but PP1-GFP was not associated with the

arc-like bridges of NDC80-mCherry representing the spindle[35] at 2 min and 4 to 5

min post activation, suggesting that PP1-GFP is only increased at the kinetochore

during initiation and termination of spindle division (**Fig 2B**).

PP1-GFP may determine apical polarity during zygote-ookinete development

and has a nuclear location on kinetochores during meiosis

After fertilisation in the mosquito midgut, the diploid zygote differentiates into an

ookinete within which the first stage of meiosis occurs. During this process DNA is

duplicated to produce a tetraploid cell with four haploid genomes within a single

nucleus in the mature ookinete. PP1 is known to be crucial during meiotic

chromosome segregation[12], and therefore we analysed the spatio-temporal

expression of PP1-GFP during ookinete development using live cell imaging. PP1-

GFP was expressed in both male and female gametes with a diffuse distribution,

along with a single focus of intense fluorescence at one end (potentially the basal

body) of each male gamete (**Fig. 3A**). Initially, the zygote also had a diffuse PP1-

GFP distribution, but after two hours (stage I) an enriched focus developed at the

periphery of the zygote, marking the point that subsequently protruded out from the

cell body and developed into the apical end of the ookinete. A strong PP1-GFP

fluorescence signal remained at this apical end throughout ookinete differentiation

(**Fig 3A**). In addition, PP1-GFP was enriched in the nucleus at four discrete foci in

mature ookinetes (**Fig 3A**). Analysis of the parasite line expressing both PP1-GFP

and NDC80-mCherry showed that these four foci correspond to the kinetochores of

meiotic development in the ookinete (**Fig 3B**).

PP1-GFP has a diffuse distribution with multiple nuclear foci during oocyst development, representing endomitosis

Upon maturation the ookinete penetrates the mosquito midgut wall and embeds in the basal lamina to form an oocyst. Over the course of 21 days sporogony occurs in which several rounds of closed endomitotic division produce a multiploid cell termed a sporoblast[38]; with subsequent nuclear division resulting in thousands of haploid nuclei and concomitant sporozoite formation. We observed multiple foci of PP1GFP along with diffused localization during oocyst development (**Fig 3C**). Dual colour imaging showed a partial co-localisation of nuclear PP1-GFP and NDC80-mCherry foci throughout oocyst development and in sporozoites (**Fig 3D**), suggesting that PP1-GFP is diffusely distributed during oocyst development but also recruited to kinetochores, as highlighted by multiple nuclear foci during oocyst development.

Generation of transgenic parasites with conditional knockdown of PP1

Our previous systematic analysis of the *P. berghei* protein phosphatases suggested that PP1 is likely essential for erythrocytic development[23], a result which was further substantiated in a recent study of *P. falciparum*[29]. Since little is known about the role of PP1 in cell division during the sexual stages of parasite development in the mosquito vector, we attempted to generate transgenic, conditional knockdown lines using either the auxin inducible degron (AID) or the promoter trap double homologous recombination (PTD) systems. Despite successful generation of a parasite line expressing PP1-AID, addition of indole-3-acetic acid (IAA) did not result in PP1 depletion (**Fig S2C**) and male gametogony was not affected (**Fig S2C**). Therefore, we used a promoter trap method that had been used previously for

functional analysis of the condensin core subunits SMC2/4 and the APC/C complex

component, APC3[39, 40]. Double homologous recombination was used to insert the

ama promoter upstream of *pp1* in a parasite line that constitutively expresses GFP[41].

The expected outcome was a parasite expressing PP1 during asexual blood stage

development, but not at high levels during the sexual stages (**Fig S2D**). This strategy

resulted in the successful generation of two independent PP1PTD parasite clones

(clones 3 and 5) produced in independent transfections, and integration confirmed by

diagnostic PCR (**Fig S2E**). The PP1PTD clones had the same phenotype and data

presented here are combined results from both clones. Quantitative real time PCR

(qRT-PCR) confirmed a downregulation of *pp1* mRNA in gametocytes by

approximately 90% (**Fig 4A**).

Knockdown of PP1 gene expression during the sexual stages blocks parasite

transmission by affecting parasite growth and development within the

mosquito vector

The PP1PTD parasites grew slower than the WT-GFP controls in the asexual blood

stage (**Fig S2E**) but produced normal numbers of gametocytes in mice injected with

phenylhydrazine before infection. Gametocyte activation with xanthurenic acid in

ookinete medium resulted in significantly fewer exflagellating PP1PTD parasites per

field in comparison to WT-GFP parasites, suggesting that male gamete formation

was severely retarded (**Fig. 4B**). For those few PP1PTD gametes that were released

and viable, zygote-ookinete differentiation following fertilization was severely affected

(**Fig 4C**), with significantly reduced numbers of fully formed, banana-shaped

ookinetes. In all PP1PTD samples analysed 24 hours post-fertilisation, the vast

majority of zygotes were still round, and there was a significant number of abnormal

retort-shaped cells with a long thin protrusion attached to the main cell body (**Fig 4C,**
D).

To investigate further the function of PP1 during parasite development in the
mosquito, *Anopheles stephensi* mosquitoes were fed on mice infected with PP1PTD
and WT-GFP parasites as a control. The number of GFP-positive oocysts on the
mosquito gut wall was counted on days 7, 14 and 21 post-infection. No oocysts were
detected from PP1PTD parasites; whereas WT-GFP lines produced normal oocysts
(**Fig 4E, F**), all of which had undergone sporogony (**Fig 4G**). Furthermore, no
sporozoites were observed in the salivary glands of PP1PTD parasite-infected
mosquitoes, in contrast to WTGFP parasite-infected mosquitoes (**Fig 4H**). This lack
of viable sporozoites was confirmed by bite back experiments that showed no further
transmission of PP1PTD parasites, in contrast to the WT-GFP lines, which showed
positive blood stage infection four days after mosquito feed (**Fig S2F**).

Ultrastructure analysis of PP1PTD gametocytes shows defects in nuclear pole
and axoneme assembly
To determine the ultrastructural consequences of reduced PP1 expression, WTGFP
and PP1PTD gametocytes were examined at 6- and 30-min post activation (pa) by
electron microscopy. In mutant PP1PTD lines, at 6-min pa microgametocytes
contained a large central nucleus with a few nuclear poles (**Fig 5a**). Basal bodies
and axonemes were often associated with the nuclear poles (**Fig 5a, b**). In cross
section, certain, but not all, axonemes displayed the 9+2 microtubular arrangement
(**Fig 5c**). These appearances were similar to those observed in the WTGFP line at 6-

min pa (**Fig 5a-c and Fig 5g-i**). However, when counts were made of the relative

number of nuclear poles, basal bodies and axonemes, quantitative differences were

observed with, PP1PTD gametocytes appearing less developed with fewer nuclear

poles (0.86 WT v 0.56 Mutant), basal bodies (1.03 WT v 0.56 Mutant) and axonemes

(2.57 WT v 1.27 Mutant); the relative numbers were based on random sections

through 30 microgametocytes in each group.

At 30-min pa, the majority of PP1PTD microgametocytes still appear similar to those

at 6-min pa, exhibiting an open nucleus with a few nuclear poles, basal bodies and

associated axonemes (**Fig 5a-c and Fig 5d-f**). The main morphological difference

was a marked increase in the number and length of the axonemes (**Fig 5a and Fig**

5d). In contrast, the majority of WTGFP microgametocytes, at 30-min pa, were at an

advanced stage of development with a number displaying nuclei with chromatin

condensation (**Fig 5j**) and others exhibiting examples of active exflagellation with the

flagellum protruding from the surface (**Fig 5k**). This was in addition to the presence

of free microgametes (**Fig 5j, l**). In summary the development of the PP1PTD

microgametocytes was severely retarded although there was an increase in

axoneme growth.

PP1PTD parasites have altered expression of genes involved in cell cycle

progression, cell motility, and apical cell polarity

To determine the consequences of PP1 knockdown on global mRNA transcription,

we performed RNA-seq in duplicate on PP1PTD and WT-GFP gametocytes, 0 min

and 30 min post-activation (**Fig 6A and B**). All replicates were clustered together

based on condition (**Fig S3A**) and totals of 13 to 32 million RNA-seq reads were

generated per sample for the global transcriptome **(Fig S3B).** In both 0 min and 30

min activated gametocytes, *pp1* was significantly down-regulated (by more than 16-

fold, q value < 0.05) in PP1PTD parasites compared to WT-GFP parasites (**Fig 6A**),

thus confirming the qPCR result (**Fig 4A**). We observed significant transcriptional

perturbation in both 0 min and 30 min activated gametocytes, affecting the

expression of 530 and 829 genes respectively (representing ~ 10% and 16% of the

total gene complement) (**Table S1**). Of the total perturbed genes, 344 and 581 were

significantly down-regulated and 186 and 248 genes were significantly up-regulated

in the PP1PTD parasites compared to WT-GFP controls in 0 min and 30 min

activated gametocytes, respectively (**Table S1**).

To explore the biological roles of the genes down-regulated following reduced *pp1*

expression, we performed gene-ontology (GO) based enrichment analysis. We

observed that many genes encoding kinases, phosphatases and motor proteins

were differentially expressed in either or in both 0 min or 30 mins activated

gametocytes (**Fig 6C**), complementing the observations from our phenotypic

analysis (**Fig 4C, D**). In addition, we also observed that many genes encoding

proteins involved in cell motility, cell-cycle progression, host-cell entry and ookinete

development were significantly affected in activated PP1PTD gametocytes.

Transcript levels measured by RNA-seq were further validated by qRT-PCR for a

few selected genes modulated in gametocytes at 0 min and 30 min, and involved in

regulation of cell cycle, cell motility, and ookinete invasion (**Fig 6D**).

PP1–GFP interacts with similar proteins in schizonts and gametocytes, but

with a predominance of microtubule motor kinesins in gametocytes

Previous studies have analysed the PP1 interactome in *P. falciparum* schizonts,

revealing several interacting partners[31, 42]. Here, we analysed the PP1 interactome in

P. berghei schizonts and gametocytes to establish whether there were differences

that might reflect distinct functions in the two stages. We immunoprecipitated PP1-

GFP from lysates of schizonts following parasite culture *in vitro* for 10 hours and 24

hours and from lysates of gametocytes 10 to 11 min post activation because of high

PP1-GFP abundance at these stages (**Fig 7A, Table S2**). Mass spectrometric

analysis of these pulldowns identified several proteins common to both schizont and

gametocyte stages, suggesting similar protein complexes in both stages (**Fig 7A, B**).

In addition, we also identified in the pulldown from gametocyte lysates a number of

microtubule proteins that are associated with the spindle or axoneme, including

kinesin-8B, kinesin-15, kinesin-13 and PF16 (**Fig 7A, B)**. These proteins are specific

to male gametocytes and may have important roles in axoneme assembly and male

gametogony[37, 43].

Discussion

Reversible protein phosphorylation is crucial for cell cycle progression in eukaryotes

and is tightly controlled by a variety of protein kinases and phosphatases[44, 45].

Plasmodium possesses a set of divergent protein kinases and protein

phosphatases, which regulate many processes during cell division and parasite

development throughout different stages of the life-cycle[23, 46], and PP1 is

quantitatively one of the most important protein phosphatases that hydrolyse

serine/threonine linked phosphate ester bonds[47]. It is expressed in all cells and

highly conserved in organisms throughout the Eukaryota including *Plasmodium*[23, 48].

It forms stable complexes with several PP1-interacting proteins (PIPs) that are

diverse and differ in various organisms, control its location and function, and assist in

processes throughout the cell cycle[8, 49, 50, 51].

The cell cycle in *Plasmodium* differs from that of many other eukaryotes with atypical

mitotic and meiotic divisions throughout its life cycle[35, 37, 52]. It lacks several classical

cell cycle regulators including the protein phosphatases CDC14 and CDC25[23], and

key protein kinases including polo-like kinases[46, 53]. Therefore, the role of PP1, or

indeed other PPPs in the absence of these protein phosphatases and kinases may

be more crucial during cell division in *Plasmodium*. In the present study, we focused

on the location and functional role of PP1 in mitotic division, particularly during male

gametogony, and in meiosis during the zygote to ookinete transition. Both processes

are essential for the sexual stage of the *Plasmodium* life cycle and are required for

transmission by the mosquito vector. Our previous systematic functional analysis of

the *Plasmodium* protein phosphatome showed that PP1 has an essential role during

asexual blood stage development; this finding was supported by a recent study,

which showed its role in merozoite egress from the host erythrocyte[29]. Our findings

here reveal that PP1 is also expressed constitutively and co-localises with the

kinetochore protein NDC80 during different stages of *Plasmodium* asexual and

sexual development, hinting at a role during atypical chromosome segregation. Our

conditional PP1 gene knockdown suggests that it plays a crucial role in mitotic

division during male gametogony; and regulates cell polarity in meiosis during zygote

to ookinete transformation.

Male gametogony in *Plasmodium* is a rapid process with DNA replication and three

mitotic divisions followed by karyokinesis and cytokinesis to form eight flagellated

male gametes within 15 min of gametocyte activation[35, 52]. The expression and

localization profiles show that although the protein is diffusely distributed throughout

the cytoplasm and the nucleus, there is a cyclic enrichment of PP1-GFP at the

kinetochore, associating with NDC80-mCherry during all three successive genome

replication and closed mitotic divisions without nuclear division[35]. The accumulation

of PP1-GFP at the kinetochore at the start of nuclear division and subsequent

decrease upon completion is similar to the situation in other eukaryotes where the

activity of PP1 increases in G2 phase, reduces during prophase and metaphase, and

increases again during anaphase[45]. This behaviour suggests a role for PP1 in

regulating rapid mitotic entry and exit during male gametogony. Further analysis of

PP1 gene knockdown showed a significant decrease in male gamete formation (i.e.

during male gametogony), and ultrastructural analysis revealed fewer nuclear poles

and basal bodies associated with axonemes, with a concomitant absence of

chromosome condensation in male gametocytes of PP1PTD parasites, suggesting a

role for PP1 during chromosome segregation and gamete formation (i.e. flagella

formation). A similar phenotype was observed in our recent study of a divergent

Plasmodium cdc2-related kinase (CRK5)[54]. CRK5 deletion resulted in fewer nuclear

poles, and no chromatin condensation, cytokinesis or flagellum formation,

suggesting that there may be a coordinated activity of PP1 and CRK5 in reversible

phosphorylation. A recent phosphoregulation study across the short time period of

male gametogony showed tightly controlled phosphorylation events mediated by

several protein kinases including ARK2, CRK5, and NEK1[55]; however, the reciprocal

protein phosphatases that reverse these events are unknown. The expression

pattern and cyclic enrichment of PP1 at the kinetochores suggest a key reciprocal

role of PP1 in reversing the protein phosphorylation mediated by these kinases. This

interpretation is supported by our global transcriptome analysis of the PP1PTD

parasite, which showed a modulated expression of several serine/threonine protein

kinases, protein phosphatases and motor proteins. The similar pattern of reduced

expression of some of these protein phosphatases and kinases suggests they have

a coordinated role in reversible phosphorylation. Of note, phosphoregulation of motor

proteins in *Plasmodium* has been described previously[55] and it plays an important

role in spindle assembly and axoneme formation during male gametogony[35, 37].

The proteomics analysis showed that PP1 interacts with a conserved set of proteins

in both asexual and sexual stages of development, and with additional proteins in

gametocytes, implicating major motor proteins that may be required for spindle and

axoneme assembly during male gametogenesis. Several of the conserved

interacting partners also form complexes with PP1 in other eukaryotes but the

gametocyte specific proteins such as kinesin-8B, kinesin-15, kinesin-13, and PF16

are unique PIPs in *Plasmodium*[9, 42]. These findings are consistent with our

transcriptomic analysis of PP1PTD showing modulated expression of motor proteins

in male gametocytes.

Analysis of PP1 during the zygote to ookinete transformation showed an additional

location that defines the cell's polarity: at the apical end during the early stages of

ookinete development. These data suggest that PP1 may also regulate cell shape

during ookinete differentiation. This suggestion is supported by the consequence of

PP1PTD gene knock down in which numerous underdeveloped ookinetes with long,

thin protrusions attached to the main cell body were observed. This is substantiated

by a recent study that showed PP1-mediated and SDS22 dependent

dephosphorylation of LGL, an actomyosin-associated protein, controls apical-basal

polarity in *Drosophila*[56]. Our transcriptomic analysis of PP1PTD-gametocytes

showed the modulation of several proteins involved in gliding motility including IMC

proteins, MyoA, GAPs, GAPMs and ookinete specific proteins. These proteins are

important for maintenance of cell shape, gliding motility, and mosquito gut wall

invasion by *Plasmodium* ookinetes. This suggests that PP1 is involved in the

phospho-regulation of proteins involved in defining polarity and maintenance of

ookinete shape.

In conclusion, PP1 is a constitutively expressed phosphatase, distributed throughout

the cell but enriched in the nucleus, and associated with the kinetochore during

mitosis and meiosis, with a role in the regulation of mitosis and meiosis throughout

the *Plasmodium* life-cycle.

Material and Methods

Ethics statement

The animal work performed in the UK passed an ethical review process and was

approved by the United Kingdom Home Office. Work was carried out under UK

Home Office Project Licenses (40/3344 and 30/3248) in accordance with the United

Kingdom 'Animals (Scientific Procedures) Act 1986'. Six- to eight-week-old female

CD1 outbred mice from Charles River laboratories were used for all experiments in

the UK.

Generation of transgenic parasites

GFP-tagging vectors were designed using the p277 plasmid vector and

transfected as described previously[23]. A schematic representation of the

endogenous *pp1* locus (PBANKA_1028300), the constructs and the recombined *pp1*

locus can be found in **Fig S1A**. For GFP-tagging of PP1 by single crossover

homologous recombination, a region of *pp1* downstream of the ATG start codon was

used to generate the construct. For the genotypic analyses, a diagnostic PCR

reaction was performed as outlined in **Fig. S1A**. Primer 1 (intP6tg) and primer 2

(ol492) were used to determine correct integration of the *gfp* sequence at the

targeted locus. For western blotting, purified gametocytes were lysed using lysis

buffer (10 mM TrisHCl pH 7.5, 150 mM NaCl, 0.5 mM EDTA and 1% NP-40). The

lysed samples were boiled for 10 min at 95 ºC after adding Laemmli buffer and were

centrifuged at maximum speed (13000g) for 5 min. The samples were

electrophoresed on a 4–12% SDS-polyacrylamide gel. Subsequently, resolved

proteins were transferred to nitrocellulose membrane (Amersham Biosciences).

Immunoblotting was performed using the Western Breeze Chemiluminescence Anti-

Rabbit kit (Invitrogen) and anti-GFP polyclonal antibody (Invitrogen) at a dilution of

1:1250, according to the manufacturer's instructions.

To study the function of PP1, we used two conditional knock down systems; a

promoter exchange/trap using double homologous recombination (PP1PTD) and an

auxin inducible degron (PP1AID) system. The PP1AID construct was derived from

the p277 plasmid, where the GFP sequence was excised following digestion with

*Age*I and *Not*I restriction enzymes and replaced with an AID/HA coding sequence.

The AID-HA sequence was PCR amplified (using primers: 5'-

CCCCAGACGTCGGATCCAATGATGGGCAGTGTCGAGCT-3' and 5'-

ATATAAGTAAGAAAAACGGCTTAAGCGTAATCTGGA-3') from the GW-AID/HA

plasmid (http://plasmogem.sanger.ac.uk/). Fragments were assembled following the

Gibson assembly protocol to generate the PP1-AID/HA transfection plasmid that was

transfected in the 615 line. Conditional degradation of PP1-AID/HA was performed

as described previously[57]. A schematic representation of the endogenous *pp1* locus

(PBANKA_1028300), the constructs and the recombined *pp1* locus can be found in

Fig S2A. A diagnostic PCR was performed for *pp1* gene knockdown parasites as

outlined in **Fig. S2A**. Primer 1 and Primer 3 were used to determine successful

integration of the targeting construct at the 3' gene locus **(Fig S2B)**. Primer 1 and

Primer 2 were used as controls **(Fig S2B)**.

The conditional knockdown construct PP1-PTD was derived from P_{ama1} (pSS368))

where *pp1* was placed under the control of the *ama1* promoter, as described

previously[58]. A schematic representation of the endogenous *pp1* locus, the

constructs and the recombined *pp1* locus can be found in **Fig S2D**. A diagnostic

PCR was performed for *pp1* gene knockdown parasites as outlined in **Fig. S2D**.

Primer 1 (5'-intPTD36) and Primer 2 (5'-intPTD) were used to determine successful

integration of the targeting construct at the 5' gene locus. Primer 3 (3'-intPTD) and

Primer 4 (3'-intPTama1) were used to determine successful integration for the 3' end

of the gene locus **(Fig. S2E)**. All the primer sequences can be found in **Table S3**. *P.*

berghei ANKA line 2.34 (for GFP-tagging) or ANKA line 507cl1 expressing GFP (for

the knockdown construct) parasites were transfected by electroporation[41].

Purification of schizonts and gametocytes

Blood cells obtained from infected mice (day 4 post infection) were cultured for 11h

and 24 h at 37°C (with rotation at 100 rpm) and schizonts were purified the following

day on a 60% v/v NycoDenz (in PBS) gradient, (NycoDenz stock solution: 27.6% w/v

NycoDenz in 5 mM Tris-HCl, pH 7.20, 3 mM KCl, 0.3 mM EDTA).

The purification of gametocytes was achieved using a protocol described

previously[59] with some modifications.   Briefly, parasites were injected into

phenylhydrazine treated mice and enriched by sulfadiazine treatment after 2 days of

infection. The blood was collected on day 4 after infection and gametocyte-infected

cells were purified on a 48% v/v NycoDenz (in PBS) gradient (NycoDenz stock

solution: 27.6% w/v NycoDenz in 5 mM Tris-HCl, pH 7.20, 3 mM KCl, 0.3 mM

EDTA). The gametocytes were harvested from the interface and activated.

Live cell imaging

To examine PP1-GFP expression during erythrocyte stages, parasites growing in

schizont culture medium were used for imaging at different stages (ring, trophozoite,

schizont and merozoite) of development. Purified gametocytes were examined for

GFP expression and localization at different time points (0, 1-15 min) after activation

in ookinete medium[37]. Zygote and ookinete stages were analyzed throughout 24 h of

culture. Images were captured using a 63x oil immersion objective on a Zeiss Axio

Imager M2 microscope fitted with an AxioCam ICc1 digital camera (Carl Zeiss, Inc).

Generation of dual tagged parasite lines

The PP1-GFP parasites were mixed with NDC80-cherry and MyoA-cherry parasites

in equal numbers and injected into mice. Mosquitoes were fed on mice 4 to 5 days

after infection when gametocyte parasitaemia was high. These mosquitoes were

checked for oocyst development and sporozoite formation at day 14 and day 21 after

feeding. Infected mosquitoes were then allowed to feed on naïve mice and after 4 - 5

days, and the mice were examined for blood stage parasitaemia by microscopy with

Giemsa-stained blood smears. In this way, some parasites expressed both PP1-GFP

and NDC80-cherry; and PP1-GFP and MyoA-cherry in the resultant gametocytes,

and these were purified and fluorescence microscopy images were collected as

described above.

Parasite phenotype analyses

Blood containing approximately 50,000 parasites of the PP1PTD line was injected intraperitoneally (i.p.) into mice to initiate infections. Asexual stages and gametocyte production were monitored by microscopy on Giemsa-stained thin smears. Four to five days post infection, exflagellation and ookinete conversion were examined as described previously [60] with a Zeiss AxioImager M2 microscope (Carl Zeiss, Inc) fitted with an AxioCam ICc1 digital camera. To analyse mosquito transmission, 30–50 *Anopheles stephensi* SD 500 mosquitoes were allowed to feed for 20 min on anaesthetized, infected mice with an asexual parasitaemia of 15% and a comparable number of gametocytes as determined on Giemsa-stained blood films. To assess mid-gut infection, approximately 15 guts were dissected from mosquitoes on day 14 post feeding, and oocysts were counted on an AxioCam ICc1 digital camera fitted to a Zeiss AxioImager M2 microscope using a 63x oil immersion objective. On day 21 post-feeding, another 20 mosquitoes were dissected, and their guts crushed in a loosely fitting homogenizer to release sporozoites, which were then quantified using a haemocytometer or used for imaging. Mosquito bite back experiments were performed 21 days post-feeding using naive mice, and blood smears were examined after 3-4 days.

Electron microscopy

Gametocytes activated for 6 min and 30 min were fixed in 4% glutaraldehyde in 0.1 M phosphate buffer and processed for electron microscopy as previously described[61]. Briefly, samples were post fixed in osmium tetroxide, treated *en bloc* with uranyl acetate, dehydrated and embedded in Spurr's epoxy resin. Thin sections

were stained with uranyl acetate and lead citrate prior to examination in a JEOL

JEM-1400 electron microscope (JEOL Ltd, UK)

Quantitative Real Time PCR (qRT-PCR) analyses

RNA was isolated from gametocytes using an RNA purification kit (Stratagene).

cDNA was synthesised using an RNA-to-cDNA kit (Applied Biosystems). Gene

expression was quantified from 80 ng of total RNA using a SYBR green fast master

mix kit (Applied Biosystems). All the primers were designed using the primer3

software (https://primer3.ut.ee/). Analysis was conducted using an Applied

Biosystems 7500 fast machine with the following cycling conditions: 95°C for 20 s

followed by 40 cycles of 95°C for 3 s; 60°C for 30 s. Three technical replicates and

three biological replicates were performed for each assayed gene. The *hsp70*

(PBANKA_081890) and *arginyl-t RNA synthetase* (PBANKA_143420) genes were

used as endogenous control reference genes. The primers used for qPCR can be

found in **Table S1.**

Transcriptome study using RNA-seq

For RNA extraction, parasite samples were passed through a plasmodipur column to

remove host DNA contamination prior to RNA isolation. Total RNA was extracted

from activated gametocytes and schizonts of WT-GFP and PP1PTD parasites (two

biological replicates each) using an RNeasy purification kit (Qiagen). RNA was

vacuum concentrated (SpeedVac) and transported using RNA-stable tubes

(Biomatrica). Strand-specific 354mRNA sequencing was performed on total RNA

and using TruSeq stranded mRNA sample prep 355kit LT (Illumina), as previously

described[62]. Libraries were sequenced using an Illumina Hiseq 4000 sequencing

platform with paired-end 150 bp read chemistry. The quality of the raw reads was

assessed   using   FATSQC

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc). Low-quality reads and

Illumina adaptor sequences from the read ends were removed using Trimmomatic

R[63]. Processed reads were mapped to the *P. beghei ANKA* reference genome

(release 40 in PlasmoDB - http://www.plasmoddb.org) using Hisat2[64] (V 2.1.0) with

parameter "—rna-strandness FR". Counts per feature were estimated using

FeatureCounts[65]. Raw read counts data were converted to counts per million (cpm)

and genes were excluded if they failed to achieve a cpm value of 1 in at least one of

the three replicates performed. Library sizes were scale-normalized by the TMM

method using EdgeR software[66] and further subjected to linear model analysis using

the voom function in the limma package[67]. Differential expression analysis was

performed using DeSeq2[68]. Genes with a fold-change greater than two and a false

discovery rate corrected p-value (Benjamini-Hochberg procedure) < 0.05 were

considered to be differentially expressed. Functional groups shown in Figure 6C

were   inferred   from   annotations   available   in   PlasmoDB: Release   49

(https://plasmodb.org/plasmo/app).

Immunoprecipitation and Mass Spectrometry

Schizonts, following 11 hours and 24 hours, respectively in *in vitro* culture, and

male gametocytes 11 min post activation were used to prepare cell lysates.

Purified parasite pellets were crosslinked using formaldehyde (10 min incubation

with 1% formaldehyde, followed by 5 min incubation in 0.125M glycine solution

and   3   washes   with   phosphate   buffered   saline   (PBS)   (pH,   7.5).

Immunoprecipitation was performed using crosslinked protein and a GFP-Trap[®]_A

Kit (Chromotek) following the manufacturer's instructions. Proteins bound to the

GFP-Trap[®]_A beads were digested using trypsin and the peptides were analysed

by LC-MS/MS.   Briefly, to prepare samples for LC-MS/MS, wash buffer was

removed, and ammonium bicarbonate (ABC) was added to beads at room

temperature. We added 10 mM TCEP (Tris-(2-carboxyethyl) phosphine

hydrochloride) and 40 mM 2-chloroacetamide (CAA) and incubation was performed

for 5 min at 70° C. Samples were digested using 1 µg Trypsin per 100 µg protein at

room temperature overnight followed by 1% TFA addition to bring the pH into the

range of 3-4 before mass spectrometry.

Statistical analysis

All statistical analyses were performed using GraphPad Prism 8 (GraphPad

Software). An unpaired t-test and two-way anova test were used to examine

significant differences between wild-type and mutant strains for qRT-PCR and

phenotypic analysis accordingly.

Data Availability. RNA Sequence reads have been deposited in the NCBI

Sequence gene expression omnibus with the accession number GSE164175.

"The mass spectrometry proteomics data have been deposited to the

ProteomeXchange Consortium with the dataset identifier PXD023571 and

10.6019/PXD023571.

Author contributions.

RT and MZ conceived and designed all experiments. RT, MZ, RP, DB, DSG, GK

performed the GFP tagging and conditional knockdown with promoter trap

experiments. RR and MBr generated and characterised the PP1-AID/HA line. MZ,

RP, GK, DB and RT performed protein pull-down experiments. ARB performed mass

spectrometry. AS, RN and AP performed RNA sequencing (RNA-seq). DJPF and

SV performed electron microscopy. RT, AAH, MZ, DSG, AS, RN, AP and DJPF

analyzed the data. MZ, DSG and RT wrote the original draft. RT, AAH, AP, DJPF,

MB edited and reviewed the manuscript and all other contributed to it.

Acknowledgments. We thank Julie Rodgers for helping to maintain the insectary

and other technical works, and the personnel at the Bioscience Core Laboratory

(BCL) in KAUST for sequencing the RNA samples and producing the raw datasets.

Funding

This work was supported by: MRC UK (G0900278, MR/K011782/1, MR/N023048/1)

and BBSRC (BB/N017609/1) to RT and MZ; the Francis Crick Institute (FC001097),

the Cancer Research UK (FC001097), the UK Medical Research Council

(FC001097), and the Wellcome Trust (FC001097) to AAH; the Swiss National

Science Foundation project grant 31003A_179321 to MBr. MBr is an INSERM and

EMBO young investigator; a faculty baseline fund (BAS/1/1020-01-01) and a

Competitive Research Grant (CRG) award from OSR (OSR-2018-CRG6-3392) from

the King Abdullah University of Science and Technology to AP.

Figure legends

Fig. 1. Location of PP1 during asexual blood stage schizogony and its

association with kinetochore (Ndc80) and glideosome (MyosinA)

(A) Live cell imaging of PP1-GFP (Green) showing its location during different stages

of intraerythrocytic development and in free merozoites.   DIC: Differential

interference contrast; Hoechst: stained DNA (blue); Merge: green and blue images

merged. A schematic guide showing the locations of PP1GFP foci during

segmentation of merozoites is depicted in right hand panel. **(B)** Live cell imaging

showing location of PP1–GFP (green) in relation to the kinetochore marker NDC80-

mCherry (red) and DNA (Hoechst, blue). Merge: green, red and blue images

merged. A schematic guide showing the locations of PP1GFP foci with NDC80-

mCherry during segmentation of merozoites is depicted in right hand panel. **(C)** Live

imaging showing the location of PP1–GFP (green) in relation to inner membrane

complex marker MyoA-mCherry (red) and DNA (Hoechst, blue) during different

stages of intraerythrocytic development and in extracellular merozoites. A schematic

guide showing the locations of PP1GFP foci with NDC80-mCherry and MyoA-

mCherry during segmentation of merozoites is depicted in the three right-hand

panels. Merge: green, red and blue images merged. Sch-E (Early schizont), Sch-M

(Middle schizont), Sch-L (Late schizont) Sch-S (Segmented schizont). In all panels,

scale bar = 5 µm.

Fig. 2. The location of PP1 and its association with the kinetochore during

chromosome segregation in male gametogony.

(A) Live-cell imaging of PP1-GFP during male gametogony showing an initial

diffused localization before activation and focal points after activation in the later

stages (shown as minutes post activation). Panels are DIC, Hoechst (blue, DNA),

PP1-GFP (green) and Merge (green and blue channels). **(B)** Live-cell imaging of

parasite line expressing both PP1-GFP and NDC80-mCherry showing location of

PP1 (green) and NDC80 (red) in male gametocytes at different time points after

activation**.** A schematic guide showing the locations of PP1GFP foci with DNA and

NDC80-mCherry during male gametogony is depicted in the right panel. Merge/DNA

is green, red and blue (Hoechst, DNA) channels. In both panels, scale bar = 5 µm.

Fig. 3. PP1-GFP localization during zygote formation, ookinete development,

and sporogony inside the mosquito gut

(A) Live cell imaging showing PP1-GFP location in male and female gametes,

zygote and during ookinete development (stages I to V and mature ookinete). A cy3-

conjugated antibody, 13.1, which recognises the P28 protein on the surface of

zygotes and ookinetes, was used to mark these stages. Panels: DIC (differential

interference contrast), PP1-GFP (green, GFP), 13.1 (red), Merged: Hoechst (blue,

DNA), PP1-GFP (green, GFP) and P28 (red). Scale bar = 5 µm. Insets show, at

higher magnification, the PP1-GFP signal on the zygote and developing apical end

of early stage 'retorts', and in the nucleus of late retorts and ookinete stage. **(B)** Live

cell imaging of PP1-GFP (green) in relation to NDC80-mCherry (red) and Hoechst

staining (blue, DNA) in zygote and ookinete stages. **(C)** Live cell imaging of PP1-

GFP in developing oocysts in mosquito guts at 7-, 14- and 21-days post-infection

and in a sporozoite. Panels: DIC, Hoechst (blue, DNA), PP1-GFP (green), Merged

(blue and green channels). **(D)** Live cell imaging of PP1-GFP in relation to NDC80 in

developing oocysts and in a sporozoite. Panels: DIC (differential interference

contrast), PP1-GFP (green), NDC80-mCherry (red), Merge (Hoechst, blue, DNA;

PP1-GFP, green; NDC80-mCherry, red). Scale bar = 5 µm

Fig. 4. PP1 has an important role during male gamete formation and zygote -

ookinete development (A) qRT-PCR analysis of *pp1* transcription in PP1PTD and

WT-GFP parasites, showing the downregulation of *pp1*. Each bar is the mean of

three biological replicates ± SD. **(B)** Male gametogony (exflagellation) of PP1PTD

line (black bar) and WT-GFP line (white bar) measured as the number of

exflagellation centres per field. Mean ± SD; n=3 independent experiments. **(C)**

Ookinete conversion as a percentage for PP1PTD and WT-GFP parasites.

Ookinetes were identified using P28 antibody as a surface marker and defined as

those cells that differentiated successfully into elongated 'banana shaped' ookinetes.

Round cells show zygotes that did not start to transform and 'retorts' could not

differentiate successfully in ookinetes. Mean ± SD; n=3 independent experiments.

(D) Representative images of round cells, retorts and fully differentiated ookinetes.

(E) Total number of GFP-positive oocysts per infected mosquito in PP1PTD and WT-

GFP parasites at 7-, 14- and 21-days post infection (dpi). Mean ± SD; n=3

independent experiments. **(F)** Representative images of mosquito midguts on day 14

showing them full of oocysts in WTGFP and no oocyst in PP1PTD **(G)** Total number

of sporozoites in oocysts of PP1PTD and WT-GFP parasites at 14 and 21 dpi. Mean

± SD; n=3 independent experiments. **(H)** Total number of sporozoites in salivary

glands of PP1PTD and WT-GFP parasites. Bar diagram shows mean ± SD; n=3

independent experiments. Unpaired t-test was performed for statistical analysis.

*p<0.05, **p<0.01, ***p<0.001.

Figure 5: Ultrastructure analysis of PP1PTD gametocytes shows defects in

nuclear pole and axoneme assembly during male gametogony

Electron micrographs of PP1PTD (a-f) and WTGFP (g-l) male gametocytes at 6 min

(a-c, g-i) and 30 min (d-f, j-l) post activation. Bars represent 1 μm (a, d, g, j) and 100

nm in all other micrographs. **(a)** Low power of mutant showing the central nucleus

(N) with a nuclear pole and associated basal body (enclosed). **(b)** Enlargement of

the enclosed area showing the nuclear pole (NP) with adjacent basal body (B) and

associated axoneme (A). **(c)** Cross section through an axoneme showing the 9+2

arrangement of microtubules. **(d)** Low power of a mutant at 30 min showing the

central nucleus (N) with a nuclear pole and an increased number of axoneme

profiles (A) within the cytoplasm. **(e)** Detail of the periphery of a nucleus showing the

nuclear pole (NP), basal body (B) and associated axoneme (A). **(f)** Cross section of

an axoneme showing the 9+2 arrangement. **(g)** Low power of a WT

microgametocyte illustrating 2 nuclear poles (NP) associated with the nucleus (N).

(h) Enlargement of the enclosed area showing the nuclear pole (NP) with adjacent

basal body (B) and associated axoneme (A). **(i)** Cross section of an axoneme

showing the 9+2 arrangement. **(j)** Low power of a late microgametocyte showing the

nucleus with areas of condensed chromatin. Note the cross sectioned free

microgametes (Mg). A – axoneme. **(k)** Periphery of a microgametocyte undergoing

exflagellation with the nucleus (N) associated with flagellum (F) protruding from the

surface. **(l)** Cross section of a free microgamete showing the 9+2 flagellum (F) and

electron dense nucleus (N). These observations are based on random sections

through 30 microgametocytes.

Figure 6: Transcriptome of PP1PTD mutant reveals important roles of PP1 in

parasite cell cycle, motor protein function, and cell polarity during gametocyte

biology

Volcano plots showing significantly down-regulated (blue Log_2 fold ≤ -1, q value <

0.05) and up-regulated genes (brown Log_2 fold ≥ 1, q value < 0.05) in PP1PTD

compared to wild-type lines in non (0 min)-activated gametocytes **(A)** and 30 min

activated gametocytes **(B).** Non-differentially regulated genes are represented as

black dots. **(C)** Expression heat maps showing affected genes from specific

functional classes of proteins such as kinases, phosphatases, motor proteins and

proteins associated with parasite motility, entry into the host cell, ookinete surface

and cell cycle. Genes are ordered based on their differential expression pattern in

non-activated gametocytes. **(D)** Validation by qRT-PCR of a few genes randomly

selected based on the RNA-seq data. All the experiments were performed three

times in duplicate with two biological replicates. *p ≤05.

Fig. 7. Interacting partners of PP1 during asexual schizont and sexual

gametocyte stages

(A) List of proteins interacting with PP1 during schizont and gametocyte stages. **(B)**

Venn diagram showing common interacting partners in schizonts and gametocytes

with some additional proteins specific to gametocytes.

Supplementary figures

Fig. S1. Generation and genotypic analysis of PP1GFP parasites

(A) Schematic representation for 3'-tagging of *pp1* gene with green fluorescent

protein (GFP) sequence via single homologous recombination. **(B)** Integration PCR

showing correct integration of tagging construct. **(C)** Western blot showing expected

size of PP1-GFP protein.

Fig. S2. Generation and genotype analysis of conditional knockdown

PP1 parasites

(A) Schematic representation of auxin inducible degron (AID) strategy to generate

PP1AID parasites. **(B)** Integration PCR of the PP1AID construct in the *pp1* locus.

Primer 1 and Primer 2 were used for control PCR while primer 1 and primer 3 were

used to determine successful integration of AID-HA sequence and selectable marker

at 3'-end of *pp1* locus **(C)** PP1AID protein expression level as measured by western

blotting upon addition of auxin to mature purified gametocytes; α-tubulin serves as a

loading control. Auxin treatment of PP1AID showed no defect in exflagellation (error

bars show standard deviation from the mean; technical replicates from three

independent infections. **(D)** Schematic representation of the promoter swap strategy

(PP1PTD, placing *pp1* under the control of the *ama1* promoter) by double

homologous recombination. Arrows 1 and 2 indicate the primer positions used to

confirm 5' integration and arrows 3 and 4 indicate the primers used for 3' integration.

(E) Integration PCR of the promotor swap construct into the *pp1* locus. Primer 1 (5'-

IntPTD36) with primer 2 (5'-IntPTD) were used to determine successful integration of

the selectable marker. Primer 3 (3'-intPTama1) and primer 4 (3'-IntPTD36) were

used to determine the successful integration of ama1 promoter. Primer 1 (5'-

IntPTD36) and primer 4 (3'-IntPTD36) were used to show complete knock-in of the

construct and the absence of a band at 2.1 kb (endogenous) resulting in complete

knock-in of the construct. **(F)** Parasitaemia during blood stage schizogony showing a

significant slow growth of PP1PTD compared to WTGFP parasites. Experiment was

done with three mice each with 1000 parasites per mice injected intraperitonially.

***$P<0.001$ **(G)** Bite back experiments show no transmission of PP1PTD parasites

(black bar) from mosquito to mouse, while successful transmission was shown by

WT-GFP parasites. Mean ± SD; n= 3 independent experiments.

Fig. S3. RNA seq analysis:

(A) Clustered dendrogram of two biological replicates of WTGFP and PP1PTD

mutant parasite lines during gametocyte stage using hierarchical clustering

algorithm. Analysis was performed on normalized count data**. (B)** RNA-seq read

statistics.

Supplementary Tables

Table S1. Differentially expressed genes in PP1PTD parasites

Table S2. List of proteins pulled down by PP1GFP

Table S3. Primers used in this study

References

1.   Marston AL, Amon A. Meiosis: cell-cycle controls shuffle and deal. *Nat Rev*

Mol Cell Biol **5**, 983-997 (2004).

2.   Sullivan M, Morgan DO. Finishing mitosis, one step at a time. *Nat Rev Mol*

Cell Biol **8**, 894-903 (2007).

3.   Novak B, Kapuy O, Domingo-Sananes MR, Tyson JJ. Regulated protein

kinases and phosphatases in cell cycle decisions. *Curr Opin Cell Biol* **22**, 801-

808 (2010).

4.   Afshar K, Werner ME, Tse YC, Glotzer M, Gonczy P. Regulation of cortical

contractility and spindle positioning by the protein phosphatase 6 PPH-6 in

one-cell stage C. elegans embryos. *Development* **137**, 237-247 (2010).

5.   Boutros R, Dozier C, Ducommun B. The when and wheres of CDC25 phosphatases. *Curr Opin Cell Biol* **18**, 185-191 (2006).

6.   Chen F*, et al.* Multiple protein phosphatases are required for mitosis in Drosophila. *Curr Biol* **17**, 293-303 (2007).

7.   Stegmeier F, Amon A. Closing mitosis: the functions of the Cdc14 phosphatase and its regulation. *Annu Rev Genet* **38**, 203-232 (2004).

8.   Bollen M, Gerlich DW, Lesage B. Mitotic phosphatases: from entry guards to exit guides. *Trends Cell Biol* **19**, 531-541 (2009).

9.   Heroes E, Lesage B, Gornemann J, Beullens M, Van Meervelt L, Bollen M. The PP1 binding code: a molecular-lego strategy that governs specificity. *FEBS J* **280**, 584-595 (2013).

10.   Holder J, Poser E, Barr FA. Getting out of mitosis: spatial and temporal control of mitotic exit and cytokinesis by PP1 and PP2A. *FEBS Lett* **593**, 2908-2924 (2019).

11.   Nilsson J. Protein phosphatases in the regulation of mitosis. *J Cell Biol* **218**, 395-409 (2019).

860   12.   Hattersley N*, et al.* A Nucleoporin Docks Protein Phosphatase 1 to Direct

861   Meiotic Chromosome Segregation and Nuclear Assembly. *Dev Cell* **38**, 463-

862   477 (2016).

864   13.   Cheeseman IM. The kinetochore. *Cold Spring Harb Perspect Biol* **6**, a015826

865   (2014).

867   14.   Nagpal H, Fukagawa T. Kinetochore assembly and function through the cell

868   cycle. *Chromosoma* **125**, 645-659 (2016).

870   15.   Varma D, Salmon ED. The KMN protein network--chief conductors of the

871   kinetochore orchestra. *J Cell Sci* **125**, 5927-5936 (2012).

873   16.   Lampert F, Westermann S. A blueprint for kinetochores - new insights into the

874   molecular mechanics of cell division. *Nat Rev Mol Cell Biol* **12**, 407-412

875   (2011).

877   17.   Wu JQ*, et al.* PP1-mediated dephosphorylation of phosphoproteins at mitotic

878   exit is controlled by inhibitor-1 and PP1 phosphorylation. *Nat Cell Biol* **11**,

879   644-651 (2009).

881   18.   Dohadwala M*, et al.* Phosphorylation and inactivation of protein phosphatase

882   1 by cyclin-dependent kinases. *Proc Natl Acad Sci U S A* **91**, 6408-6412

883   (1994).

885   19.   Bouchoux C, Uhlmann F. A quantitative model for ordered Cdk substrate

886   dephosphorylation during mitotic exit. *Cell* **147**, 803-814 (2011).

888   20.   Grallert A*, et al.* A PP1-PP2A phosphatase relay controls mitotic progression.

889   *Nature* **517**, 94-98 (2015).

891   21.   Mochida S, Hunt T. Protein phosphatases and their regulation in the control of

892   mitosis. *EMBO Rep* **13**, 197-203 (2012).

894   22.   Howick VM*, et al.* The Malaria Cell Atlas: Single parasite transcriptomes

895   across the complete Plasmodium life cycle. *Science* **365**, (2019).

897   23.   Guttery David S*, et al.* Genome-wide Functional Analysis of Plasmodium

898   Protein Phosphatases Reveals Key Regulators of Parasite Development and

899   Differentiation. *Cell Host & Microbe* **16**, 128-140 (2014).

901   24.   Bhattacharyya MK, Hong Z, Kongkasuriyachai D, Kumar N. Plasmodium

902   falciparum protein phosphatase type 1 functionally complements a glc7

903   mutant in Saccharomyces cerevisiae. *Int J Parasitol* **32**, 739-747 (2002).

25. Zhang M*, et al.* Uncovering the essential genes of the human malaria parasite Plasmodium falciparum by saturation mutagenesis. *Science* **360**, (2018).

26. Bushell E*, et al.* Functional Profiling of a Plasmodium Genome Reveals an Abundance of Essential Genes. *Cell* **170**, 260-272 e268 (2017).

27. Yokoyama D, Saito-Ito A, Asao N, Tanabe K, Yamamoto M, Matsumura T. Modulation of the growth of Plasmodium falciparum in vitro by protein serine/threonine phosphatase inhibitors. *Biochem Biophys Res Commun* **247**, 18-23 (1998).

28. Khalife J, Freville A, Gnangnon B, Pierrot C. The Multifaceted Role of Protein Phosphatase 1 in Plasmodium. *Trends Parasitol*, (2020).

29. Paul AS*, et al.* Co-option of Plasmodium falciparum PP1 for egress from host erythrocytes. *Nat Commun* **11**, 3532 (2020).

30. Fardilha M, Esteves SL, Korrodi-Gregorio L, da Cruz e Silva OA, da Cruz e Silva FF. The physiological relevance of protein phosphatase 1 and its interacting proteins to health and disease. *Curr Med Chem* **17**, 3996-4017 (2010).

927   31.   Hollin T*, et al.* Essential role of GEXP15, a specific Protein Phosphatase type

928   1 partner, in Plasmodium berghei in asexual erythrocytic proliferation and

929   transmission. *PLoS Pathog* **15**, e1007973 (2019).

931   32.   Gerald N, Mahajan B, Kumar S. Mitosis in the human malaria parasite

932   Plasmodium falciparum. *Eukaryot Cell* **10**, 474-482 (2011).

934   33.   Guttery DS, Roques M, Holder AA, Tewari R. Commit and Transmit:

935   Molecular Players in Plasmodium Sexual Development and Zygote

936   Differentiation. *Trends Parasitol* **31**, 676-685 (2015).

938   34.   Matthews H, Duffy CW, Merrick CJ. Checks and balances? DNA replication

939   and the cell cycle in Plasmodium. *Parasit Vectors* **11**, 216 (2018).

941   35.   Zeeshan M*, et al.* Real-time dynamics of Plasmodium NDC80 reveals unusual

942   modes of chromosome segregation during parasite proliferation. *J Cell Sci*

943   **134**, (2020).

945   36.   Arnot DE, Ronander E, Bengtsson DC. The progression of the intra-

946   erythrocytic cell cycle of Plasmodium falciparum and the role of the centriolar

947   plaques in asynchronous mitotic division during schizogony. *Int J Parasitol* **41**,

948   71-80 (2011).

950   37.   Zeeshan M*, et al.* Plasmodium kinesin-8X associates with mitotic spindles and

951   is essential for oocyst development during parasite proliferation and

952   transmission. *PLoS Pathog* **15**, e1008048 (2019).

954   38.   Vaughan JA. Population dynamics of Plasmodium sporogony. *Trends*

955   *Parasitol* **23**, 63-70 (2007).

957   39.   Pandey R*, et al.* Plasmodium Condensin Core Subunits SMC2/SMC4 Mediate

958   Atypical Mitosis and Are Essential for Parasite Proliferation and Transmission.

959   *Cell Rep* **30**, 1883-1897 e1886 (2020).

961   40.   Wall RJ*, et al.* Plasmodium APC3 mediates chromosome condensation and

962   cytokinesis during atypical mitosis in male gametogenesis. *Sci Rep* **8**, 5610

963   (2018).

965   41.   Janse CJ*, et al.* High efficiency transfection of Plasmodium berghei facilitates

966   novel selection procedures. *Molecular and biochemical parasitology* **145**, 60-

967   70 (2006).

969   42.   Hollin T, De Witte C, Lenne A, Pierrot C, Khalife J. Analysis of the interactome

970   of the Ser/Thr Protein Phosphatase type 1 in Plasmodium falciparum. *BMC*

971   *Genomics* **17**, 246 (2016).

43.   Straschil U*, et al.* The Armadillo repeat protein PF16 is essential for flagellar

structure and function in Plasmodium male gametes. *PloS one* **5**, e12901

(2010).

44.   Gelens L, Qian J, Bollen M, Saurin AT. The Importance of Kinase-

Phosphatase Integration: Lessons from Mitosis. *Trends Cell Biol* **28**, 6-21

(2018).

45.   Nasa I, Rusin SF, Kettenbach AN, Moorhead GB. Aurora B opposes PP1

function in mitosis by phosphorylating the conserved PP1-binding RVxF motif

in PP1 regulatory proteins. *Sci Signal* **11**, (2018).

46.   Tewari R*, et al.* The Systematic Functional Analysis of Plasmodium Protein

Kinases Identifies Essential Regulators of Mosquito Transmission. *Cell Host &*

Microbe **8**, 377-387 (2010).

47.   Ceulemans H, Bollen M. Functional diversity of protein phosphatase-1, a

cellular economizer and reset button. *Physiol Rev* **84**, 1-39 (2004).

48.   Lesage B, Qian J, Bollen M. Spindle checkpoint silencing: PP1 tips the

balance. *Curr Biol* **21**, R898-903 (2011).

49.　Peggie MW, MacKelvie SH, Bloecher A, Knatko EV, Tatchell K, Stark MJ.

Essential functions of Sds22p in chromosome stability and nuclear localization

of PP1. *J Cell Sci* **115**, 195-206 (2002).

50.　Rodrigues NT, Lekomtsev S, Jananji S, Kriston-Vizi J, Hickson GR, Baum B.

Kinetochore-localized PP1-Sds22 couples chromosome segregation to polar

relaxation. *Nature* **524**, 489-492 (2015).

1002

1003　51.　Verbinnen I, Ferreira M, Bollen M. Biogenesis and activity regulation of protein

1004　　　phosphatase 1. *Biochem Soc Trans* **45**, 89-99 (2017).

1005

1006　52.　Sinden RE. Mitosis and meiosis in malarial parasites. *Acta Leidensia* **60**, 19-

1007　　　27 (1991).

1008

1009　53.　Solyakov L*, et al.* Global kinomic and phospho-proteomic analyses of the

1010　　　human malaria parasite Plasmodium falciparum. *Nat Commun* **2**, 565 (2011).

1011

1012　54.　Balestra AC*, et al.* A divergent cyclin/cyclin-dependent kinase complex

1013　　　controls the atypical replication of a malaria parasite during gametogony and

1014　　　transmission. *Elife* **9**, (2020).

1015

1016　55.　Invergo BM, Brochet M, Yu L, Choudhary J, Beltrao P, Billker O. Sub-minute

1017　　　Phosphoregulation of Cell Cycle Systems during Plasmodium Gamete

1018　　　Formation. *Cell Rep* **21**, 2017-2029 (2017).

1019

1020 56. Moreira S, Osswald M, Ventura G, Goncalves M, Sunkel CE, Morais-de-Sa E.

1021   PP1-Mediated Dephosphorylation of Lgl Controls Apical-basal Polarity. *Cell*

1022   *Rep* **26**, 293-301 e297 (2019).

1023

1024 57. Philip N, Waters AP. Conditional Degradation of Plasmodium Calcineurin

1025   Reveals Functions in Parasite Colonization of both Host and Vector. *Cell Host*

1026   *Microbe* **18**, 122-131 (2015).

1027

1028 58. Sebastian S*, et al.* A Plasmodium calcium-dependent protein kinase controls

1029   zygote development and transmission by translationally activating repressed

1030   mRNAs. *Cell Host Microbe* **12**, 9-19 (2012).

1031

1032 59. Beetsma AL, van de Wiel TJ, Sauerwein RW, Eling WM. Plasmodium berghei

1033   ANKA: purification of large numbers of infectious gametocytes. *Exp Parasitol*

1034   **88**, 69-72 (1998).

1035

1036 60. Guttery DS*, et al.* A putative homologue of CDC20/CDH1 in the malaria

1037   parasite is essential for male gamete development. *PLoS Pathog* **8**, e1002554

1038   (2012).

1039

1040 61. Ferguson DJ*, et al.* Maternal inheritance and stage-specific variation of the

1041   apicoplast in Toxoplasma gondii during development in the intermediate and

1042   definitive host. *Eukaryot Cell* **4**, 814-826 (2005).

1043

1044   62.   Roques M*, et al.* Plasmodium centrin PbCEN-4 localizes to the putative

1045   MTOC and is dispensable for malaria parasite proliferation. *Biol Open* **8**,

1046   (2019).

1047

1048   63.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina

1049   sequence data. *Bioinformatics* **30**, 2114-2120 (2014).

1050

1051   64.   Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low

1052   memory requirements. *Nature methods* **12**, 357-360 (2015).

1053

1054   65.   Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose

1055   program for assigning sequence reads to genomic features. *Bioinformatics*

1056   **30**, 923-930 (2014).

1057

1058   66.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for

1059   differential expression analysis of digital gene expression data. *Bioinformatics*

1060   **26**, 139-140 (2010).

1061

1062   67.   Ritchie ME*, et al.* limma powers differential expression analyses for RNA-

1063   sequencing and microarray studies. *Nucleic acids research* **43**, e47 (2015).

1064

1065   68.   Love MI, Huber W, Anders S. Moderated estimation of fold change and

1066   dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

1067

1068

Fig 1

Fig 2

Fig 3

Fig 4

Fig 5

Fig 6

Fig 7

A

P. berghei Gene ID	Product Description	Gene names	MW (kDa)	Schizonts- 24 h				Schizonts-10 h				Gametocytes-10 min			
				PP1-GFP			GFP	PP1-GFP			GFP	PP1-GFP			GFP
PBANKA_1028300	serine/threonine protein phosphatase PP1	PP1	34963	31	26	23	0	37	26	25	0	23	33	26	0
PBANKA_1218500	protein phosphatase inhibitor 2, putative	PPI2	16664	11	7	9	0	9	8	7	0	10	9	9	0
PBANKA_0516600	leucine-rich repeat protein	LRR1	36735	33	26	26	0	22	29	29	0	11	22	25	0
PBANKA_0601600	conserved protein, unknown function	NEMF-like	201925	36	17	35	0	21	37	21	0	12	22	24	0
PBANKA_0515500	protein phosphatase inhibitor 3, putative	PPI3	13065	3	3	3	0	0	4	2	0	2	2	3	0
PBANKA_0515400	conserved Plasmodium protein	CD2BP2	77377	17	10	18	0	7	22	12	0	3	2	9	0
PBANKA_0310800	RTR1 domain-containing protein, putative	RTR1-like	97344	15	8	11	0	19	32	19	0	0	0	7	0
PBANKA_1458800	kinesin, putative	kinesin-15	164999	0	0	0	0	0	0	0	0	2	7	2	0
PBANKA_0202700	kinesin-8, putative	kinesin-8B	168815	0	0	0	0	0	0	0	0	2	4	2	0
PBANKA_1458300	kinesin-13, putative	kinesin-13	117181	0	0	0	0	0	0	0	0	0	3	3	0
PBANKA_0917400	armadillo repeat protein PF16	PF16	57748	0	0	0	0	0	0	0	0	0	2	2	0

B

Fig S1

A

Wild-type genomic locus

C-terminal GFP fusion tag construct

Endogenous GFP tagged locus

B

Integration-PCR   Control-PCR

WT   PP1-GFP   WT   PP1-GFP

Kb

10 -

4 -

1 -

0.6 -

0.2 -

← 1.2
← 1.0

C

WTGFP   P6GFP

kDa M

190 —

80 —

25 —

← 62 kDa
← 29 kDa

Fig S2

Fig S3

A

Activated gametocytes 0 min

Activated gametocytes 30 min

B

	Wild type (in millions)		PP1PTD (in millions)	
Gametocytes	Replicate 1	Replicate 2	Replicate 1	Replicate 2
0 min	45.99	25.5	58.6	19.8
30 min	22.57	31.57	16.33	20.65