Conditions under which distributions of edge length ratios on phylogenetic trees can be used to order evolutionary events

EDWARD SUSKO[1,2], MIKE STEEL[3] AND ANDREW J. ROGER[1,4]

[1] *Centre for Comparative Genomics and Evolutionary Bioinformatics, Dalhousie University*
[2] *Department of Mathematics and Statistics, Dalhousie University,*
Halifax, Nova Scotia, Canada B3H 4R2
[3] *Biomathematics Research Centre, University of Canterbury, Christchurch, New Zealand 8041*
[4] *Department of Biochemistry and Molecular Biology, Dalhousie University,*
Halifax, Nova Scotia, Canada B3H 4R2

Corresponding Author: Edward Susko, Department of Mathematics and Statistics, Dalhousie University, Halifax, Nova Scotia, Canada B3H 3J5; Phone: (902) 494-8865; Fax: (902) 494-5130; E-mail: edward.susko@gmail.com

Abstract

Two recent high profile studies have attempted to use edge (branch) length ratios from large sets of phylogenetic trees to determine the relative ages of genes of different origins in the evolution of eukaryotic cells. This approach can be straightforwardly justified if substitution rates are constant over the tree for a given protein. However, such strict molecular clock assumptions are not expected to hold on the billion-year timescale. Here we propose an alternative set of conditions under which comparisons of edge length distributions from multiple sets of phylogenies of proteins with different origins can be validly used to discern the order of their origins. We also point out scenarios where these conditions are not expected to hold and caution is warranted.

Main

The origin of eukaryotic cells from prokaryotic precursors - eukaryogenesis - remains one of the more mysterious major evolutionary transitions in the history of life on Earth. This transition involved a host cellular lineage related to asgard Archaea (Eme *et al.* 2017) that, at some point prior to the last eukaryotic common ancestor (LECA), took up an endosymbiotic alphaproteobacterium that became the mitochondrion, an integrated energy-producing organelle within eukaryotic cells (Dacks *et al.* 2016; Roger *et al.* 2017; Porter 2020). Genes in LECA, therefore, have multiple possible origins: either they were inherited from the host lineage, acquired from the mitochondrial symbiont by endosymbiotic gene transfer, transferred from potentially many other prokaryotic donors by lateral gene transfer (Rochette *et al.* 2014; Pittis and Gabaldón 2016a), or arose *de novo* during eukaryogenesis. Regardless of their origin, many genes were extensively duplicated during this period, as many new cellular traits including the cytoskeleton, nucleus, endomembrane system evolved

in the proto-eukaryote lineage. Determining the order of these events remains a major

roadblock in our understanding of eukaryogenesis.

In 2016, Pittis and Gabaldón introduced a novel approach to approximating the relative

ages of genes of different origins that were acquired during eukaryogenesis (Pittis and Ga-

baldón 2016a). The approach relies on the notion that edge lengths on phylogenetic trees

estimated from aligned genes or proteins, represent expected numbers of amino acid substi-

tutions along the edge and are proportional to the product of rates of substitution along the

edge and the time span of the edge. Under a strict molecular clock assumption the relative

lengths of edges are proportional to time spans of the edges.

To characterize the relative timespan a gene has been resident in the proto-eukaryote

lineage and LECA, Pittis and Gabaldón focused on the edge in each gene tree between the

LECA node and the node representing the common ancestor of the closest prokaryotic sister

group and the eukaryote lineage (Fig. 1), an edge they call the stem the length of which is

denoted here as L_s. All genes from the same origin, O, are expected to have a stem edge

that corresponds to the same time span (T_{s*}^g is constant for $g \in O$). A serious complication

arises here almost immediately. For genes in the proto-eukaryote genome that were inherited

from its common ancestor with the closest sampled asgard archaeon, the time span of the

stem edge in the protein tree T_s^g is the same as the timespan it has been resident during

eukaryogenesis, T_{s*}^g. However, for genes that were laterally acquired during eukaryogenesis

either *via* the mitochondrial symbiont or from other prokaryotic sources, T_s^g is expected

to be larger than T_{s*}^g (Fig. 1). This is because the sampled taxa are unlikely to include

representatives from the actual immediate prokaryotic sister group of the donor lineage of

the gene(s). Reasons for this include inadequacy in sampling of living prokaryote lineages

but, more likely, it is because the actual sister group, as opposed to the sampled one, went

extinct. In what follows, we assume that for all comparisons $T_s^g \approx T_{s*}^g$, but it is important to

recognize the caveats accompanying conclusions coming from stem-length methods applied

to comparisons amongst acquired genes. For instance, a claim that the time of acquisition

of a group B is earlier than that of an acquired group A, is more directly an inference that

the closest sampled sister lineage of group B diverged earlier than that of group A.

Figure 1: **The correspondence between edges on the phylogeny of a gene acquired during eukaryogenesis and edges on the geological time tree of life.** The top tree (blue box) shows the part of the geological tree of life depicting the closest asgard archaeal sister group relationship with the 'host' lineage of eukaryotes. FECA represents the *first eukaryotic common ancestor* and LECA is the *last eukaryotic common ancestor*. At some point along the eukaryogenesis edge between FECA and LECA, gene g was acquired by the proto-eukaryote genome from a prokaryotic lineage. The timespan that gene g was present during eukaryogenesis was T_{s*}^g and from LECA to the present is T_e. Below (green box) is the estimated phylogeny of protein g and its orthologs. The length of the stem edge is L_s^g and corresponds to timespan T_s^g in the time tree. The length of the segment of the latter edge post-acquisition by the proto-eukaryote lineage (green diamond to purple circle) is L_{s*}^g and corresponds to timespan T_{s*}^g. Note that T_s^g, is an upper bound on the timespan of the desired stem edge post-acquisition, T_{s*}^g, because any gene transfer to the proto-eukaryote from a prokaryotic source must have occurred after speciation of the donor lineage from its closest sampled extant sister group. This discrepancy between the time of origin of a gene and timespan of its stem edge only occurs for genes acquired during eukaryogenesis (e.g. mitochondrial and other laterally acquired genes). The median length of all possible paths between the LECA node (purple circle) and eukaryote leaf node is med(L_e^g). The normalized stem length, $sl^g = L_s^g / \text{med}(L_e^g)$.

⁵⁸   To remove the effect of different overall rates of evolution in different genes, R_g, Pittis and
⁵⁹ Gabaldón normalize the original (raw) stem edge length, by a path length, L_e, for a path,
⁶⁰ e (e for eukaryote), that corresponds to a constant time span over genes, denoted T_e, giving
⁶¹ the time from LECA to the present (Fig. 1). This path length is the sum of consecutive edge
⁶² lengths over all edges j in the path, $L_e = \sum_{j \in e} L_j$. Because for a given protein tree there
⁶³ are multiple paths from LECA to the present, and to exclude potential outliers, the path of
⁶⁴ median length was chosen as the normalization factor so that the normalized stem length is:
⁶⁵ $sl_g = L_s^g / \mathrm{med}(L_e^g)$. Following Pittis and Gabaldón, we refer to sl_g as a 'stem length' even
⁶⁶ though it is actually a normalized edge length.

⁶⁷   Under the molecular clock model, for any path p, $L_p = R_g T_p$, where T_p is the accumulated
⁶⁸ time for the path and R_g is the rate of substitution that is constant over time but may vary
⁶⁹ across genes. Thus, under the molecular clock model, the stem length satisfies that

⁷⁰
$$sl_g = R_g T_s / \mathrm{med}(R_g T_e) = T_s^g / T_e$$

⁷¹   Pittis and Gabaldón (2016a) compared the distributions of estimated stem lengths, $\widehat{sl}_g$ for
⁷² proteins of different origins. They found that $\widehat{sl}_g$ distributions from archaeal origin proteins
⁷³ ($g \in R$) were, based on Mann-Whitney U tests, significantly shifted to be larger than those of
⁷⁴ bacterial ($g \in C$) origin which were, in turn, significantly greater than alphaproteobacterial
⁷⁵ proteins ($g \in M$) (the latter are assumed to correspond to genes that originated with the
⁷⁶ mitochondrial symbiont). Since the times are constant within groups (for instance, $T_s^g = T_s^R$
⁷⁷ for $g \in R$), they interpret this as evidence that $T_s^R > T_s^C > T_s^M$. An important conclusion
⁷⁸ of their study was, therefore, that the mitochondrial symbiosis took place much later in eu-
⁷⁹ karyogenesis than suggested in mitochondria early hypotheses (e.g., Lane and Martin 2010).
⁸⁰ More recently, Vosseberg and colleagues (Vosseberg *et al.* 2020) have extended this approach
⁸¹ to address the relative timings of the mitochondrial symbiosis and gene duplication events
⁸² for a variety of functional classes of protein families that expanded during eukaryogenesis.

⁸³   Pittis and Gabaldón's approach was strongly criticized by Martin and colleagues (Martin
⁸⁴ *et al.* 2017) who argued that the results were meaningless because the method depends
⁸⁵ on the assumption that a molecular clock should hold over evolutionary time spans on
⁸⁶ the billion-year time scale. They investigated a number of the individual phylogenies from
⁸⁷ Pittis and Gabaldón study and showed that variation in edge lengths within gene trees
⁸⁸ were substantial and not consistent with a molecular clock. Pittis and Gabaldón have since
⁸⁹ countered by arguing that their approach does not assume a molecular clock and demonstrate
⁹⁰ its ability to successfully recover correct orderings of more recent evolutionary divergences in
⁹¹ eukaryotes (Pittis and Gabaldón 2016b). However, they did not provide a detailed theoretical
⁹² justification for why the method should give reliable evolutionary orderings in the absence

of a molecular clock

Here we show that assuming a molecular clock is not necessary for the method to work. In
what follows, we will show that if proteins a and b in two groups $a \in A$ and $b \in B$ of different
origins evolve independently according to the same time-dependent stochastic substitution
rate process, the distributions of the normalized stem lengths of phylogenies of proteins B
will be systematically larger than A, i.e. $\mathbb{P}[sl_b - sl_a > 0] > 1/2$, if and only if $T_s^B > T_s^A$. We
show that restrictions on the stochastic substitution rate process and the data set required
for the foregoing result to hold are surprisingly few, but do include a requirement that there
are no systematic differences between groups A and B at any given time point. This result
is shown to hold even when estimation of edge lengths is taken into account, although a
modified version of the Mann-Whitney U test will be required for testing when the variances
in edge length estimates are systematically different between the groups. We also show that
these methods will work in cases in which the proteins within the groups being compared
have ranges of different ages. In the latter cases, however, we suggest that statistical test
rejection is difficult to meaningfully interpret. Finally, we outline scenarios in which the
required assumptions of the method will not hold and caution is warranted.

Borrowing on relaxed molecular clock theory that assumes that the rate of substitution
varies stochastically over the tree (Bromham *et al.* 2019), suppose that the rate of substitu-
tion at any point along a path p can be represented as $R_g r_p^g(t)$, where $\{r_p^g(t)\}$ is a continuous
time stochastic process (R_g is the overall fixed rate of gene g as before). Assuming a con-
ventional Markov substitution model, the probability of substituting state i with state j in
$(t, t+h]$ is $q_{ij} R_g r_p(t) h + o(h)$, for some state transition rate q_{ij}. Some constraint is required to
identify parameters and we assume, without loss of generality, that $E[R_g] = 1$ as well as the
conventional constraint, $\sum_i \sum_{j \neq i} \pi_i q_{ij} = 1$. Since the chance of two or more substitutions is
small relative to h, $o(h)$, the expected number of substitutions in $(t, t + h]$, $E[N(t, t + h)]$, is
$\sum_i \sum_{j \neq i} \pi_i q_{ij} R_g r_p(t) h + o(h) = R_g r_p(t) h + o(h)$. Taking $u_0 = 0, u_1 = t_0/N, \ldots, u_N = t_0$, the
number of substitutions in the time period $(0, t_0]$ is the sum, $\sum_{k=1}^N N(u_{k-1}, u_k)$ of the substi-
tutions over the intervals $(0, u_1], \ldots, (u_{N-1}, u_N]$. Since this is true for any N, the expected
number of substitutions along path p and over time period $(0, t_0]$ is

$$E[N(0, t_0)] = \lim_N \sum_{k=1}^N E[N(u_{k-1}, u_k)]$$

$$= \lim_N \{\sum_{k=1}^N R_g r_p(u_{k-1}) t_0/N + N o(1/N)\} = R_g \int_0^{t_0} r_p(t) dt.$$

123   Thus the stem length for protein g is

124
$$sl_g = \int_{T_e}^{T_s+T_e} r_s^g(t) \; dt / \mathrm{med}(\int_0^{T_e} r_e^g(t) \; dt)$$

125   Suppose that comparison is between group A and B and that $T_s^B > T_s^A$. Let

126
$$Y_g = \mathrm{med}(\int_0^{T_e} r_e^g(t) \; dt), \; X_g = \int_{T_e}^{T_s^A+T_e} r_s^g(t) \; dt \text{ and } V_g = \int_{T_s^A+T_e}^{T_s^B+T_e} r_s^g(t) \; dt. \tag{1}$$

Then $sl_g = X_g/Y_g$ for $g \in A$ and $sl_g = (X_g + V_g)/Y_g$ for $g \in B$.

We now make the assumption that for any given path p and any two proteins g and g'

in $A \cup B$, the rate processes are probabilistically equivalent on $[0, T_s^A]$:

For any $0 \le t_1 < \cdots t_m \le T_s^A$, the joint probability distribution of $[r_p^g(t_1), \ldots, r_p^g(t_m)]$ and

$[r_p^{g'}(t_1), \ldots, r_p^{g'}(t_m)]$ are the same.

Note that this assumption allows possibly radical rate changes throughout the tree and

across proteins. Moreover, the rate processes need not be stationary nor Markov processes.

What is required, however, is that there be no systematic differences between the two groups.

Thus, for instance, the model allows that as a consequence of a radical environmental change

or change in population size at time t, for some particular lineage l, the distribution of $r_l^g(t)$

is skewed to the right of the distribution of the rate $r_l^g(t')$ at some other time t'. But that

difference in distributions is expected to apply whether $g \in A$ or $g \in B$. For simplicity, we

also make the assumption that that the rate process is bounded such that $r_p^g(t) \in [\beta, \gamma]$ for

some $\beta > 0$ and $\gamma < \infty$. This assumption is reasonable as we do not expect substitution

rates to go to 0 or to increase without bound. In any case, this assumption can be loosened

but some sort of assumption is required to avoid having a stem lengths that are almost 0

due entirely to having extremely low average rates along the stem or extremely high average

rates from LECA to present. Finally, we assume that the rate processes are independent

over genes.

With the assumptions above, if the eukaryotic taxa sampled are the same for groups

A and B, then Y_g will have the same distribution whether $g \in A$ or $g \in B$. Note that in

Pittis and Gabaldón the same taxa were not necessarily present in any two groups of proteins

being compared. We argue below that, for the approach to work, it is best if Y_g has the same

distribution for $g \in A$ or $g \in B$, otherwise differences in normalized stem lengths between

the groups may be due to unusual rates for eukaryote taxa present in one group but not the

other. Thus, it may be desirable to take means or medians over the set of eukaryote taxa

present in both groups. Nevertheless, medians are not as likely to be affected by outlying

rates (which was one of the original motivations for using them), so if taxon sampling is

comparable for the two groups, the distributions of Y_g are expected to be approximately the
same for the two groups under the assumptions above.

Consider now $\mathbb{P}[sl_b - sl_a > u]$ for fixed u, $a \in A$ and $b \in B$. In terms of the random
variables above this can be expressed as $\mathbb{P}[U + V_b/Y_b > u]$ where $U = X_b/Y_b - X_a/Y_a$. Since
$r_p^g(t) \in [\beta, \gamma]$, the smallest V_b/Y_b could be is $w := \beta(T_s^B - T_s^A)/(\gamma T_e) > 0$. Thus

$$\mathbb{P}[sl_b - sl_a > u] \geq \mathbb{P}[U + w > u] = \mathbb{P}[U > u - w] \tag{2}$$

161 Similarly

$$\mathbb{P}[sl_a - sl_b \geq u] = \mathbb{P}[-U - V_b/Y_b \geq u] \leq \mathbb{P}[-U - w \geq u] = \mathbb{P}[-U \geq u + w] \tag{3}$$

163 With the assumptions above X_b/Y_b and X_a/Y_a have the same distribution. Thus $U =$
164 $X_b/Y_b - X_a/Y_a$ has a symmetric distribution around 0. Consequently,

$$\mathbb{P}[sl_a - sl_b \geq u] \leq \mathbb{P}[U \geq u + w] \leq \mathbb{P}[U > u - w] \leq \mathbb{P}[sl_b - sl_a > u] \tag{4}$$

166 where the first inequality is from (3) and the third from (2). The inequalities are strict
167 unless U does not have mass in $(u - w, u + w]$. Since X_b/Y_b and X_a/Y_a have the same
168 distribution then U is sure to have positive probability in $(-w, w)$. Thus with $u = 0$ and
169 since $\mathbb{P}[sl_a - sl_b \geq 0] = \mathbb{P}[sl_b - sl_a \leq 0] = 1 - \mathbb{P}[sl_b - sl_a > 0]$, then we have

$$0 < \mathbb{P}[sl_b - sl_a > 0] - \mathbb{P}[sl_a - sl_b \geq 0] = 2\mathbb{P}[sl_b - sl_a > 0] - 1 \tag{5}$$

or $\mathbb{P}[sl_b - sl_a > 0] > 1/2$.

We have shown that under the alternative hypothesis that $T_s^B > T_s^A$ then we have that
$\mathbb{P}[sl_b - sl_a > 0] > 1/2$. Under the null hypothesis that $T_s^B = T_s^A$, the distributions of sl_b and
sl_a are the same. Thus if the actual normalized stem lengths were used for the two groups,
the null and alternative hypotheses of interest imply the null and alternative hypotheses of
the Mann-Whitney U test. However, the actual stem lengths are not known for the two
groups; only estimates of these quantities from sequence data are available. This raises the
question: Will the null and alternative hypotheses $T_s^B = T_s^A$ and $T_s^B > T_s^A$ correspond to
Mann-Whitney U test null and alternative hypotheses if estimated stem length distributions
are used?

Assume that the number of sites is sufficiently large for each gene that asymptotic likeli-
hood theory gives a good approximation to the sampling distributions of the stem lengths.
That theory implies that $\hat{L}_p^g$ is approximately normal with mean L_p^g. It follows from delta-
method arguments (cf. §5.3.2 of Bickel and Doksum 2007) that $\hat{L}_s^g/\hat{L}_e^g$ is approximately
normal with mean L_s^g/L_e^g for any path e from LECA to a eukaryotic taxon. Because there

are finitely many paths, for relatively large sequence lengths, the path, e^* say, corresponding

to the median $\hat{L}^g_e$ should coincide with the path corresponding the median L^g_e. Thus $\widehat{sl}_g$ will

be $\hat{L}^g_s/\hat{L}^g_{e*}$ for the path e^* corresponding to the median L^g_e. Since $\hat{L}^g_s/\hat{L}^g_{e*}$ is approximately

normal with mean L^g_s/L^g_{e*}, $\widehat{sl}_g = sl_g + \epsilon_g$ where ϵ_g has a normal distribution that is sym-

metric around 0. Consequently $\mathbb{P}[\widehat{sl}_b - \widehat{sl}_a > 0] = \mathbb{P}[sl_b - sl_a > \epsilon_b - \epsilon_a]$. As a difference

of independent, symmetric normals, $\epsilon_b - \epsilon_a$ is symmetric normal. Denote the probability

density function of the latter as $p(u)$. Then

193
$$\mathbb{P}[\widehat{sl}_b - \widehat{sl}_a > 0] = \int_{-\infty}^{\infty} \mathbb{P}[sl_b - sl_a > u]\ p(u)\ du$$

194
$$= \int_0^{\infty} \mathbb{P}[sl_b - sl_a > u]\ p(u)\ du + \int_{-\infty}^0 \mathbb{P}[sl_b - sl_a > u]\ p(u)\ du$$

195
$$= \int_0^{\infty} \mathbb{P}[sl_b - sl_a > u]\ p(u)\ du + \int_{-\infty}^0 \mathbb{P}[sl_a - sl_b < -u]\ p(u)\ du$$

196
$$= \int_0^{\infty} \{\mathbb{P}[sl_b - sl_a > u] + \mathbb{P}[sl_a - sl_b < u]\}\ p(u)\ du$$

197
$$= \int_0^{\infty} \{\mathbb{P}[sl_b - sl_a > u] + 1 - \mathbb{P}[sl_a - sl_b \geq u]\}\ p(u)\ du \tag{6}$$

198 By (4), under the alternative hypothesis, $\mathbb{P}[sl_b - sl_a > u] - \mathbb{P}[sl_a - sl_b \geq u] \geq 0$ with strict

199 inequality in a neighbourhood of 0. Thus

200
$$\mathbb{P}[\widehat{sl}_b - \widehat{sl}_a > 0] > \int_0^{\infty} p(u)\ du = 1/2 \tag{7}$$

201 We see that the alternative hypothesis of interest corresponds to $\mathbb{P}[\widehat{sl}_b - \widehat{sl}_a > 0] > 1/2$

202 as required for the Mann-Whitney U test. However, the situation under the null is a little

203 more problematic. Under the null hypothesis, $\mathbb{P}[sl_b - sl_a > u] - \mathbb{P}[sl_a - sl_b \geq u] \leq 0$, so (6)

204 gives that

205
$$\mathbb{P}[\widehat{sl}_b - \widehat{sl}_a > 0] \leq \int_0^{\infty} p(u)\ du = 1/2$$

However, the Mann-Whitney U test requires that the distributions of $\widehat{sl}_b$ and $\widehat{sl}_a$ be the

same. Although the distributions of the sl_a and sl_b are the same and the distributions of

ϵ_a and ϵ_b are both symmetrically normal, their variances need not be comparable. These

variances reflect precision of estimation and reasons that they might differ include that

numbers of sites in alignments tend to differ substantially for one group versus the other.

The null distribution used by the Mann-Whitney U test is not correct in such settings.

Indeed, Kyusa (2000) shows that if the distributions for the two groups considered by the

Mann-Whitney test are normal but with differing variances, the type I error of the test can

be inflated. Nevertheless, Chung and Romano (2015) provide an alternative test that can

²¹⁵ be used under the null hypothesis, $\mathbb{P}[\widehat{sl}_b - \widehat{sl}_a > 0] \leq 1/2$ but $\widehat{sl}_b$ and $\widehat{sl}_a$ do not have the
²¹⁶ same distribution and we recommend use of this test as a safeguard. That being said, if the
²¹⁷ variability of estimation is comparable for the two groups, a Mann-Whitney U test should
²¹⁸ give reasonable results.

²¹⁹ Much of the preceding discussion considers a case arising in both the analyses of Pittis and
²²⁰ Gabaldón (2016a), and Vosseberg and colleagues (2020) where the times of origin associated
²²¹ with a stem length are constant for proteins within a group (for instance because they all
²²² derived from a mitochondrial symbiont or were all inherited from the archaeal host). Pittis
²²³ and Gabaldón, and Vosseberg and colleagues, however, also compared groups made up of
²²⁴ proteins of different bacterial origins and considered functional classes of proteins as groups.
²²⁵ In these cases, proteins within a group are not expected to have a single time of origin (i.e.,
²²⁶ T_s^g will vary within a group).

²²⁷ To allow for stem times that are not constant within groups, we assume a model in which
²²⁸ T_s^g are independent across genes and independent of the rate variation processes $r_p^g(t)$. With
²²⁹ the previous assumptions, the null hypothesis (that for $a \in A$ and $b \in B$, T_s^a and T_s^b have
²³⁰ the same distribution) implies that sl_a has the same distribution as sl_b. The alternative
²³¹ hypothesis of greatest interest is that there is no overlap in the T_s^a and T_s^b distributions: that
²³² $\mathbb{P}[T_s^a < T_s^b] = 1$. With this assumption, the arguments assuming fixed $T_s^A < T_s^B$, apply for
²³³ the conditional distribution of $sl_s^a - sl_s^b$, given T_s^a and T_s^b. Averaging over T_s^a and T_s^b give
²³⁴ that $\mathbb{P}[sl_b - sl_a > 0] \leq 1/2$ under the null hypothesis and $\mathbb{P}[sl_b - sl_a > 0] > 1/2$ under the
²³⁵ alternative hypothesis.

²³⁶ One difficulty with analyses when the T_s^g vary within groups is that we have no control
²³⁷ over the alternative hypothesis. The desired alternative conclusion is that $\mathbb{P}[T_s^a < T_s^b] = 1$
²³⁸ and we have argued above that such an alternative relationship leads to a Mann-Whitney U
²³⁹ test null and alternative hypotheses for $\widehat{sl}_b - \widehat{sl}_a$. But suppose now that

$$\mathbb{P}[T_s^b - T_s^a > z] > \mathbb{P}[T_s^a - T_s^b > z], \text{ all } z \tag{8}$$

²⁴¹ This condition is related to the hypothesis of interest but might not be very meaningful. For
²⁴² instance, if $\log(T_s^b) \sim N(\mu_B, \sigma^2)$ and $\log(T_s^a) \sim N(\mu_A, \sigma^2)$ with $\mu_B > \mu_A$, (8) holds but if
²⁴³ σ^2 is large then there is a substantial chance that any given T_s^a is larger than a given T_s^b.
²⁴⁴ In other words, if substantial numbers of proteins in a given group A have an older origin
²⁴⁵ than many of the proteins in group B, then what should we conclude from rejecting the null
²⁴⁶ hypothesis that the group A distribution is shifted to be older than the group B distribution?
²⁴⁷ More broadly, the rationale for grouping proteins together to test hypotheses about timings
²⁴⁸ of origin is unclear if the age ranges across proteins in the groups heavily overlap and there
²⁴⁹ is large variation amongst them.

250   We now show that (8) can lead to $\mathbb{P}[\widehat{sl}_b > \widehat{sl}_a] > 1/2$. Assume for simplicity that

251   $\mathbb{P}[r_p(t) = 1] = 1$. Then the arguments leading to (4) apply with random $W = (T_s^b - T_s^a)/T_e$

252   and exact equality:

$$\mathbb{P}[sl_b - sl_a > u] \quad = \quad \mathbb{P}[W > u - U] = \mathbb{P}[T_s^b - T_s^a > T_e(u - U)]$$

$$= \quad \mathbb{P}[T_s^b - T_s^a > T_e(u + U)] \tag{9}$$

$$\mathbb{P}[sl_a - sl_b > u] \quad = \quad \mathbb{P}[T_s^a - T_s^b > T_e(u + U)] \tag{10}$$

256   where the last equality in (9) and the equality in (10) follows from independence and the

257   symmetric distribution of U. Letting $Z = T_e(u - U)$, then

$$\mathbb{P}[sl_b - sl_a > u] - \mathbb{P}[sl_a - sl_b > u] = \int_{-\infty}^{\infty} \{\mathbb{P}[T_s^b - T_s^a > z] - \mathbb{P}[T_s^a - T_s^b > z]\}p(z)\,dz > 0$$

It follows as before that $\mathbb{P}[\widehat{sl}_b > \widehat{sl}_a] > 1/2$. Since the Mann-Whitney U test or the Chung and

Romano robust alternative are designed to detect $\mathbb{P}[\widehat{sl}_b > \widehat{sl}_a] > 1/2$ vs $\mathbb{P}[\widehat{sl}_b > \widehat{sl}_a] \leq 1/2$,

whatever the cause, rejection could correspond to less meaningful alternatives like those

discussed above.

Another concern arises specifically for groups of genes that were laterally acquired during

eukaryogenesis from a prokaryotic lineage. As discussed above and shown in Fig. 1, the actual

stem-length time T_{s*} for these genes is less than the stem-length time T_s for the observed

tree. Throughout the preceding discussion we have assumed that $T_{s*}^g \approx T_s^g$. Suppose now

that the two groups A and B have different prokaryotic origins and that $\mathbb{P}[T_{s*}^B > T_{s*}^A] = 1$.

Will the Mann-Whitney U test be likely to reject in this case? Let $K_g = T_s^g/T_{s*}^g \geq 1$.

Recall that $T_s^g > T_{s*}^g$ is expected because an immediate extant sister group to the actual

prokaryotic transfer lineage is unlikely to be among the sampled taxa because of extinction.

If the extinction processes are roughly the same for the two prokaryotic origin groups, then

it is plausible that K_g will have the same distribution for the two groups. We thus make the

additional assumption that the K_g have the same distribution for the two groups and are

independent of the rate process below.

We now argue that the Mann-Whitney U test is indeed likely to reject when two such

acquired groups of genes A and B have different single prokaryotic origins and origin times

are well separated: $\mathbb{P}[T_{s*}^B > T_{s*}^A] = 1$. We condition on fixed T_{s*}^A and T_{s*}^B in what follows.

Because the result below holds for all fixed T_{s*}^B and T_{s*}^A, then averaging with respect to the

distribution of $[T_{s*}^A, T_{s*}^B]$ gives the result for random T_{s*}^A and T_{s*}^B.

Similarly as when $T_{s*} = T_s$ was assumed, $sl_g = X_g/Y_g$ for $g \in A$ and $sl_g = (X_g + V_g)/Y_g$

for $g \in B$, where Y_g is as in (1) but now

$$X_g = \int_{T_e}^{K_g T_{s*}^A + T_e} r_s^g(t)\,dt \text{ and } V_g = \int_{K_g T_{s*}^A + T_e}^{K_g T_{s*}^B + T_e} r_s^g(t)\,dt.$$

For $a \in A$ and $b \in B$, observe first that, because the $\{r_b(t)\}$ and $\{r_b(t)\}$ processes are probabilistically equivalent, and because K_a and K_b have the same distributions, then X_b/Y_b and X_a/Y_a have the same distribution. Second, because $K_g \geq 1$, the smallest V_b/Y_b can be is then $w = \beta(T_{s*}^B - T_{s*}^A)/(\gamma T_e)$. These two properties were what was used in the arguments leading to (2)–(5) and so those results hold in this setting. Here (5) gives the conclusion required for the Mann-Whitney U test, that $\mathbb{P}[sl_b - sl_a > 0] > 1/2$ and (2) is the key inequality that can be used, exactly as before, to show (7), that, even with estimation, $\mathbb{P}[\widehat{sl_b} - \widehat{sl_a} > 0] > 1/2$.

Note that the above assumption that K_g has the same distribution across groups is violated for some comparisons. For instance, comparisons of distributions from groups of acquired genes (where $T_{s*}^g \leq T_s^g$ and, possibly, broad ranges of T_s^g values within the group) with genes inherited from the asgard archaeon-eukaryote common ancestor or genes that originate by duplication (for which $T_{s*}^g = T_s^g$ in both cases). In the simplest case of fixed T_s^g values for the genes in an acquired group, the inferred age of that group will be biased to be younger than its true age in comparison with genes inherited from the asgard-eukaryote common ancestor or groups of duplicated genes.

We have shown that validity of the edge length ratio methods introduced by Pittis and Gabaldón (2016a), and extended by Vosseberg and colleagues (2020) do not require a molecular clock. They can be justified in much more general settings where substitution rates in a protein stochastically vary over the tree. Indeed, the only restrictions are that the stochastic substitution rate process is bounded away from 0 and infinity and that genes in groups of different origins (or functional classes) in a genome are all independently evolving according to this same process (i.e. the rate process for different genes are probabilistically equivalent). In terms of biological realism, it is this latter assumption that may not always hold. For example, it is well known that proteins may periodically experience episodes of rapid adaptive evolution due to acquisition of novel functions and/or loss of ancestral functions (Studer, Dessailly and Orengo, 2013). If this functional divergence differentially affected proteins within groups of different origins or functional classes, then stem length distributions of one group of proteins versus another will likely reflect this episodic shift in rates in one group, and cannot be used to test if they originated at different times. When applying these methods, it is therefore important to investigate evidence for systematic differences in the evolutionary dynamics of one group of proteins versus another.

It is also important to select edge lengths from phylogenies for different gene groups to be compared in the same manner to ensure that no biases are introduced. For example, when extending the approach to genes duplicated during eukaryogenesis, Vosseberg and colleagues (2020) were faced with deciding how to deal with the multiple possible edges or paths to

LECA nodes created by duplications (Fig. 2). Their approach was to always select the minimum of all possible edge or path lengths for calculations of stem lengths or duplication lengths (all duplication lengths were calculated as $dl_g = L_d^g/\mathrm{med}(L_e^g)$). For two groups D and S of duplicated or acquired genes (for which stem lengths can include duplicated edges), the alternative hypothesis of greatest interest is H_{DS}: $\mathbb{P}[T_{di} > T_{sj}] = 1$ for $d \in D$ and $s \in S$, regardless of i and $j \in \{1, 2\}$.

Figure 2: **Example of the multiple possible stem paths and duplication edges in the phylogeny of a protein that has been duplicated during eukaryogenesis.** This gene was acquired by the proto-eukaryote genome from a prokaryotic lineage (green box) and then duplicated (magenta box) prior to LECA (the LECA nodes of each duplicate are shown as purple circles). As a result there are two possible stem paths with lengths L_{s1}^g and L_{s2}^g (green arrows), two possible duplication edges with lengths L_{d1}^g and L_{d2}^g and two possible median paths within the eukaryote subtrees (purple boxes) with lengths $\mathrm{med}(L_{e1}^g)$ and $\mathrm{med}(L_{e2}^g)$. To calculate stem lengths and duplication lengths, Vosseberg and colleagues (2020) used the edges/paths with minimum values (minimums shown as thicker edges).

Assume that evolution is independent post duplication. Then, if all duplicated lengths are included in a comparison involving minima, the arguments above apply and the Mann-Whitney U test would be likely to reject when H_{DS} holds. One possible motivation for using minima is that it could potentially alleviate bias due to the functional divergence phenomenon alluded to above because functional divergence is more likely to have occurred in the duplicate with the longer L_{di}^g over $i \in \{1, 2\}$. Unfortunately, taking minimums leads to some loss of information and could lead to a bias which we illustrate assuming no functional divergence. Similar arguments as those above related to the median path lengths, imply

that the Mann-Whitney U test would have substantial probability of rejection whenever
$\mathbb{P}[\min(T_{d1}, T_{d2}) > \min(T_{s1}, T_{s2})] = 1$. If H_{DS} holds and evolution is independent, then this
hypothesis will hold, so the approach should work. In less ideal alternative hypothesis
scenarios where there is overlap in the distributions of T_{di} and T_{sj}, biases can occur. As an
illustrative example, suppose that S is a stem-length group of acquired genes, that $T_e = 1.5$
billion years ago, that $T_{sj} = 0.4$ billion years and that, independently, $T_{di} \sim U(0, 1)$. Then

$$\mathbb{P}[T_{di} > T_{sj}] = \mathbb{P}[T_{di} > 0.4] = 1 - 0.4 = 0.6,$$

consistent with longer duplication lengths. But

$$\mathbb{P}[\min(T_{d1}, T_{d2}) > \min(T_{s1}, T_{s2})] = \mathbb{P}[T_{d1} > 0.4, T_{d2} > 0.4] = 0.6^2 = 0.36$$

Consequently, based on the Mann-Whitney U test, one would conclude that the age of the
duplicated group of genes D tended to be less than the acquired genes S when, in fact, 60%
of the duplication group genes, D, duplicated prior to their acquisition.

Another complication arises in the comparison of duplication groups and stem-length
groups. For duplication groups, duplication lengths are all minima. The stem-length groups,
however, are usually a mix of stem lengths, some of which are chosen to be minima (Fig. 2)
and some of which did not require a minimum (Fig. 1) (Vosseberg *et al.* 2020). In this case
biases might arise even under the strong hypothesis, $\mathbb{P}[\min(T_{d1}, T_{d2}) > \min(T_{s1}, T_{s2})] = 1$ for
$d \in D$ and $s \in S$, making it difficult to reject when H_{DS} holds. Such problems can be averted
by including all duplication and stem lengths rather than minima. There are potentially
other ways of addressing functional divergence that may be less likely to introduce bias
(e.g., identifying functionally divergent sites using methods reviewed in Studer, Dessailly
and Orengo (2013) and removing them prior to analysis).

In summary, although there are a number of caveats, if the assumptions of the methods
we have elaborated above are met by the data, these edge length ratio methods have the
potential to provide important new insights into the roles of gene duplication and gene
invention in different cellular systems and clarify the relative contributions of host, symbiont
and lateral transfers to a lineage of interest.

Acknowledgements

This research was supported a grant (Award ID: 735923LPI) awarded to A.J.R. and E.S.
as part of the Moore-Simons Project on the Origin of the Eukaryotic Cell. E.S. and A.J.R.
also acknowledge partial support for this work from Discovery grants awarded to them by
the Natural Sciences and Engineering Research Council of Canada. A.J.R. acknowledges the
Allan Wilson Centre in New Zealand for sabbatical support received in 2006 when some of
these ideas were first discussed.

References

Bickel, P.J. and Doksum, K.A. (2007). Mathematical Statistics: Basic ideas and selected topics. Pearson, New Jersey.

Bromham, L., Duchne, S., Hua, X., Ritchie, A.M., Duchne, D.A., and Ho, S.Y.W. (2018). Bayesian molecular dating: opening up the black box. *Biol. Rev. Camb. Philos. Soc.* 93, 11651191.

Chung E.Y. and Romano, J.P. (2015). Asymptotically valid and exact permutation tests based on two-sample U-statistics. *J. Statist. Plann. and Inf.* 168:97–105.

Dacks, J.B., Field, M.C., Buick, R., Eme, L., Gribaldo, S., Roger, A.J., Brochier-Armanet, C., and Devos, D.P. (2016). The changing view of eukaryogenesis - fossils, cells, lineages and how they all come together. *J. Cell Sci.* 129, 36953703.

Eme, L., Spang, A., Lombard, J., Stairs, C.W., and Ettema, T.J.G. (2017). Archaea and the origin of eukaryotes. *Nat. Rev. Microbiol.* 15, 711723.

Kasuya, E. (2001). Mann-Whitney U test when variances are unequal. *An. Behav.* 61:1247–1249.

Lane N. and Martin W.F. (2010) The energetics of genome complexity *Nature* 467:929-934.

Martin, W.F., Roettger, M., Ku, C., Garg, S.G., Nelson-Sathi, S., and Landan, G. (2017). Late Mitochondrial Origin Is an Artifact. *Genome Biol. Evol.* 9:373-379.

Pittis, A.A. and Gabaldón, T. (2016a). Late acquisition of mitochondria by a host with chimaeric prokaryotic ancestry. *Nature.* 531:101-104.

Pittis, A.A. and Gabaldón, T. (2016b) On phylogenetic branch lengths distribution and the late acquisition of mitochondria. *bioRxiv doi: https://doi.org/10.1101/064873*

Porter, S.M. (2020). Insights into eukaryogenesis from the fossil record. *Interface Focus* 10, 20190105.

Rochette, N.C., Brochier-Armanet, C., and Gouy, M. (2014). Phylogenomic test of the hypotheses for the evolutionary origin of eukaryotes. *Mol. Biol. Evol.* 31:832-845.

Roger, A.J., Muñoz-Gómez, S. and Kamikawa, R. (2017). The origin and diversification of mitochondria. *Curr. Biol.* 27:R1177-R1192.

Vosseberg, J., van Hooff, J.J.E, Marcet-Houben, M., van Vlimmeren, A., van Wijk, L.M.,
Gabaldón, T. and Snel, B. (2020). Timing the origin of eukaryotic cellular complexity
with ancient duplications. *Nat. Ecol. Evol.* 5:92-100.

Studer, R.A., Dessailly, B.H., and Orengo, C.A. (2013). Residue mutations and their
impact on protein structure and function: detecting beneficial and pathogenic changes.
Biochem. J. 449:581-594.