The variability in potential biomarkers for cochlear synaptopathy after recreational noise exposure

Tine Vande Maele[1,*], Sarineh Keshishzadeh[2,*], Nele De Poortere[1], Ingeborg Dhooge[3,4], Hannah Keppler[1,4], Sarah Verhulst[2]

[1]*Ghent University, Department of Rehabilitation Sciences – Audiology*

[2] *Ghent University, Dept. of Information Technology – Hearing Technology @ WAVES*

[3] *Ghent University, Department of Head and Skin*

[4] *Ghent University Hospital, Department of Ear, Nose and Throat, Belgium*

[*]*These authors contributed equally to this work.*

Financial disclosures/conflicts of interest: This work was supported by UGent BOF-IOP project: "Portable Hearing Diagnostics: Monitoring of Auditory-nerve Integrity after Noise Exposure (EarDiMon)" (T.V.M., N.D.P., I.D., H.K. and S.V.) and European Research Council (ERC) under the Horizon 2020 Research and Innovation Program, grant agreement no. 678120 RobSpear (SK and SV). There are no conflicts of interest, financial, or otherwise.

All correspondence should be addressed to:

Sarineh Keshishzadeh, *Dept. of Information Technology – Hearing Technology @ WAVES,*

Ghent University, Technologiepark 126, Zwijnaarde 9052, Belgium.

Email: sarineh.keshishzadeh@ugent.be, s.verhulst@ugent.be

ABSTRACT

Objectives. Speech-in-noise tests and suprathreshold auditory evoked potentials are promising biomarkers to diagnose cochlear synaptopathy (CS) in humans. This study investigated whether these biomarkers changed after recreational noise exposure. **Design.** The baseline auditory status of 19 normal hearing young adults was analyzed using questionnaires, pure-tone audiometry, speech audiometry and auditory evoked potentials. Nineteen subjects attended a music festival and completed the same tests again at day 1, day 3 and day 5 after the music festival. **Results.** No significant relations were found between lifetime noise-exposure history and the hearing tests. Changes in biomarkers from the first session to the follow-up sessions were non-significant, except for speech audiometry, that showed a significant training effect (performance improvement). **Conclusions.** Despite the individual variability in pre-festival biomarkers, we did not observe changes related to the noise-exposure dose caused by the attended event. This can indicate the absence of noise-exposure-driven cochlear synaptopathy in the study cohort, or reflect that biomarkers were not sensitive enough to detect mild CS. Future research should include a more diverse study cohort, dosimetry and results from test-retest reliability studies to provide more insight into the relationship between recreational noise exposure and cochlear synaptopathy.

KEYWORDS

INTRODUCTION

Noise exposure in young adults mainly takes place during leisure time activities such as watching movies or plays, visiting nightclubs, attending music festivals or concerts, and listening to personal music players. In this population, concerts and music festivals are the loudest activities (Degeest, Clays, et al., 2017; Keppler, Dhooge, et al., 2015b; Petrescu, 2008; Smith et al., 2000). Sound level measurements during such events confirm that noise levels can exceed 100 dBA (Derebery et al., 2012; Mercier et al., 2003; Ryberg, 2009; Yassi et al., 1993). Despite the potential risk of noise-induced hearing loss (NIHL), the majority of young adults do not wear hearing protection devices (HPDs) when attending concerts (73.8%; (Degeest, Keppler, et al., 2017) or performances at music festivals (86%; (Mercier et al., 2003). After attending these music venues, 36-86% of young adults experience hearing related symptoms (HRS) such as temporary hearing loss, dullness or tinnitus (Keppler, Dhooge, et al., 2015b; Mercier et al., 2003; Smith et al., 2000).

The reduction of hearing sensitivity followingnoise exposure has extensively been investigated in animal and human studies (Bramhall et al., 2019). The incidence of NIHL depends on many factors, amongst others the noise characteristics: the intensity, frequency and duration of the exposure (Clark, 1991). NIHL can cause permanent threshold shifts (PTS) due to mechanical damage of the cochlear hair cells or supporting cells and metabolic changes that cause cell degeneration and cell death (Wang et al., 2002; Yamane et al., 1995). The reduction of hearing sensitivity after noise exposure can also be reversible. Mechanical damage of outer-hair-cell (OHC) stereocilia, reversible changes in supporting cells, or a swelling of auditory-nerve-fiber (ANF) terminals (Wang et al., 2002) can result in a temporary threshold shift (TTS) which recovers within minutes or hours (Emmerich et al., 2002; Trzaskowski et al., 2014) to weeks (Le Prell et al., 2012; Ryan et al., 2016). Research investigating the pathophysiology of TTS in rodents

indicates that excessive glutamate release excitotoxicity is the source of ANF terminal swelling

following noise exposure (Hakuba et al., 2000; Puel et al., 1998). Recent studies have shown that

TTS may not be as benign as previously thought, since it can be accompanied by permanent

deficits at the synapse level where type-I ANF terminals connect to the inner hair cells (IHCs).

Over the past years, this synapse loss has been named cochlear synaptopathy (CS) and has been

linked to noise exposure and aging in mice (Furman et al., 2013; Kujawa & Liberman, 2009;

Sergeyenko et al., 2013).

A single IHC contains 10 to 30 presynaptic ribbon-like structures surrounded with glutamate

vesicles. Each ribbon connects with a postsynaptic glutamate receptor patch at the ANF terminal

side (Liberman et al., 1990). ANFs with low and medium spontaneous firing rates have higher

thresholds and connect at the modiolar side of the IHC. They can be distinguished from ANFs

with high spontaneous firing rates and lower thresholds, that connect to the pillar side of the

IHC (Liberman, 1982). This kind of synapse is unique for the cochlea and retina, and allows fast

synaptic vesicle fusion with glutamate release, and fast recovery. Besides encoding a wide

dynamic range of sound intensities, the synaptic efficiency enables phase-locking to the

stimulus, i.e. action potentials follow the temporal precision of the stimulus fine structure and

its envelope. The phase-locked spikes are further encoded in the subcortical and cortical

pathway and are important for suprathreshold listening tasks, e.g., speech intelligibility and

sound localization (Henry et al., 2016; Safieddine et al., 2012).

Damage to the IHC synapses due to noise exposure, as seen in CS, can occur immediately after

noise exposure and can precede a slow degeneration of spiral ganglion cells (Lin et al., 2011).

Synapses that connect to ANFs with low spontaneous rates were shown to be most vulnerable,

whereas ANFs with higher spontaneous rates are more robust to noise damage (Furman et al.,

2013). This susceptibility of low-spontaneous-rate ANFs, combined with the distribution pattern

wherein multiple ANFs couple to a single IHC, could be the reason why auditory thresholds

recover or remain unaffected by permanent synaptic damage. CS could hence be a possible

explanation for the pathophysiology of hidden hearing loss, where normal auditory thresholds

are accompanied by suprathreshold hearing deficits such as decreased speech intelligibility in

noise (Bharadwaj et al., 2014; Smith et al., 2019). Furthermore, it has been argued that this

pathology may play a role in the origin of symptoms such as tinnitus or hyperacusis (Liberman &

Kujawa, 2017; Schaette & McAlpine, 2011).

Animal studies and computational auditory modeling studies have attempted to define

biomarkers for diagnosing CS noninvasively and have shown the potential of auditory evoked

potentials (AEPs). More specifically, the auditory brainstem response (ABR) and envelope

following response (EFR) are promising tests, given that ABR- and EFR-based derived metrics

relate directly to the number of histologically-verified synapse counts in animal models (Kujawa

& Liberman, 2009; Möhrle et al., 2016; Parthasarathy & Kujawa, 2018; Shaheen et al., 2015).

Suprathreshold ABRs are generated by the synchronous firing of ANFs and brainstem neurons in

response to the rapid onset of a brief stimulus (Hall, 2007). In rodents with CS, a reduced

amplitude of wave I or reduced wave I/V amplitude ratio was observed, which captures the

decreased integrity of the peripheral auditory nerve (Furman et al., 2013; Kujawa & Liberman,

2009, 2015; Lin et al., 2011). The EFR is elicited by continuous, amplitude-modulated signals and

the strength of the response to the stimulus envelope modulation frequency reflects temporal

coding precision in the subcortical neurons (Bidelman et al., 2015; Purcell et al., 2004). Animals

or models with CS showed reduced EFR-strength, which reflects a deficit in temporal coding

fidelity of the auditory system (Bharadwaj et al., 2014; Parthasarathy et al., 2018; Shaheen et

al., 2015; Vasilkov & Verhulst, 2019; Verhulst et al., 2018), even when hearing sensitivity recovered or remained normal.

Relating these AEP biomarkers to CS in living humans is far more difficult. Comorbidity with OHC loss and inter-subject variability of AEP measures in humans complicate the use of these methods. Moreover, CS diagnosis through direct synapse counts is not possible in living humans and only few human temporal bone studies have been conducted in relation to CS (Makary et al., 2011; Viana et al., 2015). Therefore, other methods have to be used to evaluate how AEP markers could relate to CS in living humans. Some studies were able to relate ABR and EFR strength to noise-exposure history (Bramhall et al., 2017; Grose et al., 2017; Paul et al., 2017; Skoe & Tufts, 2018), aging (Garrett & Verhulst, 2019; Konrad-Martin et al., 2012; Vasilkov & Verhulst, 2019) or tinnitus (Bramhall et al., 2018; Schaette & McAlpine, 2011), but other studies could not confirm these findings (Fulbright et al., 2017; Guest et al., 2018; Prendergast, Guest, et al., 2017; Spankovich, Le Prell, et al., 2017). Longitudinal studies in humans, where AEP-measurements are used to evaluate auditory-nerve integrity before and after noise exposure, are scarce. One study reported unchanged compound action potential (AP) amplitudes in electrocochleography, the analogue of ABR wave I amplitude, in relation to recreational noise exposure (Grinn et al., 2017).

This study investigates the relationship between recreational noise exposure and markers of CS and OHC damage to understand whether this exposure results in TTS, PTS, CS and/or speech in noise deficits. By adopting a test battery that includes pure tone audiometry (PTA) at conventional and extended high frequencies (EHFs), speech audiometry tests in quiet (SPiQ) and in noise (SPiN) and AEP measurements, we hypothesize that speech intelligibility and AEP measurements may deteriorate due to noise exposure, even if PTA is not affected or recovered

shortly after noise exposure. Multiple speech audiometry conditions, ABR and EFR stimuli were

used to study the relation of outcome measures to the individual noise-exposure history (Part I)

and to analyze the variability in these measures before and in a period of five days after

attending a music festival (Part II).

METHODS

PARTICIPANTS

Young adults aged 18 to 25 years who planned to attend a music festival during the summer of

2019 were recruited for this study. Volunteers completed a recruiting questionnaire, which was

used to exclude subjects with known hearing pathologies, history of ear surgery or tinnitus.

Study participation consisted of four measurement sessions: one baseline measurement session

(session 1) between 1 to 3 days before the start of the festival and 3 follow-up sessions: 1 day,

3 days and 5 days after the end of the music festival. Further on, these follow-up sessions will

respectively be referred to as session 2, session 3 and session 4. Subjects were asked not to

expose themselves to noise other than that of the festival during the follow-up period and not

to use party drugs. Participants were free to use hearing protection.

Twenty subjects participated in the first session, 8 males and 12 females (mean age 21.5 years

± 2.24 SD; standard deviation). Of those subjects, the best ear was chosen as the test ear based

on tympanometry and PTA. This was the right ear for 7 subjects (4 males) and the left ear for 13

subjects (4 males). One male subject (right testing ear) dropped out of the study for the last

measurement session, resulting in 19 participating subjects for the follow-up analysis in Part II,

with mean age of 21.6 years (SD 2.27). At every test session, subjects completed a test battery

consisting of questionnaires, PTA, SPiQ and SPiN tests, and AEP measurements. The test protocol

had a duration of maximum 3 hours per session (including breaks and information) and tests

were conducted in the same order for every subject and at every session.

This study was approved by the ethical committee of the Ghent University Hospital (Belgium)

and was performed following the statement of ethical principles of the Declaration of Helsinki.

All subjects were informed about the testing procedures and signed an informed consent.

Subjects received a financial compensation for their participation.

QUESTIONNAIRES

The questionnaire of the baseline session consisted of four parts and was completed online or

during the first session. The first part consisted of sociodemographic questions, questions

regarding present HRS, illness, recent drug use and noise exposure. Finally, the general use of

hearing protection devices and the type of hearing protection was queried.

The second part consisted of questions regarding noise-exposure history based on a Dutch

translation of Jokitulppo et al. (2006) by Keppler, Dhooge, et al. (2015b). A list of relevant,

noisy recreational activities was presented. First, subjects estimated the frequency of noise

exposure caused by each activity and the duration of each exposure in hours. Second, the

exposure level (Laeq) was estimated using a relative scale based on communication effort,

ranging from 60 dBA (sound level of a normal conversation) to 100 dBA (sound level which

makes communication impossible). The use of HPD was asked and converted in the equation

as a Laeq-reduction of 10 dB. Finally, for each activity, the total number of years that this

activity was present in their lives on the one hand, and for the last three years on the other,

was asked. Cumulative lifetime-noise-exposure history (Laeq,life) was calculated using the

formula of Jokitulppo et al. (2006). An analogous calculation was made for recent noise-

exposure history (Laeq,recent), which represents cumulative noise exposure over the past

three years.

The third and fourth part respectively consisted of the Youth Attitude to Noise Scale (YANS),

(Widen et al., 2006) and Beliefs about Hearing Protection and Hearing Loss (BAHPHL),

(Svensson et al., 2004) as modified by Keppler et al. (Keppler, Dhooge, Degeest, et al., 2015;

Keppler, Dhooge, et al., 2015a; Keppler & Dhooge, 2010). Both questionnaires use a five-

degree Likert Scale ranging from "totally disagree" to "totally agree" to determine the

attitudes and beliefs towards noise, hearing protection and hearing loss. A higher score

reflects more positive attitudes towards noise, less concerns about hearing loss and less belief

of benefit of HPDs.

At sessions 2, 3 and 4 a questionnaire that evaluates HRS, recent drug and alcohol

consumption, and between-session noise exposure was completed. The first follow-up session

contained additional questions regarding the festival duration and noise exposure. For each

level of communication effort, the subject was asked to estimate a percentage of exposure

time and whether HPDs were used in this listening situation to determine the festival noise

exposure (Laeq, festival) based on a 40-hour-week noise exposure calculation (Jokitulppo et

al., 2006; Keppler, Dhooge, et al., 2015b).

OTOSCOPY AND TYMPANOMETRY

Otoscopic inspection of the ear canal and tympanic membrane was performed at the baseline

session. All test ears had normal otoscopic results.

Middle-ear admittance was bilaterally measured at the baseline session, using a GSI TympStar

(Grason-Stadler) tympanometer with a 226 Hz, 85 dB sound pressure level (SPL) probe tone.

All test ear tympanograms had an ear canal volume between 0.6 and 2.0 mmho, a static

acoustic admittance level within a range of 0.3 to 1.7 mmho and a middle-ear pressure

between -100 and 100 daPa, and were defined as normal type-A tympanograms.

PURE-TONE AUDIOMETRY

PTA thresholds were measured while subjects were seated in a double-walled sound-

attenuating measurement booth. At the baseline session, both ears were tested to select the

test ear. Air-conduction thresholds were measured at conventional octave frequencies 0.125,

0.250, 0.500, 1, 2, 4 and 8 kHz and half-octave frequencies 3 and 6 kHz, and at EHFs of 10,

12.5, 14 and 16 kHz using the modified Hughson-Westlake procedure. An Equinox

Interacoustics audiometer was used and stimuli were transmitted using Interacoustics TDH-

39 headphones and Sennheiser HAD-200 headphones for conventional frequencies and EHF,

respectively. If no threshold could be obtained at the maximum output level, the threshold

was considered as this maximum stimulus level for data analysis. The ear with better

thresholds on conventional frequencies was chosen as a test ear, provided that these

thresholds were all 20 dB HL or better and that this ear had a normal otoscopic evaluation and

a type A tympanogram. Since air-conduction thresholds on conventional frequencies never

exceeded 20 dB HL and type-A tympanograms showed normal middle-ear admittance in all

test ears, bone conduction measurements were not performed.

SPEECH INTELLIGIBILITY IN QUIET AND IN NOISE

At each session, SPiQ and SPiN tests were performed at the test ear using the Flemish Matrix

sentence test (Luts et al., 2014) in Apex 3 software (Francart et al., 2008). Sentences were

presented in a relatively quiet room using a laptop connected to a Fireface UCX soundcard

(RME) and HDA-300 (Sennheiser) headphones.

At each session, five randomly chosen experimental test lists were presented in five differently

filtered conditions, presented in a random order. By filtering speech stimuli, different auditory

mechanisms can be targeted. Low-pass (LP) filtered speech understanding is thought to mainly

rely on temporal fine-structure cues, whereas high-pass (HP) filtered speech would mostly rely

on envelope coding of the auditory system (Garrett et al., 2020). Therefore, the five conditions

consisted of two conditions in quiet, where speech was filtered with a zero-phase 256th-order

FIR LP filter or with a zero-phase 1024th-order FIR HP filter with cutoff values of 1500 and 1650

Hz, respectively. Three conditions were presented in a speech-shaped noise with a fixed level

of 70 dB SPL: a broadband condition (BB) where no filtering was applied, a LP and a HP

condition, where speech as well as noise signals were filtered using the same filter cutoff

values as for LP and HP condition in quiet, respectively. Due to the filtering, the SPL levels of

0 dB SNR LP and HP conditions were 70 and 53 dB SPL, respectively. During the first session,

two training lists were presented in the BB-in-noise condition to minimize the known learning

effect of this test (Kollmeier et al., 2015).

For all test lists, subjects were asked to repeat the five-word sentences in a closed, forced-

choice setting (10 options for each word were given). Speech was presented in an adaptive

procedure using a staircase paradigm to determine the speech-reception threshold (SRT) with

a minimal step size of 0.1 dB. The mean signal level or mean SNR of the six last reversals was

used as the SRT for the SPiQ and SPiN-tests, respectively. Lower SRT-values reflect better

speech intelligibility.

AEP MEASUREMENTS

At each session, AEP measurements were performed with the Universal Smart Box (Intelligent

Hearing Systems) using SEPCAM software. Subjects were seated in a comfortable armchair in

a quiet room and were watching a muted video during the recordings. The skin preparation

was performed using NuPrep gel and Ambu Neuroline snap electrodes were placed on vertex,

nasion and bilateral earlobes. The electrode impedances never exceeded 3 kΩ. For one

subject, a mastoid electrode configuration was used due to an immovable earring. All stimuli

were presented using ER-2 probes (Etymotic Research). Both ears were covered with earmuffs

to minimize noise interaction and all inessential electrical devices were turned off during the

measurements. Subjects were instructed to lean their head back and to relax as much as

possible.

ABRs were collected using 5000 alternating polarity sweeps of six stimuli types: broadband

80-µs click stimuli and narrowband 500-ms toneburst (TB) stimuli. Slow click rates of 11 Hz as

well as fast click rates of 120 Hz were both presented at levels of 80 dB and 100 dB peak

equivalent SPL (peSPL). Faster click rates cause minimal neural recovery time and have earlier

been proposed for neurodiagnostic purposes (Valderrama et al., 2012). TBs with center

frequencies of 1 and 4 kHz, which provide more frequency specific information, were

presented at 70 dB SPL at a rate of 20 Hz. The raw signals were stored and processed offline

in Matlab R2018b using the custom-made "sepcam2mat" function. Recorded ABR traces were

bandpass filtered between 100 and 1500 Hz with an 800th order FIR filter "filtfilt" Matlab

function to remove the filter delay. Twenty millisecond epochs were extracted relative to the

stimulus onset and baseline correction was applied by subtracting the mean-value of each

epoch. Two hundred epochs, equal number of each polarity, with the highest peak-to-trough

values were rejected and the remaining 4800 epochs were averaged (Keshishzadeh, Garrett,

& Verhulst, 2020). Mean ABR level series, starting from 100 dB-peSPL and down, were plotted

to visually identify the peaks based on the mean latency intervals by Table 8-1 in Picton (2011).

Accordingly, ABR wave I, III and V peaks and latencies were identified manually and were

confirmed by visual inspection of a second audiologist. **Figure 1** shows exemplar ABRs to 80

and 100 dB peSPL, 11-Hz clicks. Identified peaks are specified with O (80 dB peSPL ABR) and X

(100 dB peSPL ABR). The wave I/V amplitude ratio was defined as the ratio of peak-to-baseline

amplitudes of wave I and wave V, respectively.

EFRs were evoked using 1000 sweeps of two stimulus types. Both stimuli consisted of a 4-kHz

amplitude-modulated tone and are distinguished by their modulation waveform. Specifically,

sinusoidal amplitude modulation (SAM) was applied in the first stimulus and rectangular

amplitude modulation (RAM) in the second (25% duty cycle). The latter was adopted from

Vasilkov et al. (2021), as this stimulus was found to yield strong EFRs which are sensitive to

detect CS in auditory model simulations. Both stimuli had a modulation frequency of 110Hz, a

modulation depth of 100%, and were presented at 70dB SPL. Stimuli had a duration of 500 ms

and were presented at a rate of 2 Hz.

The EFR processing was performed offline in Matlab R2018b. First, recorded EFRs were

bandpass filtered using the same bandpass FIR filter as applied to ABRs, but with low and high

cutoff frequencies of 50 Hz and 5000 Hz, respectively. Then, 400-ms epochs were extracted

from the 100 to 500-ms time-interval, relative to the stimulus onset and 20 epochs with the

highest peak-to-trough values were rejected. The remaining 800 epochs were averaged and

the corresponding magnitude spectrum was constructed using the fast Fourier Transform

(FFT). Additionally, a bootstrapping approach was adopted in the frequency domain to

estimate the noise-floor and variability of the EFR. Exemplar SAM and RAM EFR spectra and

corresponding noise-floors are shown in **Figure 2**A and C, respectively. For detailed

explanation of the bootstrapping procedure see Keshishzadeh, Garrett and Verhulst (2020).

EFR strength was defined as the summation of the signal-to-noise spectral magnitude at

fundamental frequency and the following three harmonics, i.e. 110, 220, 330 and 440 Hz, if

they were above the noise-floor (Vasilkov et al., 2021). Noise-floor corrected SAM and RAM

EFRs are shown in **Figure 2**B and D. Arrows represent the EFR magnitudes at modulation

frequency (110 Hz) and existent spectral peaks at corresponding harmonics.

DATA ANALYSIS

Statistical data analysis was performed in two stages using IBM SPSS Statistics 25. Descriptive

parameters were calculated and tests of normality – i.e. Shapiro-Wilk test, histograms, Q-Q

plots, and box and whisker plots - were applied.

First, the baseline session was analyzed. Hearing-test results of audiometry, speech

audiometry SRT, EFR strength, ABR-amplitudes and -latencies of waves I, III and V and I/V

amplitude ratio were analyzed in relation to lifetime noise-exposure history. Although

Laeq,life was normally distributed as defined by descriptive parameters and the Shapiro-Wilk

test, Spearman's ranked-order correlation coefficients (r_s) were reported as outliers and/or

not-normally distributed test parameters were present in the data for one or more

audiometric frequencies on the one hand, and for one or more test conditions of speech

audiometry, EFR or ABR on the other. Two-tailed significance level was reached for

correlations when $p < 0.05$, suggesting a monotonic and significant relationship between

Laeq,life and the considered hearing-test parameter.

Second, intersession variability was analyzed. As there were outliers and the data was not

normally distributed for some sessions or test conditions, the assumptions for using repeated

measures analysis of variance (ANOVA) were not met and a non-parametric related-samples

Friedman's two-way ANOVA by ranks was conducted to determine statistically significant

changes in all hearing test parameters over sessions. If significance level ($p < 0.05$) was reached,

post-hoc analysis of least-square means was performed for intersession differences between

session 1 on the one hand and session 2, 3 and 4 on the other. The two-tailed significance

level was reached if p<0.017 when applying a Bonferroni correction for multiple comparisons.

For all testing results, the shift from the first to the second session was calculated. Non-

parametric Man-Whitney-U tests were conducted to compare those shifts between groups of

subjects that reported HRS after the musical event and those who did not. Significance level

was reached when p<0.05, suggesting a significant difference between those two groups.

RESULTS PART I

QUESTIONNAIRES

Frequencies of Laeq,life and Laeq,recent are shown in **Figure 3**. Mean Laeq,life was 70.17 dBA

(SD 7.83, range 56.46 – 90.16) when corrected for HPD-use. Attending a discotheque or dance

party were reported as the loudest activity (m 79.5 dBA, SD 8.87, range 60.0 – 90.0), followed

by attending a concert or a festival (m 74.2 dBA, SD 5.07, range 70.0 – 80.0). All but one subject

had attended such events before they participated in this study. Noise-exposure history for

the last three years, as defined by Laeq,recent, had a mean of 66.29 dBA (SD 7.92, range 54.41

– 84.94). Pearson correlation showed a strong positive, statistically significant correlation

between Laeq,recent and Laeq,life (r(20)=0.973, p<0.001). Mean total YANS-score was 2.81

(SD 0.67, range 1.26 – 4.05) and mean total BAHPHL-score was 2.42 (SD 0.47, range 1.50 –

3.38). Twelve subjects (60%) reported commonly wearing HPDs when exposed to loud noises,

five subjects (25%) used customized ear plugs.

BASELINE HEARING STATUS

Distribution of hearing thresholds for conventional frequencies and EHF of the test ear are

shown in **Figure 4**. The mean pure-tone average of 3, 4, 6 and 8 kHz (PTA3-8kHz) was 4.88dB

HL (SD 2.75, median 5.00, range -1.25 – 8.75) and the mean pure-tone average of 10, 12.5, 14

and 16 kHz (PTA10-16kHz) was 5.94 (SD 9.59, median 2.50, range -10.00 – 25.00). Neither

PTA3-8kHz nor PTA10-16kHz significantly correlated to lifetime noise-exposure history

(p>0.05). Spearman's correlation coefficients ranged from -0.28 to 0.09.

BASELINE SPEECH AUDIOMETRY

Figure 5A and B represent the SRT-values of SPiQ and SPiN respectively. No monotonic relation

was found between SPiQ and Laeq,life, as Spearman's correlation coefficients were 0.24

(p=0.30) for the LP filtered speech and -0.20 (p=0.41) for the HP filtered speech, and both were

statistically insignificant. Spearman's correlations of SPiN to Laeq,life showed weak positive

correlation coefficients, ranging from 0.02 to 0.44 with the highest correlation coefficient for

the LP condition. However, none of the correlation coefficients were statistically significant

(p>0.05).

BASELINE AEP-MEASUREMENTS

ABR latencies and amplitudes of waves I, III and V in the baseline session are given in Table 1.

As expected from literature, latencies of all waves increased with lowering the stimulus level

(i.e. 100dB peSPL vs 80dB peSPL), faster stimulus presentation (i.e. 120 Hz vs 11 Hz) or

narrower stimulus-spectrum (i.e. TB vs click stimuli; (Picton, 2011)). A statistically significant

relation with Laeq,life was found for the ABR wave V latency using a 11Hz 100dB click stimulus

(rs=0.66, p=0.002). No other significant relations of ABR latency with Laeq,life were found.

Spearman's correlation coefficients ranged from 0.02 (p=0.92) to 0.20 (p=0.40) for wave I,

from 0.01 (p=0.99) to 0.37 (p=0.11) for wave III and from 0.02 (p=0.93) to 0.37 (p=0.11) for

wave V latencies.

Second, wave I, III and V ABR amplitudes showed no statistically significant monotonic relation

to Laeq,life. For wave I, Spearman's correlation coefficients of the different stimulus types

ranged from -0.16 (p=0.51) to 0.14 (p=0.56), for wave III from -0.34 (p=0.15) to 0.27 (p=0.25)

and for wave V from -0.25 (p=0.29) to 0.003 (p=0.99).

Finally, no significant correlation was found between the ABR I/V amplitude ratio and Laeq,life

in all conditions. Spearman's correlation coefficients ranged from -0.41 (p=0.87) to 0.32

(p=0.18) for the various ABR stimuli.

EFR strength in relation to Laeq,life is shown in **Figure 6**. For all but one subject, higher EFR-

strength was found using the RAM (mean 0.085, SD 0.029, median 0.088, range 0.007 – 0.136)

compared to the SAM-stimulus (mean 0.030, SD 0.010, median 0.028, range 0.012 – 0.056),

as was predicted by auditory models (Vasilkov et al., 2021). A negative, but weak and

statistically non-significant Spearman's correlation coefficient was found in relation to

Laeq,life for SAM-stimulus (r_s=-0.34, p=0.15) as well as for RAM-stimulus (r_s=-0.17, p=0.49).

RESULTS PART II

QUESTIONNAIRE

The mean duration of the music festival attendance was 2.68 days (SD 1.29, range 1 - 5), where

subjects self-reported to be exposed to music 8.9 hours a day (SD = 2.39, range 5 - 12).

Estimation of noise-exposure levels during the festival, HPD-attenuation taken into account,

resulted in an average Laeq,w,festival of 76.2 dBA (SD 7.82, range 59.35 – 88.19).

Thirteen subjects (68.4%) reported to have worn HPDs during the music festivals and only 7

participants (36.8%) had worn them 50% of the time during noise exposure. Eight subjects

(42.1%) experienced HRS after the noise exposure: six subjects (31.6%) experienced dullness

after the music event, 2 persons (10.5%) experienced a subjective hearing loss and 3 persons

(15.8%) reported tinnitus. Dullness and tinnitus were experienced both unilaterally or

bilaterally and disappeared after a maximum of 24 hours after the noise exposure, whereas a

subjective hearing loss was experienced in both ears and disappeared after 30 minutes. Only

one of the test subjects that had worn HPDs more than 50% of the noise exposure time

reported HRS. Baseline testing results of PTA, SPiQ and SPiN, and EFR did not differ

significantly between subjects who reported HRS after the event, and subjects who did not.

ABR results only showed between-group differences in the rate of 120 Hz, 100 dB click

condition only. For this condition, the group with HRS had a significantly (p=0.01) smaller P3

latency (m 5.30 ms, range 4.60 – 5.50) than the group without HRS (m 5.70 ms, range 5.20 –

6.10). Second, wave-I amplitudes were significantly (p=0.005) higher in the group with HRS (m

0.13 μV, range 0.02 – 0.22) than the group without (m 0.06 μV, range 0.03 – 0.09). Finally, a

significantly higher I/V amplitude ratio was found in the HRS group (m 0.40, range 0.27 – 1.55)

compared to the non-HRS group (m 0.19, range 0.09 – 0.49).

LONGITUDINAL ANALYSIS HEARING STATUS

In the second session, i.e. one day after noise exposure, a TTS was only found in one subject

(0.05%). Specifically, this person had an average threshold elevation of 10 dB or more on 2, 3

and 4 kHz, as described by Occupational Safety and Health Administration standards (OSHA,

1974). For this subject, an average elevation of 11.7 dB was found in session 2 compared to

session 1, which recovered to values below 10 dB in sessions 3 and 4. A residual shift of 6.7dB

from session 1 was present in the fourth session. This subject experienced bilaterally tinnitus

(24h) but reported no subjective hearing loss nor dullness. All other subjects had an average

threshold shift from session 1 to session 2 at 2, 3 and 4 kHz ranging from -3.3 to 5.0 dB. At

single conventional audiogram frequencies, a threshold elevation of 10 dB or more was found

in eight subjects (42.1%) during the second session. At extended high frequencies, threshold

shifts of 10 dB or higher were found in nine subjects (47.3%) between session 1 and session 2.

All frequencies included, a threshold shift of 10 dB or more was found in 15 subjects (78.9%).

Pure tone audiometry and χ^2 results are displayed in Table 2. Significant intersession

differences were found for auditory thresholds of 500 Hz (p=0.033) and 10 kHz (p=0.005).

However, post-hoc analysis with Bonferroni adjustment showed no significant threshold shift

from the first session to a follow-up session (p>0.017). PTAs revealed statistically insignificant

between-session differences (p>0.05). Even though subjects who experienced HRS showed a

median positive shift, i.e. higher values, in PTA3-8kHz (3.13 dB HL, range -2.50 – 10.00) and

PTA10-16kHz (3. 75 dB HL, range -3.75 – 6.25) in the second session and subjects without HRS

did not (respectively 0.00 dB HL, range -3.75 – 3.75 and 0.00 dB HL, range -6.25 – 11.2),

between-group differences were not significant (p=0.18 and 0.44, respectively).

LONGITUDINAL ANALYSIS SPEECH AUDIOMETRY

Figure 7A displays the results of the SPiQ test. SPiQ showed slightly increased SRT-values from

session 1 to session 2 for both LP- and HP-filtered speech conditions. Friedman's ANOVA

showed statistically significant intersession differences only for the HP filtered condition

(χ^2(3)=29.46, p<0.001). Although post-hoc analysis with Bonferroni-correction showed

statistically insignificant changes between session 1 and session 2, a significant improvement

(p=0.008) in speech intelligibility from session 1 (med =22.67, IQR 8.04) to session 4

(med=20.78, IQR 5.28) was found.

The same trend of improved speech intelligibility over sessions was found for SPiN, as shown

in **Figure 7**B. Friedman's ANOVA showed statistically significant intersession differences for

the BB as well as the LP condition (χ^2(3)=29.72, p<0.001) and the HP condition (χ^2(3)=27.97,

p<0.001). Post-hoc pairwise comparisons with Bonferroni-correction showed a significant

improvement of SNR from session 1 to 4 for all conditions (BB p=0.002, LP and HP p<0.001)

and a significant improvement from session 1 to 3 for the LP condition (p=0.006). However,

no statistical significant differences between session 1 and session 2 were found for all test

conditions.

Between-group comparisons of persons with or without HRS, did not show significant

differences in SRT-shift from session1 to session 2 for both SPiQ (p range 0.11 – 0.41) and SPiN

(p range 0.06 – 0.77).

LONGITUDINAL ANALYSIS AEP-MEASUREMENTS

Table 3 represents all results and χ^2-values of AEP-measurements.

ABR-latencies showed statistically insignificant changes between sessions for any wave or

stimulus type. ABR wave III showed significant intersession differences only for the 11-Hz 100-

dB peSPL stimulus (χ^2(3)=7.93, p=0.48) and wave V changed significantly over time for the 11-

Hz 80-dB peSPL stimulus (χ^2(3)=12.92, p=0.005), but post-hoc comparisons with Bonferroni

correction showed no significant differences between the first session and sessions 2, 3 or 4

(p>0.017). Amplitude ratios I/V did not reveal any significant intersession changes for the

considered stimulus types (p>0.05).

EFR strength of RAM stimuli was consistently higher than that of SAM-stimuli. Even though

median EFR strength showed a slight reduction from the first to the second session for both

SAM and RAM stimuli, EFR strength did not significantly change across session (p>0.05).

ABR-shifts from session 1 to session 2, compared between groups who did or did not

experience HRS, respectively, can be found in Table 4. Shifts in ABR latencies, amplitudes and

I/V amplitude showed some diffuse significant group differences for certain stimuli, e.g. ABR

wave I latency shift for 11-Hz 80-dB click stimulus (p=0.02), wave-III latency (p=0.04), ABR

wave-I (p=0.04) and wave-V (p=0.02) amplitude shift for TB 4-kHz stimulus, and I/V-amplitude

ratio for the 120-Hz 100-dB stimulus (p=0.03). However, no consistent trends across different

stimulus types were observed.

EFR-strength showed a negative shift, i.e. an EFR-strength reduction from session 1 to session

2 for both stimulus types within the group that reported HRS (SAM: med = -0.004 range = -

0.03 - 0.02; RAM: med = -0.007; range = -0.06 – 0.05). In the group without HRS, a negative

shift was only found with the SAM stimulus (med = -0.006; range = -0.01 – 0.1), but not for the

RAM stimulus (med = 0.008; range = -0.06 – 0.03). No significant group differences were found

between groups with or without HRS in EFR-strength-shifts, neither for SAM (U=49.00, p=0.71)

nor for RAM-stimuli (U=52.00, p=0.88).

DISCUSSION

PART I

The first aim of this study consisted of a baseline evaluation of hearing status in relation to the

noise-exposure history of 19 young, normal-hearing adults. Our inclusion criteria, specifically

normal auditory thresholds and the absence of HRS, eliminated ears with OHC-dysfunction as

much as possible.

As other studies failed to find a relation between young adults' hearing thresholds or

distortion product otoacoustic emissions (DPOAEs) and their lifetime noise-exposure history

(Degeest, Clays, et al., 2017; Keppler, Dhooge, et al., 2015b; Wei et al., 2017; Williams et al.,

2015), no substantial OHC-loss due to noise exposure was expected in this population. Indeed,

PTA10-16kHz was never worse than 25 dB HL and showed no significant correlation with

lifetime noise-exposure history, indicating relatively intact functioning of the OHCs. The

assumption of a normal functioning cochlear amplifier facilitates interpretation of the other

hearing test results in the context of hidden hearing loss or CS.

CS has been thought to influence suprathreshold hearing function and to affect speech

understanding, especially in difficult listening situations (Kujawa & Liberman, 2009; Lin et al.,

2011). Therefore, it could be hypothesized that SPiQ- or SPiN-metrics would relate to lifetime

noise-exposure history if CS was present. However, no relation was found in this study

population. This supports the results of the majority of studies investigating speech

intelligibility in relation to noise-exposure history of normal-hearing persons, where no

relation was found either (Couth et al., 2020; Fulbright et al., 2017; Grinn et al., 2017; Guest

et al., 2018; Prendergast, Millman, et al., 2017; Yeend et al., 2017). In contrast, some studies

have reported a link between noise-exposure history and speech intelligibility (Hope et al.,

2013; Kumar et al., 2012; Liberman et al., 2016). The inclusion of subjects that were regularly

exposed to occupational noise or a music education could have contributed to a higher noise-

exposure dose in these latter studies and increased the risk of CS in the tested population.

However, OHC damage could not be ruled out in all these studies since, respectively,

audiometric thresholds were only measured up to 4kHz (Hope et al., 2013), or speech

intelligibility deterioration was accompanied with audiometric loss in EHFs (Liberman et al.,

2016). A recent study by Prendergast et al. (2019) found a link between noise-exposure history

and speech intelligibility albeit showing a better performance in subjects with higher lifetime

noise-exposure history at levels above 85 dBA. As many of their highly-exposed subjects

worked in the music industry, listening strategies and attention training on complex

soundscapes are proposed by the authors to explain this difference.

The baseline AEP-measurements in the present study did not show significant correlations to

lifetime noise-exposure history. First, ABR wave-I, -III and -V amplitudes and the ABR I/V

amplitude ratio were not significantly correlated to noise-exposure history. These

observations corroborate results from a number of human studies that did not show a clear

link between ABR wave-I amplitudes and recreational noise exposure (Couth et al., 2020;

Fulbright et al., 2017; Grinn et al., 2017; Liberman et al., 2016; Prendergast et al., 2018; Ridley

et al., 2018; Skoe & Tufts, 2018; Spankovich, Le Prell, et al., 2017). However, two studies

performed by Bramhall et al. (2017) and Stamper and Johnson (2015a) revealed significant

wave-I amplitude reductions in subjects with higher noise-exposure history, although for the

latter study this was only true for female subjects (Stamper & Johnson, 2015b). Furthermore,

Valderrama et al. (2018) reported reduced wave-I amplitudes in relation to recreational and

occupational noise exposure, but this relation was not significant after removal of outliers.

Amplitudes of wave III have less commonly been analyzed and have not shown significant

relations to noise exposure (Ridley et al., 2018; Skoe & Tufts, 2018). Furthermore, wave-V

amplitudes did not show a relation to recreational noise exposure in literature (Prendergast

et al., 2018; Ridley et al., 2018; Skoe & Tufts, 2018; Spankovich, Bishop, et al., 2017) and

neither did I/V amplitude ratios (Prendergast, Guest, et al., 2017; Spankovich, Bishop, et al.,

2017). One study did report a reduction in I/V ratios in subjects that frequently attended music

concerts (Grose et al., 2019). On the contrary, a recent study by Couth et al. (2020) showed

higher I/V-amplitude ratios in musicians compared to non-musicians. Even though noise-

exposure history was similar between both groups in that study and noise-exposure history

did not show a significant relation to I/V amplitude, the highly-exposed subjects showed OHC-

dysfunction.

Second, regarding ABR latencies, no relation to noise-exposure history was found in the

present study. This is in line with several studies, where ABR wave-I latencies (Couth et al.,

2020; Spankovich, Bishop, et al., 2017) or wave-V latencies (Spankovich, Bishop, et al., 2017)

did not relate to recreational noise exposure. On the contrary, Skoe and Tufts (2018) did find

a delay in waves I, III and V for subjects with a higher noise-exposure based on one-week

dosimetry measurements. The more highly-exposed subjects in this study were mostly

students who participated in music ensembles on campus. A delay in wave V latency with

increasing noise exposure was also found by Couth et al. (2020), but only for males, and by

Prendergast, Guest, et al. (2017) where significance disappeared after correcting for age.

Finally, EFR strength did not correlate to lifetime noise-exposure history in the present study.

This is in line with other studies in normal-hearing subjects (Grose et al., 2017) and in normal-

hearing subjects with tinnitus (Guest et al., 2017). Prendergast, Guest, et al. (2017) found a

significant negative correlation between EFR strength and noise exposure. However, this was

only true for males and after controlling for age, this correlation was no longer significant. A

study by the same lab could not find noise exposure as a predictor of EFR- strength

(Prendergast et al., 2019). In contrast, animal studies (Möhrle et al., 2016; Parthasarathy and

Kujawa, 2018; Shaheen et al., 2015) and those that used computational modelling of the

auditory periphery or human data have shown the possible value of EFRs in diagnosing CS

(Bharadwaj et al., 2015; Garrett & Verhulst, 2019; Keshishzadeh, Garrett, Vasilkov, et al., 2020;

Paul et al., 2017; Vasilkov et al., 2021).

As described above, studies trying to demonstrate the presence of CS in humans by relating

possible biomarkers to noise-exposure history, e.g. SPiN, ABR-amplitudes or amplitude ratios,

ABR-latencies or EFR strength, report contradictory results (see Bramhall et al., 2019 for

review). The disagreement between those studies can be caused by multiple reasons. First,

study population characteristics such as age vary across studies. Age-related CS could be

dominant over noise-induced CS and therefore hiding the effects of noise-exposure history.

The use of different age ranges between studies implies that in some study cohorts mostly

occupational noise exposure was studied (Hope et al., 2013; Kumar et al., 2012; Prendergast

et al., 2019; Valderrama et al., 2018), whereas other studies focused on recreational noise

exposure. Second, methodological differences could lead to different conclusions. Based on

(the absence of) EHF audiometry- or OAE-results, OHC-deficits cannot be ruled out in some

study cohorts (Liberman et al., 2016). Furthermore, methods to calculate or measure noise-

exposure history differ between studies. As objective measurements of noise exposure such

as dosimetry is hard to be collected during a lifetime period, subjective and retrospective

questionnaires or interviews are used to estimate noise-exposure history. Beach et al. (2012)

stated that realistic loudness estimations can be made through/via questionnaires about

recent noise exposures when compared to dosimetry metrics. However, questionnaires

mostly do not take into account frequency components of noise or acoustic traumas caused

by impulse noises. Other factors such as blood type (Doğru et al., 2003), nutrition (Kopke et

al., 2005), fitness (Kolkhorst et al., 1998), smoking (Ferrite & Santana, 2005; Pouryaghoub et

al., 2007), alcohol use (Kraaijenga et al., 2018; Upile et al., 2007) and drug use (Church et al.,

2013) have been suggested to influence susceptibility to NIHL, but have not been taken into

account in studies.

PART II

The second part of this study evaluated longitudinal biomarker changes before and after

attending a music festival.

Eight subjects (42.1%) experienced HRS after the music festival, comparable to Mercier et al.

(2003), where 36% experienced HRS after attending a music festival. It has to be noted that

the majority of subjects wore HPDs, at least for some time during the concerts. However, of

those subjects who reported using HPDs, only a few wore them consistently. Only one of the

subjects who wore HPDs over 50% of the time during noise exposure experienced HRS, which

is in line with findings that HPDs reduce HRS (Kraaijenga et al., 2018) or reduce TTS (Ramakers

et al., 2016). Baseline testing results of subjects that reported HRS after the event, did not

significantly differ from those who did not, suggesting that cross-sectional use of the present

test battery does not predict the risk of developing symptoms such as tinnitus after noise

exposure.

A TTS as defined by OSHA-standards (OSHA, 1974) was only found in one subject one day after

the music festival. No significant group changes in auditory thresholds were found for any of

the tested frequencies. The difficulty of detecting a TTS in non-laboratory situations such as

attending concerts or nightclubs was earlier described (Le Prell et al., 2012). The experienced

noise-exposure is known to vary considerably between subjects attending such events, as

noise intensity levels vary depending on distance relative to the stage and the music genre

(Opperman et al., 2006). Furthermore, it is difficult to distinguish a small TTS from test-retest

variability (Le Prell et al., 2012; Schlauch & Carney, 2012), which is amongst others dependent

on participant experience and motivation (Schlauch & Carney, 2012). The observations in the

above studies contrast findings from other studies that did not observe a TTS in the majority

of subjects after attending concerts, festivals or music venues (Emmerich et al., 2002; Le Prell,

2019; Opperman et al., 2006; Ramakers et al., 2016). These studies evaluated hearing

thresholds immediately after noise exposure, whereas the greatest auditory-threshold

recovery was found to occur within two-to-four hours after the noise exposure (Emmerich et

al., 2002; Grinn et al., 2017; Le Prell et al., 2012). Therefore the possibility of a recovered TTS

in the present study population should be considered, especially in those subjects that

experienced HRS (Masterson et al., 2016; Schmuzigert et al., 2006).

SPiQ and SPiN tests showed no significant differences one day after noise exposure compared

to the baseline session. This finding is in contrast with the study of Grinn et al. (2017), where

the noise dose of a noise exposure event has been related to a decrease in speech in noise

intelligibility. However, this study showed a significant improvement in SRT-values for session

4 compared to session 1 for each condition, which could be related to a learning effect. Two

training lists were used as suggested in literature concerning the Flemish Matrix speech

material to overcome the largest training effect. However, smaller and non-significant

improvement in SRT-values are described even after the presentation of two training lists

(Kollmeier et al., 2015). As a consequence of this training effect, possible deteriorations in

speech intelligibility shortly after noise exposure could remain undetected.

No significant changes could be found between the first session and sessions 2, 3 or 4 for AEP-

measurements, as determined by analysis of ABR amplitudes and latencies of wave I, III and V

and ABR I/V amplitude ratio and EFR-amplitudes. Grinn et al. (2017), who measured ABR wave

I amplitudes by conducting electrocochleography, did not find amplitude reductions after

attending a noisy event.

DETECTING AND MONITORING CS IN NORMAL HEARING SUBJECTS

In the present study, no relation could be found between noise-exposure history and the

possible biomarkers of CS, nor did those biomarkers change over sessions surrounding a music

event in normal hearing young adults. Baseline testing results and shifts from session 1 to

session 2 did not differ between subjects that experienced HRS after the music event, and

those who did not. Thus, no strong conclusions can be made regarding the presence of noise-

induced CS in relation to noise-exposure history nor regarding the appearance of CS shortly

after visiting a music festival in this cohort of young normal-hearing subjects.

Possibly, the noise-exposure history and recreational noise exposure dose (76.2 dBA) in the

present study were not sufficiently high to cause CS. Permanent consequences of recreational

noise on hearing has been doubted before, as reviewed by Carter et al. (2014). This theory is

reflected by the lack of TTS in all but one subjects at one day after the festival. In noise-

exposed animals, smaller TTSs of 20 – 30 dB, measured at one day after a single noise

exposure, were shown not to result in CS (Fernandez et al., 2015; Jensen et al., 2015) whereas

TTSs up to 50 dB were reported for noise-exposed animals with substantial CS (Furman et al.,

2013; Kujawa & Liberman, 2009; Lin et al., 2011). It may hence be that the young normal-

hearing listeners we considered did not suffer from CS.

Second, it is possible that the biomarkers for CS are not sensitive or specific enough to detect

or to monitor CS in normal-hearing young adults. The interpretation of speech in noise

audiometry is complicated by a possible training effect. Confounding factors in AEP-

measurements include amongst others intersubject variability due to head size and sex

(Mitchell et al., 1989) and intrasubject variability due to different degrees of awakeness (part

II, Pethe, Begall et al., 1996).

In order to distinguish between the absence of CS in the study cohort and a lack of specificity

in the CS biomarkers, future studies may provide more insight. First, it would be interesting to

further investigate intersubject variability and test-retest reliability of the biomarkers of CS in

a cohort of normal-hearing subjects. Second, the use of dosimetry, where noise doses can be

measured for specific noise events, logging of HPD-use during a music event, and hearing

process evaluation immediately after the event will provide the needed information regarding

recent noise exposure doses, HPD-use and temporary versus long-term effects on hearing.

Third, as AEP-measurements can be conducted with different stimulus parameters and rates,

their sensitivity to detect CS can be further investigated by further comparing their results

within the same study cohorts (Vasilkov et al., 2020; Wilson et al., 2020). Finally, larger test

groups will allow us to compare different possibly confounding factors such as the use of HPD

and different levels of noise-exposure history.

ACKNOWLEDGMENT
This work was supported by UGent BOF-IOP project: "Portable Hearing Diagnostics:

Monitoring of Auditory-nerve Integrity after Noise Exposure (EarDiMon)" (T.V.M., N.D.P., I.D.,

H.K. and S.V.) and European Research Council (ERC) under the Horizon 2020 Research and

Innovation Program, grant agreement no. 678120 RobSpear (SK and SV).

REFERENCES

Beach, E. F., Williams, W., & Gilliver, M. (2012). The objective-subjective assessment of noise: Young

adults can estimate loudness of events and lifestyle noise. *International journal of audiology*,

51(6), 444-449. https://doi.org/10.3109/14992027.2012.658971

Bharadwaj, H. M., Masud, S., Mehraei, G., Verhulst, S., & Shinn-Cunningham, B. G. (2015). Individual

differences reveal correlates of hidden hearing deficits. *The Journal of neuroscience : the*

official journal of the Society for Neuroscience, *35*(5), 2161-2172.

https://doi.org/10.1523/JNEUROSCI.3915-14.2015

Bharadwaj, H. M., Verhulst, S., Shaheen, L., Liberman, M. C., & Shinn-Cunningham, B. G. (2014).

Cochlear neuropathy and the coding of supra-threshold sound. *Frontiers in systems*

neuroscience, *8*, 26. https://doi.org/10.3389/fnsys.2014.00026

Bramhall, N., Beach, E. F., Epp, B., Le Prell, C. G., Lopez-Poveda, E. A., Plack, C. J., Schaette, R., Verhulst,

S., & Canlon, B. (2019). The search for noise-induced cochlear synaptopathy in humans:

Mission impossible? *Hearing research*, *377*, 88-103.

https://doi.org/10.1016/j.heares.2019.02.016

Bramhall, N. F., Konrad-Martin, D., & McMillan, G. P. (2018). Tinnitus and Auditory Perception After a

History of Noise Exposure: Relationship to Auditory Brainstem Response Measures [Article].

Ear and hearing, *39*(5), 881-894. https://doi.org/10.1097/AUD.0000000000000544

Bramhall, N. F., Konrad-Martin, D., McMillan, G. P., & Griest, S. E. (2017). Auditory Brainstem Response

Altered in Humans With Noise Exposure Despite Normal Outer Hair Cell Function [Article]. *Ear*

and hearing, *38*(1), e1-e12.

http://www.embase.com/search/results?subaction=viewrecord&from=export&id=L6208749

33

Carter, L., Williams, W., Black, D., & Bundy, A. (2014). The leisure-noise dilemma: hearing loss or

hearsay? What does the literature tell us? *Ear and hearing*, *35*(5), 491-505.

https://doi.org/10.1097/01.aud.0000451498.92871.20

Church, M. W., Zhang, J. S., Langford, M. M., & Perrine, S. A. (2013). 'Ecstasy' enhances noise-induced

hearing loss. *Hearing research*, *302*, 96-106. https://doi.org/10.1016/j.heares.2013.05.007

Clark, W. W. (1991). Recent studies of temporary threshold shift (TTS) and permanent threshold shift

(PTS) in animals. *The Journal of the Acoustical Society of America*, *90*(1), 155-163.

https://doi.org/10.1121/1.401309

Couth, S., Prendergast, G., Guest, H., Munro, K. J., Moore, D. R., Plack, C. J., Ginsborg, J., & Dawes, P.

(2020). Investigating the effects of noise exposure on self-report, behavioral and

electrophysiological indices of hearing damage in musicians with normal audiometric

thresholds. *Hearing research*, 108021. https://doi.org/10.1016/j.heares.2020.108021

Degeest, S., Clays, E., Corthals, P., & Keppler, H. (2017). Epidemiology and Risk Factors for Leisure

Noise-Induced Hearing Damage in Flemish Young Adults. *Noise & health*, *19*(86), 10-19.

https://doi.org/10.4103/1463-1741.199241

Degeest, S., Keppler, H., Corthals, P., & Clays, E. (2017). Epidemiology and risk factors for tinnitus after

leisure noise exposure in Flemish young adults. *International journal of audiology*, *56*(2), 121-

129. https://doi.org/10.1080/14992027.2016.1236416

Derebery, M. J., Vermiglio, A., Berliner, K. I., Potthoff, M., & Holguin, K. (2012). Facing the music: pre-

and postconcert assessment of hearing in teenagers. *Otology & neurotology : official*

publication of the American Otological Society, American Neurotology Society [and] European

Academy of Otology and Neurotology, *33*(7), 1136-1141.

https://doi.org/10.1097/MAO.0b013e31825f2328

Doğru, H., Tüz, M., & Uygur, K. (2003). Correlation between blood group and noise-induced hearing

loss. *Acta oto-laryngologica*, *123*(8), 941-942.

Emmerich, E., Richter, F., Hagner, H., Giessler, F., Gehrlein, S., & Dieroff, H. G. (2002). Effects of

discotheque music on audiometric results and central acoustic evoked neuromagnetic

responses. *The international tinnitus journal*, *8*(1), 13-19.

Fernandez, K. A., Jeffers, P. W., Lall, K., Liberman, M. C., & Kujawa, S. G. (2015). Aging after noise
exposure: acceleration of cochlear synaptopathy in "recovered" ears. *The Journal of*
neuroscience : the official journal of the Society for Neuroscience, *35*(19), 7509-7520.
https://doi.org/10.1523/jneurosci.5138-14.2015

Ferrite, S., & Santana, V. (2005). Joint effects of smoking, noise exposure and age on hearing loss.
Occupational   medicine   (Oxford,   England),   *55*(1),   48-53.
https://doi.org/10.1093/occmed/kqi002

Francart, T., van Wieringen, A., & Wouters, J. (2008). APEX 3: a multi-purpose test platform for auditory
psychophysical experiments. *Journal of neuroscience methods*, *172*(2), 283-293.
https://doi.org/10.1016/j.jneumeth.2008.04.020

Fulbright, A. N. C., Le Prell, C. G., Griffiths, S. K., & Lobarinas, E. (2017). Effects of Recreational Noise on
Threshold and Suprathreshold Measures of Auditory Function. *Seminars in hearing*, *38*(4), 298-
318. https://doi.org/10.1055/s-0037-1606325

Furman, A. C., Kujawa, S. G., & Liberman, M. C. (2013). Noise-induced cochlear neuropathy is selective
for fibers with low spontaneous rates. *Journal of neurophysiology*, *110*(3), 577-586.
https://doi.org/10.1152/jn.00164.2013

Garrett, M., Vasilkov, V., Mauermann, M., Wilson, J. L., Henry, K. S., & Verhulst, S. (2020). Speech-in-
noise intelligibility difficulties with age: the role of cochlear synaptopathy. *bioRxiv*,
2020.2006.2009.142950. https://doi.org/10.1101/2020.06.09.142950

Garrett, M., & Verhulst, S. (2019). Applicability of subcortical EEG metrics of synaptopathy to older
listeners with impaired audiograms. *Hearing research*, *380*, 150-165.
https://doi.org/10.1016/j.heares.2019.07.001

Grinn, S. K., Wiseman, K. B., Baker, J. A., & Le Prell, C. G. (2017). Hidden Hearing Loss? No Effect of
Common Recreational Noise Exposure on Cochlear Nerve Response Amplitude in Humans.
Frontiers in neuroscience, *11*, 465. https://doi.org/10.3389/fnins.2017.00465

Grose, J. H., Buss, E., & Elmore, H. (2019). Age-Related Changes in the Auditory Brainstem Response

and Suprathreshold Processing of Temporal and Spectral Modulation. *Trends in hearing*, *23*,

2331216519839615. https://doi.org/10.1177/2331216519839615

Grose, J. H., Buss, E., & Hall, J. W. (2017). Loud Music Exposure and Cochlear Synaptopathy in Young

Adults: Isolated Auditory Brainstem Response Effects but No Perceptual Consequences

[Article].   *Trends   in   hearing*,   *21*,   2331216517737417.

https://doi.org/10.1177/2331216517737417

Guest, H., Munro, K. J., & Plack, C. J. (2017). Tinnitus with a normal audiogram: Role of high-frequency

sensitivity and reanalysis of brainstem-response measures to avoid audiometric over-matching

[Letter]. *Hearing research*, *356*, 116-117. https://doi.org/10.1016/j.heares.2017.10.002

Guest, H., Munro, K. J., Prendergast, G., Millman, R. E., & Plack, C. J. (2018). Impaired speech perception

in noise with a normal audiogram: No evidence for cochlear synaptopathy and no relation to

lifetime   noise   exposure   [Article].   *Hearing   research*,   *364*,   142-151.

https://doi.org/10.1016/j.heares.2018.03.008

Hakuba, N., Koga, K., Gyo, K., Usami, S. I., & Tanaka, K. (2000). Exacerbation of noise-induced hearing

loss in mice lacking the glutamate transporter GLAST. *The Journal of neuroscience : the official*

journal   of   the   Society   for   Neuroscience,   *20*(23),   8750-8753.

https://www.ncbi.nlm.nih.gov/pubmed/11102482

Hall,   J.   W.   (2007).   *New   Handbook   of   Auditory   Evoked   Responses*.   Pearson.

https://books.google.be/books?id=DpZrQgAACAAJ

Henry, K. S., Kale, S., & Heinz, M. G. (2016). Distorted Tonotopic Coding of Temporal Envelope and Fine

Structure with Noise-Induced Hearing Loss. *The Journal of neuroscience : the official journal of*

the Society for Neuroscience, *36*(7), 2227-2237. https://doi.org/10.1523/jneurosci.3944-

15.2016

Hope, A. J., Luxon, L. M., & Bamiou, D. E. (2013). Effects of chronic noise exposure on speech-in-noise

perception in the presence of normal audiometry. *The Journal of laryngology and otology*,

127(3), 233-238. https://doi.org/10.1017/S002221511200299X

Jensen, J. B., Lysaght, A. C., Liberman, M. C., Qvortrup, K., & Stankovic, K. M. (2015). Immediate and

delayed cochlear neuropathy after noise exposure in pubescent mice [Article]. *PLoS ONE*,

10(5). https://doi.org/10.1371/journal.pone.0125160

Jokitulppo, J., Toivonen, M., & Bjork, E. (2006). Estimated leisure-time noise exposure, hearing

thresholds, and hearing symptoms of Finnish conscripts. *Military medicine*, *171*(2), 112-116.

https://www.ncbi.nlm.nih.gov/pubmed/16578978

Keppler, H., Dhooge, I., Degeest, S., & Vinck, B. (2015). The effects of a hearing education program on

recreational noise exposure, attitudes and beliefs toward noise, hearing loss, and hearing

protector devices in young adults. *Noise & health*, *17*(78), 253-262.

https://doi.org/10.4103/1463-1741.165028

Keppler, H., Dhooge, I., & Vinck, B. (2015a). Hearing in young adults. Part I: The effects of attitudes and

beliefs toward noise, hearing loss, and hearing protector devices. *Noise & health*, *17*(78), 237-

244. https://doi.org/10.4103/1463-1741.165024

Keppler, H., Dhooge, I., & Vinck, B. (2015b). Hearing in young adults. Part II: The effects of recreational

noise exposure. *Noise & health*, *17*(78), 245-252. https://doi.org/10.4103/1463-1741.165026

Keppler, H., & Dhooge, I. J. M. c. (2010). *Optimization of the diagnosis of noise-induced hearing loss*

with otoacoustic emissions 2010.]. http://lib.ugent.be/catalog/rug01:001438088

Keshishzadeh, S., Garrett, M., Vasilkov, V., & Verhulst, S. (2020). The derived-band envelope following

response and its sensitivity to sensorineural hearing deficits. *Hearing research*, *392*, 107979.

https://doi.org/https://doi.org/10.1016/j.heares.2020.107979

Keshishzadeh, S., Garrett, M., & Verhulst, S. (2020). Towards personalized auditory models: predicting

individual sensorineural-hearing-loss profiles from recorded human auditory physiology.

bioRxiv.

Kolkhorst, F. W., Smaldino, J. J., Wolf, S. C., Battani, L. R., Plakke, B. L., Huddleston, S., & Hensley, L. D.

(1998). Influence of fitness on susceptibility to noise-induced temporary threshold shift.

Medicine and science in sports and exercise, *30*(2), 289-293.

https://www.ncbi.nlm.nih.gov/pubmed/9502359

Kollmeier, B., Warzybok, A., Hochmuth, S., Zokoll, M. A., Uslar, V., Brand, T., & Wagener, K. C. (2015).

The multilingual matrix test: Principles, applications, and comparison across languages: A

review. *International journal of audiology*, *54 Suppl 2*, 3-16.

https://doi.org/10.3109/14992027.2015.1020971

Konrad-Martin, D., Dille, M. F., McMillan, G., Griest, S., McDermott, D., Fausti, S. A., & Austin, D. F.

(2012). Age-related changes in the auditory brainstem response. *Journal of the American*

Academy of Audiology, *23*(1), 18-35; quiz 74-15. https://doi.org/10.3766/jaaa.23.1.3

Kopke, R., Bielefeld, E., Liu, J., Zheng, J., Jackson, R., Henderson, D., & Coleman, J. K. (2005). Prevention

of impulse noise-induced hearing loss with antioxidants. *Acta oto-laryngologica*, *125*(3), 235-

243. https://www.ncbi.nlm.nih.gov/pubmed/15966690

Kraaijenga, V. J. C., van Munster, J., & van Zanten, G. A. (2018). Association of Behavior With Noise-

Induced Hearing Loss Among Attendees of an Outdoor Music Festival: A Secondary Analysis of

a Randomized Clinical Trial. *JAMA otolaryngology-- head & neck surgery*, *144*(6), 490-497.

https://doi.org/10.1001/jamaoto.2018.0272

Kujawa, S. G., & Liberman, M. C. (2009). Adding insult to injury: cochlear nerve degeneration after

"temporary" noise-induced hearing loss. *The Journal of neuroscience : the official journal of*

the Society for Neuroscience, *29*(45), 14077-14085. https://doi.org/10.1523/JNEUROSCI.2845-

09.2009

Kujawa, S. G., & Liberman, M. C. (2015). Synaptopathy in the noise-exposed and aging cochlea: Primary

neural degeneration in acquired sensorineural hearing loss. *Hearing research*, *330*(Pt B), 191-

199. https://doi.org/10.1016/j.heares.2015.02.009

Kumar, U. A., Ameenudin, S., & Sangamanatha, A. V. (2012). Temporal and speech processing skills in

normal hearing individuals exposed to occupational noise. *Noise & health*, *14*(58), 100-105.

https://doi.org/10.4103/1463-1741.97252

Le Prell, C. G. (2019). Effects of noise exposure on auditory brainstem response and speech-in-noise

tasks: a review of the literature. *International journal of audiology*, *58*(sup1), S3-S32.

https://doi.org/10.1080/14992027.2018.1534010

Le Prell, C. G., Dell, S., Hensley, B., Hall, J. W., 3rd, Campbell, K. C. M., Antonelli, P. J., Green, G. E.,

Miller, J. M., & Guire, K. (2012). Digital music exposure reliably induces temporary threshold

shift   in   normal-hearing   human   subjects.   *Ear   and   hearing*,   *33*(6),   e44-e58.

https://doi.org/10.1097/AUD.0b013e31825f9d89

Liberman, M. C. (1982). Single-neuron labeling in the cat auditory nerve. *Science (New York, N.Y.)*,

216(4551), 1239-1241. https://doi.org/10.1126/science.7079757

Liberman, M. C., Dodds, L. W., & Pierce, S. (1990). Afferent and efferent innervation of the cat cochlea:

quantitative analysis with light and electron microscopy. *The Journal of comparative*

neurology, *301*(3), 443-460. https://doi.org/10.1002/cne.903010309

Liberman, M. C., Epstein, M. J., Cleveland, S. S., Wang, H., & Maison, S. F. (2016). Toward a differential

diagnosis   of   hidden   hearing   loss   in   humans   [Article].   *PLoS   ONE*,   *11*(9).

https://doi.org/10.1371/journal.pone.0162726

Liberman, M. C., & Kujawa, S. G. (2017). Cochlear synaptopathy in acquired sensorineural hearing loss:

Manifestations   and   mechanisms   [Review].   *Hearing   research*,   *349*,   138-147.

https://doi.org/10.1016/j.heares.2017.01.003

Lin, H. W., Furman, A. C., Kujawa, S. G., & Liberman, M. C. (2011). Primary neural degeneration in the

Guinea pig cochlea after reversible noise-induced threshold shift. *Journal of the Association*

for Research in Otolaryngology : JARO, *12*(5), 605-616. https://doi.org/10.1007/s10162-011-

0277-0

Luts, H., Jansen, S., Dreschler, W., & Wouters, J. (2014). Development and normative data for the

Flemish/Dutch Matrix test.

Makary, C. A., Shin, J., Kujawa, S. G., Liberman, M. C., & Merchant, S. N. (2011). Age-related primary

cochlear neuronal degeneration in human temporal bones. *Journal of the Association for*

Research in Otolaryngology : JARO, *12*(6), 711-717. https://doi.org/10.1007/s10162-011-

0283-2

Masterson, E. A., Themann, C. L., Luckhaupt, S. E., Li, J., & Calvert, G. M. (2016). Hearing difficulty and

tinnitus among U.S. workers and non-workers in 2007. *American journal of industrial medicine*,

59(4), 290-300. https://doi.org/10.1002/ajim.22565

Mercier, V., Luy, D., & Hohmann, B. W. (2003). The sound exposure of the audience at a music festival.

Noise & health, *5*(19), 51-58. http://www.noiseandhealth.org/article.asp?issn=1463-

1741;year=2003;volume=5;issue=19;spage=51;epage=58;aulast=Mercier

Mitchell, C., Phillips, D. S., & Trune, D. R. (1989). Variables affecting the auditory brainstem response:

Audiogram, age, gender and head size. *Hearing research*, *40*(1), 75-85.

https://doi.org/https://doi.org/10.1016/0378-5955(89)90101-9

Möhrle, D., Ni, K., Varakina, K., Bing, D., Lee, S. C., Zimmermann, U., Knipper, M., & Rüttiger, L. (2016).

Loss of auditory sensitivity from inner hair cell synaptopathy can be centrally compensated in

the young but not old brain. *Neurobiology of Aging*, *44*, 173-184.

https://doi.org/https://doi.org/10.1016/j.neurobiolaging.2016.05.001

Opperman, D. A., Reifman, W., Schlauch, R., & Levine, S. (2006). Incidence of spontaneous hearing

threshold shifts during modern concert performances. *Otolaryngology--head and neck surgery*

: official journal of American Academy of Otolaryngology-Head and Neck Surgery, *134*(4), 667-

673. https://doi.org/10.1016/j.otohns.2005.11.039

OSHA, U. S. D. o. L.-O. S. a. H. A. (1974). https://www.osha.gov/laws-

regs/regulations/standardnumber/1910/1910.95

Parthasarathy, A., Bartlett, E. L., & Kujawa, S. G. (2018). Age-related Changes in Neural Coding of
Envelope Cues: Peripheral Declines and Central Compensation. *Neuroscience*.
https://doi.org/10.1016/j.neuroscience.2018.12.007

Parthasarathy, A., & Kujawa, S. G. (2018). Synaptopathy in the aging cochlea: Characterizing early-
neural deficits in auditory temporal envelope processing [Article]. *Journal of Neuroscience*,
38(32), 7108-7119. https://doi.org/10.1523/JNEUROSCI.3240-17.2018

Paul, B. T., Waheed, S., Bruce, I. C., & Roberts, L. E. (2017). Subcortical amplitude modulation encoding
deficits suggest evidence of cochlear synaptopathy in normal-hearing 18-19 year olds with
higher lifetime noise exposure. *The Journal of the Acoustical Society of America*, *142*(5), El434.
https://doi.org/10.1121/1.5009603

Petrescu, N. (2008). Loud music listening. *McGill journal of medicine : MJM : an international forum for
the advancement of medical sciences by students*, *11*(2), 169-176.
https://www.ncbi.nlm.nih.gov/pubmed/19148318

Picton, T. W. (2011). *Human Auditory Evoked Potentials*. Plural Pub.
https://books.google.be/books?id=z05QAQAAIAAJ

Pouryaghoub, G., Mehrdad, R., & Mohammadi, S. (2007). Interaction of smoking and occupational
noise exposure on hearing loss: a cross-sectional study. *BMC public health*, *7*, 137.
https://doi.org/10.1186/1471-2458-7-137

Prendergast, G., Couth, S., Millman, R. E., Guest, H., Kluk, K., Munro, K. J., & Plack, C. J. (2019). Effects
of Age and Noise Exposure on Proxy Measures of Cochlear Synaptopathy. *Trends in hearing*,
23, 2331216519877301. https://doi.org/10.1177/2331216519877301

Prendergast, G., Guest, H., Munro, K. J., Kluk, K., Leger, A., Hall, D. A., Heinz, M. G., & Plack, C. J. (2017).
Effects of noise exposure on young adults with normal audiograms I: Electrophysiology.
Hearing research, *344*, 68-81. https://doi.org/10.1016/j.heares.2016.10.028

Prendergast, G., Millman, R. E., Guest, H., Munro, K. J., Kluk, K., Dewey, R. S., Hall, D. A., Heinz, M. G.,
& Plack, C. J. (2017). Effects of noise exposure on young adults with normal audiograms II:

Behavioral measures [Article]. *Hearing research*, *356*, 74-86. https://doi.org/10.1016/j.heares.2017.10.007

Prendergast, G., Tu, W., Guest, H., Millman, R. E., Kluk, K., Couth, S., Munro, K. J., & Plack, C. J. (2018). Supra-threshold auditory brainstem response amplitudes in humans: Test-retest reliability, electrode montage and noise exposure [Article]. *Hearing research*, *364*, 38-47. https://doi.org/10.1016/j.heares.2018.04.002

Puel, J. L., Ruel, J., Gervais d'Aldin, C., & Pujol, R. (1998). Excitotoxicity and repair of cochlear synapses after noise-trauma induced hearing loss. *Neuroreport*, *9*(9), 2109-2114. https://doi.org/10.1097/00001756-199806220-00037

Ramakers, G. G., Kraaijenga, V. J., Cattani, G., van Zanten, G. A., & Grolman, W. (2016). Effectiveness of Earplugs in Preventing Recreational Noise-Induced Hearing Loss: A Randomized Clinical Trial. *JAMA otolaryngology-- head & neck surgery*, *142*(6), 551-558. https://doi.org/10.1001/jamaoto.2016.0225

Ridley, C. L., Kopun, J. G., Neely, S. T., Gorga, M. P., & Rasetshwane, D. M. (2018). Using Thresholds in Noise to Identify Hidden Hearing Loss in Humans [Article]. *Ear and hearing*, *39*(5), 829-844. https://doi.org/10.1097/aud.0000000000000543

Ryan, A. F., Kujawa, S. G., Hammill, T., Le Prell, C., & Kil, J. (2016). Temporary and Permanent Noise-induced Threshold Shifts: A Review of Basic and Clinical Observations. *Otology & neurotology : official publication of the American Otological Society, American Neurotology Society [and] European Academy of Otology and Neurotology*, *37*(8), e271-275. https://doi.org/10.1097/mao.0000000000001071

Ryberg, J. B. (2009). A national project to evaluate and reduce high sound pressure levels from music. *Noise & health*, *11*(43), 124-128. https://doi.org/10.4103/1463-1741.50698

Safieddine, S., El-Amraoui, A., & Petit, C. (2012). The auditory hair cell ribbon synapse: from assembly to function. *Annual review of neuroscience*, *35*, 509-528. https://doi.org/10.1146/annurev-neuro-061010-113705

Schaette, R., & McAlpine, D. (2011). Tinnitus with a normal audiogram: physiological evidence for

hidden hearing loss and computational model. *The Journal of neuroscience : the official journal*

of the Society for Neuroscience, *31*(38), 13452-13457.

https://doi.org/10.1523/JNEUROSCI.2156-11.2011

Schlauch, R. S., & Carney, E. (2012). The challenge of detecting minimal hearing loss in audiometric

surveys. *American journal of audiology*, *21*(1), 106-119. https://doi.org/10.1044/1059-

0889(2012/11-0012)

Schmuzigert, N., Fostiropoulos, K., & Probst, R. (2006). Long-term assessment of auditory changes

resulting from a single noise exposure associated with non-occupational activities.

International journal of audiology, *45*(1), 46-54. https://doi.org/10.1080/14992020500377089

Sergeyenko, Y., Lall, K., Charles Liberman, M., & Kujawa, S. G. (2013). Age-related cochlear

synaptopathy: An early-onset contributor to auditory functional decline [Article]. *Journal of*

Neuroscience, *33*(34), 13686-13694. https://doi.org/10.1523/JNEUROSCI.1783-13.2013

Shaheen, L. A., Valero, M. D., & Liberman, M. C. (2015). Towards a Diagnosis of Cochlear Neuropathy

with Envelope Following Responses. *Journal of the Association for Research in Otolaryngology*

: JARO, *16*(6), 727-745. https://doi.org/10.1007/s10162-015-0539-3

Skoe, E., & Tufts, J. (2018). Evidence of noise-induced subclinical hearing loss using auditory brainstem

responses and objective measures of noise exposure in humans [Article]. *Hearing research*,

361, 80-91. https://doi.org/10.1016/j.heares.2018.01.005

Smith, P. A., Davis, A., Ferguson, M., & Lutman, M. E. (2000). The prevalence and type of social noise

exposure in young adults in England. *Noise & health*, *2*(6), 41-56.

Smith, S. B., Krizman, J., Liu, C., White-Schwoch, T., Nicol, T., & Kraus, N. (2019). Investigating peripheral

sources of speech-in-noise variability in listeners with normal audiograms. *Hearing research*,

371, 66-74. https://doi.org/10.1016/j.heares.2018.11.008

Spankovich, C., Bishop, C., Johnson, M. F., Elkins, A., Su, D., Lobarinas, E., & Le Prell, C. G. (2017).

Relationship between dietary quality, tinnitus and hearing level: data from the national health

and nutrition examination survey, 1999-2002. *International journal of audiology*, *56*(10), 716-

722. https://doi.org/10.1080/14992027.2017.1331049

Spankovich, C., Le Prell, C. G., Lobarinas, E., & Hood, L. J. (2017). Noise History and Auditory Function

in Young Adults With and Without Type 1 Diabetes Mellitus. *Ear and hearing*, *38*(6), 724-735.

https://doi.org/10.1097/AUD.0000000000000457

Stamper, G. C., & Johnson, T. A. (2015a). Auditory function in normal-hearing, noise-exposed human

ears. *Ear and hearing*, *36*(2), 172-184. https://doi.org/10.1097/aud.0000000000000107

Stamper, G. C., & Johnson, T. A. (2015b). Letter to the Editor: Examination of Potential Sex Influences

in . Auditory Function in Normal-Hearing, Noise-Exposed Human Ears, Ear Hear, 36, 172-184.

Ear and hearing, *36*(6), 738-740. https://doi.org/10.1097/aud.0000000000000228

Svensson, E. B., Morata, T. C., Nylen, P., Krieg, E. F., & Johnson, A. C. (2004). Beliefs and attitudes among

Swedish workers regarding the risk of hearing loss. *International journal of audiology*, *43*(10),

585-593. https://doi.org/Doi 10.1080/14992020400050075

Trzaskowski, B., Jędrzejczak, W. W., Piłka, E., Cieślicka, M., & Skarżyński, H. (2014). Otoacoustic

emissions before and after listening to music on a personal player. *Medical science monitor :*

international medical journal of experimental and clinical research, *20*, 1426-1431.

https://doi.org/10.12659/MSM.890747

Upile, T., Sipaul, F., Jerjes, W., Singh, S., Nouraei, R., Maaytah, M., Andrews, P., Graham, J., Hopper, C.,

& Wright, A. (2007). The acute effects of alcohol on auditory thresholds. *BMC ear, nose, and*

throat disorders, *7*, 4. https://doi.org/10.1186/1472-6815-7-4

Valderrama, J. T., Alvarez, I., de la Torre, A., Segura, J. C., Sainz, M., & Vargas, J. L. (2012). Recording of

auditory brainstem response at high stimulation rates using randomized stimulation and

averaging. *Journal of the Acoustical Society of America*, *132*(6), 3856-3865.

https://doi.org/10.1121/1.4764511

Valderrama, J. T., Beach, E. F., Yeend, I., Sharma, M., Van Dun, B., & Dillon, H. (2018). Effects of lifetime

noise exposure on the middle-age human auditory brainstem response, tinnitus and speech-

in-noise intelligibility. *Hearing research*, *365*, 36-48.

https://doi.org/10.1016/j.heares.2018.06.003

Vasilkov, V., Garrett, M., Mauermann, M., & Verhulst, S. (2021). Enhancing the sensitivity of the

envelope-following response for cochlear synaptopathy screening in humans: The role of

stimulus envelope. *Hear Res*, *400*, 108132. https://doi.org/10.1016/j.heares.2020.108132

Vasilkov, V., & Verhulst, S. (2019). Towards a differential diagnosis of cochlear synaptopathy and outer-

hair-cell deficits in mixed sensorineural hearing loss pathologies. *medRxiv*, 19008680.

https://doi.org/10.1101/19008680

Verhulst, S., Altoe, A., & Vasilkov, V. (2018). Computational modeling of the human auditory periphery:

Auditory-nerve responses, evoked potentials and hearing loss. *Hearing research*, *360*, 55-75.

https://doi.org/10.1016/j.heares.2017.12.018

Viana, L. M., O'Malley, J. T., Burgess, B. J., Jones, D. D., Oliveira, C. A. C. P., Santos, F., Merchant, S. N.,

Liberman, L. D., & Liberman, M. C. (2015). Cochlear neuropathy in human presbycusis:

Confocal analysis of hidden hearing loss in post-mortem tissue [Article]. *Hearing research*, *327*,

78-88. https://doi.org/10.1016/j.heares.2015.04.014

Wang, Y., Hirose, K., & Liberman, M. C. (2002). Dynamics of noise-induced cellular injury and repair in

the mouse cochlea. *Journal of the Association for Research in Otolaryngology : JARO*, *3*(3), 248-

268. https://doi.org/10.1007/s101620020028

Wei, W., Heinze, S., Gerstner, D. G., Walser, S. M., Twardella, D., Reiter, C., Weilnhammer, V., Perez-

Alvarez, C., Steffens, T., & Herr, C. E. W. (2017). Audiometric notch and extended high-

frequency hearing threshold shift in relation to total leisure noise exposure: An exploratory

analysis. *Noise & health*, *19*(91), 263-269. https://doi.org/10.4103/nah.NAH_28_17

Widen, S. E., Holmes, A. E., & Erlandsson, S. I. (2006). Reported hearing protection use in young adults

from Sweden and the USA: effects of attitude and gender. *International journal of audiology*,

45(5), 273-280. https://doi.org/10.1080/14992020500485676

Williams, W., Carter, L., & Seeto, M. (2015). Pure tone hearing thresholds and leisure noise: Is there a

relationship? *Noise & health*, *17*(78), 358-363. https://doi.org/10.4103/1463-1741.165066

Yamane, H., Nakai, Y., Takayama, M., Iguchi, H., Nakagawa, T., & Kojima, A. (1995). Appearance of free

radicals in the guinea pig inner ear after noise-induced acoustic trauma. *European archives of*

oto-rhino-laryngology : official journal of the European Federation of Oto-Rhino-Laryngological

Societies (EUFOS) : affiliated with the German Society for Oto-Rhino-Laryngology - Head and

Neck Surgery, *252*(8), 504-508. https://doi.org/10.1007/bf02114761

Yassi, A., Pollock, N., Tran, N., & Cheang, M. (1993). Risks to hearing from a rock concert. *Canadian*

family physician Medecin de famille canadien, *39*, 1045-1050.

https://www.ncbi.nlm.nih.gov/pubmed/8499785

https://www.ncbi.nlm.nih.gov/pmc/PMC2379666/

Yeend, I., Beach, E. F., Sharma, M., & Dillon, H. (2017). The effects of noise exposure and musical

training on suprathreshold auditory processing and speech perception in noise [Article].

Hearing research, *353*, 224-236. https://doi.org/10.1016/j.heares.2017.07.006

1000

1001

1002

1003

1004

1005

1006

1007

Figures

1009

1010 ***Figure 1***: *Exemplary ABR waveforms to 80 dB peSPL (dark grey) and 100 dB peSPL (blue), 11 Hz clicks. Respective ABR wave I,*
1011 *III and V peaks are specified by O and X for 80 and 100 dB peSPL click stimuli, respectively.*

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024

1025

Figure 2: *Exemplary EFR spectra. (A) SAM EFR spectrum (grey) and estimated noise-floor (pink). (B) Noise-floor corrected SAM EFR after bootstrapping. (C) RAM EFR spectrum (grey) and estimated noise-floor (pink). (D) Noise-floor corrected RAM EFR after bootstrapping. . Arrows in panels (B) and (D) show the magnitudes of EFR at modulation frequency (110 Hz) and harmonics.*

1026
1027
1028
1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041   *Figure 3*: *Frequency distribution of Laeq,life (A) and Laeq,recent (B).*

1042

1043

1044

1045

1046

1047

1048

1049

1050

Figure 4: Boxplots of baseline audiometric thresholds.

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066

1067

1068   ***Figure 5:*** *Boxplot of SPiQ (A) and SPiN (B) results at baseline session.*

1069

1070

1071

1072

1073

1074

1075

1076

1077

1078

1079

1080

Figure 6: *Scatterplot of EFR strength in relation to Laeq,life for SAM-stimuli (black triangles) and RAM-stimuli (grey triangles).*

1081

1082
1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100

1101

Figure 7: *Boxplots of speech audiometry in quiet (A) and in noise (B) for all conditions and all sessions. Dark grey boxplots represent SRTs for BB stimuli, light grey and white boxplots represent SRTs for HP and LP conditions, respectively. Intersession differences, compared to session 1, were significant if p<0.017 and are highlighted with one, two or three asterisks (resp. p<0.017, p<0.003 and p<0.0003). Both SPiN and SPIQ showed no significant differences between sessions 1 and 2.*

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116 **Tables**

Table 1: ABR data for ABR- and EFR responses of all stimulus types. Mean, SD, median and IQR are presented for ABR latencies and amplitudes of waves I, III and V and I/V amplitude ratio and EFR-strength.								
stimulus			11Hz 100dB$_{peSPL}$ click	11Hz 80dB$_{peSPL}$ click	120Hz 100dB$_{peSPL}$ click	120Hz 80dB$_{peSPL}$ click	TB$_{4kHz}$	TB$_{1 kHz}$
latencies	I	mean	2.67	3.32	3.14	3.70	3.96	4.36
		SD	0.19	0.27	0.39	0.27	0.44	0.53
		median	2.70	3.35	3.10	3.70	3.90	4.45
		IQR	0.20	0.27	0.40	0.20	0.75	0.67
	III	mean	4.94	5.46	5.47	6.01	6.15	6.48
		SD	0.23	0.31	0.33	0.33	0.38	0.70
		median	4.90	5.50	5.45	6.00	6.05	6.50
		IQR	0.20	0.35	0.40	0.27	0.65	0.75
	V	mean	6.73	7.18	7.32	7.90	7.94	5.57
		SD	0.38	0.43	0.38	0.43	0.35	0.43
		median	6.70	7.25	7.30	8.00	7.90	8.60
		IQR	0.47	0.35	0.47	0.37	0.40	0.45
amplitudes	I	mean	0.15	0.08	0.09	0.05	0.02	0.02
		SD	0.10	0.06	0.06	0.04	0.04	0.05
		median	0.17	0.07	0.08	0.04	0.03	0.02
		IQR	0.14	0.11	0.08	0.09	0.05	0.06
	III	mean	0.07	0.03	0.13	0.15	0.02	-0.01
		SD	0.07	0.07	0.10	0.09	0.06	0.07
		median	0.05	0.03	0.11	0.14	0.01	-0.01
		IQR	0.13	0.10	0.13	0.15	0.05	0.10
	V	mean	0.31	0.27	0.27	0.23	0.13	0.10
		SD	0.10	0.11	0.11	0.08	0.08	0.07
		median	0.30	0.27	0.26	0.22	0.13	0.09
		IQR	0.12	0.10	0.13	0.08	0.09	0.10
amplitude ratios	I/V	mean	0.47	0.34	0.39	0.22	-0.96	0.44
		SD	0.30	0.31	0.34	0.37	3.80	1.14
		median	0.44	0.30	0.31	0.21	0.15	0.17
		IQR	0.46	0.38	0.26	0.42	0.37	0.55
stimulus			SAM			RAM		
EFR-strength		mean	0.029			0.085		
		SD	0.010			0.029		
		median	0.028			0.088		
		IQR	0.011			0.037		

1117

1118

1119

1120

Table 2: χ2-values of related-samples Friedman's ANOVA for PTA. Significant results are highlighted with *(p<0.05), **(p<0.01) or ***(p<0.001). Median test values and IQRs are shown for each test session.

	Frequency	χ^2	Session 1 (pre)		Session 2 (1st post)		Session 3 (2nd post)		Session 4 (3d post)	
			med	IQR	med	IQR	med	IQR	med	IQR
Auditory thresholds (dB$_{HL}$)	125Hz	$\chi^2(3)$=5.56	5	5	10	10	5	10	5	10
	250Hz	$\chi^2(3)$=3.47	5	10	5	5	5	10	5	10
	500Hz	$\chi^2(3)$=8.73*	0	5	5	10	0	5	0	5
	1kHz	$\chi^2(3)$=4.73	5	5	0	5	0	5	0	10
	2kHz	$\chi^2(3)$=0.26	0	10	0	10	0	10	0	10
	3kHz	$\chi^2(3)$=0.22	0	5	0	5	0	5	0	10
	4kHz	$\chi^2(3)$=2.83	0	5	0	5	0	5	0	10
	6kHz	$\chi^2(3)$=2.03	10	5	10	15	10	15	10	10
	8kHz	$\chi^2(3)$=6.33	10	5	10	10	5	10	10	10
	10kHz	$\chi^2(3)$=12.69**	5	5	5	10	5	10	5	10
	12kHz	$\chi^2(3)$=4.88	5	15	10	20	5	15	5	15
	14kHz	$\chi^2(3)$=4.76	5	15	5	10	0	15	0	5
	16kHz	$\chi^2(3)$=7.39	10	30	10	20	10	25	5	25
	PTA$_{3-8kHz}$	$\chi^2(3)$= 3.84	5	5	5	8.75	3.75	5	5	8.75
	PTA$_{10-16kHz}$	$\chi^2(3)$=6.97	2.5	12.5	7.5	13.75	6.25	10	5	7.5

1121

1122

1123

1124

1125

1126

1127

1128

1129

1130

1131

1132

1133

1134

1135

1136

Table 3: χ^2-values of related-samples Friedman's ANOVA for AEP measurements. Significant results are highlighted with *(p<0.05), **(p<0.01) or ***(p<0.001). Median test values and IQRs are shown for each test session.

	Frequency	χ^2	Session 1 (pre)		Session 2 (1st post)		Session 3 (2nd post)		Session 4 (3d post)	
			med	IQR	med	IQR	med	IQR	med	IQR
ABR latency wave I (ms)	11Hz 100dB$_{peSPL}$	$\chi^2(3)=6.78$	2.70	0.20	2.70	0.10	2.70	0.10	2.70	0.20
	11Hz 80 dB$_{peSPL}$	$\chi^2(3)=0.88$	3.40	0.30	3.30	0.20	3.30	0.40	3.30	0.20
	120Hz 100 dB$_{peSPL}$	$\chi^2(3)=3.55$	3.10	0.40	3.20	0.40	3.00	0.80	3.20	0.50
	120Hz 80 dB$_{peSPL}$	$\chi^2(3)=0.88$	3.70	0.20	3.70	1.50	3.70	0.20	3.70	0.50
	TB$_{4kHz}$	$\chi^2(3)=2.27$	3.90	0.80	4.00	0.90	4.30	0.90	3.90	0.60
	TB$_{1kHz}$	$\chi^2(3)=0.62$	4.50	0.70	4.30	0.90	4.50	0.70	4.40	0.50
ABR latency wave III (ms)	11Hz 100dB$_{peSPL}$	$\chi^2(3)=3.16$	4.90	0.20	4.90	0.20	4.90	0.20	4.90	0.20
	11Hz 80 dB$_{peSPL}$	$\chi^2(3)=3.85$	5.50	0.30	5.70	0.30	5.50	0.30	5.50	0.30
	120Hz 100 dB$_{peSPL}$	$\chi^2(3)=1.72$	5.50	0.40	5.40	0.40	5.40	0.40	5.40	0.30
	120Hz 80 dB$_{peSPL}$	$\chi^2(3)=0.36$	6.00	0.30	6.10	0.30	6.00	0.30	6.10	0.50
	TB$_{4kHz}$	$\chi^2(3)=1.42$	6.00	0.55	6.20	0.65	6.20	0.45	6.20	0.70
	TB$_{1kHz}$	$\chi^2(3)=6.95$	6.50	0.70	6.80	0.40	6.60	0.50	6.50	0.40
ABR latency wave V (ms)	11Hz 100dB$_{peSPL}$	$\chi^2(3)=3.12$	6.70	0.50	6.70	0.30	6.70	0.30	6.70	0.30
	11Hz 80 dB$_{peSPL}$	$\chi^2(3)=1.76$	7.30	0.40	7.20	0.40	7.20	0.30	7.20	0.30
	120Hz 100 dB$_{peSPL}$	$\chi^2(3)=2.17$	7.30	0.50	7.40	0.20	7.30	0.30	7.40	0.40
	120Hz 80 dB$_{peSPL}$	$\chi^2(3)=4.11$	8.00	0.40	8.00	0.40	7.90	0.50	8.00	0.40
	TB$_{4kHz}$	$\chi^2(3)=0.14$	8.00	0.45	8.00	0.50	7.90	0.55	7.90	0.30
	TB$_{1kHz}$	$\chi^2(3)=1.06$	8.60	0.40	8.60	0.40	8.60	0.80	8.70	0.70
ABR amplitude wave I (µV)	11Hz 100dB$_{peSPL}$	$\chi^2(3)=1.61$	0.178	0.152	0.135	0.161	0.134	0.163	0.137	0.197
	11Hz 80 dB$_{peSPL}$	$\chi^2(3)=3.88$	0.068	0.11	0.061	0.12	0.087	0.15	0.091	0.14
	120Hz 100 dB$_{peSPL}$	$\chi^2(3)=1.11$	0.081	0.080	0.096	0.072	0.096	0.102	0.092	0.070
	120Hz 80 dB$_{peSPL}$	$\chi^2(3)=0.28$	0.042	0.096	0.075	0.096	0.046	0.088	0.070	0.125
	TB$_{4kHz}$	$\chi^2(3)=0.95$	0.022	0.060	0.027	0.036	0.023	0.068	0.008	0.044
	TB$_{1kHz}$	$\chi^2(3)=4.71$	0.013	0.065	0.006	0.077	0.029	0.062	0.015	0.061

ABR amplitude wave III (µV)	11Hz 100dB$_{peSPL}$	$\chi^2(3)=7.93^*$ p=0.48	0.044	0.137	0.057	0.130	0.073	0.100	0.103	0.112
	11Hz 80 dB$_{peSPL}$	$\chi^2(3)=2.43$	0.022	0.103	0.028	0.101	0.034	0.088	0.024	0.107
	120Hz 100 dB$_{peSPL}$	$\chi^2(3)=1.11$	0.114	0.122	0.096	0.079	0.133	0.072	0.112	0.081
	120Hz 80 dB$_{peSPL}$	$\chi^2(3)=3.44$	0.139	0.155	0.104	0.064	0.100	0.103	0.108	0.104
	TB$_{4kHz}$	$\chi^2(3)=7.52$	0.009	0.062	0.024	0.079	0.058	0.126	0.026	0.044
	TB$_{1kHz}$	$\chi^2(3)=3.82$	-0.020	0.101	-0.026	0.068	-0.001	0.076	-0.002	0.101
ABR amplitude wave V (µV)	11Hz 100dB$_{peSPL}$	$\chi^2(3)=6.35$	0.303	0.103	0.340	0.757	0.303	0.107	0.304	0.092
	11Hz 80 dB$_{peSPL}$	$\chi^2(3)=12.92^{**}$ p=0.005	0.269	0.108	0.287	0.113	0.254	0.122	0.279	0.092
	120Hz 100 dB$_{peSPL}$	$\chi^2(3)=3.82$	0.274	0.144	0.269	0.137	0.267	0.111	0.260	0.126
	120Hz 80 dB$_{peSPL}$	$\chi^2(3)=3.38$	0.222	0.085	0.248	0.152	0.242	0.136	0.231	0.079
	TB$_{4kHz}$	$\chi^2(3)=2.18$	0.111	0.113	0.129	0.070	0.130	0.078	0.134	0.091
	TB$_{1kHz}$	$\chi^2(3)=1.80$	0.091	0.103	0.100	0.066	0.062	0.083	0.095	0.059
ABR I/V amplitude ratio	11Hz 100dB$_{peSPL}$	$\chi^2(3)=1.67$	0.43	0.47	0.39	0.51	0.41	0.46	0.41	0.48
	11Hz 80 dB$_{peSPL}$	$\chi^2(3)=4.71$	0.29	0.39	0.23	0.37	0.33	0.62	0.27	0.59
	120Hz 100 dB$_{peSPL}$	$\chi^2(3)=1.04$	0.29	0.25	0.31	0.47	0.39	0.44	0.32	0.21
	120Hz 80 dB$_{peSPL}$	$\chi^2(3)=0.54$	0.21	0.43	0.24	0.47	0.23	0.37	0.29	0.45
	TB$_{4kHz}$	$\chi^2(3)=2.15$	0.05	0.38	0.22	0.47	0.16	0.59	0.05	0.29
	TB$_{1kHz}$	$\chi^2(3)=0.51$	0.15	0.49	0.05	0.85	0.24	0.86	0.02	0.73
EFR strength (µV)	SAM	$\chi^2(3)=1.13$	0.028	0.011	0.024	0.010	0.027	0.013	0.032	0.014
	RAM	$\chi^2(3)=2.87$	0.088	0.037	0.080	0.059	0.087	0.040	0.098	0.054

1137

1138

1139

1140

1141

1142

1143

1144

		no HRS (n=11)			HRS (n=9)			Man-Whitney U	
	Stimulus	Median	Minimum	Maximum	Median	Minimum	Maximum	U	p
latency wave I	11 Hz 80 dB	-0,1000	-0,20	0,10	0,0500	-0,10	0,40	78,00	0,03 *
	11 Hz 100 dB	-0,1000	-0,20	0,20	0,0000	0,00	0,60	79,50	0,02 *
	120 Hz 80 dB	0,0000	-0,30	0,20	0,1500	-0,70	1,20	66,00	0,23
	120 Hz 100 dB	0,1000	-0,30	1,20	-0,0500	-0,70	0,20	27,50	0,10
	TB 1 kHz	0,0000	-1,10	1,40	-0,1000	-0,60	1,00	49,00	1,00
	TB 4 kHz	-0,2000	-1,00	1,90	-0,1500	-1,00	0,80	39,50	0,66
Latency wave III	11 Hz 80 dB	0,1000	-0,10	0,80	0,1500	-0,10	1,00	50,50	0,94
	11 Hz 100 dB	0,0000	-0,80	0,10	0,0000	-0,10	0,10	57,50	0,55
	120 Hz 80 dB	0,0000	-0,30	1,20	-0,0500	-0,60	0,20	35,00	0,30
	120 Hz 100 dB	0,0000	-0,80	0,20	0,0500	-0,20	0,70	72,00	0,10
	TB 1 kHz	0,1000	-0,70	1,50	0,2000	-1,20	0,70	48,00	0,94
	TB 4 kHz	0,2000	-0,40	1,30	0,0000	-0,70	0,40	16,50	0,02 *
Latency wave V	11 Hz 80 dB	-0,1000	-0,20	0,10	0,0000	-0,10	1,40	73,00	0,08
	11 Hz 100 dB	0,0000	-0,10	0,40	0,0500	-1,30	0,20	53,00	0,82
	120 Hz 80 dB	0,0000	-0,10	0,40	-0,0500	-0,50	1,10	46,50	0,82
	120 Hz 100 dB	-0,1000	-0,20	0,70	0,1000	-0,10	1,10	73,00	0,08
	TB 1 kHz	0,1000	-0,40	1,40	-0,0500	-0,60	0,10	29,00	0,13
	TB 4 kHz	0,0500	-0,50	2,00	-0,1000	-0,20	0,40	32,00	0,32
Amplitude wave I	11 Hz 80 dB	-0,0076	-0,12	0,07	0,0096	-0,05	0,11	52,00	0,88
	11 Hz 100 dB	0,0076	-0,15	0,13	0,0338	-0,20	0,20	56,00	0,66
	120 Hz 80 dB	0,0189	-0,08	0,17	-0,0042	-0,18	0,17	45,00	0,77
	120 Hz 100 dB	0,0184	-0,08	0,10	-0,0214	-0,13	0,08	31,00	0,18
	TB 1 kHz	-0,0115	-0,08	0,24	-0,0592	-0,12	0,23	35,00	0,30
	TB 4 kHz	-0,0154	-0,07	0,04	0,0580	-0,05	0,16	71,00	0,04 *
Amplitude wave III	11 Hz 80 dB	0,0210	-0,04	0,13	0,0103	-0,10	0,05	32,00	0,20
	11 Hz 100 dB	-0,0424	-0,13	0,14	-0,0266	-0,13	0,16	55,00	0,71

Table 4: *U-values and significance levels of Man Whitney U test for AEP measurements. Median test shifts from session 1 to session 2, minima and maxima are shown for subjects that have experienced Hearing Related Symptoms (HRS) and those who did not. Significant between-group differences are highlighted with *(p<0.05), **(p<0.01) or ***(p<0.001).*

	120 Hz 80 dB	-0,0125	-0,46	0,13	-0,0298	-0,28	0,14	55,00	1,00
	120 Hz 100 dB	-0,0029	-0,29	0,10	-0,0330	-0,24	0,10	47,00	0,88
	TB 1 kHz	-0,0264	-0,17	0,08	0,0318	-0,04	0,24	71,00	0,11
	TB 4 kHz	-0,0186	-0,15	0,10	0,0316	-0,08	0,12	53,00	0,55
Amplitude wave V	11 Hz 80 dB	0,0108	-0,09	0,27	0,0202	-0,15	0,30	50,00	1,00
	11 Hz 100 dB	0,0590	-0,15	0,15	0,0108	-0,04	0,23	55,00	0,71
	120 Hz 80 dB	0,0071	-0,15	0,19	0,1152	-0,15	0,22	61,00	0,41
	120 Hz 100 dB	0,0085	-0,30	0,15	0,0240	-0,09	0,27	65,00	0,26
	TB 1 kHz	0,0114	-0,08	0,06	-0,0039	-0,08	0,03	45,00	0,77
	TB 4 kHz	0,0426	-0,10	0,15	-0,0429	-0,16	0,04	17,00	0,02 *
I/V amplitude ratio	11 Hz 80 dB	-0,0364	-0,67	0,40	-0,0373	-1,01	0,42	48,00	0,78
	11 Hz 100 dB	0,0817	-0,54	0,45	0,1142	-0,95	0,39	50,00	0,66
	120 Hz 80 dB	0,0847	-0,55	1,00	-0,0123	-0,95	0,25	31,00	0,31
	120 Hz 100 dB	0,0926	-0,36	8,57	-0,0568	-0,77	0,05	17,00	0,03 *
	TB 1 kHz	-0,1842	-4,04	63,34	-0,3445	-1,16	3,85	35,00	0,49
	TB 4 kHz	0,0700	-0,45	16,34	0,5031	-0,21	13,93	50,00	0,41

1146