Canopy spectral reflectance detects oak wilt at the landscape scale using phylogenetic discrimination

Authors

Gerard Sapes[1*], Cathleen Lapadat[1], Anna K. Schweiger[1,2,3], Jennifer Juzwik[4], Rebecca Montgomery[5], Hamed Gholizadeh[6], Philip A. Townsend[7], John A. Gamon[8,9], and Jeannine Cavender-Bares[1*].

Affiliations

[1] Department of Ecology, Evolution, and Behavior, University of Minnesota, Saint Paul, Minnesota 55108, USA

[2] Institut de recherche en biologie végétale and département de sciences biologiques, Université de Montréal, Montréal, Québec, Canada

[3] Department of Geography, University of Zurich, Zurich, Switzerland

[4] Northern Research Station, USDA Forest Service, St. Paul, MN

[5] Department of Forest Resources, University of Minnesota, St. Paul, MN, 55108, USA

[6] Center for Applications of Remote Sensing, Department of Geography, Oklahoma State University, Stillwater, Oklahoma, 74078, USA

[7] Department of Forest and Wildlife Ecology, University of Wisconsin-Madison, 1630 Linden Drive, Madison, WI 53706, USA

[8] School of Natural Resources, University of Nebraska – Lincoln, Lincoln, NE, USA 68583-0961

20   [9] Departments of Earth & Atmospheric Sciences and Biological Sciences, University of Alberta,

21   Edmonton, Alberta, Canada T6G 2E3

ORCID:

Gerard Sapes: https://orcid.org/0000-0002-6017-2053

Anna K. Schweiger: https://orcid.org/0000-0002-5567-4200

Hamed Gholizadeh: https://orcid.org/0000-0002-4770-7893

Philip A. Townsend: http://orcid.org/0000-0001-7003-8774

John A. Gamon: https://orcid.org/0000-0002-8269-7723

Jeannine Cavender-Bares: https://orcid.org/0000-0003-3375-9630

Author contribution statement

J.C.B. Conceived of and managed the project. G.S., C.L., A.S., J.J., R.M. and J.C.B conceived the experimental design and data analysis. C.L., A.S., and J.J. collected ground data. H.G., P.T., and J.G. collected and processed airborne data. G.S. analyzed the data. G.S., J.C.B., and J.J., wrote the manuscript with contributions from C.L., A.S., R.M., H.G., P.T., and J.G. J.C.B, J.J., and R.M. obtained funding for this project.

Corresponding authors:

Gerard Sapes (gsapes@umn.edu) and Jeannine Cavender-Bares (cavender@umn.edu)

Phone: (406) 207-7955

Department of Ecology, Evolution, and Behavior

University of Minnesota

1987 Upper Buford Circle 55108, Saint Paul, Minnesota, USA

Abstract

The oak wilt disease caused by the invasive fungal pathogen *Bretziella fagacearum* is one of the greatest

threats to oak-dominated forests across the Eastern United States. Accurate detection and monitoring

over large areas are necessary for management activities to effectively mitigate and prevent the spread

of oak wilt. Canopy spectral reflectance contains both phylogenetic and physiological information across

the visible near-infrared (VNIR) and short-wave infrared (SWIR) ranges that can be used to identify

diseased red oaks. We develop partial least square discriminant analysis (PLS-DA) models using airborne

hyperspectral reflectance to detect diseased canopies and assess the importance of VNIR, SWIR,

phylogeny, and physiology for oak wilt detection. We achieve high accuracy through a three-step

phylogenetic process in which we first distinguish oaks from other species (90% accuracy), then red oaks

from white oaks (*Quercus macrocarpa*) (93% accuracy), and, lastly, infected from non-infected trees

(80% accuracy). Including SWIR wavelengths increased model accuracy by ca. 20% relative to models

based on VIS-NIR wavelengths alone; using a phylogenetic approach also increased model accuracy by

ca. 20% over a single-step classification. SWIR wavelengths include spectral information important in

differentiating red oaks from other species and in distinguishing diseased red oaks from healthy red

oaks. We determined the most important wavelengths to identify oak species, red oaks, and diseased

red oaks. We also demonstrated that several multispectral indices associated with physiological decline

can detect differences between healthy and diseased trees. The wavelengths in these indices also

tended to be among the most important wavelengths for disease detection within PLS-DA models,

indicating a convergence of the methods. Indices were most significant for detecting oak wilt during late

August, especially those associated with canopy photosynthetic activity and water status. Our study

suggests that coupling phylogenetics, physiology, and canopy spectral reflectance provides an

interdisciplinary and comprehensive approach that enables detection of forest diseases at large scales.

These results have potential for direct application by forest managers for detection to initiate actions to

mitigate the disease and prevent pathogen spread.

Keywords: Oak wilt, photosynthetic decline, spectral reflectance, disease response, water content,

remote sensing, physiology, phylogenetic classification.

1. Introduction

Invasive tree pathogens are a major threat to forest diversity and function (Evans et al., 2010; Hulcr

and Dunn, 2011). The damage caused by invasive species can have negative consequences for

ecosystem processes and services, including air and water quality maintenance, nutrient and carbon

cycling, wood and food provision, and climate regulation (Cavender-Bares et al., 2019; Díaz et al., 2019;

Waller et al., 2020). In North American forests, invasive pathogens and pests that infect trees have had

devastating impacts over the last century due to multiple factors, including global trade and climate

change (Bergot et al., 2004; Liebhold et al., 1995; Sturrock et al., 2011), leading to the loss or potential

loss of multiple foundational canopy species such as American chestnut (*Castanea dentata*), elm and ash

species (*Ulmus* and *Fraxinus* spp.), and eastern hemlock (*Tsuga canadiensis*). Protecting ecosystems

from the threats of invasive species resulting from globalization and a changing climate is one of the

most pressing challenges of our times (Díaz et al., 2019; Liebhold et al., 1995; Waller et al., 2020).

The oak genus (*Quercus*) is under threat from multiple pathogens and is of critical management

interest due to its dominance in temperate forests of the Eastern US (Johnson et al., 2019). Oaks rank

among the most diverse and important tree lineages in the United States, with 91 oak species

comprising nearly 30% of biomass in US temperate forests (Cavender-Bares, 2019). Among the

pathogens affecting oaks, oak wilt caused by the fungus *Bretziella fagacearum* (de Beer et al., 2017) is

considered one of the most destructive threat to oaks (Appel, 1995; Haight et al., 2011; Wilson and

Lindsey, 2005). The oak wilt fungus spreads below-ground from diseased trees to neighboring oaks

through networks of grafted roots, thus forming centers (i.e., pockets or foci) of diseased oaks. The

pathogen is also transmitted above-ground by nitidulid beetles (family *Nitidulidae*) and oak bark beetles

(*Pityophthorus spp*) (Gibbs and French 1980).   Multiple species of nitidulid beetles are attracted to

spore-producing fungal mats that form on branches and main stems of recently wilted red oaks (Gibbs

and French, 1980; Juzwik et al., 2011; Juzwik and French, 1983). On a land parcel to larger scale, oak wilt

can be most effectively controlled when oak wilt centers are detected and appropriately treated (Juzwik

et al., 2011; Koch et al., 2010). This prevents spread or minimizes disease intensification within a stand

or the surrounding landscape. Surveys of large, forested areas to identify suspect diseased trees are

time-intensive and require expert training. Such surveys are needed for landscape level oak wilt

management efforts. Current operational surveillance of forest landscapes in the Upper Midwest USA

utilize aerial surveys conducted with fixed wing aircraft, helicopters, and UAVs (Juzwik, 2009). Other

airborne imaging spectrometry offers potential for accurate detection of oak wilt at landscape scales.

Canopy spectral reflectance can potentially be used to detect the physiological decline resulting

from oak wilt fungus infection, and thus provide forest managers with a powerful tool. Airborne spectral

reflectance and indices derived from reflectance spectra have successfully been used to detect other

diseases and insect damage, such as Rapid Ohia Death, Emerald Ash Borer, bark beetles, olive decline

due to *Xylella fastidiosa*, and both drought- and *Phytophtora*-induced holm oak decline (Asner et al.,

2018; Hornero et al., 2021; Lausch et al., 2013; Ogaya et al., 2015; Pontius et al., 2008, 2005; Zarco-

Tejada et al., 2018). To date, spectral indices for oak wilt detection have only been developed for oak

seedlings (Fallon et al., 2020). Oaks respond to oak wilt infection by forming balloon-like structures

called tyloses that occlude vessels within the xylem (Juzwik and Appel, 2016; Yadeta and Thomma, Bart,

2013). Vessel occlusion potentially blocks or slows the spread of the pathogen but also reduces water

transport and limits canopy physiological performance by reducing transpiration and photosynthesis and

potentially causing photoinhibition (Fallon et al., 2020; Juzwik and Appel, 2016; Struckmeyer et al.,

1954). In red oak species, the fungus is rapidly spread internally in the transpiration stream through

large diameter springwood vessels before tylose formation limits the pathogen's spread. However, the

tyloses formed contribute to the development of wilt symptoms. Blockage of vascular conduits by

tyloses and metabolites produced by the fungus can lead to declines in transpiration and canopy water

content as water supply to the canopy is significantly impaired. Changes in photosynthetic activity, foliar

pigment pool sizes, and water status can be detected from canopy spectra (Hanavan et al., 2015;

Pontius et al., 2005; Serbin et al., 2015). Changes in foliar pigments, particularly carotenoids, chlorophyll,

and those involved in the xanthophyll cycle, have recently been shown to be early markers of holm oak

decline (Encinas-Valero et al., 2021). Fallon et al. (2020) identified several spectral wavelengths

predictive of oak wilt in greenhouse seedlings that were related to leaf photosynthesis and water status.

However, spectral properties of seedlings grown and measured under greenhouse conditions may differ

significantly from adult trees grown under natural conditions due to growing conditions (e.g., sun,

shade, humidity, and selective filtering of solar radiation by glasshouse materials), ontogeny, canopy

position, degree of canopy emergence and other factors (Cavender-Bares et al., 2020; Fernandes et al.,

2020; Ollinger, 2011). Hence, it is important to explicitly test the extent to which we can detect oak wilt

in natural populations of adult trees using spectral reflectance. Identification of wavelengths associated

with physiological function may enable detection of isolated trees with oak wilt that would otherwise

remain undetected until oak wilt damage is more extensive.

Oak lineages vary in susceptibility to oak wilt. Consequently, identifying vulnerable oak subgenera is

crucial to detect the disease and prevent its spread. White oaks (*Quercus* section *Quercus*), such as *Q.*

alba and *Q. macrocarpa*, have narrower vessels (Cavender-Bares and Holbrook, 2001) and may produce

tyloses efficiently (Cochard and Tyree, 1990) and in a more targeted manner in response to fungal

infection (cf. Yadeta and Thomma, Bart, 2013). This slows the spread or compartmentalizes (cf. Shigo,

1984) the pathogen in infected white oak species (Jacobi and MacDonald, 1980; Koch et al., 2010;

Schoenweiss, 1959). Thus, symptoms of oak wilt in white oaks appear as scattered wilt or as dieback in

the crown that develops over several to many years. In contrast, red oaks (*Quercus* section *Lobatae*),

such as *Q. ellipsoidalis* and *Q. rubra*, have larger diameter springwood vessels and tend to delay tylose

formation in response to fungal infection, limiting their effectiveness in halting the spread of the fungus

through the vascular system (Juzwik and Appel, 2016; Struckmeyer et al., 1954; Yadeta and Thomma,

Bart, 2013). Thus, crown wilt symptoms in red oaks progress rapidly and lead to tree death within the

same season or early in the subsequent growing season. The comparatively rapid mortality of red oaks,

the common occurrence of intra-specific root grafts, and their common production of spore mats on

recently wilted trees contribute to the importance of the red oak lineage in driving disease epidemics in

the landscape (Menges and Loucks, 1984). Distinguishing red oaks from white oaks and other species

across the landscape is therefore a critical step towards large-scale management of oak wilt. Leaf level

and canopy-level modeling approaches using spectroscopic data have previously been successful in

distinguishing these lineages in experimental systems and manipulated forest communities (Cavender-

Bares et al., 2016; Fallon et al., 2020; Williams et al., 2020). We thus anticipate that it is possible to

detect red oaks across the landscape remotely by mapping lineage identities from classification

algorithms using airborne spectroscopic imagery. Here, we outline a stepwise phylogenetic approach to

remote sensing of oak wilt that entails: 1) identifying trees belonging to the oak genus, 2) identifying

oaks belonging to the red oak section, and 3) identifying red oaks infected with oak wilt. The

phylogenetic approach consists of three sequential steps: first, pixels are classified as either "oak" or

"other species" using an algorithm specifically trained to discriminate oaks and filter out pixels classified

as other species. Second, pixels classified as "oak" are classified as either "red oak" or "white oak" using

an algorithm specifically trained to discriminate between red oaks and white oaks. After filtering out

pixels classified as white oaks, pixels classified as "red oak" are classified as either "diseased" or "healthy

red oak" using an algorithm specifically trained to distinguish diseased red oaks from healthy read oaks.

The goal of this study is to identify the optimal spectral range for detection of oak wilt in red oak

species (*Q. ellipsoidalis* and *Q. rubra*) across landscapes, find key wavelengths indicative of oak wilt and

its host species, and identify common spectral indices that distinguish diseased from healthy read oaks

based on physiological processes. Specifically, we compare both full-range (visible, near-infrared,

shortwave infrared, VSWIR, 400-2500 nm) and VNIR (visible, near-infrared, 400-1000 nm) imaging

spectroscopy and assess their accuracy of oak wilt detection. While the VNIR is sensitive to

photosynthetic activity and pigments (Curran et al., 1995; Gamon and Surfus, 1999; Ustin et al., 2009),

use of the SWIR provides structural and phenotypic information (Townsend et al., 2013) that is strongly

coupled with phylogenetic information (Meireles et al., 2020a) including mesophyll integrity, chemical

composition, and canopy water content (Jacquemoud and Ustin, 2001; Ramirez et al., 2015; Romero et

al., 2012; Sims and Gamon, 2003). Then, we test the efficacy of spectral vegetation indices known to be

sensitive to physiological decline and disease response for their ability to differentiate healthy and

diseased trees (Pontius, 2014; Pontius 2020) (Table S1). Spectral indices can increase flexibility in the

detection approach because they use only a handful of wavelengths and can be easily calculated across

platforms as long as the same wavelengths are present (Pontius, 2014). Spectroscopic models that

require hundreds of wavelengths can have limited applicability across platforms when sensor

measurement characteristics vary (Castaldi et al., 2018; Crucil et al., 2019; Nouri et al., 2017).

Here, we develop statistical models for oak wilt detection at the landscape scale using airborne

spectroscopic imagery collected by two airborne systems (AISA Eagle and AVIRIS-NG) (Gholizadeh et al.,

2019; Hamlin et al., 2010) covering different ranges of wavelengths (VNIR and VSWIR, respectively). We

coupled on-ground tree identification and status surveys with airborne imaging spectroscopy data to

assess the capacity of airborne spectroscopy to detect oak wilt in a temperate, mixed hardwood forest

that included adult red oak populations. In doing so, we tested the following hypotheses:

i)   Canopy reflectance from airborne spectroscopic imagery can accurately detect oak wilt

infected trees in a natural forest landscape;

ii)   Detecting red oaks susceptible to oak wilt prior to diseased trees based on spectral features

specific to their phylogenetic lineage increases oak wilt detection by removing species

outside the oak genus and red oak lineage;

iii)   A broad spectral range (VNIR+SWIR) exhibits greater detection accuracy than a narrower

spectral range (VNIR only) due to additional spectral information related to phylogenetic

identity and plant structure; and

iv)   Spectral indices including wavelengths associated with photosynthetic activity, carotenoid,

chlorophyll, and xanthophyll pigment content, and canopy water status differentiate

diseased red oaks from healthy red oaks.

2. Methods

<u>2.1 Study area</u>

The study area was the University of Minnesota Cedar Creek Ecosystem Science Reserve (CCESR)

(N 45º40'21", W 93º19'94"). Located in central Minnesota at approx. 280 m above sea level, CCESR has

a continental climate with cold winters (January mean -10 °C), hot summers (July mean 22.2 °C), and a

mean annual precipitation of 660 mm, spread fairly evenly throughout the year. The vegetation is

comprised of a mosaic of uplands dominated by oak savanna, prairie, mixed hardwood forest, and

abandoned agricultural fields, with lowlands comprised of ash and cedar swamps, acid bogs, marshes,

and sedge meadows. Oak savanna used to be one of the dominant vegetation types in the Cedar Creek

area before the European settlement (Grigal et al., 1974). The savannas mainly include two species

within the red oak group (*Quercus* section Lobatae, (Denk et al., 2017)), *Quercus ellipsoidalis* and *Q.*

rubra (northern pin oak and northern red oak, respectively) and a bur oak, *Q. macrocarpa* (within the

white oak group, *Quercus* section Quercus). These two oak groups have different sensitivities to oak

wilt, with red oaks being more susceptible, dying within one or two years of infection. The presence of

oak wilt fungus has been documented in central Minnesota since the 1940's where it has led to

widespread mortality in forests not treated for the disease. At CCESR, the fungal pathogen oak wilt has

spread rapidly in the last decade, leading to exponential increases in the number of standing dead trees

as a result of recent mortality (Pellegrini et al., 2021). The diversity of tree species and the widespread

presence of active oak wilt make CCESR well suited to assess the capacity of airborne spectroscopy to

detect oak wilt in red oaks.

<u>2.2 Airborne data collection and tree survey</u>

We collected two airborne imaging spectroscopy datasets across the whole study area on two

dates in 2016. The first dataset was collected on 07/22/2016 between 9:08 am and 10:24 am local time

using "CHAMP" (the CALMIT Hyperspectral Airborne Monitoring Platform), the University of Nebraska –

Lincoln's (UNL) aircraft operated by UNL's Center for Advanced Land Management Information

Technologies (CALMIT) and equipped with a pushbroom imaging spectrometer (AISA Eagle, Specim,

Oulu, Finland). Data were collected at an average flight altitude of 1150 m above ground level in the

northwest-southeast direction, yielding a spatial resolution of 0.75 m. The AISA Eagle comprises 488

spectral channels covering 400-982 nm with a spectral resolution of 1.25 nm and a field of view of 37.7°

under nadir viewing conditions. To increase the signal-to-noise-ratio of the data, spectral on-chip

binning was applied. The final product had 63 bands at ca. 9 nm intervals. The AISA Eagle images were

geometrically corrected using aircraft GPS and IMU data in Specim's CaliGeoPRO software. Radiance

data were converted to reflectance using the empirical line correction (Conel et al., 1987) on reflectance

measurements collected from three calibration tarps (white, grey and black, with approx. 5%, 10%, and

40% reflectance, respectively; Odyssey, Ennis Fabrics, Edmonton, Alberta, Canada) with a portable

spectroradiometer (SVC HR-1024i, Spectra Vista Corporation, Poughkeepsie, NY, USA; 350 – 2500 nm)

simultaneous to the overflights. SVC reflectance data were resampled to match the wavelength of

airborne data and then used in the empirical line correction approach. The second dataset was collected

using the Airborne Visible/Infrared Imaging Spectrometer - Next Generation (AVIRIS-NG) by the National

Aeronautics and Space Administration (NASA) on 08/22/2016 starting at 03:43 PM local time at an

average flight altitude of 1210 m above ground level in the near West-East direction, yielding a spatial

resolution of 0.9 m. AVIRIS-NG comprises 432 spectral channels covering 380-2510 nm with a spectral

resolution of 5 nm and a field of view of 36° under nadir viewing conditions. We measured the three

calibration tarps with our portable spectroradiometer (SVC HR-1024i, Spectra Vista Corporation,

Poughkeepsie, NY, USA; 350 – 2500 nm) during the overflights for empirical line correction. AVIRIS-NG

images that were delivered by the NASA Jet Propulsion Laboratory (JPL) were orthorectified and

atmospherically corrected to obtain apparent surface reflectance using a radiative transfer approach

following Thompson et al. (2015), while the AISA Eagle images were corrected using empirical line

correction. Because the aircraft images were acquired from different platforms on different dates, with

different instruments and atmospheric correction approaches, our objective was to test the relative

capacities of each system rather than to integrate the results from each. Specifically, absolute values

from the two sensors are not directly comparable. However, if images from each sensor are processed

consistently (see Wang et al., 2021) the results from each set of analyses to the different datasets can be

compared. While using two different sensors simultaneously is not necessary to detect oak wilt, not all

forest managers have access to all sensor types. Testing two different sensors allows us to find

wavelengths and indices predictive of oak wilt for multiple sensors and provides alternative tools for oak

wilt management.

About one year after collecting airborne data, between June-August of 2017, we tagged 423

mature trees of 12 species with no visual symptoms of oak wilt in woodland and savanna areas (see

Table S2 for a description of the range of heights and diameters at breast height, and the number of

trees included for each species) including 47 *Quercus ellipsoidalis* E.J. Hill (red oak section, particularly

vulnerable to oak wilt). In addition to the 423 healthy trees, we tagged 41 adult *Q. ellipsoidalis* trees

(total of 464 trees) with foliar symptoms characteristic of oak wilt (e.g., leaf epinasty, leaf bronzing

discoloration starting from apex and lateral margins and progressing to mid-rib and base of the leaf) (Fig.

S1). If any of such symptoms were observed—whether a few branches or most of the crown—the tree

was considered positive for oak wilt. Note that the number of trees covered by each flight slightly varied

due to different flight paths (Table S2). Current season crown wilt in 2017 suggested that crown wilt was

present during mid to late August 2016 when airborne spectral data were collected. We georeferenced

the canopy center of each tagged tree using a high-precision Trimble Pro6H GPS (Trimble, Sunnyvale,

CA, USA) during the leaf-off stage the following winter 2017-2018. Finally, we georeferenced an

additional 83 oaks (48 *Q. ellipsoidalis* and 35 *Q. macrocarpa*) that were not included in the training or

testing steps of our models (see below). Instead, we used these oaks to further validate our models.

<u>2.3 Canopy spectra extraction</u>

We built a 1m-radius circular buffer around each canopy center using ArcGIS (version 10.6.1,

ESRI, 2011) to sample fully sunlit canopy pixels per individual tree (Table S2), which were then linked to

the respective species and oak wilt status (i.e., healthy, diseased). The number of pixels per tree ranged

from four to nine—depending on the position of the buffer center in respect to that of the pixels within

the buffer—for both AISA Eagle and AVIRIS-NG datasets with the AISA Eagle dataset yielding, on

average, one to two more pixels per buffer due to its smaller pixel size. We applied the segmentation

method designed to discard shaded canopy pixels and keep sunlit pixels only. We visually assessed

reflectance in canopy pixels of a subset of trees for every species within our study and determined

reflectance values near 552 nm, 671 nm, and 800 nm representative of sunlit canopy pixels (Malenovský

et al., 2006). Then, we excluded pixels with values below such thresholds (0.01, 0.01, and 0.15 for 552

nm, 671 nm, and 800 nm, respectively) from our datasets. Spectral data processing employed the

package *spectrolab* (Meireles et al. 2018) in R (version 3.6.0, R Development Core Team, 2020). First, we

resampled the extracted spectral data to 410-980 nm for AISA Eagle and 410-2400 nm for AVIRIS-NG

(both at 5 nm resolution to match wavelengths across sensors within the VNIR range) to remove noisy

wavelengths at the range ends of the sensors and reduce the number of bands in the analyses. For

AVIRIS-NG data only, we removed atmospheric water absorption bands between 1335-1430 nm and

1770-1965 nm and corrected artifacts at the sensor overlap region around 950 nm. Finally, for both

datasets we unit vector-normalized reflectance values to reduce illumination differences among spectra

(i.e., standardize differences in amplitude) (Feilhauer et al., 2010) while preserving differences in the

shape of spectra that are important for species classification (Meireles et al., 2020b). After processing

spectra, we calculated 21 spectral indices commonly used in the literature related to plant

photosynthetic activity (e.g., RDVI, SIPI, SIF), water status (e.g., WBI, NDWI), and photoprotective stress

(e.g., PRI, CRI700, NPQI) (see Table S1 for full index list). In cases where an index required a wavelength

that was not a multiple of 5 and therefore missing in our spectra, we approximated the reflectance

value of that wavelength based on the reflectance of the neighboring wavelengths either by using the

nearest wavelength if the difference was ≤ 1 nm or otherwise by interpolation between the two nearest

wavelengths. We assessed whether vector normalization affected the capacity of spectral indexes to

detect oak wilt infected trees and found no major differences in spectral index performance (Appendix

S1).

2.4 Statistical analyses

All statistical analyses were performed in R (version 3.6, R Development Core Team, 2020). To

assess the capacity of canopy spectral reflectance to distinguish healthy trees from those infected with

oak wilt, we performed partial least square discriminant analyses (PLS-DAs) (Barker and Rayens, 2003)

using AISA Eagle (410-980 nm), AVIRIS-NG VNIR (410-980 nm), AVIRIS-NG SWIR (985-2400 nm) and

AVIRIS-NG VSWIR (410-2400 nm). Performing PLS-DAs for each spectral range allowed us to assess the

importance of each range of wavelengths for accurate detection. We treated each pixel as an

observation because oak wilt disease does not manifest uniformly across the canopy of a tree, especially

during early stages of infection. At early stages, the fungus may have infected only a fraction of the

vessels within the tree trunk. Thus, curtailing the water supply to a few branches that become

symptomatic while others remain asymptomatic. Treating pixels -rather than the whole tree- as

observations is critical to prevent false negatives that result from early infected trees displaying a small

number of symptomatic pixels. Thus, averaging pixels across a canopy composed of mostly healthy

pixels may hide the signal from the infected pixels and lead to lower true positive classification rates as

evidenced by the reduced performance of whole-tree level PLS-DAs aimed at distinguishing diseased

from healthy trees (Fig. S2). In all PLS-DAs, we used ANOVA to compare models with different numbers

of components and to identify the minimal number of components that maximized Kappa, a model

performance statistic that quantifies model performance compared to random classification (Cohen,

1960). PLS-DAs were then run with the optimal number of components and the "Bayes" option to

account for differences in prior probability distributions among classes (Brereton and Lloyd, 2014). The

optimal number of components varied by model and are reported in the results section.

We tested the extent to which distinguishing red oaks from other species before oak wilt status

classification improved the predictive performance of our models by evaluating two approaches for oak

wilt detection: a modelling pipeline that did not consider species identities ("direct" approach) and one

that differentiated red oaks from other species first ("phylogenetic" approach) (Fig. 1). Both approaches

were applied to each sensor type and spectral range. In the direct approach, we ignored species

identities and split the data within each class ("diseased" and "other") into 75:25 randomly sampled

subsets for model training and testing, respectively (Brereton and Lloyd, 2014; Fallon et al., 2020). We

used the *caret* and *pls* packages in R (Kuhn 2008, Mevik et al. 2018) to assess model performance

(accuracy, sensitivity, specificity, kappa) and obtain model-predicted values for each class (Congalton,

2001; Fassnacht et al., 2006). The random sampling, model training, model testing, performance

assessment loop was iterated 10,000 times to generate 10,000 different training and test subsets,

classification models, and corresponding performance estimates. We assessed overall performance of

the direct approach by calculating the average and standard deviation of the performance outputs

across all iterations.

Direct Approach

Diseased?

All species

10,000 x

75% Train

25% Test

Phylogenetic Approach

Figure 1: Workflow of the direct and phylogenetic modeling approaches used to classify diseased red

oaks. In the phylogenetic approach, data were randomly split into 75% and 25% for model training and

testing, respectively. The training set was used iteratively to train three sets of 100 models for

distinguishing oaks from other species, red oaks from white oaks, and diseased red oaks from healthy

red oaks. The trained models were coupled to filter out any observations that do not belong to the red

oak group before running the disease detection step. This filtering process was tested using the initial

25% withheld test data. The whole process was iterated 100 times using different subsets of data to

generate uncertainty around the performance estimates of the model. All classification results

presented in the text utilize the 25% withheld data sets. See Table S3 for sample sizes within each step.

In the phylogenetic approach, we chained three distinctive PLS-DA types to solve the oak wilt

classification problem sequentially through the steps illustrated in Fig. 1. First, we split our data into

75:25 randomly sampled subsets and left the 25% aside to test the overall performance of the

phylogenetic approach at the end of the process (see below). Second, we used the 75% to train three

types of PLS-DAs specifically aimed to distinguish 1) oaks from other species, 2) red oaks from white

oaks, and 3) diseased red oaks from healthy red oaks. Accordingly, each model type had a different data

structure: data from all species for PLS-DAs that distinguished oaks from other species, data belonging

to the red and white oak group only for PLS-DAs that distinguished red from white oaks, and data

including only putative red for PLS-DAs that distinguished diseased from healthy red oaks. All three PLS-

DA types were performed following the same iterative approach described above by randomly sampling

a subset of the 75% of the data for training, testing against the unused data of the subset (a 25% of the

75%), and assessing predictive performance of each PLS-DA. The purpose of these iterations was not to

average model coefficients but rather to test how well PLS-DA types perform on average by generating

confidence intervals for model performance estimates. We assessed performance of each PLS-DA type

by calculating the average and standard deviation of the performance estimates across all iterations. We

ran a total of 100 iterations for each PLS-DA type, thus obtaining 100 separate models of each type

capable of distinguishing either oaks from other species, red oaks from white oaks, or diseased red oaks

from healthy red oaks.

Finally, as an independent validation, we sequentially applied the 100 models of each PLS-DA

type to the 25% of data originally set aside through another 100 iterations. During each iteration, the

25% subset containing all species was first split into 75:25 randomly sampled subsets (stratified by class,

i.e., taxonomic grouping or health status) and only the 75% of the data were used with the aim of

generating variation among iterations. In the first step of the phylogenetic pipeline, the selected data—

which included all species —were classified as either oak or "other species" using the oak discrimination

model. Then, the data classified as oak were classified as either "red" or "white oak" using the red oak

discrimination model. Lastly, the data classified as red oak were classified as either "diseased" or

"healthy red oak" using the disease discrimination model. Data classified as "other species", "white

oak", or "healthy red oak" were later reclassified as "other" and their predicted classes were compared

to their true identities to evaluate predictive performance. The full phylogenetic approach was iterated

100 times to ensure that the initial 75% split reflected all the existing variability within the dataset.

Hence, we report performance across a total of 10,000 (100x100) models of each type (Fig. 1, see Table

S3 for sample sizes and performance). We assessed overall performance of the phylogenetic approach

by calculating the average and standard deviation of the multistep classification performance outputs

across all iterations. Additionally, we applied the oak, and red oak PLS-DA models to the additional 83

oaks (48 *Q. ellipsoidalis* and 35 *Q. macrocarpa*) that were not included in any of the training or testing

steps of our models as a second validation step.

Finally, we performed 100 direct PLS-DAs to classify the 12 dominant species present in our

study area to identify those potentially causing misclassification of red oaks.

To determine which combination of wavelengths was most useful for detection of oak wilt, we

extracted wavelength importance factors from PLS-DAs corresponding to AISA Eagle and AVIRIS-NG

VSWIR and for both direct and phylogenetic approaches using the varImp() function in *caret* (Kuhn,

2008). We focused on these four PLS-DAs because they included the full range of wavelengths covered

by each sensor with and without considering species identity. For simplicity, we limited our selection to

the top 20 wavelengths with the highest average importance across all iterations within each model. We

also extracted wavelength importance factors from AVIRIS-NG VNIR models to compare important

wavelengths with the AISA Eagle models and identify shared wavelengths between both sensors. Lastly,

we cross-applied AISA Eagle and AVIRIS-NG VNIR models to their respective testing datasets to assess

the extent to which models developed with one sensor can be applied to data obtained from other

sensors with similar range of wavelengths (i.e., sensor specificity).

To assess whether reflectance indices associated with physiology could distinguish healthy red

oaks from those infected with oak wilt, we used ANOVA to perform pairwise comparisons between

healthy and diseased red oaks across all spectral reflectance indices and for both AISA Eagle and AVIRIS-

NG. Finally, we compared the effect sizes of these pairwise comparisons using Cohen's *d* statistic

(Cohen, 1988) to assess differences in the detectability of oak wilt between late July and late August.

2.5 PLS-DA mapping

As proof of concept, we mapped PLS-DA model outputs for predicted oak pixels, red oak pixels, and

diseased red oak pixels across the landscape using an AISA Eagle flight line. We used AISA Eagle rather

than AVIRIS-NG because the lower number of wavelengths significantly reduced the computational

resources and time needed to generate the maps. First, we resampled and vector-normalized every

pixel with the flight line using the same functions from the *spectrolab* package to match the spectral

data used to train and test oak, red oak, and diseased red oak AISA Eagle PLS-DA models. For each

model type (oak, red oak, and diseased red oak), we applied the coefficients of 100 randomly chosen

model iterations of the 10,000 produced to each pixel of the resampled and normalized flight line using

the function "predict" from package *car* (Fox and Weisberg, 2019). The result was 100 values per pixel—

and thus 100 maps- describing the probability of being an oak, red oak, and a diseased red oak. We

averaged the 100 values to obtain a single map with probability values representative of the average

prediction. Then, we used a probability threshold of 0.5 to determine pixels with high certainty of being

oaks, red oaks, and a 0.8 threshold for diseased red oaks to counter false positives resulting from the

AISA model and generate masks to select pixels of the targeted classes. Lastly, we applied the masks to

the original flight line both seen through true color and a combination of CMS, CI, and VOG2 indices in

the red, green, and blue channels, respectively, to obtain landscape maps of oaks, red oaks, and

diseased red oaks.

3. Results

All classification accuracy results are reported for the sets of 25% of samples withheld from the PLS-DA

modeling steps, with the standard deviation calculated across the 10,000 iterations performed. All

classification results are reported in Table S3.

<u>3.1 Tree species identities can be predicted from canopy spectral reflectance</u>

The tree species classification PLS-DA demonstrated that it is possible to accurately identify

most of our 12 study species from spectral reflectance (AISA Eagle: 82.0% (±1.2%) correctly identified,

AVIRIS-NG VSWIR: 90.0% (±1.3%), Appendix S2). Models correctly classified and differentiated white

oaks (*Q. macrocarpa*) (AISA Eagle: 85.4% (±3.1%), AVIRIS-NG VSWIR: 94.7% (±1.3%)) and red oaks (*Q.*

ellipsoidalis and *Q. rubra*) (AISA Eagle: 91.0% (±3.7%), AVIRIS-NG VSWIR: 99.4% (±1.4%)). However,

models classifying the oak genus as a whole had higher accuracies (90.1% (±4.9%) and 98.6% (±2.9%) for

AISA Eagle and AVIRIS-NG VSWIR, respectively, Appendix S3) than individual species models (Appendix

S4), similar to results from leaf level spectra (Cavender-Bares et al., 2016). Accordingly, the validation

steps also had higher performance when considering the oak genus as a whole (85.7% (±3.8%) and

94.7% (±2.5%) for AISA Eagle and AVIRIS-NG VSWIR, respectively, Appendix S3) instead of white and red

oaks separately (59.9% (±3.7%) and 85.1% (±4.7%) for AISA Eagle and AVIRIS-NG VSWIR, respectively,

Appendix S4).

<u>3.2 Spectral reflectance models detected diseased red oaks</u>

Spectral reflectance models did not accurately distinguish diseased red oaks from other trees

unless red oaks were first distinguished from other species (Table S3). In the direct approach, overall

model accuracy was significantly better than expected by chance (AISA Eagle: 58.9% (±5.0%),

components (k) = 18; AVIRIS-NG VSWIR: 68.8% (±8.5%), k = 29, Fig. 2), but only healthy trees (true

negatives) were correctly classified with high accuracy (AISA Eagle: 98.7% (±0.77%), AVIRIS-NG VSWIR:

97.5% (±1.4%), indicating high model specificity). Diseased red oaks were correctly classified (true

positives) only 19.1% (±9.13%) and 40.0% (±15.5%) of the AISA Eagle and AVIRIS-NG VSWIR cases,

respectively, indicating low model sensitivity (Fig. 2). As a result, isolating oaks and then red oaks

through a stepwise phylogenetic PLS-DA model prior to disease detection increased correct

classifications and improved the overall performance of both AISA Eagle and AVIRIS-NG VSWIR models

(AISA Eagle: 75.4% (±6.4%), k = oaks: 16, red oaks: 10, diseased red oaks: 21; AVIRIS-NG VSWIR: 84.2%

(±7.7%), k = oaks: 15, red oaks: 10, diseased red oaks: 16; Appendix S3-5, Table S3). The increase in

performance was mostly due to a major increase in correct classification (true positives) of diseased red

oaks (AISA Eagle: 54.6% (±11.5%), AVIRIS-NG VSWIR: 71.0% (±14.0%)) (Fig. 2, Appendix S5) resulting in

increased   model   sensitivity   compared   to   the   direct   PLS-DA   approach.

Figure 2: A stepwise phylogenetic classification approach enhanced detection of oak wilt in red oaks. Models that included both VNIR and SWIR wavelengths (AVIRIS-NG VSWIR) showed better prediction capacity than models including VNIR only. The short-wave infrared (SWIR) range is responsible for the increased predictive performance of AVIRIS-NG VSWIR relative to AVIRIS-NG VNIR PLS-DA models in the direct and phylogenetic modelling approaches. Blue and red circles represent correct and incorrect classifications, respectively. The size and color intensity of the circle represent the average percentage of classifications into each group based on the 25% of data withheld from 10,000 model-fitting iterations, one standard deviation is shown in parentheses. Grey boxes describe the overall predictive performance for a given approach and dataset. Red and blue circles in colored inset boxes above each phylogenetic model describe the performance of the steps within the phylogenetic model at discriminating oaks (gold), red oaks (red), and diseased red oaks (purple), respectively. The number of components used for each model or model step (O = oaks, R = red oaks, D = diseased) is given at the top left corner of the plot. See Appendices S3, S4, and S5 and Table S3 for detailed performance of the phylogenetic steps).

All steps within the phylogenetic PLS-DA model showed high performance (Table S3). The oak detection step showed high accuracy (AISA Eagle: 91% (±5.55%), k = 16; AVIRIS-NG VSWIR: 97% (±6.5%), k = 15) and correctly classified oaks in 90.7% (±6.2%) and 95.1% (±3.6%) of the AISA Eagle and AVIRIS-NG VSWIR cases, respectively (Appendix S3). Similarly, the red oak detection step showed high accuracy (AISA Eagle: 88% (±3.4%), k = 10; AVIRIS-NG VSWIR: 97% (±1.35%), k = 10) and correctly classified red oaks in 91.0% (±3.7%) and 99.4% (±1.4%) of the AISA Eagle and AVIRIS-NG VSWIR cases, respectively (Appendix S4). Finally, the diseased red oak detection step also showed high accuracy (AISA Eagle: 84% (±3.2%), k = 21; AVIRIS-NG VSWIR: 91% (±3.45%), k = 16) and correctly classified diseased red oaks in 78.4% (±3.1%) and 86.5% (±3.4%) of the AISA Eagle and AVIRIS-NG VSWIR cases, respectively (Appendix

S5). The complexity of the model in this last step was higher in AISA Eagle (k = 21) than in AVIRIS-NG

VSWIR models (k = 16). We note, however, that models were sensor-specific. AISA Eagle models failed

to correctly distinguish classes when challenged with AVIRIS-NG VNIR data (Table S4). Similarly, AVIRIS-

NG VNIR models failed to correctly distinguish classes when challenged with AISA Eagle data (Table S4).

<u>3.3 VNIR and SWIR ranges are both important in detecting oak wilt</u>

AVIRIS-NG SWIR models showed slightly higher classification accuracy (true positive rate) of

diseased trees than AISA Eagle and AVIRIS-NG VNIR models in both direct (AVIRIS-NG SWIR: 63.7%

(±7.75%), AISA Eagle: 58.9% (±5.0%), AVIRIS-NG VNIR: 55.6% (±5.45%)) and phylogenetic approaches

(AVIRIS-NG SWIR: 78.9% (±8.4%), AISA Eagle: 75.4% (±6.4%), AVIRIS-NG VNIR: 72.6% (±8.65%)) (Fig. 2).

When both AVIRIS-NG VNIR and SWIR were used together, models outperformed those using either

VNIR or SWIR only. This was the case under both direct (AVIRIS-NG VSWIR: 68.8% (±8.45)) and

phylogenetic approaches (AVIRIS-NG VSWIR: 84.2% (±7.7)).

When differentiating oaks from other species using AISA Eagle models, important wavelengths

were clustered within the 500-560 nm, 600-620 nm, 660-690 nm, and the 970-980 nm regions of the

VNIR range (Fig. 3). From these, wavelengths located around 550 nm and 675 nm where also important

in AVIRIS-NG VNIR models. However, in AVIRIS-NG models that included both VNIR and SWIR, the

importance of these regions was outweighed by regions 1200-1300 nm, 1440 nm, 1600-1750 nm, and

2230-2400 nm within the SWIR range. When differentiating red oaks from white oaks using AISA Eagle

models, we found important wavelengths clustered within the 400-425 nm, 700-770 nm, and 925-980

nm regions. From these, wavelengths located at 420 nm and around 725 nm and 940 nm where also

important in AVIRIS-NG VNIR models. In AVIRIS-NG VSWIR models, the importance of these VNIR

regions remained high, but several regions within the SWIR range showed similar degree of importance.

Within the SWIR, the important wavelengths were clustered at 1200 nm, 1440 nm, around 1490-1550

nm, and around 1700 nm. When differentiating healthy red oaks from diseased red oaks using AISA

Eagle models, important wavelengths appeared at 725-750 nm and across the 800-980 nm region of the

VNIR range. Wavelengths located around 725 nm, at 810 nm, 860, around 950 nm, and at 980 nm where

also important in AVIRIS-NG VNIR models. AVIRIS-NG VSWIR models also identified as important

wavelengths around 810 nm and 970 nm, but also identified important wavelengths within the SWIR

range such as wavelengths around 1160 nm, 1260 nm, 1600 to 1750 nm, 1975 to 2050 nm and 2350 to

2400 nm. Most importantly, the twenty most important wavelengths for detection of oaks, red oaks,

and diseased red oaks rarely overlapped in the AVIRIS-NG VSWIR models (Fig. 3). This was not the case

for AISA Eagle models where most of the important wavelengths for red oak detection were the same

than for diseased red oak detection.

Figure 3: The twenty most important wavelengths—based on variable importance in projection (VIP)—differed among steps discriminating oaks (gold) from other species, red oaks (red) from white oaks, and diseased red oaks (purple) from healthy red oaks, and among models using either VNIR range (AISA Eagle) or both VNIR and SWIR ranges (AVIRIS-NG VSWIR). Vertical lines with numbers indicate wavelengths used in spectral indices associated with photosynthetic capacity (green), photoprotective pigment content (yellow), and water status (blue) that showed

significant differences between healthy and oak wilt-infected trees. Numbers indicate spectral indices

SIPI (1), PRIM4 (2), TCARI/OSAVI (3), CMS (4), SR_{SIF} (5), CRI700 (6), VOG2 (7), CI (8), RDVI (9), SR (10),

NDWI (11), WBI (12), WBI-SWIR (13), PRIm1 (14), CCI (15).

<u>3.4 Declines in photosynthetic capacity and water status signal oak wilt</u>

Overall, spectral indices calculated from the AISA Eagle dataset collected during July showed

slightly more pronounced differences between healthy and diseased red oaks than those calculated

from the AVIRIS-NG dataset collected in August (Fig. 4). However, confidence intervals overlapped

between both sensors for all photosynthetic and water status indices. Spectral indices associated with

canopy photosynthetic capacity showed significant differences between healthy and diseased red oaks

for both sensors and time periods. Within the AISA Eagle dataset, all indices associated with

photosynthetic capacity were significantly different between healthy and diseased red oaks (Table S4).

Within the AVIRIS-NG dataset, all indices associated with photosynthetic capacity except NPCI were

significantly different between healthy and diseased red oaks. Most spectral indices associated with

photoprotective stress only showed significant differences between healthy and diseased red oaks

during the earliest flight (AISA Eagle). Those that were always significant—Photochemical Reflectance

Index (PRIm4) and Carotenoid Reflectance Index (CRI700)—share wavelengths with indices of

photosynthetic capacity, such as the SR and Transformed Chlorophyl Absorption in Reflectance

Index/Optimized Soil-Adjusted Vegetation Index (TCARI/OSAVI) indices. Indices associated with canopy

water status also showed significant differences between healthy and diseased red oaks. The effect sizes

of the differences were comparable to those of indices associated with photosynthetic capacity.

Figure 4: Spectral indices associated with photosynthetic (green) and water status (blue) differentiated

diseased red oaks from healthy red oaks. Indices associated with photoprotective pigments (gold) only

do so when collected in July or when they also include wavelengths associated with photosynthetic

capacity. Each point represents the magnitude of the difference between healthy and diseased trees—

shown by the absolute value of the Cohen's d—for a given index and sensor type (AISA Eagle collected in

July or AVIRIS-NG collected in August). Effect size can be understood as the amount of overlap between

the distributions of two groups. For an effect size of 0, the mean of group 2 falls within the 50th

percentile of group 1, and the distributions overlap completely, meaning there is no difference between

them. For an effect size of 0.8, the mean of group 2 falls within the 79th percentile of group 1; thus, an

average sample from group 2 would have a higher value than 79% of all samples from group 1 (Sullivan

and Feinn, 2012). Lines represent 95% confidence intervals. Effect sizes are significantly different from

zero when their confidence intervals do not overlap with the red zero line.

4. **Discussion**

The negative impacts of oak wilt and its rate of spread across North American ecosystems calls for detection tools that accurately identify trees affected by oak wilt at landscape scales (Haight et al., 2011; Hulcr and Dunn, 2011; Juzwik et al., 2011). We show that PLS-DA models developed from airborne spectroscopic imagery can detect oak wilt-infected red oaks. We demonstrate an approach to identify oak wilt-infected red oaks, which takes advantage of the physiological and phylogenetic information embedded in their reflectance spectra (Cavender-Bares et al., 2016; Meireles et al., 2020a). By first differentiating oaks from non-oaks, and then identifying red oaks—which are highly susceptible to rapid disease development—classification models based on spectral reflectance data can be used to distinguish oak-wilt affected and healthy red oaks with high accuracy. We also found that spectral indices associated with plant photosynthesis, water status, and photoprotective pigments are potentially sensitive to disease progression through physiological decline. Spectral indices also provide a mechanistic basis for understanding and tracking the physiological changes that allow classification models to detect oak wilt and are consistent with findings from other oak decline systems (Encinas-Valero et al. 2021).

4.1 Including short wave infrared reflectance improves model accuracy

Including SWIR wavelengths in spectral reflectance models increases oak wilt detection accuracy. We observed higher oak wilt detectability in direct AVIRIS-NG SWIR and VSWIR than direct AISA Eagle and AVIRIS-NG VNIR models. Direct PLS-DAs using AVIRIS-NG VNIR showed similar performance to that of AISA Eagle VNIR models (Fig. 2). We can therefore attribute the greater performance of direct AVIRIS-NG VSWIR models to the addition of SWIR wavelengths. Direct AISA Eagle models rely on many of the same wavelengths to distinguish red oaks from white oaks and to distinguish diseased and healthy red oaks (Fig. 3). As such, they often misclassify diseased red oaks as white oaks

(Fig. 2, Appendix S4). However, even direct models show much higher accuracy when both VNIR and

SWIR ranges are included (AVIRIS-NG VSWIR). The additional information-rich SWIR wavelengths allow

models to use different wavelength regions to distinguish oaks from other species, red oaks from white

oaks, and diseased red oaks from healthy red oaks (Fig. 2). Consequently, the critical wavelengths to

identify oaks, red oaks, and diseased red oaks overlap less, which reduces the chances of confusion

among classes (Fig. 3 and validation steps in Appendices S3 & 4). Most likely, including SWIR reflectance

provides temporally stable spectral features containing phylogenetic information associated with plant

structural traits (Cavender-Bares et al., 2020; Meireles et al., 2020a) that serve to reduce

misclassification of diseased red oaks as white oaks and other species. Indeed, we find that the SWIR

wavelengths were more often represented than VNIR wavelengths in oak models and as represented as

VNIR wavelengths in red oak models (Fig. 3). In particular, 17 and 11 of the most important wavelengths

for identifying oaks and red oaks, respectively, fell within the SWIR range. The SWIR was also important

for distinguishing diseased from healthy red oaks. Among the most important SWIR wavelengths were

those associated to plant water content and leaf chemistry such as protein, sugars, lignin, and cellulose

content (Asner et al., 2018; Fourty et al., 1996). Based on our results, the SWIR range appears to contain

disease-specific and phylogenetic information highly relevant to detecting symptoms of oak wilt and to

identifying its hosts. Hence, similar to previous work combining VNIR reflectance with SIF or thermal

data (Zarco-Tejada et al., 2018, 2016), when SWIR wavelengths are combined with VNIR in oak wilt

detection models, detection rates are maximized.

4.2 A multi-step phylogenetic approach increases accuracy

Partitioning the classification process into simple binary steps within a phylogenetic framework

reduces potential misclassification and increases model accuracy. We used a hierarchical classification

approach (Allen and Walsh, 1996; Townsend et al., 2009; Wolter et al., 1995) aimed towards

distinguishing the more susceptible red oaks from white oaks and other species that are less susceptible

to oak wilt. During the first step, phylogenetic models distinguish between oaks and other species

because the reflectance spectrum shows phylogenetic conservatism among the oaks (Cavender-Bares et

al., 2016; Cavender-Bares, 2019), including those infected by oak wilt. The model is not required to

distinguish between healthy and diseased conspecifics in this first step, thus simplifying the task.

Reducing the number of potential classes becomes increasingly important as the individuals become

more phylogenetically related—and hence more phenotypically similar—which makes correct

classification more challenging (Meireles et al., 2020b). Removing non-oak species significantly reduces

variation in phylogenetically conserved regions of the spectra, allowing the model to be trained on

spectral differences that distinguish white and red oaks and subsequently on the spectral variation that

distinguishes diseased and healthy red oaks. Because of these filtering steps, the disease detection

algorithm is highly accurate (84%; Fig. 2, appendix S5) and significantly more accurate than a single-step,

direct approach. While the phylogenetic approach gains complexity in terms of number of steps, each

binary classification step is simple and requires few independent components. Although each step

generates classification errors that propagate through the modeling pipeline, these errors are captured

by the overall performance metrics, indicating that the increase in accuracy gained through the

phylogenetic filtering outweighs the propagated errors. The phylogenetic approach increases accuracy

by reducing the number of classes to compare while inclusion of the SWIR range increases accuracy by

increasing the number of informative wavelengths.

While the PLS-DA models are sensor-specific (Table S4), the benefits of implementing a stepwise

phylogenetic approach or using the widest range of wavelengths possible do seem common across

hyperspectral products. Implementing a stepwise phylogenetic approach boosted model performance

to a similar extent in models that had the same range of wavelengths (i.e., AISA Eagle and AVIRIS-NG

VNIR models). Similarly, reducing the range of wavelengths in AVIRIS-NG models to the VNIR range

reduced model performance to similar values observed in AISA Eagle models. The similar performance

between datasets from different sensors and time periods could suggest that the benefits of adding

SWIR wavelengths and a stepwise phylogenetic approach are insensitive to sensor type or time of the

year. Furthermore, despite the sensor-specificity of the models, several wavelengths were commonly

highlighted as important across sensor types in oak, red oak, and diseased red oak models (Figure S3,

see results section). These wavelengths were located around the green band and close to the red edge

in case of the oak model. In the case of the red oak and diseased red oak models, shared wavelengths

were also close to the red edge, and representative of starch and water content (Curran, 1989).

Our results highlight that species classification is critical for increasing model accuracy for a

simple reason: if the disease is host-specific, modeling can be more tractable by detecting potential

hosts first. Future studies should test whether phylogenetic models with simple binary classification

steps such as the one used here make disease detection models generic enough to be applicable across

different sites and years.

4.3 Targeted spectral indices help understand physiological changes associated with oak wilt disease

Diseased red oaks were more easily differentiated from healthy red oaks by spectral reflectance

indices associated with photosynthetic activity (Carter and Knapp, 2001; Vogelmann et al., 1993; Zarco-

Tejada et al., 2002) and water status (Ceccato et al., 2001; Serrano et al., 2000; Ullah et al., 2014) than

by photoprotective pigment content indices. Indices based on photoprotective pigment only

differentiated diseased trees from asymptomatic trees in July or when they included wavelengths also

associated with photosynthetic activity (Figs. 3 & 4). These results suggest that oak wilt infection in adult

trees in natural ecosystems first triggers changes in photoprotective pigments (similar to holm oak

decline caused by *Phytopthora spp* fungal infection and drought, Encinas-Valero et al. 2021) followed by

declines in photosynthetic rate, stomatal conductance, and water content just as in greenhouse red oak

seedlings infected with oak wilt (Fallon et al., 2020). Our results uncover an important temporal pattern

in physiological decline: photoprotective pigment content ceases to be a good predictor of oak wilt by

the end of summer. By July, indices of photoprotective pigment content showed somewhat similar

sensitivity to oak wilt than indices of photosynthetic capacity or water status (Fig. 4). At early stages,

both tylose production (in response to infection) and plugging of vessels by metabolites of the fungus

are likely to have contributed to diminished water transport. In turn, reduced transpiration and stomatal

closure induced by reduced water supply is expected to have caused photosynthetic decline (Fallon et

al., 2020) and abnormally increased photoprotective pigment content to deal with light stress resulting

from reduced photosynthesis. By August, healthy trees may also have increased photoprotective

pigment content as summer drought becomes more prevalent and photosynthesis is partially impaired.

Yet, diseased trees have likely experienced greater water deficit due to tylose formation and their

photosynthetic capacity and water status is impaired to a greater extent than drought-stressed but

uninfected trees. This would explain why photosynthetic capacity indices still distinguished diseased and

healthy red oaks and why water status indices showed greater overall sensitivity to oak wilt than

photosynthetic activity indices by August (Fig. 4). The fact that, photosynthetic and water status indices

showed similar performance between sensors suggests that normalized spectral indices should be

comparable and applicable across sensor types to some degree due to their underlying mechanistic

basis. The results are consistent with experimental work indicating that photosynthesis is the first

physiological process to decline as stomata shut down (Fallon et al., 2020) followed by water content as

vessel occlusion develops and the fungus damages cell walls and membranes (e.g., through pathogen-

produced toxins) leading to dehydration and tissue death (Oliva et al., 2014). While other studies have

proposed the use of machine learning algorithms to identify tree species and the use of spectral indices

to detect plant diseases (Abdulridha et al., 2019; Ghosh et al., 2014; Iordache et al., 2020; Pontius, 2014;

Tang et al., 2021), we see potential in coupling both. Pairing phylogenetic approaches to map oak wilt

presence with photosynthetic activity and water status indices can provide powerful tools to delineate

oak wilt centers across areas of the landscape. The steps would entail first mapping red oaks (Fig. S4) -

and diseased red oaks if possible- and then combining spectral indices associated with photosynthetic

and water status, or indices that use oak-wilt sensitive wavelengths to detect potential oak wilt pockets

(Fig. 5). Pockets of affected trees may show a center-outward radial gradient with dry dead trees at the

center, dehydrated and photosynthetically impaired trees in the middle, and trees with slightly lower

photosynthetic capacity than expected around the edge of the pocket (i.e., early disease development

phase) (Figs. 5, S4, S5 & S5). The step of distinguishing oaks and red oaks from other species across the

landscape is highly advantageous for forest managers on its own as it allows them to identify areas at

high risk of infection in which to apply spectral indices for further diagnostic. Hence, phylogenetic

spectral models paired with indices is a novel approach to predict the risk of disease spread based on

the abundance of red oaks across the landscape and the stage of disease development based on

physiological status, thus allowing managers to better assess risk of spread and adjust the magnitude of

their interventions accordingly (Pontius and Hallett, 2014).

Figure 5. A typical oak wilt pocket observed through true color and a combination of spectral indices using the 2016 AVIRIS-NG data. A tree killed by oak wilt during 2015 (orange circle) can be observed at the center of the oak wilt pocket in true color (red as 640 nm, green as 550 nm, and blue as 470 nm). Three diseased trees (blue circles) stand next to it that cannot be detected with true color images. Both dead and diseased trees are surrounded by an outer ring of healthy trees. Diseased trees become apparent through spectral indices associated with photosynthetic function and water status -such as the Carter-Miller Stress index (CMS), Chlorophyll Index red edge (CI), and Water Band Index in the SWIR range (WBI SWIR)- placed on the red (R), green (G), and blue (B) channels.

5. Conclusions

Remote detection greatly enhances the ability of managers to prevent the enormous ecological and economical damage caused by invasive species (Juzwik, 2000; Poland et al., 2021). Airborne spectroscopic imagery enables landscape-level detection of diseases caused by invasive pathogens, like oak wilt, due to the phylogenetic and physiological information embedded in spectral reflectance. SWIR wavelengths increased model accuracy by enabling detection of disease-specific hosts, a critical step in identifying forested areas vulnerable to infection. Additionally, inference of the physiological basis of oak wilt symptom development using spectral indices associated with known spectral features points to the potential to delineate oak wilt centers using remote sensing products that monitor canopy photosynthetic capacity and water status such as VOG2, SIPI, SR_{SIF}, CMS, CI, RDVI, WBI, and NDWI among others. Importantly, in our study landscape detection was made possible by coupling airborne spectroscopic imagery with traditional knowledge from taxonomic and disease experts and high precision ground GPS reference data. While landscape detection of oak wilt will facilitate the task of detecting infected trees, there is still much work to do. Future studies should assess whether PLS-DA

models will be general enough to detect oak wilt across years and sites and whether the physiological

basis of oak wilt symptom development will be sufficient to make accurate inferences about the

presence of new oak wilt infections. Further investigation of the physiological changes that accompany

disease progression may also provide the link to scale spectral detection to regional scales via

spaceborne platforms. The work done here points to the benefit of research that might lead to an

"optimal" remote sensing system (airborne or satellite) for detecting invasive diseases. We hope that

our research motivates such work.

Acknowledgements

This project was funded by the Minnesota Invasive Terrestrial Plants and Pests Center, the National

Science Foundation (NSF) and the National Aeronautics and Space Administration (NASA) through the

Dimensions of Biodiversity program (DEB-1342872 and DEB-1342778), the Cedar Creek Long Term

Ecological Research program (1831944), and the NSF Biology Integration Institute ASCEND (DBI:

2021898). The University of Minnesota, including Cedar Creek ESR, lies on the ancestral, traditional, and

contemporary Land of the Dakota people. We would like to thank Brett Fredericksen, Erin Murdock, Kali

Hall, Lewis French and Travis Cobb for help with tarp measurement for empirical line corrections. Thanks

to Dr. Joe Knight for access to the infrastructure of the Remote Sensing and Geospatial Analysis Lab at

UMN, and to Dan Bahauddin for informatics help which facilitated flight planning at Cedar Creek. We

thank Paul Castillo, USFS for his technical assistance in the training of field technicians, and Zhihui Wang

for assistance processing AVIRIS-NG imagery. We also thank Shan Kothari, Lucy Shroeder, Anna Yang,

Austin Yantes, Clarissa Fontes, Byju Govidan, Artur Stefanski, and Adriana Castillo Castillo for their

comments in previous versions of this manuscript.

References

Abdulridha, J., Batuman, O., Ampatzidis, Y., 2019. UAV-Based Remote Sensing Technique to Detect

Citrus Canker Disease Utilizing Hyperspectral Imaging and Machine Learning. Remote Sens. 11,

1373. https://doi.org/10.3390/rs11111373

Allen, T.R., Walsh, S.J., 1996. Spatial and compositional pattern of alpine treeline, Glacier National Park,

Montana. Photogramm. Eng. Remote Sensing 62, 1261–1268.

Appel, D.N., 1995. The Oak Wilt Enigma: Perspectives from the Texas Epidemic. Annu. Rev. Phytopathol.

33, 103–118. https://doi.org/10.1146/annurev.py.33.090195.000535

Asner, P., Martin, R.E., Keith, L.M., Heller, W.P., Hughes, M.A., Vaughn, R., Hughes, R.F., Balzotti, C.,

2018. A Spectral Mapping Signature for the Rapid Ohia Death (ROD) Pathogen in Hawaiian Forests.

Remote Sens. 10, 1–14. https://doi.org/10.3390/rs10030404

Barker, M., Rayens, W., 2003. Partial least squares for discrimination. J. Chemom.

https://doi.org/10.1002/cem.785

Bergot, M., Cloppet, E., Pérarnaud, V., Déqué, M., Marçais, B., Desprez-Loustau, M.-L., 2004. Simulation

of potential range expansion of oak disease caused by Phytophthora cinnamomi under climate

change. Glob. Chang. Biol. 10, 1539–1552. https://doi.org/10.1111/j.1365-2486.2004.00824.x

Brereton, R.G., Lloyd, G.R., 2014. Partial least squares discriminant analysis: Taking the magic away. J.

Chemom. 28, 213–225. https://doi.org/10.1002/cem.2609

Carter, G.A., Knapp, A.K., 2001. Leaf optical properties in higher plants: linking spectral characteristics to

stress and chlorophyll concentration. Am. J. Bot. https://doi.org/10.2307/2657068

Castaldi, F., Chabrillat, S., Jones, A., Vreys, K., Bomans, B., Wesemael, B. Van, 2018. Soil Organic Carbon

Estimation in Croplands by Hyperspectral Remote APEX Data Using the LUCAS Topsoil Database.

Remote Sens. 10, 2–19. https://doi.org/10.3390/rs10020153

Cavender-Bares, J., Gamon, J.A., Townsend, P.A. (Eds.), 2020. Remote Sensing of Plant Biodiversity,

Remote Sensing of Plant Biodiversity. Springer International Publishing, Cham.

https://doi.org/10.1007/978-3-030-33157-3

Cavender-Bares, J., Holbrook, N.M., 2001. Hydraulic properties and freezing-induced cavitation in

sympatric evergreen and deciduous oaks with contrasting habitats. Plant. Cell Environ. 24, 1243–

1256. https://doi.org/10.1046/j.1365-3040.2001.00797.x

Cavender-Bares, J., Meireles, J.E., Couture, J.J., Kaproth, M.A., Kingdon, C.C., Singh, A., Serbin, S.P.,

Center, A., Zuniga, E., Pilz, G., Townsend, P.A., 2016. Associations of leaf spectra with genetic and

phylogenetic variation in oaks: Prospects for remote detection of biodiversity. Remote Sens. 221,

1–17. https://doi.org/10.3390/rs8030221

Cavender-Bares, J., Nelson, E., Meireles, J.E., Lasky, J., Miteva, D., Nowak, D., Pearse, W., Helmus, M.,

Zanne, A., Fagan, W., Mihiar, C., Muller, N., Kraft, N., Polasky, S., 2019. The hidden value of trees:

quantifying the ecosystem services of tree lineages and their major threats across the continental

US. EcoEvorxiv Prepr. 1–42. https://doi.org/10.32942/osf.io/gp7mt

Cavender-Bares, J., 2019. Diversification, adaptation, and community assembly of the American oaks (

Quercus), a model clade for integrating ecology and evolution. New Phytol. 221, 669–692.

https://doi.org/10.1111/nph.15450

Ceccato, P., Flasse, S., Tarantola, S., Jacquemoud, S., Grégoire, J.M., 2001. Detecting vegetation leaf

water content using reflectance in the optical domain. Remote Sens. Environ. 77, 22–33.

https://doi.org/10.1016/S0034-4257(01)00191-2

Cochard, H., Tyree, M.T., 1990. Xylem dysfunction in Quercus: vessel sizes, tyloses, cavitation and

seasonal changes in embolism. Tree Physiol. 6, 393–407. https://doi.org/10.1093/treephys/6.4.393

Cohen, J., 1988. The effect size index: d, in: Statistical Power Analysis for the Behavioral Sciences.

Conel, J.E., Green, R.O., Vane, G., Bruegge, C.J., Alley, R.E., Curtiss, B.J., 1987. AIS-2 Radiometry and a

Comparison of Methods for the Recovery of Ground Reflectance, in: Third Airborne Imaging

Spectrometer Data Analysis Workshop. Jet Propulsion Lab, Pasadena, CA, pp. 18–47.

Congalton, R.G., 2001. Accuracy assessment and validation of remotely sensed and other spatial

information. Int. J. Wildl. Fire 10, 321. https://doi.org/10.1071/WF01031

Crucil, G., Castaldi, F., Aldana-jague, E., Wesemael, B. Van, Macdonald, A., Van Oost, K., 2019. Assessing

the Performance of UAS-Compatible Multispectral and Hyperspectral Sensors for Soil Organic

Carbon Prediction. Sustainability 11, 2–18. https://doi.org/10.3390/su11071889

Curran, P.J., 1989. Remote sensing of foliar chemistry. Remote Sens. Environ. 30, 271–278.

https://doi.org/10.1016/0034-4257(89)90069-2

Curran, P.J., Windham, W.R., Gholz, H.L., 1995. Exploring the relationship between reflectance red edge

and chlorophyll concentration in slash pine leaves. Tree Physiol. 15, 203–206.

https://doi.org/10.1093/treephys/15.3.203

de Beer, Z.W., Marincowitz, S., Duong, T.A., Wingfield, M.J., 2017. Bretziella, a new genus to

accommodate the oak wilt fungus, Ceratocystis fagacearum (Microascales, Ascomycota). MycoKeys

27, 1–19. https://doi.org/10.3897/mycokeys.27.20657

Denk, T., Grimm, G., Manos, P., Deng, M., Hipp, A., 2017. An Updated Infrageneric Classification of the

Oaks: Review of Previous Taxonomic Schemes and Synthesis of Evolutionary Patterns, in: Gil-

Pelegrín, E., Peguero-Pina, J., Sancho-Knapik, D. (Eds.), Oaks Physiological Ecology. Exploring the

Functional Diversity of Genus Quercus L. Tree Physiology, pp. 13–38.

https://doi.org/https://doi.org/10.1007/978-3-319-69099-5_2

Díaz, S., Settele, J., Brondízio, E.S., Ngo, H.T., Agard, J., Arneth, A., Balvanera, P., Brauman, K.A., Butchart,

S.H.M., Chan, K.M.A., Garibaldi, L.A., Ichii, K., Liu, J., Subramanian, S.M., Midgley, G.F., Miloslavich,

P., Molnár, Z., Obura, D., Pfaff, A., Polasky, S., Purvis, A., Razzaque, J., Reyers, B., Chowdhury, R.R.,

Shin, Y.-J., Visseren-Hamakers, I., Willis, K.J., Zayas, C.N., 2019. Pervasive human-driven decline of

life on Earth points to the need for transformative change. Science (80-.). 366, eaax3100.

https://doi.org/10.1126/science.aax3100

Encinas-Valero, M., Esteban, R., Heres, A.-M., Becerril, J.-M., Garcia-Plazaola, J.-I., Artexe, U., Vivas, M.,

Solla, A., Moreno, G., Curiel-Yuste, J., 2021. Photoprotective compounds as early markers to

predict holm oak crown defoliation in declining Mediterranean savannahs. Tree Physiol. tpab006.

https://doi.org/https://doi.org/10.1093/treephys/tpab006

ESRI, 2011. ArcGIS Desktop: Release 10. Redlands, CA: Environmental Systems Research Institute.

Redlands.

Evans, E.A., Crane, J., Hodges, A., Osborne, J.L., 2010. Potential economic impact of laurel wilt disease on

the Florida avocado industry. Horttechnology 20, 234–238.

https://doi.org/10.21273/horttech.20.1.234

Fallon, B., Yang, A., Lapadat, C., Armour, I., Juzwik, J., Montgomery, R.A., Cavender-bares, J., 2020.

Spectral differentiation of oak wilt from foliar fungal disease and drought is correlated with

physiological changes. Tree Physiol. 40, 377–390. https://doi.org/10.1093/treephys/tpaa005

Fassnacht, K.S., Cohen, W.B., Spies, T.A., 2006. Key issues in making and using satellite-based maps in

ecology: A primer. For. Ecol. Manage. https://doi.org/10.1016/j.foreco.2005.09.026

Feilhauer, H., Asner, G.P., Martin, R.E., Schmidtlein, S., 2010. Brightness-normalized Partial Least Squares

Regression for hyperspectral data. J. Quant. Spectrosc. Radiat. Transf. 111, 1947–1957.

https://doi.org/10.1016/j.jqsrt.2010.03.007

Fernandes, A.M., Fortini, E.A., Müller, L.A. de C., Batista, D.S., Vieira, L.M., Silva, P.O., Amaral, C.H. do,

Poethig, R.S., Otoni, W.C., 2020. Leaf development stages and ontogenetic changes in passionfruit

(Passiflora edulis Sims.) are detected by narrowband spectral signal. J. Photochem. Photobiol. B

Biol. https://doi.org/10.1016/j.jphotobiol.2020.111931

Fourty, T., Baret, F., Jacquemoud, S., Schmuck, G., Verdebout, J., 1996. Leaf optical properties with

explicit description of its biochemical composition: Direct and inverse problems. Remote Sens.

Environ. 56, 104–117. https://doi.org/10.1016/0034-4257(95)00234-0

Fox, J., Weisberg, S., 2019. An R Companion to Applied Regression, 3rd editio. ed. Sage, Thousand Oaks,

CA.

Gamon, J.A., Surfus, J.S., 1999. Assessing leaf pigment content and activity with a reflectometer. New

Phytol. 143, 105–117. https://doi.org/10.1046/j.1469-8137.1999.00424.x

Gholizadeh, H., Gamon, J.A., Townsend, P.A., Zygielbaum, A.I., Helzer, C.J., Hmimina, G.Y., Yu, R., Moore,

R.M., Schweiger, A.K., Cavender-Bares, J., 2019. Detecting prairie biodiversity with airborne remote

sensing. Remote Sens. Environ. 221, 38–49. https://doi.org/10.1016/j.rse.2018.10.037

Ghosh, A., Fassnacht, F.E., Joshi, P.K., Koch, B., 2014. A framework for mapping tree species combining

hyperspectral and LiDAR data: Role of selected classifiers and sensor across three spatial scales. Int.

J. Appl. Earth Obs. Geoinf. 26, 49–63. https://doi.org/10.1016/j.jag.2013.05.017

Gibbs, J.N., French, D.W., 1980. The transmission of oak wilt, USDA Forest Service, North Central

Research Station.

Grigal, D.F., Chamberlain, L.M., Finney, H.R., Wroblewki, D.V., Gross, E.R., 1974. Soils of the Cedar Creek

Natural History Area, Miscelaneous report. St. Paul.

Haight, R.G., Homans, F.R., Horie, T., Mehta, S. V., Smith, D.J., Venette, R.C., 2011. Assessing the Cost of

an Invasive Forest Pathogen: A Case Study with Oak Wilt. Environ. Manage. 47, 506–517.

https://doi.org/10.1007/s00267-011-9624-5

Hamlin, L., Green, R.O., Mouroulis, P., Eastwood, M., Mccubbin, I., Wilson, D., Randall, D., Dudik, M.,

2010. Imaging Spectrometer Science Measurements for Terrestrial Ecology: AVIRIS and the Next

Generation AVIRIS Characteristics and Development Status, in: NASA Earth Science Technology

Forum. pp. 1–7.

Hanavan, R.P., Pontius, J., Hallett, R., 2015. A 10-Year Assessment of Hemlock Decline in the Catskill

Mountain Region of New York State Using Hyperspectral Remote Sensing Techniques. J. Econ.

Entomol. 1–11. https://doi.org/10.1093/jee/tou015

Hornero, A., Zarco-Tejada, P.J., Quero, J.L., North, P.R.J., Ruiz-Gómez, F.J., Sánchez-Cuesta, R.,

Hernandez-Clemente, R., 2021. Modelling hyperspectral- and thermal-based plant traits for the

early detection of Phytophthora-induced symptoms in oak decline. Remote Sens. Environ. 263,

112570. https://doi.org/10.1016/j.rse.2021.112570

Hulcr, J., Dunn, R.R., 2011. The sudden emergence of pathogenicity in insect–fungus symbioses

threatens naive forest ecosystems. Proc. R. Soc. B Biol. Sci. 278, 2866–2873.

https://doi.org/10.1098/rspb.2011.1130

Iordache, M.-D., Mantas, V., Baltazar, E., Pauly, K., Lewyckyj, N., 2020. A Machine Learning Approach to

Detecting Pine Wilt Disease Using Airborne Spectral Imagery. Remote Sens. 12, 2280.

https://doi.org/10.3390/rs12142280

Jacobi, W., MacDonald, W., 1980. Colonization of Resistant and Suceptible Oaks by Ceratocystis

Fagacearum. Phytopathology 70, 618–623.

Jacquemoud, S., Ustin, S.L., 2001. Leaf optical properties: a state of the art. Proc. 8th Int. Symp. Phys.

Meas. Signatures Remote Sens. https://doi.org/10.1017/CBO9781107415324.004

Johnson, P.S., Shifley, S.R., Rogers, R., 2019. The ecology and silviculture of oaks, The ecology and

silviculture of oaks. CABI, Wallingford. https://doi.org/10.1079/9781780647081.0000

Juzwik, J., 2009. Epidemiology and occurrence of oak wilt in Midwestern, middle, and south Atlantic

states., in: Proceedings of the National Oak Wilt Symposium. Texas Forest Service Publication,

Austin, TX, pp. 49–60.

Juzwik, J., 2000. An oak wilt primer. Int. Oaks 11, 14–20.

Juzwik, J., Appel, D.N., 2016. Diseases of trees in the Great Plains: Oak Wilt, in: Diseases of Trees in the

Great Plains. USDA, Forest Service, Rocky Mountain Forest and Range Station, Fort Collins, Colo.:,

pp. 129–133. https://doi.org/Gen. Tech. Rep. RMRS-GTR-335

Juzwik, J., Appel, D.N., MacDonald, W.L., Burks, S., 2011. Challenges and Successes in Managing Oak Wilt

in the United States. Plant Dis. 95, 888–900. https://doi.org/10.1094/PDIS-12-10-0944

Juzwik, J., French, D.W., 1983. Ceratocystis fagacearum and C. piceae on the Surfaces of Free-Flying and

Fungus-Mat-Inhabiting Nitidulids. Phytopathology 73, 1164–1168. https://doi.org/10.1094/phyto-

73-1164

Koch, K.A., Quiram, G.L., Venette, R.C., 2010. A review of oak wilt management: A summary of

treatment options and their efficacy. Urban For. Urban Green. 9, 1–8.

https://doi.org/10.1016/j.ufug.2009.11.004

Kuhn, M., 2008. Building Predictive Models in R Using the caret Package. J. Stat. Softw. 28.

https://doi.org/10.18637/jss.v028.i05

Lausch, A., Heurich, M., Gordalla, D., Dobner, H.J., Gwillym-Margianto, S., Salbach, C., 2013. Forecasting

potential bark beetle outbreaks based on spruce forest vitality using hyperspectral remote-sensing

techniques at different scales. For. Ecol. Manage. 308, 76–89.

https://doi.org/10.1016/j.foreco.2013.07.043

Liebhold, A.M., MacDonald, W.L., Bergdahl, D., Mastro, V.C., 1995. Invasion by Exotic Forest Pests: A

Threat to Forest Ecosystems. For. Sci. 41, a0001-z0001.

https://doi.org/10.1093/forestscience/41.s1.a0001

Malenovský, Z., Ufer, C., Lhotáková, Z., Clevers, J.G.P.W., Schaepman, M.E., Albrechtová, J., Cudlín, P.,

2006. a New Hyperspectral Index for Chlorophyll Estimation of a Forest Canopy : Area Under Curve

Normalised To Maximal Band Depth Between 650-725 Nm. EARSeL eProceedings 5, 161–172.

Meireles, J.E., Cavender-Bares, J., Townsend, P.A., Ustin, S., Gamon, J.A., Schweiger, A.K., Schaepman,

M.E., Asner, G.P., Martin, R.E., Singh, A., Schrodt, F., Chlus, A., O'Meara, B.C., 2020a. Leaf

reflectance spectra capture the evolutionary history of seed plants. New Phytol. nph.16771.

https://doi.org/10.1111/nph.16771

Meireles, J.E., O'Meara, B., Cavender-Bares, J., 2020b. Linking Leaf Spectra to the Plant Tree of Life, in:

Remote Sensing of Plant Biodiversity. Springer International Publishing, Cham, pp. 155–172.

https://doi.org/10.1007/978-3-030-33157-3_7

Menges, E.S., Loucks, O.L., 1984. Modeling a Disease-Caused Patch Disturbance: Oak Wilt in the

Midwestern United States. Ecology 65, 487–498. https://doi.org/10.2307/1941411

Nouri, M., Gomez, C., Gorretta, N., Roger, J.M., 2017. Clay content mapping from airborne hyperspectral

Vis-NIR data by transferring a laboratory regression model. Geoderma 298, 54–66.

https://doi.org/10.1016/j.geoderma.2017.03.011

Ogaya, R., Barbeta, A., Başnou, C., Peñuelas, J., 2015. Satellite data as indicators of tree biomass growth

and forest dieback in a Mediterranean holm oak forest. Ann. For. Sci. 72, 135–144.

https://doi.org/10.1007/s13595-014-0408-y

Oliva, J., Stenlid, J., Martinez-Vilalta, J., 2014. The effect of fungal pathogens on the water and carbon

economy of trees: implications for drought-induced mortality. New Phytol. 203, 1028–1035.

Ollinger, S. V., 2011. Sources of variability in canopy reflectance and the convergent properties of plants.

New Phytol. 189, 375–394. https://doi.org/10.1111/j.1469-8137.2010.03536.x

Pellegrini, A., Hein, A.M., Cavender-bares, J., Montgomery, R.A., Staver, A.C., Hobbie, S.E., Reich, P.B.,

2021. Disease and fire interact to influence transitions between savanna – forest ecosystems over

a multi-decadal experiment. Ecol. Lett. 24, 1007–1017. https://doi.org/10.1111/ele.13719

Poland, T.M., Patel-Weynand, T., Finch, D., Fort Miniat, C., Hayes, D.C., Lopez, V., 2021. Invasive Species

in Forests and Grasslands of the United States: A Comprehensive Science Synthesis for the United

States Forest Sector. Springer Verlag, Cham, Switzerland.

Pontius, J., 2014. A new approach for forest decline assessments: Maximizing detail and accuracy with

multispectral imagery. Int. J. Remote Sens. 35, 3384–3402.

https://doi.org/10.1080/01431161.2014.903439

Pontius, J., Hallett, R., 2014. Comprehensive methods for earlier detection and monitoring of forest

decline. For. Sci. https://doi.org/10.5849/forsci.13-121

Pontius, J., Hallett, R., Martin, M., 2005. Using AVIRIS to assess hemlock abundance and early decline in

the Catskills, New York. Remote Sens. Environ. 97, 163–173.

https://doi.org/10.1016/j.rse.2005.04.011

Pontius, J., Martin, M., Plourde, L., Hallett, R., 2008. Ash decline assessment in emerald ash borer-

infested regions: A test of tree-level, hyperspectral technologies. Remote Sens. Environ. 112,

2665–2676. https://doi.org/10.1016/j.rse.2007.12.011

R Development Core Team, 2020. R: A Language and Environment for Statistical Computing. version

3.6.0.

Ramirez, J.A., Posada, J.M., Handa, I.T., Hoch, G., Vohland, M., Messier, C., Reu, B., 2015. Near-infrared

spectroscopy (NIRS) predicts non-structural carbohydrate concentrations in different tissue types

of a broad range of tree species. Methods Ecol. Evol. 6, 1018–1025. https://doi.org/10.1111/2041-

210X.12391

Romero, A., Aguado, I., Yebra, M., 2012. Estimation of dry matter content in leaves using normalized

indexes and prospect model inversion. Int. J. Remote Sens. 33, 396–414.

https://doi.org/10.1080/01431161.2010.532819

Schoenweiss, D.F., 1959. Xylem formation as a factor in oak wilt resistance. Phytopathology 49, 335–

337.

Serbin, S.P., Singh, A., Desai, A.R., Dubois, S.G., Jablonski, A.D., Kingdon, C.C., Kruger, E.L., Townsend,

P.A., 2015. Remotely estimating photosynthetic capacity, and its response to temperature, in

vegetation canopies using imaging spectroscopy. Remote Sens. Environ. 167, 78–87.

https://doi.org/10.1016/j.rse.2015.05.024

Serrano, L., Ustin, S.L., Roberts, D.A., Gamon, J.A., Peñuelas, J., 2000. Deriving water content of

chaparral vegetation from AVIRIS data. Remote Sens. Environ. https://doi.org/10.1016/S0034-

4257(00)00147-4

Shigo, A.L., 1984. Compartmentalization: A Conceptual Framework for Understanding How Trees Grow

and Defend Themselves. Annu. Rev. Phytopathol. 22, 189–214.

932 https://doi.org/10.1146/annurev.py.22.090184.001201

933 Sims, D.A., Gamon, J.A., 2003. Estimation of vegetation water content and photosynthetic tissue area

934 from spectral reflectance: A comparison of indices based on liquid water and chlorophyll

935 absorption features. Remote Sens. Environ. https://doi.org/10.1016/S0034-4257(02)00151-7

936 Struckmeyer, B.E., Beckman, C.H., Kuntz, J.E., Riker, A.J., 1954. Plugging of vessels by tyloses and gums in

937 wilting oaks. Phytopathology 44, 148–153.

938 Sturrock, R.N., Frankel, S.J., Brown, A. V., Hennon, P.E., Kliejunas, J.T., Lewis, K.J., Worrall, J.J., Woods,

939 A.J., 2011. Climate change and forest diseases. Plant Pathol. 60, 133–149.

940 https://doi.org/10.1111/j.1365-3059.2010.02406.x

941 Sullivan, G.M., Feinn, R., 2012. Using Effect Size—or Why the P Value Is Not Enough. J. Grad. Med. Educ.

942 4, 279–282. https://doi.org/10.4300/JGME-D-12-00156.1

943 Tang, C., Uriarte, M., Jin, H., Morton, D., Zheng, T., 2021. Large-scale, image-based tree species mapping

944 in a tropical forest using artificial perceptual learning. Methods Ecol. Evol. 12, 608–618.

945 https://doi.org/10.1111/2041-210X.13549

946 Townsend, P.A., Helmers, D.P., Kingdon, C.C., McNeil, B.E., de Beurs, K.M., Eshleman, K.N., 2009.

947 Changes in the extent of surface mining and reclamation in the Central Appalachians detected

948 using a 1976–2006 Landsat time series. Remote Sens. Environ. 113, 62–72.

949 https://doi.org/10.1016/j.rse.2008.08.012

950 Townsend, P.A., Serbin, S.P., Kruger, E.L., Gamon, J.A., 2013. Disentangling the contribution of biological

951 and physical properties of leaves and canopies in imaging spectroscopy data. Proc. Natl. Acad. Sci.

952 110, E1074–E1074. https://doi.org/10.1073/pnas.1300952110

953 Ullah, S., Skidmore, A.K., Ramoelo, A., Groen, T.A., Naeem, M., Ali, A., 2014. Retrieval of leaf water

content spanning the visible to thermal infrared spectra. ISPRS J. Photogramm. Remote Sens. 93,

56–64. https://doi.org/10.1016/j.isprsjprs.2014.04.005

Ustin, S.L., Gitelson, A.A., Jacquemoud, S., Schaepman, M., Asner, G.P., Gamon, J.A., Zarco-Tejada, P.,

2009. Retrieval of foliar information about plant pigment systems from high resolution

spectroscopy. Remote Sens. Environ. 113, S67–S77. https://doi.org/10.1016/j.rse.2008.10.019

Vogelmann, J.E., Rock, B.N., Moss, D.M., 1993. Red edge spectral measurements from sugar maple

leaves. Int. J. Remote Sens. https://doi.org/10.1080/01431169308953986

Waller, L.P., Allen, W.J., Barratt, B.I.P., Condron, L.M., França, F.M., Hunt, J.E., Koele, N., Orwin, K.H.,

Steel, G.S., Tylianakis, J.M., Wakelin, S.A., Dickie, I.A., 2020. Biotic interactions drive ecosystem

responses to exotic plant invaders. Science (80-.). 368, 967–972.

https://doi.org/10.1126/science.aba2225

Wang, R., Gamon, J.A., Moore, R., Zygielbaum, A.I., Arkebauer, T.J., Perk, R., Leavitt, B., Cogliati, S.,

Wardlow, B., Qi, Y., 2021. Errors associated with atmospheric correction methods for airborne

imaging spectroscopy: Implications for vegetation indices and plant traits. Remote Sens. Environ.

265, 112663. https://doi.org/10.1016/j.rse.2021.112663

Williams, L.J., Cavender-Bares, J., Townsend, P.A., Couture, J.J., Wang, Z., Stefanski, A., Messier, C.,

Reich, P.B., 2020. Remote spectral detection of biodiversity effects on forest biomass. Nat. Ecol.

Evol. 5, 46–54. https://doi.org/10.1038/s41559-020-01329-4

Wilson, J.S., Lindsey, G.H., 2005. Socioeconomic correlates and environmental impacts of urban

development in a central Indiana landscape. J. Urban Plan. Dev.

https://doi.org/10.1061/(ASCE)0733-9488(2005)131:3(159)

Wolter, P.T., Mladenoff, D.J., Host, G.E., Crow, T.R., 1995. Improved forest classification in the northern

Lake States using multi-temporal Landsat imagery. Photogramm. Eng. Remote Sensing 61, 1129–

1143.

Yadeta, K.A., Thomma, Bart, P., 2013. The xylem as battleground for plant hosts and vascular wilt

pathogens. Front. Plant Sci. 4, 1–12. https://doi.org/10.3389/fpls.2013.00097

Zarco-Tejada, P.J., Camino, C., Beck, P.S.A., Calderon, R., Hornero, A., Hernández-Clemente, R.,

Kattenborn, T., Montes-Borrego, M., Susca, L., Morelli, M., Gonzalez-Dugo, V., North, P.R.J., Landa,

B.B., Boscia, D., Saponari, M., Navas-Cortes, J.A., 2018. Previsual symptoms of Xylella fastidiosa

infection revealed in spectral plant-trait alterations. Nat. Plants 4, 432–439.

https://doi.org/10.1038/s41477-018-0189-7

Zarco-Tejada, P.J., González-Dugo, M.V., Fereres, E., 2016. Seasonal stability of chlorophyll fluorescence

quantified from airborne hyperspectral imagery as an indicator of net photosynthesis in the

context of precision agriculture. Remote Sens. Environ. 179, 89–103.

https://doi.org/10.1016/j.rse.2016.03.024

Zarco-Tejada, P.J., Miller, J.R., Mohammed, G.H., Noland, T.L., Sampson, P.H., 2002. Vegetation Stress

Detection through Chlorophyll a + b Estimation and Fluorescence Effects on Hyperspectral Imagery

. J. Environ. Qual. https://doi.org/10.2134/jeq2002.1433