Distributed networks for auditory memory differentially contribute to recall precision

Sung-Joo Lim[1,2,3,4]*, Christiane Thiel[5], Bernhard Sehm[2], Lorenz Deserno[2], Jöran Lepsien[2], and Jonas Obleser[1,2, 6]*

Affiliations:

[1] Department of Psychology, University of Lübeck, 23562, Lübeck, Germany

[2] Max Planck Institute for Human Cognitive and Brain Sciences, 04103, Leipzig, Germany

[3] Department of Psychology, Binghamton University, State University of New York, Vestal, NY, USA

[4] Department of Speech, Language, and Hearing Sciences, Boston University, Boston, MA, USA

[5] Department of Psychology, Carl von Ossietzky University of Oldenburg, 26129, Oldenburg, Germany

[6] Center of Brain, Behavior, and Metabolism, University of Lübeck, 23562, Lübeck, Germany

*Correspondence:

Sung-Joo Lim (sungjoo@binghamton.edu)

Jonas Obleser (jonas.obleser@uni-luebeck.de)

Running title: Distributed auditory memory representations

Abstract

Re-directing attention to objects in working memory can enhance their representational

fidelity. However, how this attentional enhancement of memory representations is

implemented across distinct, sensory and cognitive-control brain network is unspecified. The

present fMRI experiment leverages psychophysical modelling and multivariate auditory-

pattern decoding as behavioral and neural proxies of mnemonic fidelity. Listeners performed

an auditory syllable pitch-discrimination task and received retro-active cues to selectively

attend to a to-be-probed syllable in memory. Accompanied by increased neural activation in

fronto-parietal and cingulo-opercular networks, valid retro-cues yielded faster and more

precise responses in recalling acoustic detail of memorized syllables. Information about the

cued auditory object was decodable from hemodynamic response patterns in superior

temporal sulcus (STS), fronto-parietal, and sensorimotor regions. However, among these

regions retaining auditory memory objects, neural fidelity in the left STS and its enhancement

through attention-to-memory best predicted individuals' gain in auditory memory recall

precision. Our results demonstrate functionally discrete brain regions differentially contribute

to maintaining and attentionally enhancing memory representations.

Introduction

Working memory is a short-term mental storage that maintains perceptual information even in

the absence of sensory input. A hallmark of working memory is in its limited capacity—that is,

perceptual information held in memory is inherently noisy (Cowan, 2001; Bays and Husain,

2008; Luck and Vogel, 2013; Ma et al., 2014).

Selective attention plays a crucial role in overcoming the bottleneck of this limited

capacity. Directing attention to the relevant information in memory can reduce inherent noise

and enhance representations of the attended memory object—a process referred to as

retrospective attention (Griffin and Nobre, 2003; Postle, 2006; Gazzaley and Nobre, 2012;

Myers et al., 2017). Thus, even without sensory information, selective attention interacts with

working memory and facilitates the maintenance of attended memory objects.

Research on the neural bases of working memory and selective attention implicates

that multiple functional brain networks support active maintenance of the attended memory

objects. Neuroimaging research across various modalities suggests that both domain-general

networks related to attention (i.e., fronto-parietal network) and cognitive control (i.e., cingulo-

opercular network), and the sensory-specific neural regions are involved in maintaining items

held in working memory (Goldman-Rakic, 1995; Harrison and Tong, 2009; Higo et al., 2011;

Ester et al., 2015). For example, auditory and verbal working memory research has

demonstrated contributions of the domain-general networks, language-related brain regions

(e.g., the inferior frontal gyrus and inferior parietal areas), and premotor cortex (Baddeley,

2003; Hickok and Poeppel, 2007; Koelsch et al., 2009; Fedorenko et al., 2011), as well as the

bilateral auditory cortices for retaining stimulus-specific features of auditory information in

memory (Linke et al., 2011; Linke and Cusack, 2015; Kumar et al., 2016).

Collectively, the converging evidence suggests that neural traces of memory

representations under focused attention are maintained across distributed brain networks.

However, little is known about whether and how the multiple networks support the benefits

from attention re-directed to working memory. Can object-based retrospective attention

highlight the representational precision of the attended object already held in memory? If so,

how do these large-scale brain networks support attention leading to neural enhancement,

and are the memory representations maintained in these brain networks related to the

precision enhancement from attention? These questions are important in elucidating the

neural mechanisms of retrospective attention, which are still under debate (Souza et al., 2014;

Souza and Oberauer, 2016).

Human speech processing presents an excellent case for examining how attention

affects auditory working memory. Speech signals are inherently variable and they evolve

rapidly on the order of tenths of a second; also, the signals are often intermixed with interfering

sound sources, which can result in imprecise representations in memory. Thus, selective

attention plays a crucial role in forming and maintaining stable auditory objects in memory

(Fritz et al., 2007; Shinn-Cunningham, 2008; Wilsch and Obleser, 2016). The traditional

working memory framework proposes that only the abstract linguistic-level of information (e.g.,

syllables or words), but not low-level acoustic features, are maintained in memory for speech

(i.e., the phonological loop; Baddeley, 1992; 2003). Yet, little is known about whether and how

retrospective attention interacts with these internal representations held in working memory

for speech and the neural implementations associated with this process.

Prior work in working memory for speech has demonstrated that retrospective attention

facilitates working memory performance by enhancing representational precision of the

attended versus unattended objects held in memory, rather than removing the unattended

object from memory (Lim et al., 2015; 2018). Importantly, such precision benefit from

retrospective attention has been reflected in an attention-induced increase in neural activities

measured via EEG (i.e., enhanced sustained negativity and neural alpha (~10 Hz) oscillatory

power), indicating that greater cognitive resource is allocated to the attended memory object.

Thus, this prior work supports the notion that retrospective auditory attention enhances the

precision of the memory item, rather than reducing cognitive load by removing the unattended

items from memory (Lim et al., 2015). However, the limited spatial resolution of EEG cannot

delineate how distinct functional networks of the brain—cognitive control vs. auditory

processing regions—contribute to precision benefit from retrospective attention.

To investigate the neural underpinning of auditory retrospective attentional benefit, we

conducted an fMRI experiment, in which participants performed a previously established

auditory working memory task—a delayed recall of syllable pitch discrimination task (Lim et

al., 2015; Alavash et al., 2018; Lim et al., 2018). In this task, listeners encoded two speech

syllable sounds into memory and then received a retroactive cue to direct attention to one of

the syllables (i.e., auditory objects) prior to recalling lower-level feature (pitch) of the object.

Here, we specifically investigated (*i*) whether selective attention to auditory memory enhances

representational fidelity by using a psychophysical modeling approach (Bays and Husain,

2008; Zhang and Luck, 2008; Murray et al., 2013); (*ii*) how retrospective attention modulates

the neural activities in domain-general and modality-specific brain regions by using univariate

fMRI analysis; and (*iii*) whether the speech memory objects were retained in aforementioned

brain regions, and which of these brain regions contribute to the attentional enhancement in

mnemonic fidelity using a multivariate classifier approach on fMRI data.

117   We hypothesized that if retrospective attention facilitates working memory
118 performance by enhancing mnemonic fidelity of the attended object, we would observe higher
119 precision in recalling the acoustic feature of the syllable object held in memory, and attention-
120 related increase in neural activations in the domain-general brain regions, reflecting the higher
121 attentional demands to maintain precise memory representations. In addition, we expect that
122 the auditory memory object representations distributed across the brain—the domain-general
123 cognitive control and sensory-specific brain regions—would contribute to the attentional
124 enhancement in mnemonic fidelity, but potentially in varying degrees given that functionally
125 discrete brain regions may retain either more abstracted or sensory-specific information in
126 working memory (Lee et al., 2013; Ester et al., 2015; Christophel et al., 2017). However, if
127 retrospective attention reduces working memory load by removing unattended items from
128 memory (Kuo et al., 2012; Souza et al., 2014), we expect to observe an attention-related
129 decrease in neural activations in the domain-general networks (i.e., fronto-parietal and
130 cingulo-opercular networks), and no direct relationship between the neural representations of
131 the auditory memory object to the mnemonic precision in recalling the object.

132   Using univariate analysis on the fMRI data, we demonstrate that selective attention to
133 auditory memory leads to increased neural activity in domain-general cognitive control brain
134 networks, but not in the auditory-sensitive superior temporal cortex. Using a multivariate
135 analysis approach, we further show that while the neural representation of the attended
136 memory object is distributed across the brain, but these regions differentially contribute to the
137 attentional gain in improving representational fidelity. Beyond the domain-general cognitive
138 control regions commonly implicated in working memory, we found that the neural fidelity and
139 its enhancement through attention-to-memory in the speech-sensitive brain region in the left
140 superior temporal sulcus are also shown to best predict the individual gain in representational
141 precision of auditory memory objects.

A Syllable pitch-discimination task

B Behavioral results

C Psychophysical modeling results

* *p* < 0.05
*** *p* < 0.0001

Fig. 1. Illustration of the syllable pitch-discrimination task and the summary of task performance and psychophysical modeling results for validly cued (V) and neutrally cued (N) trials. (**A**) Trial structure of the task. On each trial, participants encoded two syllables (each syllable 0.2 s in duration) presented in a random order separated by a 1- or 2-s inter-stimulus interval (ISI). After the first delay period (3–5 s), a visual retro-cue appeared on the screen for 1 s, which was followed by a stimulus-free retention period (with duration jittered from 5–7 s). After the retention period, participants heard one of the encoded syllables again as a probe, the pitch of which was slightly manipulated. Participants judged whether the pitch of the probe syllable was higher or lower than the corresponding syllable heard during encoding (i.e., beginning of the trial). Participants were given a maximum of 4 s to provide their response. On every trial, participants received either a valid or neutral retro-cue, which respectively informed or did not inform the identity of the to-be-probed syllable category. (**B**) The behavioral performance results as a function of retro-cues. Left, response time of the correctly responded trials. Right, response accuracy. Bar plot shows the average performance measure in the valid and neutral cue conditions. Error bars indicate ± 1 standard error of the mean (SEM). Scatter dots represent the corresponding behavioral measure with respect to the cue conditions of individual participants. The 45-degree lines represent the performances in the two retro-cue conditions are identical. The histograms represent the distributions of the cue-related behavioral difference across N=22 participants. (**C**) The parameter estimates of psychophysical modeling results. Left, an exemplary psychophysical modeling fit of a single participant's response patterns in the valid and neutral retro-cue conditions. Each dot indicates the proportion responses "high" as a function of syllable pitch change at probe, with respect to the matching reference syllable presented during the encoding phase. The lines denote the logistic model fits. Right, individual's perceptual precision estimates (ln slope *k*) with respect to the cue conditions. The illustration scheme is identical to panel B.

Results

Our study aims to understand neural underpinning of working memory representations with
respect to whether and how attention enhances mnemonic fidelity when it is redirected to
objects held in memory. To this end, we collected functional MRI while participants performed
an auditory working memory task—a syllable-pitch discrimination task with retroactive cues
(see Materials and Methods; Fig. 1A; Lim et al., 2015; Alavash et al., 2018; Lim et al., 2018).
On each trial, participants heard two syllable sounds (i.e., /da/ and /ge/) in a random order,
and maintained the syllables in memory during a 9–13 s delay. After the delay, participant
heard an auditory probe and had to judge whether the pitch of the probe syllable was higher
or lower than the corresponding syllable heard in the beginning of the trial. Critically, to direct
participants' attention to memory items selectively vs. non-selectively, one of the two types of
visual retro-cues was presented during the memory delay period. A "valid" retro-cue was
presented as a written syllable (e.g., "da") to provide information about the upcoming probe
syllable; these cues directed participants' attention to one of the syllables maintained in
memory prior to hearing the probe. A "neutral" cue (written as "xx") did not provide any useful
information about the upcoming probe; thus, it required participants to maintain both syllable
sounds in memory until hearing a probe.

The task required participants to detect a small change in the pitch (F0) of the auditory
syllables (i.e., changes of ± 0.125 and ± 0.75 semitones), and the syllable pitch varied
throughout the experiment. Thus, listeners must remember specific sound instance of the
syllable to succeed in the task. To obtain behavioral and neural proxies of memory
representational fidelity at which participants maintained and recalled the syllable sounds in
memory, we adopted psychophysical modelling and multivariate pattern decoding approaches.

Valid retro-cues facilitate syllable-pitch recall performance

We first examined the effect of retro-cues (valid vs. neutral) on behavioral performances. A
linear mixed effects model on participants' log-transformed response times (RTs) of the
correct trials revealed a significant main effect of retro-cues ($M_{Valid–Neutral} = -151.03$ ms, $\chi^2(1) =$
92.26, $p \ll 0.0001$). As shown in Fig. 1B (left), participants were significantly faster in correctly
judging the pitch of syllables when they received valid retro-cues compared to the neutral cues.
On the contrary, we did not observe robust benefit from valid retro-cues on response accuracy.
Although we observed 13/22 participants were more accurate in recalling syllable pitch with
valid than neutral retro-cues, a logistic mixed-effects model on listeners' response accuracy
did not reveal any significant effect of retro-cues ($\chi^2(1) = 1.73$, $p = 0.19$; Fig. 1B, right).

We further examined the effect of retro-cues on listeners' perceptual precision and
response bias, estimated by the slope (k) and inflection/mid-point (m) parameters of logistic
function fit from psychophysical modeling (see Materials and Methods; Fig. 1C, left). A linear
mixed-effects model on the log-transformed perceptual precision estimate (ln slope k)
revealed a significant effect of retro-cue condition ($M_{Valid-Neutral} = 0.25$, $\chi^2(1) = 4.56$, $p = 0.033$).
Consistent with the prior findings (Lim et al., 2015; 2018), valid retro-cues led to more precise
recall of syllable pitch compared to the neutral retro-cue trials (Fig. 1C, right). For the response
bias estimate (m), a linear mixed-effects model revealed no significant effect of retro-cue
condition ($\chi^2(1) = 1.22$, $p = 0.27$); moreover, neither of the cue conditions did exhibit significant
bias (one sample t-test against 0, $ts \leq 1.11$, $ps \geq 0.28$).

It is of note that for any of these measures, we did not find any significant main or
interaction effects related to the factor of no interest—that is, the order of the two separate
scanning sessions taking place for each participant (all χ^2s < 0.36, all ps > 0.55). In addition,
we performed the same analyses with an additional factor to test whether the probed syllable
category had any impact on these measures; except for the response accuracy exhibiting a
slightly higher accuracy for /da/ than /ge/ ($\chi^2(1) = 5.03$, $p = 0.025$; M ± SD = 3.91 ± 6.90%),
we did not find any significant effects related to the probed syllable category in any other
measures (all ps > 0.14).

Valid and neutral retro-cues differentially modulate domain-general networks during
memory maintenance

To identify brain regions that were modulated by retrospective auditory attention, we
compared the brain activity of valid vs. neutral retro-cue trials during the memory retention
period following visual retro-cues. As shown in Fig. 2A, we found significantly greater
recruitment of broad regions across the whole brain when participants received valid than
neutral retro-cues (Table 1). While the effect was more prevalent in the left hemisphere, these
regions include the bilateral inferior frontal gyri extending into the dorsolateral prefrontal cortex,
and the bilateral inferior parietal regions, the left superior parietal lobule, and bilateral
supramarginal gyri. In addition, we found increased activity for the valid than neutral retro-
cues trials in the supplementary motor area (SMA) extending into the bilateral superior and
medial frontal gyri, and cingulate regions, the bilateral insula as well as the putamen. Similar
patterns of the effect were also observed in the bilateral precuneus/superior parietal lobule
(BA 7). Furthermore, the greater activations in the valid vs. neutral retro-cues was
accompanied by lower activations in the brain regions, typically identified as default-mode
network. The whole-brain analysis revealed significantly less activations for the valid than

neutral retro-cue conditions in the medial prefrontal gyrus (BA 10), the anterior cingulate (BA

32) and posterior cingulate cortex (BA 31), as well as bilateral angular gyrus regions (Fig. 2A).

In addition to the whole-brain analysis, we also conducted a ROI-based analysis in

order to examine the effect of valid vs. neutral retro-cues on the independently defined brain

networks established in previous literature (Dosenbach et al., 2007; Sadaghiani and

D'Esposito, 2014). The results from the ROI-based analysis were consistent with the result

patterns revealed in the whole-brain analysis. While both retro-cue conditions recruited the

task-relevant attentional networks (i.e., fronto-parietal and cingulo-opercular ROIs; Valid: all t

> 5.32, all p < 0.0001; Neutral: all t > 2.77, all p < 0.011; Fig. 2C), the corresponding linear

mixed-effects model revealed that both of these networks exhibited a significantly higher

activation in the valid than neutral retro-cues during the follow-up memory retention period

(fronto-parietal ROI: $\chi^2(1) = 25.22$, $p < 0.0001$, $\eta_p^2 = 0.56$; cingulo-opercular ROI: $\chi^2(1) = 34.66$,

$p < 0.0001$, $\eta_p^2 = 0.63$). As expected, the BOLD response in the default mode network was

lower than baseline during the memory retention phase of the trial for both cue conditions

(one-sampled t-tests against 0: Valid: $t(21) = -4.47$, $p = 0.00021$; Neutral: $t(21) = -2.41$, $p =$

0.025), but valid retro-cues led to a significantly lower activations of the default mode network

compared to the neutral retro-cue condition ($\chi^2(1) = 22.32$, $p \ll 0.0001$, $\eta_p^2 = 0.53$).

While the broad regions along the auditory cortex (i.e., the bilateral temporal cortex)

were significantly recruited when listeners were encoding the auditory speech syllables on trial

onset (Fig. S1), these regions were not recruited during the memory retention period. While

the bilateral auditory cortical ROIs were maximally active when listeners encode the auditory

syllables, activations in these regions were significant lower than baseline when listeners were

maintaining syllable information during the memory retention period (Fig. 2B–C; one-sample

t-test against 0; Valid: M = −0.094, $t(21) = -2.25$, $p = 0.036$, Neutral: M = −0.099, $t(21) = -2.96$,

$p = 0.0074$). Furthermore, these auditory cortical regions did not exhibit significantly different

BOLD activation in the valid vs. neutral retro-cue trials during memory retention (Fig. 2B–C;

$\chi^2(1) = 0.056$, $p = 0.82$). Likewise, the whole-brain analysis did not reveal any significant

activation differences in the auditory cortex based on the cue conditions, except in the right

middle temporal gyrus (BA 21) showing a relatively greater activation in the valid than neutral

retro-cue trials; nevertheless, this region was not strongly active during the memory retention

period (one-sample t-test against 0; Valid: M = −0.049, $t(21) = 0.78$ $p = 0.45$, Neutral: M = −

0.053, $t(21) = 0.91$, $p = 0.37$).

We further examined whether the retrocue-related modulations in the BOLD activation

in the brain networks, including the fronto-parietal, cingulo-opercular, default-mode, and the

auditory networks defined by ROIs, were related to the retrocue-related enhancement of the

274 perceptual precision measure (i.e., Δ ln slope k). The step-wise model comparison approach

275 starting from the maximal model did not reveal any significant effect of BOLD modulations on

276 the cue-related perceptual precision modulations; the simplest and the best model, which was

277 revealed by this analysis, included a marginal effect of the cue-related BOLD modulation of

278 the default-mode network ROI on the recall precision enhancement ($\beta = 0.39$, $t = 1.91$, $p =$

279 0.071) explaining 11.14% of the variance of the model (i.e., adjusted R^2). The leave-one-

280 subject-out cross-validation of this final model exhibited that the average model predicted

281 outcomes did not exhibit significant relationship to the observed outcome of cue-related

282 precision modulation (Pearson's correlation $r = 0.18$, $p = 0.42$; average adjusted R^2s from 22

283 cross-validations = 0.11).

Fig. 2. Result of the univariate analysis of Valid vs. Neutral retro-cues on modulating the brain activation during the post-cue memory retention phase. (**A**) Brain regions exhibiting significant effects of retro-cues (Valid, V – Neutral, N) (FDR corrected at $p < 0.05$). ROI seeds of the four, independently-defined networks are illustrated with the spheres of the corresponding colors. L, left; R, right. (**B**) BOLD activation time courses of the Valid vs. Neutral retro-cue conditions from the cue onset. The BOLD time courses were extracted from the exemplary regions (fronto-parietal and auditory ROIs). The grey boxes denote the time windows of the BOLD activity during the cue and the following memory retention phase (4–8s in FIR; see Methods and Materials). (C) Cue-related modulation of the BOLD activation in each network ROI. Bar graphs depict the average BOLD activations (i.e., beta coefficient) during the time points (4–8 s) of the Valid and Neutral retro-cue conditions. Scatter plots represent the average BOLD activations of the corresponding ROIs with respect to the retro-cue conditions of individual participants. The 45-degree lines indicate identical magnitude of activation of the two cue conditions. The histograms depict the distributions of the cue-related BOLD activity differences across N=22 participants. *** indicates $p < 0.0001$ (FDR corrected). Error bars in panels B and C indicate ± 1 SEM.

304 **Table 1. Brain regions exhibiting a significant contrast of Valid vs. Neutral retro-cue**
305 **trials during the cue and following memory retention period**

Brain region	MNI peak coordinate			Z-score	Voxels
	x	y	z		
Valid > Neutral					
L Pre-SMA	−6	18	54	7.93	2639
L Inferior frontal gyrus (BA 44)	−48	8	26	7.72	
L Dorsolateral frontal gyrus (BA 46)	−40	−32	26	7.56	
R Dorsolateral frontal gyrus (BA 46)	38	42	32	5.50	
R Middle FG (BA 6)	44	2	54	5.44	
L Insula/Inferior frontal gyrus (BA 24)	−34	24	6	6.79	
R Insula /Inferior frontal gyrus	36	24	2	7.17	689
L Putamen	−18	6	8	5.76	
R Putamen	20	8	6	5.16	54
L Inferior parietal lobule (BA 40)	−36	52	50	7.40	808
L Superior parietal lobule	−32	−60	54		
L Supramarginal gyrus	−54	−44	34	5.03	
R Inferior parietal lobule	36	−46	48	5.16	254
R Supramarginal gyrus	54	−42	30	4.67	
R Precuneus / Superior parietal lobule	8	−66	56	5.93	96
L Cerebellum	−34	−66	−24	5.02	103
R Cerebellum	38	−60	−28	5.89	202
R Cerebellum (semi lunar lobule)	24	−70	−48	4.36	34
R Middle temporal gyrus	54	−34	−4	4.62	42
Neutral > Valid					
R Medial frontal gyrus (BA 10)	12	66	8	−6.27	1118
L Anterior cingulate (BA 32)	−4	26	−10	−6.26	
L Posterior cingulate gyrus (BA 31) / Precuneus	−6	−46	32	−5.54	272
L Intraparietal lobule / Angular gyrus	−42	−72	38	−4.51	49
R Angular gyrus	54	−66	36	−4.17	14

306 Note: L, left; R, right; BA, Brodmann area

Valid retro-cues enhance neural representations of syllable category in memory

Using the multivariate classification analysis, we investigated whether the speech syllable object retained in memory can be decoded from the patterns of neural activation during the stimulus-free memory retention period upon the presentation of retro-cues. We trained the classifier to discriminate auditorily presented syllables (i.e., /da/ vs. /ge/) when participants heard them as a probe (i.e., one syllable presentation) only in the neutral cue trials in order to prevent confounds from listeners' expectation about the upcoming probe syllable informed by valid retro-cues. We then tested generalization of the trained classifier whether it could decode auditory syllable object from patterns from neural activation activity patterns during the cue–retention phase (i.e., in the absence of auditory presentation) of the task.

The multivariate analysis revealed the brain regions, in which attended syllable category objects can be decoded from neural activities during the cue–retention phase when listeners received valid retro-cues (Fig. 3; Table 2; Table S1). Specifically, we found that neural activity patterns in the left inferior frontal gyrus/dorsolateral prefrontal cortex, an anterior portion of the left superior temporal sulcus (STS), and right precentral gyrus adjoining the lateral Heschl's gyrus exhibited decoding accuracy significantly above chance when a valid retro-cue was presented. In addition, activity patterns in the right precentral region and the left parietal cortical region can accurately distinguish syllable category representations during the cue–memory retention phase. Furthermore, because retro-cues were presented visually on the screen prior to the memory retention period, we found that brain regions along the visual processing pathway, such as the left inferior temporal/fusiform gyrus region exhibited above-chance decoding accuracy for syllable category information with valid retro-cues, as expected.

However, the same analysis did not reveal any significant brain clusters exhibiting above-chance decoding accuracy for syllable objects under neutral retro-cue trials. Furthermore, as expected, among the brain regions that reliably retained syllable category information (Fig. 3), none of the areas did exhibit better decoding accuracies in the neutral than valid retro-cue conditions (Δ decoding accuracies Valid – Neutral = [2.67 – 5.90%]). Note that when we performed a parametric test of the multivariate analysis against the chance level decoding (i.e., 50% against 1-tailed test), we found a qualitatively similar finding as the results derived from the non-parametric permutation-based approach (Fig. 3).

340
341
Fig. 3. Multivariate classification results on decoding attended syllable object held in
memory. The classifier was trained to decode syllable categories based on the auditorily
presented probe sounds in the neutral retro-cue trials; and the classifier was tested to decode the
neural activity patterns during the memory retention phase following retro-cues. The brain regions
exhibit above-chance classifier accuracies on decoding the cued syllable category (i.e., indicated
by valid retro-cues; see Methods and Materials for further details) against non-parametric testing.
Each scatter plot depicts individuals' decoding accuracies of the valid vs. neutral retro-cue
conditions extracted from the corresponding brain regions that retain neural representations of
syllable category. The distribution of decoding accuracy of each cue condition is shown next to the
corresponding axis of the scatter plot. The histogram represents the distributions of the cue-related
decoding accuracy modulations (Valid – Neural) of N=22 participants. The 45-degree line would
denote identical decoding accuracies in the two cue conditions. Dashed lines are shown to indicate
hypothetical chance level (50%) decoding accuracy.

Table 2. MNI coordinates of significant clusters within the task-relevant brain regions exhibiting significantly above chance decoding accuracy for auditory syllable categories

Brain region	MNI peak coordinate		
	x	*y*	*z*
L Inferior frontal gyrus (BA 9)	−54	18	26
R Dorsolateral prefrontal cortex	48	14	26
R Anterior inferior frontal gyrus	50	42	8
R Superior frontal gyrus	32	−10	60
L Superior parietal lobule	−34	−60	62
L Insula	−40	−10	−6
L Superior temporal sulcus (BA 21)	−54	−6	−16
R Heschl's gyrus	60	0	0
R Postcentral gyrus	66	−22	44
R Precental gyrus	62	0	8
R Angular gyrus	60	−52	36
R Supramarginal gyrus	48	−42	30
L Inferior temporal gyrus (BA 20)	−52	−36	−22

Note: L, left; R, right; BA, Brodmann area

Based on the set of task-relevant neural regions that exhibited significantly accurate decoding of syllable categories during the memory retention phase (Fig. 3; Table 2), we further examined the relationship to the behavior. Specifically, we examined whether cue-related modulation of decoding accuracy in these brain regions was related to the representational precision benefit from valid retro-cues (i.e., ln slope *k* Valid vs. ln slope *k* Neutral). As done for the ROI-based univariate analysis (BOLD), we used a stepwise multiple linear regression analysis approach to start and deduce from the 13 task-relevant brain regions (Table 2; see Materials and Methods for details). Among these brain regions, this analysis revealed significant contributions of nine brain regions in predicting the degree of perceptual precision benefit from valid retro-cues (adjusted R^2 = 0.49; $F(10,11)$ = 3.03; p = 0.0421; Fig. 4A). The significant contributions of the brain regions are listed in Table 3. We found that the neural decoding accuracy modulations in the left STS, right postcentral gyrus, and right SMG exhibited significant relationships to the precision gain from valid retro-cues. We also found significant relationships to the precision benefit with the neural decoding modulations in the right dorsolateral prefrontal cortex, right precental gyrus, as well as left insula. However, we

did not find any significant contributions of the inferior frontal gyri, right superior temporal gyrus,
and the left inferior temporal gyrus.

To confirm the robustness of the model, we performed 22 iterations of a leave-one-
participant-out cross-validation of the final model to predict each subject's precision
modulation (ln k Valid vs. ln k Neutral) based on the remaining 21 participants' neural
modulations. We found that this model-predicted precision modulation had a significant
relationship to the observed outcome of the cue-related precision modulation (Fig. 4B;
Pearson's correlation $r = 0.47$, $p = 0.027$; average adjusted R^2 (M ± SD) = 0.49 ± 0.073).

Finally, based on the model estimates, we used a Wald statistic (Z_{Wald}) to directly
compare whether the neural decoding modulations of these brain regions differentially
contribute to the recall precision gain from valid retro-cues (Fig. 4A; Table 3). For a given pair
of model estimates, we computed a Z_{Wald} of their difference to test whether the two estimates
are significantly different from each other. We found that the behaviour-predicting contribution
of the neural decoding modulation of the left STS was significantly different from those of all
other brain regions (all $Z_{Wald} > 2.13$; all $p < 0.026$; FDR adjusted), including the right
dorsolateral prefrontal cortex ($Z_{Wald} = 4.04$; $p = 0.00017$), the left insula ($Z_{Wald} = 3.88$; $p = $
0.00023). We also found that the contributions of the right dorsolateral prefrontal cortex and
the left insula were significantly different from each other ($Z_{Wald} = 2.47$; $p = 0.012$). Table S2
reports Z_{Wald} of the differences between the model estimates.

Table 3. Multiple linear regression modeling result on predicting cue-related
modulation of perceptual precision from the neural decoding accuracy modulations
from retro-cues

Predictors	β	t	p
L Superior temporal sulcus (BA 21)	1.93	3.67	0.0037
R Postcentral gyrus	0.97	3.07	0.011
R Supramarginal gyrus	0.65	2.91	0.014
R Dorsolateral prefrontal cortex	−0.95	−3.85	0.0027
L Insula	−1.85	−3.83	0.0028
R Precental gyrus	−1.18	−3.49	0.0051
R Superior frontal gyrus	−0.98	−3.03	0.011
L Superior parietal lobule	−0.91	−3.03	0.011
R Angular gyrus	−0.75	−2.36	0.038
R Anterior inferior frontal gyrus	−0.39	−1.83	0.095

402 Note: L, left; R, right; BA, Brodmann area
403

404
405
Fig. 4. The relationship between the neural decoding for syllable category representations
in memory and behavioral modulations. (A) Estimated contributions of the brain regions' neural
decoding modulations are resulted from the step-wise multiple linear regression model, which
significantly predicted the extent of perceptual precision modulations (quantified as log-
transformed slope *k* from psychophysical modeling) from Valid vs. Neutral cue conditions; * $p <$
0.05, ** $p < 0.01$, *** $p < 0.005$. **(B)** Correlation between observed and the cross-validated model-
predicted cue-related modulations in perceptual precision (ln slope *k*) of the two cue conditions (Δ:
Valid – Neutral). Individual's model-predicted perceptual precision modulation obtained from cross-
validation (model fitted for nth participant from remaining *n*=21 participants). All data points were
z-scored prior to the model fitting.

416
417

Discussion

How can selective attention directed to an auditory object held in memory enhance the

representational precision of the object? How are multiple brain networks engaged in directing

attention to auditory memory object, and how do the memory representations held in these

networks contribute to enhancing the representational precision from attention? The present

study investigated the role of various neocortical brain regions, including the sensory-specific

and modality-general brain areas, in supporting the benefit from attention directed to working

memory objects. We here have utilized psychophysical modeling and a combination of

univariate and multivariate analyses on the fMRI data collected during an auditory retro-cueing

working memory task with speech syllables.

Attention directed to auditory memory objects enhances task performance and
representational precision

We found that retrospective attention facilitated auditory working memory performance.
Consistent with previous work on how retrospective attention benefits memory recall (Griffin
and Nobre, 2003; Lepsien et al., 2005; Lepsien and Nobre, 2007; Backer et al., 2015), valid
retro-cues led to faster responses than uninformative neutral cues. Furthermore, replicating
the prior findings with working memory for speech (Lim et al., 2015; 2018), our psychophysical
modeling results demonstrate that directing attention to a specific auditory object in memory
enhances the representational precision of the attended object even under adverse listening
conditions (i.e., auditory syllables presented under background noise). Overall, our findings
are in line with the notion that retrospective attention enhances memory representations
(Lepsien et al., 2011; Rerko and Oberauer, 2013).

Notably, we observed the facilitatory effects of retrospective attention even though
valid retro-cues provided information orthogonal to behavioral response planning. Valid cues
in the current study provided information only about the upcoming probe syllable category (i.e.,
object-based linguistic unit), whereas the task required listeners to make decisions on the
acoustic feature (i.e., pitch) of the cued object; nevertheless, we found that valid retro-cues
led to faster and more precise responses. Our findings are consistent with the object-based
account of attention (Duncan, 1984; Desimone and Duncan, 1995; Alain and Arnott, 2000;
Shinn-Cunningham, 2008). Sound features are integrated into a perceptual object, and
attention directed to one feature also enhances the processing of other features comprising
that object. Similarly, the performance benefits from valid retro-cues that we found suggest
that selective attention directed to a specific memory object enhances its representation
across varying levels of information, including the lower-level features comprising that object.
As we will argue below, our fMRI results indicate that retrospective auditory attention actively
engages neural resources to maintain the cued objects in memory, and the distributed neural
representations of the attended objects support the attentional gain in memory
representational fidelity.

Selective attention directed to memory objects engages domain-general cognitive
control networks

Our univariate results revealed that retrospective attention to auditory objects held in memory
recruited domain-general task control brain regions. Compared to neutral cue trials, valid retro-
cues led to significantly higher activations in broad cortical areas, including bilateral frontal
and parietal cortices, as well as SMA/ACC and bilateral insula. These brain regions largely

overlap with the well-known top-down networks: (1) the frontoparietal network—comprised of
prefrontal and parietal cortices—has been implicated in executive control, including selective
attention and working memory maintenance across various modalities (Rowe et al., 2000;
Owen et al., 2005; Bledowski et al., 2009); (2) the cingulo-opercular network, consisted of
SMA/ACC and bilateral anterior insula, is known to play a role in sustained maintenance of
task control and goals (Dosenbach et al., 2007; Seeley et al., 2007).

Prior neuroimaging work has observed increased activations in these domain-general
networks with greater cognitive demands. For instance, increased activation in the cingulo-
opercular network is observed when detecting unpredictable vs. predictable events
(Sadaghiani and D'Esposito, 2014), or during effortful comprehension of highly degraded
speech (Erb et al., 2013; Vaden et al., 2013). Similarly, the fronto-parietal network activity also
scales with task difficulty. For example, the fronto-parietal network activity increases when
more items are held in working memory and when there are increased attentional demands
(Todd and Marios, 2004; Xu and Chun, 2006; Emrich et al., 2013). Based on these pieces of
evidence, our results may seem counterintuitive—that is, while valid retro-cues consistently
facilitated task performance, we observed increased activations in these networks in the valid
vs. neutral cue conditions. This pattern of the result may suggest a potential neural mechanism
by which retrospective attention facilitates memory recall: increased activations in these
domain-general cognitive control networks suggest that selective attention directed to a
specific memory object does not reduce working memory load by removing unattended
objects; instead, maintaining the attended object in memory actively engages cognitive
resources.

The recent models of working memory, predominantly established in vision, posits that
the items held in memory can be maintained via the synaptic weights even in the absence of
persistent neural activity (Stokes, 2015; Wolff et al., 2017; Manohar et al., 2019). When an
item in memory is in the focus of attention (i.e., via retro-cueing), its representation becomes
re-activated, whereas the unattended memory items are still maintained in activity-silent
storage (Lewis-Peacock et al., 2012; Zokaei, 2014; Sprague et al., 2016). Because the current
study focused on understanding the benefits from selective vs. non-selective attention re-
directed to auditory working memory, we cannot provide evidence about the representations
of the unattended (hence un-cued) auditory memory object. Nevertheless, our univariate
results are consistent with the prediction of the recent working memory model: neural systems
maintain the items held in working memory regardless of the focus of attention, but selective
attention re-directed to an object changes its representation to be in an active state by

engaging the additional cognitive resources manifested in recruiting the general-purpose
functional brain areas, such as prefrontal cortex (Manohar et al., 2019).

Furthermore, it is worth noting that our findings and interpretations are also consistent
with prior EEG findings. Using the very similar experimental paradigm, our prior EEG study
demonstrated that valid retro-cues engaged neural resources to actively maintain the cued
item, and the greater resources allocated to the cued item enhanced its representational
precision of speech memory object (Lim et al., 2015). Overall, our results suggest that
retrospective attention engages additional neural processes (Griffin and Nobre, 2003; Lepsien
et al., 2005; Lepsien and Nobre, 2007), and this resource allocation is manifested in the
involvement of fronto-parietal and cingulo-opercular networks for selecting and prioritizing the
cued item in memory to be readily available for upcoming tasks (Wallis et al., 2015; Myers et
al., 2017). Although we did not find a direct relationship between the brain activations and the
enhancement of the representational precision of the cued syllable, the active engagement of
the neural resources allocated to the attended item in memory might be indirectly related to
enhancing its representational precision.

Distributed neural representations of auditory memory object differentially contribute
to representational precision benefit from attention
The multivariate analysis revealed that attended syllable held in memory was represented in
distributed areas of the brain. When listeners directed their attention to a particular speech
syllable guided by a valid retro-cue, we found that auditory representations of the cued syllable
category could be decoded from the activity patterns in both modality-general and sensory-
specific brain areas (Fig. 3). These areas include lateral prefrontal and parietal cortical areas
and insula region, part of the domain-general attention and cognitive control networks known
to support both visual and auditory working memory (Duncan and Owen, 2000; Postle et al.,
2000; Todd and Marios, 2004; Noyce et al., 2017; Uluç et al., 2018), as well as the
supramarginal gyrus (SMG), inferior frontal gyrus, and the sensorimotor cortices, classically
implicated in verbal working memory (Awh et al., 1996; Smith et al., 1998; Baddeley, 2003)
and phonological processing (Price et al., 1997; Hartwigsen et al., 2010; Qi et al., 2019).
Beyond these brain regions that have been commonly implicated in working memory, we also
found that speech-sensitive cortical area along the left superior temporal sulcus (STS)
maintained the cued syllable information during memory retention. Consistent with the
distributed account of working memory representations (Ester et al., 2015; Christophel et al.,
2017), we found that multiple brain regions, in the frontal, parietal, and sensory-specific areas
represent information maintained in memory.

Interestingly, our results further suggest that widely distributed memory
representations do not uniformly contribute to retrospective attentional benefit in enhancing
the representation of the cued item. The regression analysis in predicting individual
differences in the memory precision gain from valid retro-cues (Δ slope k) revealed that the
extent of individuals' neural fidelity enhancements in the various brain regions—including the
left STS, right SMG, the prefrontal cortex, and left insula—exhibited significant relationships
to the behavioral enhancement of representational precision in recalling the cued syllable in
memory. However, no such relationships were found with the bilateral IFG, and the left inferior
temporal cortex (Fig. 4). This pattern indicates that the distributed memory representations
across broad cortical regions work together in enhancing representational precision of the
attended object held in working memory; however, they do so by exerting differential
contributions.

One potential explanation of our findings is that while the multiple brain regions
redundantly represent the objects held in working memory, these regions may not necessarily
maintain the same level of representations of the objects. This explanation might be consistent
with a view that there is a division of labor between sensory-specific and domain-general
cortices in efficient maintenance of working memory (Spitzer et al., 2014; Sreenivasan et al.,
2014; Christophel et al., 2017). That is, our results may suggest that those frontal cortical
regions hold object-level representations of speech sounds (i.e., syllable category), whereas
the speech-sensitive area in the left STS as well as language-related brain regions, such as
right SMG, may retain both the object-level syllable category and the acoustic featural
information bound to the object.

This interpretation is in line with the functional specializations of these cortical regions.
Many prior works demonstrate that frontal areas are sensitive to between-category, but not
within-category differences (Freedman et al., 2001; Myers et al., 2009). In addition, the frontal
cortex might maintain memory representations in an abstract format that can be generalized
across sensory modalities (Spitzer et al., 2014; Uluç et al., 2018) and be flexibly transformed
based on the task-relevant goals (Lee et al., 2013; Stokes et al., 2013; Long and Kuhl, 2018).
On the contrary, as we will discuss below, the left STS has been prominently implicated in
speech sound processing, categorization, and learning (Formisano et al., 2008; Leech et al.,
2009; Obleser and Eisner, 2009; Mesgarani et al., 2014; Lim et al., 2019), which may be
functionally suitable for retaining acoustic details of speech sounds held in memory. Overall,
our results suggest that while both domain-general fronto-parietal networks and domain-
specific regions maintain the working memory contents (Ester et al., 2015; Bettencourt and
Xu, 2016), the representations held in the distinct regions might be differentially sensitive to

top-down task-related goals vs. bottom-up stimulus-specific features (Long and Kuhl, 2018),
thereby exerting differential influences in the attentional enhancement of representational
precision.

Note that among the brain regions retaining syllable memory objects (Fig. 3; Table 2),
none of the regions did exhibit higher decoding accuracy in the neutral vs. valid retro-cue
condition. While we considered this pattern as evidence of neural fidelity enhancement from
selective (valid) vs. non-selective (neutral) retrospective attention, one might question that
whether lower decoding accuracy in the neutral vs. valid cue condition might result from
retaining a mixture of robust representations of both syllable sounds in memory rather than
reflecting attentional enhancement of memory representations. If this possibility were to be
true, we expected to observe higher neural activations in the above brain regions in the neutral
compared to the valid condition to actively maintain strong memory representations. We tested
this possibility using the extent of BOLD responses as a proxy for active representations of
memory objects. The neutral condition did not lead to greater activation compared to the valid
condition, for example, in the left STS (e.g., mean BOLD difference in Valid – Neutral = 0.048).
Thus, we conclude that this alternative possibility—maintenance of more robust
representations of two syllables in the neutral compared to valid condition—is unlikely (Emrich
et al., 2013).

The role of the left STS in maintaining auditory memory representations

Apart from the cognitive-control brain regions typically implicated in verbal working memory,
we also found that the auditory/speech-sensitive brain regions along the superior temporal
cortex contribute to attentionally enhancing auditory memory representations. During memory
maintenance of auditory syllables, we neither observed significant engagement of the
auditory/speech-sensitive brain regions along the superior temporal cortex, nor found
differential recruitment of these regions during memory maintenance in the valid vs. neutral
cue condition. However, the MVPA result revealed that the left STS retained the cued syllable
item held in memory, which seems to be directly related to the extent of precision
enhancement gained from retrospective attention (i.e., precision slope k measure; Fig. 4).
Although there is conflicting evidence about the engagement of the superior temporal cortex
during auditory working memory (Gaab et al., 2003; Linke et al., 2011; Linke and Cusack,
2015; Kumar et al., 2016), our results are generally in line with the prior findings that sensory-

specific brain regions retain working memory contents, irrespective of its activations (Harrison
and Tong, 2009; Serences et al., 2009; Riggall and Postle, 2012; Emrich et al., 2013).

One difference to note is that while prior work found the primary auditory cortex to
maintain pure tones in memory (Linke et al., 2011; Kumar et al., 2016), we found that the left
STS maintained the attended auditory speech syllables. This discrepancy might lie in the
nature of the stimulus objects held in memory. Although listeners had to judge the pitch of
speech syllables, the current task cannot be performed simply by extracting pitch information
from syllables without maintaining syllable category representation in memory. And this task
demand might have induced the auditory memory representations to reside in the left STS.

Prior neuroimaging studies in speech perception demonstrate that the left superior
temporal regions functionally subserve speech category representation and processing
(Formisano et al., 2008; Obleser and Eisner, 2009; Lim et al., 2019). Especially, the anterior
STS/STG is shown to be sensitive to tonal speech categories and speech vowels that contain
talkers' voice and/or pitch (Bachorowski and Owren, 1999; Obleser et al., 2006; Feng et al.,
2018). Thus, accessing the auditory syllable mnemonic representations in the left STS might
allow the readout of the featural-level information (e.g., pitch) of the object, which can be
enhanced by attention directed to the syllable object in memory. The contribution of the
speech-sensitive brain areas seems consistent with emerging evidence that speech working
memory is not independent of sensory/auditory processing; rather, both auditory cortical
regions, responsible for speech perception, and sensorimotor cortical areas support speech
working memory (Jacquemot and Scott, 2006; Perrachione et al., 2017; Scott and Perrachione,
2019).

Potential limitations
At first glance, one might be concerned with a potential circularity in finding brain regions that
retained auditory syllable memory representations. We defined these regions based on the
neural activity patterns in brain areas retaining the cued object information provided by the
visual valid cues. These regions might be biased to exhibit higher decoding accuracies for
memory content in the valid than the neutral cue conditions. Nevertheless, the extent of
individuals' neural decoding accuracy modulations (i.e., valid vs. neutral) in predicting the
extent of behavioral (precision slope k) modulations cannot be solely driven by the decoding
accuracy in the valid cue condition. Furthermore, our main focus was on examining the relative

contributions of distributed brain regions' enhanced representational precision, which should

be relatively unaffected by the overall bias that favored the valid cue condition.

The second potential issue might be on the chosen retro-cues themselves. Because

valid retro-cues were written syllables, all activation and representation differences observed

between valid and neutral cues might at least in parts be by-products of reading or imagery

evoked by written text. To minimize this concern, our classification approach specifically aimed

to find the brain regions maintaining the auditory traces of speech syllable objects held in

memory. The classifier was trained to learn syllable sounds when they were clearly presented

as auditory probes, and tested to decode the neural activity patterns during memory

maintenance (i.e., no auditory input). Furthermore, the classifier was trained only on the

auditory probes of the neutral cue trials to further avoid potential bias or expectations of a

particular syllable informed by valid retro-cues. Of course, we cannot completely rule out the

possibility that reading written syllables on valid cues itself yields reliable neural decoding. For

instance, we found that the left inferior temporal gyrus, known to be sensitive to written texts

(Cohen et al., 2002), exhibited above-chance accuracy in decoding syllable category

maintained in memory. Nevertheless, listeners must maintain the auditory-form of the speech

syllables held in memory to succeed in the task. Thus, relying on the visually presented

syllable category information alone or the potential evoked imagery from the written text

cannot fully explain consistent facilitation in the task performance from valid retro-cues.

Supporting this possibility, the attentional enhancement of the neural representations in the

left inferior temporal gyrus (ITG), sensitive to reading texts, was not related to the extent of

the cue-related enhancement of representational precision (slope k; Fig. 4).

Lastly, as our main research focus was on examining the benefits of the object-based

auditory retrospective attention in highlighting the attended memory representations, and

because our current design was not optimal to neurally decode the exact pitch of the syllables,

we performed the MVPA analysis to decode the abstracted linguistic-level of the attended

speech sounds (i.e., syllable category information) instead of the low-level pitch (F0) of the

syllables. While it is interesting that our findings suggest that high-level object-based neural

representation of syllables can significantly contribute to the behavioral facilitation in recall

precision of low-level pitch information (Fig. 4), our MVPA results cannot fully reveal which of

the exact features (i.e., higher-level vs. lower-level) underlie enhancement in neural decoding

accuracy during memory maintenance. Given the fact that complex auditory objects (such as

speech sounds) are comprised of varying levels of information, future studies are necessary

to investigate whether and how different levels of auditory memory representation—low-level

acoustic features vs. high-level linguistic information—are maintained in auditory working memory, and can be enhanced by selective attention.

Conclusions

As demonstrated here, working memory representations across the brain, ranging from sensory-specific and domain-general cognitive-control regions differentially contribute to the retrospective attentional gain in the fidelity of memory representations. Here, we show that while re-directing attention to the relevant auditory memory objects mainly recruits domain-general cognitive-control networks, the attentional enhancement in the neural fidelity in the superior temporal sulcus and frontal cortical regions was related to the individual gain in recall precision of auditory objects from memory, with the superior temporal sulcus best predicting the attentional gain. Together with previous research on working memory, our findings provide evidence that discrete, functionally specialized brain regions collectively contribute to maintaining and attentionally enhancing working memory representations.

Materials and Methods

Participants

Twenty-two participants (mean age: 27.9 years; SD: 2.75 years; age range: 25–35; 12 females) took part in the experiment that has been a part of a previously published study (Alavash et al., 2018). Participants were recruited from the participant database of the Max Planck Institute for Human Cognitive and Brain Sciences (Leipzig, Germany). Two additional participants underwent the study, but were excluded from data analyses due to excessive head movements (>5 mm). All subjects reported normal hearing, and no history of neurological or psychiatric disorders. None of the participants were under medication. All participants except one were right handed. Based on our prior EEG and behavioral work adopting the similar auditory retro-cueing task (Lim et al., 2015, 2018), we determined that with 17 participants we would achieve at least 80% power to detect the within-subject effect of retro-cues even under simultaneous noise in the background.

All participants gave their informed written consent, and received financial compensation for their participation. The study procedure was in accordance with the Declaration of Helsinki. The local ethics committee (University of Leipzig) approved the study.

Stimuli

The stimulus material was identical to the ones used in previously published studies (*26-28*). In brief, we used two German syllables, /da/ and /ge/. Each syllable category consisted of six naturally spoken tokens, truncated from German words recorded by a trained female speaker. All sound tokens were digitized at 44.1 kHz, and 200-ms in duration. For each token, 3-ms and 30-ms linear ramps were applied at sound onset and offset, respectively.

For each syllable category, two tokens served as to-be-probed syllables in the syllable-pitch discrimination task. The pitch dimension (i.e., the fundamental frequency; F0) of each token was parametrically manipulated in four steps: ± 0.125 and ± 0.75 semitones from its F0; we chose these relatively small step sizes in order to closely examine whether selective attention to speech objects held in memory benefits precise recall of low-level (pitch) information of the cued object. The original token was presented during the syllable encoding phase, and the F0-manipulated tokens were presented as probes. The average F0's of the to-be-probed /da/ tokens were 162.8 Hz [min = 154.0, max = 171.7] and 175.5 Hz [min = 169.4, max = 183.5] for /ge/ tokens. To increase acoustic variability during the syllable encoding phase, we created additional 12 tokens for each syllable category based on the six utterances from the same speaker. These sounds served as unprobed tokens: they were presented during encoding, but were not used for discriminating pitch change at probe. On average, unprobed tokens of /da/ and /ge/ had F0's of 163.7 Hz [min= 152.8, max = 181.7] and 175.9 Hz [min = 163.7, max = 186.9], respectively. These unprobed sounds were used to ensure that participants were exposed to acoustically-variable sounds beyond the fixed set of to-be-probed tokens, but within the similar F0-range such that to-be-probed and unprobed sounds were indistinguishable during the encoding phase.

All stimuli were normalized to equivalent amplitude (root-mean-squared dB full scale; RMS dBFS). Praat version 5.3 was used to manipulate the F0 dimension.

Experimental Design

In the MRI scanner, participants performed a previously established auditory working memory task—a syllable-pitch discrimination task with retroactive cues (*26-28*). Figure 1A illustrates the task structure. On each trial, participants first heard two syllable tokens (i.e., /da/ and /ge/) in a random order sequence, separated by a 1- or 2-s inter-stimulus interval (ISI). After a delay period of 3–5 s, participants saw a visual retro-cue for 1 s displayed on the screen. With an additional delay of 5–7 s following the visual cue, participants heard a probe syllable. The probe was one of the two syllables heard during encoding with a slight manipulation in the pitch (F0). Within a 4-s time window, participants responded whether the pitch of the probe

syllable was higher or lower than the corresponding syllable (i.e., the same category) heard

during encoding in the beginning of the trial. Response feedback was provided on the screen

for 0.5 s. Trials were separated by a 3–7 s silence period with a fixation cross on the screen.

During the retention period on each trial, participants saw one of the two retro-cues:

valid or neutral. A valid retro-cue indicated the identity of the probe syllable category; a written

syllable (e.g., "da") was presented on the screen for 1 s in order to direct participants' attention

to one of the syllables held in memory prior to hearing the probe. On the neutral retro-cue

trials, participants were presented with "xx" on the screen. This cue did not provide any

information about the upcoming probe syllable. Thus, listeners must maintain both syllable

sounds in memory until hearing a probe.

The pitch variations of the two to-be-encoded syllables were drawn from the similar

F0-range. On average, pitch (F0) difference between the two syllables presented during

encoding was 4.13 Hz (cf. note that a wider pitch range (>200 Hz) of auditory pure tones were

used in previous auditory working memory studies; e.g., Link et al., 2011; Kumar et al., 2016),

and the F0 difference was not significantly different between the two retro-cue conditions (p =

0.998). Nevertheless, F0's of the syllables presented during the encoding phase varied

continuously across trials; thus, listeners must remember specific sound instance of the

syllable in order to perform the task.

Participants performed a total of eight runs of 16 trials. On each run, the valid and

neutral retro-cue trials appeared on an equal probability (i.e., 8 trials per cue condition); thus,

participants were unaware of trial type until seeing a visual retro-cue. For every two runs,

unique combinations of 2 syllable categories × 2 syllable positions during encoding being

probed × 2 retro-cue types × 4 probed pitch step sizes were presented.

Procedure

Prior to fMRI scanning, all participants completed a short practice session of the syllable pitch

discrimination task on a laptop in a separate behavioral testing room. The practice session

involved 18 trials (9 trials for each cue type), which only probed easily detectable syllable pitch

changes (i.e., ±1.25 semitones). This practice session was given to ensure that participants

understood the task.

In the scanner, participants used left and right index fingers to provide "high" and "low"

responses. The mapping between the responses and hands were counterbalanced across

participants. Experimental trials were controlled with Presentation software (Neurobehavioral

Systems). Auditory stimuli were delivered through MR-compatible headphones (MrConfon,

Magdeburg, Germany), and Music safe pro earplugs (Alpine Hearing Protection) provided an

additional attenuation (on average 20-dB across 125–8000 Hz range). To ensure that

participants could hear the auditory syllables during task in the scanner, we conducted a brief

hearing test; participants heard four randomly selected speech stimuli with an on-going

functional (T2*-sequence) scanner noise, and verbally reported the syllable categories of the

sounds.

After the eight scanning runs of the syllable-pitch discrimination task, participants went

through two additional scanning runs of an auditory habituation (~16 min), involving listening

to a sequence of four syllable tokens while watching a silent movie. However, excessive head

movements were observed in at least 7/22 participants (> 4.5 mm) during this task; therefore,

data from this task was not further reported.

Note that the data and results reported here have been acquired as part of a larger,

pharmacological fMRI study such that all participants went through two identical fMRI sessions,

separated by at least one week (EudraCT number 2015-002761-33). The study design was a

double-blind, counterbalanced repeated-measures of the intake of levodopa (L-dopa) and

placebo. Because our current research focus is on the neural basis of retrospective attention

to auditory working memory (rather than the effect of L-dopa), we report the results only from

the placebo session. Of significant note, in all our analyses, we controlled a potential confound

in the placebo session order differences across participants. The effect of L-dopa on auditory

working memory, corresponding functional neural organization, and the detailed procedure of

the pharmacological fMRI study has been described elsewhere and can be found in Alavash

et al. (*28*).

Statistical Analysis

Behavioral measures and psychophysical modeling analysis

Response time (RT) of the correct trials and accurate syllable pitch judgment (i.e., a binary

response accuracy measure) served as overall behavioral performance measures. To ensure

normality, RT data were log-transformed prior to analysis. These single-trial behavioral

measures were separately analyzed using a (generalized) linear mixed-effects model

framework (*lme4*, ver. 3.3) in R environment. The model included a fixed factor of retro-cue

(Valid vs. Neutral), which was the main variable of interest. The model also included the

session order as a nuisance factor. The random-effects terms included by-participant

intercepts. The significance of the effects was determined based on Wald χ^2 tests (at $p < 0.05$).

Beyond these two behavioral performance measures, we also obtained a more fine-

grained perceptual precision measure using a psychophysical modeling approach used in the

visual domain (Bays and Husain, 2008; Zhang and Luck, 2008; Murray et al., 2013). We

estimated individual participant's perceptual precision in judging syllable pitch change by fitting

each participant's response patterns along the varying levels of pitch changes occurred at

probe. Following the previous auditory working memory work (*26,27*), we used a nonlinear

least square curve fitting procedure (MATLAB *lsqcurvefit*) to fit individual's response patterns

with a logistic (sigmoid) function,

$$y = \frac{1}{1 + e^{-k(x-m)}}$$

where *x* indicates the amount of pitch changes occurred at probe relative to the syllable (4

step sizes), *y* indicates the proportion of "high" responses, *k* indicates the slope of the curve,

and *m* indicates the mid-point/inflection point of the function along the x-axis. The parameter

m estimates bias in each participant's response, and the slope *k* provides an estimate of the

perceptual precision in pitch judgment in the task: higher perceptual precision is indicated by

steeper slope (see Fig. 1C, left for a single subject exemplary model fit).[1] As done for analyses

of the behavioral measures, the psychophysical modeling estimates of perceptual precision

slope *k* and bias *m* were separately analyzed using a linear mixed-effects model. The linear

mixed-effects models included a fixed factor of retro-cue (Valid vs. Neutral) and the nuisance

factor of the session order. The random-effects term included by-participant intercepts. The

significance of the effects was determined based on Wald χ^2 tests (at $p < 0.05$).

Functional MRI acquisition and analysis

Imaging data were acquired on a 3-Tesla Siemens MAGNETOM Prisma[fit] scanner (TimTrio

Upgrade) at the Max Planck Institute for Human Cognitive and Brain Sciences. Whole-brain

functional images, sensitive to the blood-oxygen-level-dependent (BOLD) contrast, were

acquired with a 20-channel head/neck coil using a T2*-weighted echo-planar imaging (EPI)

sequence (repetition time (TR) = 2000 ms; echo time (TE) = 26 ms; flip angle (FA) = 90°).

Each volume had 40 oblique axial slices (ascending) parallel to the anterior commissure–

posterior commissure (AC–PC) line (acquisition matrix = 64 × 64; field of view (FOV) = 192

mm × 192 mm; voxel size = 3 × 3 × 3 mm; interslice gap = 0.3 mm). For each block of the

syllable-pitch discrimination task runs, a total of 181 volumes were acquired.

Structural images of (*n* = 15) participants were available from the Institute's brain

database acquired from previous MR studies. A magnetization prepared rapid gradient echo

(MP-RAGE) T1-weighted sequence was used to acquire the structural images (TR = 2300 ms;

TE = 2.01–2.98 ms; FA = 9°; 176 saggital slices with 1 × 1 × 1 mm resolution). For the

---

[1] Note that the slope *k* measure is analogous to the inverse of response variability transformed based on a cumulative distribution function.

participants (*n* = 7) whose structural images did not exist in the database, a high-resolution

T1-weighted structural image was acquired using a MP-RAGE sequence (TR = 2300 ms; TE

= 2.98 ms; FA = 9°; 176 saggital slices with 1 × 1 × 1 mm resolution) at the end of functional

scans.

Imaging data were preprocessed using the Analysis of Functional NeuroImages (AFNI;

ver. 01-2017) software (Cox, 1996). The first two volumes of each run were discarded to allow

the scanner to approach net magnetization equilibrium. Preprocessing steps included 1) slice-

time correction, 2) motion correction of all volumes based on the first volume of the syllable-

pitch discrimination task run using a rigid-body alignment, 3) spatial alignment of functional

data to each participant's structural anatomy, and 4) the signal for each voxel was scaled to a

global mean of 100. For univariate analysis, images were spatially smoothed with an 8-mm

full-width at half maximum (FWHM) Gaussian kernel. For multivariate classification analysis,

data were not smoothed. Spatial normalization of functional images to Montreal Neurological

Images (MNI–152) was performed prior to the group-level statistical analyses.

Univariate analysis. For each participant, a voxel-wise general linear model (GLM)

was constructed to estimate hemodynamic responses during different phases of the syllable-

pitch discrimination task across all scanning runs. The BOLD response to the trial onset (i.e.,

the syllable encoding phase) was modeled using a canonical gamma variate function (GAM

by AFNI) along with a pairwise amplitude-modulated regressor that accounted for the ISI

between presentation of the two syllables. The second task regressor captured the entire

hemodynamic response of the visual retro-cue and the following stimulus-free retention phase

up to auditory probe phase using a finite impulse response (FIR) function (TENT by AFNI) as

a change in BOLD activation across 0–20 s from the retro-cue onset. To account for the

varying durations of the retention phase (Fig. 1A), a parametric amplitude-modulated

regressor was entered as a trial-wise covariate. The third regressor captured the BOLD

response related to the participants' response and visual feedback, we used a canonical GAM.

All task regressors were separately modelled for the valid and neutral retro-cue trials in the

GLM. The baseline activity was modelled by trends from linear up to fourth-degree

polynomials in order to remove the slow scanner signal drift (i.e., high-pass filter with an

approximate cut-off of 240 s). The six additional continuous head motion parameters were

entered as regressors of no interest.

The primary purpose of the univariate analysis was to identify brain regions that were

differentially modulated by selective vs. non-selective attention directed to auditory working

memory objects informed by retro-cues. Thus, we conducted a group-level analysis to contrast

the neural activities during the cue and the following retention phase in the valid vs. neutral

retro-cue conditions. For each subject, the cue-retention phase neural activity was quantified
by averaging the β-parameter estimate of the FIR during 4–8 s from the cue onset, separately
for the valid and neutral retro-cue conditions (Fig. 2B). The group-level analysis was
conducted using a voxel-wise linear mixed-effects modeling approach (3dLME by AFNI). The
model involved one within-subjects fixed factor of interest (retro-cue: Valid vs. Neutral) with
subjects as a random factor, and regressor of no interest (i.e., placebo session order) was
centered, and entered as covariates in the model. Planned contrast was established to
compare BOLD activations during the cue–retention phase in the valid vs. neutral retro-cue
conditions. The group-level maps were corrected for multiple comparisons using false
discovery rate (FDR $p < 0.05$; Benjamini and Hochberg, 1995).

Region-of-interest (ROI)-based analysis. We conducted an additional ROI-based
analysis (*i*) to complement and confirm the whole-brain univariate analysis as to whether the
valid vs. neutral retro-cues modulate the brain activities in the well-known domain-general
cognitive networks; and, more importantly, (*ii*) to investigate whether the neural activity
modulations in these distinct networks are related to behavior (see below). To ensure
independence of defining the domain-general network ROIs and the retro-cue-related BOLD
activity modulations, we defined the ROIs based on the brain regions organized by intrinsic
connectivity networks established in previous literature (Dosenbach et al., 2007; 2008; Power
et al., 2011; Sadaghiani and D'Esposito, 2014). To this end, we placed spherical seeds (radius
6 mm) at the centers of each brain network coordinates defined in previous work (Fig. 2A); for
the fronto-parietal network, the seeds were placed over the bilateral dorsolateral prefrontal
cortex (MNI coordinates: [–44, 27, 33] and [46, 28, 31]) and the bilateral inferior parietal lobules
([-53 –50 39], and [54 –44 43]) as defined in Dosenbach et al. (34) and Power et al. (*94*). ROIs
representing the cingulo-opercular network were placed based on Dosenbach et al. (*34*) in the
bilateral insula/frontal operculum ([–36, 18, 2] and [38, 21, –1]), and the supplementary motor
area (SMA)/dorsal anterior cingulate cortex at [0, 15, 45]. Default mode network seeds were
placed at the posterior cingulate cortex [1, –51, 29] and medial prefrontal cortex [–1, 61, 22]
based on previous findings (Raichle et al., 2001). In order to observe neural activity modulated
by the retro-cues in the sensory processing (i.e., auditory) regions, we also placed spherical
seeds, centered over the peak activations during encoding of the two syllables at the onset of
trial in the current experiment; these were placed in the bilateral auditory regions (superior
temporal gyrus/sulcus) at [–58, –24, 12] and [62, –22, 2].

To test whether and how the retro-cues modulate brain activity in each of the four
network ROIs, we first extracted the average BOLD responses during the cue–memory
retention phase (i.e., average β-estimates during 4–8 s time window of the FIR; Fig. 2B) from

the voxels within the seeds representing each network. The average β estimate data of each

network were submitted to a linear mixed-effects model with a fixed factor of retro-cue (Valid

vs. Neutral), a nuisance regressor of the session order, and subject-wise random intercept.

We corrected for multiple comparisons using FDR ($p < 0.05$).

Multivariate analysis. Our main goal was to understand how object-based attention

directed to auditory working memory improves the representations held in memory. To this

end, we conducted a multi-voxel pattern analysis (MVPA) to examine (*i*) whether the attended

syllable object information held in memory can be decoded from patterns of neural activity

during the memory retention phase, and (*ii*) whether the decoding performance for the

attended auditory object differed between the valid vs. neutral retro-cue conditions. Here, we

performed a whole-brain searchlight analysis with a searchlight sphere of 10 mm radius (~123

voxels; Kriegeskorte et al., 2006; Haynes et al., 2007). For each retro-cue condition, we used

a linear support vector machine classifier, implemented in LIBSVM (Chang and Lin, 2011),

with neural activation estimates in voxels in a searchlight sphere as features.

Because two syllable sounds presented during encoding not only varied throughout

the experiment, but also were not so differentiated by the pitch dimension (F0 difference

varying between 0.64–7.86 Hz across trials), we trained the classifiers to decode syllable

category information (i.e., /da/ vs. /ge/) instead decoding the syllable pitch. To identify brain

regions that retain auditory-based form of the attended syllable category during memory

retention, we trained SVM classifiers to decode syllable categories from the brain activity

patterns when participants heard one of the two syllables at the end of the trial (i.e., the

auditory probe phase) so that neural activity patterns used for training was not contaminated

by other auditory events. We then tested whether the trained classifier could successfully

decode syllable category information from the brain activity patterns during the memory

maintenance phase (i.e., the retro-cue and the following stimulus-free retention period). Of

important note, it is possible that in valid retro-cue trials, neural activity evoked by auditory

probes might exhibit syllable category-specific bias informed by the valid cues. To avoid this

potential bias, classifier training was performed only on the auditory probe phase of the neutral

retro-cue trials, in which participants did not have any expectation about the upcoming probe

syllable. Thus, training and testing phases for the classifier were separate and temporally

independent from each other (especially, for the valid retro-cue trials).

To estimate neural activity patterns during the cue-retention and the auditory probe

phases used for the MVPA, we first constructed a single subject-level GLM for each functional

run. As in the univariate analysis, a 0–20 s FIR was used to capture the hemodynamic

responses ranging from the retro-cue onset, the following retention, to the auditory probe

phases of the task. Four, 20-s FIRs were included in the GLM to separately model the

hemodynamic response of each condition: 2 cue types (valid and neutral) × 2 to-be-probed

syllable categories (i.e., /da/ and /ge/). From each run-wise GLM, condition-specific activation

across voxels were estimated with *t*-estimates for the MVPA (Misaki et al., 2010). The neural

activity during the cue-retention phase was quantified by averaging the *t*-estimates of the FIR

during 4–8 s from the cue onset; and the neural activity evoked by the auditory probe was

quantified by averaging the *t*-estimates during 10–12 s of the FIR. These time windows were

selected to account for about 4–6 s delay of the hemodynamic responses to these trial events

(see Fig. 2B for the average activations during all time points of the FIR). The GLM also

included additional canonical regressors to account for BOLD responses during encoding of

two speech syllables during trial onset, and during participants' response and visual feedback,

as well as continuous regressors to model head motion and scanner drift.

For cross-validation of the generalization of classifier performance, eight functional

runs were divided into training and test datasets. The classifier was trained on the auditory

probe phase activity of the neutral-cue trials of five randomly selected functional runs, and the

classifier was tested on the cue–retention phase activity of each cue condition from the

remaining three functional runs. The cross-validation iteration was performed 28 times (i.e., a

half of all possible combinations of five training and three test datasets), and the decoding

accuracy was calculated based on the average performances across all iterations, separately

for each cue condition. Searchlight MVPA was performed on unsmoothed functional data in

each participant's native space. The resulting decoding accuracy map of each participant was

centered at zero (i.e., accuracy minus 50% chancel level), and normalized to standard MNI

space. The group-level accuracy map was determined by averaging the classification

accuracy across participants.

To determine statistical assessment of classification performance above chance, we

adopted a non-parametric permutation approach in order (Stelzer et al., 2013; Allefeld et al.,

2016). To this end, we repeated the whole-brain searchlight analysis with randomly assigned

classification labels for 1000 times for each participant. We then generated a permuted group-

level accuracy map by averaging one randomly selected map out of the 1000 permuted maps

drawn from each participant (with replacement); this process was repeated 10^5 times, which

resulted in 10^5 permuted group-level accuracy maps. Based on the null-hypothesis distribution

(permutation) maps of chance level decoding, we computed a map of voxel-wise significance

value for the un-permuted group accuracy map, and corrected for multiple comparisons (FDR

at $p < 0.05$). From these regions that survived the statistical threshold, we extracted decoding

accuracies for the valid and neutral retro-cue conditions.

Brain–Behavior relationships. In order to understand how the network of task-
relevant brain regions contribute to the representational precision benefits from valid retro-
cues (i.e., Valid – Neutral: Δ ln slope k), we performed a multiple linear regression analysis to
examine the relationship between the degrees of neural modulations and the representational
precision modulations from the retro-cues (i.e., Valid vs. Neutral) across individuals. The two
types of neural modulation measures, one from the univariate and the other from the
multivariate results, were analyzed separately to explain the extent of the precision
enhancement with valid (vs. neutral) retro-cues.

Starting from a model that includes all factors representing neural modulations of
multiple brain regions, we used a step-wise model comparison to find the best model that
explains the precision modulations from valid vs. neutral cues. The model comparison was
based on the AIC of the model (*StepAIC* from the package *MASS* ver. 7.3-49 in R) as well as
χ^2 testing to deduce into a simplest model that explains the cue-related modulations of the
perceptual precision.

In analyzing the univariate BOLD data, we performed the model comparison, starting
from the full model—that is, the neural activity modulations derived from the four ROIs,
including the bilateral auditory region (Fig. 2), to explain the degree of the perceptual precision
modulations (Δ ln slope k):

Δ ln slope $k = \Delta$ Auditory $+ \Delta$ Fronto-Parietal $+ \Delta$ Cingulo-Opercular $+ \Delta$ Default-mode
where Δ represents the degree of cue-related modulation (i.e., Valid – Neutral).

Using a similar approach, we analyzed the multivariate data to understand whether the
neural decoding accuracy modulations in the task-relevant brain regions were related to the
extent of precision modulation from valid vs. neutral cues. To this end, we started the model
comparison from the task-relevant cortical regions that exhibited significantly accurate
decoding of to-be-probed syllable categories during the memory retention period (Fig. 3; Table
2). These regions were selected based on whether they fell within the language processing
regions (Baddeley, 2003; Hickok and Poeppel, 2007; Koelsch et al., 2009), auditory regions
in the temporal cortex, domain-general attentional and cognitive control network regions (i.e.,
fronto-parietal and cingulo-opercular networks). Starting from the full-factor model that
included the neural decoding accuracy modulations from 13 brain regions (Table 2), we used
a step-wise model comparison to deduce from the full-factor model to find the simplest/best
model that explains the extent of precision modulation (Δ ln slope k: ln slope k_{Valid} – ln slope
k_{Neutral}).

Note that all of the cue-related modulation data (i.e., the perceptual precision
modulation and neural modulations across the brain regions) were z-scored prior to the model

testing. In order to ensure the validity of the final model resulted from the model comparison

step, we performed a leave-one-subject-out cross-validation (i.e., 22 CV's). On each CV step,

we used n=21 subjects' data to fit the model parameters, and tested on the remaining one

subject's data. We compared the cross-validated fitted values vs. the actual data outcome

using a correlation test; and we report the average adjusted R^2 from 22 cross-validation steps.

1017

Acknowledgements

This work was supported by the Max Planck Society (Max Planck research grant to JO) and

the European Research Council (ERC Consolidator grant AUDADAPT, no. 646696, to JO).

We thank Steven Kalinke, Sören Nikolaus, and Dunja Kunke for their help in data collection

of the study. We would also like to thank Mandy Jochemko, Anke Kummer, and Nicole

Pampus for their assistance at the MRI scanner.

1024

Competing interests

1026 The authors declare that they have no competing interests.

1027

References

Alain, C., & Arnott, S. R. (2000). Selectively attending to auditory objects. *Frontiers in*
Biosciences, *5*, d202–212.

Alavash, M., Lim, S.-J., Thiel, C., Sehm, B., Deserno, L., & Obleser, J. (2018). Dopaminergic
modulation of hemodynamic signal variability and the functional connectome during
cognitive performance. *NeuroImage, 172*, 341–356.
http://doi.org/10.1016/j.neuroimage.2018.01.048

Allefeld, C., Görgen, K., & Haynes, J.-D. (2016). Valid population inference for information-
based imaging: From the second-level t-test to prevalence inference. *NeuroImage*, *141*,
378–392. http://doi.org/10.1016/j.neuroimage.2016.07.040

Awh, E., Jonides, J., Smith, E. E., Schumacher, E. H., Koeppe, R. A., & Katz, S. (1996).
Dissociation of Storage and Rehearsal in Verbal Working Memory: Evidence From
Positron Emission Tomography:. *Psychological Science*, *7*(1), 25–31.
http://doi.org/10.1111/j.1467-9280.1996.tb00662.x

Bachorowski, J.-A., & Owren, M. J. (1999). Acoustic correlates of talker sex and individual
talker identity are present in a short vowel segment produced in running speech. *Journal*
of the Acoustical Society of America, *106*(2), 1054–1063.
http://doi.org/10.1121/1.427115

Backer, K. C., Binns, M. A., & Alain, C. (2015). Neural dynamics underlying attentional
orienting to auditory representations in short-term memory. *Journal of Neuroscience*,
35(3), 1307–1318. http://doi.org/10.1523/JNEUROSCI.1487-14.2015

Baddeley, A. (1992). Working memory. *Science*, *255*(5044), 556–559.

Baddeley, A. (2003). Working memory: looking back and looking forward. *Nature Reviews.*
Neuroscience, *4*(10), 829–839. http://doi.org/10.1038/nrn1201

Bays, P. M., & Husain, M. (2008). Dynamic shifts of limited working memory resources in
human vision. *Science*, *321*(5890), 851–854. http://doi.org/10.1126/science.1160575

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B*
(Methodological), *57*(1), 289–300. http://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Bettencourt, K. C., & Xu, Y. (2016). Decoding the content of visual short-term memory under
distraction in occipital and parietal areas. *Nature Reviews. Neuroscience*, *19*(1), 150–
157. http://doi.org/10.1038/nn.4174

Bledowski, C., Rahm, B., & Rowe, J. B. (2009). What "Works" in Working Memory?
Separate Systems for Selection and Updating of Critical Information. *Journal of*
Neuroscience, 29(43), 13735–13741. http://doi.org/10.1523/JNEUROSCI.2547-09.2009

Chang, C.-C., & Lin, C.-J. (2011). LIBSVM: A Library for Support Vector Machines. *ACM*
Transactions on Intelligent Systems and Technology, *2*(3), 27:1–27:27.

Christophel, T. B., Klink, P. C., Spitzer, B., Roelfsema, P. R., & Haynes, J.-D. (2017). The
Distributed Nature of Working Memory. *Trends in Cognitive Sciences*, *21*(2), 111–124.
http://doi.org/10.1016/j.tics.2016.12.007

Cohen, L., Lehericy, S., Chochon, F., Lemer, C., Rivaud, S., & Dehaene, S. (2002).
Language-Specific Tuning of Visual Cortex? Functional Properties of the Visual Word
Form Area. *Brain*, *125*, 1054–1069. http://doi.org/doi.org/10.1093/brain/awf094

Cowan, N. (2001). Metatheory of storage capacity limits. *Behavioral and Brain Sciences*,
24(01), 154–176.

Cox, R. W. (1996). AFNI: software for analysis and visualization of functional magnetic
resonance neuroimages. *Computers and Biomedical Research, an International Journal*,
29(3), 162–173.

Desimone, R., & Duncan, J. (1995). Neural Mechanisms of Selective Visual Attention.
Annual Review of Neuroscience, *18*(1), 193–222.
http://doi.org/10.1146/annurev.ne.18.030195.001205

Dosenbach, N. U. F., Fair, D. A., Cohen, A. L., Schlaggar, B. L., & Petersen, S. E. (2008). A
dual-networks architecture of top-down control. *Trends in Cognitive Sciences*, *12*(3), 99–
105. http://doi.org/10.1016/j.tics.2008.01.001

Dosenbach, N. U. F., Fair, D. A., Miezin, F. M., Cohen, A. L., Wenger, K. K., Dosenbach, R.
A. T., et al. (2007). Distinct brain networks for adaptive and stable task control in
humans. *Proceedings of the National Academy of Sciences*, *104*(26), 11073–11078.

Duncan, J. (1984). Selective Attention and the Organization of Visual Information. *Journal of*
Experimental Psychology. General, *113*(4), 51–517.

Duncan, J., & Owen, A. M. (2000). Common regions of the human frontal lobe recruited by
diverse cognitive demands. *Trends in Neurosciences*, *23*(10), 475–483.
http://doi.org/10.1016/S0166-2236(00)01633-7

Emrich, S. M., Riggall, A. C., LaRocque, J. J., & Postle, B. R. (2013). Distributed Patterns of
Activity in Sensory Cortex Reflect the Precision of Multiple Items Maintained in Visual
Short-Term Memory. *Journal of Neuroscience*, *33*(15), 6516–6523.
http://doi.org/10.1523/JNEUROSCI.5732-12.2013

Erb, J., Henry, M. J., Eisner, F., & Obleser, J. (2013). The brain dynamics of rapid
perceptual adaptation to adverse listening conditions. *Journal of Neuroscience*, *33*(26),
10688–10697. http://doi.org/10.1523/JNEUROSCI.4596-12.2013

Ester, E. F., Sprague, T. C., & Serences, J. T. (2015). Parietal and Frontal Cortex Encode

Stimulus-Specific Mnemonic Representations during Visual Working Memory. *Neuron,*

87(4), 893–905. http://doi.org/10.1016/j.neuron.2015.07.013

Fedorenko, E., Behr, M. K., & Kanwisher, N. (2011). Functional specificity for high-level

linguistic processing in the human brain. *Proceedings of the National Academy of*

Sciences, *108*(39), 16428–16433. http://doi.org/10.1073/pnas.1112937108

Feng, G., Yi, H.-G., & Chandrasekaran, B. (2018). The Role of the Human Auditory

Corticostriatal Network in Speech Learning. *Cerebral Cortex, 29*(10), 4077–4089.

http://doi.org/10.1093/cercor/bhy289

Formisano, E., De Martino, F., Bonte, M., & Goebel, R. (2008). "Who" is saying "what?"

Brain-based decoding of human voice and speech. *Science, 322*(5903), 970–973.

http://doi.org/10.1126/science.1164318

Freedman, D. J., Riesenhuber, M., Poggio, T., & Miller, E. K. (2001). Categorical

representation of visual stimuli in the primate prefrontal cortex. *Science, 291*(5502),

312–316. http://doi.org/10.1126/science.291.5502.312

Fritz, J. B., Elhilali, M., David, S. V., & Shamma, S. A. (2007). Auditory attention — focusing

the searchlight on sound. *Current Opinion in Neurobiology, 17*(4), 437–455.

http://doi.org/10.1016/j.conb.2007.07.011

Gaab, N., Gaser, C., Zaehle, T., Jancke, L., & Schlaug, G. (2003). Functional anatomy of

pitch memory—an fMRI study with sparse temporal sampling. *NeuroImage, 19*(4),

1417–1426. http://doi.org/10.1016/S1053-8119(03)00224-6

Gazzaley, A., & Nobre, A. C. (2012). Top-down modulation: bridging selective attention and

working memory. *Trends in Cognitive Sciences, 16*(2), 129–135.

http://doi.org/10.1016/j.tics.2011.11.014

Goldman-Rakic, P. S. (1995). Cellular Basis of Working Memory Review. *Neuron, 14*, 477–

485.

Griffin, I. C., & Nobre, A. C. (2003). Orienting Attention to Locations in Internal

Representations. *Journal of Cognitive Neuroscience, 15*(8), 1176–1194.

Harrison, S. A., & Tong, F. (2009). Decoding reveals the contents of visual working memory

in early visual areas. *Nature, 458*(7238), 632–635. http://doi.org/10.1038/nature07832

Hartwigsen, G., Baumgaertner, A., Price, C. J., Koehnke, M., Ulmer, S., & Siebner, H. R.

(2010). Phonological decisions require both the left and right supramarginal gyri.

Proceedings of the National Academy of Sciences, 107(38), 16494–16499.

http://doi.org/10.1073/pnas.1008121107

Haynes, J.-D., Sakai, K., Rees, G., Gilbert, S., Frith, C., Passingham, R. E. (2007). Reading

hidden intentions in the human brain. *Current Biology 17*, 323–328.

Hickok, G., & Poeppel, D. (2007). The cortical organization of speech processing. *Nature*

Reviews. Neuroscience, *8*(5), 393–402. http://doi.org/10.1038/nrn2113

Higo, T., Mars, R. B., Boorman, E. D., Buch, E. R., & Rushworth, M. F. S. (2011). Distributed

and causal influence of frontal operculum in task control. *Proceedings of the National*

Academy of Sciences of the United States of America, *108*(10), 4230–4235.

http://doi.org/10.1073/pnas.1013361108

Jacquemot, C., & Scott, S. K. (2006). What is the relationship between phonological short-

term memory and speech processing? *Trends in Cognitive Sciences*, *10*(11), 480–486.

http://doi.org/10.1016/j.tics.2006.09.002

Koelsch, S., Schulze, K., Sammler, D., Fritz, T., Müller, K., & Gruber, O. (2009). Functional

architecture of verbal and tonal working memory: an FMRI study. *Human Brain*

Mapping, *30*, 859–873. http://doi.org/10.1002/hbm.20550'

Kriegeskorte, N., Goebel, R., & Bandettini, P. (2006). Information-based functional brain

mapping. *Proceedings of the National Academy of Sciences*, *103*(10), 3863–3868.

http://doi.org/10.1073/pnas.0600244103

Kumar, S., Joseph, S., Gander, P. E., Barascud, N., Halpern, A. R., & Griffiths, T. D. (2016).

A Brain System for Auditory Working Memory. *Journal of Neuroscience*, *36*(16), 4492–

4505. http://doi.org/10.1523/JNEUROSCI.4341-14.2016

Kuo, B.-C., Stokes, M. G., & Nobre, A. C. (2012). Attention modulates maintenance of

representations in visual short-term memory. *Journal of Cognitive Neuroscience*, *24*(1),

51–60. http://doi.org/10.1162/jocn_a_00087

Lee, S.-H., Kravitz, D. J., & Baker, C. I. (2013). Goal-dependent dissociation of visual and

prefrontal cortices during working memory. *Nature Neuroscience*, *16*(8), 997–999.

http://doi.org/10.1038/nn.3452

Leech, R., Holt, L. L., Devlin, J. T., & Dick, F. (2009). Expertise with artificial nonspeech

sounds recruits speech-sensitive cortical regions. *Journal of Neuroscience*, *29*(16),

5234–5239. http://doi.org/10.1523/JNEUROSCI.5758-08.2009

Lepsien, J., & Nobre, A. C. (2007). Attentional Modulation of Object Representations in

Working Memory. *Cerebral Cortex*, *17*(9), 2072–2083.

http://doi.org/10.1093/cercor/bhl116

Lepsien, J., Griffin, I. C., Devlin, J. T., & Nobre, A. C. (2005). Directing spatial attention in

mental representations: Interactions between attentional orienting and working-memory

load. *NeuroImage*, *26*(3), 733–743. http://doi.org/10.1016/j.neuroimage.2005.02.026

Lepsien, J., Thornton, I., & Nobre, A. C. (2011). Modulation of working-memory maintenance

by directed attention. *Neuropsychologia*, *49*(6), 1569–1577.

http://doi.org/10.1016/j.neuropsychologia.2011.03.011

Lewis-Peacock, J. A., Drysdale, A. T., Oberauer, K., & Postle, B. R. (2012). Neural Evidence

for a Distinction between Short-term Memory and the Focus of Attention. *Journal of*

Cognitive Neuroscience, *24*(1), 61–79. http://doi.org/10.1162/jocn_a_00140

Lim, S.-J., Fiez, J. A., & Holt, L. L. (2019). Role of the striatum in incidental learning of sound

categories. *Proceedings of the National Academy of Sciences*, 1–10.

http://doi.org/10.1073/pnas.1811992116

Lim, S.-J., Wöstmann, M., & Obleser, J. (2015). Selective Attention to Auditory Memory

Neurally Enhances Perceptual Precision. *Journal of Neuroscience*, *35*(49), 16094–

16104. http://doi.org/10.1523/JNEUROSCI.2674-15.2015

Lim, S.-J., Wöstmann, M., Geweke, F., & Obleser, J. (2018). The Benefit of Attention-to-

Memory Depends on the Interplay of Memory Capacity and Memory Load. *Frontiers in*

Psychology, *9*, 146. http://doi.org/10.3389/fpsyg.2018.00184

Linke, A. C., & Cusack, R. (2015). Flexible Information Coding in Human Auditory Cortex

during Perception, Imagery, and STM of Complex Sounds. *Journal of Cognitive*

Neuroscience, *27*(7), 1322–1333. http://doi.org/10.1162/jocn_a_00780

Linke, A. C., Vicente-Grabovetsky, A., & Cusack, R. (2011). Stimulus-specific suppression

preserves information in auditory short-term memory. *Proceedings of the National*

Academy of Sciences of the United States of America, *108*(31), 12961–12966.

http://doi.org/10.1073/pnas.1102118108

Long, N. M., & Kuhl, B. A. (2018). Bottom-up and top-down factors differentially influence

stimulus representations across large-scale attentional networks. *Journal of*

Neuroscience, *38*(10), 2724–17–2504. http://doi.org/10.1523/JNEUROSCI.2724-

17.2018

Luck, S. J., & Vogel, E. K. (2013). Visual working memory capacity: from psychophysics and

neurobiology to individual differences. *Trends in Cognitive Sciences*, *17*(8), 391–400.

http://doi.org/10.1016/j.tics.2013.06.006

Ma, W. J., Husain, M., & Bays, P. M. (2014). Changing concepts of working memory. *Nature*

Reviews. Neuroscience, *17*(3), 347–356. http://doi.org/10.1038/nn.3655

Manohar, S. G., Zokaei, N., Fallon, S. J., Vogels, T. P., & Husain, M. (2019). Neural

mechanisms of attending to items in working memory. *Neuroscience and Biobehavioral*

Reviews, *101*, 1–12. http://doi.org/10.1016/j.neubiorev.2019.03.017

Mesgarani, N., Cheung, C., Johnson, K., & Chang, E. F. (2014). Phonetic feature encoding
in human superior temporal gyrus. *Science*, *343*(6174), 1006–1010.
http://doi.org/10.1126/science.1245994
Misaki, M., Kim, Y., Bandettini, P. A., & Kriegeskorte, N. (2010). Comparison of multivariate
classifiers and response normalizations for pattern-information fMRI. *NeuroImage*,
53(1), 103–118. http://doi.org/10.1016/j.neuroimage.2010.05.051
Murray, A. M., Nobre, A. C., Clark, I. A., Cravo, A. M., & Stokes, M. G. (2013). Attention
Restores Discrete Items to Visual Short-Term Memory. *Psychological Science : a*
Journal of the American Psychological Society / APS, *24*(4), 550–556.
http://doi.org/10.1177/0956797612457782
Myers, E. B., Blumstein, S. E., Walsh, E., & Eliassen, J. (2009). Inferior frontal regions
underlie the perception of phonetic category invariance. *Psychological Science*, *20*(7),
895–903. http://doi.org/10.1111/j.1467-9280.2009.02380.x
Myers, N. E., Stokes, M. G., & Nobre, A. C. (2017). Prioritizing Information during Working
Memory: Beyond Sustained Internal Attention. *Trends in Cognitive Sciences*, *21*(6),
449–461. http://doi.org/10.1016/j.tics.2017.03.010
Noyce, A. L., Cestero, N., Michalka, S. W., Shinn-Cunningham, B. G., & Somers, D. C.
(2017). Sensory-Biased and Multiple-Demand Processing in Human Lateral Frontal
Cortex. *Journal of Neuroscience*, *37*(36), 8755–8766.
http://doi.org/10.1523/JNEUROSCI.0660-17.2017
Obleser, J., & Eisner, F. (2009). Pre-lexical abstraction of speech in the auditory cortex.
Trends in Cognitive Sciences, *13*(1), 14–19. http://doi.org/10.1016/j.tics.2008.09.005
Obleser, J., Boecker, H., Drzezga, A., Haslinger, B., Hennenlotter, A., Roettinger, M., et al.
(2006). Vowel sound extraction in anterior superior temporal cortex. *Human Brain*
Mapping, *27*(7), 562–571. http://doi.org/10.1002/hbm.20201
Owen, A. M., McMillan, K. M., Laird, A. R., & Bullmore, E. (2005). N-back working memory
paradigm: A meta-analysis of normative functional neuroimaging studies. *Human Brain*
Mapping, *25*(1), 46–59. http://doi.org/10.1002/hbm.20131
Perrachione, T. K., Ghosh, S. S., Ostrovskaya, I., Gabrieli, J. D. E., & Kovelman, I. (2017).
Phonological Working Memory for Words and Nonwords in Cerebral Cortex. *Journal of*
Speech, Language, and Hearing Research, *60*(7), 1959–1979.
http://doi.org/10.1044/2017_JSLHR-L-15-0446
Postle, B. R. (2006). Working memory as an emergent property of the mind and brain.
Neuroscience, *139*(1), 23–38. http://doi.org/10.1016/j.neuroscience.2005.06.005

Postle, B. R., Stern, C. E., Rosen, B. R., & Corkin, S. (2000). An fMRI Investigation of Cortical Contributions to Spatial and Nonspatial Visual Working Memory. *NeuroImage*, *11*(5), 409–423. http://doi.org/10.1006/nimg.2000.0570

Power, J. D., Cohen, A. L., Nelson, S. M., Wig, G. S., Barnes, K. A., Church, J. A., et al. (2011). Functional Network Organization of the Human Brain. *Neuron*, *72*(4), 665–678. http://doi.org/10.1016/j.neuron.2011.09.006

Price, C. J., Moore, C. J., Humphreys, G. W., & Wise, R. J. S. (1997). Segregating semantic from phonological processes during reading. *Journal of Cognitive Neuroscience*, *9*(6), 727–733. http://doi.org/10.1162/jocn.1997.9.6.727

Qi, Z., Han, M., Wang, Y., de Los Angeles, C., Liu, Q., Garel, K., et al. (2019). Speech processing and plasticity in the right hemisphere predict variation in adult foreign language learning. *NeuroImage*, *192*, 76–87. http://doi.org/10.1016/j.neuroimage.2019.03.008

Raichle, M. E., MacLeod, A. M., Snyder, A. Z., Powers, W. J., Gusnard, D. A., & Shulman, G. L. (2001). A default mode of brain function. *Proceedings of the National Academy of Sciences*, *98*(2), 676–682.

Rerko, L., & Oberauer, K. (2013). Focused, unfocused, and defocused information in working memory. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *39*(4), 1075–1096. http://doi.org/10.1037/a0031172

Riggall, A. C., & Postle, B. R. (2012). The Relationship between Working Memory Storage and Elevated Activity as Measured with Functional Magnetic Resonance Imaging. *Journal of Neuroscience*, *32*(38), 12990–12998. http://doi.org/10.1523/JNEUROSCI.1892-12.2012

Rowe, J. B., Toni, I., Josephs, O., Frackowiak, R. S., & Passingham, R. E. (2000). The prefrontal cortex: response selection or maintenance within working memory? *Science*, *288*(5471), 1656–1660.

Sadaghiani, S., & D'Esposito, M. (2014). Functional Characterization of the Cingulo-Opercular Network in the Maintenance of Tonic Alertness. *Cerebral Cortex*, 1–11. http://doi.org/10.1093/cercor/bhu072

Scott, T. L., & Perrachione, T. K. (2019). Common cortical architectures for phonological working memory identified in individual brains. *NeuroImage*, *202*, 116096. http://doi.org/10.1016/j.neuroimage.2019.116096

Seeley, W. W., Menon, V., Schatzberg, A. F., Keller, J., Glover, G. H., Kenna, H., et al. (2007). Dissociable Intrinsic Connectivity Networks for Salience Processing and

Executive Control. *Journal of Neuroscience, 27*(9), 2349–2356.

http://doi.org/10.1523/JNEUROSCI.5587-06.2007

Serences, J. T., Ester, E. F., Vogel, E. K., & Awh, E. (2009). Stimulus-specific delay activity

in human primary visual cortex. *Psychological Science, 20*(2), 207–214.

http://doi.org/10.1111/j.1467-9280.2009.02276.x

Shinn-Cunningham, B. G. (2008). Object-based auditory and visual attention. *Trends in*

Cognitive Sciences, 12(5), 182–186. http://doi.org/10.1016/j.tics.2008.02.003

Smith, E. E., Jonides, J., Marshuetz, C., & Koeppe, R. A. (1998). Components of verbal

working memory: Evidence from neuroimaging. *Proceedings of the National Academy of*

Sciences, 95(3), 876–882. http://doi.org/10.1073/pnas.95.3.876

Souza, A. S., & Oberauer, K. (2016). In search of the focus of attention in working memory:

13 years of the retro-cue effect. *Attention, Perception, & Psychophysics, 78*(7), 1839–

1860. http://doi.org/10.3758/s13414-016-1108-5

Souza, A. S., Rerko, L., & Oberauer, K. (2014). Unloading and reloading working memory:

Attending to one item frees capacity. *Journal of Experimental Psychology. Human*

Perception and Performance, 40(3), 1237–1256. http://doi.org/10.1037/a0036331

Spitzer, B., Fleck, S., & Blankenburg, F. (2014). Parametric Alpha- and Beta-Band

Signatures of Supramodal Numerosity Information in Human Working Memory. *Journal*

of Neuroscience, 34(12), 4293–4302. http://doi.org/10.1523/JNEUROSCI.4580-13.2014

Sprague, T. C., Ester, E. F., & Serences, J. T. (2016). Restoring Latent Visual Working

Memory Representations in Human Cortex. *Neuron, 91*(3), 694–707.

http://doi.org/10.1016/j.neuron.2016.07.006

Sreenivasan, K. K., Gratton, C., Vytlacil, J., & D'Esposito, M. (2014). Evidence for working

memory storage operations in perceptual cortex. *Cognitive, Affective & Behavioral*

Neuroscience, 14(1), 117–128. http://doi.org/10.3758/s13415-013-0246-7

Stelzer, J., Chen, Y., & Turner, R. (2013). Statistical inference and multiple testing correction

in classification-based multi-voxel pattern analysis (MVPA): Random permutations and

cluster size control. *NeuroImage, 65*, 69–82.

http://doi.org/10.1016/j.neuroimage.2012.09.063

Stokes, M. G. (2015). "Activity-silent" working memory in prefrontal cortex: a dynamic coding

framework. *Trends in Cognitive Sciences, 19*(7), 394–405.

http://doi.org/10.1016/j.tics.2015.05.004

Stokes, M. G., Kusunoki, M., Sigala, N., Nili, H., Gaffan, D., & Duncan, J. (2013). Dynamic

Coding for Cognitive Control in Prefrontal Cortex. *Neuron, 78*(2), 364–375.

http://doi.org/10.1016/j.neuron.2013.01.039

Todd, J. J., & Marios, R. (2004). Capacity limit of visual short-term memory in human
posterior parietal cortex. *Nature Neuroscience*, *428*, 751–754.

Uluç, I., Schmidt, T. T., Wu, Y.-H., & Blankenburg, F. (2018). Content-specific codes of
parametric auditory working memory in humans. *NeuroImage*, *183*, 254–262.

Vaden, K. I., Kuchinsky, S. E., Cute, S. L., Ahlstrom, J. B., Dubno, J. R., & Eckert, M. A.
(2013). The Cingulo-Opercular Network Provides Word-Recognition Benefit. *Journal of*
Neuroscience, *33*(48), 18979–18986. http://doi.org/10.1523/JNEUROSCI.1417-13.2013

Wallis, G., Stokes, M., Cousijn, H., Woolrich, M., & Nobre, A. C. (2015). Frontoparietal and
Cingulo-opercular Networks Play Dissociable Roles in Control of Working Memory.
Journal of Cognitive Neuroscience, *15*, 1–16. http://doi.org/10.1162/jocn_a_00838

Wilsch, A., & Obleser, J. (2016). What works in auditory working memory? A neural
oscillations perspective. *Brain Research*, *1640* (Part B), 193–207.
http://doi.org/10.1016/j.brainres.2015.10.054

Wolff, M. J., Jochim, J., Akyürek, E. G., & Stokes, M. G. (2017). Dynamic hidden states
underlying working-memory-guided behavior. *Nature Reviews. Neuroscience*, *20*(6),
864–871. http://doi.org/10.1038/nn.4546

Xu, Y., & Chun, M. M. (2006). Dissociable neural mechanisms supporting visual short-term
memory for objects. *Nature*, *440*(7080), 91–95. http://doi.org/10.1038/nature04262

Zhang, W., & Luck, S. J. (2008). Discrete fixed-resolution representations in visual working
memory. *Nature*, *453*(7192), 233–235. http://doi.org/10.1038/nature06860

Zokaei, N. (2014). Flexibility of representational states in working memory, *Frontiers in Human*
Neuroscience, 8, 853.

1325