Development of larvae of the Australian blowfly, *Calliphora augur* (Diptera: Calliphoridae), at constant temperatures

Donnah M. Day[1]

Nathan J. Butterworth[2*], ORCID: 0000-0002-5679-2700, Twitter: @Butterworthbugs

Anirudh Tagat[3], ORCID: 0000-0002-7707-453X, Twitter: @inhouseeconomist

Gregory Markowsky[3], ORCID: 0000-0003-1656-337X

James F. Wallman[1,4], ORCID: 0000-0003-2504-6075

[1]*Centre for Sustainable Ecosystem Solutions, School of Earth, Atmospheric and Life Sciences, University of Wollongong, NSW 2522, Australia*

[2]*School of Life Sciences, University of Technology Sydney, Ultimo, NSW 2007, Australia*

[3]*School of Mathematics, Monash University, VIC 3800, Australia*

[4]*Faculty of Science, University of Technology Sydney, Ultimo, NSW 2007, Australia*

* Corresponding author

E-mail address: nathan.butterworth@uts.edu.au

Keywords: Forensic entomology, post-mortem interval, prediction interval

Abstract

Calliphora augur (Diptera: Calliphoridae) is a common carrion-breeding blowfly of forensic, medical and agricultural importance in eastern Australia. Despite this, detailed information on the developmental biology of *C. augur* is lacking. Here, we present the first comprehensive study on the development of all three larval instars of *C. augur*, fed on sheep's liver, at constant temperatures of 15, 20, 25, 30 and 35°C. We provide thermal summation models describing instar duration, as well as 95% prediction intervals for larval length at each constant temperature, enabling the age of larvae of *C. augur* to be estimated from their developmental stage and their average length. These data provide a basis for the application of this species to forensic investigations.

Introduction

Forensic entomology is the study of insects as applied to legal issues. It may relate to medicolegal events, especially murders, for the purpose of uncovering information useful to an investigation [1]. This is because decomposing vertebrate remains provide a temporary micro-habitat and food source that is attractive to a variety of insects and other invertebrates [2]. Arthropods constitute a major element of this fauna, with insects, especially flies and their larvae, being the predominant group. Blowflies (Diptera: Calliphoridae) are ubiquitous and are typically the first insects to visit a dead body, often well before it is discovered by humans [3].

Data on the development of blowflies and other important carrion flies can help provide an estimate of the minimum time since death (or minimum post-mortem interval: mPMI), especially when combined with information on the succession patterns of insects on corpses[4]. Current approaches to age estimation are based on species-specific rates of development such as length, weight, and developmental duration in relation to temperature. While measures based on length and weight are valuable, developmental duration is the most widely used method of age estimation for two primary reasons; 1) because the duration of each stage of larval development (referred to as 'instars') is less variable than larval size (particularly in response to differences in diet)[5]; and 2) because instar stage is not affected by specimen shrinkage during preservation[6]. Once these measurements are obtained for the entire period of larval development, a variety of models can be used to estimate mPMI; including curvilinear regression [8], isomegalen and isomorphen diagrams [9], and thermal summation models [10,11]. Of these, the most widely used are thermal summation models that incorporate temperature (usually as degree-hours) and time spent in each developmental stage [12].

Calliphora augur Fabricius is a common blowfly species found in eastern Australia [13], the larvae of which are among those most commonly collected from forensic cases, including from inside buildings [14]. It is ovoviviparous [15], and also an agent of myiasis in sheep and other animals [16–18]. Adult females prefer fresh carcasses [19], but will also larviposit on carcasses in active decomposition (i.e., 10-15 days old) [20]. While previous studies have considered the larval development of *C. augur* [16,20–24] none have made fine-scale size measurements across a range of constant temperatures, or used thermal summation models to estimate mPMI. This lack of accurate developmental data means that at present, *C. augur* cannot be used reliably in a forensic context. To ameliorate this, we present the first comprehensive analysis of larval growth at constant temperatures of this important species. Our investigations are based on measurement of larval length throughout development, as well as

64 observation of larval instar duration. Most importantly, we provide thermal summation models based
65 on instar duration, and calculate prediction intervals based on larval length – for the first time allowing
66 the application of this species to forensic investigations.

Materials and methods

Fly culture

A culture of *C. augur* was established from adults trapped at Wollongong, New South Wales (34° 25'
S, 150°53' E). Experiments commenced using larvae of the F3 generation and were continued for 11
generations. This culture was refreshed regularly with wild-type (F_0) individuals from the same source
population as above and maintained with mixed ages of females and males. The average fecundity of
each female was estimated by separating 20 females and transferring them to 50 mL specimen pots
containing approximately 10 g of freshly defrosted sheep's liver (defrosted to room temperature
immediately prior to experiments). Larvae were counted the following day. Larvae and adults were
kept in a temperature-controlled room at 25±3.5˚C and ambient humidity, with a 12:12 light:dark
regime incorporating a 15 minute 'dusk' and 'dawn' transition period of low light. Adult flies were
maintained in square plastic cages measuring 330 mm long, 220 mm wide and 250 mm high (external
dimensions). The flies in each cage were provided with water and sugar *ad libitum*, and chopped
sheep's liver in a small plastic weigh boat for larviposition as required. Larvae were reared in square
white 2 L plastic containers, measuring 170 mm wide along each side and 90 mm deep. Most of the
centre of the lid of each container was cut away and replaced with fine mesh to permit ventilation. In
each container, larvae were sustained on sheep's liver in the weigh boats placed atop a layer of wheaten
chaff approximately 20 mm deep to enable pupariation. After pupariation, pupae were manually sorted
from the chaff and transferred to a cage. Sheep's liver was provided just prior to eclosion to provide
females with a constant source of protein for ovary maturation [20].

Development at constant temperatures

To assess rates of development at constant temperatures, freshly laid larvae were randomly collected
from the main culture. Preliminary studies on larval density indicated that some endogenous heat
generation occurred with as few as 25 larvae growing on 50 g of sheep's liver. To ensure that the
larvae developed at the ambient temperature being examined, each replicate (a plastic weigh boat
containing larvae and liver defrosted to room temperature immediately prior to the experiment) was
limited to 10 larvae. The growth period (between 0 and 600 hours) of each replicate of 10 larvae at a
particular temperature was chosen by lottery. Between three and six replicates were used for each time

interval (Supplementary Material 1). These replicates were grown at specific constant temperature regimes (15, 20, 25, 30 and 35˚C ± 1.0 °C) and 60 ± 5% relative humidity, in an Axyos (Brisbane, Queensland, Australia) environmental cabinet. To ensure that the temperature inside the cabinet matched the set value, each replicate also contained a temperature logger (iButton; accuracy ±1.0 °C, resolution 0.5 °C; Maxim Integrated Products, Sunnyvale, CA, USA). Since only one cabinet was available, growth at each temperature was studied in turn. Larvae were left undisturbed in darkness in the environmental cabinet until collection, because disturbance can delay pupariation [20,24]. To account for growth rates increasing with temperature, we sampled more frequently at higher temperatures. For 15°C, sampling was conducted every twenty-four hours, up to a total of 600 hours. For 20°C, sampling was conducted every six hours, up to a total of 216 hours. For 25, 30 and 35°C sampling was conducted every four hours, up to a total of 204 hours. The choice of these specific sampling frequencies was guided by the growth curves of O'Flynn[24,25], and frequencies were kept relatively high to minimise error in the calculation of the prediction intervals.

Sample collection, handling and preservation

At collection, all larvae were killed and fixed immediately by immersion in boiling water, dried with paper towel and preserved in 80% EtOH. Larvae were placed into glass vials posteriad to prevent head-curling [26] and length and instar were recorded.

Larval length measurements

The body lengths of larvae were measured with the aid of a dissecting microscope and Mitutoyo Absolute digimatic digital callipers (Kawasaki, Japan) after the larvae had been in preservative for a minimum of ten days [27]. Body length was measured as the distance, viewed laterally, between the most distal parts of the head and the last abdominal segment. The lengths of pupae were also measured with callipers. The dataset consists of the measurements of the lengths of larvae grown at temperatures of 15°C, 20°C, 25°C, 30°C, and 35°C. Since no larvae were measured more than once, the length data for the various time intervals should be regarded as independent.

Larval instar durations

The instar of each larva was also recorded when it was measured as either 1st, 2nd, or 3rd, allowing for the calculation of thermal summation models based on instar duration. We estimated a thermal summation model separately for each stage of development. We estimated two parameters of the thermal summation model given by ($DT = k + t\,D$), where DT is the product of D, development duration (in hours) and T, environmental temperature in degrees celsius. The parameters, k (sum of

effective temperatures) and t (lower developmental threshold) were estimated using ordinary least squares regression as well as reduced major axis regression (RMA) (Ikemoto and Takai, 2000) using the statistical software Stata 16.1. The coefficient on development duration is recovered as *t* and the constant is recovered as *k*.

Results

On average, gravid females laid 58 ± 14 eggs. In total, 3,200 individual larvae (320 replicates of 10 larvae) were measured across a range of temperatures (15, 20, 25, 30 and 35°C ± 1.0°C) (Figure 1). Regarding the timing of developmental stages, the duration of each instar varied substantially between temperatures (Figure 2). As was expected, the average development time of each instar decreased with increasing temperature up to 30°C but increased slightly for third instars at 35°C. To determine the precise relationships between instar duration and degree-hours, thermal summation models were calculated using ordinary least squares regression (Figure 3) and reduced major axis regression (Figure 4). From these models, the sum of effective temperatures (*k*) and lower developmental threshold (*t*) were calculated for each instar for both the ordinary least squares regression (Table 1) and reduced major axis regression (Table 2).

From our measurements of larval size, we also provide 95% prediction intervals that can assist in the estimation of age (Supplementary Material 2). Growth curves revealed that first-instar larvae were largest on average when grown at 25°C. Similarly, second instars were also largest when grown at 25°C, but they also displayed a much smaller variation at this temperature compared with the other temperatures examined. Third-instar larvae were largest on average when grown at 20°C, but were above the overall mean (for third-instars) when growing at 15, 20, 25 and 30°C. The only temperature at which the average length of third instars dropped below their overall mean was at 35°C. Post-feeding third instars were largest on average when grown at 20°C.

Discussion

This study provides the first detailed dataset of the development of the Australian carrion-breeding blowfly *Calliphora augur*. The results we provide here reinforce the previous work done on this species but go substantially further by providing a level of statistical confidence in the application of developmental data in forensic investigations.

Fecundity

Robust and highly active first-instar larvae of *C. augur* hatch promptly after deposition, giving this

species an ecological advantage over oviparous species for exploiting small, quickly perishable

carcasses [17]. Johnston & Tiegs [30] reported that *C. augur* is capable of depositing eggs or larvae, with

the eggs usually hatching within six hours. In the present study, we noted that *C. augur* will normally

deposit living larvae but will also lay soft infertile eggs for some days before normal larviposition

occurs – similar to the observations of Mackerras [20] and O'Flynn [25]. The average number of larvae

per female has previously been noted as 50 [20] or 58 larvae [17]. Callinan [31] recorded a range of between

22.5 and 98.3 larvae per female in a controlled environment and a range of between 0.3 and 45.2 larvae

per female in a field environment. The average number of larvae deposited in the present study was

58 ± 14 per gravid female, which is consistent with these previous data.

Development at constant temperatures

The outcomes of some other developmental studies have been limited by a less thorough approach

than taken here. For example, Byrd and Butler [32] noted that their sampling method selected for the

fastest growing larvae, which often indicates a shorter period than could otherwise be deduced from

normal collection techniques at a crime scene. The influence of size extremes upon variance has also

been considered by Wells & LaMotte [7]. In the present study, we included larvae of all sizes to

demonstrate the full range that may be encountered in crime scene samples. Variation in larval body

length appeared to decrease steadily from 15 to 30°C and then increased again dramatically at 35°C.

The decrease in variation with increasing temperature may relate to a more even growth rate at higher

temperatures. However, the subsequent increase in variation at 35°C likely reflects the fact that the

larvae are reaching their thermal limit, at which point some larvae may begin to suffer from increased

heat and oxidative stress, hindering their growth and increasing the perceived variation in size[33]. The

variation in overlap of the growth stages, particularly the transitional forms between instars,

emphasises the importance of assessing crime scene samples for instar and/or transitional forms, and

collecting an adequate number of specimens. While there has been comment on the value of

transitional forms [34], for most blowfly species information on transitional forms is lacking.

The reader will note an interesting feature of the plots related to larval length, namely that the measured

lengths increase linearly with time to a certain point, but then begin to decrease after that. This late-

stage reduction in larval length is a result of post-feeding shrinkage that calliphorid larvae undergo

after leaving the carcass, prior to pupariation. This post-feeding shrinkage can cause some difficulty

in interpreting the age of larvae, because the smaller length of post-feeding larvae makes them appear

younger. However, this can be solved if investigators record where larvae are collected, as post-feeding
larvae will depart the carcass and move into the surrounding substrate. If larvae are collected away
from the carcass, it can be presumed that they are in the post-feeding stage and as such, post-feeding
confidence intervals should be used to infer their age. Furthermore, post-feeding larvae will rapidly
evacuate the crop prior to pupariation, and thus crop dissection can be performed to check for the
presence of food particulates. If no food particulates are found, then it can be assumed that the larvae
are post-feeding. For a more thorough discussion of this technique see Archer et al. [35].

While some other previous studies have included data on the development of *C. augur* [16,20–24], they
were usually conducted as parts of larger projects and provide insufficient detail for inferring an mPMI.
For example, Fuller [16] only reported the average length of the maggot when full grown (18 mm), with
which our findings concur. Mackerras [20] referred only to complete development, with 21-22 d recorded
for growth in summer conditions in an insectary and 18-20 d for development in a room at 20°C. Levot
et al. [23] examined weights of *C. augur* larvae at 27-28°C and found the time to reach maximum larval
weight to be 65.5 hours. While direct comparisons cannot be drawn with this, we observed that the
time to reach maximum larval length at 30°C was 72 h. The techniques of Levot et al. [23] were very
different to ours, with a large mass of larvae feeding on excess liver and each sampling event removing
some individuals. It is possible that the larvae in their experiment were actually exposed to higher
temperatures than the ambient of 27-28°C via massing, and that the feeding mass was better able to
liquefy the substrate, and therefore grow more quickly, than the larvae in the current study.
Comprehensive work by O'Flynn [24,25] examined larval growth at temperatures between 5 and 45°C
but did not produce sufficient data for explicit mPMI predictions. At 5°C, the minimum period from
larviposition to commencement of wandering was 54 d; some larvae survived for 110 d but none
pupated.

Larval temperature preference
It is well known that blowfly larvae show species-specific temperature preferences, which are
behavioural adaptations that maximise developmental rate, growth quality, and survival [36].
Examination of instar size and duration across temperatures in our study indicates that the larvae of *C.*
augur also experience optimal growth at specific temperatures related to their growth stage. In terms
of size, the only temperature at which the average length of third instars dropped below their overall
mean was at 35°C, and post-feeding third instars were largest on average when grown at 20°C.
Regarding instar duration, the shortest developmental times were observed at 35°C for first and second
instars, and at 30°C for third instars – but the most substantial reduction in development time for all

instars was seen between 15 and 20°C. In addition, it appears that the lower developmental threshold
(*t*) for *C. augur* lies between 11 and 20°C (depending on instar). Altogether, this suggests that the
development of *C. augur* is likely optimal around 30°C, but suffers at temperatures above this.
Importantly, the species may be capable of completing development at temperatures as low as 11°C
and can comfortably complete development at 20°C – which is in line with this species remaining
active during the Australian winter.

Environmental conditions and larval growth

Importantly, it is well established that the growth and survival of blowfly larvae is dependent upon
their food source. For example, studies on *Calliphora vicina* and *Calliphora vomitoria* have shown
substantial differences in larval growth between mixed minced meats and beef liver [37] and there are
differences in larval growth between livestock and human tissue[38]. While the measurements we
provide here are based on *C. augur* fed on sheep's liver, these data are assumed to apply to the
equivalent growth in human bodies – an assumption that holds true for comparisons between *C. vicina*
larvae fed on porcine and human tissue[5]. Nevertheless, this assumption remains to be tested with *C.*
augur, and age calculations based on data from animal tissue may lead to inaccurate mPMI estimates
[37,38]. For most forensically important species, including *C. augur*, there remains a pressing need for
broad-scale studies that compare growth rates between a range of tissues from different animal taxa[37,5].

In addition to this, it is important to note that the larvae reared in our developmental experiments were
kept in darkness without a day/night light cycle. It is well known that photoperiod can influence larval
growth rate and development[39,40], and as such, our data may not reflect the developmental rate that
would occur under a 12:12 day night cycle. Nevertheless, very few studies have explored the effect of
complete darkness on insect larval development, so it is unclear to what extent this would influence
developmental rate. In addition, many forensic cases concerning entomology involve corpses found in
secluded places such as abandoned buildings, houses, and basements[41] which are often exposed to very
low levels of light (and in certain cases complete darkness). Future studies of *C. augur* development,
and indeed blowflies more generally, will benefit from investigating how larval development differs
under a range of lighting conditions (including full darkness, artificial light, and natural light).

Conclusions

263 Overall, we provide prediction intervals based on larval length and thermal summation models based
264 on instar duration, at a range of constant temperatures. This comprehensive dataset forms a basis for

265 estimating the age of larvae of *C. augur*, for the first time enabling the application of this species to
266 forensic investigations.
267

Acknowledgements

269 We thank the University of Wollongong for financial assistance with this study.

Disclosure statement

272 The authors have no conflict of interest to declare.
273

Data availability statement

275 All data has been made available as supplementary material.
276
277

References

1. Catts EP & Goff ML (1992) Forensic entomology in criminal investigations. Annual Review
Entomology 37: 253-272.
2. Smith KGV (1986) A manual of forensic entomology. British Museum (Natural History), London
3. Greenberg B (1991) Flies as forensic indicators. Journal of Medical Entomology 28: 565-577.
4. Sharma R, Garg RK, Gaur JR (2015) Various methods for the estimation of the post-mortem interval
from Calliphoridae: A review. Egyptian Journal of Forensic Sciences 5:1-12.
5. Bernhardt V, Schomerus C, Verhoff MA, Amendt J (2017) Of pigs and men – comparing the
development of *Calliphora vicina* (Diptera: Calliphoridae) on human and porcine tissue.
International Journal of Legal Medicine 131:847-853.
6. Harvey ML, Gasz NE, Voss SC (2016) Entomology-based methods for estimation of postmortem
interval. Research and Reports in Forensic Medical Sciences 6:1-9.
7. Wells JD & LaMotte LR (1995) Estimating maggot age from weight using inverse prediction.
Journal of Forensic Sciences 40: 585-590.
8. Ullyett GC (1950) Competition for food and allied phenomena in sheep-blowfly populations.
Philosophical Transactions of the Royal Society London B: Biological Sciences. 234:77-174
9. Grassberger M & Reiter C (2001) Effect of temperature on *Lucilia sericata* (Diptera: Calliphoridae)
development with special reference to the isomegalen- and iosmorphen-diagram. Forensic
Science International 120: 32-36.

10. Jakubec P (2016) Thermal summation model and instar determination of all developmental stages
of necrophagous beetle, Sciodrepoides watsoni (Spence) (Coleoptera: Leiodidae: Cholevinae).
PeerJ 4:e1944

11. Richards CS, Crous KL & Villet MH (2009) Models of development for blowfly sister species
Chrysomya chloropyga and *Chrysomya putoria*. Medical and Veterinary Entomology 23: 56-61.

12. Harvey, ML, Gasz NE, Voss SC (2016) Entomology-based methods for estimation of postmortem
interval. Research and Reports in Forensic Medical Science 6:1-9.

13. Wallman JF & Adams M (1997) Molecular systematics of Australian carrion-breeding blowflies
of the genus *Calliphora* (Diptera: Calliphoridae). Aust Journal of Zoology 45: 337-356.

14. Levot G (2003) Insect fauna used to estimate the post-mortem interval of deceased persons.
General and Applied Entomology 32: 31-39.

15. Meier R, Kotrba M & Ferrar P (1999) Ovoviviparity and viviparity in the Diptera. Biological
Reviews 74: 199-258.

16. Fuller ME (1934) The Insect Inhabitants of Carrion: A Study in Animal Ecology. Bulletin of the
Council for Scientific and Industrial Research 82: 1-63.

17. Norris KR (1959) The ecology of sheep blowflies in Australia. Monographiae Biologicae:
Biogeography and Ecology in Australia (ed. by A Keast, RL Crocker & CS Christian) Junk, The
Hague, The Netherlands, pp. 514-544.

18. Lee DJ (1968) Human myiasis in Australia. Medical Journal of Australia 1: 170-173.

19. Monzu N (1977) Coexistence of Carrion Breeding Calliphoridae (Diptera) in Western Australia.
PhD Thesis, University of Western Australia, Crawley, WA, Australia.

20. Dawson BM, Barton PS, Wallman JF (2020) Contrasting insect activity and decomposition of pigs
and humans in an Australian environment: A preliminary study. Forensic Science International
316:110515

21. Johnston TH & Hardy GH (1923) Observations regarding the life-cycle of certain Australian
blowflies. Proceedings of the Royal Society of Queensland 35: 21-42.

22. Mackerras MJ & Freney MR (1933) Observations on the nutrition of maggots of Australian blow-
flies. Journal of Experimental Biology 10: 237-246.

23. Levot GW, Brown KR & Shipp E (1979) Larval growth of some calliphorid and sarcophagid
Diptera. Bulletin of Entomological Research 69: 469-475.

24. O'Flynn MA (1983) The succession and rate of development of blowflies in carrion in southern
Queensland and the application of these data to forensic entomology. Journal of the Australian
Entomological Society 22: 137-148.

25. O'Flynn MA (1980) Studies on Blowflies and Related Flies from Queensland. PhD Thesis,
University of Queensland, Brisbane, Qld, Australia.

26. Day DM & Wallman JF (2006) Width as an alternative measurement to length for postmortem
interval estimations from *Calliphora augur* (Diptera: Calliphoridae) larvae. Forensic Science
International 159: 158-167.

27. Day DM & Wallman JF (2008) Effect of preservative solutions on preservation of *Calliphora*
augur and *Lucilia cuprina* larvae (Diptera: Calliphoridae) with implications for post-mortem
interval estimates. Forensic Science International 179: 1-10.

28. Geisser S & Johnson W (2006) Modes of Parametric Statistical Inference, Wiley, New York, NY,
USA.

29. Lumley T, Diehr P, Emerson S, Chen L (2002) The importance of the normality assumption in
large public health datasets. Annual Review of Public Health 23:151-169

30. Johnston TH & Tiegs OW (1922) Notes on the biology of some of the more common Queensland
muscoid flies. Proceedings of the Royal Society of Queensland 34: 77-104.

31. Callinan APL (1980) Aspects of the ecology of Calliphora augur (Fabricus) (Diptera:
Calliphoridae), a native Australian blowfly. Australian Journal of Zoology 28:679-684

32. Byrd JH & Butler JF (1998) Effects of temperature on *Sarcophaga haemorrhoidalis* (Diptera:
Sarcophagidae) development. Journal of Medical Entomology 35: 694-698.

33. Rivers DB, Thompson C, Brogan R (2011) Physiological trade-offs of forming maggot masses by
necrophagous flies on vertebrate carrion. Bulletin of Entomological Research 101:599-611.

34. Catts EP (1991) Analyzing entomological data. Entomology and Death: a procedural guide (ed.
by EP Catts & NH Haskell) Joyce's Print Shop Inc, Clemson, SC, USA, pp. 124-137.

35. Archer MS, Jones SD, Wallman JF (2017) Delayed reception of live blowfly (*Calliphora vicina*
and *Chrysomya rufifacies*) larval samples: implications for minimum postmortem interval
estimates. Forensic Sciences Research 3:27-39.

36. Aubernon C, Boulay J, Hédouin V, Charabidzé D (2016) Thermoregulation in gregarious dipteran
larvae: evidence of species-specific temperature selection. Entomologia Experimentalis et
Applicata 160:101-108.

37. Neideregger S, Wartenberg N, Spiess R, Mall G (2013) Influence of food substrates on the
development of the blowflies *Calliphora vicina* and *Calliphora vomitoria* (Diptera,
Calliphoridae). Parasitology Research 112:2847-2853.

38. Wallman JF & Archer MS (2020) The application of insects to the estimation of the time since
death. In J Hayman & M Oxenham, Estimation of the Time Since Death (pp. 57-74). Elsevier,
London, UK

39. Saunders DS (1972) Circadian control of larval growth rate in Sarcophaga argyrostoma.
Proceedings of the National Academy of Science USA 69:2738-2740.

40. Nabity PD, Higley LG, Heng-Moss TM (2007) Light-induced variability in development of
forensically important blow fly Phormia regina (Diptera: Calliphoridae). Journal of Medical
Entomology 44:351-358

41. Michalski M, Nadolski J (2018) Thermal conditions in selected urban and semi-natural habitats,
important for the forensic entomology. Forensic Science International 287:153-162

409 **Table 1. Summary of development constants for C. augur for three developmental stages,**
410 **based on ordinary least squares regression.**
411 Sum of effective temperatures (k) and lower developmental threshold (t) shown as means with
412 standard errors (coefficient of determination (R^2) and degrees of freedom (Df) and p-values are
413 provided).

Stage	Temperature range	R^2	Df	p-value	k ($\pm SE$)	t ($\pm SE$)
1	15−35	0.9045	526	< 0.001	19.5135 ± 2.5992	19.202 ± 0.2717
2	15−35	0.7744	586	< 0.001	361.332 ± 10.889	11.543 ± 0.257
3	15−35	0.7379	2068	< 0.001	1072.928 ± 27.335	13.283 ± 0.174

415 **Table 2. Summary of development constants for C. augur for three developmental stages based**
416 **on the reduced major axis regression.** Sum of effective temperatures (k) and lower developmental
417 threshold (t) shown as means. Coefficient of determination (R^2) and degrees of freedom (Df) and p-
418 values are provided.
419

Stage	Temperature range	R^2	Df	p-value	k	t
1	15−35	0.951	526	< 0.001	14.216	20.188
2	15−35	0.7744	586	< 0.001	303.242	13.114
3	15−35	0.7379	2068	< 0.001	792.078	15.462

Figure 1. Growth curves of *Calliphora augur* larvae at constant temperatures. Data points represent individual larvae at each treatment time. First-instar larvae are represented by the ×, second-instar by the +, and third-instar by the o. The first plot (15°C) has a different scale on the x-axis.

Figure 2. Observed range of instar durations of *Calliphora augur* larvae over five experimental treatments (15, 20, 25, 30, 35°C). The horizontal lines within boxes represent median values. The upper and lower boxes indicate 75th and 25th percentiles.

Figure 3. Bar plots of mean instar duration (hours) of all three instars of *Calliphora augur* across five experimental treatments (15, 20, 25, 30, 35°C).

Figure 4. Ordinary least squares regression for the duration of development of each instar stage of *Calliphora augur*. DT represents the time in hours to reach the stage multiplied by the constant rearing temperature.

Figure 5. Reduced major axis regressions for the duration of development of each instar stage of *Calliphora augur*. DT represents the time in hours to reach the stage multiplied by the constant rearing temperature.