Development of larvae of the Australian blowfly, *Calliphora augur* (Diptera: Calliphoridae), at constant temperatures

Donnah M. Day[1], Nathan J. Butterworth[2*,] Anirudh Tagat[3], Gregory Markowsky[3], and James F. Wallman[1,2]

[1]*Centre for Sustainable Ecosystem Solutions, School of Earth, Atmospheric and Life Sciences, University of Wollongong, NSW 2522, Australia*

[2]*School of Life Sciences, University of Technology Sydney, Ultimo, NSW 2007, Australia*

[3]*School of Mathematics, Monash University, VIC 3800, Australia*

* Corresponding author

E-mail address: nathan.butterworth@uts.edu.au

Keywords: Forensic entomology, post-mortem interval, prediction interval

Abstract

Calliphora augur (Diptera: Calliphoridae) is a common carrion-breeding blowfly of forensic, medical and agricultural importance in eastern Australia. Despite this, detailed information on the developmental biology of *C. augur* is lacking. Here, we present the first comprehensive study on the development of all three larval instars of *C. augur,* fed on sheep's liver, at constant temperatures of 15, 20, 25, 30 and 35°C. We highlight decreasing variation in final larval length at sampling periods up to 30°C, although this variation increased again when larvae were grown at 35°C. 95% prediction intervals are provided for each constant temperature, enabling the age of larvae of *C. augur* to be estimated from their average length. These data will assist with the application of this species to forensic investigations.

Introduction

Forensic entomology is the study of insects as applied to legal issues. It may relate to medicolegal events, especially murders, for the purpose of uncovering information useful to an investigation (Catts & Goff, 1992). This is because decomposing vertebrate remains provide a temporary micro-habitat and food source that is attractive to a variety of insects and other invertebrates (Smith, 1986). Arthropods constitute a major element of this fauna, with insects, especially flies and their larvae, being the predominant group. Blowflies (Diptera: Calliphoridae) are ubiquitous and are typically the first insects to visit a dead body, often well before it is discovered by humans (Greenberg, 1991).

Data on the development of blowflies and other important carrion flies can help provide an estimate of the minimum time since death (or minimum post-mortem interval: mPMI), especially when combined with information on the succession patterns of insects on corpses (Sharma et al., 2015). The traditional approach adopted in developmental studies has been to generate growth data based on larval size or weight, and then use tables (Kamal, 1958; Nishida et al., 1986) or plots incorporating various models (Wells & LaMotte, 1995; Byrd & Butler, 1997; Grassberger & Reiter 2001; Donovan et al., 2006; Richards et al., 2009) to estimate the age of larvae collected from a corpse. This information would then be used to derive the time of egg or larval deposition and thus an estimate of the mPMI. However, many studies have neglected to provide mathematically explicit mPMI predictions, meaning that reliable predictions are still lacking for many calliphorid species. It is now generally suggested that studies of larval development provide mPMI predictions with a range of values rather than a single value, and that this range should be defined as a confidence set (Wells & LaMotte 2017).

Calliphora augur Fabricius is a common blowfly species found in eastern Australia (Wallman & Adams, 1997), the larvae of which are among those most commonly collected from forensic cases, including from inside buildings (Levot, 2003). It is ovoviviparous (Meier et al., 1999), and also an agent of myiasis in sheep and other animals (Fuller, 1932; Norris, 1959; Lee, 1968). While it has a preference for fresh carcasses (Monzu, 1977), it will also larviposit on putrid liver (Mackerras, 1933). While previous studies have considered the larval development of *C. augur* (Johnston & Hardy, 1923; Fuller, 1932; Mackerras, 1933; Mackerras & Freney, 1933; Levot et al., 1979; O'Flynn, 1983) none have made fine scale measurements across a range of constant temperatures, or have attempted to provide mathematically explicit mPMI predictions. This lack of accurate developmental data means that at present, *C. augur* cannot be used reliably in a forensic context. To ameliorate this, we present the first comprehensive analysis of larval growth at constant temperatures of this important species.

Our investigations are based on measurement of larval length throughout development, including observations on size overlap between larval instars. Most importantly, we calculate prediction intervals that can be used to mathematically infer larval age from size.

Materials and methods

Fly culture

A culture of *C. augur* was established from adults trapped at Wollongong, New South Wales (34° 25' S, 150°53' E). Experiments commenced using larvae of the F3 generation and were continued for 11 generations. The culture was refreshed regularly with wild-type (F_0) individuals from the same source population as above and maintained with mixed ages of females and males. The average fecundity of each female was estimated by separating 20 females and transferring them to 50 mL specimen pots containing approximately 10 g of sheep's liver. Larvae were counted the following day. Larvae and adults were kept in a temperature-controlled room at 25±3.5˚C and ambient humidity, with a 12:12 light:dark regime incorporating a 15 minute 'dusk' and 'dawn' transition period of low light. Adult flies were maintained in square plastic cages measuring 330 mm long, 220 mm wide and 250 mm high (external dimensions). The flies in each cage were provided with water and sugar *ad libitum*, and chopped sheep's liver in a small plastic weigh boat for larviposition as required. Larvae were reared in square white 2 L plastic containers, measuring 170 mm wide along each side and 90 mm deep. Most of the centre of the lid of each container was cut away and replaced with fine mesh to permit ventilation. In each container, larvae were sustained on sheep's liver in the weigh boats placed atop a layer of wheaten chaff approximately 20 mm deep to enable pupariation. After pupariation, pupae were manually sorted from the chaff and transferred to a cage. Sheep's liver was provided just prior to eclosion to provide females with a constant source of protein for ovary maturation (Mackerras, 1933).

Development at constant temperatures

Each replicate consisted of freshly laid larvae randomly collected from the main culture. Preliminary studies on larval density indicated that some endogenous heat generation occurred with as few as 25 larvae growing on 50 g of sheep's liver. To ensure that the larvae developed at the ambient temperature being examined, each replicate (a plastic weigh boat containing larvae and meat) was limited to 10 larvae. These replicates were grown at specific constant temperatures of 15, 20, 25, 30 and 35˚C, and 60±5% humidity, in an Axyos (Brisbane, Queensland, Australia) environmental cabinet. Since only one cabinet was available, growth at each temperature was studied in turn. Larvae were left

undisturbed in darkness in the environmental cabinet until collection, because disturbance can delay

pupariation (Mackerras, 1933; O'Flynn, 1983). To account for growth rates increasing with

temperature, we sampled more frequently at higher temperatures.

The growth period of each replicate of 10 larvae at a particular temperature was chosen by lottery. For

the first three to six days, six-hourly sampling per sample was done at 15°C, while four-hourly

sampling was conducted at 20, 25, 30 and 35°C. Six replicates were used for the first sample period,

and every 48 hours thereafter, and three replicates were used for every other time interval (either 12-

or 24-hourly, depending on temperature). The choice of these specific sampling frequencies was

guided by the growth curves of O'Flynn (1980; 1983), and frequencies were kept relatively high to

minimise error in the calculation of the prediction intervals.

Sample collection, handling and preservation

At collection, all larvae were killed and fixed immediately by immersion in boiling water, dried with

paper towel and preserved in 80% EtOH. Larvae were placed into glass vials posteriad to prevent

head-curling (Day & Wallman, 2006) and length and instar were recorded.

Measurement and statistics

The body lengths of larvae were measured with the aid of a dissecting microscope and Mitutoyo

Absolute digimatic digital callipers (Kawasaki, Japan) after the larvae had been in preservative for a

minimum of ten days (Day & Wallman, 2008). Body length was measured as the distance, viewed

laterally, between the most distal parts of the head and the last abdominal segment. The lengths of

pupae were also measured with callipers. The ambient temperature within the temperature-controlled

cabinet was monitored with data loggers (iButtons: Maxim Integrated Products Inc., Sunnyvale, USA).

The dataset consists of the measurements of the lengths of larvae grown at temperatures of 15°C, 20°C,

25°C, 30°C, and 35°C. Since no larvae were measured more than once, the length data for the various

time intervals should be regarded as independent. As such, the variable of interest for this study was

the variation in growth rate, as measured by length, at the different temperature treatments.

The instar of each larva was also recorded when it was measured as either 1st, 2nd, or 3rd. As the

dataset is rather large (more than 3000 datapoints), presenting it in its entirety would be impractical,

and we therefore display the data in scatterplots for each of the temperatures (the small number of

intermediate instars were removed from these graphs in order to aid readability). Note that the intervals

differ for the various temperatures; the intervals were chosen to accurately capture the growth of the

larvae, with the higher temperatures naturally necessitating more frequent sampling: the larvae at 15°C

were measured every 24 hours, those at 20°C every 6 hours, and the rest every 4 hours.

Results

The average numbers of eggs laid were 58±14 per gravid female. First-instar larvae were largest on

average when grown at 25°C. Similarly, second instars were also largest when grown at 25°C, but

they also displayed a much smaller variation at this temperature compared with the other temperatures

examined. Third-instar larvae were largest on average when grown at 20°C, but were above the overall

mean (for third-instars) when growing at 15, 20, 25 and 30°C. The only temperature at which the

average length of third instars dropped below their overall mean was at 35°C. Post-feeding third instars

were largest on average when grown at 20°C.

Utilizing the dataset

Our dataset can provide potentially valuable information to an investigator by simply comparing the

ratios of the various instars found at a crime scene with those obtained in our experiment. To this end,

we present graphs showing the association between size and time for each of the temperatures

measured (Figure 1).

For instance, supposing that a batch of larvae is recovered from a body from a location where the

average temperature has been 15°C, and these larvae are roughly evenly split between instars 2 and 3,

then consulting Figure 1 would give a range of the mPMI between 72 and 120 hours, with 96 hours

being the best estimate. On the other hand, if only larvae in instar 1 are recovered from a body at an

average temperature of 35°C, then the mPMI is confidently less than 12 hours, and so forth. This should

help to establish the mPMI in certain circumstances, in particular when the percentages of the various

instars are specific to a relatively small time interval, but is clearly not good enough in other instances,

most notably those corresponding to longer mPMIs in which one is likely to find only larvae in instar

3. These cases require a more refined analysis, which we now provide.

It is assumed that an investigator is able to procure at least five larvae from a crime scene and is able

to correctly record their average length, as we refer to in the 'measurement and statistics' section. We

will provide 95% prediction intervals for the average length of the larvae for each combination of time,

instar, and temperature; this is analogous to but different than giving confidence intervals, since we

are predicting the mean value of a sample rather than the population mean. A detailed explanation of

the process can be found in Geisser and Johnson (2006), but for the benefit of the reader we briefly

review the theory behind the method. Those unfamiliar with some of the standard distributions and

facts from statistics theory used below are advised to consult almost any appropriate statistics textbook

or online resource.

Suppose that, for a given time, instar, and temperature, the length of a maggot is normally distributed

with mean μ and variance σ^2. We regard our experimental data as a sample from this population of

size N, and wish to test the hypothesis that a sample procured from a crime scene by an investigator

was taken from the same population. The crime scene sample is regarded as independent from the

experimental sample, and let m be the number of larvae taken from the time scene. Let us label the

sample mean from our experiment as $\bar{x}_{\exp}$ and from the crime scene as $\bar{x}_{\mathrm{cri}}$.

As is well known, $\bar{x}_{\exp}$ and $\bar{x}_{\mathrm{cri}}$ will be normally distributed random variables, each with mean μ but

with variances σ^2/N and σ^2/m respectively. Since $\bar{x}_{\exp}$ and $\bar{x}_{\mathrm{cri}}$ are independent random variables, their

difference $\bar{x}_{\exp}$ - $\bar{x}_{\mathrm{cri}}$ will also be normally distributed, but now with mean 0 and variance $\sigma^2/N + \sigma^2/m$.

It follows that $Z = (\bar{x}_{\exp}$ - $\bar{x}_{\mathrm{cri}})/\sqrt{\sigma^2(\frac{1}{N} + \frac{1}{m})}$ is a standard normal (mean 0, variance 1). If we knew the

population variance σ^2, we could use a standard normal table to create 95% prediction intervals, but

since we do not (as is typical) we will use the sample variance s^2 from our experiment in its place.

Now, $\frac{(N-1)s^2}{\sigma^2}$ will have a $\chi^2(N\text{-}1)$ distribution, and will be independent of Z (it is independent of $\bar{x}_{\mathrm{cri}}$ by

assumption, and of $\bar{x}_{\exp}$ by the standard independence of sample mean and variance for normal random

variables). Thus, $\frac{Z}{\sqrt{s^2/\sigma^2}}$ will follow a $t_{n\text{-}1}$ distribution. Replacing Z by the expression that defined it and

cancelling the σ^2 in the numerator and denominator, we see that $(\bar{x}_{\exp}$ - $\bar{x}_{\mathrm{cri}})/\sqrt{s^2(\frac{1}{N} + \frac{1}{m})} \sim t_{n\text{-}1}$. Our

95% prediction interval for $\bar{x}_{\mathrm{cri}}$ is therefore $\bar{x}_{\exp} \pm t_{.025,N\text{-}1} \sqrt{s^2(\frac{1}{N} + \frac{1}{m})}$.

A few comments on the normality assumption made above are in order. It is commonly accepted that

measurements such as the size of organisms are generally modelled well by normal distributions;

indeed, we ran the standard Kolmogorov-Smirnov and Shapiro-Wilk normality tests on the data, and

found that most of the data subsets for each time, instar, and temperature are consistent with an

underlying normal distribution, with a small number of exceptions. Furthermore, even in the case of

the exceptions, the *t*-test is known to be highly robust against non-normality, for the reason that even

when the population does not follow a normal distribution the sample means $\bar{x}_{\text{exp}}$ and $\bar{x}_{\text{cri}}$ will still be

approximately normally distributed (Lumley et al 2002). This assures us that the method described

above is general and not dependent upon the normality assumption.

We present the prediction intervals in table form below (Table 1), with a separate section for each

combination of temperature and instar. Note that we have not provided tables for the intermediate

stages (first- to second-instar, or second- to third-instar) since we feel that they do not appear in

sufficient numbers to provide valuable prediction intervals; we have also removed from Table 1 the

times at which there fewer than five larvae of a particular instar present, for the same reason. The

prediction intervals are given in the form (*a,b*), which is an equivalent notation to $a \leq \bar{x}_{\text{cri}} \leq b$. An

investigator should make use of these tables by calculating the mean length of their sample of larvae,

and then comparing this with the prediction intervals given in the table corresponding to the correct

instar and average temperature; if the mean of the sample lies within the prediction interval for a given

time, then that time should be regarded as a possible age for the larvae. We have provided prediction

intervals for crime scene sample sizes *m = 5* and *m = 30*; the *m* = 5 interval should be used for sample

sizes between 5 and 29, and the *m = 30* interval for any *m* over 30.

Figure 1. Growth curves of *Calliphora augur* larvae at constant temperatures. Data points represent individual larvae at each treatment time. First-instar larvae are represented by the red ×, second-instar by the blue +, and third-instar by the green o. The first plot (15°C) has a different scale on the x-axis.

Table 1. The 95% prediction intervals based on the developmental curves of *Calliphora augur* larvae at five different constant temperatures (15, 20, 25, 30, 35°C). n = sample number, $\bar{x}_{exp}$ = sample mean, s^2 = sample variance. $m = 5$ interval is the prediction interval to be used for sample sizes between 5 and 29 larvae, $m = 30$ interval is the prediction interval to be used for sample sizes larger than 30 larvae.

Time (hours)	N	$\bar{x}_{exp}$	s^2	$m = 5$ interval	$m = 30$ interval
Instar 1, Temp 15 ºC					
0	59	2.1725	0.072	(1.989,2.356)	(2.052,2.293)
24	25	4.0148	0.240	(3.636,4.393)	(3.741,4.289)
Instar 2, Temp 15 ºC					

48	65	5.4798	0.530	(4.986,5.974)	(5.159,5.801)
72	26	7.3438	0.782	(6.666,8.022)	(6.856,7.832)
96	21	8.2752	1.286	(7.366,9.184)	(7.602,8.948)

Instar 3, Temp 15 ºC

96	21	11.7024	0.632	(11.065,12.340)	(11.231,12.174)
120	17	11.9724	4.935	(10.096,13.849)	(10.543,13.402)
144	51	14.3841	10.767	(12.105,16.663)	(12.868,15.901)
168	25	17.3384	5.108	(15.593,19.084)	(16.075,18.602)
192	42	16.7124	13.344	(14.117,19.308)	(14.949,18.476)
216	19	17.7626	1.108	(16.899,18.627)	(17.114,18.411)
240	43	17.9709	3.062	(16.731,19.211)	(17.131,18.811)
264	20	17.149	2.339	(15.909,18.389)	(16.225,18.073)
288	31	17.0787	1.668	(16.119,18.038)	(16.403,17.754)
312	32	16.5694	1.934	(15.542,17.597)	(15.849,17.290)
336	29	16.0203	2.029	(14.950,17.090)	(15.260,16.780)
360	8	16.5700	0.607	(15.696,17.444)	(15.837,17.303)
384	41	14.7044	4.616	(13.173,16.236)	(13.661,15.748)
408	25	14.8204	4.069	(13.263,16.378)	(13.693,15.948)

Instar 1, Temp 20 ºC

0	45	2.5538	0.164	(2.268,2.839)	(2.361,2.746)
6	11	3.3409	0.104	(3.027,3.655)	(3.088,3.594)
12	13	4.2562	0.135	(3.919,4.593)	(3.990,4.522)

Instar 2, Temp 20 ºC

24	18	5.1522	0.519	(4.553,5.752)	(4.699,5.605)
30	13	5.5415	0.538	(4.869,6.214)	(5.011,6.072)
36	20	7.0295	0.865	(6.276,7.783)	(6.468,7.591)
42	19	7.1032	0.404	(6.581,7.625)	(6.712,7.495)
48	22	9.0773	0.772	(8.380,9.774)	(8.564,9.590)
66	10	8.5380	0.643	(7.727,9.349)	(7.876,9.200)

Instar 3, Temp 20 ºC

54	18	14.4444	3.65	(12.855,16.034)	(13.243,15.646)
60	15	11.6647	1.906	(10.456,12.874)	(10.728,12.601)
66	14	14.8379	10.474	(11.943,17.733)	(12.575,17.101)
72	27	14.6637	3.818	(13.177,16.151)	(13.598,15.729)
84	38	15.6639	3.861	(14.249,17.079)	(14.692,16.636)
96	64	17.7798	2.149	(16.784,18.776)	(17.132,18.428)
108	24	18.245	1.689	(17.233,19.257)	(17.509,18.981)
120	18	18.1456	2.534	(16.821,19.470)	(17.144,19.147)
132	27	18.6196	1.475	(17.695,19.544)	(17.957,19.282)
144	44	18.2527	2.302	(17.181,19.325)	(17.528,18.977)

156	19	18.3826	2.562	(17.069,19.696)	(17.397,19.369)
168	18	17.2128	2.393	(15.926,18.500)	(16.240,18.186)
180	16	18.6719	0.627	(17.992,19.352)	(18.149,19.194)
192	54	18.1091	1.673	(17.216,19.002)	(17.518,18.700)
204	16	16.8313	3.503	(15.223,18.439)	(15.596,18.066)
216	14	18.0257	2.066	(16.740,19.311)	(17.021,19.031)

Instar 1, Temp 25 ºC

0	48	2.9033	0.121	(2.660,3.147)	(2.740,3.066)
4	27	3.6081	0.037	(3.462,3.754)	(3.503,3.713)
8	11	3.81	0.048	(3.597,4.023)	(3.638,3.982)
12	30	4.268	0.214	(3.923,4.613)	(4.024,4.512)
16	6	4.2917	0.178	(3.732,4.852)	(3.807,4.777)

Instar 2, Temp 25 ºC

20	23	7.143	0.291	(6.719,7.567)	(6.833,7.453)
24	28	7.31	0.352	(6.862,7.758)	(6.990,7.630)
28	21	7.5662	0.864	(6.821,8.311)	(7.015,8.118)
32	18	7.9783	0.203	(7.603,8.353)	(7.695,8.262)
36	16	8.275	0.712	(7.550,9.000)	(7.718,8.832)
40	16	8.6106	0.564	(7.965,9.256)	(8.115,9.106)
44	17	8.7971	0.609	(8.138,9.456)	(8.295,9.299)

Instar 3, Temp 25 ºC

48	34	13.6715	2.144	(12.600,14.743)	(12.925,14.418)
52	19	12.1405	2.881	(10.747,13.534)	(11.095,13.186)
56	29	15.8128	2.342	(14.663,16.962)	(14.996,16.629)
60	27	16.0007	4.955	(14.307,17.695)	(14.787,17.214)
64	27	14.7056	5.339	(12.947,16.464)	(13.446,15.966)
72	21	16.5886	5.495	(14.710,18.467)	(15.197,17.980)
84	24	17.3142	5.99	(15.409,19.220)	(15.928,18.701)
96	39	17.7841	2.745	(16.595,18.973)	(16.970,18.599)
108	24	18.8704	3.099	(17.500,20.241)	(17.873,19.868)
120	27	17.2167	1.342	(16.335,18.098)	(16.585,17.848)
132	28	17.1954	1.806	(16.180,18.211)	(16.471,17.920)
144	45	17.0431	2.249	(15.986,18.100)	(16.331,17.755)
156	33	16.5845	2.917	(15.329,17.840)	(15.707,17.462)
168	21	16.7257	0.818	(16.001,17.451)	(16.189,17.262)
180	7	16.8843	0.592	(15.956,17.812)	(16.094,17.675)
192	6	15.6000	2.257	(13.606,17.594)	(13.873,17.327)

Instar 1, Temp 30 ºC

0	60	2.5583	0.104	(2.338,2.779)	(2.414,2.703)
4	28	3.5518	0.153	(3.256,3.847)	(3.341,3.763)

8	29	3.9228	0.168	(3.615,4.231)	(3.704,4.141)
Instar 2, Temp 30 ºC					
12	14	5.6279	0.278	(5.156,6.100)	(5.259,5.997)
16	29	7.0290	0.106	(6.784,7.274)	(6.855,7.203)
20	21	7.6262	0.946	(6.847,8.406)	(7.049,8.203)
24	13	9.0131	0.288	(8.521,9.505)	(8.625,9.401)
Instar 3, Temp 30 ºC					
28	13	10.5392	0.349	(9.998,11.081)	(10.112,10.967)
32	23	11.2465	1.399	(10.317,12.176)	(10.567,11.926)
36	25	13.1956	2.810	(11.901,14.490)	(12.259,14.132)
40	25	15.5992	2.760	(14.316,16.882)	(14.671,16.528)
44	30	16.1283	2.987	(14.838,17.419)	(15.216,17.041)
48	38	16.0895	3.748	(14.695,17.484)	(15.131,17.048)
52	27	16.4419	2.032	(15.357,17.527)	(15.665,17.219)
54	6	16.4567	0.141	(15.958,16.955)	(16.025,16.888)
56	25	17.7292	2.825	(16.431,19.027)	(16.790,18.669)
60	28	18.4364	0.625	(17.839,19.034)	(18.010,18.863)
64	34	17.4650	2.535	(16.300,18.630)	(16.654,18.276)
68	28	18.0279	1.991	(16.961,19.094)	(17.267,18.789)
72	23	18.0283	0.908	(17.280,18.777)	(17.481,18.576)
84	28	17.4275	1.600	(16.471,18.384)	(16.746,18.109)
96	53	16.3130	4.688	(14.815,17.811)	(15.320,17.306)
120	22	15.7395	1.246	(14.854,16.625)	(15.088,16.391)
132	12	15.7417	4.745	(13.689,17.795)	(14.104,17.379)
144	28	15.6121	3.940	(14.112,17.112)	(14.542,16.682)
156	10	15.3620	2.889	(13.642,17.082)	(13.958,16.766)
168	9	15.5289	0.800	(14.581,16.477)	(14.745,16.313)
Instar 1, Temp 35 ºC					
0	63	2.2735	0.082	(2.079,2.468)	(2.147,2.400)
4	16	3.3800	0.074	(3.146,3.614)	(3.201,3.559)
8	19	3.8411	0.129	(3.546,4.136)	(3.62,4.062)
Instar 2, Temp 35 ºC					
12	18	6.0122	0.462	(5.447,6.578)	(5.585,6.440)
16	27	6.8485	0.261	(6.460,7.237)	(6.570,7.127)
20	21	8.3519	0.920	(7.583,9.121)	(7.783,8.921)
24	8	6.8000	2.611	(4.988,8.612)	(5.280,8.320)
Instar 3, Temp 35 ºC					
28	5	10.4360	1.655	(8.480,12.392)	(8.711,12.161)
32	25	13.0576	3.477	(11.618,14.498)	(12.015,14.100)
36	20	11.9110	9.849	(9.367,14.455)	(10.015,13.807)

44	19	14.5695	8.228	(12.215,16.924)	(12.803,16.336)
48	27	15.2704	5.556	(13.477,17.064)	(13.985,16.556)
52	15	10.0413	9.612	(7.327,12.756)	(7.939,12.144)
56	20	16.1190	6.689	(14.022,18.216)	(14.556,17.682)
60	17	13.4012	22.225	(9.418,17.384)	(10.367,16.435)
68	6	15.8800	11.792	(11.322,20.438)	(11.932,19.828)
72	22	16.7327	3.414	(15.267,18.198)	(15.654,17.811)
84	14	15.7243	5.512	(13.624,17.824)	(14.083,17.366)
96	28	16.5311	0.826	(15.844,17.218)	(16.041,17.021)
108	14	15.7743	5.267	(13.721,17.827)	(14.170,17.379)
120	10	15.2280	1.652	(13.928,16.528)	(14.166,16.290)
132	13	13.9485	1.780	(12.726,15.171)	(12.983,14.914)
144	11	12.2182	5.073	(10.025,14.411)	(10.449,13.987)
156	13	12.9208	1.246	(11.898,13.944)	(12.113,13.728)
168	6	12.6067	2.463	(10.523,14.69)	(10.803,14.411)
180	9	13.4256	3.840	(11.349,15.502)	(11.708,15.143)
192	7	11.9214	5.784	(9.021,14.821)	(9.451,14.392)

272 The investigator may also be interested in forming prediction intervals for time, temperature, and instar

273 combinations that do not appear above. The table below (Table 2) may be of some use if the

274 combination appears there, but there are still some combinations (for example, temperature 35°, instar

275 3, time 76 hours) for which we have no data. As a simple rule of thumb, in this case, if the nearest

276 times for which there are data (here 72 and 84 hours) include $\bar{x}_{\text{cri}}$ then the time in question should be

277 regarded as possible. If the investigator requires a precise prediction interval then we recommend

278 interpolating the means and variances linearly from the nearest points in time to estimate the mean and

279 variance at the point in question, and taking the degrees of freedom to be equal to the minimum of the

280 degrees of freedom at the nearest points (this is a slight simplification of the method used in Wells &

281 Lamotte, 1995). For example, in our case of temperature 35°C, instar 3, time 76 hours, we look at the

282 closest time points for the given temperature and instar, which are at 72 and 84 hours. The means are

283 at 16.733 and 15.724, respectively, and the variances are 3.414 and 5.512. We therefore estimate the

284 mean and variance at 76 hours to be $\left(\frac{2}{3}\right) 16.733 + \left(\frac{1}{3}\right) 15.724 = 16.397$ and $\left(\frac{2}{3}\right) 3.414 +$

285 $\left(\frac{1}{3}\right) 5.512 = 4.113$; note that the values at 72 hours are weighted double those at 84 hours, because 76

286 is twice as far from 84 as from 72. There are 22 measurements at 72 hours and 14 measurements at 84

287 hours, so we take as our degrees of freedom *min* (22-1,14-1) = 13. We therefore obtain prediction

288 intervals by the same method as before of (14.497,18.297) for $m = 5$ and (14.979,17.815) for $m = 30$.

For completeness, Table 2 provides the sample means and variances (when they exist) for the cases
for which we had between one and five measurements (we have not formulated prediction intervals in
these cases). Any remaining time, temperature, and instar combinations that do not appear in either the
above or below tables are ones for which we have no measurements, and for these the interpolation
method described above should be used to form prediction intervals, if necessary.

Table 2. The sample means and variances of the growth of *Calliphora augur* larvae for cases which
included between one and five measurements. Dots represent variances that could not be calculated
due to there being only one sample.

Instar	Temp (ºC)	Time	n	$\bar{x}_{exp}$	s^2
1	15	48	1	3.910	.
2	-	120	1	7.020	.
3	-	432	4	14.203	3.890
-	-	456	2	13.340	0.168
-	-	480	4	15.385	5.512
-	-	504	1	17.680	.
-	-	576	1	13.400	.
-	-	600	1	15.230	.
1	20	18	2	4.395	0.026
-	-	30	1	4.350	.
2	-	54	3	8.707	0.584
3	-	48	1	11.570	.
1	25	52	1	6.250	.
2	-	16	3	5.093	0.099
-	-	52	3	7.490	0.381
3	-	36	3	10.150	0.373
-	-	40	2	10.955	2.856
-	-	44	3	10.940	0.125
-	-	216	4	16.050	3.518
1	30	12	1	2.960	.
2	-	28	1	9.100	.
3	-	180	1	15.030	.
-	-	192	4	13.663	5.704
-	-	204	4	15.003	0.241
2	35	28	1	6.61 0	.
3	-	24	4	11.023	0.723
-	-	204	2	13.760	0.003

Discussion

This study provides the first detailed dataset of the development of the Australian carrion-breeding blowfly *Calliphora augur*. The results we provide here reinforce the previous work done on this species but go substantially further by providing a level of statistical confidence in the application of developmental data in forensic investigations.

Fecundity

Robust and highly active first-instar larvae of *C. augur* hatch promptly after deposition, giving this species an ecological advantage over oviparous species for exploiting small, quickly perishable carcasses (Norris, 1959). Johnston & Tiegs (1922) reported that *C. augur* is capable of depositing eggs or larvae, with the eggs usually hatching within six hours. In the present study, we noted that *C. augur* will normally deposit living larvae but will also lay soft infertile eggs for some days before normal larviposition occurs – similar to the observations of Mackerras (1933) and O'Flynn (1980). The average number of larvae per female has previously been noted as 50 (Mackerras, 1933) or 58 larvae (Norris, 1959). Callinan (1980) recorded a range of between 22.5 and 98.3 larvae per female in a controlled environment and a range of between 0.3 and 45.2 larvae per female in a field environment. The average numbers of eggs laid in the present study were 58±14 per gravid female, which is consistent with these previous data.

Development at constant temperatures

The outcomes of some other developmental studies have been limited by a less thorough approach than taken here. For example, Byrd and Butler (1998) noted that their sampling method selected for the fastest growing larvae, which often indicates a shorter period than could otherwise be deduced from normal collection techniques at a crime scene. The influence of size extremes upon variance has also been considered by Wells & LaMotte (1995). In the present study, we included larvae of all sizes to demonstrate the full range that may be encountered in crime scene samples. Variation in larval body length appeared to decrease steadily from 15 to 30°C and then increased again dramatically at 35°C. The decrease in variation with increasing temperature may relate to a more even growth rate at higher temperatures. However, the subsequent increase in variation at 35°C likely reflects the fact that the larvae are reaching their thermal limit, at which point some larvae may begin to suffer from increased heat and oxidative stress, hindering their growth and increasing the perceived variation (Rivers et al. 2011). The variation in overlap of the growth stages, particularly the transitional forms between instars, emphasises the importance of assessing crime scene samples for instar and/or transitional forms, and

collecting an adequate number of specimens. While there has been comment on the value of transitional forms (Catts, 1991), few published papers include this information and some workers disregard their influence on size ranges (e.g. Wells & LaMotte, 1995).

The reader will note an interesting feature of the plots, namely that the measured lengths increase linearly with time to a certain point, but then begin to decrease after that. This late-stage reduction in larval length is a result of post-feeding shrinkage that calliphorid larvae undergo after leaving the carcass, prior to pupariation. This post-feeding shrinkage can cause some difficulty in interpreting the age of larvae, because the smaller length of post-feeding larvae makes them appear younger. However, this can be solved if investigators record where larvae are collected, as post-feeding larvae will depart the carcass and move into the surrounding substrate. If larvae are collected away from the carcass, it can be presumed that they are in the post-feeding stage and as such, post-feeding confidence intervals should be used to infer their age. Furthermore, post-feeding larvae will rapidly evacuate the crop prior to pupariation, and thus crop dissection can be performed to check for the presence of food particulates. If no food particulates are found, then it can be assumed that the larvae are post-feeding. For a more thorough discussion of this technique see Archer et al. (2017).

While some other previous studies have included data on the development of *C. augur* (Johnston & Hardy, 1923; Fuller, 1932; Mackerras, 1933; Mackerras & Freney, 1933; Levot et al., 1979; O'Flynn, 1983), they were usually conducted as parts of larger projects and provide insufficient detail for inferring an mPMI. For example, Fuller (1932) only reported the average length of the maggot when full grown (18 mm), with which our findings concur. Mackerras (1933) referred only to complete development, with 21-22 d recorded for growth in summer conditions in an insectary and 18-20 d for development in a room at 20°C. Levot et al. (1979) examined weights of *C. augur* larvae at 27-28°C and found the time to reach maximum larval weight to be 65.5 hours. While direct comparisons cannot be drawn with this, we observed that the time to reach maximum larval length at 30°C was 72 h. The techniques of Levot et al. (1979) were very different to ours, with a large mass of larvae feeding on excess liver and each sampling event removing some individuals. It is possible that the larvae in their experiment were actually exposed to higher temperatures than the ambient of 27-28°C via massing, and that the feeding mass was better able to liquefy the substrate, and therefore grow more quickly, than the larvae in the current study. Comprehensive work by O'Flynn (1980; 1983) examined larval growth at temperatures between 5 and 45°C but did not produce sufficient data for explicit mPMI predictions. At 5°C, the minimum period from larviposition to commencement of wandering was 54 d; some larvae survived for 110 d but none pupated.

Larval temperature preference

Archer and Elgar (2003) studied blowfly larval feeding strategies in carrion and found that site preference changes with the length of time the carcass was exposed to the elements. Examination of instar size and temperature in our study (Figure 1) indicates that larvae also appear to have temperature preferences that relate to their growth stage. The only temperature at which the average length of third instars dropped below their overall mean was at 35°C. This drop appears to be further evidence that this is around the upper thermal limit for this species. Post-feeding third instars were largest on average when grown at 20°C. These results indicate that the influence of temperature on larval growth is more complex than it appears. Detailed studies are needed to investigate movement and temperature preferences of individual larvae growing in a mass.

Substrate and larval growth

Importantly, it is well established that the growth and survival of blowfly larvae is dependent upon their food source. For example, studies on *Calliphora vicina* and *Calliphora vomitoria* have shown substantial differences in larval growth between mixed minced meats and beef liver (Neideregger et al. 2013) and there are almost certainly differences in larval growth between livestock and human tissue (Wallman and Archer 2020). While the prediction intervals we provide are based on *C. augur* fed on sheep's liver, these data are assumed to apply to the equivalent growth in human bodies – an assumption that holds true for comparisons between *C. vicina* larvae fed on porcine and human tissue (Bernhardt et al. 2017). Nevertheless, this assumption may not hold true for all *Calliphora* species, and as such there remains a pressing need for studies that compare the growth rates and fitness of blowflies reared on tissues from different animals.

Conclusions

Other workers in forensic entomology have explored various models to infer larval age, including techniques based on weight (Wells & LaMotte, 1995; von Zuben et al., 1998). Such approaches have obvious limitations, as it is difficult to control for the moisture content of the larvae after their collection. Previous studies based on larval size have predominantly used plots based upon models involving isomorphen or isomegalen diagrams (Grassberger & Reiter, 2001) or thermal summation (ADH) (Donovan et al., 2006; Higley & Haskell, 2010). Unlike many former approaches, the technique we outline here allows a clear statement of the statistical rigour underlying a maggot age estimate, while avoiding difficulties inherent in thermal summation, especially the reliance on the linear portion of the development curve and the estimation of the minimum developmental threshold (Nabity et al., 2006). Although inferences of larval age from our data must be based on the specific temperatures

402 used in our study, the predicted average developmental temperature in forensic cases will, nonetheless,

403 often be no more precise than the nearest 5°C because of uncertainty about the amount of heat

404 generated by the feeding larvae. Overall, the 95% prediction intervals we provide for each constant

405 temperature, enable the age of larvae of *C. augur* to be estimated from their average length, for the

406 first time enabling the reliable application of this species to forensic investigations.

Acknowledgements

409 We thank the University of Wollongong for financial assistance with this study.

References

Archer MS, Jones SD, Wallman JF (2017) Delayed reception of live blowfly (*Calliphora vicina* and

Chrysomya rufifacies) larval samples: implications for minimum postmortem interval estimates.

Forensic Sciences Research 3:27-39.

Bernhardt V, Schomerus C, Verhoff MA, Amendt J (2017) Of pigs and men – comparing the

development of *Calliphora vicina* (Diptera: Calliphoridae) on human and porcine tissue.

International Journal of Legal Medicine 131:847-853.

Bornemissza GF (1957) An analysis of arthropod succession in carrion and the effect of its

decomposition on the soil fauna. Australian of Journal Zoology 5: 1-12.

Byrd JH & Butler JF (1997) Effects of temperature on *Chrysomya rufifacies* (Diptera: Calliphoridae)

development. Journal of Medical Entomology 34: 353-358.

Byrd JH & Butler JF (1998) Effects of temperature on *Sarcophaga haemorrhoidalis* (Diptera:

Sarcophagidae) development. Journal of Medical Entomology 35: 694-698.

Callinan APL (1980) Aspects of the ecology of Calliphora augur (Fabricus) (Diptera: Calliphoridae),

a native Australian blowfly. Australian Journal of Zoology 28:679-684

Catts EP & Goff ML (1992) Forensic entomology in criminal investigations. Annual Review

Entomology 37: 253-272.

Catts EP (1991) Analyzing entomological data. Entomology and Death: a procedural guide (ed. by

EP Catts & NH Haskell) Joyce's Print Shop Inc, Clemson, SC, USA, pp. 124-137.

Day DM & Wallman JF (2006) Width as an alternative measurement to length for postmortem interval

estimations from *Calliphora augur* (Diptera: Calliphoridae) larvae. Forensic Science

International 159: 158-167.

Day DM & Wallman JF (2008) Effect of preservative solutions on preservation of *Calliphora augur*
and *Lucilia cuprina* larvae (Diptera: Calliphoridae) with implications for post-mortem interval
estimates. Forensic Science International 179: 1-10.

Donovan SE, Hall, MJR, Turner BD, Moncrieff & CB (2006) Larval growth rates of the blowfly,
Calliphora vicina, over a range of temperatures. Medical and Veterinary Entomology 20: 106-
114.

Draper NR & Smith H (1981) Applied Regression Analysis, 2nd edn. Wiley, New York, NY, USA.

Early M & Goff ML (1986) Arthropod succession patterns in exposed carrion on the island of O'Ahu,
Hawaiian Islands, U.S.A. Journal of Medical Entomology 23: 520-531.

Fuller ME (1934) The Insect Inhabitants of Carrion: A Study in Animal Ecology. Bulletin of the
Council for Scientific and Industrial Research 82: 1-63.

Geisser S & Johnson W (2006) Modes of Parametric Statistical Inference, Wiley, New York, NY,
USA.

Goff ML (1991) Comparison of insect species associated with decomposing remains recovered inside
dwellings and outdoors on the island of Oahu, Hawaii. Journal of Forensic Sciences 36:748-753.

Grassberger M & Reiter C (2001) Effect of temperature on *Lucilia sericata* (Diptera: Calliphoridae)
development with special reference to the isomegalen- and iosmorphen-diagram. Forensic
Science International 120: 32-36.

Greenberg B (1991) Flies as forensic indicators. Journal of Medical Entomology 28: 565-577.

Haskell NH (1991) Procedures in the entomology laboratory. Entomology and Death: a procedural
guide (ed. by EP Catts & NH Haskell) Joyce's Print Shop Inc, Clemson, SC, USA, pp. 111-123.

Hastie TJ & Tibshirani RJ (1990) Generalized Additive Models. Chapman and Hall, London, UK.

Higley LG & Haskell NH (2010) Insect development and forensic entomology. Forensic Entomology:
the Utility of Arthropods in Legal Investigations (ed. by JH Byrd & JL Castner) CRC Press,
Boca Raton, FL, USA, pp. 389-405.

Johnston TH & Hardy GH (1923) Observations regarding the life-cycle of certain Australian blowflies.
Proceedings of the Royal Society of Queensland 35: 21-42.

Johnston TH & Tiegs OW (1922) Notes on the biology of some of the more common Queensland
muscoid flies. Proceedings of the Royal Society of Queensland 34: 77-104.

Kamal AS (1958) Comparative study of thirteen species of sarcosaprophagous Calliphoridae and
Sarcophagidae (Diptera). I. Bionomics. Annals of the Entomological Society of America 51:
261-271.

Lane RP (1975) An investigation into blowfly (Diptera: Calliphoridae) succession on corpses. Journal
of Natural History 9: 581-588.

Lee DJ (1968) Human myiasis in Australia. Medical Journal of Australia 1: 170-173.

Levot G (2003) Insect fauna used to estimate the post-mortem interval of deceased persons. General
and Applied Entomology 32: 31-39.

Levot GW, Brown KR & Shipp E (1979) Larval growth of some calliphorid and sarcophagid Diptera.
Bulletin of Entomological Research 69: 469-475.

Lumley T, Diehr P, Emerson S, Chen L (2002) The importance of the normality assumption in large
public health datasets. Annual Review of Public Health 23:151-169

Mackerras MJ (1933) Observations on the life-histories, nutritional requirements and fecundity of
blowflies. Bulletin of Entomological Research 24: 353-362.

Mackerras MJ & Freney MR (1933) Observations on the nutrition of maggots of Australian blow-flies.
Journal of Experimental Biology 10: 237-246.

Meier R, Kotrba M & Ferrar P (1999) Ovoviviparity and viviparity in the Diptera. Biological Reviews
74: 199-258.

Monzu N (1977) Coexistence of Carrion Breeding Calliphoridae (Diptera) in Western Australia. PhD
Thesis, University of Western Australia, Crawley, WA, Australia.

Nabity PD, Higley LG & Heng-Moss TM (2006) Effects of temperature on development of Phormia
regina (Diptera: Calliphoridae) and use of developmental data in determining time intervals in
forensic entomology. Journal of Medical Entomology 43: 1276-1286.

Neideregger S, Wartenberg N, Spiess R, Mall G (2013) Influence of food substrates on the
development of the blowflies *Calliphora vicina* and *Calliphora vomitoria* (Diptera,
Calliphoridae). Parasitology Research 112:2847-2853.

Newman LJ & Clark J (1926) Trapping blowflies. Journal of the Department of Agriculture, Western
Australia 2: 382-391.

Nishida K, Shinonaga &, Kano R (1986) Growth tables of larvae for the estimation of post mortem
interval. Ochanomizu Medical Journal 34: 157-172.

Norris KR (1959) The ecology of sheep blowflies in Australia. Monographiae Biologicae:
Biogeography and Ecology in Australia (ed. by A Keast, RL Crocker & CS Christian) Junk, The
Hague, The Netherlands, pp. 514-544.

O'Flynn MA & Moorhouse DE (1980) Identification of early immature stages of some
common Queensland carrion flies. Journal of the Australian Entomological Society 19:
53-61.

O'Flynn MA (1976) A Study of Blowflies Breeding in Sheep and in Carrion. MAgSc Thesis,
University of Queensland, Brisbane, Qld, Australia.

O'Flynn MA (1980) Studies on Blowflies and Related Flies from Queensland. PhD Thesis, University
501   of Queensland, Brisbane, Qld, Australia.

502   O'Flynn MA (1983) The succession and rate of development of blowflies in carrion in southern
503   Queensland and the application of these data to forensic entomology. Journal of the Australian
504   Entomological Society 22: 137-148.

505   Richards CS, Crous KL & Villet MH (2009) Models of delevelopment for blowfly sister species
506   *Chrysomya chloropyga* and *Chrysomya putoria*. Medical and Veterinary Entomology 23: 56-61.

507   Rivers DB, Thompson C, Brogan R (2011) Physiological trade-offs of forming maggot masses by
508   necrophagous flies on vertebrate carrion. Bulletin of Entomological Research 101:599-611.

509   Sharma R, Garg RK, Gaur JR (2015) Various methods for the estimation of the post mortem interval
510   from Calliphoridae: A review. Egyptian Journal of Forensic Sciences 5:1-12.

511   Smith KGV (1986) A manual of forensic entomology. British Museum (Natural History), London

512   Tarone AM & Foran DR (2008) Generalized additive models and *Lucilia sericata* growth: assessing
513   confidence intervals and error rates in forensic entomology. Journal of Forensic Sciences 53:
514   942-948.

515   von Zuben CJ, Bassanezi RC & von Zuben FJ (1998) Theoretical approaches to forensic entomology.
516   II. Mathematical model of larval development. Journal of Applied Entomology 122: 275-278.

517   Wallman JF & Adams M (1997) Molecular systematics of Australian carrion-breeding blowflies of
518   the genus *Calliphora* (Diptera: Calliphoridae). Aust Journal of Zoology 45: 337-356.

519   Wallman JF & Archer MS (2020) The application of insects to the estimation of the time since death.
520   In J Hayman & M Oxenham, Estimation of the Time Since Death (pp. 57-74). Elsevier, London,
521   UK

522   Wallman JF (2001a) A key to the adults of species of blowflies in southern Australia known or
523   suspected to breed in carrion carrion [corrigendum in Medical and Veterinary Entomology 16:
524   223]. Medical and Veterinary Entomology 15: 433-437.

525   Wallman JF (2001b) Third-instar larvae of common carrion-breeding blowflies of the genus
526   *Calliphora* (Diptera: Calliphoridae) in South Australia. Invertebrate Taxonomy 15: 37-51.

527   Wallman JF (2004) The application of forensic entomology to criminal investigations. The practice of
528   crime scene investigation (ed. J Horswell) CRC Press, Boca Raton, FL, USA, pp 347-359.

529   Wells JD & Kurahashi H (1994) *Chrysomya megacephala* (Fabricius) (Diptera: Calliphoridae)
530   development: rate, variation and the implications for forensic entomology. Japanese Journal of
531   Sanitary Zoology 45: 303-309.

532   Wells JD & LaMotte LR (1995) Estimating maggot age from weight using inverse prediction. Journal
533   of Forensic Sciences 40: 585-590.

534   Wells JD & LaMotte LR (2017) The role of a PMI-prediction model in evaluation forensic entomology
535   experimental design, the importance of covariates, and the utility of response variables for
536   estimating time since death. Insects 8:47-54.