Title

Transcripts repressed at the stop of phloem unloading highlight the energy efficiency of sugar

import in the ripening *V. vinifera* fruit.

Running Title

Energetics of grapevine berry ripening

Authors

Stefania Savoi[1], Laurent Torregrosa[1], Charles Romieu[1*]

Affiliation

[1]AGAP, Montpellier University, CIRAD, INRAe, Institut Agro-Montpellier SupAgro, 34060

Montpellier, France

Email

Stefania Savoi: stefania.savoi@supagro.fr

Laurent Torregrosa: laurent.torregrosa@supagro.fr

Charles Romieu: charles.romieu@inrae.fr

Corresponding author

Charles Romieu, AGAP, Montpellier University, CIRAD, INRAe, Montpellier SupAgro, 2 Place

Pierre Viala, 34060 Montpellier, France.

Tel. +33 04 99 61 28 88

Abstract

Transcriptomic changes at the cessation of sugar accumulation in the pericarp of *Vitis vinifera* were addressed on single berries re-synchronized according to their individual growth patterns. The net rates of water, sugars and K^+ accumulation inferred from individual growth and solute concentration confirmed that these inflows stopped simultaneously in the ripe berry, while the small amount of malic acid remaining at this stage was still being oxidized at a low rate. Re-synchronized individual berries displayed negligible variations in gene expression among triplicates. RNA-Seq studies revealed sharp reprogramming of cell wall enzymes and structural proteins, associated with an 80% repression of specific sugar transporters and aquaporins on the plasma or tonoplast membranes, at the stop of phloem unloading in the three genotypes and two environments investigated. The prevalence of SWEET transporters suggests that electrogenic transporters would just play a minor role on the plasma membrane of SE/CC complex, and the one of the flesh, while sucrose/H^+ exchangers dominate on its tonoplast. *Cis*-regulatory elements present in their promoters allowed to sort these transporters in different groups, also including specific TIPs and PIPs paralogs, and cohorts of cell wall related genes. These results lead us to propose which structural, developmental and energy adaptations would give this fruit such a power of attraction for water and photoassimilates.

Introduction

During ripening, fleshy fruits shift from a repulsive to an attractive and nutritionally rewarding status, thanks to the induction of a finely orchestrated transcriptomic reprogramming in their pericarp, triggering convergent softening, colouration, accumulation of soluble sugars, and aroma development in unrelated species[1]. Fleshy fruits are divided in climacteric (e.g. tomato, apple, pear, etc.) and non-climacteric ones (e.g. grape, strawberry, citrus, etc.) depending on the occurrence of an autocatalytic emission of ethylene and a respiratory burst facilitating starch hydrolysis and cell wall degradation at the onset of ripening[2]. Grape, as the model of non-climacteric fruits, does not store any starch reserves and necessarily ripens on the vine, at the expense of a massive phloem unloading of sucrose[3] tightly connected with malate breakdown and fruit expansion[4]. It was recently advanced that the high rate of sugar accumulation would raise an energy challenge considering the limited oxidative capacity of grapevine, which could be solved by the discharge of phloem-vectored K^+ through the *VviK3.1* channel[5]. However, the ATP needed for sugar unloading critically depends on the respective expression and activities of

SWEETs, sugar/H^+ symporters and antiporters on the serial membrane interfaces from sieve

elements to pericarp vacuoles, which remain to be elucidated even though the grapevine gene

families of SWEETs, sucrose, and hexose transporters have been already described. Single berry

kinetic data now argue in favour of a global sucrose/H^+ exchange at the tonoplast[4].

Transcriptomics approaches have become frequently employed in grapevine physiological

studies focusing either on berry development or in responses to abiotic or biotic stresses. Several

developmental studies targeted the incipit of berry ripening[6,7] or the late phases of berry

development[8–11] for finding master regulators of key transitions[12]. However, albeit the clustering

of RNA-Seq samples largely depended on sugar concentrations, in none of these works, there

was a hint on the origin of the arrest of sugar phloem unloading in the ripening berry.

The grape berry development follows a double sigmoid curve[13]. The major events

associated to each phase can be summarised as (i) a first growth phase resulting from cell

division and expansion, depending on the accumulation of tartaric and malic acids as major

osmotica in the green berry; (ii) a lag phase with no growth; (iii) the ripening phase initiated by

berry softening, during which 1 M of glucose plus fructose accumulates at the expense of

imported sucrose. During this last phase, malic acid is consumed, and the berry volume nearly

doubles due to water influx. Before ripening, the xylem sap is the core source of water and

minerals but phloem mass flow prevails after the onset of ripening, due to a switch from the

symplasmic to the apoplastic unloading pathway[14]. One may consider that the berry is

physiologically ripe when phloem unloading definitively stops, whereupon sugar concentration

continues to increase only due to evaporation. It was accepted that sugar unloading might

continue following growth cessation, so that excess water in phloem mass flow must be rejected

by reverse xylem back-flow during late ripening[3,15,16]. However, such disconnection between

sugar and water flows can be an artefact resulting from mixing asynchronised berries[4].

Difficulties emerge from the two to three weeks delay between ripening berries, which proves as

long as the duration of the second growth phase itself, precluding the date of phloem arrest from

being unambiguously defined on usual asynchronous samples[4,17]. Transcriptomic studies may

help to find out the regulation of sugar concentration at phloem arrest, a first step for

understanding the impact of climatic change on grape composition.

Attempts were made to sort berries by softening and density since the earliest

transcriptomic studies[18] to unravel key regulatory genes implicated in berry ripening. In 2008, a

pioneering study[19] investigated individual berries sorted by firmness and colour to study the

early ripening process. In another work, individual berries were also considered to assess if the
ripening program was accelerated in "delayed" berries[20]. Following density sorting, attempts
were made to identify genes expressed or repressed at the late stages of ripening, which may be
used as indicators of maturity for oenological scope[8,21]. Individual berries from vines that
underwent heat stress or leaf-roll virus infection were re-synchronized based on their primary
metabolites content[22–24]. This brief survey shows that the awareness of berry heterogeneity is a
rising topic, although usual samples aiming to represent the whole diversity at the experimental
plot still predominate[7,25].

In this work, we tackle the long-standing issue of sugar accumulation mechanisms in
grapes upon elucidating which transcriptomic reprogramming would trigger phloem switch-off at
ripe stage. Pervasive bias caused by averaging asynchronised fruits was discarded upon sorting
single berries according to their actual rate of water and sugars accumulation. Prevalent
transcripts stably identified in different genotypes and environments highlight the functional
bases of the accumulation of 1 M hexose in this non-climacteric fruit.

Results

Experimental setting and monitoring of single grape berry growth during ripening

The experiment was conducted with three genotypes of *V. vinifera*: cv. Syrah, grown
outdoors, and two microvines (MV032 and MV102), grown in a greenhouse, resulting in
drastically different environments (Fig. S1). Although MV032 and MV102 were classified as
high sugar accumulator genotypes among a segregating microvine population[26], they showed a
35% lower sugar concentration at phloem arrest (Fig. 1, Table S1). Moreover, MV032 was
ranked as a low-acid genotype while in comparison MV012 displayed +13% and +69% malic
acid at the beginning of fruit ripening and at the phloem arrest (Fig. 1, Table S1). Finally,
MV102 carried the rpv1/run1 locus which confers a high level of tolerance to downy and
powdery mildews whereas this locus was absent in the MV032 susceptible genotype.

Fruit growth was monitored through bi-weekly pictures and image analysis, starting from
berry softening (stage V for veraison, the beginning of sugar accumulation) (Fig. 1a). Syrah,
MV032, and MV102 berries were sampled at three specific stages. The first sampling, called
stage G for growing, included berries growing at full speed, whose volume increased by roughly
50% within two weeks after softening. The second sampling, called stage P for peak, represented
berries that were as close as possible to the peak volume, and therefore growth arrest. Lastly, the

third sampling, called stage S for shrivelling, represented those berries that have peaked two weeks ago, and thus have reached the shrivelling period (Fig 1a). The changes in the total amounts of solutes per fruit (concentration x volume) confirmed that the net influx of water, photoassimilates, and K^+ simultaneously and definitively stopped at P stage (Fig. 1, Table S1).

Figure 1. Growth and solute accumulation during the ripening of synchronised grapevine single berries of MV032, MV102, and Syrah. (a) Relative growth, (b) glucose + fructose, (c) potassium, (d) malic acid, and (e) tartaric acid. Results in b and c were normalised in order to consider 1 g of pericarp at phloem arrest. The developmental stage is indicated by colours: V for veraison in

green, G for growing in yellow, P for peak in pink, S for shrivelling in purple. Letters indicate significant differences according to the Tukey ad hoc test.

Single berry primary metabolites analysis

Sugars are translocated by phloem in the form of sucrose in *V. vinifera*, and the cleaved glucose and fructose (Fig. 1b, Table S1) accumulate in equimolar amounts from stage V as the berry begins to soften. The amount of glucose + fructose in the berry dramatically increased from stage V to stage P indicating maximal activity of the apoplastic pathway of phloem unloading in G stage, accompanied by permanent cellular expansion and cell wall extension in all berries, as monitored by growth (Fig. 1a). The amount of sugar quantified in each berry did not increase any longer after the cessation of berry growth, indicating the stop of the phloem unloading. Moreover, in field conditions, the Syrah sugar content decreased in stage S, probably due to increased respiration during a heatwave recorded in those weeks (Fig. S1), while no significant decrease occurred in the greenhouse. K^+ accumulation inside the berry (Fig. 1c, Table S1) showed the same behaviour as sugars, though its intensity was, on average, 30 and 15 times lower in Syrah and in the two microvines. K^+ increased during the active growth phase mostly before the G stage in MV032, MV102, and after it in Syrah. There was thus no clear relationship in the timing and intensities of K^+ and sugar unloading in the berries, except that when the phloem stopped, the amount of both solutes remained constant whatever the genotype.

Malic acid (Fig. 1d) displayed the typical pattern of malate breakdown in ripening *V. vinifera* berries, with a faster reduction between stage V and G in all genotypes. Two weeks after the cessation of phloem import (stage S), it remained between 20% (Syrah and MV032) and 40% (MV102) of the initial malic content measured in V stage. To notice that MV102 classified as high-acid genotype kept a higher amount of malic acid in shrivelling berries. It is widely accepted that tartaric acid is not metabolised during berry ripening. Indeed, in Fig. 1e, one can observe that Syrah and MV032 kept more than 90% of their initial tartaric acid content, while tartrate was reduced by less than 20% in MV102.

Transcriptomic overview

For RNA-sequencing, nine berries per genotype (3 berries for each stage G, P, and S) were carefully selected (Fig. S2) according to their individual growth rate and primary metabolites contents. A principal component analysis (PCA) was performed over the 27 berry transcriptomes

(Fig. 2a). The first and the second principal component (PC1 and PC2) accounted for 37% and
21% of the variance in gene expression among samples. Sample distribution was driven by both
the developmental stage and genotype, with a clear distinction between Syrah and the two
microvines. G stage samples were resolved from those at P and S stage in Syrah and MV032,
while in MV102 stage distinction was less obvious. Except for one MV032 sample, stages P and
S samples were mainly clustered in all genotypes. Cluster dendrogram analysis (Fig. 2b) divided
the samples into four main clades. Genotypes fell in different branches that were subdivided
according to the stage, except for MV102-P-2 that was more close to MV102-G-3.

Overall, the sampling strategy based on image analysis and primary metabolites
concentration resulted in the adequate clustering of the biological triplicates with very few
exceptions found especially in the most difficult physiological stage to detect, stage P.

181
Figure 2. Overview of the transcriptomic results: (a) principal component analysis, and (b)
sample dendrogram. Colours and symbols indicate genotypes and stages, respectively: MV032 in
yellow, MV102 in red, Syrah in green; a circle for stage G, a square for P, and a triangle for S.

Time significant genes

A comparable number of genes displayed significant (P<0.05) developmental changes in
expression in the three genotypes (7025, 6629, and 6505 in Syrah, MV032, and MV102, Table
S2a). They were further grouped according to their expression profiles (Fig. S3), and the cluster
number was restricted to four upon visual analysis. Cluster A and B included the majority of the
genes increasing or decreasing in expression from G to S (Fig. 3a). The genes commonly
regulated in the three genotypes, as shown in the Venn diagram in Fig 3b, were then screened for

enrichment in GO categories in the Biological Process (BP), Cellular Component (CC), and Molecular Function (MF) groups. Ultimately, clusters C and D showed genes up-peaking or down-peaking in stage P (Fig. S4a). Stage P was the most problematic stage to detect, as highlighted by the Venn diagram (Fig. S4b) where very few genes if not zero were in common between two genotypes, probably because the gene expression peak just represented small erratic variations. Growth and phloem switch-off should be a fast transition; therefore P stage detection critically relies on the accuracy of the growth measurement method and can be only characterised as *a posteriori*. The time needed to screen berries by image analysis forced us to sample the next morning when some metabolic processes were already stopped or partially modified.

Clusters A included genes whose expression increased as growth and phloem arrested and berries started to shrivel (Fig. 3a, Table S3). Six-hundred-two genes were commonly modulated in the different genotypes (Fig. 3b), whose enriched GO categories are listed in Fig. 3c. Of interest, we observed catabolic process-related categories and RNA metabolic process for BP, encoding proteins with a principal localisation in the intracellular space, but also nucleus and organelle for CC, and several binding categories were over-represented in MF. Diverse transcription factors belonging to the NAC (7 up-regulated genes), ERF/AP2 (7), bZIP (7), WRKY (7), LBD (2) multigenic families together with RNA helicases, translation initiation and splicing factor subunits and chromosome organization-related genes were up-regulated. Consistently, chromosome organisation and regulation of transcription were already reported as up-regulated in late stages of ripening[10].

Cluster B encompassed genes decreasing in expression during development (Fig. 3a, Table S3), showing a higher number of commonly modulated genes in the three genotypes (846) (Fig. 3b). Noticeably, the BP enriched GO categories included "Water transport", "Cell wall organisation", "Polysaccharide and Carbohydrate metabolic process", while for CC several categories were related to membranes, and "Water channel activity" and "Water transmembrane transporter activity", were highly represented in MF (Fig. 3c). Among all, seven PIPs (*VviPIP1.3*, *VviPIP2.5*, *VviPIP2.3*, *VviPIP1.4*, *VviPIP2.7*, *VviPIP2.4*, *VviPIP1.2a*) and three TIPs (*VviTIP1.2*, *VviTIP1.3*, *VviTIP2.1*), sugar transporters such as *VviHT6*, *VviSWEET10*, the vacuolar invertase *VviGIN2*, the alcohol dehydrogenase *VviADH2* were highly down-regulated in the three genotypes. Besides, numerous genes related to the cuticle or the cell wall metabolism encompassing cellulose, pectin metabolism, and several expansins were strongly down-

regulated. As examples, we observed *VviCER3-like*, *VviPAS2-like,* and a *VviCYP716A-like*,
which are involved in the cuticular aliphatic wax biosynthesis[27], cellulose synthases A and
cellulose synthase-like, a pectate lyase, several fasciclin-like arabinogalactan proteins, pectin
methyl esterases, polygalacturonases, and expansins A or expansins A-like such as *VviEXPA04*,
VviEXPA11, *VviEXPA18*, *VviEXLA01*. Moreover, several genes related to the
glycolysis/gluconeogenesis pathways, such as a UTP-glucose-1-phosphate uridyltransferase, two
triose-phosphate isomerases, a glyceraldehyde-3-phosphate dehydrogenase, a phosphoglycerate
mutase, one enolase were down-regulated together with malate dehydrogenases.

Figure 3. Significant time-course genes expression of (a) cluster A and B representing genes
increasing or decreasing respectively in expression at phloem arrest. (b) Venn diagram showing
genes displaying common or genotype specific changes in expression at phloem arrest. (c)
Significantly enriched GO categories in modulated genes, divided in biological process (BP),
cellular component (CC), and molecular function (MF).

Pairwise differential gene expression (S vs G)

Other important differentially expressed genes (DEGs) could be identified in Syrah and
only one microvine due to discrepancies at P stage. Since the variance among P stage samples
was certainly exacerbated by abrupt changes in physiology and gene expression at phloem arrest,
worsened by sampling delay, data were tested in pairwise mode analysing stage S versus stage
G.

A total number of 6585, 6453 and 5621 DEGs were identified in Syrah, MV032, and

MV102 (Table S2b). The commonly modulated genes increased to 899 up-regulated genes and

1155 down-regulated ones, yielding to 2054 DEGs (Table S3). Others 411 genes were modulated

in the three genotypes but with a discordant gene trajectory. Compared to the previous analysis,

this second comparison gave a more comprehensive list of commonly modulated DEGs (before

they were 1448). These two lists shared 809 genes. In this new DEGs list, there were down-

regulated genes like *VviTMT2*, *VviHT2*, *VviGIN1, VviH$^+$ATPase, VviEXP19* and other cell wall-

related genes (Table S3).

Rates of gene expression: ranking genes by transcription priority

Gene expression burdens cells by consuming resources and energy for transcription,

translation, and the synthesis and degradation of hydrophobic proteins like transporters is

particularly expensive[28]. In this respect, the transcription of transporters must stop very fast once

they are no longer required. As a proxy for energy cost, we merged and ranked the two lists of

differentially expressed genes according to their absolute change in expression (CPM, i.e.

uncorrected for transcript length) during the stop of water, sugar and K$^+$ influx in the berry (S

versus G), as summarised in Table 1 (full list in Table S3). This straightforward procedure

proved extremely efficient in the detection of channels and transporters virtually turned off

simultaneously with phloem together with cell wall-related genes. Amazingly, in Syrah, among

solute transporters, *VviHT6* exhibited the largest decay in expression from G to S stages with

80% inhibition, indicating tight synchronism with phloem arrest. This was also verified for

VviSWEET10, *VviPIP1.3* and *VviPIP2.5*, which decreased by 90% and 83%. Moreover, a pectate

lyase was virtually shut down (-98%) with several expansins (*VviEXPA19*, *VviEXPA14*,

VviEXPA16, -73%, -92%, -99%). On the other hand, genes annotated as hydroperoxide lyase

(+244%), metallothionein (+213%), and polyubiquitin (+73%) appeared as the most induced

ones.

Table 1 List of thirty-five selected genes related to cell growth, water and sugar transport sorted by absolute difference in Syrah G versus P.

Gene ID is reported as VCost.V3 and RefSeq code. Gene expression is expressed as count per million (CPM).

VCost.V3	RefSeq	Syrah G (CPM)	Syrah P (CPM)	Syrah S (CPM)	Variation Syrah (%)	Difference Syrah	Rank Syrah	Annotation
Vitvi05g01947	LOC100264738	5380	974	376	-93	-5003.9	1	early nodulin-75-like
Vitvi04g00442	LOC100252479	4387	720	241	-95	-4146.8	2	Uncharacterized
Vitvi02g00310	LOC100233001	4818	1336	814	-83	-4004.5	3	[Aquaporin] VviPIP1-3
Vitvi16g01800	LOC100232884	8635	6627	5693	-34	-2941.5	4	[Cell wall - SWITCH] invertase/pectin methylesterase inhibitor
Vitvi13g00605	LOC100233002	2773	971	462	-83	-2311.3	5	[Aquaporin] VviPIP2-5
Vitvi18g00189	LOC100245385	2812	1299	751	-73	-2061.0	8	[Cell wall] Expansin A (VvEXPA19)
Vitvi13g00172	LOC100244917	1941	414	161	-92	-1780.1	13	[Cell wall] Expansin A (VvEXPA14)
Vitvi18g00056	LOC100232977	1609	345	302	-81	-1307.6	21	[Sugar] Tonoplast Monosaccharide Transporter HT6
Vitvi05g00953	LOC100232902	1143	72	17	-98	-1125.5	24	[Cell wall] Pectate lyase
Vitvi08g01038	LOC100233094	1045	105	74	-93	-971.6	26	[Aquaporin] VviPIP2-3
Vitvi12g02137	LOC100233113	1185	499	250	-79	-934.2	27	[Cell wall] Pectin methylesterase
Vitvi06g01329	LOC100265471	1947	1129	1070	-45	-876.4	29	[Cell wall - SWITCH] xyloglucan endotransglucosylase/hydrolase 32
Vitvi04g00465	LOC100254408	1092	388	308	-72	-783.6	32	[Cell wall] Cellulose synthase CESA1
Vitvi08g01602	LOC100233004	823	163	101	-88	-721.4	34	[Aquaporin] VviTIP1-2
Vitvi05g00548	LOC100245945	605	52	45	-93	-559.8	40	Sulfate transporter 3.1
Vitvi15g01110	LOC100240701	827	360	322	-61	-504.8	47	[Aquaporin] VviPIP1-4
Vitvi17g00070	LOC100267921	511	71	48	-91	-462.8	52	[Sugar] bidirectional sugar transporter SWEET10
Vitvi15g00643	LOC100253046	1015	413	554	-45	-460.7	53	[Cell wall - SWITCH] Expansin B (VvEXPB04)
Vitvi03g00247	LOC100264011	594	212	143	-76	-451.2	55	[Sugar] Tonoplast Monosaccharide Transporter TMT2
Vitvi18g00008	LOC100242715	441	80	72	-84	-369.5	72	[Cell wall] Cellulose synthase CESA2
Vitvi03g00155	LOC100233027	434	139	70	-84	-363.9	73	[Aquaporin] VviPIP2-7
Vitvi11g00272	LOC100233140	541	259	177	-67	-363.6	74	Malate dehydrogenase (NADP+)

Vitvi04g01668	LOC100232854	1434	786	1084	-24	-350.1	78	Alcohol dehydrogenase 2
Vitvi16g00713	LOC100256970	351	24	15	-96	-335.9	82	[Sugar] Vacuolar Invertase GIN1
Vitvi14g01977	LOC100244103	297	2	3	-99	-294.2	93	[Cell wall] Expansin A (VvEXPA16)
Vitvi18g00397	LOC100232961	333	119	52	-84	-280.7	105	[Sugar] Hexose transporter HT2
Vitvi03g00209	LOC100258952	286	35	25	-91	-260.4	113	[Cell wall] Expansin A-like (VvEXLA01)
Vitvi18g01628	LOC100255463	351	133	106	-70	-245.8	124	[Cell wall] Cellulose synthase CESA2
Vitvi11g00835	LOC100248411	277	102	47	-83	-229.9	131	[Cuticle metabolism] CER3like - Aliphatic wax
Vitvi12g00721	LOC100242544	237	9	9	-96	-227.4	134	fasciclin-like arabinogalactan protein 2
Vitvi13g00255	LOC100233003	245	48	32	-87	-213.6	143	[Aquaporin] VviTIP1-3
Vitvi13g01731	LOC100256811	289	126	133	-54	-156.0	186	Cellulose synthase CESA3
Vitvi01g01805	LOC100232906	157	7	3	-98	-153.9	189	Xyloglucan endotransglucosylase/hydrolase protein B
Vitvi17g01251	LOC100261426	136	7	2	-99	-134.2	225	Expansin A (VvEXPA18)
Vitvi06g00281	LOC100233093	126	23	12	-91	-113.9	281	[Aquaporin] VviPIP2-4

    Promoter analysis

Sixty selected genes related to water, sugar, and growth down-regulated at the stop of

phloem were distributed in different regulatory networks when clustered according to the Z-

score[29] which accounts for length, number and position of their promoter motifs (Fig. 4).

C2C2dof, NAC, G2like, Homeobox, WRKY, MYB, and AP2/EREB families were driving the

major differences. Interestingly, SWEETs were unambiguously separated from electrogenic

transporters such as *VviHT6*: AP2/EREB and NAC binding site motifs were mainly absent from

VviSWEET10 promoter but present in *VviHT6*. Conversely, WRKY motifs absent from *VviHT6*

were present in *VviSWEET10*. While MYB motifs were close to TSS in SWEETs, they were far

from *VviHT6* one. Moreover, differences appeared among PIP or TIP isogenes: by example,

VviPIP2.4 and *VviPIP2.5* belonged to the same branch as SWEET while *VviPIP1.4* and

VviPIP2.3 showed a different profile. The different regulatory networks may indicate that these

down-regulated genes are expressed in separated cell types or tissues inside the pericarp, such as

phloem SE/CC, hypertrophied cells in the flesh or subepidermal tissues where respiration occurs.

Figure 4 Genes inhibited at phloem stop belong to different regulatory networks. Genes and transcription factors were clustered according to the Z-score of the motifs present within 1500 bp before the TSS. Red and blue colours indicate proximity or distance of the promoter sequence to the TSS. White colour denotes the absence of a specific TF in the promoter of a gene. Horizontal and vertical colour bars grouped promoter and genes in promoter families and gene functions as indicated in the legend. The two examples discussed in the text regarding SWEETs and HT6 with reference to NAC, MYB, and AP2/EREB are marked by black squares.

Discussion

For the first time, non-destructive monitoring of single berry growth allowed us to sample synchronised fruits within an original population marked by a significant delay in the development of individual fruits. Sorting berries according to their net growth (or water import) rate warranted that berries still growing and importing water were not mixed with shrivelling (over-ripe) ones. Real-time individual growth monitoring outperformed our previous sampling procedure which just relied on solute concentration[4,17] upon adding the lacking kinetic dimension. We showed here that the net imports of water, sugar, and K^+ simultaneously stop when the ripening berry reaches its maximum volume, which confirms that sugar accumulation at constant volume indicating intense xylem back-flow during late ripening[3,16] resulted from an averaging artefact on asynchronous samples[4]. This was verified in contrasted thermo-hydric conditions (field – higher temperature and water demand, versus greenhouse – non-limiting water supply) and when phloem unloading arrested at 0.8 M of sugar in microvines (as already observed for these genotypes[30]) and at 1.2 M in Syrah. At phloem arrest, with a malate concentration below 100 mEq (39 in Syrah, 48 and 93 mEq in MV032 and MV102), the berry already consumed between 60 and 80% of the malic acid accumulated before softening. These differences in acidity that would principally originate from the green stage accumulation period[26] did not apparently interfere with phloem arrest.

Our study, performed in two environments and in three genotypes displaying different ripening features, reports for the first time robust gene trajectories associated with the end of the sugar and water accumulation processes, with the most intensively down-regulated genes related to cell wall modification (arrest of growth), aquaporins and sugar transporters (arrest of phloem unloading). The prevalence of plasma-membrane channels and transporters genes virtually turned off at phloem arrest re-emphasizes the central role of the apoplasmic pathway in ripening berry. It clearly illustrates that a strategy addressing the single berry level can lead to more comprehensive insights on fruit developmental biology than random samplings.

The arrest of growth is modulated by the down-regulation of cell wall associated genes

The large cohort of cell wall related genes being repressed at growth arrest yields an extremely dynamic image of the rearrangements accompanying cell wall extension and cellular expansion. This cell wall life seemed to be extremely reduced after growth cessation and phloem arrest where many genes were down-regulated. The present single berry profiling greatly affirms

the connection between aquaporins, cell wall and sugar transport. No obvious marker of cell

death was detected here, which is often described later in the shrivelling process of Syrah

berries[31].

Here we reported that genes involved in cellulose metabolism (15 cellulose synthases),

hemicellulose metabolism (4 endo-1,4-beta-glucanases, 7 xyloglucan endo-transglucosylase /

hydrolase proteins), pectin metabolism (4 pectate lyase, 4 polygalacturonases, 6 ß-galactosidases,

6 fasciclin-like arabinogalactan protein), and expansins genes (9) were strongly down-regulated

at the arrest of phloem (Table S3). In particular, we emphasise that precisely such expansins

(*VviEXPA19*, *VviEXLA1*, *VviEXPB4*, *VviEXPA14*) previously linked to growth and cellular

expansion during the second growth phases both in the flesh and in the skin[32] were down-

regulated concomitantly with growth cessation. A link between these expansins and aquaporins

has been envisaged in a pre-genomic study reporting a down-regulating trend two or three weeks

before the attainment of maximum berry size on asynchronised berry samples[33]. This so-called

"unexpected" link was recently confirmed through a gene coexpression network analysis[34].

The arrest of phloem is linked with strongly down-regulated aquaporins

Aquaporins are transmembrane channels facilitating the transport of water and small

solutes from cell to cell and between cell compartments. Noticeably, seven of the ten plasma

membrane intrinsic proteins (PIPs) identified in *V. vinifera*[34] were drastically inhibited (Table 1,

Table S3) simultaneously with growth and phloem arrest. Among them, *VviPIP1.3*

(*Vitvi02g00310 - LOC100233001*), *VviPIP2.5* (*Vitvi13g00605 - LOC100233002*), and *VviPIP2.3*

(*Vitvi08g01038 - LOC100233094*) were the most intensively repressed transcripts (ranked 3rd,

5th, and 26th according to their absolute expression changes in Syrah) with a percentage of

inhibition ranging from 83 to 93% in Syrah. Data on grapevine aquaporins[34], including the gene

expression atlas[25] and various other RNA-Seq datasets, showed that these three genes exhibited

quite a common expression pattern and actually decreased at ripe or harvest stages (defined as 20

°Brix) but the inhibition continued in post-harvest withering phases (see figure 3a,b in[34]). Present

results showed that this progressive decrease does not occur in individual fruits, but may

represent an emerging property on asynchronous samples. While the *PIP1* gene family was more

associated with dynamic changes in membrane permeability, *PIP2* was recently allocated to

sieve element plasma membrane in poplar, a symplastic loader[35]. Much work remains to be done

regarding their exact localisation in an apoplastic unloader such as grapevine berry insofar direct

orthologous relationship between *Vitis*, *Populus*, and *Arabidopsi*s are difficult to establish[34].

Finally, among the eleven tonoplast intrinsic proteins (TIPs), only three were significantly

inhibited. The TIPs expression level was lower than in the three previously discussed PIPs. In

particular, the predominant *VviTIP1.2* (*Vitvi08g01602 - LOC100233004*) was six times less

expressed than *VviPIP1.3* but exhibited a huge inhibition upon phloem arrest (88%).

Phloem unloading is stopped by the down-regulation of specific sugar transporters

Phloem unloading via the apoplastic pathway involves sugar transport through sequential

interfaces from the sieve element-companion cell complex (SE/CC) to sink tissues such as the

hypertrophied cells of the pericarp till its final storage compartment, the vacuole.

Sugar Will Eventually be Exported Transporters (SWEETs) are sugar uniporter proteins

recently identified in plants. We show here that *Vitvi17g00070* (*LOC100267921*) was the only

SWEET gene highly down-regulated at the arrest of phloem. This gene, annotated as SWEET14

at NCBI, corresponds to *VviSWEET10* (*VIT_17s0000g00830*) in [36], the ortholog of *AtSWEET10*,

a plasma membrane sucrose transporter of the clade III SWEETs. These authors[36] evidenced a

strong induction of *VviSWEET10* at veraison, without berry sorting nor compositional data, and

confirmed its expression at the plasma membrane. Ectopic expression in tomato increased its

sugar content, while promoter-GUS fusions showed preferential expression in the vascular

bundles and the flesh. These authors privileged a *VviSWEET10* role in hexose loading in the

flesh, but it may preferentially export sucrose from SE/CC to the pericarp apoplast as reported

for *AtSWEET11* and *AtSWEET12*, two other *Arabidopsis* genes belonging to the same clade III

SWEETs[37]. Noticeably, preferential expression of *VviSWEET10* was confirmed during ripening

in Riesling and Cabernet Sauvignon with a decrement at 100-110 days after veraison[38].

However, the same authors evidenced a higher expression of *VviSWEET15* in Petit Manseng,

that was not followed by a marked decrease at phloem arrest[38]. Our RNA-Seq data show that

VviSWEET15 displays a similar expression level than *VviSWEET10* during the active phloem

unloading without then being inhibited at phloem stop (data not shown) as reported in [36]. This

might suggest that *VviSWEET10* would be preferentially expressed at the phloem unloading site

while *VviSWEET15* would facilitate sugar transfer between the core and the periphery of the

fruit, where respiration occurs (Fig. 5).

Conversely to SWEET, sugar/H^+ antiporters and symporters can transport sugar against its

concentration gradient using the proton motive force. *VviHT6* (*Vitvi18g00056 - LOC100232977*)

was the sugar transporter with the highest decrement in expression at the stop of phloem. *VviHT6*

induction during Syrah ripening[39] was reported before the release of the *Vitis* genome[40] together

with its complete ORF (DQ017393.1), as confirmed in other cultivars[41]. The prevalence of

VviHT6 expression over those of *VviSUT*s and *VviHT*s was pointed out[42], but its synchronism

with sugar accumulation remained particularly elusive due to excessive sampling period and

absence of sugar measurements. The expected targeting of *VviHT6* to the tonoplast membrane

and its accumulation during ripening were masterly confirmed through proteomic analysis of

purified vacuoles[43], while present ectopic expression of *VviHT6::GFP* fusion in the hairy root

homologous system rules out its presence on grapevine plasma membrane (Fig. S5). Finally, the

recent re-evaluation of malate and sugar fluxes in synchronised berries provided first quantitative

evidence that an overall H^+/sucrose exchange parallels the predominant expression of *VviHT6* in

ripening berries[4]. In line with accumulating evidence that *VviHT6* orthologs should be renamed

tonoplast sucrose transporter (TST1/2) in *Arabidopsis*, apple, taproot, and melon[44–47], this

important result enlights the long-standing question of the origin of the malate/sugar relation in

grapevine berry. *VviTMT2* (*Vitvi03g00247 - LOC100264011*), an isogene three times less

expressed than *VviHT6* at G stage, decreased by 76, 74 and 54% in Syrah, MV032, and MV102;

however, at the protein level, *VviTMT2* and *VviHT6* display comparable expression[43]. Moreover,

the repression of *VviTMT2* was recently pointed out in low-sugar accumulating berries affected

by the ripening physiological disorder berry shrivel[48]. The third gene of this clade, *VviTMT3*

(*Vitvi07g01898 - LOC100243856*)[41,49] was not expressed (data not shown) although it was the

first one localised at the tonoplast[49].

The probable PM hexose symporter *VviHT2* (Vitvi18g00397 - *LOC100232961*)[41] also

exhibited a considerable increase during ripening, followed by a 35% decrease in two weeks[50].

Synchronised berries show here that more than 50% inhibition occurs at growth arrest, reaching

after that 84% decrease in Syrah. *VviHT2* was already associated with the induction of *VviHT6* at

veraison both in flesh and skin[51] and in ABA, and GA$_3$ treated berry[42,52], but data were lacking

regarding its inhibition at ripe stage. To the best of our knowledge, *VviHT2* localisation is still

unknown. Unfortunately, more information is available for *VviHT1,* but it was not commonly

modulated in the genotypes studied here.

Do different CREs belong to different tissues?

Several transcription factors (TF) were induced at P and S stages which may play a
decisive role in the stop of berry growth and phloem unloading. TFs regulate the expression of
genes involved in different biological processes by binding to specific nucleotide sequences in
their promoter. The genes commonly inhibited at the stop of phloem fell in different groups
when classified for the presence of specific CRE (Fig. 4), which may suggest they belong to
different cell types or tissues.

The presence of target CREs in the promoter of important down-regulated genes was
consistent with the parallel induction of some TFs. Among the most up-regulated NACs, which
binding sites were present in *VviHT6* but not in *VviSWEET10*, we notice *VviNAC18*
(Vitvi19g00271 - LOC100248744) and *VviNAC03 (Vitvi10g00437 - LOC100244547),* previously
identified as the two closest tomato NOR homologous[53], together with *VviNAC60*
(*Vitvi08g01842 - LOC100255512*) and *VviNAC11* (*Vitvi14g01985 - LOC100250998*) putatively
associated with the beginning of ripening process[53]. AP2/EREB motifs were also associated with
VviHT6 and not *VviSWEET10,* and besides a DREB2G down-regulated, *VviERF003*
(*Vitvi19g00558 - LOC100264172*) and *VviAP2* (*Vitvi07g01706 - LOC100268051*) were the most
changing AP2/EREB TFs. To the other side, among WRKY, which motifs were absent from
VviHT6 promoter but present in *VviSWEET10*, *VviWRKY47* (*Vitvi07g00523 - LOC100264000*)
showed the biggest change in expression. This gene was recently reported to participate in the
sugar and ABA signaling[54]. Finally, two LBD proteins (*VviLBD1a3 - Vitvi15g00735,*
LOC100255815 and *VviLBD1d6 - Vitvi06g00772, LOC100262112*) up-regulated at the stop of
phloem, were already identified as molecular markers of the ripening stage[55,56]; the LBD binding
motif was located in several genes with no specificities to particular branches/gene classes.
These exemplary TFs may possibly be involved in the regulation of the last stage of berry
ripening and in particular in the stop of sugar unloading, water and sugar transport, and cell
growth.
454
An energetically optimised sugar accumulation pathway operates in berries?
A ripening berry can just produce 3-6 ATP by oxidative phosphorylation while it
accumulates one hexose[4]. Thus, a phloem unloading pathway passing through H^+-coupled sugar
transporters on the three membrane interfaces from SE/CC complex to flesh vacuoles, combined
with the functioning of apoplasmic invertase, would waste within 50% of cellular ATP for H^+
recycling, in a pure loss as the sucrose gradient between both compartments is

thermodynamically favourable. Present results (DEGs, promoter analysis) lead to propose a more

efficient design of sugar import, which refines the previous ones[4,57] upon integrating key

genes/functions observed in this study (Fig. 5).

VviSWEET10, located on the SE/CC plasma membrane, would be the major player in the

export of sucrose from the phloem into flesh apoplasm. The simplest hypothesis is that sucrose

would then be transported by *VviSWEET15* on flesh cell plasma membrane, even though further

work is needed to precise its substrate specificity. Sucrose hydrolysis in the apoplasm would

require its resynthesis in flesh cells to allow sucrose vacuolar transport by TST (see discussion

above) which could be compatible with the huge induction of SPS during ripening[23,57] and with

SPS, SuSy, AI, NI *in vitro* activities[38] in five thousand excess, at least, with sugar accumulation

rate. However, except for AI, the corresponding genes were not strongly inhibited at phloem

arrest, which may indicate that sucrose resynthesis could be restricted to intercellular sugar

exchanges needed for respiration in fruit periphery, which continues after phloem arrest. In fact,

Vitvi07g00353 and Vitvi11g00542 annotated as sucrose synthase and sucrose phosphate

synthase (Table S4) were rather constant during the three stages in the three genotypes.

Sugar transport by SWEETs is energetically silent and the sink strength for sugar

accumulation is driven by the key electrogenic antiporters (proton-coupled, i.e. energy

consuming) *VviHT6* and *VviTMT2* that transport sucrose across the tonoplast. At the beginning

of ripening, the vacuolar acidity within 400 and 450 mEq (Table S1) at pH 2.7, provides a

considerable proton motive force for the transport of sucrose into the vacuole, as long as the

counterion malate is available. In tandem with vacuolar invertase, this system is susceptible to

restrict cytosolic sucrose below the nanomolar range (Table S5). Once in the cytoplasm, the

exchanged malic acid will serve as a substrate for respiration or gluconeogenesis, according to

the reactions: $2\,H^+ + malate + 3\,O_2 = 4\,CO_2 + 3\,H_2O$ and $2\,malate + 4\,H^+ \rightarrow 1\,hexose + 2\,CO_2$

that progressively vanish with the exhaustion of malic acid. Between 22 (Syrah, exposed to

stress) and 40% (MV032, greenhouse) of the H^+ exchanged with sucrose at the tonoplast (Table

S1) are scavenged from the cytoplasm through these reactions, but the remaining ones need to be

pumped back in the vacuole at the expense of a progressive activation of glycolysis, aerobic

fermentation (ADH2), vacuolar H^+/ATPase and H^+/PPiase[58]. It is noticeable that also *VviADH2*

is inhibited at phloem arrest, along with vacuolar invertase. Results on the respiration of single

grapevine berry, which provide strong arguments in this direction, will be detailed in a next

publication. The only PM H^+ symporter encountered here (*VviHT2*) displayed the highest

expression in Syrah, which also needed more ATP at the tonoplast (Table S1), when compared to microvines. This suggests that energy-requiring transports may be recruited to increase sink strength in unfavourable environmental conditions.

Figure 5 Hypothetical structure of the energy-efficient apoplastic pathway in the ripening berry. The figure is divided in three panels: early ripening, late ripening, and shrivelling with the stop

of phloem between the two latter (marked by a black circle). Genes written in red or blue italics

refer to those respectively expressed or repressed at specific stages, those in black are

constitutive.

The energetically-optimised model proposed here alleviates the hypothetical opening of an

AKT2-like channel that would play a major role in driving sugar transport upon compensating an

energetic deficit during phloem unloading[5]. Indeed, this last mechanism would require the

simultaneous activation of quantitatively similar flows of sugars and K^+ in ripening berries

which was never observed as confirmed here on single berry. In previous studies, where the

determination of the sampling stages was only based on Brix monitoring and not with precise

physiological landmarks, observed correlations likely resulted from a confusion between

accumulation and concentration mechanisms. Indeed, when the phloem unloading stops, berries

shrivel inducing concomitant concentrations of all soluble solids without any need of co-

transport mechanisms. In future studies, it would be interesting to understand which aspects of

this energetically optimised mechanism proposed here are conserved among cultivated and

ancestral cultivars or if a genetic diversity prevails, for example in extreme cultivars such as

acidless mutants or acidic varieties.

Materials and methods

Plants

Three genotypes were characterised in this study: the cv. *V. vinifera* Syrah, a traditional

variety adapted to warm climates and used for wine production, and two hermaphroditic

microvines offspring, named MV032 and MV102. Both microvines displayed a semi-dwarf

stature[30], resulting from five pseudo backcrosses between *Muscadinia rotundifolia* G52 and *V.*

vinifera genotypes. The last cross involved the 04c023V0003 female microvine[59] and the 3197-

81B hybrid[60]. Genetically, microvines used here can be considered very close to *V. vinifera* (1%

or less of *M. rotundifolia* background). Syrah vines grafted on SO4 were in an open field at the

experimental vineyard of Montpellier Supagro planted in 2000 while the two-years-old

microvines were grown in pots in a greenhouse under semi-controlled condition (25/15°C

day/night temperature, VPD 1 kPa, photoperiod 12 h).

Single berry sampling

Single berry growth and development were monitored through biweekly pictures of
selected clusters (up to 8 for each genotype) starting a few days before softening day until two
weeks after maximum berry growth. Pictures were taken using a Lumix FZ100 camera
(Panasonic), keeping the focal range and cluster to camera lens distance (30cm) constant. The
volume increment of selected berries was calculated by analysing the pictures with ImageJ[61]. The
software, after calibrating the images using the 1 cm scale present in each of them, automatically
counted the pixels enclosed in each targeted berry area, measured as an ellipsoid. The estimated
berry volume was then calculated accordingly using the radius of the previously calculated area.
To eliminate the intra- and inter- cultivars variability in berry sizes and compare the changes in
volume among berries of different clusters, vines, and genotypes, each berry growth profile was
normalised to the softening volume, set to 1.

According to their own volume changes, 10 to 15 berries were sampled for each genotype
at different dates during the ripening growing phase (stage G), closest to the berry growth peak
(stage P) and two weeks after the maximum growth (stage S) in the shrivelling phase.
Concerning stage P, one must notice that *a-posteriori* it was found that phloem and, obviously,
growth, were already blocked at this stage, which should, technically speaking, be called P+1.
Berries were sampled at the same time of the day, between 9 am to 11 am, to avoid circadian
cycle influences. Berries without pedicel were rapidly deseeded before freezing (1-2 min after
harvest) in liquid N_2 and stored at -80 until further analysis. Single berries were ground to a fine
powder under liquid nitrogen using a ball mixer mill (Retsch MM400).

Primary metabolites analysis

Glucose, fructose, malate and tartrate were analysed by HPLC. For each berry, 100 mg of
powder was weighed, diluted 5x with a solution of HCl 0.25 N, well shacked, and left overnight
at room temperature. Samples were then centrifuged for 10 min at maximum speed, and a
supernatant aliquot was diluted 10x with a solution of H_2SO_4 containing as internal standard 600
µM of acetic acid. Samples were injected according to [23]. For K^+ analysis, the previous samples
in HCl were further diluted 10x with MilliQ water and analysed as in [26]. Tukey.HSD post-hoc
test was used as a statistical method to find significant differences among metabolites
concentration over time.

RNA extraction and sequencing

Based on growth increment, internal sugars and organic acids, three berries per stage G, P,

and S for each genotype were selected for individual RNA extraction and library preparation, as

in [23]. Samples were sequenced on an Illumina HiSeq3000 in paired-end mode, 2x150 bp reads, at

the Genotoul platform of INRAe-Toulouse.

Data analysis

Raw reads were trimmed for quality and length with Trimmomatic, version 0.38[62]. Reads

were aligned against the reference grapevine genome PN40024 12X2[63], using the software

Hisat2, version 2.1.0[64] yielding an average of 31.6 M sequence per sample (Table S6). Aligned

reads were counted using the last available annotation VCost.v3 with HTSeq-count (version

0.9.1) with the "nonunique all" flag[65]. Genes were filtered by applying an RPKM>1 cut-off in at

least one experimental condition (Table S4), and the variance stabilising transformation was

applied for data visualisation.

Genes (TMM normalised) were tested for multi time-series significance for finding genes

with significant ($P<0.05$) temporal expression changes using the R package MaSigPro[66], with

Syrah dataset used as a control. A quadratic regression model (rqs=0.7) was applied. Significant

genes over time were clustered using STEM (short time-series expression miner) software[67] with

the maximum number of model profiles set to 24. Filtering or normalisation was not applied by

the software since input data were already filtered and normalised, but a mean-centred scale was

applied to overcome cluster mismatches due to different expression levels. Genes showing the

same profiles pattern among the significant clusters were grouped as follows: cluster A = cluster

7 & 9; cluster B = cluster 16, 17 & 14; cluster C = cluster 8; cluster D = cluster 15, (Fig. S3).

Venn diagrams were drawn with the R package VennDiagram. GO term profiler was retrieved

from the web tool gProfiler by applying a significant p-value of 0.001.

Pairwise differential expressed genes (FDR < 0.05) analysis was performed with the R

package DeSeq2[68] testing S/G samples. As before, genes were filtered by RPKM>1.

In silico functional CREs analysis

Promoter sequences (1.5 kb upstream of TSS) were extracted using bedtools from the

12X2 genome assembly[63]. Gene correspondence file at

https://urgi.versailles.inra.fr/content/download/5723/43038/file/list_genes_vitis_correspondences

V3_1.xlsx and in the related gff3 shows that the 5'UTR was frequently lost in the VCost.v3

annotation. Therefore, we decided to take the longest sequence between VCost.v3 and NCBI

(Vitvi and LOC references, respectively) as the most probable TSS. *Cis*-regulatory elements

(CREs) from a selected list of genes involved in the stop of the phloem were searched for an

exact match based on the DAP-Seq motif library[69] with HOMER[70]. For each gene, each CRE

was normalised for positional bias score by calculating the Z-score, which considers the motifs

length, its frequency and position[29].

Data Availability

Acknowledgements

We would like to thank the Poupelain Foundation, the Comité Interprofessionnel des Vins de

Bordeaux (CIVB) and the Agence Nationale de la Recherche (G2WAS project, ANR-19-CE20-

0024) for providing the financial support of this study. Thanks to Marc Farnos for plant

management, Sylvain Santoni and Muriel Latreille for RNA library preparation, Gauthier Sarah

for bioinformatic support and Yannick Sire for cation analysis.

Conflict of interests

The authors declare no conflict of interest.

Contributions

SS monitored and collected the grape samples, extracted the RNA and the metabolites,

performed the data analysis, interpreted the results, and drafted the manuscript. LT provided the

microvine lines and supervised the experiments, interpreted the results, and edited the

manuscript. CR conceived and designed the study, coordinated and supervised the experiments,

interpreted the results, and drafted the manuscript. All authors read and approved the final

version of the manuscript.

References

1.   Nevo, O., Razafimandimby, D., Jeffrey, J. A. J., Schulz, S. & Ayasse, M. Fruit scent as an

evolved signal to primate seed dispersal. *Sci. Adv.* **4**, eaat4871 (2018).

2. Colombié, S. *et al.* Respiration climacteric in tomato fruits elucidated by constraint-based
modelling. *New Phytol.* **213**, 1726–1739 (2017).

3. Keller, M. & Shrestha, P. M. Solute accumulation differs in the vacuoles and apoplast of
ripening grape berries. *Planta* **239**, 633–642 (2014).

4. Shahood, R., Torregrosa, L., Savoi, S. & Romieu, C. First quantitative assessment of growth,
sugar accumulation and malate breakdown in a single ripening berry. *OENO One* **54**, 1077–
1092 (2020).

5. Nieves☐Cordones, M. *et al.* Characterization of the grapevine Shaker K+ channel VvK3.1
supports its function in massive potassium fluxes necessary for berry potassium loading and
pulvinus-actuated leaf movements. *New Phytol.* **222**, 286–300 (2019).

6. Massonnet, M. *et al.* Ripening transcriptomic program in red and white grapevine varieties
correlates with berry skin anthocyanin accumulation. *Plant Physiol.* pp.00311.2017 (2017)
doi:10.1104/pp.17.00311.

7. Fasoli, M. *et al.* Timing and order of the molecular events marking the onset of berry
ripening in grapevine. *Plant Physiol.* **178**, 1187–1206 (2018).

8. Guillaumie, S. *et al.* Transcriptional analysis of late ripening stages of grapevine berry. *BMC*
Plant Biol. **11**, 165 (2011).

9. Cramer, G. R. *et al.* Transcriptomic analysis of the late stages of grapevine (*Vitis vinifera* cv.
Cabernet Sauvignon) berry ripening reveals significant induction of ethylene signaling and
flavor pathways in the skin. *BMC Plant Biol.* **14**, 1–21 (2014).

10. Ghan, R. *et al.* The common transcriptional subnetworks of the grape berry skin in the late
stages of ripening. *BMC Plant Biol.* **17**, (2017).

11. Cramer, G. R., Cochetel, N., Ghan, R., Destrac-Irvine, A. & Delrot, S. A sense of place:
transcriptomics identifies environmental signatures in Cabernet Sauvignon berry skins in the
late stages of ripening. *BMC Plant Biol.* **20**, 41 (2020).

12. Palumbo, M. C. *et al.* Integrated network analysis identifies fight-club nodes as a class of
hubs encompassing key putative switch genes that induce major transcriptome
reprogramming during grapevine development. *Plant Cell* **26**, 4617–4635 (2014).

13. Coombe, B. G. Research on development and ripening of the grape berry. *Am. J. Enol. Vitic.*
43, 101–110 (1992).

14. Zhang, X. Y. *et al.* A shift of phloem unloading from symplasmic to apoplasmic pathway is
involved in developmental onset of ripening in grape berry. *Plant Physiol.* **142**, 220–232

(2006).

15. Coombe, B. G. & McCarthy, M. G. Dynamics of grape berry growth and physiology of

ripening. *Aust. J. Grape Wine Res.* **6**, 131–135 (2000).

16. Zhang, Y. & Keller, M. Discharge of surplus phloem water may be required for normal

grape ripening. *J. Exp. Bot.* **68**, 585–595 (2017).

17. Bigard, A. *et al.* The kinetics of grape ripening revisited through berry density sorting.

OENO One **53**, (2019).

18. Terrier, N., Ageorges, A., Abbal, P. & Romieu, C. Generation of ESTs from grape berry at

various developmental stages. *J. Plant Physiol.* **158**, 1575–1583 (2001).

19. Lund, S. T., Peng, F. Y., Nayar, T., Reid, K. E. & Schlosser, J. Gene expression analyses in

individual grape (*Vitis vinifera* L.) berries during ripening initiation reveal that pigmentation

intensity is a valid indicator of developmental staging within the cluster. *Plant Mol. Biol.* **68**,

301–315 (2008).

20. Gouthu, S. *et al.* A comparative study of ripening among berries of the grape cluster reveals

an altered transcriptional programme and enhanced ripening rate in delayed berries. *J. Exp.*

Bot. **65**, 5889–5902 (2014).

21. Carbonell-Bejerano, P. *et al.* Reducing sampling bias in molecular studies of grapevine fruit

ripening: transcriptomic assessment of the density sorting method. *Theor. Exp. Plant*

Physiol. **28**, 109–129 (2016).

22. Rienth, M. *et al.* Day and night heat stress trigger different transcriptomic responses in green

and ripening grapevine (*Vitis vinifera*) fruit. *BMC Plant Biol.* **14**, 108 (2014).

23. Rienth, M. *et al.* Temperature desynchronizes sugar and organic acid metabolism in ripening

grapevine fruits and remodels their transcriptome. *BMC Plant Biol.* **16**, 164 (2016).

24. Ghaffari, S., Reynard, J. S. & Rienth, M. Single berry reconstitution prior to RNA-

sequencing reveals novel insights into transcriptomic remodeling by leafroll virus infections

in grapevines. *Sci. Rep.* **10**, 12905 (2020).

25. Fasoli, M. *et al.* The grapevine expression atlas reveals a deep transcriptome shift driving the

entire plant into a maturation program. *Plant Cell* **24**, 3489–3505 (2012).

26. Bigard, A., Romieu, C., Sire, Y. & Torregrosa, L. *Vitis vinifera* L. diversity for cations and

acidity is suitable for breeding fruits coping with climate warming. *Front. Plant Sci.* **11**,

(2020).

27. Dimopoulos, N. *et al.* Drought stress modulates cuticular wax composition of the grape

berry. *J. Exp. Bot.* **71**, 3126–3141 (2020).

28. Frumkin, I. *et al.* Gene architectures that minimize cost of gene expression. *Mol. Cell* **65**,

142–153 (2017).

29. Ma, S., Shah, S., Bohnert, H. J., Snyder, M. & Dinesh-Kumar, S. P. Incorporating motif

analysis into gene co-expression networks reveals novel modular expression pattern and new

signaling pathways. *PLoS Genet* **9**, e1003840 (2013).

30. Torregrosa, L. J.-M., Rienth, M., Romieu, C. & Pellegrino, A. The microvine, a model for

studies in grapevine physiology and genetics. *OENO One* **53**, (2019).

31. Tilbrook, J. & Tyerman, S. D. Cell death in grape berries: varietal differences linked to

xylem pressure and berry weight loss. *Funct. Plant Biol.* **35**, 173–184 (2008).

32. Dal Santo, S. *et al.* Genome-wide analysis of the expansin gene superfamily reveals

grapevine-specific structural and functional characteristics. *PLOS ONE* **8**, e62206 (2013).

33. Schlosser, J. *et al.* Cellular expansion and gene expression in the developing grape (*Vitis*

vinifera L.). *Protoplasma* **232**, 255 (2008).

34. Wong, D. C. J., Zhang, L., Merlin, I., Castellarin, S. D. & Gambetta, G. A. Structure and

transcriptional regulation of the major intrinsic protein gene family in grapevine. *BMC*

Genomics **19**, 248 (2018).

35. Stanfield, R. C., Hacke, U. G. & Laur, J. Are phloem sieve tubes leaky conduits supported

by numerous aquaporins? *Am. J. Bot.* **104**, 719–732 (2017).

36. Zhang, Z. *et al. VvSWEET10* mediates sugar accumulation in grapes. *Genes* **10**, 255 (2019).

37. Chen, L.-Q. *et al.* Sucrose efflux mediated by SWEET proteins as a key step for phloem

transport. *Science* **335**, 207–211 (2012).

38. Ren, R. *et al.* Coexpression of sucrose synthase and the SWEET transporter, which are

associated with sugar hydrolysis and transport, respectively, increases the hexose content in

Vitis vinifera L. grape berries. *Front. Plant Sci.* **11**, (2020).

39. Terrier, N. *et al.* Isogene specific oligo arrays reveal multifaceted changes in gene expression

during grape berry (*Vitis vinifera* L.) development. *Planta* **222**, 832–847 (2005).

40. Jaillon, O. *et al.* The grapevine genome sequence suggests ancestral hexaploidization in

major angiosperm phyla. *Nature* **449**, 463–467 (2007).

41. Afoufa-Bastien, D. *et al.* The *Vitis vinifera* sugar transporter gene family: phylogenetic

overview and macroarray expression profiling. *BMC Plant Biol.* **10**, 245 (2010).

42. Murcia, G., Pontin, M. & Piccoli, P. Role of ABA and Gibberellin A_3 on gene expression

pattern of sugar transporters and invertases in *Vitis vinifera* cv. Malbec during berry

ripening. *Plant Growth Regul.* **84**, 275–283 (2018).

43. Kuang, L., Chen, S., Guo, Y. & Ma, H. Quantitative proteome analysis reveals changes in

the protein landscape during grape berry development with a focus on vacuolar transport

proteins. *Front. Plant Sci.* **10**, (2019).

44. Schulz, A. *et al.* Proton-driven sucrose symport and antiport are provided by the vacuolar

transporters *SUC4* and *TMT1/2*. *Plant J.* **68**, 129–136 (2011).

45. Wei, X., Liu, F., Chen, C., Ma, F. & Li, M. The *Malus domestica* sugar transporter gene

family: identifications based on genome and expression profiling related to the accumulation

of fruit sugars. *Front. Plant Sci.* **5**, (2014).

46. Jung, B. *et al.* Identification of the transporter responsible for sucrose accumulation in sugar

beet taproots. *Nat. Plants* **1**, 1–6 (2015).

47. Cheng, J. *et al.* Overexpression of the tonoplast sugar transporter *CmTST2* in melon fruit

increases sugar accumulation. *J. Exp. Bot.* **69**, 511–523 (2018).

48. Savoi, S., Herrera, J. C., Forneck, A. & Griesser, M. Transcriptomics of the grape berry

shrivel ripening disorder. *Plant Mol. Biol.* **100**, 285–301 (2019).

49. Zeng, L., Wang, Z., Vainstein, A., Chen, S. & Ma, H. Cloning, localization, and expression

analysis of a new tonoplast monosaccharide transporter from *Vitis vinifera* L. *J. Plant*

Growth Regul. **30**, 199–212 (2011).

50. Fillion, L. *et al.* Cloning and expression of a hexose transporter gene expressed during the

ripening of grape berry. *Plant Physiol.* **120**, 1083–1094 (1999).

51. Lijavetzky, D. *et al.* Berry flesh and skin ripening features in *Vitis vinifera* as assessed by

transcriptional profiling. *PLoS ONE* **7**, e39547 (2012).

52. Murcia, G. *et al.* ABA and GA$_3$ increase carbon allocation in different organs of grapevine

plants by inducing accumulation of non-structural carbohydrates in leaves, enhancement of

phloem area and expression of sugar transporters. *Physiol. Plant.* **156**, 323–337 (2016).

53. Zenoni, S., D'Incà, E. & Tornielli, G. B. Genetic dissection of grape berry ripening control:

defining a role for NAC transcription factors. *Acta Hortic.* 387–402 (2019).

54. Huang, T., Yang, J., Yu, D., Han, X. & Wang, X. Bioinformatics analysis of WRKY

transcription factors in grape and their potential roles prediction in sugar and abscisic acid

signaling pathway. *J. Plant Biochem. Biotechnol.* (2020) doi:10.1007/s13562-020-00571-y.

55. Grimplet, J., Pimentel, D., Agudelo-Romero, P., Martinez-Zapater, J. M. & Fortes, A. M.

The LATERAL ORGAN BOUNDARIES Domain gene family in grapevine: genome-wide

characterization and expression analyses during developmental processes and stress

responses. *Sci. Rep.* **7**, 15968 (2017).

56. Agudelo-Romero, P. *et al.* Search for transcriptional and metabolic markers of grape pre-

ripening and ripening and insights into specific aroma development in three Portuguese

cultivars. *PLoS ONE* **8**, e60422 (2013).

57. Sarry, J.-E. *et al.* Grape berry biochemistry revisited upon proteomic analysis of the

mesocarp. *Proteomics* **4**, 201–215 (2004).

58. Terrier, N., Sauvage, F.-X., Ageorges, A. & Romieu, C. Changes in acidity and in proton

transport at the tonoplast of grape berries during development. *Planta* **213**, 20–28 (2001).

59. Chaïb, J. *et al.* The grape microvine – a model system for rapid forward and reverse genetics

of grapevines. *Plant J.* **62**, 1083–1092 (2010).

60. Escudier, J.-L. *et al.* De la vigne au vin□: des créations variétales adaptées au changement

climatique et résistant aux maladies cryptogamiques Partie 1/2□: La résistance variétale.

Rev. Oenologues Tech. Vitivinic. Oenologiques 16–18 (2017).

61. Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image

analysis. *Nat. Methods* **9**, 671–675 (2012).

62. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

63. Canaguier, A. *et al.* A new version of the grapevine reference genome assembly (12X.v2)

and of its annotation (VCost.v3). *Genomics Data* **14**, 56–62 (2017).

64. Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low memory

requirements. *Nat. Methods* **12**, 357–360 (2015).

65. Anders, S., Pyl, P. T. & Huber, W. HTSeq—a Python framework to work with high-

throughput sequencing data. *Bioinformatics* **31**, 166–169 (2015).

66. Nueda, M. J., Tarazona, S. & Conesa, A. Next maSigPro: updating maSigPro bioconductor

package for RNA-Seq time series. *Bioinformatics* **30**, 2598–2602 (2014).

67. Ernst, J. & Bar-Joseph, Z. STEM: a tool for the analysis of short time series gene expression

data. *BMC Bioinformatics* **7**, 191 (2006).

68. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

69. O'Malley, R. C. *et al.* Cistrome and epicistrome features shape the regulatory DNA

landscape. *Cell* **165**, 1280–1292 (2016).

70. Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime cis-

regulatory elements required for macrophage and B cell identities. *Mol. Cell* **38**, 576–589

(2010).

Figure legends

Figure 1. Growth and solute accumulation during the ripening of synchronised grapevine single

berries of MV032, MV102, and Syrah. (a) Relative growth, (b) glucose + fructose, (c) potassium,

(d) malic acid, and (e) tartaric acid. Results in b and c were normalised in order to consider 1 g of

pericarp at phloem arrest. The developmental stage is indicated by colours: V for veraison in

green, G for growing in yellow, P for peak in pink, S for shrivelling in purple. Letters indicate

significant differences according to the Tukey ad hoc test.

Figure 2. Overview of the transcriptomic results: (a) principal component analysis, and (b)

sample dendrogram. Colours and symbols indicate genotypes and stages, respectively: MV032 in

yellow, MV102 in red, Syrah in green; a circle for stage G, a square for P, and a triangle for S.

Figure 3. Significant time-course genes expression of (a) cluster A and B representing genes

increasing or decreasing respectively in expression at phloem arrest. (b) Venn diagram showing

genes displaying common or genotype specific changes in expression at phloem arrest. (c)

Significantly enriched GO categories in modulated genes, divided in biological process (BP),

cellular component (CC), and molecular function (MF).

Figure 4 Genes inhibited at phloem stop belong to different regulatory networks. Genes and

transcription factors were clustered according to the Z-score of the motifs present within 1500 bp

before the TSS. Red and blue colours indicate proximity or distance of the promoter sequence to

the TSS. White colour denotes the absence of a specific TF in the promoter of a gene. Horizontal

and vertical colour bars grouped promoter and genes in promoter families and gene functions as

indicated in the legend. The two examples discussed in the text regarding SWEETs and HT6

with reference to NAC, MYB, and AP2/EREB are marked by black squares.

Figure 5 Hypothetical structure of the energy-efficient apoplastic pathway in the ripening berry.

The figure is divided in three panels: early ripening, late ripening, and shrivelling with the stop

of phloem between the two latter (marked by a black circle). Genes written in red or blue italics

refer to those respectively expressed or repressed at specific stages, those in black are

constitutive.

Table 1 List of thirty-five selected genes related to cell growth, water and sugar transport sorted

by absolute difference in Syrah G versus P. Gene ID is reported as VCost.V3 and RefSeq code.

Gene expression is expressed as count per million (CPM).

Supplementary materials legend

Figure S1. Meteorological data showing temperature max and min recorded in the greenhouse

(dotted lines) and in the field (solid line) during the grape growing season. Syrah sampling dates

(V, G, P, and S) based on Julian days are reported on the top.

Figure S2. Glucose plus fructose (mM) in relation to malic acid (mEq) in Syrah, MV032, and

MV102. Different shapes denote different developmental stages: circle, square and triangle

represent stage G, P, and S. Coloured points in red, yellow and green represent the samples

selected for RNA-Seq analysis in G, P, and S.

Figure S3. Overview of the model profiles interface (from the STEM software) where each box

corresponds to a model expression profile, manually set to 24. Red lines show genes distribution.

Coloured profiles have a significant number of genes assigned and therefore were considered for

the following analysis.

Figure S4. Significant time-course genes expression of clusters C and D representing genes up

or down peaking respectively in expression at phloem arrest and Venn diagram showing genes

displaying common or cultivar specific changes in expression at phloem arrest.

Figure S5. Subcellular localization of *VviHT6* proteins to the tonoplast after stable expression in

grapevine hairy roots. The complete ORF of the *VviHT6* gene without the stop codon was

amplified from Syrah cDNA (ripening berry) using the following primer pair: HT6for

CACCATGAACGGAGCTGTGCT, HT6rev GTCATTCTTTGCTGCAGTAAC before blunt

end topo cloning, transfer to the pH7FWG2 vector and stable transformation of grapevine with

agrobacterium rhyzogenes A4 according to Gomez et al., 2009.

Table S1. Evolution of sugars, acid and cations. (a) Amounts per fruit (concentration x volume);

(b) concentrations and (c) net H^+/sucrose exchange escaping V-ATPase at the tonoplast**.**

Table S2. List of genes displaying time significant expression ($P<0.05$) in the three genotypes.

(a) Time significant gene identified with MaSigPro (G-P-S); (b) pairwise significant gene

identified with DeSeq2 (S/G).

Table S3. Merged list of commonly regulated genes ranked by absolute difference in CPM

expression between G and S.

Table S4. Genes that passed the filtering threshold (RPKM>1) in at least one group of sample

triplicates.

Table S5. Exemplary calculation of cytosolic sucrose at equilibrium.

Table S6. RNA-Sequencing data metrics.