Title: Shape analysis of gamma rhythm supports a superlinear inhibitory regime in an inhibition-stabilized network

Running title: Shape analysis of gamma rhythm

Authors: R (first name) Krishnakumaran (last name)[1], Mohammed (first name) Raees (last name)[2] and Supratim (first name) Ray (last name)[1,2,*].

Affiliations:

[1]IISc Mathematics Initiative, Department of Mathematics, Indian Institute of Science, Bangalore, India, 560012

[2]Centre for Neuroscience, Indian Institute of Science, Bangalore, India, 560012, Telephone +91 80 2293 3437, Facsimile +91 80 2360 3323

[*]**Corresponding Author:** sray@iisc.ac.in

Keywords: Gamma, Harmonics, Wilson-Cowan model, Phase analysis

Conflict of interest: The authors declare no competing financial interests.

Funding Disclosure: This work was supported by Wellcome Trust/DBT India Alliance (Senior fellowship IA/S/18/2/504003 to SR) and DBT-IISc Partnership Programme.

Acknowledgements: We thank Dr. Adam Kohn for his helpful comments on the paper.

Abstract

Visual inspection of stimulus-induced gamma oscillations (30-70 Hz) often reveals a non-sinusoidal shape. Such distortions are a hallmark of non-linear systems and are also observed in mean-field models of gamma oscillations. A thorough characterization of the shape of the gamma cycle can therefore provide additional constraints on the operating regime of such models. However, the gamma waveform has not been quantitatively characterized, partially because the first harmonic of gamma, which arises because of the non-sinusoidal nature of the signal, is typically weak and gets masked due to a broadband increase in power related to spiking. To address this, we recorded spikes and local field potential (LFP) from the primary visual cortex (V1) of two awake female macaques while presenting full-field gratings or iso-luminant chromatic hues that produced huge gamma oscillations with prominent peaks at harmonic frequencies in the power spectra. We found that gamma and its first harmonic always maintained a specific phase relationship, resulting in a distinctive shape with a sharp trough and a shallow peak. Interestingly, a Wilson-Cowan (WC) model operating in an inhibition stabilized mode could replicate the findings, but only when the inhibitory population operated in the super-linear regime, as predicted recently. However, another recently developed model of gamma that operates in a linear regime driven by stochastic noise failed to produce salient harmonics or the observed shape. Our results impose additional constraints on models that generate gamma oscillations and their operating regimes.

Significance Statement

Gamma rhythm is not sinusoidal. Understanding these distortions could provide clues about the cortical network that generates the rhythm. Here, we use harmonic phase analysis to describe these waveforms quantitatively, and show that the gamma rhythm in macaque V1, during the presentation of fullscreen plain-hues and achromatic-gratings, has a signature arch-shaped waveform, despite the variation in power and frequency reported earlier. We further demonstrate using population rate models that the non-sinusoidal waveform is dependent on the operating domain of the system generating it. Consequently, shape analysis provides additional constraints on cortical models and their operating regimes.

Introduction

Gamma rhythm refers to oscillatory neural activity in the 30-70 Hz range that changes in response to different stimuli and cognitive states (Buzsaki, 2006). In the primary visual cortex (V1), the gamma rhythm has been studied extensively using achromatic gratings, wherein gamma power and center-frequency have been shown to vary systematically with the properties of the grating (Jia et al., 2013; Murty et al., 2018). For instance, peak-frequency of gamma increases with the contrast of gratings (Ray and Maunsell, 2010; Jia et al., 2013) and, gamma power increases and frequency decreases with stimulus size (Gieselmann and Thiele, 2008; Ray and Maunsell, 2011). More recently, chromatic stimuli have also been explored, which, for low-wavelength (reddish) hues, can generate huge gamma oscillations that are an order of magnitude stronger than gamma produced by achromatic gratings (Shirhatti and Ray, 2018; Bartoli et al., 2019; Peter et al., 2019).

Gamma oscillations are thought to reflect the push-pull activity of interconnected excitatory and inhibitory neurons, which has been demonstrated in different spiking network models (Tiesinga and Sejnowski, 2009; Mazzoni et al., 2015; Chariker et al., 2018; Keeley et al., 2019). However, since such large-scale network models have several parameters to be tuned and can become hard to interpret, simplified population rate models are often used. One pioneering model was proposed by Wilson and Cowan (WC; 1972), in which excitatory and inhibitory neurons were grouped into populations, and the dynamics of these populations were characterized. Variants of WC models have recently been used to explain some properties of the gamma rhythm. For example, based on the observation that gamma rhythm appears in short bursts and its phase does not vary linearly (Burns et al., 2010, 2011; Xing et al., 2012), a WC based model was proposed in which the activation function (input-output relationship) was linear but driven by Poisson noise (Kang et al., 2009; Xing et al., 2012). A variant of this model, in which an additional global excitatory population was added, was used to explain the stimulus

dependence of gamma oscillations (Jia et al., 2013). On the other hand, Jadi and Sejnowski

(2014) used a WC model with a non-linear (sigmoidal) activation function and showed that

constraining the model to operate in an inhibition stabilized mode (Tsodyks et al., 1997) and

the inhibitory population to operate in superlinear domain can also reproduce the size and

contrast dependence of gamma rhythm.

Although both these models can explain the stimulus dependence of gamma, the shape

of the rhythm likely depends on the presence (and type) of non-linearity. The shape could be

non-sinusoidal (see, for example, Figure 11 of Wilson and Cowan, 1972), which is represented

in the spectral domain as peaks at harmonics of the fundamental frequency. Such distortions

have been observed in various brain oscillations such as theta rhythm (for a review, see Cole

and Voytek (2017)). For gamma rhythm, a visual inspection of raw traces reveals some

distortion (Henrie and Shapley, 2005; Jia and Kohn, 2011; Jia et al., 2011; Ray and Maunsell,

2015), which is also corroborated by the presence of peaks at the first harmonic of the

fundamental gamma peak in the power spectra (Gieselmann and Thiele, 2008; Ray and

Maunsell, 2010; Shirhatti and Ray, 2018). However, a quantitative study of the gamma

waveform, which could potentially constrain the type and operating regime of models, has not

been undertaken. This is partly because the noise in the harmonic range could offset its phase

estimates if the harmonic is not prominent enough. To address this, we studied gamma

oscillations produced by presenting full-screen hues that generated salient gamma oscillations

with prominent harmonics (Shirhatti and Ray, 2018), characterized the shape, and tested

models that could replicate the shape.

Materials and Methods

<u>Data Acquisition</u>

We used LFP data recorded by Shirhatti and Ray (2018) from V1 of two female macaque subjects, whom we refer to as M1 and M2 (correspond to M1 and M3 in the earlier paper). Both monkeys were fitted with a titanium headpost and trained to perform a visual passive fixation task, after which a Utah array (96 and 81 electrodes for M1 and M2) was implanted in V1 (details of the surgery and implants are provided in Shirhatti and Ray (2018)). The raw signals from microelectrodes were recorded using the 128-channel Cerebus neural signal processor (Blackrock Microsystems). LFP was obtained by filtering the raw signals online between 0.3 Hz and 500 Hz (Butterworth filters; first-order analog and fourth-order digital respectively), and recorded at 2 kHz sampling rate and 16-bit resolution. No further offline filtering was performed on this data before analysis.

Multiunit activity was also extracted from the raw signal by filtering online between 250 Hz and 7.5 kHz (Butterworth filters; fourth-order digital and third-order analog respectively), and subjecting the resultant signal to an amplitude threshold of ~5 SD of the signal. The recorded units were found to have receptive fields located in the lower left quadrant of the visual field with respect to fixation in both monkeys, at an eccentricity of ~3°–4.5° in M1 and ~3.5°– 4.5° in M2. Full-field iso-luminant hues did not drive the neurons well, and therefore we did not get usable spiking activity from most electrodes. Therefore, as in our previous report (Shirhatti and Ray, 2018), all the analyses were performed only using LFP data (see Supplementary Figure 1 of Shirhatti and Ray, 2018, for description of spiking activity).

<u>Experimental setup and behavior</u>

During the experiment, the monkey was seated in a monkey chair with its head held stationary by the headpost. The monkey viewed a monitor (BenQ XL2411, LCD, 1,280 × 720 resolution, 100 Hz refresh rate) placed ~50 cm from its eyes. The monkey and the display setup

were housed in a Faraday enclosure with a dedicated grounding separate from the main supply

ground to provide isolation from external electrical noise. The monitor was calibrated and

gamma-corrected using i1Display Pro (x-rite PANTONE) to obtain a mean luminance of 60

cd/m^2 on its surface and to obtain a gamma of unity for each of the three primaries of the color

gamut, which had the following CIE chromaticity xy coordinates: red, (0.644, 0.331); green,

(0.327, 0.607); blue, (0.160, 0.062). The white point was at (0.345, 0.358).

Each monkey performed a passive fixation task, which required them to fixate at a small

dot of 0.05°–0.10° radius at the center of the screen throughout the trial (3.3 or 4.8 s duration;

fixation spot was displayed throughout). Each trial began with fixation, following which an

initial blank grey screen of 1,000 ms was displayed, and then, two to three stimuli were shown

for 800 ms each with an interstimulus interval of 700 ms. The monkey was rewarded with juice

for maintaining fixation within 2° from the fixation point. Trials in which fixation was broken

were not considered in our analyses. Eye position data was recorded as horizontal and vertical

coordinates using the ETL-200 Primate Eye Tracking System (ISCAN) and monitored

throughout the task using custom software running on macOS, which also controlled the task

flow, generated stimuli, and randomized stimuli presentation.

<u>Stimuli</u>

The stimuli consisted of 36 hues and 1 achromatic grating. The hues were equally

spaced along the circular hue space of the standard HSV nomenclature (0° hue to 350° hue,

where 0°, 120°, and 240° represent red, green, and blue respectively), which were displayed

full screen and at full saturation and value. The achromatic grating was at an orientation of 90°

and had a spatial frequency of 4 cpd for M1 and 2 cpd for M2. These grating parameters were

optimized to capture strong fast gamma with minimal slow gamma (see Murty et al., 2018).

The full-screen stimuli, in our setup, subtended a visual angle of ∼56° in the horizontal

direction and ∼33° in the vertical direction.

<u>Electrode selection</u>

As with our previous report (Shirhatti and Ray, 2018), electrodes were considered for

analysis only if they gave consistent stimulus-induced changes and reliable receptive field

estimates across sessions, determined by a receptive field mapping protocol that was run across

multiple days (Dubey and Ray, 2019). Further, we discarded signals from electrodes with

unusable or inconsistent signals, a high degree of crosstalk with other electrodes, or impedances

outside the range of 250–2,500 KΩ for monkey M1 and 125–2,500 KΩ for M2. This resulted

in 64 and 16 usable electrodes for M1 and M2, respectively.

<u>Data Analysis</u>

Stimulus presentations with excessive artifacts (<5.9% and <5.0% of presentations of

each stimulus in M1 and M2) were discarded for each session, yielding 19.9±6.2 repeats in M1

and 20.2±1.0 repeats in M2 per stimulus.

<u>Spectral analysis of LFP</u>

For each stimulus, LFP recorded from -500 to 0 ms from stimulus onset was taken as

the 'baseline period' and 250 to 750 ms from stimulus onset was taken as the 'stimulus period'

to avoid the transient responses to the stimulus onset. This yielded a frequency resolution of 2

Hz in the Power Spectral Density (PSD). PSD was computed using the Multitaper method

using the Chronux toolbox (Bokil et al., 2010), with three tapers. The change in power was

calculated as 10 times the difference between base-10 logarithm of PSDs at stimulus period

and baseline period, expressed in decibels (dB). Estimation of peak frequencies was done on

these baseline-corrected PSDs.

Gamma range was taken as 30-70 Hz, and 'gamma peak frequency' was estimated as

the highest peak within this range. In most stimulus conditions tested, a discernible 'second

bump' was observed in the baseline-corrected PSD. The peak frequency of the second bump

was estimated as the highest peak occurring beyond 12 Hz after the estimated gamma peak

frequency up to 140 Hz, to exclude higher frequency bumps.

<u>Analysis of gamma and harmonic phases</u>

To compute the phase difference of gamma and its first harmonic, gamma and harmonic

signals were extracted from LFP during the stimulus period by bandpass filtering using separate

Butterworth filters (zero-phase; order 4). The passband for gamma was 20 Hz wide, centered

around the gamma peak frequency, identified from trial-averaged PSD for a given stimulus and

an electrode. The passband for the first harmonic of gamma was also 20 Hz wide but centered

around twice the corresponding gamma peak frequency. The phases of these signals (φ_{gamma}

and $\varphi_{harmonic}$) were then computed using Hilbert transform. The phase difference between

gamma and its harmonic was calculated as:

Phase difference $= |2*\varphi_{gamma} - \varphi_{harmonic}|$ 	(1)

The above phase difference estimate was computed at each timepoint within the

stimulus period of every trial for each electrode.

Effect of harmonic phase on gamma waveform

To illustrate the effect of the gamma-harmonic phase relationship on the shape of

gamma waveform, gamma and its first harmonic were mimicked using sinusoids and the initial

phase of gamma (φ_G) was varied. The formulation of gamma and harmonics were as follows:

Gamma wave, $G = \cos(2\pi\ 45\ t + \varphi_G)$

Harmonic wave, $H = \frac{1}{4}*\cos(2\pi\ 90\ t)$

Sum of gamma and harmonic = G + H   (2)

These waveforms are displayed in Figure 4C along with the phase difference (Figure

4D) obtained by applying equation (1).

<u>Experimental Design and Statistical Analysis</u>

First, we tested our hypothesis that the second bump in PSD accompanying gamma was

indeed its harmonic by computing the ratio of harmonic to gamma peak-frequencies, obtained

from the trial-averaged change in power (dB) from baseline spectra for each electrode. The

ratios were subsequently subjected to a non-parametric Wilcoxon signed rank test (Null

hypothesis: median ratio = 2). The standard error (SE) of the median of the gamma-harmonic

frequency ratio was estimated by bootstrapping over N iterations (where N is the number of

datapoints). This involved random sampling with replacement of the ratio data N times and

estimating their median each time, which resulted in N medians, whose standard deviation (SD)

is reported as the standard error (SE).

Circular statistics on the gamma-harmonic phase difference data were computed in

Matlab using the Circular Statistics toolbox (Berens, 2009). The mean phase differences are

reported as MEAN ± CI in Figure 5, where MEAN is the circular mean and CI is the 95%

confidence interval of MEAN, under a Von Mises distribution (implemented in *circ_mean* and

circ_confmean functions of the Circular Statistics toolbox).

To validate that the distribution of gamma-harmonic phase differences was non-

uniform and, thus, assess the validity of circular mean estimates in Figure 5, we subjected trial-

averaged phase-differences gathered from all electrodes for a given stimulus to a Rayleigh test

of non-uniformity (Null hypothesis: uniform distribution of phases). When the gamma activity

recorded for a given stimulus has a specific waveform across electrodes, the corresponding

distribution of gamma-harmonic phase-differences will be unimodal. We considered this to be

the case when the p-value of the Rayleigh test was less than 0.01.

<u>Studying non-sinusoidal waveforms in mean-field models of gamma</u>

To understand the factors resulting in the characteristic waveform of gamma, we

investigated the gamma-harmonic phase relationship emerging in mean-field models

introduced in earlier works, which produced gamma oscillations with power and frequency

trends as observed in experiments. We explored the waveform in these models by identifying

input regimes that produced a first-harmonic with a specific phase difference from the

fundamental of gamma, as required for its arch-shape. We assessed the presence of such

regimes in two models. The first model operated linearly and produced oscillations by virtue

of stochastic time-varying inputs (Jia-Xing-Kohn or JXK model; Jia et al., 2013), while the

second model had non-linear dynamics, which gave rise to gamma frequency limit cycles in

response to constant inputs (Jadi and Sejnowski or JS model; Jadi and Sejnowski, 2014).

<u>Jia-Xing-Kohn (JXK) model</u>

Jia et al. (2013) defined a linear EI rate-model and extended it by adding a Global

Excitatory Population (G) to approximate recurrent excitatory feedback within V1. The model,

when subjected to step input, acts as a damped oscillator but produces oscillations by virtue of

constant perturbation from the Poisson inputs. The original model, formulated in the paper,

incorporates detailed stimulus descriptions, namely Masked Noise (MN) level and stimulus

size (r). The stimulus size parameter (r) scales the extent of Network recurrence in steady-state

proportionally as larger size stimuli excite a larger area of cortical cells, increasing the global

feedback. In our simulations, the Masking Noise level (MN) has been set to 0, resulting in the

following reduced formulation.

$\tau_E \dfrac{dr_E}{dt} = -r_E + W_{EE}\lfloor r_E \rfloor - W_{EI}\lfloor r_I \rfloor + W_{EG}\lfloor r_G \rfloor + E_{inp}$

$\quad \tau_I \dfrac{dr_I}{dt} = -r_I + W_{IE}\lfloor r_E \rfloor - W_{II}\lfloor r_I \rfloor + W_{IG}\lfloor r_G \rfloor + I_{inp}$

$\quad \tau_G \dfrac{dr_G}{dt} = -r_G + r^2 . W_{GE}\lfloor r_E \rfloor$

where $\lfloor x \rfloor = x \; if \; x > 0 \; and \; 0 \; otherwise,$ and E_{inp} and I_{inp} approximate a Poisson process

with average rates IE and II respectively, which vary with stimulus contrast (c) as:

$\quad I_E = 40. \dfrac{c^2}{c^2 + 0.3^2}; \; I_I = 32. \dfrac{c^2}{c^2 + 0.3^2}$ (3)

Parameter values of this model are given in Table 1.

<u>Jadi-Sejnowski (JS) model</u>

Jadi and Sejnowski (2014) used a simple rate model consisting of an excitatory and an

inhibitory population with sigmoidal activation, operating as an Inhibition Stabilized Network

(ISN) and constrained the input drives to the populations to reproduce the increase in power

and decrease in peak frequency of gamma with increasing stimulus size as earlier studies have

observed in V1 (Gieselmann and Thiele, 2008; Ray and Maunsell, 2011).

The model defines the population firing rates of Excitatory and Inhibitory populations

as follows:

$\quad \tau_E \dfrac{dr_E}{dt} = -r_E + \sigma_E(W_{EE}r_E - W_{EI}r_I + I_E)$

$\quad \tau_I \dfrac{dr_I}{dt} = -r_I + \sigma_I(W_{IE}r_I - W_{II}r_I + I_I)$

$\quad \sigma_P(x) = \dfrac{1}{1+exp(\,m_P.(\theta_P - x)\,)} - \dfrac{1}{1+exp(m_P.\theta_P)}$ (4)

In this model, larger stimuli cause an increased inhibitory drive to the population, owing to

suppression from the surrounding populations. Since the model operated close to a supercritical

Hopf bifurcation, the amplitude and frequency of oscillations in firing rates could be closely

approximated by linearization of the model. The authors deduced and demonstrated that the

model gave rise to the observed trends in the gamma when the inputs were such that the

inhibitory population was strongly 'superlinear'. This means that the summed inputs to the

inhibitory population (from recurrent and external sources; argument of σ_I in equation 4) must

lie in a certain range of values where the activation function σ_I curves upwards (increasing in

slope with increasing summed input). For the sigmoidal activation function used in equation 4,

the summed inputs must operate in the lower half of the sigmoid. Superlinear activation of the

excitatory population, on the other hand, was antagonistic (not strongly superlinear). The set

of such inputs constitute the operating regime of the model, which we refer to as the

'superlinear' regime.

Identifying the operational input regimes using gamma-harmonic phase difference

Because the gamma-harmonic relationship was studied using LFP recordings in real

data while the model simulations yielded firing rates of excitatory and inhibitory populations,

these firing rates had to be converted to a 'proxy' LFP. The relationship between the two is

complex (Hasenstaub et al., 2005; Einevoll et al., 2013; Mazzoni et al., 2015). For simplicity,

we used the negative sum of population firing rates of the excitatory and inhibitory populations

in the models as a proxy for LFP, since the LFP, being the extracellular potential, could be

approximated to vary inversely with the depolarization (excitability) of these populations. We

also approximated the LFP as the negative of just the excitatory or inhibitory populations,

which yielded qualitatively similar results. We elaborate this further in the Discussion section.

The environment used for simulation was Matlab 2019b (Mathworks,

RRID:SCR_001622), where the models were simulated by a forward Euler method, with

parameters as in Table 1. Simulations were run for a duration of 2 seconds in time-steps of 0.1

ms, on all pairs of 41 excitatory and 41 inhibitory input values sampled from the ranges

specified in Table 1. In each case, PSD of the LFP proxy was computed between 1-2 seconds

to avoid the initial slow transient. The gamma and harmonic peak frequencies were identified

as frequencies containing maximum power in the 30-70 Hz range and twice the gamma peak

frequency, respectively. For further analyses in the JS model, only those input combinations

were considered for which gamma and harmonic frequency amplitudes were greater than 1e-3

and 1e-6 units respectively, to ensure phase analysis of oscillatory activity was not dominated

by simulation errors or noise (grey region in Figure 7F). The LFP proxy signal was filtered

using 20 Hz passbands centered at each frequency identified above. The gamma-harmonic

phase difference (equation 1) was computed, just as for LFP in macaque data. An input point

(pair of values supplied as excitatory and inhibitory input drives; stimulus parameters in the

case of JXK) was said to be 'in-regime' if the gamma-harmonic phase difference was within

22.5 degrees from 180-degrees (ideal for arch shape; Figure 4).

Since JXK takes stochastic inputs, to assess if there is indeed a unique gamma-harmonic phase

difference at each of the mean input-drive combinations tested and whether the model can

indeed exhibit in-regime behavior, we ran the simulation 50 times (analogous to 50 different

trials) for each stimulus condition. In each iteration, we considered the mean phase-difference

over the 1-2 second interval. For every selected input combination, the mean phase difference

over this interval in each iteration was taken and the resultant pool of 50 mean phase differences

was subjected to Rayleigh test of uniformity to identify those inputs that have a 'consistent'

gamma-harmonic phase relationship across iterations (p-value < 0.01). Among the qualifying

input combinations, those input drive pairs for which the circular mean of mean phase

differences across iterations lay within 180 ± 22.5 degrees were deemed to be 'in-regime'. For

JXK, the gamma-harmonic phase reported in Figure 6 is the circular average of the phase

differences from all the iterations. The amplitudes of gamma and its first harmonic in the figure

are estimated from the average PSD (examples in Figure 6C-D), computed as the mean of the

PSDs obtained from 50 iterations for a given stimulus combination.

Results

We collected spikes and LFP from 96 and 81 electrodes from two monkeys, M1 and M2, while they viewed full-screen color patches of different hues. Limiting our analysis to electrodes with reliable estimates of RF centers (see "Electrode selection" section in Methods) yielded 64 and 16 electrodes from the two monkeys.

<u>First harmonic of gamma oscillation</u>

In Figure 1A-D, the top row shows the power spectral density (PSD; averaged across all trials and subsequently across all electrodes) of the LFP signal in subject M1 during baseline period (-500 to 0 ms from stimulus onset; black trace) and during stimulus period (250 to 750 ms from stimulus onset; color/grey trace; the color indicates the hue that was presented and grey trace represents gratings). The corresponding change in power (dB) from baseline (Figure 1A-D bottom row) shows a prominent peak in the gamma range, along with another prominent peak near twice the frequency. Figure 1E-H show the same plots from subject M2. We first tested whether this second peak was indeed at twice the frequency. For each electrode and stimulus, we measured the gamma peak frequency from trial-averaged baseline-corrected PSDs as the highest peak within the 30-70 Hz band, and the peak of the second bump as the highest peak occurring after 12 Hz past the identified gamma peak frequency but before 140 Hz (the identified peaks are highlighted by black crosses in Figure 1).

Figure 2 presents the medians of the frequency ratios of second bump to gamma peak frequencies for each stimulus (hues indicated by color, gratings by grey color), computed across all electrodes in each subject, with error bars indicating the standard error of the median computed by bootstrapping. The stimuli are arranged along the horizontal axis by the average gamma power across electrodes. The frequency ratios were tightly clustered around 2, especially for stimuli that produced strong gamma oscillations. To test whether the ratio for a given stimulus was significantly different from 2, the frequency ratios computed in different

electrodes for the stimulus were subject to a Wilcoxon signed rank test. Stimuli for which the ratios were found to be significantly different from 2 (p < 0.01) are indicated by open circles. Estimates of peak-frequency ratios for stimuli producing less gamma power were more susceptible to noise and showed larger deviations in frequency ratios. However, frequency ratios were distributed more narrowly around 2 for stimuli inducing stronger gamma. The median ratio value for M1 was 2.00±0.021, not significantly different from 2 (p = 0.26 for M1; Wilcoxon signed rank test). For M2, the median ratio was 1.95±0.026, significantly different from 2 (p = 9.96e-6; Wilcoxon signed rank test). However, note that because we computed PSDs over a 500 ms window, the frequency resolution was 2 Hz, which introduced some error in the estimated ratio. For example, if the true gamma and harmonic peaks are at 41 and 82 Hz, the estimated gamma will either be at 40 or 42 Hz, yielding a ratio of either 82/40 = 2.05 or 82/42 = 1.95. These margins of error are shown as dotted lines in Figure 2. Most of the points lay within these error margins, especially for electrodes with higher gamma power. When we restricted the analysis to the top ten hues for each monkey in terms of gamma power, the ratios were 1.98±0.046 (p=1; Wilcoxon signed rank test) and 1.99±0.014 (p=0.38; Wilcoxon signed rank test). These results confirm that the second peak was indeed the first harmonic of gamma. We also note that achromatic gratings failed to produce a salient harmonic in M2 (Figure 1H), yielding a ratio that was much less than 2 (Figure 2). Although the frequency ratios for achromatic gratings in most electrodes were concentrated near 1.5 (as indicated by the very small errorbars), a few electrodes (2 out of 16) gave ratios greater than 2, causing the Wilcoxon signed rank test to fail in rejecting the Null hypothesis (albeit with a small p-value of 0.042).

Shape of the gamma waveform

Figure 3 shows LFP traces from example trials corresponding to each stimulus case presented in Figure 1. These traces revealed a characteristic arch shape of gamma waveform, featuring narrower troughs separated by much broader crests, hinting at a characteristic

alignment of gamma fundamental and its harmonic in these cases (best observed in M2 when colored patches were shown).

To visualize how phase differences between gamma and its first harmonic components affected the shape of the summed signal, we added two sinusoids to emulate the fundamental and the first harmonic of gamma, at frequencies 45 Hz and 90 Hz respectively, and varied the initial phase of the 45Hz component (as described in equation 2; shown in Figure 4A). Figure 4B shows the waveforms produced at each of these phases, and Figure 4C indicates the corresponding values of our phase difference measure (computed as in equation 1) between gamma and its harmonic. Using our convention (equation 1), a phase difference of 180 degrees (Figure 4A-C row 3) gave the desired shape as troughs of both sinusoids aligned to produce a steeper overall trough.

Gamma waveform is similar in different hues

In Figure 5, we show the distribution of trial-averaged gamma-harmonic phase differences in all electrodes, for each stimulus (indicated by color), arranged horizontally in order of average gamma power produced. For each stimulus, the trial-averaged phase differences from all electrodes were subject to Rayleigh test, and the stimuli with non-uniform phase difference distributions (p-value < 0.01) are indicated by filled circles. The error bars in each case represent the 95% confidence interval of mean phase differences. The phase differences at higher gamma generating stimuli were distributed close to 180° in both subjects, as predicted from the arch-shaped waveforms (Figure 4). Stimuli that produced low power had high dispersion of phase differences, likely originating from higher influence of noise, as is visible from the larger confidence intervals in M1, resulting in some hues with nearly uniform phase difference distributions (unfilled circles; some cases lack error bars as confidence intervals could not be computed due to their near-uniform distributions). For hues that generated high gamma power, gamma-harmonic phase differences were concentrated slightly

higher than 180° for M1 and close to 180° for M2. The circular mean of electrode-averaged

phase difference between gamma and its first harmonic across those stimuli which showed

unimodal distribution (filled circles in Figure 5; 29 stimuli in M1 and all stimuli in M2) was

185.33±16.66 degrees for M1 (176.60±16.32 degrees if all stimuli were considered) and

153.90±5.56 degrees for M2. When restricted to the ten hues with the highest gamma power

for each monkey, the phases were 206.02±4.65 degrees for M1 and 170.23±3.00 degrees for

M2.

<u>Gamma waveform in a linear population rate model with stochastic inputs</u>

To test whether the stereotypical gamma waveforms observed in the data could be

modeled, we focused on two recently developed rate models by Jia, Xing and Kohn (2013;

abbreviated as the JXK model) and Jadi and Sejnowski (2014; JS model), both of which can

explain the contrast and size dependence of gamma oscillations (see Methods for details).

The JXK model, which is a piecewise linear model, operates as a damped oscillator in

its linear domain and produces sustained oscillations by virtue of time-varying input drives

generated by a Poisson process causing the responses to be stochastic as well. Hence, we

simulated the model over 2 seconds repeatedly for 50 iterations and analyzed the interval from

1-2 seconds ('analysis' window). Figure 6A shows the gamma peak amplitude obtained from

the average PSD of the LFP proxy across iterations for each stimulus size and contrast; Figure

6B shows the corresponding mean gamma peak frequencies. These plots show the model

replicating the stimulus size/contrast effects on gamma as demonstrated by Jia et al. (2013):

gamma frequency decreased and power increased as stimulus size increased, and gamma peak

frequency increased when contrast increased. Figure 6C shows the average PSD from all

iterations for different stimulus sizes (r) at a fixed contrast (c) of $10^{-0.25}$, showing a distinct

bump of activity in gamma range, whose peak frequency decreases but peak power increases

as we go from small to larger values of stimulus size (r). Likewise, Figure 6D shows the average

PSD for different contrasts and a fixed size of 4.375. The gamma peak frequency in the LFP proxy increased with contrast as demonstrated by Jia et al. (2013). However, the gamma bumps were broad, as expected of a noisy pseudoperiodic signal, and no prominent harmonic band activity was discernible in the PSDs. We performed phase analysis, considering the activity in the band centered around twice the gamma peak frequency (Figure 6B) as the harmonic band activity. Figure 6E shows the harmonic amplitudes and Figure 6F shows the mean gamma-harmonic phase differences of LFP proxy traces from different iterations. We searched for stimulus size and contrasts which yielded arch-shaped gamma, or equivalently a gamma-harmonic phase differences close to 180 degrees (Figure 4C, middle) consistently across iterations. Such domains of inputs are shown by black contours in Figure 6F and are identified as 'in-regime' stimuli (see Methods; *Identifying the operational input regimes using gamma-harmonic phase difference*). Such regimes were few and scattered sparsely. An example LFP proxy trace for the highest contrast condition in Figure 6D, which was in-regime, is shown in Figure 6G. The top panel shows the LFP proxy trace and the bottom panel shows the gamma (blue) and harmonic (red) band signals. The LFP traces exhibited bursts of gamma oscillations with randomly distorted waveforms, with no consistent relationship between gamma and the harmonic.

Gamma waveform in JS model

Jadi and Sejnowski (2014) simulated visually evoked gamma using a firing rate model with only an Excitatory and an Inhibitory population and demonstrated gamma as self-sustained oscillations in response to steady constant input drives. As a result, any waveform shape found in a cycle of the oscillations is repeated throughout. We studied whether this model could produce realistic gamma waveforms by identifying the input regime (pairs of input drives to Excitatory and Inhibitory population) that would give rise to gamma-harmonic phase difference around 180° in LFP.

We implemented the model with the parameters specified in Jadi and Sejnowski (2014) for different input drives as listed in Table 1, and computed the LFP proxy in each case. Figure 7A-B shows the gamma peak amplitude and frequency respectively computed from the LFP proxy for each input-combination. The region within the white contour indicates an inhibition-stabilized network in which the inhibitory inputs are superlinear (for details, see Jadi and Sejnowski (2014)). Figure 7C shows the PSDs of LFP proxies obtained for different inhibitory input drives and a constant excitatory drive (input-combinations marked by same colored markers in Figures 7A-B), simulating the variation of stimulus size at a fixed contrast, as simulated in Jadi and Sejnowski (2014). The PSDs demonstrate the decrease in peak frequency and increase in power of gamma in our LFP proxy signal with increasing stimulus size. To demonstrate the contrast effect, we chose a set of input-combinations where both excitatory and inhibitory input drives linearly increased with contrast (colored markers in Figures 7A-B). The PSDs of LFP proxy generated for increasing contrasts are shown in Figure 7D. The gamma frequency increased for higher contrasts mimicking gamma in LFP recordings (Jia et al., 2013). Importantly, this model showed prominent harmonics as well.

Figure 7F plots the gamma-harmonic phase differences in the LFP proxy trace generated for each input combination. Regions in the input drives where the phase difference fell within 22.5° from 180° are enclosed by contours (Figure 7F, black contours) and identified as 'in-regime'. Interestingly, this identified regimes fell inside the region predicted by Jadi and Sejnowski, (2014) where inhibition was superlinear (Figure 7F, white contours), and hence both validated their earlier findings and constrained the model to operate in more restricted regions. Figure 7G presents an example LFP proxy trace in the top panel, corresponding to the largest size condition in Figure 7C (in-regime), whose gamma fundamental (blue) and first harmonic (red) components are shown in the bottom panel. The troughs of gamma and harmonic were indeed aligned, adding up to a wider crest and a sharper trough overall.

Figure 8 shows the gamma-harmonic phase differences and regimes found in both the

models when the activity of only one population was taken as the LFP proxy. Figure 8A show

the phase differences in the JXK model using -rE as LFP proxy and Figure 8B shows the results

using -rI. As before, no connected region of in-regime stimuli could be found (similar to the

case in Figure 6F). Figure 8C-D show the phase differences found in the JS model using these

LFP proxies. While the location and number of in-regime inputs identified were different, the

superlinear regime was still found to contain a connected region of in-regime inputs in either

case. This suggests that an LFP proxy chosen as any non-negative linear combination of -rE

and -rI, would give rise to 'in-regime' inputs within the superlinear regime of the JS model.

Discussion

We show quantitatively that gamma oscillations produced by different hues possess a

distinctive arch-shaped waveform, which leads to a distinct peak at the first harmonic of gamma

in the PSD, and a specific phase relationship between gamma and its harmonic. Further, we

show that a linear, stochastically forced model proposed by Jia et al. (2013) does not produce

distinct harmonics with consistent phase relationship and hence does not retain a specifically-

shaped gamma waveform. On the other hand, a non-linear, self-oscillating model proposed by

Jadi and Sejnowski (2014) produces prominent harmonics, and in a subdomain of inputs within

the previously identified superlinear regime, the model generates the observed gamma

waveform as well.

One important point to note is that this arch shaped gamma waveform with prominent

harmonics were categorized for hue induced gamma, whereas most previous reports have used

achromatic stimuli. In the achromatic stimulus used here, harmonics were not salient,

especially for M2. However, although earlier experimental studies using achromatic gratings

did not describe or quantify the gamma waveform, visual inspection of raw LFP traces

containing gamma bursts reveal a similar shape in some studies (Jia and Kohn, 2011, Fig. 2;

Jia et al., 2011, Fig. 1A; Xing et al., 2012, Fig. 2; Brunet et al., 2014, Fig. 2D) showing that the

arch-shape of gamma is observed for at least some achromatic stimuli. The arch shape

described here is also visible when monkeys viewed colored natural images (Brunet et al.,

2015, Fig. 2). Interestingly, some papers (Gieselmann and Thiele, 2008, Fig. 1; Jia et al., 2011,

Fig. 9) have shown traces with an inverted arch shape. It is unclear why this is the case;

potential reasons could be due to differences in the position of the reference wire, cortical depth

of recording, or differences in the conductance level. For example, arch-shaped gamma

waveform is observed in biophysical ING-PING and spiking models as well (Mazzoni et al.,

2015, Fig. 4; Lowet et al., 2016, Fig. 4), in which non-sinusoidal gamma arises from the

dynamics of the voltage-gated conductances, and the specific shape produced by the model

varies when the conductances are varied. LFP waveform also changes with the recording depth

(Figure 4B versus 4C of Mazzoni et al., 2015). Non-sinusoidal waveforms are common in

neural rhythms, such as the sawtooth-shaped theta rhythm in the hippocampus and comb-

shaped beta rhythm, arising from a variety of different biophysical phenomena (Cole and

Voytek, 2017).

<u>Gamma oscillations in the JXK model</u>

JXK model, like another model proposed by Kang and colleagues (Kang et al., 2010),

explains the "bursty" nature of gamma oscillations, which was shown in previous studies

(Burns et al., 2010; Xing et al., 2012). These studies observed that gamma band power varied

during a trial, with short intervals (~130 ms) of high gamma power, interspersed by durations

of no prominent power. Based on these, Xing et al. (2012) suggested that gamma rhythms were

the outcome of cortical network resonating to the stochastic component in the inputs. During

periods in a trial when stochastic variations in input are less, the cortical network would tend

to asymptotically stabilize and oscillations would get weaker over time. Xing et al. (2012)

demonstrated this mechanism in a piecewise linear asymptotically stable model with noisy

inputs. The JXK model (Jia et al., 2013) emulated gamma burst-activity exhibiting the observed

dependencies of gamma power and frequency on stimulus parameters such as size and contrast

by varying input drives and recurrent excitation to the neural populations. However, the JXK

model does not produce a consistent waveform shape, because despite the presence of a

rectification component, the model operates mostly in a linear (post-rectification) domain for

all inputs simulated. In this domain the step response would be sinusoidal oscillations damping

exponentially over time. Although the presence of Poisson inputs distorted the waveform and

prolonged oscillations for a longer duration, these distortions are not consistent across different

cycles of oscillation. This was reflected as wide band of activity around the gamma band in the

PSD (Figure 6C-D), and a lack of discernible harmonic activity. Therefore, while these models

adequately explained the bursty nature of gamma oscillations, it is not surprising that they

failed to produce a specific gamma waveform.

However, a recent study has shown that the duration of gamma 'burst' activity reported

in Xing et al. (2012) could have been underestimated (Chandran KS et al., 2017). This error

arises because the spectral estimator that is used to compute gamma power is itself noisy,

causing the rhythm to appear bursty in the spectral domain (see Chandran KS et al. (2017) for

details). Chandran KS et al. (2017) used Matching Pursuit algorithm with a Gabor dictionary,

which chooses the best-matching template to estimate the duration of a gamma burst in the

time domain itself to show that gamma bursts during presentation of achromatic grating were

longer than reported earlier (median of ~300 ms, although the mode was ~100 ms). Along with

the results presented here, this suggests that the gamma rhythm could have more temporal

regularity than what is predicted by these stochastic models.

However, we note that the non-sinusoidal nature of gamma rhythm described here could

be introduced in the JXK model by changing the activation function. Specifically, if a non-

linear activation function is used in the place of the piece-wise linear rectifier, the step response

of the system would itself be a specific non-sinusoidal waveform in subsequent cycles of the

resulting damped oscillation. When stochastic perturbations are added, the resultant bursts of

oscillatory activity will contain both irregular distortions caused by the noisily fluctuating

inputs and more regular distortions induced by non-linearity at specific parts of the trajectory

in each cycle. Therefore, although we show that our results are more consistent with JS instead

of the JXK model, it is possible that the JXK model with a suitable activation function can also

replicate our findings. Indeed, while the JS model generates the proper gamma shape within a

sub-regime, the duration of gamma is as long as the duration of the stimulus itself. Therefore,

some stochasticity may be necessary to replicate all features of gamma rhythmicity, including

duration and shape.

In the JXK model, there were a few inputs that qualified our phase-difference criteria

for 'in-regime' behavior. However, a close examination of the 'in-regime' trace (example in

Figure 6G) revealed that, in addition to the irregularly distorted gamma cycles, these inputs

had occasional isolated steep troughs. These steep troughs could be approximated by a negative

impulse, which could be decomposed as a series of sinusoids with their troughs aligned,

resulting in sporadic 180-degree phase-difference estimates. Since these distortions are of

lower frequency than gamma, the JXK model does not operate 'in-regime' for any input

combination.

Relationship between firing rates and LFP

JXK and JS models used excitatory cell activity to study the power and frequency trends

of gamma. Here, we converted these firing rates to a "proxy" LFP to be compatible with the

real data. Because the extracellular potential is generated due to the spatial separation of

transmembrane currents (generating dipoles or multi-poles), as well as alignment of such

dipoles across neurons, LFPs are thought to mainly reflect the transmembrane currents of

pyramidal neurons (Einevoll et al., 2013). But Hasenstaub et al. (2005), in their in-vivo study

of ferret prefrontal cortex, found that the postsynaptic potentials and the firing of regular-

spiking excitatory neurons and fast-spiking interneurons were both synchronized to the

extracellular gamma rhythm, but the activation of interneurons was more aligned with the

troughs of the oscillation. A modelling study (Mazzoni et al., 2015) simulated LFP using

various outputs of a Leaky Integrate-and-Fire (LIF) network (such as firing rates, membrane

potential and synaptic currents) and found that the sum of absolute values of synaptic currents

(with both AMPA and GABA currents weighted nearly equally) served as the best proxy for

LFP. As far as gamma oscillations are concerned, some studies have shown that spikes tend to

occur just before the trough of gamma (Chalk et al., 2010; Vinck et al., 2010; Das and Ray,

2018), with the pyramidal cells leading the interneurons by a few milliseconds (Csicsvari et

al., 2003; Hasenstaub et al., 2005; Vinck et al., 2013). In the JS model as well, we found that

the E population led the I population by about 50º or 2-3 ms in the superlinear regime. Further,

the results were qualitatively similar when just E or I or the sum of both were used as the LFP

proxy (Figure 7 and Figure 8).

<u>Gamma and Harmonics vs slow/fast gamma</u>

Murty et al., (2018) showed that fullscreen achromatic gratings produced two different

oscillations simultaneously – a fast gamma (40-70 Hz), which has been observed previously

for smaller sized gratings, and a slow gamma (20-40 Hz) which is prominent only for large

stimuli. Even though this also generates two bums in the PSD, fast gamma is not a harmonic

of the slow gamma. First, the center frequency of fast gamma was not twice that of slow

gamma. Second, slow and fast gamma were not co-tuned: they had distinct orientation,

contrast, temporal frequency and size tuning preferences.

The arch-shape shown here poses additional problems for phase coding schemes in

which spike position relative to the gamma phase is used to code information (Fries et al., 2007;

Vinck et al., 2010), because spikes tend to occur near the trough of the rhythm and having a

sharper trough reduces the operating range. Other studies have proposed that gamma

oscillations may not play a role but could be a useful marker/indicator of cortical processing

(Ray and Maunsell, 2015). In that framework, properties such as shape and duration along with

stimulus tuning could provide additional clues about the underlying circuitry.

References

Bartoli E, Bosking W, Chen Y, Li Y, Sheth SA, Beauchamp MS, Yoshor D, Foster BL

(2019) Functionally Distinct Gamma Range Activity Revealed by Stimulus Tuning in

Human Visual Cortex. Curr Biol 29:3345-3358.e7.

Berens P (2009) CircStat: A MATLAB Toolbox for Circular Statistics. J Stat Softw 31:1–21.

Bokil H, Andrews P, Kulkarni JE, Mehta S, Mitra PP (2010) Chronux: A platform for

analyzing neural signals. J Neurosci Methods 192:146–151.

Brunet N, Bosman CA, Roberts M, Oostenveld R, Womelsdorf T, De Weerd P, Fries P

(2015) Visual Cortical Gamma-Band Activity During Free Viewing of Natural

Images. Cereb Cortex 25:918–926.

Brunet NM, Bosman CA, Vinck M, Roberts M, Oostenveld R, Desimone R, De Weerd P,

Fries P (2014) Stimulus repetition modulates gamma-band synchronization in primate

visual cortex. Proc Natl Acad Sci U S A 111:3626–3631.

Burns SP, Xing D, Shapley RM (2011) Is Gamma-Band Activity in the Local Field Potential

of V1 Cortex a "Clock" or Filtered Noise? J Neurosci 31:9658–9664.

Burns SP, Xing D, Shelley MJ, Shapley RM (2010) Searching for autocoherence in the

cortical network with a time-frequency analysis of the local field potential. J Neurosci

Off J Soc Neurosci 30:4033–4047.

Buzsaki G (2006) Rhythms of the brain. Oxford University Press, USA.

Chalk M, Herrero JL, Gieselmann MA, Delicato LS, Gotthardt S, Thiele A (2010) Attention

reduces stimulus-driven gamma frequency oscillations and spike field coherence in

V1. Neuron 66:114–125.

Chandran KS S, Seelamantula CS, Ray S (2017) Duration analysis using matching pursuit

algorithm reveals longer bouts of gamma rhythm. J Neurophysiol 119:808–821.

Chariker L, Shapley R, Young L-S (2018) Rhythm and Synchrony in a Cortical Network

Model. J Neurosci 38:8621–8634.

Cole SR, Voytek B (2017) Brain Oscillations and the Importance of Waveform Shape.

Trends Cogn Sci 21:137–149.

Csicsvari J, Jamieson B, Wise KD, Buzsáki G (2003) Mechanisms of Gamma Oscillations in

the Hippocampus of the Behaving Rat. Neuron 37:311–322.

Das A, Ray S (2018) Effect of Stimulus Contrast and Visual Attention on Spike-Gamma

Phase Relationship in Macaque Primary Visual Cortex. Front Comput Neurosci 12

Available at: https://www.frontiersin.org/articles/10.3389/fncom.2018.00066/full

[Accessed March 15, 2019].

Einevoll GT, Kayser C, Logothetis NK, Panzeri S (2013) Modelling and analysis of local

field potentials for studying the function of cortical circuits. Nat Rev Neurosci

14:770–785.

Fries P, Nikolić D, Singer W (2007) The gamma cycle. Trends Neurosci 30:309–316.

Gieselmann MA, Thiele A (2008) Comparison of spatial integration and surround

suppression characteristics in spiking activity and the local field potential in macaque

V1. Eur J Neurosci 28:447–459.

Hasenstaub A, Shu Y, Haider B, Kraushaar U, Duque A, McCormick DA (2005) Inhibitory

Postsynaptic Potentials Carry Synchronized Frequency Information in Active Cortical

Networks. Neuron 47:423–435.

Henrie JA, Shapley R (2005) LFP Power Spectra in V1 Cortex: The Graded Effect of

Stimulus Contrast. J Neurophysiol 94:479–490.

Jadi MP, Sejnowski TJ (2014) Regulating Cortical Oscillations in an Inhibition-Stabilized

Network. Proc IEEE Inst Electr Electron Eng 102.

Jia X, Kohn A (2011) Gamma rhythms in the brain. PLoS Biol 9:e1001045.

Jia X, Smith MA, Kohn A (2011) Stimulus Selectivity and Spatial Coherence of Gamma

Components of the Local Field Potential. J Neurosci 31:9390–9403.

Jia X, Xing D, Kohn A (2013) No consistent relationship between gamma power and peak

frequency in macaque primary visual cortex. J Neurosci Off J Soc Neurosci 33:17–25.

Kang K, Shelley M, Henrie JA, Shapley R (2009) LFP spectral peaks in V1 cortex: network

resonance and cortico-cortical feedback. J Comput Neurosci Available at:

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Ci

tation&list_uids=19862612.

Kang K, Shelley M, Henrie JA, Shapley R (2010) LFP spectral peaks in V1 cortex: network

resonance and cortico-cortical feedback. J Comput Neurosci 29:495–507.

Keeley S, Byrne Á, Fenton A, Rinzel J (2019) Firing rate models for gamma oscillations. J

Neurophysiol 121:2181–2190.

Lowet E, Roberts MJ, Bonizzi P, Karel J, Weerd PD (2016) Quantifying Neural Oscillatory

Synchronization: A Comparison between Spectral Coherence and Phase-Locking

Value Approaches. PLOS ONE 11:e0146443.

Mazzoni A, Lindén H, Cuntz H, Lansner A, Panzeri S, Einevoll GT (2015) Computing the

Local Field Potential (LFP) from Integrate-and-Fire Network Models. PLOS Comput

Biol 11:e1004584.

Murty DVPS, Shirhatti V, Ravishankar P, Ray S (2018) Large Visual Stimuli Induce Two

Distinct Gamma Oscillations in Primate Visual Cortex. J Neurosci Off J Soc Neurosci

38:2730–2744.

Peter A, Uran C, Klon-Lipok J, Roese R, van Stijn S, Barnes W, Dowdall JR, Singer W, Fries

P, Vinck M (2019) Surface color and predictability determine contextual modulation

of V1 firing and gamma oscillations Colgin L, ed. eLife 8:e42101.

Ray S, Maunsell JH (2011) Different origins of gamma rhythm and high-gamma activity in

macaque visual cortex. PLoS Biol 9.

Ray S, Maunsell JHR (2010) Differences in Gamma Frequencies across Visual Cortex

Restrict Their Possible Use in Computation. Neuron 67:885–896.

Ray S, Maunsell JHR (2015) Do gamma oscillations play a role in cerebral cortex? Trends

Cogn Sci 19:78–85.

Shirhatti V, Ray S (2018) Long-wavelength (reddish) hues induce unusually large gamma

oscillations in the primate primary visual cortex. Proc Natl Acad Sci 115:4489–4494.

Tiesinga P, Sejnowski TJ (2009) Cortical enlightenment: are attentional gamma oscillations

driven by ING or PING? Neuron 63:727–732.

Tsodyks MV, Skaggs WE, Sejnowski TJ, McNaughton BL (1997) Paradoxical effects of

external modulation of inhibitory interneurons. J Neurosci Off J Soc Neurosci

17:4382–4388.

Veit J, Hakim R, Jadi MP, Sejnowski TJ, Adesnik H (2017) Cortical gamma band

synchronization through somatostatin interneurons. Nat Neurosci 20:951–959.

Vinck M, Lima B, Womelsdorf T, Oostenveld R, Singer W, Neuenschwander S, Fries P

(2010) Gamma-phase shifting in awake monkey visual cortex. J Neurosci Off J Soc

Neurosci 30:1250–1257.

Vinck M, Womelsdorf T, Buffalo EA, Desimone R, Fries P (2013) Attentional Modulation of

Cell-Class-Specific Gamma-Band Synchronization in Awake Monkey Area V4.

Neuron 80:1077–1089.

Wilson HR, Cowan JD (1972) Excitatory and inhibitory interactions in localized populations

of model neurons. Biophys J 12:1–24.

Xing D, Shen Y, Burns S, Yeh C-I, Shapley R, Li W (2012) Stochastic Generation of

Gamma-Band Activity in Primary Visual Cortex of Awake and Anesthetized

Monkeys. J Neurosci 32:13873–13880a.

Figures

Figure 1: Visual Stimulation induces narrowband activity in gamma and its harmonic bands. (A-D, *top row*) PSDs for baseline (-500 to 0 ms; 0 indicates stimulus onset) and stimulus (250 to 750 ms) periods for different stimuli, averaged across trials and electrodes, from monkey 1 (M1). The baseline PSDs are plotted in black, while the stimulus period PSDs are colored (corresponding to the hue presented) or grey (for fullscreen grating). (A-D, *bottom row*) show the change in power (dB) in the stimulus period from the baseline, computed from the PSDs in the top row. The gamma power computed from the change in power spectrum is

702   represented by vertical dashed line). The peaks in gamma range and the second bump activity

703   are marked in each plot. (E-H) Same as A-D, for monkey M2.

Figure 2: Second bump in PSD corresponds to the first harmonic of gamma. Median of peak frequency ratios of the second bump with respect to the gamma band, scattered against average gamma power for each stimulus (circles are colored to represent the presented hues; grating stimulus is represented by the grey circle). Error bars indicate the standard error of median (estimated by bootstrapping). Open circles mark stimuli whose ratios were found to be significantly different from 2 (p-value < 0.01 without any Bonferroni correction; Wilcoxon signed-rank test). The dotted horizontal lines correspond to the least-error margin 2±0.05.

Figure 3: Gamma waveform has consistent non-sinusoidal shape under different stimuli.

(A-D) present LFP traces for different stimuli, shown in Figure 1, from an example trial and

electrode in M1. (E-H) shows LFP traces obtained from M2 for corresponding stimuli.

Figure 4: Observed arch-shape of gamma corresponds to gamma-harmonic phase

difference of 180°. (A) Gamma (blue) and harmonic (red) components, mimicked using

sinusoids as in equation (2) with different initial gamma phases (φ_G) in each row. (B) Gamma

waveform resulting from summation of gamma and its harmonic from the corresponding row

in (B). (C) Phase difference of gamma and its first harmonic, as given in equation (1), computed

from filtering and Hilbert transforming gamma and harmonic bands from the summed

waveform in (B) in each row. Note that the arch-shaped gamma observed in recordings arises

from a phase difference of 180° (middle row).

Figure 5: Gamma-harmonic phase differences in different stimuli are concentrated near 180°. Circular mean of trial-averaged gamma-harmonic phase differences (across all electrodes) are scattered against the average gamma power for each stimulus (circles are colored to represent the presented hues; grey for grating stimulus). For each stimulus, the pool of trial-averaged phase differences from all electrodes is subject to Rayleigh test of non-uniformity to check if phase-difference is consistent across electrodes so that circular mean estimates are reliable (p-value < 0.01). If found reliable (even though it could be different from 180°), the stimulus is represented by a filled circle. Error bars represent the 95% confidence interval of circular mean.

Figure 6: Gamma properties for the JXK model (A) Gamma peak amplitude and (B)

frequency found in the average PSD from 50 iterations at each stimulus size (r) and contrast

(c) combination. (C) Average PSD of LFP proxy traces generated at fixed contrast and different

sizes, across all iterations. (D) Average PSD of LFP proxy traces generated at different

contrasts at a fixed size. The stimuli values that generated these PSDs are indicated in plots

(A), (B) and (E) using colored markers (same colors as the plots in C and D). (E) Harmonic

band peak amplitude in the average PSD. The harmonic is taken to be twice of gamma

frequency. (F) Mean gamma-harmonic phase difference (equation (1)) across multiple

iterations, with in-regime inputs encircled by black contours. (G) Example LFP proxy activity

(top panel) and its gamma and first harmonic components (bottom panel) generated at the

highest contrast condition in (D) (identified as in-regime). This trace shows occasional steeper

troughs (bottom trace; near 1.35 s, 1.52 s and 1.58 s timepoints) although there is no consistent

trend between gamma and the harmonic.

Figure 7: JS model produces sustained arch-shaped gamma oscillations within the previously identified superlinear inhibitory regime. (A) Peak gamma amplitude and (B)

Peak gamma frequency in the LFP proxy generated by the JS model for each input-drive combination. (C) PSD of LFP proxy generated for different stimulus sizes. (D) PSD of LFP proxy generated for different contrast. Input combinations used in (C) and (D) are indicated by markers, with same colors as traces, in (A), (B) and (E). (E) Harmonic amplitude identified from the PSD of LFP proxy for each stimulus condition. (F) Phase difference of gamma and its harmonic (equation (1)) computed for each simulated input drive pair. The black contours encircle input drives identified to be 'in-regime' from their phase differences. The white contour in each figure enclose the input domain identified by Jadi and Sejnowski (2014) to replicate gamma power increase and peak frequency decrease in response to increasing stimulus size, and it can be seen to enclose a significant portion of the 180° phase difference regime (black contour). (G) LFP proxy activity trace corresponding to the largest stimulus size shown in Figure 7C. The dotted horizontal line shown is equidistant from the minima and maxima of the traces. A sinusoidal oscillation would have the same crest and trough width on this line. The bottom panels show the gamma and its first-harmonic components, plotted in blue and red respectively, filtered from the LFP proxy traces.

Figure 8: Gamma-harmonic phase difference in rate models using different LFP proxies.

(A) Phase difference of gamma and its harmonic in the JXK model computed for each stimulus

condition (as in Figure 6F) but using only E population activity to compute the LFP proxy. (B)

Gamma-harmonic phase differences in JXK using I population activity alone. (C) Phase

differences in JS model using E population activity only. (D) Phase differences in JS using I

population activity only. Black contours encircle stimulus conditions identified as 'in-regime'.

White contours in (C) and (D) mark the superlinear regime of the JS model.

783   Tables:

784   **Table 1: Parameter values used in rate models.**

JS							
W_{EE}	W_{EI}	W_{IE}	W_{II}	τ_E, τ_I (ms)			
16	26	20	1	20,10			
me,mi	thetae,thetai	I_E,I_I					
1,1	5,20	0 to 20 (steps of 2.5)					
JXK							
W_{EE}	W_{EI}	W_{IE}	W_{II}	W_{EG}	W_{IG}	W_{GE}	τ_E, τ_I, τ_G (ms)
1.5	3.25	3.5	2.5	0.25	0.5	0.6	6,15,19
MN	r	c					
0	1 to 5 (steps of 0.125)	0.01 to 1 (logarithmic steps of 0.5)					

43