Title: Shape analysis of gamma rhythm supports a superlinear inhibitory

regime in an inhibition-stabilized network

Running title: Shape analysis of gamma rhythm

Authors: R (first name) Krishnakumaran (last name)[1], Mohammed (first name) Raees (last

name)[2] and Supratim (first name) Ray (last name)[1,2,*]

Affiliations:

[1]IISc Mathematics Initiative, Department of Mathematics, Indian Institute of Science,

Bangalore, India, 560012

[2]Centre for Neuroscience, Indian Institute of Science, Bangalore, India, 560012,

Telephone +91 80 2293 3437, Facsimile +91 80 2360 3323

*__Corresponding Author:__ sray@iisc.ac.in

Keywords: Gamma, Harmonics, Phase analysis, Wilson-Cowan model

Conflict of interest: The authors declare no competing financial interests.

Funding Disclosure: This work was supported by Wellcome Trust/DBT India Alliance

(Senior fellowship IA/S/18/2/504003 to SR) and DBT-IISc Partnership Programme.

Acknowledgements: We thank Dr. Adam Kohn for his helpful comments on the paper.

Abstract

Visual inspection of stimulus-induced gamma oscillations (30-70 Hz) often reveals a non-sinusoidal shape. Such distortions are a hallmark of non-linear systems and are also observed in mean-field models of gamma oscillations. A thorough characterization of the shape of the gamma cycle can therefore provide additional constraints on the operating regime of such models. However, the gamma waveform has not been quantitatively characterized, partially because the first harmonic of gamma, which arises because of the non-sinusoidal nature of the signal, is typically weak and gets masked due to a broadband increase in power related to spiking. To address this, we recorded local field potential (LFP) from the primary visual cortex (V1) of two awake female macaques while presenting full-field gratings or iso-luminant chromatic hues that produced huge gamma oscillations with prominent peaks at harmonic frequencies in the power spectra. We found that gamma and its first harmonic always maintained a specific phase relationship, resulting in a distinctive shape with a sharp trough and a shallow peak. Interestingly, a Wilson-Cowan (WC) model operating in an inhibition stabilized mode could replicate this shape, but only when the inhibitory population operated in the super-linear regime, as predicted recently. However, another recently developed model of gamma that operates in a linear regime driven by stochastic noise failed to produce salient harmonics or the observed shape. Our results impose additional constraints on models that generate gamma oscillations and their operating regimes.

Introduction

Gamma rhythm refers to oscillatory neural activity in the 30-70 Hz range that changes in response to different stimuli and cognitive states [1]. In the primary visual cortex (V1), the gamma rhythm has been studied extensively using achromatic gratings, wherein gamma power and center-frequency have been shown to vary systematically with the properties of the grating [2,3]. For instance, peak-frequency of gamma increases with the contrast of gratings [2,4] and, gamma power increases and frequency decreases with stimulus size [5,6]. More recently, chromatic stimuli have also been explored, which, for low-wavelength (reddish) hues, can generate huge gamma oscillations that are an order of magnitude stronger than gamma produced by achromatic gratings [7–9]. Further, these plain colored blobs of different sizes exhibited trends in gamma and firing rate analogous to that of achromatic gratings of different sizes [8].

Gamma oscillations are thought to reflect the push-pull activity of interconnected excitatory and inhibitory neurons, which has been demonstrated in different spiking network models [10–13]. However, since such large-scale network models have several parameters to be tuned and can become hard to interpret, simplified population rate models are sometimes used. Although these models have limitations because of fewer free parameters, they are analytically more tractable. One pioneering model was proposed by Wilson and Cowan (WC) [14], in which excitatory and inhibitory neurons were grouped into populations, and the dynamics of these populations were characterized. Variants of WC models have recently been used to explain some properties of the gamma rhythm. For example, based on the observation that gamma rhythm appears in short bursts and its phase does not vary linearly [15–17], a WC based model was proposed in which the activation function (input-output relationship) was linear but driven by Poisson noise [17,18]. A variant of this model by Jia, Xing and Kohn (JXK) [2], in which an additional global excitatory population was added, was used to explain

the stimulus dependence of gamma oscillations. On the other hand, Jadi and Sejnowski (JS)[19]

used a WC model with a non-linear (sigmoidal) activation function and showed that

constraining the model to operate in an inhibition stabilized mode [20] and the inhibitory

population to operate in superlinear domain can also reproduce the size and contrast

dependence of gamma rhythm. In this study, we focus on JXK and JS models since these are

both WC based models which can explain the size and contrast-based stimulus dependencies.

More complex rate models that explain stimulus dependencies and potentially also the gamma

shape are elaborated in the Discussion section.

Although both JXK and JS models can explain the stimulus dependence of gamma, the

shape of the rhythm likely depends on the presence (and type) of non-linearity. The shape could

be non-sinusoidal (see, for example, Figure 11 of [14]), which is represented in the spectral

domain as peaks at harmonics of the fundamental frequency. Such distortions have been

observed in various brain oscillations such as theta rhythm (for a review, see [21]). For gamma

rhythm, a visual inspection of raw traces reveals some distortion [22–24], which is also

corroborated by the presence of peaks at the first harmonic of the fundamental gamma peak in

the power spectra [4,5,7]. However, a quantitative study of the gamma waveform, which could

potentially constrain the type and operating regime of models, has not been undertaken. This

is partly because the noise in the harmonic range could offset its phase estimates if the harmonic

is not prominent enough. To address this, we studied gamma oscillations produced by

presenting full-screen hues that generated salient gamma oscillations with prominent

harmonics [7], characterized the shape, and tested models that could replicate the shape.

Results

We collected spikes and LFP from 96 and 81 electrodes from two monkeys, M1 and

M2, while they viewed full-screen color patches of different hues. Limiting our analysis to

electrodes with reliable estimates of RF centers (see "Electrode selection" section in Methods

and Models) yielded 64 and 16 electrodes from the two monkeys.

<u>First harmonic of gamma oscillation</u>

In Figure 1A-D, the top row shows the power spectral density (PSD; averaged across

all trials and subsequently across all electrodes) of the LFP signal in subject M1 during baseline

period (-500 to 0 ms from stimulus onset; black trace) and during stimulus period (250 to 750

ms from stimulus onset; color/grey trace; the color indicates the hue that was presented and

grey trace represents gratings). The corresponding change in power (dB) from baseline (Figure

1A-D bottom row) shows a prominent peak in the gamma range, along with another prominent

peak near twice the frequency. Figure 1E-H show the same plots from subject M2. We first

tested whether this second peak was indeed at twice the frequency. For each electrode and

stimulus, we measured the gamma peak frequency from trial-averaged baseline-corrected

PSDs as the highest peak within the 30-70 Hz band, and the peak of the second bump as the

highest peak occurring after 12 Hz past the identified gamma peak frequency but before 140

Hz (the identified peaks are highlighted by black crosses in Figure 1).

Figure 2 presents the medians of the frequency ratios of second bump to gamma peak

frequencies for each stimulus (hues indicated by color, gratings by grey color), computed

across all electrodes in each subject, with error bars indicating the standard error of the median

computed by bootstrapping. The stimuli are arranged along the horizontal axis by the average

gamma power across electrodes. The frequency ratios were tightly clustered around 2,

especially for stimuli that produced strong gamma oscillations. To test whether the ratio for a

given stimulus was significantly different from 2, the frequency ratios computed in different

electrodes for the stimulus were subject to a Wilcoxon signed rank test. Stimuli for which the

ratios were found to be significantly different from 2 ($p < 0.01$) are indicated by open circles.

Estimates of peak-frequency ratios for stimuli producing less gamma power were more

susceptible to noise and showed larger deviations in frequency ratios. However, frequency ratios were distributed more narrowly around 2 for stimuli inducing stronger gamma. The median ratio value for M1 was 1.95±0.014, significantly different from 2 ($p = 0.018$ for M1; Wilcoxon signed rank test). For M2, the median ratio was 1.95±0.018, significantly different from 2 ($p = 1.01e-5$; Wilcoxon signed rank test). However, note that because we computed PSDs over a 500 ms window, the frequency resolution was 2 Hz, which introduced some error in the estimated ratio. For example, if the true gamma and harmonic peaks are at 41 and 82 Hz, the estimated gamma will either be at 40 or 42 Hz, yielding a ratio of either $82/40 = 2.05$ or $82/42 = 1.95$. These margins of error are shown as dotted lines in Figure 2. Most of the points lay within these error margins, especially for electrodes with higher gamma power. When we restricted the analysis to the top five hues for each monkey in terms of gamma power, the ratios were 1.96±0.042 ($p=0.63$; Wilcoxon signed rank test) and 1.95±0.026 ($p=0.25$; Wilcoxon signed rank test). These results confirm that the second peak was indeed the first harmonic of gamma. The overall power at peak gamma and first-harmonic frequencies were also highly correlated across electrodes in each monkey (Pearson correlation coefficient was 0.85, $p = 4.86e-19$ for M1 and 0.86, $p = 1.55e-5$ for M2), and the results appeared similar if the power at first harmonic was used instead of main gamma power in Fig 2.

For hues with low gamma power, the peak frequency ratios were more frequently less than 2 in both monkeys. Inspection of their PSDs showed some overlap between the gamma and the harmonic bumps, resulting in non-zero change in power (dB) between the two. When the actual gamma and harmonic peaks were not prominent enough, the overlapping region often had higher power, leading to an overestimation of gamma peak frequency and an underestimation of the harmonic peak frequency, both resulting in a decrease in the peak-frequency ratio.

 Shape of the gamma waveform

Figure 3 shows LFP traces from example trials corresponding to each stimulus case

presented in Figure 1. These traces revealed a characteristic arch shape of gamma waveform,

featuring narrower troughs separated by much broader crests, hinting at a characteristic

alignment of gamma fundamental and its harmonic in these cases (best observed in M2 when

colored patches were shown).

To visualize how phase differences between gamma and its first harmonic components

affected the shape of the summed signal, we added two sinusoids to emulate the fundamental

and the first harmonic of gamma, at frequencies 45 Hz and 90 Hz respectively, and varied the

initial phase of the 45Hz component (as described in equation (2); shown in Figure 4A). Figure

4B shows the waveforms produced at each of these phases, and Figure 4C indicates the

corresponding values of our phase difference measure (computed as in equation (1)) between

gamma and its harmonic. Using our convention (equation (1)), a phase difference of 180

degrees (Figure 4A-C row 3) gave the desired shape as troughs of both sinusoids aligned to

produce a steeper overall trough.

Gamma waveform is similar for different hues

In Figure 5, we show the distribution of trial-averaged gamma-harmonic phase

differences in all electrodes, for each stimulus (indicated by color), arranged horizontally in

order of average gamma power produced. For each stimulus, the trial-averaged phase

differences from all electrodes were subject to Rayleigh test, and the stimuli with non-uniform

phase difference distributions (p-value < 0.01) are indicated by filled circles. The error bars in

each case represent the 95% confidence interval of mean phase differences. The phase

differences at higher gamma generating stimuli were distributed close to $180°$ in both subjects,

as predicted from the arch-shaped waveforms (Figure 4). Stimuli that produced low power had

high dispersion of phase differences, likely originating from higher influence of noise, as is

visible from the larger confidence intervals in M1, resulting in some hues with nearly uniform

phase difference distributions (unfilled circles; some cases lack error bars as confidence

intervals could not be computed due to their near-uniform distributions). For hues that

generated high gamma power, gamma-harmonic phase differences were concentrated slightly

higher than 180° for M1 and close to 180° for M2. The circular mean of electrode-averaged

phase difference between gamma and its first harmonic across those stimuli which showed

unimodal distribution (filled circles in Figure 5; 28 stimuli in M1 and all stimuli in M2) was

197.33±14.44 degrees for M1 (194.00±17.93 degrees if all stimuli were considered) and

161.79±6.79 degrees for M2. When restricted to the five hues with the highest gamma power

for each monkey, the phases were 213.94±6.56 degrees for M1 and 183.92±2.26 degrees for

M2. Interestingly, although the harmonic-band activity was not saliently visible in the PSD in

the achromatic stimulus used here (especially for M2), the phase analysis nonetheless revealed

a consistent phase difference of 227.04±6.81 degrees for M1 (Rayleigh test; p-value = 6.10e-

58) and 164.38±9.57 degrees for M2 (p-value = 2.11e-29; grey filled circle in Fig 5). The slight

difference in distributions of gamma harmonic phase-differences across the two subjects could

arise from unavoidable differences in the recording depths and intrinsic biological differences.

Phase differences for stimuli producing lower gamma powers appeared slightly less

than those at higher powers. For a second set of five stimuli with the highest gamma power

below 10 dB for M1 and below 15 dB for M2, the phases were about 20º less (195.18±31.09

degrees and 143.06±17.25 degrees, respectively). To test the possibility of this reduction

arising from any systematic bias in the phase analyses due to noise, we performed two tests.

First, we repeated the phase analyses on simulated data (see Methods and models; *Effect of*

harmonic phase on gamma waveform) by adding Gaussian noise of different variances to the

summed waveform in equation (2). Secondly, we chose trials in M2 which had the highest

gamma power, and created a synthetic signal which was a weighted average of the signal

segments in the stimulus and baseline periods. In both simulations, the gamma power in the

simulated signal could be reduced by increasing the variance of Gaussian noise and the fraction

of baseline signal, respectively. In both tests, the phase-difference estimates obtained from our

analysis were not biased by higher noise levels, but had larger confidence intervals as observed

in the data at lower gamma powers.

<u>Gamma waveform in a linear population rate model with stochastic inputs</u>

To test whether the stereotypical gamma waveforms observed in the data could be

modeled, we focused on two recently developed rate models by Jia, Xing and Kohn (2013;

abbreviated as the JXK model) [2] and Jadi and Sejnowski (2014; JS model) [19], both of

which can explain the contrast and size dependence of gamma oscillations (see Methods and

Models for details).

The JXK model, which is a piecewise linear model, operates as a damped oscillator in

its linear domain and produces sustained oscillations by virtue of time-varying input drives

generated by a Poisson process causing the responses to be stochastic as well. Hence, we

simulated the model over 2 seconds repeatedly for 50 iterations and analyzed the interval from

1-2 seconds ('analysis' window). Figure 6A shows the gamma peak amplitude obtained from

the average PSD of the LFP proxy across iterations for each stimulus size and contrast; Figure

6B shows the corresponding mean gamma peak frequencies. These plots show the model

replicating the stimulus size/contrast effects on gamma as demonstrated by Jia and colleagues

[2]: gamma frequency decreased and power increased as stimulus size increased, and gamma

peak frequency increased when contrast increased. Figure 6C shows the average PSD from all

iterations for different stimulus sizes (r) at a fixed contrast (c) of $10^{-0.25}$, showing a distinct

bump of activity in gamma range, whose peak frequency decreases but peak power increases

as we go from small to larger values of stimulus size (r). Likewise, Figure 6D shows the average

PSD for different contrasts and a fixed size of 4.375. The gamma peak frequency in the LFP

proxy increased with contrast as demonstrated by Jia and colleagues [2]. However, the gamma

bumps were broad, as expected of a noisy pseudoperiodic signal, and no prominent harmonic

band activity was discernible in the PSDs. We performed phase analysis, considering the

activity in the band centered around twice the gamma peak frequency (Figure 6B) as the

harmonic band activity. Figure 6E shows the harmonic amplitudes and Figure 6F shows the

mean gamma-harmonic phase differences of LFP proxy traces from different iterations. We

searched for stimulus size and contrasts which yielded arch-shaped gamma, or equivalently a

gamma-harmonic phase differences close to 180 degrees (Figure 4C, middle) consistently

across iterations. Such domains of inputs are shown by black contours in Figure 6F and are

identified as 'in-regime' stimuli (see Methods and Models; *Identifying the operational input*

regimes using gamma-harmonic phase difference). Such regimes were few and scattered

sparsely. An example LFP proxy trace for the highest contrast condition in Figure 6D, which

was in-regime, is shown in Figure 6G. The top panel shows the LFP proxy trace and the bottom

panel shows the gamma (blue) and harmonic (red) band signals. The LFP traces exhibited

bursts of gamma oscillations with waveforms showing varied distortions.

<u>Gamma waveform in JS model</u>

Jadi and Sejnowski [19] simulated visually evoked gamma using a firing rate model

with only an Excitatory and an Inhibitory population and demonstrated gamma as self-

sustained oscillations in response to steady constant input drives. As a result, any waveform

shape found in a cycle of the oscillations is repeated throughout. We studied whether this model

could produce realistic gamma waveforms by identifying the input regime (pairs of input drives

to Excitatory and Inhibitory population) that would give rise to gamma-harmonic phase

difference around 180° in LFP.

We implemented the model with the parameters specified in [19] for different input

drives as listed in Table 1, and computed the LFP proxy in each case. Figure 7A-B shows the

gamma peak amplitude and frequency respectively computed from the LFP proxy for each input-combination. The region within the white contour indicates an inhibition-stabilized network in which the inhibitory inputs are superlinear (for details, see [19]). Figure 7C shows the PSDs of LFP proxies obtained for different inhibitory input drives and a constant excitatory drive (input-combinations marked by same-colored markers in Figures 7A-B), simulating the variation of stimulus size at a fixed contrast, as simulated in [19]. The PSDs demonstrate the decrease in peak frequency and increase in power of gamma in our LFP proxy signal with increasing stimulus size. To demonstrate the contrast effect, we chose a set of input-combinations where both excitatory and inhibitory input drives linearly increased with contrast (colored markers in Figures 7A-B). The PSDs of LFP proxy generated for increasing contrasts are shown in Figure 7D. The gamma frequency increased for higher contrasts mimicking gamma in LFP recordings [2]. Importantly, unlike JXK, this model showed power at harmonic frequencies as well.

Figure 7F plots the gamma-harmonic phase differences in the LFP proxy trace generated for each input combination. Regions in the input drives where the phase difference fell within 22.5° from 180° are enclosed by contours (Figure 7F, black contours) and identified as 'in-regime'. Interestingly, this identified regimes fell inside the region predicted by Jadi and Sejnowski [19] where inhibition was superlinear (Figure 7F, white contours), and further constrained the model to operate in more restricted regions. Note that small regions lying close to the grey region on the plot were also occasionally characterized as in-regime, but they are likely spuriously generated by simulation noise and the corresponding gamma and harmonic powers were very low (Figure 7A and 7E) and were therefore not considered. Figure 7G presents an example LFP proxy trace in the top panel, corresponding to the largest size condition in Figure 7C (in-regime), whose gamma fundamental (blue) and first harmonic (red) components are shown in the bottom panel. The troughs of gamma and harmonic were indeed

aligned, adding up to a wider crest and a sharper trough overall. However, we note that the

overall magnitude of the harmonic was smaller than what we observed in real data. This could

be a general limitation of mean field models, as discussed later.

The LFP proxy used here involves both r_E and r_I with equal contributions, as suggested

by a computational modelling study of LFP proxies by Mazzoni and colleagues [10] (see

Relationship between firing rates and LFP under Discussion for further details and discussion

of other proxies). However, the contributions to LFP by different populations could vary with

depth of recording, proximity to the neurons since they determine the relative distribution of

different synapses in the recording neighborhood [25]. To test whether the ability of the models

to generate arch-shaped waveforms depends on the relative contribution of the two populations,

we tested proxies containing contribution from only one population. Figure 8 shows the

gamma-harmonic phase differences and regimes found in both the models for these proxies.

Figure 8A show the phase differences in the JXK model using -rE as LFP proxy and Figure 8B

shows the results using -rI. As before, no connected region of in-regime stimuli could be found

(similar to the case in Figure 6F). Figure 8C-D show the phase differences found in the JS

model using these LFP proxies. While the location of in-regime inputs identified was different,

the superlinear regime was still found to contain a connected region of in-regime inputs in

either case, although the operating regions were very small now. Nevertheless, this suggests

that an LFP proxy chosen as any non-negative linear combination of -rE and -rI, would give

rise to 'in-regime' inputs within the superlinear regime of the JS model.

Discussion

We show quantitatively that gamma oscillations produced by different hues possess a

distinctive arch-shaped waveform, which leads to a distinct peak at the first harmonic of

gamma in the PSD, and a specific phase relationship between gamma and its harmonic. Further,

we show that a linear, stochastically forced model proposed by Jia and colleagues [2] does not

produce distinct harmonics with consistent phase relationship and hence does not retain a

specifically-shaped gamma waveform. On the other hand, a non-linear, self-oscillating model

proposed by Jadi and Sejnowski [19] produces prominent harmonics, and in a subdomain of

inputs within the previously identified superlinear regime, the model generates the observed

gamma waveform as well.

One important point to note is that this arch shaped gamma waveform with prominent

harmonics were characterized for hue induced gamma, whereas most previous reports have

used achromatic stimuli. In the achromatic stimulus used here, harmonics were not salient,

especially for M2. However, although earlier experimental studies using achromatic gratings

did not describe or quantify the gamma waveform, visual inspection of raw LFP traces

containing gamma bursts reveal a similar shape in some studies (Fig. 1A of [26] , Fig. 2 of

[23] and Fig. 2 of [17]) showing that the arch-shape of gamma is observed for at least some

achromatic stimuli. The arch shape described here is also visible when colored natural images

were presented (Fig. 2 of [27] and Fig. 2 of [9]). Interestingly, some papers (Fig. 1 of [5]; Fig.

9 of [26]) have shown traces with an inverted arch shape. It is unclear why this is the case;

potential reasons could be due to differences in the position of the reference wire, cortical depth

of recording, or differences in the conductance level. For example, arch-shaped gamma

waveform is observed in biophysical ING-PING and spiking models as well (Fig. 4 of [10];

Fig. 4 of [28]), in which non-sinusoidal gamma arises from the dynamics of the voltage-gated

conductances, and the specific shape produced by the model varies when the conductances are

varied. LFP waveform also changes with the recording depth (Figure 4B versus 4C of [10]).

Non-sinusoidal waveforms are common in neural rhythms, such as the sawtooth-shaped theta

rhythm in the hippocampus and comb-shaped beta rhythm, arising from a variety of different

biophysical phenomena [21].

Generalizing the arch shape and the associated phase-difference between gamma and

its first harmonic observed in the fullscreen iso-luminant chromatic stimuli used in this study

to other visual stimuli would be challenging, especially for stimuli that produce less gamma

powers such as achromatic gratings of smaller sizes. Indeed, for stimuli such as small

achromatic gratings that do not produce strong gamma but produce high firing rate (leading to

high broadband 'pedestal'), estimation of the phase of the gamma or harmonic may not be

possible at all, even though the two models used in this study had been designed to explain the

firing rates and gamma observed when achromatic gratings were presented. This is because

unlike quantities such as change in power, for which the 'pedestal' can be subtracted out by

either using the power spectral density at baseline period (as done here) or by explicitly

modeling the pedestal [29], the overall phase at a particular frequency has contributions from

both the true gamma signal and the broadband pedestal. Phase analysis is therefore susceptible

to errors and increased variability caused by high broadband noise, especially at low gamma

powers (as evident from the larger error bars at low powers in Figure 2B).

Although our results may not be generalizable to stimuli that do not produce very strong

gamma, they may nevertheless be useful when comparing two different schemes or principles

that produce oscillations. As discussed below, the two models use two very different principles

– the JXK model is a damped oscillator which is driven by stochastic noise and therefore cannot

produce gamma that is structured over long timescales, while oscillations in the JS arise from

a resonance phenomenon which can maintain a consistent shape over multiple cycles.

Therefore, by showing that gamma oscillations have a consistent arch shape (even for some

classes of stimuli), a resonant type architecture can be favored over a noise-driven architecture.

<u>Gamma oscillations in the JXK model</u>

The JXK model, like another model proposed by Kang and colleagues [18], explains

the "bursty" nature of gamma oscillations, which was shown in previous studies [15,17]. These

studies observed that gamma band power varied during a trial, with short intervals (~130 ms)

of high gamma power, interspersed by durations of no prominent power. Based on these, Xing

and colleagues [17] suggested that gamma rhythms were the outcome of cortical network

resonating to the stochastic component in the inputs. During periods in a trial when stochastic

variations in input are less, the cortical network would tend to asymptotically stabilize and

oscillations would get weaker over time. Xing and colleagues [17] demonstrated this

mechanism in a piecewise linear asymptotically stable model with noisy inputs. The JXK

model [2] emulated gamma burst-activity exhibiting the observed dependencies of gamma

power and frequency on stimulus parameters such as size and contrast by varying input drives

and recurrent excitation to the neural populations. However, the JXK model does not produce

a consistent waveform shape, because despite the presence of a rectification component, the

model operates mostly in a linear (post-rectification) domain for all inputs simulated. In this

domain the step response would be sinusoidal oscillations damping exponentially over time.

Although the presence of Poisson inputs distorted the waveform and prolonged oscillations for

a longer duration, these distortions are not consistent across different cycles of oscillation. This

was reflected as wide band of activity around the gamma band in the PSD (Figure 6C-D), and

a lack of discernible harmonic activity. Therefore, while these models adequately explained

the bursty nature of gamma oscillations, it is not surprising that they failed to produce a specific

gamma waveform.

However, a recent study has shown that the duration of gamma 'burst' activity reported

in Xing and colleagues [17] could have been underestimated [30]. This error arises because the

spectral estimator that is used to compute gamma power is itself noisy, causing the rhythm to

appear bursty in the spectral domain (see [30] for details). Chandran KS and colleagues [30]

used Matching Pursuit algorithm with a Gabor dictionary, which chooses the best-matching

template to estimate the duration of a gamma burst in the time domain itself to show that

gamma bursts during presentation of achromatic grating were longer than reported earlier

(median of ~300 ms, although the mode was ~100 ms). Along with the results presented here,

this suggests that the gamma rhythm could have more temporal regularity than what is

predicted by these stochastic models.

However, we note that the non-sinusoidal nature of gamma rhythm described here

could be introduced in the JXK model by changing the activation function. Specifically, if a

non-linear activation function is used in the place of the piece-wise linear rectifier, the step

response of the system would itself be a specific non-sinusoidal waveform in subsequent cycles

of the resulting damped oscillation. When stochastic perturbations are added, the resultant

bursts of oscillatory activity could contain both irregular distortions caused by the noisily

fluctuating inputs and more regular distortions induced by non-linearity at specific parts of the

trajectory in each cycle. Therefore, although we show that our results are reproduced by the JS

model within the superlinear regime, it is possible that the JXK model with a suitable activation

function can also replicate our findings. Further, in the JXK model, we have used Poisson noise

as in the original paper [2] to study the inherent propensity to produce arch-shaped waveforms.

However, a more realistic input would be colored noise [31], and it would be interesting to

study the modulatory effects on the waveform shape in different cycles. Furthermore, the JXK

model, despite having a rectification in its activation, operates in the linear region in almost all

gamma cycles. A substitution of a smoother non-linearity with similar slope might still induce

distorted waveforms while still requiring noisy inputs to sustain oscillations. Indeed, while the

JS model generates the proper gamma shape within a sub-regime, the duration of gamma is as

long as the duration of the stimulus itself. Therefore, some stochasticity may be necessary to

replicate all features of gamma rhythmicity, including duration and shape.

In the JXK model, there were a few inputs that satisfied our phase-difference criteria

for 'in-regime' behavior. However, a close examination of the 'in-regime' trace (example in

Figure 6G) revealed that, in addition to the irregularly distorted gamma cycles, these inputs

had occasional isolated steep troughs. These steep troughs could be approximated by a negative

impulse, which could be decomposed as a series of sinusoids with their troughs aligned,

resulting in sporadic 180-degree phase-difference estimates. Since these distortions are of

lower frequency than gamma, the JXK model does not operate 'in-regime' for any input

combination.

<u>Gamma oscillations in the JS model</u>

The JS model is a two-dimensional WC model that operates in ISN mode, where the

strong recurrent excitation would lead to instability if the inhibitory population were

inactivated [20]. Using a WC model operating as ISN, Tsodyks and colleagues [20] were able

to model firing rate modulation and phase shift of CA1 interneurons in theta rhythm cycles,

and further demonstrated a paradoxical behaviour where reduction in external drive to

inhibitory population resulted in a transient increase in inhibitory activity. In V1, Ozeki and

colleagues [32] modelled surround suppression as an increase in inhibitory drive and found

that the ISN model showed a simultaneous reduction in network excitation and inhibition, as

observed in their experiments.

Jadi and Sejnowski [19] modelled gamma rhythm as limit cycles generated for a range

of inputs by a local instability and observed that the observed stimulus dependencies of gamma

amplitude and frequency could be simulated while the model operated in a certain range of

inputs, which resulted in the local activation of inhibitory population to be strongly superlinear.

Superlinear activation in individual neurons have been used to explain normalization and size-

tuning in a network model [33]. Stronger super linearity of inhibitory population could have

contributions from their higher firing rates and a higher convergence of lateral excitatory

inputs.

In the current study, we could observe that the superlinear regime contains a subregion

where the waveform shape of limit cycles is the same arch shape observed during presentations

of our fullscreen hues and gratings. Further, we note that for proxies $-r_E-r_I$ and $-r_E$, the in-regime

input combinations include the higher values of inhibitory drive (I_I) (see Figure 7 and Figure

8), which in the model, corresponds to surround suppression during presentation of larger size

stimuli such as the full-field stimuli used in our experiments.

<u>Limitations and future directions</u>

While the LFP simulated in the JS model exhibited the desired gamma-harmonic phase

differences and size-contrast effects, there are several limitations. First, the regime for the arch

shape was obtained for a highly restrictive set of inputs. Second, the JS oscillations showed

very weak distortions from a sinusoid (low harmonic power). Finally, the results were tested

only for a particular type of activation function as described in the JS model. We found that

changing the type of activation function had a drastic effect on both JS and JXK models (data

not shown). While it is possible that another type of activation function can produce stronger

harmonics over a larger input space, we have tried to remain as close as possible to the original

models and tested whether these models can satisfy the new constraints that are imposed by

the shape of gamma and the presence of harmonics.

It is likely that more complex models such as PING models [10] and rate models with

synaptic dynamics [11] can explain these results, since they often exhibit strong arch-

shaped waveforms (see Fig. 4 of [10] and Fig. 1 and Fig. 2 in [11]; the rate and synaptic

variables show inverted arch-shape with sharp crests, which yield arch-shaped

gamma in our LFP proxy from equation (1)). Indeed, a typical weak PING model

involves the activation of the E population that then excites the I population within a

few milliseconds, and the resulting inhibition shuts down both E and I cells, with the

cycle repeating once the E cells recover from the inhibition and fire again [12,34].

Some experimental studies have indeed shown that spikes tend to occur just before the trough

of gamma [35–37], with the pyramidal cells leading the interneurons by a few milliseconds

[38–40]. In the JS model as well, we found that the E population led the I population by about

50º or 2-3 ms in the superlinear regime. Further, the results were qualitatively similar when

just E or I or the sum of both were used as the LFP proxy (Figure 7 and Figure 8).

<u>Relationship between firing rates and LFP</u>

JXK and JS models used excitatory cell activity to study the power and frequency trends

of gamma. Here, we converted these firing rates to a "proxy" LFP to be compatible with the

real data. Because the extracellular potential is generated due to the spatial separation of

transmembrane currents (generating dipoles or multi-poles), as well as alignment of such

dipoles across neurons, LFPs are thought to mainly reflect the transmembrane currents of

pyramidal neurons [41]. Hasenstaub and colleagues [38], in their in-vivo study of ferret

prefrontal cortex, found that the postsynaptic potentials and the firing of regular-spiking

excitatory neurons and fast-spiking interneurons were both synchronized to the extracellular

gamma rhythm, but the activation of interneurons was more aligned with the troughs of the

oscillation. A modelling study [10] simulated a Leaky Integrate-and-Fire (LIF) network and

computed LFP by injecting the simulated synaptic currents on corresponding synapses

distributed over a population of morphologically realistic neuron models. The LFP, thus

simulated, was compared against various outputs of the LIF network (such as firing rates,

membrane potential and synaptic currents). They found that the sum of absolute values of

synaptic currents (with both AMPA and GABA currents weighted nearly equally) served as

the best proxy for LFP. This paradox of both depolarizing (sink) and hyperpolarizing (source)

postsynaptic currents adding up in the same polarity could be explained considering the spatial

organization of the synapses. In a cortical pyramidal cell, inhibitory synapses are typically

concentrated towards the soma, whereas excitatory synapses are distributed less densely along
the apical and the basal dendrites. In such an arrangement, a large proportion of the
depolarizing and the hyperpolarizing postsynaptic currents would themselves constitute
'synaptic' dipoles [25], whose effect on the LFP would be in the polarity of the source or sink
closest to the electrode. Consequently, the polarity (sign) of LFP proxy would vary with
recording depth (see Fig 4 in [10]). The arrays in our dataset were 1 mm long and therefore are
expected to be in superficial/granular layers where the firing of both E and I cells are expected
to produce a downward deflection in the LFP [10]. Hence in our simulations, the LFP proxy
was taken as the sum of excitatory and inhibitory contributions, both with a negative polarity.

<u>Gamma and Harmonics vs slow/fast gamma</u>

Murty and colleagues [3] showed that fullscreen achromatic gratings produced two
different oscillations simultaneously – a fast gamma (40-70 Hz), which has been observed
previously for smaller sized gratings, and a slow gamma (20-40 Hz) which is prominent only
for large stimuli. Even though this also generates two bumps in the PSD, fast gamma is not a
harmonic of the slow gamma. First, the center frequency of fast gamma was not twice that of
slow gamma. Second, slow and fast gamma were not co-tuned: they had distinct orientation,
contrast, temporal frequency and size tuning preferences.

The arch-shape shown here poses additional problems for phase coding schemes in
which spike position relative to the gamma phase is used to code information [36,42], because
spikes tend to occur near the trough of the rhythm and having a sharper trough reduces the
operating range. Other studies have proposed that gamma oscillations may not play a role but
could be a useful marker/indicator of cortical processing [24]. In that framework, properties
such as shape and duration along with stimulus tuning could provide additional clues about the
underlying circuitry.

Methods and Models

<u>Data Acquisition</u>

We used LFP data recorded by Shirhatti and Ray [7] from V1 of two female macaques, whom we refer to as M1 and M2 (correspond to M1 and M3 in the earlier paper). Data from the third monkey were not considered in this paper since it showed relatively weaker gamma and harmonic power and very strong broadband high-gamma power, which caused systematic biases in the harmonic peak-frequency estimation. However, even for this monkey, the phase relationship between gamma and its harmonic (as shown in Fig 2B) was around 180º, similar to the other two monkeys.

Both monkeys were fitted with a titanium headpost and trained to perform a visual passive fixation task, after which a Utah array (96 and 81 electrodes for M1 and M2) was implanted in V1 (details of the surgery and implants are provided in [7]). The raw signals from microelectrodes were recorded using the 128-channel Cerebus neural signal processor (Blackrock Microsystems). LFP was obtained by filtering the raw signals online between 0.3 Hz and 500 Hz (Butterworth filters; first-order analog and fourth-order digital respectively), and recorded at 2 kHz sampling rate and 16-bit resolution. No further offline filtering was performed on this data before analysis.

Multiunit activity was also extracted from the raw signal by filtering online between 250 Hz and 7.5 kHz (Butterworth filters; fourth-order digital and third-order analog respectively), and subjecting the resultant signal to an amplitude threshold of ~5 SD of the signal. The recorded units were found to have receptive fields located in the lower left quadrant of the visual field with respect to fixation in both monkeys, at an eccentricity of ~$3°$–$4.5°$ in M1 and ~$3.5°$– $4.5°$ in M2. Full-field iso-luminant hues did not drive the neurons well, and therefore we did not get usable spiking activity from most electrodes (see Supplementary Figure 1 of [7] for description of spiking activity). In our current study, all the analyses were performed only on LFP data.

  Experimental setup and behavior

During the experiment, the monkey was seated in a monkey chair with its head held

stationary by the headpost. The monkey viewed a monitor (BenQ XL2411, LCD, 1,280 × 720

resolution, 100 Hz refresh rate) placed ∼50 cm from its eyes. The monkey and the display

setup were housed in a Faraday enclosure with a dedicated grounding separate from the main

supply ground to provide isolation from external electrical noise. The monitor was calibrated

and gamma-corrected using i1Display Pro (x-rite PANTONE) to obtain a mean luminance of

60 cd/m^2 on its surface and to obtain a gamma of unity for each of the three primaries of the

color gamut, which had the following CIE chromaticity xy coordinates: red, (0.644, 0.331);

green, (0.327, 0.607); blue, (0.160, 0.062). The white point was at (0.345, 0.358).

Each monkey performed a passive fixation task, which required them to fixate at a small

dot of 0.05°–0.10° radius at the center of the screen throughout the trial (3.3 or 4.8 s duration;

fixation spot was displayed throughout). Each trial began with fixation, following which an

initial blank grey screen of 1,000 ms was displayed, and then, two to three stimuli were shown

for 800 ms each with an interstimulus interval of 700 ms. The monkey was rewarded with juice

for maintaining fixation within 2° from the fixation point. Trials in which fixation was broken

were not considered in our analyses. Eye position data was recorded as horizontal and vertical

coordinates using the ETL-200 Primate Eye Tracking System (ISCAN) and monitored

throughout the task using custom software running on macOS, which also controlled the task

flow, generated stimuli, and randomized stimuli presentation. Although the monkeys were

required to maintain fixation within 2° of the fixation spot, they were able to hold their fixation

well within these limits (standard deviation < 0.4° across all sessions for all monkeys) during

the task. Since the stimuli were all full screen, the small deviations are unlikely to affect the

results.

 Stimuli

The stimuli consisted of 36 hues and 1 achromatic grating. The hues were equally

spaced along the circular hue space of the standard HSV nomenclature (0° hue to 350° hue,

where 0°, 120°, and 240° represent red, green, and blue respectively), which were displayed

full screen and at full saturation and value. The achromatic grating was at an orientation of 90°

and had a spatial frequency of 4 cpd for M1 and 2 cpd for M2. These grating parameters were

optimized to capture strong fast gamma with minimal slow gamma (see [3]). The full-screen

stimuli, in our setup, subtended a visual angle of ∼56° in the horizontal direction and ∼33° in

the vertical direction.

Electrode selection

As with our previous report [7], electrodes were considered for analysis only if they

gave consistent stimulus-induced changes and reliable receptive field estimates across sessions,

determined by a receptive field mapping protocol that was run across multiple days [43].

Further, we discarded signals from electrodes with unusable or inconsistent signals, a high

degree of crosstalk with other electrodes, or impedances outside the range of 250–2,500 KΩ

for monkey M1 and 125–2,500 KΩ for M2. This resulted in 64 and 16 usable electrodes for

M1 and M2, respectively.

Data Analysis

Stimulus presentations with excessive recording artifacts (<5.9% and <5.0% of

presentations of each stimulus in M1 and M2) were discarded for each session. The typical

artifacts were large deflections in the recorded traces observed across electrodes, usually

accompanying occasional physical movements from the subjects (jaw movements, for

instance). A trial was rejected from our analyses if the corresponding LFP trace showed

deflections about the mean greater than 6 times the standard deviation of the traces during the

stimulus period. This process removed prominent artifacts and yielded 19.9±6.2 repeats in M1

and 20.2±1.0 repeats in M2 per stimulus.

Spectral analysis of LFP

For each stimulus, LFP recorded from -500 to 0 ms from stimulus onset was taken as

the 'baseline period' and 250 to 750 ms from stimulus onset was taken as the 'stimulus period'

to avoid the transient responses to the stimulus onset. This yielded a frequency resolution of 2

Hz in the Power Spectral Density (PSD). PSD was computed using the Multitaper method

using the Chronux toolbox [44], with three tapers. The change in power was calculated as 10

times the difference between base-10 logarithm of PSDs at stimulus period and baseline period,

expressed in decibels (dB). Estimation of peak frequencies was done on these baseline-

corrected PSDs.

Gamma range was taken as 30-70 Hz, and 'gamma peak frequency' was estimated as

the highest peak within this range. In most stimulus conditions tested, a discernible 'second

bump' was observed in the baseline-corrected PSD. The peak frequency of the second bump

was estimated as the highest peak occurring beyond 12 Hz after the estimated gamma peak

frequency up to 140 Hz, to exclude higher frequency bumps.

Analysis of gamma and harmonic phases

To compute the phase difference of gamma and its first harmonic, gamma and harmonic

signals were extracted from LFP during the stimulus period by bandpass filtering using

separate Butterworth filters (zero-phase; order 4). The passband for gamma was 20 Hz wide,

centered around the gamma peak frequency, identified from trial-averaged PSD for a given

stimulus and an electrode. The passband for the first harmonic of gamma was also 20 Hz wide

but centered around twice the corresponding gamma peak frequency. The phases of these

signals (φ_{gamma} and $\varphi_{harmonic}$) were then computed using Hilbert transform. The phase difference

between gamma and its harmonic was calculated as:

Phase difference $= |2*\varphi_{gamma} - \varphi_{harmonic}|$ (1)

The above phase difference estimate was computed at each timepoint within the

stimulus period of every trial for each electrode. Note that the 20 Hz passband for gamma and

its harmonic-band filters is wide enough so that errors in gamma peak-frequency detection do

not affect the phase difference estimates.

<u>Effect of harmonic phase on gamma waveform</u>

To illustrate the effect of the gamma-harmonic phase relationship on the shape of

gamma waveform, gamma and its first harmonic were mimicked using sinusoids and the initial

phase of gamma (φ_G) was varied. The formulation of gamma and harmonics were as follows:

Gamma wave, $G = \cos(2\pi\, 45\, t + \varphi_G)$

Harmonic wave, $H = \frac{1}{4}* \cos(2\pi\, 90\, t)$

Sum of gamma and harmonic $= G + H$ (2)

These waveforms are displayed in Figure 4C along with the phase difference (Figure

4D) obtained by applying equation (1).

<u>Experimental Design and Statistical Analysis</u>

First, we tested our hypothesis that the second bump in PSD accompanying gamma was

indeed its harmonic by computing the ratio of harmonic to gamma peak-frequencies, obtained

from the trial-averaged change in power (dB) from baseline spectra for each electrode. The

ratios were subsequently subjected to a non-parametric Wilcoxon signed rank test (Null

hypothesis: median ratio = 2). The standard error (SE) of the median of the gamma-harmonic

frequency ratio was estimated by bootstrapping over N iterations (where N is the number of

datapoints). This involved random sampling with replacement of the ratio data N times and

estimating their median each time, which resulted in N medians, whose standard deviation (SD)

is reported as the standard error (SE).

Circular statistics on the gamma-harmonic phase difference data were computed in

Matlab using the Circular Statistics toolbox [45]. The mean phase differences are reported as

MEAN ± CI in Figure 5, where MEAN is the circular mean and CI is the 95% confidence

interval of MEAN, under a Von Mises distribution (implemented in *circ_mean* and

circ_confmean functions of the Circular Statistics toolbox).

To validate that the distribution of gamma-harmonic phase differences was non-

uniform and, thus, assess the validity of circular mean estimates in Figure 5, we subjected trial-

averaged phase-differences gathered from all electrodes for a given stimulus to a Rayleigh test

of non-uniformity (Null hypothesis: uniform distribution of phases). When the gamma activity

recorded for a given stimulus has a specific waveform across electrodes, the corresponding

distribution of gamma-harmonic phase-differences will be unimodal. We considered this to be

the case when the p-value of the Rayleigh test was less than 0.01.

<u>Studying non-sinusoidal waveforms in mean-field models of gamma</u>

To understand the factors resulting in the characteristic waveform of gamma, we

investigated the gamma-harmonic phase relationship emerging in two mean-field models

introduced in earlier works, which produced gamma oscillations with power and frequency

trends as observed in experiments. We explored the waveform in these models by identifying

input regimes that produced a first-harmonic with a specific phase difference from the

fundamental of gamma, as required for its arch-shape. We assessed the presence of such

regimes in two models. The first model operated linearly and produced oscillations by virtue

of stochastic time-varying inputs (Jia-Xing-Kohn or JXK model; [2]), while the second model

had non-linear dynamics, which gave rise to gamma frequency limit cycles in response to

constant inputs (Jadi and Sejnowski or JS model; [19]).

Jia-Xing-Kohn (JXK) model

Jia and colleagues [2]defined a linear EI rate-model and extended it by adding a Global

Excitatory Population (G) to approximate recurrent excitatory feedback within V1[2]. The

model, when subjected to step input, acts as a damped oscillator but produces oscillations by

virtue of constant perturbation from the Poisson inputs. The original model, formulated in the

paper, incorporates detailed stimulus descriptions, namely Masked Noise (MN) level and

stimulus size (r) and contrast (c). The stimulus size parameter (r) scales the extent of Network

recurrence (G) in steady-state proportionally as larger size stimuli excite a larger area of

cortical cells, increasing the global feedback. The stimulus contrast (c) proportionally increases

the drives to the excitatory and inhibitory populations, I_E and I_I respectively. In our simulations,

the Masking Noise level (MN) has been set to 0, resulting in the following reduced formulation.

The resultant reduced JXK model formulation is as follows:

$$\tau_E \frac{dr_E}{dt} = -r_E + W_{EE}\lfloor r_E \rfloor - W_{EI}\lfloor r_I \rfloor + W_{EG}\lfloor r_G \rfloor + E_{inp}$$

$$\tau_I \frac{dr_I}{dt} = -r_I + W_{IE}\lfloor r_E \rfloor - W_{II}\lfloor r_I \rfloor + W_{IG}\lfloor r_G \rfloor + I_{inp}$$

$$\tau_G \frac{dr_G}{dt} = -r_G + r^2.W_{GE}\lfloor r_E \rfloor$$

where $\lfloor x \rfloor = x\ if\ x > 0\ and\ 0\ otherwise,$ and E_{inp} and I_{inp} approximate a Poisson

process with average rates IE and II respectively, which vary with stimulus contrast (c) as:

$$I_E = 40.\frac{c^2}{c^2+0.3^2} ; I_I = 32.\frac{c^2}{c^2+0.3^2} \qquad (3)$$

Parameter values of this model are given in Table 1.

Jadi-Sejnowski (JS) model

Jadi and Sejnowski [19] used a simple rate model to qualitatively describe stimulus size
and contrast dependent modulation of gamma. The model consisted of an excitatory and an
inhibitory population with sigmoidal activation, operating as an Inhibition Stabilized Network
(ISN), and was constrained the input drives to the populations to reproduce the increase in
power and decrease in peak frequency of gamma with increasing stimulus size as earlier studies
have observed in V1 [5,6].

The model defines the population firing rates of Excitatory and Inhibitory populations
as follows:

$$\tau_E \frac{dr_E}{dt} = -r_E + \sigma_E(W_{EE}r_E - W_{EI}r_I + I_E)$$

$$\tau_I \frac{dr_I}{dt} = -r_I + \sigma_I(W_{IE}r_E - W_{II}r_I + I_I)$$

$$\sigma_P(x) = \frac{1}{1+exp(m_P.(\theta_P-x))} - \frac{1}{1+exp(m_P.\theta_P)} \tag{4}$$

In this model, larger stimuli were simulated by an increased inhibitory drive (I_I) to the
population, owing to suppression from the surrounding populations [32], whereas stimuli of
different contrasts could be simulated by applying co-varying drives to inhibitory (I_I) and
excitatory (I_E) populations [46].

Oscillations in this model emerged after the system underwent a supercritical Hopf
bifurcation at sufficiently high input drives. Since the model operated close to the Hopf
bifurcation, the amplitude and frequency of oscillations in firing rates could be closely
approximated by linearization of the model (For the interested readers, we refer to Appendix-
B in [19] for their detailed discussion). The authors deduced and demonstrated that the model

gave rise to the observed trends in the gamma when the inputs were such that the inhibitory

population was strongly 'superlinear'. This means that the summed inputs to the inhibitory

population (from recurrent and external sources; argument of σ_I in equation (4)) must lie in a

certain range of values where the activation function σ_I curves upwards (increasing in slope

with increasing summed input). For the sigmoidal activation function used in equation (4), the

summed inputs must operate in the lower half of the sigmoid. Superlinear activation of the

excitatory population, on the other hand, was antagonistic and, hence, excitatory activation

should not be strongly superlinear. The set of such inputs constitute the operating regime of

the model, which we refer to as the 'superlinear' regime. In Figure 7C-D, we have presented

examples of the size and contrast effects on gamma rhythm in the model. The example input

combinations (indicated by markers in Figure 7A-B) were chosen such that they lay within the

'superlinear' regime and were varied based on the above criteria to represent different sizes

and contrasts.

<u>Identifying the operational input regimes using gamma-harmonic phase difference</u>

Because the gamma-harmonic relationship was studied using LFP recordings in real

data while the model simulations yielded firing rates of excitatory and inhibitory populations,

these firing rates had to be converted to a 'proxy' LFP. The relationship between the two is

complex [10,38,41]. For simplicity, we used the negative sum of population firing rates of the

excitatory and inhibitory populations in the models as a proxy for LFP, since the LFP, being

the extracellular potential, could be approximated to vary inversely with the depolarization

(excitability) of these populations. Furthermore, the spatial organization of the synapses are

likely to produce strong dipole contributions with negative polarity to the LFP recorded in our

setup. We elaborate this further in the Discussion section. We also approximated the LFP as

the negative of just the excitatory or inhibitory populations, which yielded qualitatively similar

results.

The environment used for simulation was Matlab 2019b (Mathworks,

RRID:SCR_001622), where the models were simulated by a forward Euler method, with

parameters as in Table 1. Simulations were run for a duration of 2 seconds in time-steps of 0.1

ms, on all pairs of 41 excitatory and 41 inhibitory input values sampled from the ranges

specified in Table 1. In each case, PSD of the LFP proxy was computed between 1-2 seconds

to avoid the initial slow transient. The gamma and harmonic peak frequencies were identified

as frequencies containing maximum power in the 30-70 Hz range and twice the gamma peak

frequency, respectively. For further analyses in the JS model, only those input combinations

were considered for which gamma and harmonic frequency amplitudes were greater than 1e-3

and 1e-6 units respectively, to ensure phase analysis of oscillatory activity was not dominated

by simulation errors or noise (grey region in Figure 7F). The LFP proxy signal was filtered

using 20 Hz passbands centered at each frequency identified above. The gamma-harmonic

phase difference (equation (1)) was computed, just as for LFP in macaque data. An input point

(pair of values supplied as excitatory and inhibitory input drives; stimulus parameters in the

case of JXK) was said to be 'in-regime' if the gamma-harmonic phase difference was within

22.5 degrees from 180-degrees (ideal for arch shape; Figure 4). Since JXK takes stochastic

inputs, to assess if there is indeed a unique gamma-harmonic phase difference at each of the

mean input-drive combinations tested and whether the model can indeed exhibit in-regime

behavior, we ran the simulation 50 times (analogous to 50 different trials) for each stimulus

condition. In each iteration, we considered the mean phase-difference over the 1-2 second

interval. For every selected input combination, the mean phase difference over this interval in

each iteration was taken and the resultant pool of 50 mean phase differences was subjected to

Rayleigh test of uniformity to identify those inputs that have a 'consistent' gamma-harmonic

phase relationship across iterations (p-value < 0.01). Among the qualifying input combinations,

those input drive pairs for which the circular mean of mean phase differences across iterations

lay within 180±22.5 degrees were deemed to be 'in-regime'. For JXK, the gamma-harmonic

phase reported in Figure 6 is the circular average of the phase differences from all the iterations.

The amplitudes of gamma and its first harmonic in the figure are estimated from the average

PSD (examples in Figure 6C-D), computed as the mean of the PSDs obtained from 50 iterations

for a given stimulus combination.

References

1. Buzsáki G. Rhythms of the brain. New York, NY, US: Oxford University Press; 2006.

xv, 448 p. [Rhythms of the brain].

2. Jia X, Xing D, Kohn A. No consistent relationship between gamma power and peak

frequency in macaque primary visual cortex. J Neurosci. 2013 Jan 2;33(1):17–25.

3. Murty DVPS, Shirhatti V, Ravishankar P, Ray S. Large Visual Stimuli Induce Two

Distinct Gamma Oscillations in Primate Visual Cortex. J Neurosci. 2018

14;38(11):2730–44.

4. Ray S, Maunsell JHR. Differences in Gamma Frequencies across Visual Cortex Restrict

Their Possible Use in Computation. Neuron. 2010 Sep 9;67(5):885–96.

5. Gieselmann MA, Thiele A. Comparison of spatial integration and surround suppression

characteristics in spiking activity and the local field potential in macaque V1. Eur J

Neurosci. 2008 Aug;28(3):447–59.

6. Ray S, Maunsell JH. Different origins of gamma rhythm and high-gamma activity in

macaque visual cortex. PLoS biology. 2011;9(4).

7. Shirhatti V, Ray S. Long-wavelength (reddish) hues induce unusually large gamma

oscillations in the primate primary visual cortex. PNAS. 2018 Apr 24;115(17):4489–94.

8. Peter A, Uran C, Klon-Lipok J, Roese R, van Stijn S, Barnes W, et al. Surface color and

predictability determine contextual modulation of V1 firing and gamma oscillations.

Colgin L, editor. eLife. 2019 Feb 4;8:e42101.

9.   Bartoli E, Bosking W, Chen Y, Li Y, Sheth SA, Beauchamp MS, et al. Functionally

Distinct Gamma Range Activity Revealed by Stimulus Tuning in Human Visual Cortex.

Current Biology. 2019 Oct 21;29(20):3345-3358.e7.

10.   Mazzoni A, Lindén H, Cuntz H, Lansner A, Panzeri S, Einevoll GT. Computing the

Local Field Potential (LFP) from Integrate-and-Fire Network Models. PLOS

Computational Biology. 2015 Dec 14;11(12):e1004584.

11.   Keeley S, Byrne Á, Fenton A, Rinzel J. Firing rate models for gamma oscillations.

Journal of neurophysiology. 2019;121(6):2181–90.

12.   Tiesinga P, Sejnowski TJ. Cortical Enlightenment: Are Attentional Gamma Oscillations

Driven by ING or PING? Neuron. 2009 Sep 24;63(6):727–32.

13.   Chariker L, Shapley R, Young L-S. Rhythm and Synchrony in a Cortical Network

Model. J Neurosci. 2018 Oct 3;38(40):8621–34.

14.   Wilson HR, Cowan JD. Excitatory and inhibitory interactions in localized populations

of model neurons. Biophys J. 1972 Jan;12(1):1–24.

15.   Burns SP, Xing D, Shelley MJ, Shapley RM. Searching for Autocoherence in the

Cortical Network with a Time-Frequency Analysis of the Local Field Potential. J

Neurosci. 2010 Mar 17;30(11):4033–47.

16.   Burns SP, Xing D, Shapley RM. Is Gamma-Band Activity in the Local Field Potential

of V1 Cortex a "Clock" or Filtered Noise? J Neurosci. 2011 Jun 29;31(26):9658–64.

17.   Xing D, Shen Y, Burns S, Yeh C-I, Shapley R, Li W. Stochastic Generation of Gamma-

Band Activity in Primary Visual Cortex of Awake and Anesthetized Monkeys. J

Neurosci. 2012 Oct 3;32(40):13873–13880a.

18.   Kang K, Shelley M, Henrie JA, Shapley R. LFP spectral peaks in V1 cortex: network

resonance and cortico-cortical feedback. J Comput Neurosci. 2010 Dec 1;29(3):495–

507.

19.   Jadi MP, Sejnowski TJ. Regulating Cortical Oscillations in an Inhibition-Stabilized

Network. Proc IEEE Inst Electr Electron Eng. 2014 Apr 21;102(5).

20.   Tsodyks MV, Skaggs WE, Sejnowski TJ, McNaughton BL. Paradoxical effects of

external modulation of inhibitory interneurons. J Neurosci. 1997 Jun 1;17(11):4382–8.

21.   Cole SR, Voytek B. Brain Oscillations and the Importance of Waveform Shape. Trends

Cogn Sci (Regul Ed). 2017;21(2):137–49.

22.   Henrie JA, Shapley R. LFP Power Spectra in V1 Cortex: The Graded Effect of Stimulus

Contrast. Journal of Neurophysiology. 2005 Jul 1;94(1):479–90.

23.   Jia X, Kohn A. Gamma rhythms in the brain. PLoS Biol. 2011 Apr;9(4):e1001045.

24.   Ray S, Maunsell JHR. Do gamma oscillations play a role in cerebral cortex? Trends in

Cognitive Sciences. 2015 Feb 1;19(2):78–85.

25.   Buzsáki G, Anastassiou CA, Koch C. The origin of extracellular fields and currents--

EEG, ECoG, LFP and spikes. Nat Rev Neurosci. 2012 May 18;13(6):407–20.

26.   Jia X, Smith MA, Kohn A. Stimulus Selectivity and Spatial Coherence of Gamma

Components of the Local Field Potential. J Neurosci. 2011 Jun 22;31(25):9390–403.

27.   Brunet N, Bosman CA, Roberts M, Oostenveld R, Womelsdorf T, De Weerd P, et al.

Visual cortical gamma-band activity during free viewing of natural images. Cereb

Cortex. 2015 Apr;25(4):918–26.

28. Lowet E, Roberts MJ, Bonizzi P, Karel J, Weerd PD. Quantifying Neural Oscillatory

Synchronization: A Comparison between Spectral Coherence and Phase-Locking Value

Approaches. PLOS ONE. 2016 Jan 8;11(1):e0146443.

29. Donoghue T, Haller M, Peterson EJ, Varma P, Sebastian P, Gao R, et al. Parameterizing

neural power spectra into periodic and aperiodic components. Nat Neurosci. 2020

Dec;23(12):1655–65.

30. Chandran KS S, Seelamantula CS, Ray S. Duration analysis using matching pursuit

algorithm reveals longer bouts of gamma rhythm. Journal of Neurophysiology. 2017

Nov 8;119(3):808–21.

31. Lerchner A, Ahmadi M, Hertz J. High-conductance states in a mean-field cortical

network model. Neurocomputing. 2004 Jun 1;58–60:935–40.

32. Ozeki H, Finn IM, Schaffer ES, Miller KD, Ferster D. Inhibitory stabilization of the

cortical network underlies visual surround suppression. Neuron. 2009 May

28;62(4):578–92.

33. Rubin DB, Van Hooser SD, Miller KD. The Stabilized Supralinear Network: A

Unifying Circuit Motif Underlying Multi-Input Integration in Sensory Cortex. Neuron.

2015 Jan 21;85(2):402–17.

34. Buzsáki G, Wang X-J. Mechanisms of Gamma Oscillations. Annual Review of

Neuroscience. 2012;35(1):203–25.

35. Chalk M, Herrero JL, Gieselmann MA, Delicato LS, Gotthardt S, Thiele A. Attention

reduces stimulus-driven gamma frequency oscillations and spike field coherence in V1.

Neuron. 2010 Apr 15;66(1):114–25.

36. Vinck M, Lima B, Womelsdorf T, Oostenveld R, Singer W, Neuenschwander S, et al.

815   Gamma-Phase Shifting in Awake Monkey Visual Cortex. J Neurosci. 2010 Jan

816   27;30(4):1250–7.

817   37. Das A, Ray S. Effect of Stimulus Contrast and Visual Attention on Spike-Gamma Phase

818   Relationship in Macaque Primary Visual Cortex. Front Comput Neurosci. 2018;12:66.

819   38. Hasenstaub A, Shu Y, Haider B, Kraushaar U, Duque A, McCormick DA. Inhibitory

820   Postsynaptic Potentials Carry Synchronized Frequency Information in Active Cortical

821   Networks. Neuron. 2005 Aug 4;47(3):423–35.

822   39. Csicsvári J, Jamieson B, Wise KD, Buzsáki G. Mechanisms of gamma oscillations in

823   the hippocampus of the behaving rat. Neuron. 2003 Jan 23;37(2):311–22.

824   40. Vinck M, Womelsdorf T, Buffalo EA, Desimone R, Fries P. Attentional modulation of

825   cell-class-specific gamma-band synchronization in awake monkey area v4. Neuron.

826   2013 Nov 20;80(4):1077–89.

827   41. Einevoll GT, Kayser C, Logothetis NK, Panzeri S. Modelling and analysis of local field

828   potentials for studying the function of cortical circuits. Nat Rev Neurosci. 2013

829   Nov;14(11):770–85.

830   42. Fries P, Nikolić D, Singer W. The gamma cycle. Trends Neurosci. 2007 Jul;30(7):309–

831   16.

832   43. Dubey A, Ray S. Cortical Electrocorticogram (ECoG) Is a Local Signal. J Neurosci.

833   2019 May 29;39(22):4299–311.

44.   Bokil H, Andrews P, Kulkarni JE, Mehta S, Mitra PP. Chronux: A platform for

analyzing neural signals. Journal of Neuroscience Methods. 2010 Sep 30;192(1):146–

51.

45.   Berens P. CircStat: A MATLAB Toolbox for Circular Statistics. Journal of Statistical

Software. 2009 Sep 23;31(1):1–21.

46.   Priebe NJ. Mechanisms of Orientation Selectivity in the Primary Visual Cortex. Annual

Review of Vision Science. 2016;2(1):85–107.

Figures

Figure 1: Visual Stimulation induces narrowband activity in gamma and its harmonic bands. (A-D, *top row*) PSDs for baseline (-500 to 0 ms; 0 indicates stimulus onset) and stimulus (250 to 750 ms) periods for different stimuli, averaged across trials and electrodes, from monkey 1 (M1). The baseline PSDs are plotted in black, while the stimulus period PSDs are colored (corresponding to the hue presented) or grey (for fullscreen grating). (A-D, *bottom row*) show the change in power (dB) in the stimulus period from the baseline, computed from the PSDs in the top row. The gamma power computed from the change in power spectrum is

 represented by vertical dashed line). The peaks in gamma range and the second bump activity

 are marked in each plot. (E-H) Same as A-D, for monkey M2.

 Figure 2: Second bump in PSD corresponds to the first harmonic of gamma. Median of

 peak frequency ratios of the second bump with respect to the gamma band, scattered against

 average gamma power for each stimulus (circles are colored to represent the presented hues;

 grating stimulus is represented by the grey circle). Error bars indicate the standard error of

 median (estimated by bootstrapping). Open circles mark stimuli whose ratios were found to be

 significantly different from 2 (p-value < 0.01 without any Bonferroni correction; Wilcoxon

 signed-rank test). The dotted horizontal lines correspond to the least-error margin 2±0.05.

Figure 3: Gamma waveform has consistent non-sinusoidal shape under different stimuli.

(A-D) present LFP traces for different stimuli, shown in Figure 1, from an example trial and

electrode in M1. (E-H) shows LFP traces obtained from M2 for corresponding stimuli.

Figure 4: Observed arch-shape of gamma corresponds to gamma-harmonic phase

difference of 180°. (A) Gamma (blue) and harmonic (red) components, mimicked using

sinusoids as in equation (2) with different initial gamma phases (φ_G) in each row. (B) Gamma

waveform resulting from summation of gamma and its harmonic from the corresponding row

in (B). (C) Phase difference of gamma and its first harmonic, as given in equation (1), computed

from filtering and Hilbert transforming gamma and harmonic bands from the summed

waveform in (B) in each row. Note that the arch-shaped gamma observed in recordings arises

from a phase difference of 180° (middle row).

Figure 5: Gamma-harmonic phase differences in different stimuli are concentrated near
180°. Circular mean of trial-averaged gamma-harmonic phase differences (across all
electrodes) are scattered against the average gamma power for each stimulus (circles are
colored to represent the presented hues; grey for grating stimulus). For each stimulus, the pool
of trial-averaged phase differences from all electrodes is subject to Rayleigh test of non-
uniformity to check if phase-difference is consistent across electrodes so that circular mean
estimates are reliable (p-value < 0.01). If found reliable (even though it could be different from

180º), the stimulus is represented by a filled circle. Error bars represent the 95% confidence

interval of circular mean.

Figure 6: Gamma properties for the JXK model (A) Gamma peak amplitude and (B)

frequency found in the average PSD from 50 iterations at each stimulus size (r) and contrast

(c) combination. (C) Average PSD of LFP proxy traces generated at fixed contrast and different

sizes, across all iterations. (D) Average PSD of LFP proxy traces generated at different

contrasts at a fixed size. The stimuli values that generated these PSDs are indicated in plots

(A), (B) and (E) using colored markers (same colors as the plots in C and D). (E) Harmonic

band peak amplitude in the average PSD. The harmonic is taken to be twice the gamma

frequency. (F) Mean gamma-harmonic phase difference (equation (1)) across multiple

iterations, with in-regime inputs encircled by black contours. (G) Example LFP proxy activity

(top panel) and its gamma and first harmonic components (bottom panel) generated at the

highest contrast condition in (D) (identified as in-regime). This trace shows occasional steeper

troughs (bottom trace; near 1.35 s, 1.52 s and 1.58 s timepoints) although there is no consistent

trend between gamma and the harmonic.

Figure 7: JS model produces sustained arch-shaped gamma oscillations within the previously identified superlinear inhibitory regime. (A) Peak gamma amplitude and (B)

Peak gamma frequency in the LFP proxy generated by the JS model for each input-drive

combination. (C) PSD of LFP proxy generated for different stimulus sizes. (D) PSD of LFP

proxy generated for different contrast. Input combinations used in (C) and (D) are indicated by

markers, with same colors as traces, in (A), (B) and (E). (E) Harmonic amplitude identified

from the PSD of LFP proxy for each stimulus condition. (F) Phase difference of gamma and

its harmonic (equation (1)) computed for each simulated input drive pair. The black contours

encircle input drives identified to be 'in-regime' from their phase differences. The white

contour in each figure enclose the input domain identified by Jadi and Sejnowski (2014) to

replicate gamma power increase and peak frequency decrease in response to increasing

stimulus size, and it can be seen to enclose a significant portion of the 180° phase difference

regime (black contour). (G) LFP proxy activity trace corresponding to the largest stimulus size

shown in Figure 7C. The dotted horizontal line shown is equidistant from the minima and

maxima of the traces. A sinusoidal oscillation would have the same crest and trough width on

this line. The bottom panels show the gamma and its first-harmonic components, plotted in

blue and red respectively, filtered from the LFP proxy traces.

Figure 8: Gamma-harmonic phase difference in rate models using different LFP proxies.
(A) Phase difference of gamma and its harmonic in the JXK model computed for each stimulus condition (as in Figure 6F) but using only E population activity to compute the LFP proxy. (B) Gamma-harmonic phase differences in JXK using I population activity alone. (C) Phase differences in JS model using E population activity only. (D) Phase differences in JS using I population activity only. Black contours encircle stimulus conditions identified as 'in-regime'. White contours in (C) and (D) mark the superlinear regime of the JS model.

934 **Table 1: Parameter values used in rate models.**

JS

W_{EE}	W_{EI}	W_{IE}	W_{II}	τ_E, τ_I (ms)
16	26	20	1	20,10

m_E, m_I	θ_E, θ_I	I_E, I_I		
1,1	5,20	0 to 20 (steps of 0.5)		

JXK

W_{EE}	W_{EI}	W_{IE}	W_{II}	W_{EG}	W_{IG}	W_{GE}	τ_E, τ_I, τ_G (ms)
1.5	3.25	3.5	2.5	0.25	0.5	0.6	6,15,19

MN	r	c					
0	1 to 5 (steps of 0.125)	0.01 to 1 (logarithmic steps of 0.5)					