Evolution of phenotypic variance provides insights into the genetic basis of

adaption

Author list:

Wei-Yun Lai[1,2], Viola Nolte[1], Ana Marija Jakšić[1,2,3] and Christian Schlötterer[1*]

[1]Institut für Populationsgenetik, Vetmeduni Vienna, Vienna, Austria.

[2]Vienna Graduate School of Population Genetics, Vetmeduni Vienna, Vienna, Austria.

[3]Current affiliation: École polytechnique fédérale de Lausanne, Lausanne, Switzerland

[*]Correspondence: christian.schloetterer@vetmeduni.ac.at; Tel.: +43-1-25077-4300.

Keywords:

Phenotypic variance, temperature adaptation, *Drosophila simulans*, experimental evolution

Abstract

Most traits are polygenic and the contributing loci can be identified by GWAS. Their

adaptive architecture is, however, poorly characterized. Here, we propose a new approach to

study the adaptive architecture, which does not depend on genomic data. Relying on

experimental evolution we measure the phenotypic variance in replicated populations during

adaptation to a new environment. Extensive computer simulations show that the evolution of

phenotypic variance in a replicated experimental evolution setting is a powerful approach to

distinguish between oligogenic and polygenic adaptive architectures. We apply this new

method to gene expression variance in male *Drosophila simulans* before and after 100

generations of adaptation to a novel hot environment. The variance change in gene

expression was indistinguishable for genes with and without a significant change in mean

expression after 100 generations of evolution. We conclude that adaptive gene expression

evolution is best explained by a highly polygenic adaptive architecture. We propose that the

evolution of phenotypic variance provides a powerful approach to characterize the adaptive

architecture, in particular when combined with genomic data.

Introduction

It is widely accepted that most complex traits have a polygenic or even infinitesimal basis (Ayroles et al., 2009; Boyle et al., 2017; Liu et al., 2019). Nevertheless, it is difficult to predict which of these loci are responding to selection when a population is exposed to a new selection regime. If pleiotropic constraints are strong, only a small subset of the genes constituting are free to respond to respond to selection. Hence, the genetic basis of the adaptive response of a complex trait (i.e. adaptive architecture (Barghi et al., 2020)) may differ substantially from the genetic architecture. Since even for large phenotypic changes the genetic basis of an adaptive response is difficult to study when more than a handful of genes are contributing, we introduce a new approach to study the complexity of the adaptive architecture. Rather than aiming to map the contributing loci, we propose to study the evolution of phenotypic variance in an experimental evolution framework.

The phenotypic variance of a quantitative trait is a key determinant for its response to selection. It can be decomposed into genetic and environmental components (Falconer and Mackay, 1963). Over the past years, mathematical models have been developed which describe the expected genetic variance of a quantitative trait under selection and its maintenance in evolving populations (Bulmer, 1972; Chevalet, 1994; Kimura and Crow, 1964; Turelli, 1984). For infinitely large populations and traits controlled by many independent loci with infinitesimal small effect, changes in trait optimum are not expected to affect the phenotypic variance (Lande, 1976). A much more complex picture is expected when the effect sizes are not equal, the population size is finite, or the traits have a simpler genetic basis (Barton and Keightley, 2002; Barton and Turelli, 1987; Franssen et al., 2017; Hayward and Sella, 2019; Jain and Stephan, 2015; Keightley and Hill, 1989). For instance, for a trait with oligogenic architectures, the genetic variance could drop dramatically during

adaptation, while with polygenic architectures, only minor effects on the variance are

expected (Barton et al., 2017; Franssen et al., 2017; Jain and Stephan , 2015). These studies

suggest that a time-resolved analysis of phenotypic variance has the potential to shed light

onto the complexity of the underlying adaptive architecture.

Despite its potential importance for the understanding of adaption, we are faced with the

situation that very few empirical data are available for the evolution of phenotypic variance.

The use of natural populations to study changes in phenotypes and even more so phenotypic

variances is limited, as the environmental heterogeneity cannot be controlled. A

complementary approach to study the evolution of phenotypic variance is experimental

evolution (Kawecki et al., 2012). With replicated populations starting from the same founders

and evolving under tightly controlled environmental conditions, experimental evolution

provides an enormous potential to study the evolution of phenotypic variance.

Most experimental evolution studies in sexual populations focused on the evolution of

phenotypic means, rather than variance (Burke et al., 2010; Chippindale et al., 1996; Jakšić et

al., 2020; Mallard et al., 2018). A notable exception is a study which applied fluctuating,

stabilizing and disruptive selection to a small number of traits (wing shape) and observed a

change of the phenotypic variance (Pélabon et al., 2010). Instead of looking at a preselected

subset of phenotypes which limits the generality, we will focus on gene expression, a set of

molecular phenotypes, which can be easily quantified since microarrays and, more recently,

RNA-Seq have become available. Importantly, the expression levels of genes exhibit the

same properties (e.g.: continuality and normality) as other complex quantitative traits

(Mackay et al., 2009). Thus, gene expression has also been widely employed to search for

putative adaptive traits of locally adapted populations (Romero et al., 2012; Signor and

Nuzhdin, 2018; Sork, 2017) or ancestral and evolved populations in the context of

experimental evolution (Ferea et al., 1999; Huang and Agrawal, 2016; Lenski et al., 1994;

Mallard et al., 2018).

In this study, we used forward simulations that match not only essential design features of

typical experimental evolution studies, but also incorporate realistic parameters of the genetic

architecture. We recapitulate the classic expectations that even a moderately polygenic

architecture is associated with a high stability of the phenotypic variance of a selected trait

across different phases of adaptation. Applying this insight to a real dataset (Barghi et al.,

2019; Hsu et al., 2020; Jakšić et al., 2020), we show that a considerable set of genes changed

their mean expression, but their expression variance was indistinguishable from genes

without changes in mean expression. We propose that this pattern reflects that adaptive gene

expression evolution generally has a polygenic basis.

Results

The central idea of this study is that the complexity of an adaptive trait can be estimated from

the trajectory of the phenotypic variance during adaptation: the phenotypic variance remains

relative stable for a trait with polygenic (infinitesimal) architecture while it changes across

generations for a trait with oligogenic architecture. Although this prediction has been

illustrated in multiple theoretical and simulation studies (e.g.: Barton et al., 2017; Franssen et

al., 2017; Jain and Stephan, 2015), as the first step of this study, we explored to what extent it

can be generalized to a typical E&R setting considering a broader parameter space. Assuming

additivity and a negative correlation between ancestral allele frequency and effect size (Otte

et al., 2020) (Figure 1b), we simulated traits adapting to a mild/distant shift in trait optimum

with weak/intermediate/strong stabilizing selection (Figure 1a and Figure 1 – Figure

supplement 1). With three different distributions of effect size (Figure 1c) we investigate how

the number of contributing loci affects the phenotypic variance.

We monitored the change in phenotypic variance over 200 generations, which was sufficient

to reach the trait optimum for most parameter combinations (Figure 2 – Figure supplement 4

and 5). We compared the change in variance relative to the start of the experiment in

populations with and without selection. First, we studied a mild (one standard deviation of

the ancestral phenotypic distribution) shift in trait optimum (Figure 2 and Figure 2 – Figure

supplement 2). As expected for a founder population derived from a substantially larger

natural population, we find that even under neutrality the phenotypic variance does not

remain constant, but gradually decreases during 200 generations of experimental evolution

(Figure 2). This loss of variance is best explained by the fixation of variants segregating in

the founder population and we did not simulate new mutations, as they do not contribute to

adaptation in such short time scales (Burke et al., 2010). Our simulations show that even

experimental evolution studies with moderate population sizes and linkage very nicely

recapitulate the patterns from previous studies (Barton et al., 2017; Franssen et al., 2017; Jain

and Stephan , 2015). A pronounced drop in phenotypic variance is observed while a trait is

approaching a shifted optimum with few contributing loci (Figure 2). When more loci are

contributing to the selected phenotype, the difference to neutrality becomes very small

(Figure 2 and Figure 2 - Figure supplement 2). In addition to the number of contributing loci,

also the heterogeneity in effect size among loci and the shape of the fitness function have a

major impact. The larger the difference in effect size is, the more pronounced was the

influence of the number of contributing loci (Figure 2). The opposite effect was seen for the

width of the fitness function – a larger variance decreased the influence of the number of

contributing loci (Figure 2). Importantly, these patterns were not affected by the duration of

the experiment-qualitatively identical patterns were seen at different time points until

generation 200.

For a more distant trait optimum (three standard deviations of the ancestral phenotypic

distribution), we noticed some interesting dynamics that were not apparent for a closer trait

optimum (Figure 2 – Figure supplement 1 and 3). The most striking one was the temporal

heterogeneity of the phenotypic variance for few contributing loci of unequal effects. During

the early stage of adaptation, the variance increased and dropped later below the variance in

the founder population. With an increasing number of contributing loci, this pattern

disappeared and closely matched the neutral case (Figure 2 – Figure supplement 1).

Modifying dominance did not change the overall patterns-with a large number of contributing

loci the variance fitted the neutral pattern best (Figure 2 – Figure supplement 6). Overall, our

simulations indicate that only for a small set of parameters, the variance will increase during

the early stage of adaptation - in particular scenarios based on a few contributing loci with

very different effect sizes. The large influence of key parameters of the adaptive architecture,

in particular the number of contributing loci and their effect sizes on the temporal phenotypic

variance dynamics, suggest that it should be possible to exploit this for a test of polygenic

adaptation, which is independent from genomic data.

As a proof of principle, we studied the evolution of gene expression variance in replicated

populations evolving in a new hot temperature regime (Barghi et al., 2019; Hsu et al., 2020;

Jakšić et al., 2020). The evolved populations were derived from the same ancestral

population, but evolved independently for 100 generations in a novel temperature regime

with daily temperature fluctuations between 18 and 28°C (Figure 3a). Rather than relying on

pooled samples that allow only to estimate means, we quantified gene expression of

individuals from reconstituted ancestral populations and two evolved populations in a

common garden setup. Similar to previous studies (Ayroles et al., 2009; Huang et al., 2015),

we estimated a heritability of around 60% across the transcriptome among individual flies

from different families, which demonstrates the robustness of our experimental setup to

estimate expression variances (Method – Figure supplement 1; See Materials and methods).

Principle Component Analysis (PCA) indicated that 11.9% of the variation in gene

expression can be explained by the first PC which separates evolved and ancestral

populations, reflecting clear adaptive gene expression changes in response to the novel, hot

temperature regime (Figure 3b). The means and variances of the expression of each gene

were estimated and compared between the reconstituted ancestral populations and the two

evolved populations (Method – Figure supplement 2 and 3; See Materials and methods). Due

to the usage of different lot numbers for the RNA-Seq library preparation (Supplementary file

1), we only contrasted ancestral and evolved samples generated with the same lot number

(See Materials and methods) to avoid any unnecessary confounding results.

The comparison of ancestral and evolved populations identified 2,812 genes in the first

replicate and 2,704 genes in the second replicate which significantly changed mean

expression in the evolved flies (FDR<0.05, Supplementary file 2). With about 20% of the

genes changing mean expression, it is apparent that both populations evolved during 100

generations of exposure to a novel environment. 93.8% of the genes with a significant mean

expression change in both populations changed in the same direction, more than expected by

chance (Figure 4a, $\chi^2 = 896.34$, p-value $< 2.2 \times 10^{-16}$). This concordance suggests that

most of the altered expression means are mainly driven by selection, rather than by drift. We

quantified the expression change by relating the change in gene expression to the standard

deviation in the ancestral population. The differentially expressed genes in both replicates

showed a broad distribution of expression change, but the mean expression changed by one

standard deviation (Figure 4b). Assuming that all expression phenotypes reached trait

optimum, this reflects on average a moderate shift in trait optimum.

Consistent with computer simulations, the analysis of gene expression variance showed a

slight decrease in evolved populations relative to the ancestral ones (median F-value = 0.84 in

both replicates). Only a small subset of gene (n=125 and 97 in each replicate) experienced a

substantial reduction in variance. Because this reduction in variance is probably driven by a

different evolutionary force, we discuss them elsewhere (Lai et al., 2021). For the remaining

genes, we related the changes in gene expression variance of selected genes to non-selected

ones, we tested whether the variance changes in expression differ between the genes with

significant mean expression changes and those without. Hence, we assume that genes with

significant mean expression changes are under selection and the rest of the transcriptome has

no large effect on the fitness (neutral). Remarkably, the changes in variance of putative

adaptive genes with significant mean expression changes are indistinguishable from the genes

that do not change their mean expression (Figure 4c). This suggests that the selection on

mean expression is independent from the change in variance during adaptation in the focal

populations. While variance estimates from two time points do not provide sufficient power

to estimate the number of contributing loci in absence of more information about the adaptive

architecture, our computer simulations (Figure 2), suggest in line with previous theoretical

work (Barton et al., 2017; Jain and Stephan, 2015) that the observed stability in variance

evolution reflects a polygenic architecture underlying the adaptive gene expression evolution.

Since we only explored two time points rather than a full time series, it may be possible that

an oligogenic basis could also result in a similar phenotypic variance change as a polygenic

architecture (Figure 2 – Figure supplement 1 and 3). This can be seen in an intuitive case

when a single/few major effect allele(s) starts at a low frequency and becomes fixed (Yoo et

al., 1980). Because an oligogenic basis results in a highly parallel genomic selection response

(Figure 2 – Figure supplement 7), it is possible to distinguish polygenic and oligogenic

architectures with phenotypes from two time points only, when genomic data are available.

Because the genomic signature in the same experiment uncovered a highly heterogeneous

selection response (Barghi et al., 2019), we can exclude the unlikely explanation of an

oligogenic architecture resulting in a similar expression variance as non-selected genes.

Rather, we conclude that the adaptive response in gene expression is best explained by a

highly polygenic architecture.

Discussion

Population genetics has a long tradition to characterize adaptation based on the genomic signature of selected loci (Nielsen, 2005). Nevertheless, for selected phenotypes with a polygenic architecture, the contribution of individual loci to phenotypic change may be too subtle to be detected with classic population genetic methods (Pritchard et al., 2010). Even with an oligogenic basis, the identification of the selection targets with classic population genetic tests can be challenging.

Here, we used a conceptually different approach, which does not build on the genomic signature, to infer the adaptive architecture. Reasoning that experimental evolution is probably the best approach to obtain phenotypic time series, we performed computer simulations specifically tailored to typical experimental evolution studies with *Drosophila*. We showed that the temporal dynamics of the phenotypic variance is strongly affected by the number of contributing loci and other parameters of the adaptive architecture, such as the distribution of effect size and the underlying fitness function. Similar to the classic Castle-Wright estimator (Castle, 1921) that estimates the number of loci contributing to a quantitative trait from the phenotypic variance of the F2, we propose that the temporal heterogeneity of the phenotypic variance can be used to estimate the number of loci contributing to the adaptive response of a phenotype as well as other parameters of the adaptive architecture. Hence, unlike other approaches to characterize polygenic adaptation, the proposed estimator does not require genetic data when phenotypic time series data are available.

Because gene expression changes are constituting a major component of adaptation to a novel environment (Romero et al., 2012; Signor and Nuzhdin, 2018; Sork, 2017), it provides an

excellent model to evaluate a variance-based test for polygenicity. Gene expression is

modified by many trans-acting factors and some cis-regulatory variation. Adaptive gene

expression changes can be either driven by polymorphism in *cis*-regulatory elements or by

trans-acting variants. While interspecific differences in gene expression are predominantly

caused by *cis*-regulation (Wittkopp et al., 2004), intraspecific variation is mostly driven by

trans regulatory changes (Suvorov et al., 2013; Wittkopp et al., 2008). Adaptive gene

expression changes which are well-characterized on the molecular level typically have a

cis-regulatory basis that is not only frequently associated with the insertion of a transposable

element (e.g.: (Daborn et al., 2002)) but also sometimes with multiple regulatory variants

(Endler et al., 2018). Two lines of evidence suggest that cis-regulatory variation cannot be the

driver of adaptive gene expression changes observed in this study. First, the mutational target

size is too small to harbor a sufficiently large number of alleles segregating in the founder

population. Second, too few recombination events occur during the experiment to uncouple

regulatory variants located on a given haplotype such that they could generate a signal of

polygenic adaptation. More likely, the polygenic adaptive architecture of gene expression

change reflects the joint effects of many *trans*-acting variants.

Because we could only analyze phenotypic data from two time points, the founder population

and replicate populations evolved for 100 generations, we were not able to obtain a more

quantitative estimate of the number of contributing loci, in particular as other parameters of

the adaptive architecture are not known and need to be co-estimated. For the distinction

between an oligogenic and polygenic basis, we additionally relied on the heterogeneity of

genomic selection signatures among replicates, because for some parameter combinations the

oligogenic response can also result in a similar phenotypic variance as a polygenic one, but

with a much higher parallel response of genomic markers (Figure 2 – Figure supplement 7).

Hence, not only more time points describing the phenotypic trajectory, but also some

genomic data could contribute to infer the adaptive architecture in experimental evolution

studies. The extension of this approach to natural populations faces several challenges. First,

phenotypic time series over evolutionary relevant time scales are rare (but see (Clutton-Brock

and Pemberton, 2004)) and second, the distinction of environmental heterogeneity from

genetic changes is considerably more challenging than under controlled laboratory

conditions.

Figure 1. Simulating polygenic adaptation to a shift in trait optimum with different parameter combinations. a. For the computer simulations we consider a quantitative trait (in black) experiencing a sudden shift in trait optimum under stabilizing selection. The underlying fitness functions are illustrated in red. The new trait optimum is shifted from the ancestral trait mean by one/three standard deviation of the ancestral trait distribution. The strength of stabilizing selection is modified by changing the variance of the fitness function: 1.8, 3.6 and 5.4 standard deviations of the ancestral trait distribution. **b.** The negative correlation between the allele frequencies and the effect sizes (r = -0.7, estimated in Barghi *et*

281   *al*., 2019). We consider such negative correlation when assigning the effect sizes to variants

282   underlying a simulated trait. **c.** The distribution of effect sizes of the contributing loci is

283   determined by the shape parameter (α) of gamma sampling process ($\alpha = 0.5$, 2.5 and 100).

Figure 2. The trajectory of changes in phenotypic variance during adaptation to a mild optimum shift. The changes in phenotypic variance within 200 generations adapting to a moderate optimum shift (orange) are compared to the changes under neutrality (grey) on y axis. The change in variance (F) is calculated as the ratio of phenotypic variance between each evolved time point (generation x) and the ancestral state (σ_x^2/σ_0^2). The simulations cover traits controlled by varying numbers of loci underlying the adaptation with three different distributions of effect sizes (columns) under different strength of stabilizing selection (rows). For each scenario, 1000 runs of simulations have been performed. Only traits with the most

294   (dotted lines, 1000 loci), intermediate (dash lines, 50 loci) and the least (solid lines, 5 loci)

295   polygenic architectures are shown. In all scenarios, the variance of the trait decreases

296   drastically when the adaptation is controlled by a small number of loci (orange solid lines; 5

297   loci). While, for traits with extremely polygenic basis, the phenotypic variance stays stable

298   over time (orange dotted lines).

Figure 3. Schematic overview of the experimental procedures (a) and the divergence in gene expression during experimental evolution (b). **a.** Experimental evolution: starting from one common founder population, two replicate populations evolved for 100 generations in a hot laboratory environment fluctuating between 18 and 28°C. Common garden Experiment: after 100 generations, the two evolved replicate populations were maintained together with the reconstituted ancestral population for two generations in a hot laboratory environment fluctuating between 18 and 28°C. After this common garden procedure, about 30 males from each population were subjected to RNA-Seq. **b.** Principle Component Analysis (PCA) of the transcriptomic profiles of individuals from the ancestral population (blue) and the hot-evolved population (red). Circles indicate individuals of the first replicate (Anc. No. 27 and Evo. No. 4). Triangles represent individuals of the second replicate (Anc. No. 28 and Evo. No. 9). The two replicates were made with two different batches of library cards for RNA-Seq library preparation.

Figure 4. Evolution of phenotypic mean and variance over 100 generations of adaptation in empirical data. a. The evolution of gene expression mean during adaptation in the two replicates. For the genes with significant changes (DE, in orange), the changes are correlated between replicates (Spearman's rho = 0.53). For the genes without significant changes (non-DE, in grey), the correlation between replicates is much lower (Spearman's rho = 0.2). **b.** The evolution of gene expression mean scaled by the ancestral variation. For the DE genes (in orange), the median change falls around 1 standard deviation of the ancestral expression value, suggesting mild shift in trait optimum in the novel environment. For the

non-DE genes (in grey), the changes in expression are mostly negligible. The same pattern is seen in the second replicate. **c.** The change in expression variance during adaptation for DE and non-DE genes. In both replicates, the distribution of variance changes is indistinguishable between DE genes (orange) and non-DE genes (grey) (Wilcoxon's rank sum test, $p > 0.1$ for both replicates).

Materials and Methods

<u>Computer simulations</u>

We performed forward simulations with MimicrEE2 (Vlachos and Kofler, 2018) using the qff mode to illustrate the influence of the genetic architecture on the evolution of phenotypic variance during the adaptation to a new trait optimum. With 189 founder haplotypes (Barghi et al., 2019), we simulated quantitative traits under the control of different numbers of loci (5, 25, 50, 100, 200 and 1000) with an effective population size of 300. For each trait, we assume an additive model and a negative correlation (r = -0.7, estimated in Barghi et al., 2019) between the effect size and starting frequency (Figure 1b). The effect sizes of each locus can disperse in different levels which depend on the shape parameter of gamma sampling process (shape = 0.5, 2.5 and 100, Figure 1c). We used correlate() function implemented in "fabricatr" R package (Blair et al., 2019) to generate the effect sizes. The sum of effect sizes of each trait was normalized to 1. We assumed a heritability $h^2 = 0.6$ (from a family-based estimation in this study). To simulate stabilizing selection with trait optimum shift, we provided the Guassian fitness functions with mean of $\overline{X_{anc.}} + a\sqrt{V_{anc.}}$ and standard deviation of $b\sqrt{V_{anc.}}$, where $\overline{X_{anc.}}$ is the ancestral phenotypic mean and $V_{anc.}$ is the ancestral genetic variance (Figure 1a). Parameter "a" determines the distance of optimum shift, which is set to one (similar to the empirical case, Figure 4b) or three (Adopted from Sella et al., 2019). Parameter "b" indicates how strong the phenotypic constrain would be when the trait optimum is reached. The value 3.6 is adopted from Sella et al., 2019. In this study, we increase and decrease it by 50% to explore its impact (1.8 or 5.4). For the neutrality case, we assumed uniform fitness for all individuals. For each trait under each scenario, the phenotypic variance was estimated at different generations and compared to the ancestral phenotypic variance at generation 1 to illustrate the dynamic of phenotypic variance during the evolution. We note that we do not assume that the ancestral population has reached an

equilibrium, because the ancestral population in this study was phenotyped in the new

environment.

<u>Experimental evolution</u>

The setup of populations and evolution experiment have been described by (Barghi et al.,

2019). Briefly, ten outbred populations seeded from 202 isofemale lines were exposed to a

laboratory experiment at 28/18 °C with 12hr light/12hr dark photoperiod for more than 100

generations. Each replicate consisted of 1000 to 1250 adults at each generation.

<u>Common garden experiment</u>

The collection of samples from the evolution experiment for RNA-Seq was preceded by two

generations of common garden (CGE). The common garden experiment was performed at

generation 103 of the evolution in the hot environment and this CGE has been described in

(Barghi et al., 2019; Hsu et al., 2020, 2019; Jakšić et al., 2020). In brief, an ancestral

population was reconstituted by pooling five mated females from 184 founder isofemale lines

(Nouhaud et al., 2016). No significant allele frequency differences are expected between the

reconstituted ancestral populations and the original ancestral populations initiating the

experiment (Nouhaud et al., 2016). Because we evaluated phenotypes on the population

level, deleterious mutations will have a very limited impact. The reason is that they occur

only in a single isofemale line, which represents a very small fraction of the total population.

Two replicates of the reconstituted ancestral population and two independently evolved

populations at generation 103 were reared for two generations with egg-density control (400

eggs/bottle) at the same temperature regime as in the evolution experiment. Freshly eclosed

flies were transferred onto new food for mating. Sexes were separated under CO_2 anesthesia

at day 3 after eclosure, left to recover from CO_2 for two days, and at the age of five days

whole-body mated flies of each sex were snap-frozen at 2pm in liquid nitrogen and stored at

-80°C until RNA extraction. In this study, more than 30 individual male flies from two

reconstituted ancestral populations (replicate no. 27 and no. 28) and two evolved populations

(replicate no. 4 and no. 9) were subjected to RNA-Seq.

RNA extraction and library preparation

Whole bodies of individual male flies were removed from the -80°C freezer and immediately

homogenized in Qiazol lysis reagent (Qiagen, Hilden, Germany). The homogenate was

treated with DNase I followed by addition of chloroform, centrifugation and mixture of the

upper phase with 70% ethanol as described for the Qiagen RNeasy Universal Plus Mini Kit.

The mixture was subsequently loaded onto a RNeasy MinElute Spin column as provided by

the RNeasy Plus Micro Kit (Qiagen, Hilden, Germany), and all washing steps were

performed according to the instructions for that kit. All resulting total RNA was used to

prepare stranded mRNA libraries on the Neoprep Library Prep System (Illumina, San Diego,

USA) following the manufacturer's protocol: Neoprep runs were performed using software

version 1.1.0.8 and protocol version 1.1.7.6 with default settings for 15 PCR cycles and an

insert size of 200bp. All libraries for individuals of ancestral replicate no. 27 and evolved

replicate no. 4 were prepared with library cards of lot no. 20180170; all libraries for

individuals of ancestral replicate no. 28 and evolved replicate no. 9 were prepared with

library cards of lot no. 20178099. 50bp single-end reads were sequenced on an Illumina

HiSeq 2500. All sequencing data will be available in European Nucleotide Archive (ENA)

under the accession number PRJEB37011 upon publication.

RNA-Seq data processing and quality control

All RNA-Seq reads were trimmed using ReadTools (Gómez-Sánchez and Schlötterer, 2018)

with quality score of 20 and aligned to *Drosophila simulans* reference genome (Palmieri et

al., 2015) using GSNAP (Wu et al., 2016) with parameter setting -k 15 -N 1 -m 0.08.

Exon-aligned reads were piped into Rsubread (Liao et al., 2019) to calculate read counts of

each gene, and raw read counts of each gene were normalized with the TMM method

implemented in edgeR (Robinson et al., 2010). Samples with severe 3'- bias were removed

based on visual inspection of the gene-body coverage plot (Jakšić and Schlötterer, 2016;

Wang et al., 2012).

<u>Genetic variance in gene expression across F1 families</u>

We evaluated how much of the expression variance can be explained by genetic variation by

performing RNA-Seq on individual flies, with 3-4 individuals each from three isofemale lines

maintained at the same density and culturing conditions.

Six out of 184 founder isofemale lines from the evolution experiment and were maintained

for one generation with controlled egg density (400 eggs/bottle) in the same environment as

the main experiment (12h 28°C with light followed by 12h 18 °C with dark conditions).

Using the offspring, we generated three crosses between two of the six lines each: FL 138 x

FL 137, FL 157 x FL 112, FL 123 x FL 127: we combined 50 virgin females from one of the

lines with 50 males from the other line, let them lay eggs under density control as above and

maintained and froze their F1 offspring in the same way as in the main CGE: sexes were

separated after mating at the age of three days and snap-frozen at the age of five days at 2pm.

From each cross, we used four F1 males to prepare individual RNA-Seq libraries as described

above.

Assuming no environmental heterogeneity, we decomposed the total variance of the

expression of each gene measured in these individuals into the genetic difference among

three different F1 families and random error. The data were analyzed as follows:

Natural log-transformation was applied to CPMs of all genes to improve data normality

(Rocke and Durbin, 2003). Principal component and principal variance component analysis

(Bushel, 2019) was performed to the whole transcriptome to decompose the variance

components. We found that around 60% of the gene expression variance can be explained by

the genetic difference among the three F1 families (method – Figure supplement 1a and b).

This implies that the within-population gene expression variance is largely contributed by the

genetic components. Because we only used offspring from single vials, we may have

overestimated the heritability if the environment in the vials differs. Nevertheless, since our

heritability estimates are very similar to previous ones (Ayroles et al., 2009), we consider our

estimates reliable.

In addition to general analysis across all genes, we also tested for the genetic variance of each

gene separately using analysis of variance (ANOVA):

$$y_{ij} = \mu + \tau_i + \varepsilon_{ij},$$

Where i=1, 2, 3, …, n (the i^{th} genes); j =1, 2, 3, 4 (the j^{th} individuals in each cross). y_{ij} is the

observed expression level of a gene in a given sample, μ is the overall mean; τ_i is the

effect of genetic background and ε_{ij} is the random noise. We calculated the proportion of

total variance explained by random error using the following equation:

$$variance\ explained\ by\ random\ error = \frac{sum\ squares\ of\ error\ (SSE)}{sum\ squares\ of\ total\ (SST)}$$

Genes were binned based on their average expression value (lnCPM) which ranged from -0.8

to 4.1, by bin size of 0.1. The average proportion of variance explained by random error of

each bin was calculated.

The expression variance of genes with less than 1 count per million bases (CPM) is

dominated by technical errors (method – Figure supplement 1c). Thus, genes with less than

1 count per million base (CPM) were excluded for subsequent analysis.

<u>RNA-Seq data analysis</u>

We observed some outlier individuals and suspected that the freezing process may have led

to detachment of body parts, such as eyes or heads, in these individuals. We compared gene

expression between such outliers and all other samples and performed tissue enrichment

analysis for genes with at least 2-fold lower expression in the outlier samples. Samples with

evidence of tissue detachment were excluded. After filtering, each population remained

approximately 20 individuals (Supplementary file 1). Only genes with at least 1 count per

million base (CPM) were included in the analyses to avoid extremely lowly expressed genes.

For all RNA-Seq data we only compared samples which were prepared with library cards

from the same lot number to avoid batch effects (Replicate 1: evolved replicate 4 vs.

reconstituted ancestral population replicate 27; Replicate 2: evolved replicate 9 ss.

reconstituted ancestral population replicate 28).

For differential expression analysis on mean expression, we used the generalized linear

modeling function implemented in edgeR (Robinson et al., 2010) to fit the expression to the

model ($Y = E + \varepsilon$) in which Y stands for gene expression, E is the effect of evolution and ε

is the random error. The likelihood ratio test was performed to test the effect of evolution.

P-value adjustment was performed using the Benjamini-Hochberg false discovery rate (FDR)

correction.

For the analysis of expression variance evolution, we applied natural log transformation

(Rocke and Durbin, 2003) to eliminate the strong mean-variance dependency in RNA-Seq

data due to the nature of the negative binomial distribution (method – Figure supplement 2).

The variance of the expression of each gene (lnCPM) was estimated in each population. With

the moderate sample size, we needed to estimate the uncertainty of variance estimates.

Jackknifing was applied to measure the uncertainty of estimator (Fukunaga and Hummels,

1989). The procedure was conducted independently on four replicates and we calculated the

95% confidence interval of the estimated variance (method – Figure supplement 3). The

change of gene expression variance was determined by the F statistics calculated as the ratio

between the variance within the ancestral population and the variance within the evolved

population of each gene. To test whether selection alter the expression variance, a

comparison was made between the F statistics of genes with significant changes in mean

expression and the ones without.

Acknowledgments

Special thanks to David Houle, who provided fantastic support during the collection and

establishment of the isofemale lines in Florida. We thank all member of the Institut für

Populationsgenetik for discussion. We are grateful to Reinhard Bürger and David Houle for

helpful comments on earlier versions of the manuscript. Neda Barghi, François Mallard and

Kathrin Otte contributed to the common garden experiment. Illumina sequencing was

performed at the VBCF NGS Unit (www.vbcf.ac.at). This work was support by the Austrian

Science Funds (FWF, W1225) and the European Research Council (ERC, ArchAdapt).

Author contribution

W.Y.L and C.S. conceived the study. V.N. prepared all RNA-Seq and supervised the

maintenance of the evolution experiment. A.M.J supervised the common garden experiment.

W.Y.L performed the simulation and data analysis. W.Y.L. and C.S. wrote the manuscript.

Competing interests

The authors declare no competing interests.

Correspondence and requests for materials should be addressed to C.S.

References

Ayroles JF, Carbone MA, Stone EA, Jordan KW, Lyman RF, Magwire MM, Rollmann SM,

Duncan LH, Lawrence F, Anholt RRH, Mackay TFC. 2009. Systems genetics of

complex traits in *Drosophila melanogaster*. *Nat Genet* **41**:299–307. doi:10.1038/ng.332

Barghi N, Hermisson J, Schlötterer C. 2020. Polygenic adaptation: a unifying framework to

understand positive selection. *Nat Rev Genet* **21**:769–781.

doi:10.1038/s41576-020-0250-z

Barghi N, Tobler R, Nolte V, Jakšić AM, Mallard F, Otte KA, Dolezal M, Taus T, Kofler R,

Schlötterer C. 2019. Genetic redundancy fuels polygenic adaptation in *Drosophila*.

PLOS Biol **17**:e3000128. doi:10.1371/journal.pbio.3000128

Barton NH, Etheridge AM, Véber A. 2017. The infinitesimal model: Definition, derivation,

and implications. *Theor Popul Biol* **118**:50–73. doi:10.1016/J.TPB.2017.06.001

Barton NH, Keightley PD. 2002. Understanding quantitative genetic variation. *Nat Rev Genet*

3:11–21. doi:10.1038/nrg700

Barton NH, Turelli M. 1987. Adaptive landscapes, genetic distance and the evolution of

quantitative characters. *Genet Res* **49**:157–173. doi:10.1017/S0016672300026951

Boyle EA, Li YI, Pritchard JK. 2017. An expanded view of complex traits: From polygenic

to omnigenic. *Cell* **169**:1177–1186. doi:10.1016/J.CELL.2017.05.038

Bulmer MG. 1972. The genetic variability of polygenic characters under optimizing

selection, mutation and drift. *Genet Res* **19**:17–25. doi:10.1017/S0016672300014221

Burke MK, Dunham JP, Shahrestani P, Thornton KR, Rose MR, Long AD. 2010.

Genome-wide analysis of a long-term evolution experiment with *Drosophila*. *Nature*

467:587–590. doi:10.1038/nature09352

Bushel P. 2019. pvca: Principal variance component analysis (PVCA). *R Packag version*

1240.

Castle WE. 1921. An improved method of estimating the number of genetic factors

concerned in the cases of blending inheritance. *Science* **54**:223.

doi:10.1126/science.54.1393.223

Chevalet C. 1994. An approximate theory of selection assuming a finite number of

quantitative trait loci. *Genet Sel Evol.* **26**:379. doi:10.1186/1297-9686-26-5-379

Chippindale AK, Chu TJF, Rose MR. 1996. Complex trade-offs and the evolution of

starvation resistance in *Drosophila melanogaster.* *Evolution (N Y)* **50**:753–766.

doi:10.1111/j.1558-5646.1996.tb03885.x

Clutton-Brock TH, Pemberton JM (Josephine M). 2004. Soay sheep : population dynamics

and selection on St. Kilda. Cambridge University Press.

Daborn PJ, Yen JL, Bogwitz MR, Le Goff G, Feil E, Jeffers S, Tijet N, Perry T, Heckel D,

Batterham P, Feyereisen R, Wilson TG, ffrench-Constant RH. 2002. A single p450

allele associated with insecticide resistance in *Drosophila.* *Science* **297**:2253–6.

doi:10.1126/science.1074170

Endler L, Gibert J-M, Nolte V, Schlötterer C. 2018. Pleiotropic effects of regulatory variation

in *tan* result in correlation of two pigmentation traits in *Drosophila melanogaster. Mol*

Ecol **27**:3207–3218. doi:10.1111/mec.14781

Falconer DS, Mackay TFC. 1963. Introduction to quantitative genetics, Poultry Science.

CreateSpace Independent Publishing Platform. doi:10.3382/ps.0420547

Ferea TL, Botstein D, Brown PO, Rosenzweig RF. 1999. Systematic changes in gene

expression patterns following adaptive evolution in yeast. *Proc Natl Acad Sci U S A*

96:9721–6. doi:10.1073/pnas.96.17.9721

Franssen SU, Kofler R, Schlötterer C. 2017. Uncovering the genetic signature of quantitative

trait evolution with replicated time series data. *Heredity (Edinb)* **118**:42–51.

doi:10.1038/hdy.2016.98

Fukunaga K, Hummels DM. 1989. Leave-one-out procedures for nonparametric error

estimates. *IEEE Trans Pattern Anal Mach Intell* **11**:421–423. doi:10.1109/34.19039

Gómez-Sánchez D, Schlötterer C. 2018. *ReadTools* : A universal toolkit for handling

sequence data from different sequencing platforms. *Mol Ecol Resour* **18**:676–680.

doi:10.1111/1755-0998.12741

Hayward LK, Sella G. 2019. Polygenic adaptation after a sudden change in environment.

bioRxiv 792952. doi:10.1101/792952

Hsu S-K, Jakšić AM, Nolte V, Barghi N, Mallard F, Otte KA, Schlötterer C, Hsu S-K, Jakšić

AM, Nolte V, Barghi N, Mallard F, Otte KA, Schlötterer C. 2019. A 24 h age difference

causes twice as much gene expression divergence as 100 generations of adaptation to a

novel environment. *Genes (Basel)* **10**:89. doi:10.3390/genes10020089

Hsu S-K, Jakšić AM, Nolte V, Lirakis M, Kofler R, Barghi N, Versace E, Schlötterer C.

2020. Rapid sex-specific adaptation to high temperature in *Drosophila*. *Elife* **9**.

doi:10.7554/eLife.53237

Huang W, Carbone MA, Magwire MM, Peiffer JA, Lyman RF, Stone EA, Anholt RRH,

Mackay TFC. 2015. Genetic basis of transcriptome diversity in *Drosophila*

melanogaster. *Proc Natl Acad Sci U S A* **112**:E6010–E6019.

doi:10.1073/pnas.1519159112

Huang Y, Agrawal AF. 2016. Experimental evolution of gene expression and plasticity in

alternative selective regimes. *PLOS Genet* **12**:e1006336.

doi:10.1371/journal.pgen.1006336

Jain K, Stephan W. 2015. Response of polygenic traits under stabilizing selection and

mutation when loci have unequal effects. *G3 (Bethesda)* **5**:1065–74.

doi:10.1534/g3.115.017970

Jakšić AM, Karner J, Nolte V, Hsu S-K, Barghi N, Mallard F, Otte KA, Svečnjak L, Senti

K-A, Schlötterer C. 2020. Neuronal function and dopamine signaling evolve at high

temperature in *Drosophila*. *Mol Biol Evol* **37**:2630–2640. doi:10.1093/molbev/msaa116

Jakšić AM, Schlötterer C. 2016. The interplay of temperature and genotype on patterns of

alternative splicing in *Drosophila melanogaster*. *Genetics* **204**:315–325.

doi:10.1534/GENETICS.116.192310

Kawecki TJ, Lenski RE, Ebert D, Hollis B, Olivieri I, Whitlock MC. 2012. Experimental

evolution. *Trends Ecol Evol* **27**:547–560. doi:10.1016/J.TREE.2012.06.001

Keightley PD, Hill WG. 1989. Quantitative genetic variability maintained by

mutation-stabilizing selection balance: sampling variation and response to subsequent

directional selection. *Genet Res* **54**:45–58. doi:10.1017/S0016672300028366

Kimura M, Crow JF. 1964. The number of alleles that can be maintained in a finite

population. *Genetics* **49**:725–738.

Lai W.-Y. & Schlötterer, C. 2021. Evolution of gene expression variance during adaptation to

high temperature in *Drosophila*. *BioRxiv*.

Lande R. 1976. Natural selection and random genetic drift in phenotypic evoltion. *Evolution*

(N Y) **30**:314–334. doi:10.1111/j.1558-5646.1976.tb00911.x

Lenski RE, Travisano M, Larison B, Moritz C. 1994. Dynamics of adaptation and

diversification: a 10,000-generation experiment with bacterial populations. *Proc Natl*

Acad Sci **91**:6808–6814. doi:10.1073/pnas.91.15.6808

Liao Y, Smyth GK, Shi W. 2019. The R package Rsubread is easier, faster, cheaper and

better for alignment and quantification of RNA sequencing reads. *Nucleic Acids Res*

47:e47–e47. doi:10.1093/nar/gkz114

Liu X, Li YI, Pritchard JK. 2019. Trans effects on gene expression can drive omnigenic

inheritance. *Cell* **177**:1022-1034.e6. doi:10.1016/J.CELL.2019.04.014

Mackay TFC, Stone EA, Ayroles JF. 2009. The genetics of quantitative traits: challenges and

prospects. *Nat Rev Genet* **10**:565–577. doi:10.1038/nrg2612

Mallard F, Nolte V, Tobler R, Kapun M, Schlötterer C. 2018. A simple genetic basis of

adaptation to a novel thermal environment results in complex metabolic rewiring in

Drosophila. *Genome Biol* **19**:119. doi:10.1186/s13059-018-1503-4

Nielsen R. 2005. Molecular signatures of natural selection. *Annu Rev Genet* **39**:197–218.

doi:10.1146/annurev.genet.39.073003.112420

Nouhaud P, Tobler R, Nolte V, Schlötterer C. 2016. Ancestral population reconstitution from

isofemale lines as a tool for experimental evolution. *Ecol Evol* **6**:7169–7175.

doi:10.1002/ece3.2402

Otte KA, Nolte V, Mallard F, Schlötterer C. 2020. The adaptive architecture is shaped by

population ancestry and not by selection regime. *bioRxiv* 2020.06.25.170878.

doi:10.1101/2020.06.25.170878

Palmieri N, Nolte V, Chen J, Schlötterer C. 2015. Genome assembly and annotation of a

Drosophila simulans strain from Madagascar. *Mol Ecol Resour* **15**:372–81.

doi:10.1111/1755-0998.12297

Pélabon C, Hansen TF, Carter AJR, Houle D. 2010. Evolution of variation and variability

under fluctuating, stabilizing, and disruptive selection. *Evolution (N Y)* **64**:1912–25.

doi:10.1111/j.1558-5646.2010.00979.x

Pritchard JK, Pickrell JK, Coop G. 2010. The genetics of human adaptation: Hard sweeps,

soft   sweeps,   and   polygenic   adaptation.   *Curr   Biol*   **20**:R208–R215.

doi:10.1016/J.CUB.2009.11.055

Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. *Bioinformatics* **26**:139–

140. doi:10.1093/bioinformatics/btp616

Rocke DM, Durbin B. 2003. Approximate variance-stabilizing transformations for
gene-expression microarray data. *Bioinformatics* **19**:966–72.

Romero IG, Ruvinsky I, Gilad Y. 2012. Comparative studies of gene expression and the
evolution of gene regulation. *Nat Rev Genet* **13**:505–516. doi:10.1038/nrg3229

Signor SA, Nuzhdin S V. 2018. The evolution of gene expression in *cis* and *trans*. *Trends*
Genet **34**:532–544. doi:10.1016/j.tig.2018.03.007

Sork VL. 2017. Genomic studies of local adaptation in natural plant populations. *J Hered*
109:3–15. doi:10.1093/jhered/esx091

Suvorov A, Nolte V, Pandey RV, Franssen SU, Futschik A, Schlötterer C. 2013.
Intra-specific regulatory variation in *Drosophila pseudoobscura*. *PLoS One* **8**:e83547.
doi:10.1371/journal.pone.0083547

Turelli M. 1984. Heritable genetic variation via mutation-selection balance: Lerch's zeta
meets the abdominal bristle. *Theor Popul Biol* **25**:138–193.
doi:10.1016/0040-5809(84)90017-0

Wang L, Wang S, Li W. 2012. RSeQC: quality control of RNA-seq experiments.
Bioinformatics **28**:2184–2185. doi:10.1093/bioinformatics/bts356

Wittkopp PJ, Haerum BK, Clark AG. 2008. Regulatory changes underlying expression
differences within and between *Drosophila* species. *Nat Genet* **40**:346–350.
doi:10.1038/ng.77

Wittkopp PJ, Haerum BK, Clark AG. 2004. Evolutionary changes in *cis* and *trans* gene
regulation. *Nature* **430**:85–88. doi:10.1038/nature02698

Wu TD, Reeder J, Lawrence M, Becker G, Brauer MJ. 2016. GMAP and GSNAP for
genomic sequence alignment: Enhancements to speed, accuracy, and functionality.
Humana Press, New York, NY. pp. 283–334. doi:10.1007/978-1-4939-3578-9_15

Yoo BH, Nicholas FW, Rathie KA. 1980. Long-term selection for a quantitative character in

large replicate populations of *Drosophila melanogaster. Theor Appl Genet* **57**:113–117.

doi:10.1007/BF00253881

Figure supplements

Figure 1 – Figure supplement 1. The evolutionary scenario for distant optimum shift.
We consider the case when a quantitative trait (in black) experiences a sudden shift in trait
optimum under stabilizing selection. The imposed fitness functions (F.F.) are illustrated in
red. The new trait optimum is set away from the ancestral trait mean by three standard
deviation of the ancestral trait distribution for distal shift. To vary the strength of stabilizing
selection, the variance of the fitness function is set as 1.8, 3.6 and 5.4 standard deviation of
the ancestral trait distribution.

Figure 2 – Figure supplement 1. The trajectory of expected changes in phenotypic variance when adapting to a distant optimum shift. The changes in phenotypic variance within 200 generations adapting to a distant optimum shift (orange) are compared to the changes under neutrality (grey) on y axis. The change in variance (F) is calculated as the ratio of phenotypic variance between each evolved time point (generation x) and the ancestral state ($\sigma^2_{200}/\sigma^2_0$). The simulations cover traits controlled by varying numbers of loci underlying the adaptation with three different distributions of effect sizes (columns) and under different

strength of stabilizing selection (rows). For each scenario, 1000 runs of simulations have

been performed. Only traits with the most (dotted lines, 1000 loci), intermediate (dash lines,

50 loci) and the least (solid lines, 5 loci) polygenic architectures are shown. Unlike the

continuous decreasing pattern in the cases with moderate optimum shifts, the variance of

traits controlled by a few loci (5 loci) with largely dispersed effects would increase first and

then decrease when the effect sizes of contributing loci are dispersed (orange solid lines).

Nevertheless, for traits with extremely polygenic basis, the phenotypic variance always stays

stable over time (orange dotted lines).

Figure 2 – Figure supplement 2. The expected changes in phenotypic variance and the number of contributing loci. The changes in phenotypic variance after 200 generations adapting to a mild optimum shift (orange) are compared to the changes under neutrality (grey) on y axis. The change in variance (F) is calculated as the ratio between the evolved and ancestral phenotypic variance ($\sigma^2_{200}/\sigma^2_0$). The simulations cover traits controlled by varying numbers of loci underlying the adaptation (x-axes) with three different distributions of effect sizes (columns) and under different strength of stabilizing selection (rows). For each scenario, 1000 runs of simulations have been performed. In all scenarios, the variance of the trait

697   decreases drastically when the adaptation is controlled by a small number of loci. As the

698   number of contributing loci increases, the phenotypic variance becomes more stable.

Figure 2 – Figure supplement 3. The expected changes in phenotypic variance and the number of contributing loci. The changes in phenotypic variance after 200 generations adapting to a distant optimum shift (orange) are compared to the changes under neutrality (grey) on y axis. The change in variance (F) is calculated as the ratio between the evolved and ancestral phenotypic variance ($\sigma^2_{200}/\sigma^2_0$). The simulations cover traits controlled by varying numbers of loci underlying the adaptation (x-axes) with three different distributions of effect sizes (columns) and under different strength of stabilizing selection (rows). For each scenario, 1000 runs of simulations have been performed. In most scenarios, the variance of

709 the trait decreases drastically when the adaptation is controlled by a small number of loci. As

710 the number of contributing loci increases, the phenotypic variance becomes more stable.

711 However, exceptions can be observed when the effect sizes of the contributing loci are

712 largely dispersed. In these cases, shift in optimum does not have the strong impact on the

713 variance of traits under simple genetic control (5 contributing loci). In the extreme case,

714 highly dispersed effect sizes in combination with a relaxed phenotypic constraint removes the

715 relationship between the number of loci and the changes in phenotypic variance.

Figure 2 – Figure supplement 4. The changes in phenotypic mean when adapting to a

mild optimum shift. The changes in phenotypic mean after 200 generations adapting to a

mild optimum shift (orange) are compared to the changes under neutrality (grey) on y axis.

The changes in phenotypic mean are scaled by the standard deviation of the ancestral trait

distribution. The simulations cover traits controlled by varying numbers of loci underlying

the adaptation (x axes) with three different distributions of effect sizes (columns) and under

different strength of stabilizing selection (rows). For each scenario, 1000 runs of simulations

726   have been performed. In most cases, the traits under selection (orange) shift their means by

727   one standard deviation of the ancestral trait distribution (i.e. reaching the new trait optimum)

728   while the neutral traits (grey) stay unchanged.

Figure 2 – Figure supplement 5. The changes in phenotypic mean when adapting to a distant optimum shift. The changes in phenotypic mean after 200 generations adapting to a distant optimum shift (orange) are compared to the changes under neutrality (grey) on y axis. The changes in phenotypic mean are scaled by the standard deviation of the ancestral trait distribution. The simulations cover traits controlled by varying numbers of loci underlying the adaptation (x axes) with three different distributions of effect sizes (columns) and under different strength of stabilizing selection (rows). For each scenario, 1000 runs of simulations have been performed. In most cases, the traits under selection (orange) shift their means by

739   three standard deviation of the ancestral trait distribution (i.e. reaching the new trait optimum)

740   while the neutral traits (grey) stay unchanged.

Figure 2 – Figure supplement 6. The expected changes in phenotypic variance for traits controlled by dominant and recessive alleles. The changes in phenotypic variance after 200 generations adapting to a mild optimum shift (orange) are compared to the changes under neutrality (grey) on y axis. The change in variance (F) is calculated as the ratio between the evolved and ancestral phenotypic variance ($\sigma_{200}^2/\sigma_0^2$). This simulation covers traits controlled by varying numbers of loci underlying the adaptation (x axes) with recessive, additive and dominant effects. For each scenario, 1000 runs of simulations have been performed. No matter how the dominance varies, the variance of the trait decreases drastically when the adaptation is controlled by a small number of loci. As the number of contributing loci increases, the phenotypic variance becomes more stable.

Figure 2 – Figure supplement 7. Parallelism in genomic response across 10 evolved replicates for traits under the control of five loci and 100 loci. For the loci with allele frequency change of at least 10% in 200 generations, the average Pearson's correlation coefficient of the frequency change between all pairs of two evolved replicates was calculated to describe the parallelism of the evolution at these loci. An average across loci is used to obtain a general parallelism. 100 runs of simulations with 10 evolution replicates have been performed for each scenario. With five contributing loci, the genomic evolution of

762   the contributing loci is more parallel across the 10 evolved replicates compared to the case

763   when with 100 contributing loci.

(A)

(C)

Method – Figure supplement 1. Genetic variance in gene expression across F1 families.

(A) Principal component analysis (PCA) on the transcriptomes of F1 individuals from three

different crosses between the founder iso-female lines. Individuals from different families

clustered nicely based on the first two PCs. **(B)** Principal variance component analysis

(PVCA) on the transcriptomes of F1 individuals. 67% of the total variance in gene expression

was explained by the genetic difference between the individuals. **(C)** Gene-wise analysis of

variance (ANOVA) in gene expression. Genes were binned based on their average expression

value (lnCPM) which ranged from -0.8 to 4.1, by bin size of 0.1. The average proportion of

variance explained by random error of each bin was visualized. The expression variance of

genes with less than 1 count per million bases (CPM) is dominated by residuals.

Method – Figure supplement 2. Log-transformation eliminates the positive relationship between the changes in mean and variance of gene expression. In each panel, the changes in mean expression, log(FC) ($FC = \frac{\overline{y_{evo.}}}{\overline{y_{anc.}}}$) and in variance before **(A)** and after **(B)** the natural log-transformation of each gene were visualized ($F' = \frac{var(y_{evo.})}{var(y_{anc.})}$ and $F = \frac{var(\ln(y_{evo.}))}{var(\ln(y_{anc.}))}$). The positive correlation (r = 0.45) due to the positive mean-variance dependency of negative binomial distribution is removed by the log-transformation (r = -0.05) on gene expression level.

Method – Figure supplement 3. Robustness of the variance estimation using individual sequencing data. Jackknife method was applied to measure the uncertainty of variance estimation. Given a sample size of K, the procedure is to estimate the variance of each gene for K times, each time leaving one sample out. The procedure was conducted independently on 4 populations (anc.27, anc.28, evo.4 and evo.9). In each panel, we visualize Jackknife approximated 95% confidence interval for the variance estimates of each gene. The genes are ordered based on the average variance estimates (black dash line) on the x-axis. The upper and lower limits of the 95% confidence interval are indicated with grey curves. The salmon

line denotes the observed value of the variance estimates. In most cases, the estimates lie in

the confidence interval, suggesting robust estimation.

Titles and legend for supplementary files

Supplementary file 1. Library information of the sample in this study. This file provides

a list of all sequenced samples and the library information.

Supplementary file 2. Differential gene expression analysis of two contrasts between

ancestral and evolved populations. This file reports the results of DE analysis between anc.

27 and evo. 4 (Table S2**A**) and between anc. 28 and evo. 9 (Table S2**B**).

Supplementary file 3. F value on the gene expression of two contrasts between ancestral

and evolved populations. This file reports the results of gene expression variance

comparisons between anc. 27 and evo. 4 (Table S3**A**) and between anc. 28 and evo. 9 (Table

S3**B**).