Evolution of phenotypic variance provides insights into the genetic basis of adaptation

Author list:

Wei-Yun Lai[1,2], Viola Nolte[1], Ana Marija Jakšić[1,2,3] and Christian Schlötterer[1*]

[1]Institut für Populationsgenetik, Vetmeduni Vienna, Vienna, Austria.

[2]Vienna Graduate School of Population Genetics, Vetmeduni Vienna, Vienna, Austria.

[3]Current affiliation: École polytechnique fédérale de Lausanne, Lausanne, Switzerland

[*]Correspondence: christian.schloetterer@vetmeduni.ac.at; Tel.: +43-1-25077-4300.

Keywords:

Phenotypic variance, temperature adaptation, *Drosophila simulans*, experimental evolution

Abstract

Most traits are polygenic and the contributing loci can be identified by GWAS. Their adaptive

architecture is, however, difficult to characterize. Here, we propose to study the adaptive

architecture of traits by monitoring the evolution of their phenotypic variance during adaptation

to a new environment in well-defined laboratory conditions. Extensive computer simulations

show that the evolution of phenotypic variance in a replicated experimental evolution setting

provides power to distinguish between oligogenic and polygenic adaptive architectures. We

compared gene expression variance in male *Drosophila simulans* before and after 100

generations of adaptation to a novel hot environment. The variance change in gene expression

was indistinguishable for genes with and without a significant change in mean expression after

100 generations of evolution. We suggest that a majority of adaptive gene expression evolution

can be explained by a polygenic architecture. We propose that tracking the evolution of

phenotypic variance across generations can provide an approach to characterize the adaptive

architecture.

Introduction

It is widely accepted that most complex traits have a polygenic basis (Ayroles *et al.*, 2009; Boyle *et al.*, 2017; Liu *et al.*, 2019). Nevertheless, it is difficult to predict which of these loci are responding to selection when a population is exposed to a new selection regime. If pleiotropic constraints are strong, only a small subset of the genes constituting are free to respond to selection. Hence, the genetic basis of the adaptive response of a complex trait (i.e. adaptive architecture (Barghi *et al.*, 2020)) may differ substantially from the genetic architecture. Characterizing the adaptive architecture by mapping selected loci is not easy, in particular when more than a handful of genes are involved. To circumvent this problem, we introduce a complementary approach, analogous to the Castle-Wright estimator (Castle, 1921), to infer the complexity of the adaptive architecture (i.e. simple with few contributing loci or with a polygenic basis). We propose to study the evolution of phenotypic variance, which may provide some insights into the key parameters of the adaptive architecture.

The phenotypic variance of a quantitative trait is a key determinant for its response to selection. It can be decomposed into genetic and environmental components (Falconer & Mackay, 1963). Over the past years, mathematical models have been developed to describe the expected genetic variance of a quantitative trait under selection and its maintenance in evolving populations (Bulmer, 1972; Chevalet, 1994; Kimura and Crow, 1964; Turelli, 1984). For infinitely large populations and traits controlled by many independent loci with infinitesimal effect, changes in trait optimum are not expected to affect the phenotypic variance (Lande, 1976). A much more complex picture is expected when the effect sizes are not equal, the population size is finite, or the traits have a simpler genetic basis (Barton & Turelli, 1987; Keightley & Hill, 1989; Barton & Keightley, 2002; Jain & Stephan, 2015; Franssen *et al.*, 2017; Hayward & Sella, 2019). For instance, for traits with oligogenic architecture, the genetic variance could drop dramatically during adaptation, while with polygenic architecture, only minor effects on the variance are expected (Barton et al., 2017; Franssen et al., 2017; Jain and Stephan , 2015). These studies suggest that a time-resolved analysis of phenotypic variance has the potential to shed light onto the complexity of the underlying adaptive architecture.

Despite its potential importance for the understanding of adaptation, we are faced with the situation that few empirical data are available for the evolution of phenotypic variance. The use of natural populations to study changes in phenotypes, and even more so phenotypic

60 variances, is limited as the environmental heterogeneity cannot be controlled. A
61 complementary approach to study the evolution of phenotypic variance is experimental
62 evolution (Kawecki *et al.*, 2012). With replicated populations starting from the same founders
63 and evolving under tightly controlled environmental conditions, experimental evolution
64 provides the opportunity to study the evolution of phenotypic variance.
65

Most experimental evolution studies in sexual populations focused on the evolution of
phenotypic means, rather than variance (Chippindale *et al.*, 1996; Burke *et al.*, 2010; Mallard
et al., 2018; Jakšić *et al.*, 2020). Notable exceptions include a study which applied fluctuating,
stabilizing and disruptive selection to a small number of wing shape-related traits (Pélabon *et*
al., 2010) and other studies tracking the variance evolution of behavioral and cranial traits
under directional selection in mice (Careau *et al.*, 2015; Penna *et al.*, 2017). Instead of looking
at a preselected subset of phenotypes which limits the generality, we will focus on gene
expression, a set of molecular phenotypes, which can be easily quantified as microarrays and,
more recently, RNA-Seq have become available. Importantly, the expression levels of genes
exhibit the same properties (e.g.: continuality and normality) as other complex quantitative
traits (Mackay *et al.*, 2009). Thus, gene expression has also been widely employed to study the
adaptation of locally adapted populations (e.g.: Romero *et al.* 2012; Sork 2017; Signor and
Nuzhdin 2018) or ancestral and evolved populations in the context of experimental evolution
(e.g.: Lenski *et al.* 1994; Ferea *et al.* 1999; Huang and Agrawal 2016; Mallard *et al.* 2018).
80

In this study, we performed forward simulations that match not only essential design features
of typical experimental evolution studies, but also incorporate realistic parameters of the
genetic architecture. We recapitulate the classic expectations that even a moderately polygenic
architecture is associated with a high stability of the phenotypic variance of selected traits
across different phases of adaptation and evaluate the statistical power. Applying this insight
to a recently published dataset (Lai & Schlötterer, 2022), we show that for putatively selected
genes (DE genes), their average changes in expression variance were indistinguishable from
genes without changes in mean expression. We propose that this pattern reflects that adaptive
gene expression evolution generally has a polygenic basis.
90

Materials and Methods

<u>Computer simulations</u>

We performed forward simulations with MimicrEE2 (Vlachos & Kofler, 2018) using the qff mode to illustrate the influence of the genetic architecture on the evolution of phenotypic variance during the adaptation to a new trait optimum (Figure 1a). With 189 founder haplotypes (Barghi *et al.*, 2019), we simulated quantitative gene expression traits under the control of different numbers of loci (M = 5, 25, 50, 100, 200 and 1000) in a population with an effective population size of 300 (reflecting the estimated effective population size in an experimentally evolving population (Barghi *et al.*, 2019)). We used the empirical linkage map from *Drosophila simulans* (Howie *et al.*, 2019) to account for linkage. For each trait, we assume an additive model and a negative correlation (r = -0.7, reflecting the observed relationship from Otte *et al.* (2020)) between the effect size and starting frequency (Figure 1b). We used the *correlate()* function implemented in "fabricatr" R package (Blair *et al.*, 2019) to generate the effect sizes. The distribution of the effect sizes of loci controlling a trait can vary depending on the shape parameter of gamma sampling process (shape = 0.5, 2.5 and 100, Figure 1c). The mean of the effect sizes from 3 different shape parameters was standardized to 1/M. As the focus of our study is on expression traits, we used the distribution of heritabilities obtained from empirical gene expression data (Ayroles et al. 2009 and this study, see supplementary information) to assign the environmental/technical variance to a simulated trait (see supplementary information; Supplementary figure S1) (i.e. For each simulated trait, its heritability and associated environmental/technical variance is sampled from the empirical distribution). We do not simulate de novo mutations as they are not contributing to adaptation in this short time scale (Burke *et al.*, 2010). To simulate stabilizing selection with trait optimum shift, we provided the Gaussian fitness functions with mean of $\overline{X_{anc.}} + a\sqrt{V_{anc.}}$ and standard deviation of $b\sqrt{V_{anc.}}$, where $\overline{X_{anc.}}$ is the ancestral phenotypic mean and $V_{anc.}$ is the ancestral genetic variance (Figure 1a). Parameter "a" determines the distance of optimum shift, which is set to one (similar to the empirical case, Figure 4b) or three (Adopted from (Hayward & Sella, 2019)). Parameter "b" indicates how strong the phenotypic constraint would be when the trait optimum is reached. The value 3.6 for parameter "b" is adopted from Hayward and Sella (2019). In this study, we increase and decrease it by 50% to explore its impact (1.8 or 5.4). For the neutrality case, we assumed no variation in fitness among all individuals. For each scenario, 1000 traits were simulated, and each trait was affected by a different set of loci. We compared the phenotypic variance of selected and neutral traits for each genetic architecture. For each trait

under each scenario, the phenotypic variance was estimated at different generations and

compared to the ancestral phenotypic variance at generation 1 to illustrate the dynamic of

phenotypic variance during the evolution ($F = \sigma_x^2/\sigma_1^2$), where x stands for the number of

generations. We note that we do not assume that the ancestral population has reached an

equilibrium because the ancestral population in a typical experimental evolution study is often

phenotyped in the new environment.

<u>Power analysis</u>

We propose that the temporal dynamics of the phenotypic variance of selected traits can be

exploited to characterize the genetic basis of adaptation. This concept could be applied either

to a single trait under selection or to a group of selected traits. Depending on whether a single

trait or a group of traits is studied, different research objectives can be achieved. Hence, we

carefully examined the power and the required sample size for both types of inference based

on computer simulations.

Power estimates for the analysis of a single trait: For each simulated trait under different

genetic control, we tested whether the variance of this trait changed more than expected under

neutrality (F of 0.9 according to the neutral simulations) after 100 generations of selection.

First, we assumed that all simulated 300 individuals were phenotyped at both time points. For

the 1000 traits with each genetic architecture under each selection scenario, we calculated how

often a significant difference is detected. Since it is not possible to phenotype all individuals,

we investigated whether and how reduced sample sizes could compromise the power to infer

the adaptive architecture of a single trait. From the simulated population, we randomly selected

20 individuals for the variance estimation and tested whether the magnitude of variance change

for a given trait under selection is significantly different from neutral expectation. We

evaluated the power with the reduced sample sizes by calculating how often a significant

difference is detected among the 1000 simulated traits.

Power estimates for the analysis on a group of traits: We assessed the power to infer the

adaptive architecture for a group of selected traits under the assumption that most traits of

interest have similar level of complexity. In this case, a test on the average variance changes

(F) between a group of selected traits and another group of neutral traits is required. We

simulated 1,000 independent selected and 1,000 neutral traits for each genetic architecture and

selection regime, and estimated the variance before and after 100 generations of evolution for

the selected and neutral traits. For each trait, we measured the phenotypic variance for a sample

of 20 individuals. Next, we performed a t-test to compare the distribution of variance change

(F) between selected and neutral traits. The power was determined by 100 iterations of

sampling 20 individuals from a simulated population of 300 individuals.

<u>Experimental evolution</u>

The setup of populations and evolution experiment have been described by Barghi *et al.* (2019).

Briefly, ten outbred *Drosophila simulans* populations seeded from 202 isofemale lines were

exposed to a laboratory experiment at 28/18 °C with 12hr light/12hr dark photoperiod for more

than 100 generations. Each replicate consisted of 1000 to 1250 adults for each generation.

<u>Common garden experiment</u>

The collection of samples from the evolution experiment for RNA-Seq was preceded by two

generations of common garden (CGE). The common garden experiment was performed at

generation 103 of the evolution in the hot environment and this CGE has been described in

(Barghi *et al.*, 2019; Hsu *et al.*, 2019, 2020; Jakšić *et al.*, 2020; Lai & Schlötterer, 2022). In

brief, an ancestral population was reconstituted by pooling five mated females from 184

founder isofemale lines (Nouhaud *et al.*, 2016). No significant allele frequency differences are

expected between the reconstituted ancestral populations and the original ancestral populations

initiating the experiment (Nouhaud *et al.*, 2016). Because we evaluated phenotypes on the

population level, deleterious mutations will have a very limited impact. The reason is that

deleterious mutations are expected to occur at low frequencies in natural populations and novel

deleterious mutations occurring during the maintenance of the isofemale lines present in a

single line. In both cases, deleterious alleles are expected to represent a small fraction of the

total population. As described previously (Lai & Schlötterer, 2022), two replicates of the

reconstituted ancestral population and two independently evolved populations at generation

103 were reared for two generations with egg-density control (400 eggs/bottle) at the same

temperature regime as in the evolution experiment. Freshly eclosed flies were transferred onto

new food for mating. Sexes were separated under CO_2 anesthesia at day 3 after eclosure, left

to recover from CO_2 for two days, and at the age of five days whole-body mated flies of each

sex were snap-frozen at 2pm in liquid nitrogen and stored at -80°C until RNA extraction. More

than 30 individual male flies from two reconstituted ancestral populations (replicate no. 27 and

no. 28) and two evolved populations (replicate no. 4 and no. 9) were subjected to RNA-Seq.

The protocols of RNA extraction and library preparation are described in (Lai and Schlötterer
2022).
193
<u>RNA-Seq data analysis for mean and variance evolution</u>
The processed RNA-Seq data were obtained from Lai and Schlötterer (2022). To characterize
the adaptive architecture of gene expression evolution, we compared the evolutionary
dynamics of the variance between the genes with significant mean change and those without.
The underlying assumption is that genes with significant mean expression changes are under
selection and the rest of the transcriptome is not responding to the new environment (neutral).
The genes with/without significant mean evolution for two evolved populations were taken
from (Lai & Schlötterer, 2022).
For the quantification the variance changes during adaptation for each gene, we used the
variance estimates of the expression (lnCPM) of each gene in each population from (Lai &
Schlötterer, 2022). Due to the moderate sample size, we performed additional checks for the
uncertainty of variance estimates. Jackknifing was applied to measure the uncertainty of
estimator (Fukunaga & Hummels, 1989). The procedure was conducted independently on four
populations and we calculated the 95% confidence interval of the estimated variance
(Supplementary Figure S2). Additionally, the robustness of variance estimation was also
supported by the high correlation (rho = 0.79, Spearman's rank correlation) in variance
estimates between the two independently reconstituted ancestral populations. The change of
gene expression variance was determined by the F statistics calculated as the ratio between the
variance within the ancestral population and the variance within the evolved population of each
gene. To test whether selection on mean expression generally alters the expression variance,
we compared the F statistics of genes with significant changes in mean expression to the genes
without.
216
We note that the natural log transformation (as in Lai and Schlötterer 2022) does not attempt
to quantify and partition the noise due to the read sampling process at the gene level from the
across-individual variance. Since we were primarily interested in the latter, we evaluated the
effect of read sampling noise. We did an additional check by distinguishing the true biological
variation and the measurement error ([eq1], from User's Guide of edgeR) of each gene using
the statistical method implemented in edgeR, where the true variance across individuals
(biological coefficient of variation (BCV)) of each gene can be estimated (tag-wised
dispersion).

$$Total\ CV^2 = Technical\ CV^2 + Biological\ CV^2\ [eq1]$$

Proportion of technical CV^2 in total CV^2 was calculated. We showed that the average proportion of technical CV^2 gradually decreased with increasing sample size (Supplementary Figure S3), and for our sample size (n = 22) it is 17%. The correlation between BCV^2 and the variance estimates after log transformation is 0.98 (Spearman's rho), suggesting that the two estimates are similar. As a sanity check, we repeated the comparisons of the variance changes in selected (DE) and neutral (non-DE) genes using BCV^2. Similarly, the changes in variance of putative adaptive genes are indistinguishable from the genes that do not change their mean expression (t-test, p-value > 0.05; Supplementary Figure S3).

Results

The central idea of this study is that the complexity of adaptive traits can be inferred from the trajectory of the phenotypic variance during adaptation: the phenotypic variance remains relative stable for a trait with polygenic (infinitesimal) architecture while it changes across generations for a trait with oligogenic architecture. Although this prediction has been illustrated in multiple theoretical and simulation studies (e.g.: (Jain & Stephan, 2015; Barton *et al.*, 2017; Franssen *et al.*, 2017)), its application to empirical data has been limited. A recent experimental evolution study reporting the phenotypic variance of more than 10,000 gene expression traits before and after 100 generations of adaptation to a novel environment (Lai & Schlötterer, 2022) provides an opportunity to fill this gap. Since gene expression can be viewed as quantitative traits (Ayroles *et al.*, 2009; Hill *et al.*, 2021), we apply the theoretical predictions to the empirical data to infer the genetic basis of adaptive gene expression evolution. Hence, as the first step of this study, we explored to what extent theoretical predictions can be generalized to expression traits obtained from a typical experimental evolution setting by considering a broad parameter space and accounting for linkage. Using population genetic parameters (e.g. number of starting haplotypes, effective population size) estimated from the above-mentioned evolution experiment, we simulated 1000 neutral and 1000 selected expression traits. To account for experimental and environmental noise, we used an empirical distribution of heritabilities (estimated from gene expression measurements across F1 families; see supplementary information). The simulated selection regime consists of a mild/distant shift in trait optimum with weak/intermediate/strong stabilizing selection (Figure 1a and Supplementary Figure S4) (Hayward & Sella, 2019). We assume additivity and a negative correlation between the ancestral allele frequency and the effect size of contributing loci (Otte et al. 2020) (Figure 1b). With three different distributions of effect size (Figure 1c), we investigate how the number of contributing loci affects the evolution of gene expression variance with and without selection.

We monitored the change in phenotypic variance over 100 generations, which was sufficient to reach the trait optimum for most parameter combinations (Supplementary Figure S5). We compared the change in variance relative to the start of the experiment in populations with and without selection. First, we studied a mild (one standard deviation of the ancestral phenotypic distribution) shift in trait optimum. As expected for a founder population derived from a substantially larger natural population, we find that even under neutrality, the phenotypic variance does not remain constant, but gradually decreases during 100 generations of

experimental evolution (Figure 2). This pattern is unaffected by the genetic architecture of the
neutral traits. We explain this loss of variance by the fixation of variants segregating in the
founder population and the fact that we did not simulate new mutations, as they do not
contribute to adaptation in such short time scales (Burke *et al.*, 2010). Although our simulations
used moderate population sizes, they nicely recapitulate the patterns described for populations
without drift (Jain & Stephan, 2015; Barton *et al.*, 2017; Franssen *et al.*, 2017). A pronounced
drop in phenotypic variance is observed when a trait is approaching a new optimum with few
contributing loci (Figure 2). When more loci (with smaller effects) are contributing to the
selected phenotype, the difference to neutrality becomes very small (Figure 2). In addition to
the number of contributing loci, the heterogeneity in effect size among loci and the shape of
the fitness function have a major impact. The larger the difference in effect size is, the more
pronounced was the influence of the number of contributing loci (Figure 2). The opposite effect
was seen for the width of the fitness function – a wider fitness function decreased the influence
of the number of contributing loci (Figure 2). Importantly, these patterns were not affected by
the duration of the experiment-qualitatively identical patterns were seen at different time points
until generation 200.

For a more distant trait optimum (three standard deviations of the ancestral phenotypic
distribution), we noticed some interesting dynamics that were not apparent for a closer trait
optimum (Supplementary Figure S6). The most striking one was the temporal heterogeneity of
the phenotypic variance when few loci of unequal effects are contributing. During the early
stage of adaptation, the variance increased and dropped later below the variance in the founder
population. With an increasing number of contributing loci, this pattern disappeared and
closely matched the neutral case (Supplementary Figure S6).

Overall, our simulations indicate that with a larger number of contributing loci the variance
fitted the neutral pattern better. Modifying dominance did not change the overall patterns
(Supplementary Figure S7). The large influence of key parameters of the adaptive architecture,
in particular the number of contributing loci and their effect sizes, on the temporal phenotypic
variance dynamics suggests that it should be possible to exploit this relationship to characterize
the genetic basis of phenotypic adaptation.

A potentially interesting application of the relationship between number of contributing loci
and evolution of variance is the inference of the adaptive architecture of a given trait. For a

single selected trait, a F test contrasting the reduction of variance after adaptation against

neutral loss of variance (~0.9 after 100 generations in our simulations) can be used to infer the

number of contributing loci behind its adaptive evolution. The lower bound for the number of

contributing loci can be determined by the largest number of loci that results in a significant

difference between a selected trait and the neutral expectation in all parameter combinations.

Nevertheless, the power of this approach depends strongly on the sample size. When the entire

population (N = 300) is phenotyped for the focal trait and a neutral trait, the power to detect a

significant decrease in variance of a selected trait controlled by five contributing loci is only

44% (i.e. a lack of significant difference in variance after the adaptation cannot inform us

whether the selected traits are controlled by five or more loci). With a more realistic sample

size of 20 individuals, it is not possible to infer the adaptive architecture for a single trait when

only two time points are available (Supplementary figure S8).

Alternatively, it is possible to study multiple, presumably independent, selected phenotypes

(e.g.: transcriptomic data) together. Assuming a similar genetic architecture of the selected

traits, we showed that the distribution of variance changes of 1000 selected traits with a simple

genetic architecture (five loci) can be distinguished from neutral changes. Even with a sample

size of 20 individuals, as in our experimental data (see Materials and Methods), significant

differences from neutral expectations can be detected for some parameters with a power close

to 100% (Figure 3 and Supplementary figure S9). This suggests that oligogenic and polygenic

architectures of a group of selected traits can be distinguished experimentally even with

moderate sample sizes.

As a case study, we investigated the evolution of gene expression variance in replicated

populations evolving in a new hot temperature regime and inferred the adaptive architecture of

gene expression traits. The evolved populations were derived from the same ancestral

population but evolved independently for more than 100 generations in a novel temperature

regime with daily temperature fluctuations between 18 and 28°C (Figure 4a). Rather than

relying on pooled samples which only allow mean estimates, we quantified gene expression of

19-22 individuals from reconstituted ancestral populations (F0) and two evolved populations

(F103) in a common garden setup (Lai & Schlötterer, 2021). We applied natural log

transformation to the expression of each gene (counts per million (CPM)) to fit normal

assumption for all subsequent analyses and make mean and variance independent from each

other (see Material and Methods). Principal Component Analysis (PCA) indicated that 11.9%

of the variation in gene expression can be explained by the first PC which separates evolved and ancestral populations, reflecting clear adaptive gene expression changes in response to the novel hot temperature regime (Figure 4b). The means and variances of the expression of each gene were estimated and compared between the two reconstituted ancestral populations and the two evolved populations separately (see Materials and methods). Due to the usage of different lot numbers for the RNA-Seq library preparation (Supplementary file 1), we only contrasted ancestral and evolved samples generated with the same lot number (see Materials and methods) to avoid any unnecessary confounding effects.

The comparison of ancestral and evolved populations identified 2,775 genes in the first population and 2,677 genes in the second population which significantly changed mean expression in the evolved flies (FDR<0.05, (Lai & Schlötterer, 2022)) (Figure 5a). The concordance of both populations and the similarity to previous studies of the same evolution experiment (Hsu *et al.*, 2020; Jakšić *et al.*, 2020; Lai & Schlötterer, 2022) suggests that most of the altered expression means are mainly driven by selection, rather than by drift. We scaled the gene expression change with the standard deviation in the ancestral population to approximate the selection strength on each gene. The differentially expressed genes in both populations showed a broad distribution of expression change, but the averaged mean expression changed by one standard deviation (Figure 5b), which is unlikely to be reached by drift based on our simulations (less than 1% of the neutral traits drifted by one standard deviation of the ancestral phenotypic variance). Assuming that these putative adaptive gene expression phenotypes reached the new trait optimum, this corresponds, on average, to a mild shift in trait optimum in our computer simulations. Reasoning that genes with significant mean expression changes are under selection and the rest of the transcriptome is not responding to the new environment (neutral), we compared the evolutionary changes in expression variance of genes with and without significant changes in expression mean (DE and non-DE genes), to characterize the adaptive architecture of gene expression. The underlying hypothesis is that a simple adaptive architecture will be associated with a significant drop in the expression variance of the selected genes, while for polygenic gene expression evolution, the variance remains similar to neutral genes. Our computer simulations showed that this expectation from theoretical results (Jain & Stephan, 2015; Barton *et al.*, 2017; Franssen *et al.*, 2017) also holds for moderately sized populations and linked loci (Figure 2).

The reduction in population size from a natural population to laboratory cultures would result

in a continuous reduction in phenotypic variance over time even under neutrality, as shown in

the simulations (Figure 2). Hence, a robust inference of the adaptive architecture of gene

expression relies on the comparison of putative selected (DE) and neutral (non-DE) genes. In

both replicates the expression variance of putative neutral (non-DE) genes dropped to 84% of

the ancestral variance, which is quite close to the neutral cases in our simulations. While it

would be desirable to test each putatively selected (DE) gene independently, the moderate

sample size (~20 per population) does not provide sufficient power to do so (Supplementary

figure S8). Rather, we consider all putatively selected (DE) genes jointly and compare their

variance changes to the ones of putatively neutral (non-DE) genes. Remarkably, we find that

the changes in variance of putatively selected genes with significant mean expression changes

are indistinguishable from the genes that do not change their mean expression (t test, p-value

> 0.05; Figure 5c). The magnitude of gene expression variance changes in both sets of genes

(median F-value = 0.85 and 0.84 for DE and non-DE genes, respectively, in replicate 1, and

0.82 and 0.84 in replicate 2) is quite close to the neutral conditions in our computer simulations.

This suggests that selection on mean expression does not significantly change the expression

variance during adaptation. This is further supported by the lack of correlation between the

evolutionary changes in gene expression variance and mean (Lai & Schlötterer, 2022).

Although the adaptive architecture may differ between genes, we suggest that variation in

expression of most putatively adaptive genes has polygenic basis (i.e. more than five

contributing loci).

Since we only explored two time points (F0 and F103) rather than a full time series, it may be

possible that an oligogenic basis could also result in a similar phenotypic variance as neutral

phenotypes like what is observed for a polygenic architecture (Supplementary Figure S9). This

can be seen in an intuitive case when a single/few major effect allele(s) starts at a low frequency

and becomes fixed (Yoo *et al.*, 1980). Because an oligogenic basis results in a highly parallel

genomic selection response (Supplementary Figure S10), it is possible to distinguish polygenic

and oligogenic architectures with phenotypes from two time points only when genomic data

are available. Because the genomic signature in the same experiment uncovered a highly

heterogeneous selection response (Barghi *et al.*, 2019), we can exclude the unlikely explanation

of an oligogenic architecture resulting in a similar expression variance as non-selected genes.

Rather, we suggest that the adaptive response in gene expression could be explained by a

polygenic architecture (i.e. more than five contributing loci).

Discussion

Population genetics has a long tradition of characterizing adaptation based on the genomic signature of selected loci (Nielsen, 2005). Nevertheless, for selected phenotypes with a polygenic architecture, the contribution of individual loci to phenotypic change may be too subtle to be detected with classic population genetic methods (Pritchard *et al.*, 2010). Although identification of many small-effect loci is challenging, it may be possible to determine the number of contributing loci based on the dynamics of phenotypic variance.

Inspired by the Castle-Wright estimator (Castle, 1921) that estimates the number of loci contributing to a quantitative trait from the phenotypic variance of the F2, we propose that the temporal heterogeneity of the phenotypic variance can potentially be used to infer the number of loci contributing to the adaptive response of a phenotype as well as other parameters of the adaptive architecture. Reasoning that experimental evolution is probably the best approach to obtain phenotypic time series, we performed computer simulations specifically tailored to typical experimental evolution studies with *Drosophila*. We demonstrated that, in an experimental evolution setup, the temporal dynamics of the phenotypic variance is strongly affected by the number of contributing loci and other parameters of the adaptive architecture such as the distribution of effect size.

We acknowledge that the proposed approach for the characterization adaptive architecture critically depends on the power to distinguish the variance changes of selected traits from the neutral expectations. Since the accurate inference of gene expression variance is key, we partitioned the expression variance of each gene among three independent F1 families consist of almost identical, heterozygous individuals (see Supplementary Information). We show that on average, ~50% of expression variance can be attributed to random error - a combination of technical (library preparation, RNA-extraction, sequencing) and biological (stochastic gene expression differences among individuals) noise. We included this error in our simulations by adopting the empirical heritability distribution (heritability of every gene inferred from this experiment) such that random errors relative to the ancestral phenotypic variance were introduced to the calculation of phenotypic values at each generation for each simulated trait. The heritability estimates in our experiment may be too conservative (i.e. too high) because we assumed no environmental heterogeneity. We also used heritability estimates from another gene expression study on *Drosophila* which included more lines and accounted for

environmental heterogeneity (Ayroles et al., 2009). The results are qualitatively similar (Figure
3).

Given the limited power to infer the number of contributing loci for each expression phenotype,
we grouped all putatively selected expression traits and limited our inference of the adaptive
architecture to these genes jointly. This approach makes the implicit assumption that all
expression traits in this group are independent of each other and have similar level of
complexity in their adaptive architectures. This joint inference could potentially compromise
the detection of a minor subset of genes whose expression evolution is under simple genetic
control.

Our results may also provide some insights into the molecular basis of gene expression
evolution. Gene expression is modified by many trans-acting factors and some cis-regulatory
variation (Hill *et al.*, 2021). Adaptive gene expression changes can be either driven by
polymorphism in *cis*-regulatory elements or by *trans*-acting variants. While interspecific
differences in gene expression are predominantly caused by *cis*-regulation (Wittkopp *et al.*,
2004), intraspecific variation is mostly driven by trans regulatory changes (Wittkopp *et al.*,
2008; Suvorov *et al.*, 2013). Adaptive gene expression changes which are well-characterized
on the molecular level typically have a *cis*-regulatory basis that is not only frequently
associated with the insertion of a transposable element (e.g.: (Daborn *et al.*, 2002)) but also
sometimes with multiple regulatory variants (Endler *et al.*, 2018). Two lines of evidence
suggest that *cis*-regulatory variation cannot be the driver of adaptive gene expression changes
observed in this study. First, the mutational target size is too small to harbor a sufficiently large
number of alleles segregating in the founder population. Second, too few recombination events
occur during the experiment to uncouple regulatory variants located on a given haplotype such
that they could generate a signal of polygenic adaptation. More likely, the polygenic adaptive
architecture of gene expression change reflects the joint effects of many *trans*-acting variants.

Because we could only analyze phenotypic data from two time points, the founder population
and replicate populations evolved for 103 generations, we were not able to obtain a more
quantitative estimate of the number of contributing loci, in particular as other parameters of the
adaptive architecture are not known and need to be co-estimated. In addition, with only two
time points, for a few parameter combinations, an oligogenic response can also result in a
similar phenotypic variance change as a polygenic one but with a much higher parallel response

of genomic markers (Supplementary Figure S10). Hence, not only more time points describing
the phenotypic trajectory, but also some genomic data could contribute to inferring the adaptive
architecture in experimental evolution studies.
473
The extension of the concept proposed in this study to natural populations could face several
challenges that warrant extra caution and further investigation. First, the above-mentioned
challenge of sample size would be more pronounced in a natural setup. Second, phenotypic
time series over evolutionary relevant time scales are costly (but see (Clutton-Brock &
Pemberton, 2004)) and third, the distinction of environmental heterogeneity from genetic
changes is considerably more challenging than under controlled laboratory conditions.
480

Figure 1. Simulating polygenic adaptation to a shift in trait optimum with different parameter combinations. a. For the computer simulations we consider a quantitative trait (in black) experiencing a sudden shift in trait optimum under stabilizing selection. The underlying fitness functions are illustrated in red. The new trait optimum is shifted from the ancestral trait mean by one/three standard deviation of the ancestral trait distribution. The strength of stabilizing selection is modified by changing the variance of the fitness function: 1.8, 3.6 and 5.4 standard deviations of the ancestral trait distribution. **b.** The negative correlation between the allele frequencies and the effect sizes (r = -0.7, estimated in Otte *et al.* (2020)). We consider such negative correlation when assigning the effect sizes to variants underlying a simulated trait. **c.** The distribution of effect sizes of the contributing loci is determined by the shape parameter (α) of gamma sampling process (α = 0.5, 2.5 and 100).

Figure 2. The trajectory of changes in phenotypic variance during adaptation to a mild optimum shift. The changes in phenotypic variance within 200 generations adapting to a moderate optimum shift (orange) are compared to the changes under neutrality (grey) on y axis. The average variance changes (F) of 1000 simulated traits is calculated as the ratio of phenotypic variance between each evolved time point (generation *x*) and the ancestral state (σ_x^2/σ_1^2). The translucent band indicates the 95% confidence interval for 1000 simulated traits. The simulations cover traits controlled by varying numbers of loci underlying the adaptation with three different distributions of effect sizes (columns) under different strength of stabilizing selection (rows). Only traits with the most (dotted lines, 1000 loci), intermediate (dash lines, 50 loci) and the least (solid lines, 5 loci) polygenic architectures are shown. In all scenarios, the variance of the trait decreases drastically when the adaptation is controlled by a small

505   number of loci (orange solid lines; 5 loci). While, for traits with extremely polygenic basis, the

506   phenotypic variance stays stable over time (orange dotted lines).

Figure 3. Power to detect significant differences in variance change between groups of selected and neutral traits. For each set of the 1000 traits controlled by different numbers of loci (x-axis) with varying effect sizes (columns) under each selection strength (rows), we calculated how often the variance change differs between 1000 neutral and 1000 selected traits after 100 generations (y-axis). The same genetic architecture is assumed for all 1000 selected traits with a sample size of 20. For each parameter combination, we have almost 100% power to detect a significant difference when all traits were under simple genetic control (5 loci). The power gradually decreases with an increasing number of loci (orange). A lack of significant difference indicates polygenic architecture of the adaptive traits. The results remain unaffected when we used heritability estimates from Ayroles et al., (2009) (purple).

Figure 4. Schematic overview of the experimental procedures (a) and the divergence in
gene expression during experimental evolution (b). a. Experimental evolution: starting
from one common founder population, two replicate populations evolved for 100 generations
in a hot laboratory environment fluctuating between 18 and 28°C. Common garden Experiment:
after 100 generations, the two evolved replicate populations were maintained together with the
reconstituted ancestral population for two generations in a hot laboratory environment
fluctuating between 18 and 28°C. After this common garden procedure, about 30 males from
each population were subjected to RNA-Seq. **b.** Principle Component Analysis (PCA) of the
transcriptomic profiles of individuals from the ancestral population (blue) and the hot-evolved
population (red). Circles indicate individuals of the first replicate (Anc. No. 27 and Evo. No.
4). Triangles represent individuals of the second replicate (Anc. No. 28 and Evo. No. 9). The
two replicates were made with two different batches of library cards for RNA-Seq library
preparation. The impact of the library card batch can be seen from the separation of the
ancestral replicates which were reconstituted from the same iso-female lines.

538 **Figure 5. Evolution of phenotypic mean and variance over 100 generations of adaptation**

539 **in empirical data. a.** The evolution of gene expression means during adaptation in the two

540 replicates. For the genes with significant changes (DE, in orange), the changes are correlated

541 between replicates (Spearman's rho = 0.53). For the genes without significant changes (non-

542 DE, in grey), the correlation between replicates is much lower (Spearman's rho = 0.2). **b.** The

543 evolution of gene expression means scaled by the ancestral variation. For the DE genes (in

544 orange), the median change falls around 1 standard deviation of the ancestral expression value,

545 suggesting mild shift in trait optimum in the novel environment. For the non-DE genes (in

546 grey), the changes in expression are mostly negligible. The same pattern is seen in the second

547 replicate. **c.** The change in expression variance during adaptation for DE and non-DE genes. In

548 both replicates, the distribution of variance changes is indistinguishable between DE genes

(orange) and non-DE genes (grey) (t-test, $p > 0.05$ for both replicates). Almost identical pattern

is observed when the biological coefficient of variation (BCV; see materials and methods) is

used to estimate the expression variance (Supplementary Figure S4).

Acknowledgments

Special thanks to David Houle, who provided fantastic support during the collection and
establishment of the isofemale lines in Florida. We thank all member of the Institut für
Populationsgenetik for discussion. We are grateful to Reinhard Bürger, David Houle, Dagný
Ásta Rúnarsdóttir and anonymous reviewers for helpful comments on earlier versions of the
manuscript. Neda Barghi, François Mallard and Kathrin Otte contributed to the common
garden experiment. Illumina sequencing was performed at the VBCF NGS Unit
(www.vbcf.ac.at). This work was support by the Austrian Science Funds (FWF, W1225) and
the European Research Council (ERC, ArchAdapt).

Author contribution

W.Y.L and C.S. conceived the study. V.N. prepared all RNA-Seq and supervised the
maintenance of the evolution experiment. A.M.J supervised the common garden experiment.
W.Y.L performed the simulation and data analysis. W.Y.L. and C.S. wrote the manuscript.

Data accessibility statement

All sequencing data will be available in European Nucleotide Archive (ENA) under the
accession number PRJEB37011 upon publication.

Competing interests

The authors declare no competing interests.

Correspondence and requests for materials should be addressed to C.S.

References

Ayroles, J.F., Carbone, M.A., Stone, E.A., Jordan, K.W., Lyman, R.F., Magwire, M.M., *et al.* (2009) Systems genetics of complex traits in *Drosophila melanogaster*. *Nature Genetics*, **41**, 299–307.

Barghi, N., Hermisson, J. & Schlötterer, C. (2020) Polygenic adaptation: a unifying framework to understand positive selection. *Nature Reviews Genetics*, **21**, 769–781.

Barghi, N., Tobler, R., Nolte, V., Jakšić, A.M., Mallard, F., Otte, K.A., *et al.* (2019) Genetic redundancy fuels polygenic adaptation in *Drosophila*. *PLOS Biology*, **17**, e3000128.

Barton, N.H., Etheridge, A.M. & Véber, A. (2017) The infinitesimal model: Definition, derivation, and implications. *Theoretical Population Biology*, **118**, 50–73.

Barton, N.H. & Keightley, P.D. (2002) Understanding quantitative genetic variation. *Nature Reviews Genetics*, **3**, 11–21.

Barton, N.H. & Turelli, M. (1987) Adaptive landscapes, genetic distance and the evolution of quantitative characters. *Genetical Research*, **49**, 157–173.

Blair, G., Cooper, J., Coppock, A., Humphreys, M., Rudkin, A. & Fultz, N. (2019) fabricatr: imagine your data before you collect it.

Boyle, E.A., Li, Y.I. & Pritchard, J.K. (2017) An expanded view of complex traits: From polygenic to omnigenic. *Cell*, **169**, 1177–1186.

Bulmer, M.G. (1972) The genetic variability of polygenic characters under optimizing selection, mutation and drift. *Genetical Research*, **19**, 17–25.

Burke, M.K., Dunham, J.P., Shahrestani, P., Thornton, K.R., Rose, M.R. & Long, A.D. (2010) Genome-wide analysis of a long-term evolution experiment with *Drosophila*. *Nature*, **467**, 587–590.

Careau, V., Wolak, M.E., Carter, P.A., Garland, T. & Jr. (2015) Evolution of the additive genetic variance-covariance matrix under continuous directional selection on a complex behavioural phenotype. *Proceedings. Biological sciences*, **282**.

Castle, W.E. (1921) An improved method of estimating the number of genetic factors concerned in the cases of blending inheritance. *Science (New York, N.Y.)*, **54**, 223.

Chevalet, C. (1994) *An approximate theory of selection assuming a finite number of quantitative trait loci*. *Genetics Selection Evolution*.

Chippindale, A.K., Chu, T.J.F. & Rose, M.R. (1996) Complex trade-offs and the evolution of starvation resistance in Drosophila melanogaster. *Evolution*, **50**, 753–766.

Clutton-Brock, T.H. & Pemberton, J.M. (Josephine M. (2004) *Soay sheep : population dynamics and selection on St. Kilda*. Cambridge University Press.

Daborn, P.J., Yen, J.L., Bogwitz, M.R., Goff, G. Le, Feil, E., Jeffers, S., *et al.* (2002) A single

p450 allele associated with insecticide resistance in *Drosophila*. *Science (New York, N.Y.)*, **297**,

2253–6.

Endler, L., Gibert, J.-M., Nolte, V. & Schlötterer, C. (2018) Pleiotropic effects of regulatory

variation in *tan* result in correlation of two pigmentation traits in *Drosophila melanogaster*.

Molecular Ecology, **27**, 3207–3218.

Falconer, D.S. & Mackay, T.F.C. (1963) *Introduction to quantitative genetics. Poultry Science*.

CreateSpace Independent Publishing Platform.

Ferea, T.L., Botstein, D., Brown, P.O. & Rosenzweig, R.F. (1999) Systematic changes in gene

expression patterns following adaptive evolution in yeast. *Proceedings of the National*

Academy of Sciences of the United States of America, **96**, 9721–6.

Franssen, S.U., Kofler, R. & Schlötterer, C. (2017) Uncovering the genetic signature of

quantitative trait evolution with replicated time series data. *Heredity*, **118**, 42–51.

Fukunaga, K. & Hummels, D.M. (1989) Leave-one-out procedures for nonparametric error

estimates. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, **11**, 421–423.

Hayward, L.K. & Sella, G. (2019) Polygenic adaptation after a sudden change in environment.

bioRxiv, 792952.

Hill, M.S., Zande, P. Vande & Wittkopp, P.J. (2021) Molecular and evolutionary processes

generating variation in gene expression. *Nature Reviews Genetics*, **22**, 203–215.

Howie, J.M., Mazzucco, R., Taus, T., Nolte, V. & Schlötterer, C. (2019) DNA motifs are not

general predictors of recombination in two drosophila sister species. *Genome Biology and*

Evolution, **11**, 1345–1357.

Hsu, S.-K., Jakšić, A.M., Nolte, V., Barghi, N., Mallard, F., Otte, K.A., *et al.* (2019) A 24 h

age difference causes twice as much gene expression divergence as 100 generations of

adaptation to a novel environment. *Genes*, **10**, 89.

Hsu, S.-K., Jakšić, A.M., Nolte, V., Lirakis, M., Kofler, R., Barghi, N., *et al.* (2020) Rapid sex-

specific adaptation to high temperature in *Drosophila*. *eLife*, **9**.

Huang, Y. & Agrawal, A.F. (2016) Experimental evolution of gene expression and plasticity

in alternative selective regimes. *PLOS Genetics*, **12**, e1006336.

Jain, K. & Stephan, W. (2015) Response of polygenic traits under stabilizing selection and

mutation when loci have unequal effects. *G3 (Bethesda, Md.)*, **5**, 1065–74.

Jakšić, A.M., Karner, J., Nolte, V., Hsu, S.-K., Barghi, N., Mallard, F., *et al.* (2020) Neuronal

function and dopamine signaling evolve at high temperature in *Drosophila*. *Molecular Biology*

and Evolution, **37**, 2630–2640.

Kawecki, T.J., Lenski, R.E., Ebert, D., Hollis, B., Olivieri, I. & Whitlock, M.C. (2012)

Experimental evolution. *Trends in Ecology & Evolution*, **27**, 547–560.

Keightley, P.D. & Hill, W.G. (1989) Quantitative genetic variability maintained by mutation-

stabilizing selection balance: sampling variation and response to subsequent directional

selection. *Genetical Research*, **54**, 45–58.

Kimura, M. & Crow, J.F. (1964) The number of alleles that can be maintained in a finite

population. *Genetics*, **49**, 725–738.

Lai, W. & Schlötterer, C. (2022) Evolution of phenotypic variance in response to a novel hot

environment. *Molecular Ecology*, **31**, 934–945.

Lande, R. (1976) Natural selection and random genetic drift in phenotypic evoltion. *Evolution*,

30, 314–334.

Lenski, R.E., Travisano, M., Larison, B. & Moritz, C. (1994) Dynamics of adaptation and

diversification: a 10,000-generation experiment with bacterial populations. *Proceedings of the*

National Academy of Sciences, **91**, 6808–6814.

Liu, X., Li, Y.I. & Pritchard, J.K. (2019) Trans effects on gene expression can drive omnigenic

inheritance. *Cell*, **177**, 1022-1034.e6.

Mackay, T.F.C., Stone, E.A. & Ayroles, J.F. (2009) The genetics of quantitative traits:

challenges and prospects. *Nature Reviews Genetics*, **10**, 565–577.

Mallard, F., Nolte, V., Tobler, R., Kapun, M. & Schlötterer, C. (2018) A simple genetic basis

of adaptation to a novel thermal environment results in complex metabolic rewiring in

Drosophila. *Genome Biology*, **19**, 119.

Nielsen, R. (2005) Molecular signatures of natural selection. *Annual Review of Genetics*, **39**,

197–218.

Nouhaud, P., Tobler, R., Nolte, V. & Schlötterer, C. (2016) Ancestral population reconstitution

from isofemale lines as a tool for experimental evolution. *Ecology and Evolution*, **6**, 7169–

7175.

Otte, K.A., Nolte, V., Mallard, F. & Schlötterer, C. (2020) The adaptive architecture is shaped

by population ancestry and not by selection regime. *bioRxiv*, 2020.06.25.170878.

Pélabon, C., Hansen, T.F., Carter, A.J.R. & Houle, D. (2010) Evolution of variation and

variability under fluctuating, stabilizing, and disruptive selection. *Evolution*, **64**, 1912–25.

Penna, A., Melo, D., Bernardi, S., Oyarzabal, M.I. & Marroig, G. (2017) The evolution of

phenotypic integration: How directional selection reshapes covariation in mice. *Evolution*, **71**,

2370–2380.

Pritchard, J.K., Pickrell, J.K. & Coop, G. (2010) The genetics of human adaptation: Hard

sweeps, soft sweeps, and polygenic adaptation. *Current Biology*, **20**, R208–R215.

Romero, I.G., Ruvinsky, I. & Gilad, Y. (2012) Comparative studies of gene expression and the

evolution of gene regulation. *Nature Reviews Genetics*, **13**, 505–516.

Signor, S.A. & Nuzhdin, S. V. (2018) The evolution of gene expression in cis and trans. *Trends*

in Genetics, **34**, 532–544.

Sork, V.L. (2017) Genomic studies of local adaptation in natural plant populations. *Journal of*

Heredity, **109**, 3–15.

Suvorov, A., Nolte, V., Pandey, R.V., Franssen, S.U., Futschik, A. & Schlötterer, C. (2013)

Intra-specific regulatory variation in *Drosophila pseudoobscura. PLoS ONE*, **8**, e83547.

Turelli, M. (1984) Heritable genetic variation via mutation-selection balance: Lerch's zeta

meets the abdominal bristle. *Theoretical Population Biology*, **25**, 138–193.

Vlachos, C. & Kofler, R. (2018) MimicrEE2: Genome-wide forward simulations of Evolve

and Resequencing studies. *PLOS Computational Biology*, **14**, e1006413.

Wittkopp, P.J., Haerum, B.K. & Clark, A.G. (2004) Evolutionary changes in cis and trans gene

regulation. *Nature*, **430**, 85–88.

Wittkopp, P.J., Haerum, B.K. & Clark, A.G. (2008) Regulatory changes underlying expression

differences within and between *Drosophila* species. *Nature Genetics*, **40**, 346–350.

Yoo, B.H., Nicholas, F.W. & Rathie, K.A. (1980) Long-term selection for a quantitative

character in large replicate populations of *Drosophila melanogaster. Theoretical and Applied*

Genetics, **57**, 113–117.

Supplementary text

<u>Estimating genetic variance in gene expression across F1 families</u>

In this study, we included another experiment to evaluate how much of the expression variance

can be explained by genetic variation by performing RNA-Seq on individual flies, with 3-4

individuals each from three iso-female lines maintained at the same density and culturing

conditions (Supplementary file 1).

Six out of 184 founder iso-female lines from the evolution experiment and were maintained

for one generation with controlled egg density (400 eggs/bottle) in the same environment as

the main experiment (12h 28°C with light followed by 12h 18 °C with dark conditions). Using

the offspring, we generated three crosses between two of the six lines each: FL 138 x FL 137,

FL 157 x FL 112, FL 123 x FL 127: we combined 50 virgin females from one of the lines with

50 males from the other line, let them lay eggs under density control as above and maintained

and froze their F1 offspring in the same way as in the main CGE: sexes were separated after

mating at the age of three days and snap-frozen at the age of five days at 2pm. From each cross,

we used four F1 males to prepare individual RNA-Seq libraries as described above.

Assuming no environmental heterogeneity, we decomposed the total variance of the expression

of each gene measured in these individuals into the genetic difference among three different

F1 families and environmental/technical error. The data were analyzed as follows:

Natural log-transformation was applied to CPMs of all genes to improve data normality (Rocke

and Durbin 2003). We tested for the genetic variance of each gene separately using analysis of

variance (ANOVA):

$$y_{ij} = \mu + \tau_i + \varepsilon_{ij},$$

Where i=1, 2, 3 (the ith families); j =1, 2, 3, 4 (the jth individuals in each cross). y_{ij} is the

observed expression level of a gene in a given sample, μ is the overall mean; τ_i is the effect of

genetic background and ε_{ij} is the random noise. We calculated the proportion of total variance

explained by random error using the following equation:

$$variance\ explained\ by\ random\ error = \frac{sum\ squares\ of\ error\ (SSE)}{sum\ squares\ of\ total\ (SST)}$$

We found that on average, around 50% of the gene expression variance can be explained by
the genetic difference among the three F1 families (Supplementary Figure S1). This implies
that a substantial within-population (between individuals) gene expression variance in our focal
experiment is contributed by the genetic components. Because we only used offspring from
single vials, we may have overestimated the heritability if the environment in the vials differs.
Nevertheless, since our heritability estimates are very similar to previous ones (Ayroles *et al.*
2009), we consider our estimates reliable.
39
The technical error will go into residual in this analysis. Since variance of lowly expressed
genes are mainly dominated by technical error, we defined a cut-off of expression value for a
more reliable variance estimate. Genes were binned based on their average expression value
(lnCPM) which ranged from -0.8 to 4.1, by bin size of 0.1. The average proportion of variance
explained by random error of each bin was calculated. The expression variance of genes with
less than 1 count per million bases (CPM) is dominated by technical errors (Supplementary
Figure S9). Thus, genes with less than 1 count per million base (CPM) were excluded for
subsequent analysis.
48

Supplementary Figure S1. Genetic variance in gene expression across F1 families. (A)
Principal component analysis (PCA) on the transcriptomes of F1 individuals from three
different crosses between the founder iso-female lines. Individuals from different families
clustered nicely based on the first two PCs. (**B**) Gene-wise heritability on the transcriptomes
of F1 individuals. On average, 48.8% of the total variance in gene expression was explained
by the genetic difference between the individuals. (**C**) Genes were binned based on their
average expression value (lnCPM) which ranged from -0.8 to 4.1, by bin size of 0.1. The
average proportion of variance explained by random error of each bin was visualized. The

59 expression variance of genes with less than 1 count per million bases (CPM) is dominated by

60 residuals.

Supplementary Figure S2. Robustness of the variance estimation using individual sequencing data. Jackknife method was applied to measure the uncertainty of variance estimation within each population. Given a sample size of K, the procedure is to estimate the variance of each gene for K times, each time leaving one sample out. The procedure was conducted independently on 4 populations (anc.27, anc.28, evo.4 and evo.9). In each panel, we visualize Jackknife approximated 95% confidence interval for the variance estimates of each gene. The genes are ordered based on the average variance estimates (black dash line) on the x-axis. The upper and lower limits of the 95% confidence interval are indicated with grey curves. The salmon line denotes the observed value of the variance estimates. In most cases, the estimates lie in the confidence interval, suggesting robust estimation.

Supplementary Figure S3. The decomposition of measurement error in RNA-Seq data and its implication for the tests of variance evolution. (A) The relationship between the average proportion of measurement error among the RNA-Seq samples and the sample sizes is visualized. As the sample size increases, the measurement error of RNA-Seq data is well controlled. The results are consistent across replicate populations. For simplicity, only the result in the first replicate of ancestral population is shown. **(B)** We observed a strong correlation between the variance estimates after log-transformation and the biological coefficient of variation (BCV) estimated from edgeR, which claims to exclude the measurement error. Such strong correlation suggests that measurement error in RNA-Seq data wouldn't have strong impact on the test of variance of evolution. The results are consistent across replicate populations. For simplicity, only the result in the first replicate of ancestral

population is shown. **(C)** Using the BCV estimates, we repeated the comparison between the

variance changes ($F = \frac{BCV_{evo}^2}{BCV_{anc}^2}$) of the genes with and without significant evolution in mean

expression. In both replicates, the distribution of variance changes is indistinguishable between

DE genes and non-DE genes (t-test, p > 0.05 for both replicates).

Supplementary Figure S4. The evolutionary scenario for distant optimum shift. We

consider the case when a quantitative trait (in black) experiences a sudden shift in trait optimum

under stabilizing selection. The imposed fitness functions (F.F.) are illustrated in red. The new

trait optimum is set away from the ancestral trait mean by three standard deviation of the

ancestral trait distribution for distal shift. To vary the strength of stabilizing selection, the

variance of the fitness function is set as 1.8, 3.6 and 5.4 standard deviation of the ancestral trait

distribution.

Supplementary Figure S5. The changes in phenotypic mean when adapting to optimum shift. The changes in phenotypic mean after 100 generations adapting to a mild/distant optimum shift (orange/red) are compared to the changes under neutrality (grey) on y axis. The changes in phenotypic mean are scaled by the standard deviation of the ancestral trait distribution. The simulations cover traits controlled by varying numbers of loci underlying the adaptation (x axes) with three different distributions of effect sizes (columns) and under different strength of stabilizing selection (rows). For each scenario, 1000 traits have been performed. In most cases, the traits under selection (orange/red) shift their means by one/three standard deviation of the ancestral trait distribution (i.e. reaching the new trait optimum) while

111   the neutral traits (grey) stay unchanged. The error bar indicates the 95% confidence interval

112   for 1000 simulated traits.

Supplementary Figure S6. The trajectory of expected changes in phenotypic variance when adapting to a distant optimum shift. The changes in phenotypic variance within 200 generations adapting to a distant optimum shift (red) are compared to the changes under neutrality (grey) on y axis. The average change in variance among 1000 traits (F) is calculated as the ratio of phenotypic variance between each evolved time point (generation x) and the ancestral state (σ_x^2/σ_1^2). The translucent band indicates the 95% confidence interval for 1000 simulated traits. The simulations cover traits controlled by varying numbers of loci underlying the adaptation with three different distributions of effect sizes (columns) and under different strength of stabilizing selection (rows). For each scenario, we simulated 1000 traits. Only traits with the most (dotted lines, 1000 loci), intermediate (dash lines, 50 loci) and the least (solid lines, 5 loci) polygenic architectures are shown. Unlike the continuous decreasing pattern in

126   the cases with mild optimum shifts, the variance of traits controlled by a few loci (5 loci) with

127   largely dispersed effects would increase first and then decrease when the effect sizes of

128   contributing loci are dispersed (red solid lines). Nevertheless, for traits with extremely

129   polygenic basis, the phenotypic variance always stays stable over time (red dotted lines).

Supplementary Figure S7. The expected changes in phenotypic variance for traits controlled by dominant and recessive alleles. The changes in phenotypic variance after 100 generations adapting to a mild optimum shift (orange) are compared to the changes under neutrality (grey) on y axis. The change in variance (F) is calculated as the ratio between the evolved and ancestral phenotypic variance ($\sigma_{100}^2/\sigma_0^2$). This simulation covers traits controlled by varying numbers of loci underlying the adaptation (x axes) with recessive, additive and dominant effects. For each scenario, we simulated 1000 traits. No matter how the dominance varies, the variance of the trait decreases drastically when the adaptation is controlled by a small number of loci. As the number of contributing loci increases, the phenotypic variance becomes more stable. The error bar indicates the 95% confidence interval for 1000 simulated traits.

**Supplementary Figure S8. Power for detecting significant variance changes for a single
trait.** With each genetic architecture (x-axis) under each selection scenario, we calculated how
often we can detect a significant difference between the variance change of a single trait after
100 generations of selection and the neutral expectation (F of 0.9 according to the neutral
simulations) (y-axis). The power is largely limited by the sample size. When all 300 individuals
were considered, only 40% of power is observed under simple genetic control (5 loci). When
the sampling processes of 20 individuals from the 300 simulated individuals was considered,
there is nearly no power for the inference.

Supplementary Figure S9. Power for detecting significant different variance change between a group of selected and neutral traits under distant optimum shift. For each set of the 1000 traits controlled by different numbers of loci (x-axis) with varying effect sizes (columns) under each selection strength (rows), we calculated how often the variance changes of these 1000 selected traits differs from 1000 neutral traits after 100 generations (y-axis). The same genetic architecture is assumed for all 1000 selected traits with with a sample size of 20. In five out of the nine parameter combinations, we have nearly 100% power to detect a significant difference when all traits were under simple genetic control (5 loci). The power gradually decreases when the number of loci increased. Nevertheless, in the other parameter combinations, exception can be found for the traits controlled by a few loci (5 loci) with largely dispersed effects (see Discussion).

Supplementary Figure S10. Parallelism in genomic response across 10 evolved replicates for traits under the control of five loci and 100 loci. For the loci with allele frequency change of at least 10% in 100 generations, the average Pearson's correlation coefficient of the frequency change between all pairs of two evolved replicates was calculated to describe the parallelism of the evolution at these loci. An average across loci is used to obtain a general parallelism. 100 traits with 10 evolution replicates have been performed for each scenario. With five contributing loci, the genomic evolution of the contributing loci is more parallel across the 10 evolved replicates compared to the case when with 100 contributing loci.

Titles and legend for supplementary files

Supplementary file 1. Library information of the sample in this study. This file provides a
list of all sequenced samples and the library information.

References

Ayroles J. F., M. A. Carbone, E. A. Stone, K. W. Jordan, R. F. Lyman, *et al.*, 2009 Systems

genetics of complex traits in Drosophila melanogaster. Nat. Genet. 41: 299–307.

https://doi.org/10.1038/ng.332

Rocke D. M., and B. Durbin, 2003 Approximate variance-stabilizing transformations for gene-

expression microarray data. Bioinformatics 19: 966–72.