Title

VISUAL network analysis reveals the role of BEH3 as a stabilizer in the

secondary vascular development in Arabidopsis

Short title

BEH3 acts as a stabilizer in cambium.

Authors

Tomoyuki Furuya[1,2], Masato Saito[2], Haruka Uchimura[2], Akiko Satake[3], Shohei

Nosaki[4], Takuya Miyakawa[4], Shunji Shimadzu[1,2], Wataru Yamori[2,4], Masaru

Tanokura[4], Hiroo Fukuda[2], Yuki Kondo[1,2*].

Affiliations

[1] Department of Biology, Graduate School of Science, Kobe University, 1-1

Rokkodai, Kobe 657-8501, Japan.

[2] Department of Biological Sciences, Graduate School of Science, The University

of Tokyo, Tokyo 113-0033, Japan.

[3] Department of Biology, Faculty of Science, Kyushu University, Fukuoka, 819-

0395, Japan.

[4] Department of Applied Biological Chemistry, Graduate School of Agricultural

and Life Sciences, The University of Tokyo, Tokyo 113-8657, Japan.

Corresponding author e-mail address

* Correspondence: pkondo@tiger.kobe-u.ac.jp

Abstract

During secondary growth in plants, vascular stem cells located in the cambium continuously undergo self-renewal and differentiation throughout the lifetime. Recent cell-sorting technique enables to uncover transcriptional regulatory framework for cambial cells. However, the mechanisms underlying the robust control of vascular stem cells have not been understood yet. By coexpression network analysis using multiple transcriptome datasets of an ectopic vascular cell transdifferentiation system using Arabidopsis cotyledons, VISUAL, we newly identified a cambium-specific gene module from an alternative approach. The cambium gene list included a transcription factor BES1/BZR1 homolog 3 (BEH3), whose homolog BES1 is known to control vascular stem cell maintenance negatively. Interestingly, the vascular size of the *beh3* mutants showed a large variation, implying the role of BEH3 as a stabilizer. BEH3 almost lost the transcriptional repressor activity and functioned antagonistically with other BES/BZR members via competitive binding to the same motif BRRE. Indeed, mathematical modeling suggests that the competitive relationship among BES/BZRs leads to the robust regulation of vascular stem cells.

INTRODUCTION

Continuous stem cell activity relies on a strict balance between cell proliferation and cell differentiation. Plant vascular stem cells located in the cambium proliferate by themselves, then continuously giving rise to conductive tissues consisting of xylem and phloem cells, as bifacial stem cells (Shi et al., 2017; De Rybel et al., 2016; Fischer et al., 2019). Previous genetic studies have indicated that vascular stem cells are maintained by cell-cell communication via a peptide hormone tracheary element differentiation inhibitory factor (TDIF) and its specific receptor PHLOEM INTERCALATED WITH XYLEM (PXY) / TDIF RECEPTOR (TDR) (Ito et al., 2006; Fisher et al., 2007; Hirakawa et al., 2008). For the maintenance of vascular stem cells, the TDIF-PXY/TDR signaling promotes cell proliferation via WUS-RELATED HOMEOBOX 4 (WOX4) and WOX14 (Hirakawa et al., 2010, Etchells et al., 2010) and represses cell differentiation via BRI1-EMS-SUPPRESSOR 1 (BES1) (Kondo et al., 2014). In addition, the TDIF-PXY/TDR signaling genetically cross-talks with several peptide or hormonal signaling pathways, thereby forming a complicated regulatory network (Han et al., 2018; Etchells et al., 2012; Kondo and Fukuda, 2015). As another attempt to identify stem cell regulators, transcriptome analysis using fluorescence activated cell sorting (FACS) and fluorescence activated nuclear sorting (FANS) method has been conducted. FACS transcriptome analysis with a (pro)cambium marker *ARABIDOPSIS RESPONSE REGULATOR 15* (*ARR15*) *promoter:GFP* educed a list of cambium-expressed transcription factors, including *AINTEGUMENTA* (*ANT*), *KNOTTED-like from Arabidopsis thaliana 1* (*KNAT1*), *LOB DOMAIN-CONTAINING PROTEIN 3* and *4* (*LBD3* and *LBD4*), *SHORT VEGETATIVE PHASE* (*SVP*), and *PETAL LOSS* (*PTL*) (Zang et al. 2019). Furthermore, in the inflorescence stem, FANS transcriptome analysis using tissue-specific promoters including *PXY/TDR* promoter for proximal cambial cells and *SUPPRESSOR OF MAX2 1-LIKE PROTEIN 5* (*SMXL5)* promoter for distal cambial cells provided informative vascular tissue-specific gene expression profiles (Shi et al., 2020).

Nevertheless, the molecular mechanism enabling a robust control of vascular
stem cells remains largely unknown.

The vascular system consists of various types of vascular cells at
different developmental stages, preventing the sequential analysis of vascular
development with a focus on bifacial stem cells. To simply understand a complex
biological phenomenon such as vascular development, reconstitutive approach
has great experimental advantages with regard to molecular genetic studies
(Kondo 2018). We previously established a tissue culture system, Vascular cell
Induction culture System Using Arabidopsis Leaves (VISUAL), which can
synchronously and efficiently induce xylem and phloem cells from mesophyll cells
via vascular stem cell (Kondo et al., 2016). The loss-of-function mutant for the
phloem regulator ALTERED PHLOEM DEVELOPMENT (APL) lacks phloem
sieve element differentiation. In addition to the microarray data with the *apl*
mutants, the datasets of the FACS with a phloem marker gene, *SIEVE ELEMENT*
OCCLUSION RELATED 1 (*SEOR1*) in VISUAL enabled the construction of a
coexpression gene network from early to late phloem sieve element
differentiation process (Kondo et al. 2016). Although such a network analysis is
useful for temporal dissection of vascular differentiation process, this gene
network is still only limited to phloem sieve element differentiation. Recently, we
discovered that the loss-of-function *bes1* mutants showed a dramatic impairment
in the differentiation of xylem and phloem cells, resulting in the accumulation of
vascular stem cells in the VISUAL (Kondo et al., 2014; Saito et al., 2018; Kondo
et al., 2015). Then we further integrated *bes1* transcriptome and time-course
(every 6 hours) transcriptome datasets in order to visualize the gene network for
the entire vascular differentiation process.

In this study, we newly established a coexpression network of the
VISUAL differentiation process utilizing multiple transcriptome datasets including
the *bes1* mutants. As a result, we succeeded in the identification of gene modules
corresponding to not only phloem but also xylem and cambial cells. These
classifications were validated with a comparison to the already-existing vascular
FACS and FANS datasets. Through the network analysis, one of BES/BZR

homolog BEH3 was found as a potential vascular stem cell regulator. Indeed, a

knock-out mutant of BEH3, *beh3-1*, exhibited a high variation in vascular size

among individual samples, suggesting that BEH3 functions as a stabilizer of

vascular stem cells. Further genetic and molecular studies revealed that BEH3

has an opposite function toward other BES/BZR homologs by interfering with

other BES/BZR homologs via competitive binding to the BR response element

(BRRE). Moreover, mathematical modeling analysis showed that the competitive

relationship among BEH3 and other BES/BZR homologs enables the robust

control of vascular stem cells.

RESULTS

Coexpression network analysis using the multiple VISUAL transcriptome

datasets identifies a cambium-related gene module

To find out novel regulators for vascular stem cells, we conducted a gene network

analysis using the transdifferentiation system VISUAL (Figure 1A; Kondo et al.,

2015). First, we selected 855 VISUAL-induced genes that were up-regulated

highly (more than eight-fold) in microarray data (0 h vs 72 h after VISUAL

induction) (Figure 1B; Supplemental Data set 1). Since the selected 855 VISUAL-

induced genes should include xylem-, phloem- and cambial-related genes, we

constructed a coexpression gene network for these genes with the use of multiple

VISUAL transcriptome datasets including (1) 6-h interval time-course, (2) cell-

sorting with the phloem marker *pSEOR1:SEOR1-YFP*, (3) phloem deficient *apl*

mutants, and (4) *bes1-1* mutants (Figure 1A). Here the *bes1-1* mutant

transcriptome datasets were used for dissecting cambial cells and differentiated

xylem/phloem cells, because the *bes1-1* mutants significantly accumulate

cambial cells due to the inhibition of xylem and phloem differentiation in the

VISUAL (Figure 1A). Coexpression network using the WGCNA package divided

the 855 VISUAL-induced genes into four distinct modules (Module-I to IV) and 11

outgroup genes (Others, O) (Figure 1C). Expression of Module-III and -IV genes

were down-regulated in the *bes1-1* mutants (Figure 1C), suggesting that these

modules were considered as differentiated xylem and phloem cells-related ones.

In addition, expression of Module-III genes was high in the SEOR1-positive cell-

sorting datasets and was down-regulated in the *apl* mutants, whereas expression

of Module-IV genes was unchanged in the cell-sorting and the *apl* mutant

datasets (Figure 1D), suggesting that Module-III and -IV represents phloem-

related and xylem-related genes, respectively. Consistent with this idea, the

known phloem-related genes such as *RAB GTPASE HOMOLOG C2A* (*RABC2A*),

BREVIS RADIX (*BRX*), *GLUCAN SYNTHASE-LIKE 7* (*GSL07*), and *NAC*

DOMAIN CONTAINING PROTEIN 45 (*NAC045*), were found in Module-III,

whereas xylem-related genes such as *IRREGULAR XYLEM 3* (*IRX3*), *XYLEM*

CYSTEINE PEPTIDASE 1 (*XCP1*), and *VASCULAR-RELATED NAC-DOMAIN 6*

(*VND6*), were in Module-IV (Figure 1C). On the other hand, expression of

Module-I and -II genes was induced at earlier timing and was remained higher

even in the *bes1-1* mutants than that of Module-III and -IV genes (Figure 1B and

1D), suggesting that Module-I and -II genes correspond to (pro)cambial-related

genes. To validate these characterizations, we investigated gene expression of

each module in the transcriptome datasets of FACS in roots and FANS in stems

obtained using the cell-type specific markers (Brady et al. 2007, Zang et al. 2019,

Shi et al. 2020). Expectedly, expression of Module-III and -IV genes were quite

high in the phloem specific *APLpro* datasets and the xylem specific *S18* and

VND7pro datasets, respectively (Figure 1E; Supplemental Figure 1A). As for

Module-I and II, the genes in Module-I were relatively highly expressed in the

procambium (ARRIII ± BAP) and expression of Module-II gene is more specific

to the developing cambium and cambium (ARRI ± BAP and ARRII ± BAP) in the

FACS datasets (Figure 1E). These results suggest that Module-I and Module-II

represents procambium-related and cambium-related genes, respectively.

Consistent with this idea, expression of Module-I genes was expressed much

earlier than that of Module-II genes (Figure 1B).

We further analyzed the Module-II genes for analyzing stem cell

regulation. The Module-II consists of a total of 394 genes including some known

cambium-related genes, such as *PYX/TDR*, *SMXL5*, *ANT*, *LBD3*, *LBD4*, and

HOMEOBOX GENE 8 (*HB-8*) (Figure 1B – 1D). Of them, 32 genes encoding

transcription factors were investigated in more detail (Supplemental Data set 2).

Previous studies have revealed that vascular stem cells locate in the overlapped

region of *PXY/TDR* and *SMXL5* expression, which marks proximal/xylem and

distal/phloem side of cambium, respectively (Shi et al. 2019). According to the

FANS transcriptome datasets (Shi et al., 2020), 13 out of the 32 transcription

factor genes are expressed mainly in *PXY/TDRpro* and/or *SMXL5pro* (Figure 1F).

They included not only known cambial regulators such as *ANT* (Randall et al.,

2015) and *LBD4* (Zang et al., 2019; Smit et al., 2020）, but also novel candidates

such as *BEH3*, *DOF AFFECTING GERMINATION 1* (*DAG1*), and *At1g63100*

(Figure 1E). The analysis of these newly selected genes may provide a novel

insight into regulation of cambial activity.

BEH3 stabilizes secondary vascular development

We recently showed that BES1 and its closest homolog BRASSINAZOLE

RESISTANT1 (BZR1) redundantly function to promote vascular stem cell

differentiation in VISUAL (Saito et al., 2018). Therefore, we chose BEH3 as a

potential vascular stem cell-specific regulator for further studies. Among 6

BES/BZR homologs in *Arabidopsis thaliana*, only *BEH3* was found in the

cambium-related module. Indeed, only *BEH3* expression levels were remarkably

increased in the time course RNA-seq data of the VISUAL (Figure 2B), and the

GUS expression domain of *pBEH3:GUS* gradually expanded over time (Figure

2C). Consistent with the observation in the VISUAL, *pBEH3:GUS* was

preferentially expressed in the vascular tissues including cambium of leaves and

hypocotyls (Figure 2D and 2E). Then to investigate the function of BEH3 in

vascular development, we analyzed the BEH3 knock-out mutant, *beh3-1*, which

has a T-DNA insertion on its 1st exon (Lachowiec et al., 2018). Here we

compared vascular secondary growth in the WT and the *beh3-1* mutants grown

under short-day condition. The area inside the cambium in the *beh3-1* mutants

were slightly but significantly decreased when compared with the WT (Figure 2F

and 2G; Supplemental Figure 2). In addition, we realized that the *beh3-1* mutants

seem to have a higher variation in area inside the cambium than the WT (Figure

2H). Indeed, the values of coefficient of variation (CV) between the *beh3-1*

mutants and WT plants were significantly different in two statistical analyses:

asymptotic and Modified signed-likelihood ratio test (M-SLRT) (Feltz and Miller,

1996; Krishnamoorthy and Lee, 2014) (Figure 2I). These results suggest that the

presence of BEH3 contributes to not only promoting but also stabilizing

secondary vascular development.

BEH3 functions oppositely to BES1 in vascular development

In *Arabidopsis thaliana*, 6 BES/BZR transcription factors have the redundant

function in brassinosteroid (BR) signaling (Yin et al., 2005; Lachowiec et al.,

2018; Chen et al., 2019a; Chen et al., 2019b; Nolan et al., 2020). As previously

reported (Kondo et al., 2014; Saito et al., 2018), *bes1-D*, which has a mutation in

protein degradation PEST sequence, exhibited a reduced number of (pro)cambial

cell layers between differentiated xylem (red) and phloem (green) in hypocotyls

(Figure 3A). To compare the function of BEH3 and BES1, vascular phenotype of

gain-of-function mutant for BEH3 was analyzed. Since BEH3 does not possess

any obvious PEST domain, we generated a transgenic stable line of β-estradiol-

inducible BEH3 overexpressor (*pER8:BEH3-CFP*). In contrast to the *BES1* gain-

of-function mutants, *BEH3* overexpression lines showed an increase in the

number of (pro)cambium layers (Figure 3B). In association with this tendency,

the vascular area was also enlarged in the two independent *BEH3*

overexpression lines (Figure 3C and 3D; Supplemental Figure 3), which is not

contradictory to the *beh3-1* mutant phenotype. These results suggest that BEH3

has an opposite function to BES1 during secondary vascular development.

As described above, the *bes1-1* mutants disrupt cell differentiation from

vascular stem cells in the VISUAL (Figure 1A). To investigate the function of

BEH3 and other BES/BZR homologs, we evaluated the contribution of each

BES/BZR homolog to xylem differentiation with a semi-automatic calculation

method using a xylem indicator, BF-170 (Nurani et al., 2020) in VISUAL (Figure

4A and 4B; Supplemental Figure 4). As for the analysis with single mutants, *beh4-*

1 as well as *bes1-1* and *bzr1-1* showed reduced xylem differentiation ratio in the
VISUAL compared with the WT (Fig. 4C and 4D; Supplemental Figure 5),
indicating that BEH4 also promotes xylem differentiation. On the other hand,
xylem differentiation ratio in *beh3* as well as *beh1* and *beh2* were comparative to
the WT (Figure 4B and 4C). Next, we examined the impact of mutating a BES1
homolog in the *bes1-1* mutant background as double mutants. Only the *bes1-1*
beh3-1 double mutants showed a high xylem differentiation ratio, similar to that
observed in the WT, whereas none of the other mutants rescued the *bes1*
phenotype (Figure 4B and 4C; Supplemental Figure 5 and 6). The *bes1* mutants
disrupt the differentiation from vascular stem cells into not only xylem but also
phloem cells (Saito et al., 2018), accompanied by the decrease in expression
levels of xylem-specific (*IRX3*) and phloem-specific (*SEOR1*) marker genes
(Figure 4E). This reduction in the *bes1-1* mutants was recovered up to WT levels
by the additional *beh3-1* mutation (Figure 4E). Quantitative expression analysis
also indicated that the single *beh3-1* mutants slightly increase the expression
levels of *IRX3* and *SEOR1* at 48 h, when compared to the WT (Figure 4E;
Supplemental Table 1). These genetic data suggest that BEH3 functions
oppositely to BES1 also in the VISUAL.
245

BEH3 has a competitive relationship with other BES/BZR homologs
To unravel the molecular basis of the antagonistic function of BEH3, we
examined the functional difference among BEH3 and other BES/BZR proteins.
Amino acid sequence alignments suggested no obvious difference in the known
functional domains/motifs, such as DNA-binding domain (DBD) and EAR motif
(Yin et al., 2002; Wang et al., 2002; Oh et al., 2014). It is known that BES1 and
BZR1 repress the transcription of a BR biosynthetic gene, *DWARF4* (*DWF4*), by
directly binding to the BRRE and/or E-box motif in the *DWF4* promoter (He et al.,
2005; Sun et al., 2010; Yu et al., 2011; Chung et al., 2011; Nosaki et al., 2018).
Therefore, we tested the transcriptional control activity of BES/BZR members
against the *pDWF4: Emerald Luciferase (ELUC)* construct transiently expressed
in *Nicotiana benthamiana* leaves. The signal intensity of ELUC gradually

decreased from 3 to 9 h after activating BES1, BZR1, and BEH4 by induction

system using dexamethasone (DEX) and rat glucocorticoid receptor (GR)

(Aoyama et al., 1995), whereas BEH3 activation did not alter ELUC activity

(Figure 5A). Electrophoretic mobility shift assay (EMSA) for the BRRE core motif

and E-box motif with DBD of BES/BZR members indicated that BEH3 can bind

to the BRRE and E-box, similar to BES1, BZR1, and BEH4 (Nosaki et al., 2018)

(Figure 5B; Supplemental Figure 7). These results suggest that BEH3 binds to

the BRRE and E-box motif but exhibits much lower transcriptional repressor

activity than other BES/BZR homologs. When BEH3 and BES1 were

simultaneously expressed in *N. benthamiana* leaves, repressor activity of BES1

against *pDWF4:ELUC* was weakened (Supplemental Figure 8). By contrast, co-

expression of BZR1 with BES1 did not show such a combinatory repressive effect

(Supplemental Figure 8), suggesting the possibility that BEH3 interferes with

other BES/BZR homologs via the binding competition.

In the VISUAL, the mutation in *BES1* did not affect *BEH3* expression, and

the mutation in *BEH3* did not affect *BES1* expression in VISUAL (Supplemental

Figure 9), indicating that there seems not to be an epistatic relationship between

BEH3 and *BES1*. Then to genetically investigate the competitive relationship

among *BEH3* and other *BES/BZR* homologs, several combinations of triple

mutants were analyzed in VISUAL. Although the *beh3-1* mutant allele could

completely rescue the phenotype of the *bes1-1* single mutants (Figure 4B – 4D),

restoration of xylem differentiation by the *beh3-1* mutant allele was only partial in

the background of *bes1-1 bzr1-2* double mutants (Figure 4F and 4G). Such a

partial rescue was not detected in the *bes1-1 bzr1-2 beh4-1* triple mutants at all

(Figure 4F and G). Consistent with the xylem differentiation ratio, the *bes1-1 bzr1-*

2 beh3-1 triple mutants were intermediate between WT and the *bes1-1 bzr1-2*

double mutants in terms of expression levels of xylem and phloem marker genes

(Figure 4H). Taken together, these results suggest that BEH3 acts

antagonistically with other homologs in a competitive manner in the VISUAL.

Competitive inhibition by BEH3 may be limited to the transcriptional repressor activity of other BES/BZR homologs

Previous studies showed that a *bes1* hextuple mutants, in which all BES/BZR members are knocked-out, exhibits male sterility (Chen et al., 2019a). If BEH3 always has a competitive role against other BES/BZR homologs, *bes1 bzr1 beh1 beh2 beh4* quintuple mutants are expected to show the same phenotype as *bes1 bzr1 beh1 beh2 beh3 beh4* hextuple mutants. Then we generated a *bes1* hextuple mutants (*bes1-1 bzr1-2 beh1-2 beh2-5 beh3-3 beh4-1*) and a *bes1* quintuple mutants (*bes1-1 bzr1-2 beh1-2 beh2-5 beh4-1*) using CRISPR/Cas9 system. Consistent with the previous reports, the newly established *bes1* hextuple mutants showed shorter siliques with no seed (Figure 6A and 6B; Supplemental Figure 10). However, the *bes1* quintuple mutants successfully produced seeds in siliques (Figure 6A and 6B). In addition, while the primary root length of the *bes1* hextuple mutants were shorter than the WT, that of the *bes1* quintuple mutants were comparable to the WT (Figure 6C and 6D). These results suggest that BEH3 partially remains the same function to other BES/BZR homologs.

BES1 and BZR1 is known to function as both transcriptional activators and repressors (Yin et al., 2005; He et al., 2005; Nolan et al., 2020). Then, we investigated to which extent BEH3 affects the expression of known BES/BZR-target genes in planta. Here we used the β-estradiol-inducible *BEH3* overexpressor *pER8:BEH3-CFP,* in which expression of *BEH3* was increased upon β-estradiol treatment (Figure 6E; Supplemental Figure 11). *DWF4* and *BRASSINOSTEROID-6-OXIDASE 2* (*BR6ox2*), as BES1-direct repressed genes (He et al., 2005; Sun et al., 2010; Yu et al., 2011), were remarkably repressed in the constitutively active mutant for *BES1* and *BZR1* (*bes1-D bzr1-D*) (Figure 6F), whereas *pER8:BEH3-CFP* transgenic plants significantly increased the mRNA levels of both *DWF4* and *BR6ox2* upon β-estradiol treatment (Figure 6E; Supplemental Figure 11; Supplemental Table 2). These results suggest that the non-canonical BEH3 can passively inhibit the transcriptional repressor activity of endogenous BES1 homologs by competing for the same binding motif. However,

BEH3 overexpression did not suppress mRNA levels of BES1-direct activated
genes, *SAUR-AC1* and *INDOLE-3-ACETIC ACID INDUCIBLE 19* (*IAA19*) (Yin et
al., 2005; Sun et al., 2010; Oh et al., 2012) (Figure 6E), whose expression was
highly induced in the *bes1-D bzr1-D* double mutants (Figure 6F). Together with
the genetic results of the *bes1* high-order mutants, our results suggest that
competitive relationship among BES/BZR members may be limited mainly to their
transcriptional repressor activity.
326

BEH3 contributes to increasing robustness via the competition with other
homologs

Our observation revealed that the *beh3* mutants increased the variation in the
size of vascular area (Figure 2). Then to understand the significance of the
competitive relationship between BEH3 and other BES/BZR members in
phenotypic variation, a mathematical model for competitive binding was
constructed with a focus on the transcriptional repressor activity. In this
mathematical modeling, we set the repressor activity of BEH3 and other
homologs at 0.95× and 0.8×, respectively, when they occupied one binding site
in the target gene promoter. Here, we assumed three binding sites in the putative
promoter and predicted the output as vascular cell number, when the binding rate
of BEH3 (P_{BEH3}) and other homologs (P_O) was varied (Figure 7A). When P_{BEH3}
was increased, the average predicted outputs went up because the accessibility
of the binding sites for other homologs was decreased (Figure 7B). This
competitive modeling can account for the phenotype that *BEH3* overexpression
leads to the enlargement of vascular area (Figure 3C and 3D). On the other hand,
further modeling revealed that a reduction in P_{BEH3} decreased the means of
output value, which corresponds to smaller area inside the cambium in the *beh3-*
1 mutants when compared with the WT (Figure 2F - 2H). Moreover, because the
beh3-1 mutants exhibited higher variation in the vascular size than the WT
(Figure 2F - 2H), we also simulated the shift of the CV when the value of P_{BEH3}
was decreased. As a result, a reduction in P_{BEH3} increased the CV of the output
(Figure 7C), which mimics the vascular phenotype of the *beh3-1* mutants.

Collectively, our results indicate that the cambium-expressed BEH3 competitively

inhibits transcriptional repressor activity of other BES/BZR homologs, thereby

minimizing the fluctuation of the vascular stem cell activity.

DISCUSSION

In this study, we constructed the gene coexpression network during the VISUAL

differentiation process (Figure 1). With the help of the existing FACS and FANS

transcriptome datasets, our network analysis categorized vascular-related genes

into four distinct modules; procambium (Module-I), cambium (Module-II), phloem

(Module-III) and xylem (Module-IV). Thus, reconstitute approach using the

VISUAL is effective to dissect a hierarchical process of vascular cell

differentiation and helpful for isolating genes involved in this process. Although

the FACS datasets (Zang et al. 2019) almost share a similar tendency with our

network prediction (Figure 1; Supplemental Figure 1), the overlapping rate of

cambial genes was not so high (12.9 %, 45 genes/394 genes of Module- II). The

low rate may result from differential cell state of cambium/vascular stem cells

between "*in vivo*" and "*in vitro*" and/or technical differences between cell-sorting

and whole tissue sampling. Therefore, combinatory approach by bioinformatics

will allow the accurate and efficient identification of gene-of-interest expressed in

the vasculature, which is usually masked by its deep location. Indeed, we

successfully identified a novel cambium regulator BEH3 in the gene lists

combined the Module-II genes with the FANS datasets (Shi et al., 2020). Also in

the xylem and phloem development, there remain so many unidentified genes.

Moreover, it is poorly understood how the cambium is established after

procambium development. Therefore, our coexpression network will be

informative for further studies on the sequential vascular development.

Although previous studies indicated redundant roles of BES/BZR family

members in BR signaling (Yin et al., 2005; Lachowiec et al., 2018; Chen et al.,

2019a; Chen et al., 2019b; Nolan et al., 2020), little has been reported about their

functional differences. Here we found BEH3 as a non-canonical member of the

BES/BZR family. *BEH3* is expressed preferentially in the vascular cambium,

whereas *BES1* and *BZR1* expression is observed ubiquitously (Yin et al., 2002).

Moreover, our results indicated the opposite function between BEH3 and BES1.

In the context of *in vivo* vascular development, the gain-of-function *bes1-D*

mutants accelerate cell differentiation of vascular stem cells and consequently

decreased the number of (pro)cambial cell layer, eventually resulting in

diminishment of the vascular size (Figure 2) (Kondo et al., 2014). Conversely,

BEH3 overexpression led to the increase in the number of (pro)cambial cell layer

with enlargement of the vascular size. Consistently, the *beh3* loss-of-function

mutants showed a decreased size of vascular area. Considering that BEH3 has

a repressive effect on vascular cell differentiation in the VISUAL, these results

suggest that BEH3 has a positive role in the maintenance of vascular stem cells

probably by inhibiting excess cell differentiation (Figure 7D). Molecular analyses

suggested that BEH3 retains a binding capability to the BRRE but loses a

transcriptional repressor activity. Such a competitive action leads to the opposite

behavior of BEH3. As a similar example in mammals, JunB, a transcription factor,

exhibits weaker activity than its homolog, c-Jun, can repress c-Jun-mediated

transactivation (Deng and Karin, 1993). In this study, our modeling and genetic

analysis with the *beh3* mutants suggested that the presence of a weak or non-

active type member in a transcription factor family enables a fine-tuned control of

signaling outputs.

Taken together, we conclude that the competition among BES/BZR

family members in the cambium potentially enables the stable and robust

maintenance of vascular stem cells for ensuring continuous radial growth.

However, it remains unclear why only BEH3 partially lacks the transcriptional

repressor activity and why the competitive suppression by BEH3 is limited to the

repressor activity. Further comparative interactome analysis between BEH3 and

other BES1 homologs will uncover the mechanism underlying functional

differences among BES/BZR proteins.

METHODS

Plant materials

Seeds of *Arabidopsis bes1-1* (SALK_098634), *beh1-1* (SAIL_40_D04), *beh3-1* (SALK_017577), and *beh4-1* (SAIL_750_F08) mutants were obtained from the Arabidopsis Biological Resource Center (ABRC), Ohio, USA. The *bzr1-1* single mutants and *bes1-1 bzr1-2* double mutants were generated previously using the CRISPR system (Saito et al., 2018), and *bes1-1 beh1-1*, *bes1-1 beh3-1*, *bes1-1 beh4-1*, *bes1-1 bzr1-2 beh3-1,* and *bes1-1 bzr1-2 beh4-1* mutants were generated by crossing. Chen et al. reported that the *bes1-1* mutants expresses one of the two splice variants of *BES1* (*BES1-S*); however, the other splice variant (*BES1-L*), which is functionally more important, is completely eliminated (Chen at al., 2019b, Jiang et al., 2015). The *bes1-1 bzr1-2 beh2-5 beh4-1* quadruple mutants were generated by the CRISPR/Cas9 system using pKAMA-ITACH vector (Tsutsui et al., 2017). The single guide RNA (sgRNA; Supplemental table 3.) sequence corresponding to the target gene, *BEH2*, was cloned into the *pKIR1.1* vector. The resulting construct was transformed into the *bes1-1 bzr1-2 beh4-1* triple mutants. In the T2 generation, *Cas9*-free plants with homozygous *beh2-5* mutation were selected as described previously (Tsutsui et al., 2017). The *bes1* quintuple mutants (*bes1-1 bzr1-2 beh1-2 beh2-5 beh4-1*) were also generated using the CRISPR/Cas9 system, which targeted the *BEH1* locus in *bes1-1 bzr1-2 beh2-5 beh4-1*. To generate the *bes1-1 beh3-2* double mutants and *bes1* hextuple mutants (*bes1-1 bzr1-2 beh1-2 beh2-5 beh3-3 beh4-1*), the *BEH3* locus was edited using the CRISPR/Cas9 system in *bes1-1* single mutants and *bes1* quintuple mutants, respectively. To generate a construct for *BEH3* overexpression under the control of a β-estradiol-inducible promoter, the coding sequence of *BEH3* (At4g18890.1) was amplified from Col-0 cDNA by PCR and then cloned into pENTR/D-TOPO (Life Technologies). Using LR clonase II (Life Technologies), the region between *attL1* and *attL2* in the entry vector was cloned into the destination vector, pER8-GW-CFP, which expressed the cyan fluorescent protein (CFP)-tagged proteins upon estrogen application (Ohashi-ito et al., 2010). To generate the *pBEH3:GUS* construct, a 1,997 bp fragment upstream of *BEH3* was amplified from Col-0 genomic DNA by PCR and cloned

into pENTR/D-TOPO. The region between *attL1* and *attL2* in the entry vector was

cloned into the destination vector, pGWB434. These constructs were transformed

into Col-0 plants by the floral dip method (Clough et al., 1998). All genotypes used

in this study were in the Columbia background. Plants were grown on

conventional half-strength Murashige and Skoog (1/2 MS) agar plates (pH 5.7) at

22°C under continuous light, unless stated otherwise.

Vascular cell Induction culture System Using Arabidopsis Leaves (VISUAL)

Plants were analysed using VISUAL as described previously (Kondo et al., 2016),

with a slight modification. Briefly, cotyledons of 6-day-old seedlings grown in

liquid 1/2 MS medium were cultured under continuous light in MS-based VISUAL

induction medium containing 0.25 mg/l 2,4-D, 1.25 mg/l kinetin, and 20 µM bikinin.

To observe ectopic xylem differentiation, a xylem indicator BF-170 (Sigma-

Aldrich) was added to the induction medium (Nurani et al., 2020). At 4 days after

induction, cotyledons were fixed in ethanol: acetic acid (3:1, v/v) and mounted

with a clearing solution (chloral hydrate: glycerol: water = 8:1:2 [w/w/v]). Images

obtained using a fluorescent stereomicroscope (M165, Laica) were binarized

using the "Threshold" function of the ImageJ software (Schneider et al., 2012).

Xylem differentiation ratio was calculated based on the BF-170 positive area cut-

off by suitable thresholding per a whole cotyledon area.

Microarray Experiments

To obtain the time-course maicroarray datasets, VISUAL was performed as

described above. After VISUAL induction, cultured cotyledons (6 to 10

cotyledons) were collected every six hours. Total RNA was extracted from

cultured cotyledons using an RNeasy Plant Mini Kit (Qiagen). Microarray

experiments were performed using the Arabidopsis Gene 1.0STArray

(Affymetrix) according to the standard Affymetrix protocol.

Construction of the Coexpression Network

Based on the VISUAL time-course maicroarray data, the VISUAL-induced genes
were defined as >8-fold changed genes compared with VISUAL for 0 and 72 h.
Median normalized values in $\log_2$ scale of the VISUAL-induced genes (855
genes) were obtained from multiple transcriptome datasets: time-course, wild
type versus *bes1-1* (Saito et al. 2018)*,* wild type versus *apl* (Kondo et al. 2016),
and SEOR1 cell-sorting data (Kondo et al. 2016). As described previously (Kondo
et al., 2016), the VISUAL coexpression network was constructed using the
weighted gene coexpression network analysis (WGCNA) package (Langfelder
and Horvath, 2008). The adjacency matrix was calculated with soft thresholding
power, which was chosen based on the criterion of scale-free topology (fit index
=0.8).

Confocal laser scanning microscopy
BF-170 signals in VISUAL samples were visualized using the FV1200 confocal
laser scanning microscope (Olympus). Z-series fluorescence images were
obtained at excitation and detection wavelengths of 473 and 490–540 nm,
respectively. Z-projection images were created using the ImageJ software.

Quantitative reverse transcription PCR (qRT-PCR)
Total RNA was extracted from cotyledons subjected to VISUAL experiments or
whole seedlings in other experiments using RNeasy Plant Mini Kit (Qiagen).
Quantitative PCR was performed using a LightCycler (Roche Diagnostics).
UBQ14 (for VISUAL experiments) and *ACT2* (for other experiments) were used
as internal controls. Gene-specific primers and TaqMan Probe sets are listed in
Supplemental Table 4.

RNA-seq
Total RNA was extracted from cotyledons of Col-0 plants subjected to VISUAL
experiments (Kondo et al., 2015) at 0, 30, and 72 h after induction using RNeasy
Plant Mini Kit (Qiagen). RNA-seq analysis was carried out by Hokkaido System
Science Co., Ltd.

Expression and purification of BES/BZR family proteins

MBP-fused DNA-binding domains (DBDs) of BES1 (20T–103R) and BZR1 (21A–104R) were expressed and purified as described previously (Nosaki et al., 2018). The DBDs of BEH3 (4G–88S) and BEH4 (4G–89T) were cloned into pMAL-c2X (New England Biolabs) and subsequently transformed into *Escherichia coli* Rosetta (DE3) cells (Novagen). MBP-fused DBDs of BEH3 and BEH4 were expressed and purified in the same way as BES1 and BZR1.

Analysis of BES/BZR proteins by size exclusion chromatography (SEC)

The purified proteins were loaded onto a Superdex 200 GL 10/300 (GE Healthcare) column and eluted with a buffer containing 20 mM Tris-HCl (pH 7.5), 250 mM NaCl, 1 mM DTT, and 5% glycerol. To estimate the multimerization state of BES/BZR family proteins, the following standards were used: thyroglobulin (*Mr* 670,000), bovine globulin (*Mr* 158,000), chicken ovalbumin (*Mr* 44,000), equine myoglobin (*Mr* 17,000), and vitamin B-12 (*Mr* 1,350) (Bio-Rad).

EMSA

EMSAs of BES/BZR family proteins were performed as described previously (Nosaki et al., 2018).

Luciferase (LUC) assay

Coding sequences (CDSs) of *BES1-L* (*At1g19350.3*), *BZR1* (*At1g75080.1*), and *BEH4* (*At1g78700.1*) were cloned and fused to the ligand-binding domain of the rat glucocorticoid receptor (GR). An approximately 1,952 bp sequence upstream of the *DWF4* transcription start site was cloned and fused to ELUC with the PEST domain (Tamaki et al., 2020) (Toyobo). *Rhizobium radiobacter* GV3101 MP90 strain harboring both effector and reporter constructs as well as a p19k suppressor construct was infiltrated into *Nicotiana benthamiana* leaves. After 2 days, leaf discs excised from the transformed leaves were incubated with 200 µM D-luciferin (Wako) for 2 h in white 24-well plates (PerkinElmer). Then, 10 µM

dexamethasone (DEX) was added into each well, and LUC activity was

measured using the TriStar2 LB942 (Berthold) system, as described previously

(Tamaki et al., 2020).

Histological analysis

Cross-sections of hypocotyls of seedlings were prepared as described previously

(Kondo et al., 2014). Samples were fixed in FAA solution containing 40% (v/v)

ethanol, 2.5% (v/v) acetic acid, and 2.5% (v/v) formalin. The fixed samples were

gradually dehydrated using an ethanol dilution series and then embedded in

Technovit 7100 resin (Kulzer). Sections of 5 μm thickness were prepared using

the RM2165 microtome (Leica). Sliced samples were stained with 0.1% (w/v)

toluidine blue for 1 min.

Mathematical modeling

A mathematical model for the competitive binding between BEH3 and other

homologs was constructed based on the assumptions that 1) there are n binding

sites at the promoter, and 2) the weak-type BEH3 is competitively superior to the

strong-type homologs in terms of binding to the promoter. The weak-type BEH3

occupies the vacant sites first, followed by the strong-type homologs, which

occupy the remaining vacant sites. The binding probabilities of the weak-type

BEH3 and strong-type homologs (p_{BEH3} and p_O, respectively) are proportional

to the transcript abundance of *BEH3* and other homologs, respectively. Using

p_{BEH3} and p_O, the probability that *i* BEH3 and *j* homologs bind to the promoter is

given as the product of two binomial distributions as follows:

559
$$f(i,j|p_{BEH3}, p_O, n) = \binom{n}{i} p_{BEH3}^i (1 - p_{BEH3})^{n-i} \times \binom{n-i}{j} p_O^j (1 - p_O)^{n-i-j}.$$

Binding of BEH3 and other homologs to the binding sites suppresses the

transcription level of direct target genes leading to the repression of the

proliferation of stem cells. The effect of the binding sites of BEH3 and other

homologs on the cell proliferation rate of stem cells was designated as α and β,

respectively. Because the effect of BEH3 is weaker than that of other homologs,

$\alpha < \beta$ always holds. When *i* BEH3 and *j* homologs bind to the promoter, the cell

proliferation rate is given as $\alpha^i \beta^j$. The cell division rate should be proportional

to the area inside the cambium, which was measured in experiments because

the area inside the cambium increases with the increase in the rate of cell division.

When there are multiple individuals, the expected rate of cell division of stem cell

is calculated as follows:

571
$$E[cell\ division\ rate]\ =\ \sum_{i=0}^{n}\sum_{j=0}^{n-i}\alpha^i\beta^j\,f(i,j|p_{BEH3},p_O,n)$$

572
$$=\quad [\alpha p_{BEH3} + (1 - p_{BEH3})(\beta p_O + 1 - p_O)]^n$$

573 (1)

574 Using Equation (1), the variance of cell division rate can be calculated as:

575
$$V[cell\ division\ rate] = [\alpha^2 p_{BEH3} + (1 - p_{BEH3})(\beta^2 p_O + 1 - p_O)]^n$$

576
$$-[\alpha p_{BEH3} + (1 - p_{BEH3})(\beta p_O + 1 - p_O)]^{2n}$$

(2)

The coefficient of variation (CV) of the cell division rate was calculated as:

$$CV = \sqrt{V[cell\ dividion\ rate]}/E[cell\ division\ rate]$$

Data availability

All data is available in the manuscript or the supplementary materials. All

sequence reads were deposited in the NCBI Sequence Read Archive (SRA)

database under the following accession numbers: XXXXXXX

AUTHER CONTRIBUTIONS

T.F., M.S., and Y.K. designed the experiments, coordinated the project, and

wrote the manuscript; T.F, M.S., H.U., S.S., W.Y., S.N., T.M., and Y.K. performed

the experiments; A.S. performed mathematical model simulations; M.T. and H.F.

participated in discussions. All authors reviewed and edited the manuscript.

ACKNOWLEDGMENTS

We thank Yasuko Ozawa, Yuki Fukaya, and Akiho Suizu for technical support.

We also thank Dr. Keiko U. Torii (University of Texas, Austin, Texas) for critically

reading the manuscript and providing insightful comments. This work was funded

by the Ministry of Education, Culture, Sports, Science and Technology, Japan

(Scientific Research on Priority Areas and Scientific Research on Innovative

Areas) (17H06476 and 20H05407 to Y.K., and 19H04855 to T.M.), and by the

Japan Society for the Promotion of Science (17H05008 and 20K15815 to Y.K.,

18H06065 and 20K15813 to T.F., 16H06377 to H.F., 19K23658 to S.N., and

18KK0170 and 18H02185 to W.Y.).

REFERENCES

Shi, D., Tavhelidse, T. Thumberger, T., Wittbrodt, J., and Greb, T. (2017).

Bifacial stem cell niches in fish and plants. Curr. Opin. Genet. Dev. **45**: 28-33.

De Rybel, B., Mähönen, A. P., Helariutta, Y., and Weijers, D. (2016). Plant

vascular development: from early specification to differentiation. Nat. Rev. Mol.

Cell Biol. **17**: 30–40.

Fischer, U., Kucukoglu, M., Helariutta, Y., and Bhalerao, R. P. (2019). The

Dynamics of Cambial Stem Cell Activity. Annu. Rev. Plant Biol. **70**: 293–319.

Ito, Y., Nakanomyo, I., Motose, H., Iwamoto, K., Sawa, S., Dohmae, N., and

Fukuda, H. (2006). Dodeca-CLE peptides as suppressors of plant stem cell

differentiation. Science **313**: 842–845.

Fisher, K., and Turner, S. (2007). PXY, a receptor-like kinase essential for

maintaining polarity during plant vascular-tissue development. Curr. Biol. **17**:

1061–1066.

Hirakawa, Y., Shinohara, H., Kondo, Y., Inoue, A., Nakanomyo, I., Ogawa, M.,
Sawa, S., Ohashi-Ito, K., Matsubayashi, Y., and Fukuda, H. (2008). Non-cell-
autonomous control of vascular stem cell fate by a CLE peptide/receptor system.
Proc. Natl. Acad. Sci. USA. **105**: 15208–15213.

Hirakawa, Y., Kondo, Y.,and Fukuda, H. (2010). TDIF peptide signaling
regulates vascular stem cell proliferation via the WOX4 homeobox gene in
Arabidopsis. Plant cell **22**: 2618–2629.

Etchells, J. P., and Turner, S. R. (2010). The PXY-CLE41 receptor ligand pair
defines a multifunctional pathway that controls the rate and orientation of vascular
cell division. Development **137**: 767–774.

Han, S., Cho, H., Noh, J., Qi, J., Jung, H. J., Nam, H., Lee, S., Hwang, D.,
Greb, T., and Hwang, I. (2018). BIL1-mediated MP phosphorylation integrates
PXY and cytokinin signalling in secondary growth. Nat. plants **4**: 605–614.

Etchells, J. P., Provost, C. M., and Turner, S. R. (2012). Plant vascular cell
division is maintained by an interaction between PXY and ethylene
signalling. PLoS Genet. **8**: e1002997.

Kondo, Y., and Fukuda, H. (2015). The TDIF signaling network. Curr. Opin.
Plant Biol. **28**: 106–110.

Zhang, J., Eswaran, G., Alonso-Serra, J., Kucukoglu, M., Xiang, J., Yang, W.,
Elo, A., Nieminen, K., Damén, T., Joung, J. G., Yun, J. Y., Lee, J. H., Ragni,
L., Barbier de Reuille, P., Ahnert, S. E., Lee, J. Y., Mähönen, A. P., and
Helariutta, Y. (2019). Transcriptional regulatory framework for vascular cambium
development in Arabidopsis roots. Nat. plants **5**: 1033–1042.

Shi, D., Jouannet, V., Agustí, J., Kaul, V., Levitsky, V., Sanchez, P., Mironova,

V. V., and Greb, T. (2020). Tissue-specific transcriptome profiling of the

Arabidopsis inflorescence stem reveals local cellular signatures, The Plant Cell,

koaa019, https://doi.org/10.1093/plcell/koaa019

Kondo Y. (2018). Reconstitutive approach for investigating plant vascular

development. J. Plant Res. **131**: 23–29.

Kondo, Y., Nurani, A. M., Saito, C., Ichihashi, Y., Saito, M., Yamazaki, K.,

Mitsuda, N., Ohme-Takagi, M., and Fukuda, H. (2016). Vascular Cell Induction

Culture System Using Arabidopsis Leaves (VISUAL) Reveals the Sequential

Differentiation of Sieve Element-Like Cells. Plant cell **28**: 1250-1262.

Kondo, Y., Ito, T., Nakagami, H., Hirakawa, Y., Saito, M., Tamaki, T., Shirasu,

K., and Fukuda, H. (2014). Plant GSK3 proteins regulate xylem cell

differentiation downstream of TDIF-TDR signalling. Nat. Commun. **24**: 3504.

Saito, M., Kondo, Y., and Fukuda, H. (2018). BES1 and BZR1 Redundantly

Promote Phloem and Xylem Differentiation. Plant Cell Physiol. **59**: 590-600.

Kondo, Y., Fujita, T., Sugiyama, M., and Fukuda, H. (2015). A novel system

for xylem cell differentiation in *Arabidopsis thaliana*. Mol. Plant **8**: 612-621.

Brady, S. M., Orlando, D. A., Lee, J. Y., Wang, J. Y., Koch, J., Dinneny, J. R.,

Mace, D., Ohler, U., and Benfey, P. N. (2007). A high-resolution root

spatiotemporal map reveals dominant expression patterns. Science **318**: 801-

806.

Shi, D., Lebovka, I., López-Salmerón, V., Sanchez, P., and Greb, T. (2019).

Bifacial cambium stem cells generate xylem and phloem during radial plant

growth. Development **146**: dev171355

Randall, R. S., Miyashima, S., Blomster, T., Zhang, J., Elo, A., Karlberg, A., Immanen, J., Nieminen, K., Lee, J. Y., Kakimoto, T., Blajecka, K., Melnyk, C. W., Alcasabas, A., Forzani, C., Matsumoto-Kitano, M., Mähönen, A. P., Bhalerao, R., Dewitte, W., Helariutta, Y., Murray, J. A. (2015). AINTEGUMENTA and the D-type cyclin CYCD3;1 regulate root secondary growth and respond to cytokinins. Biol Open. **4**: 1229-36.

Smit, M. E., McGregor, S. R., Sun, H., Gough, C., Bågman, A. M., Soyars, C. L., Kroon, J. T., Gaudinier, A., Williams, C. J., Yang, X., Nimchuk, Z. L., Weijers, D., Turner, S. R., Brady, S. M., and Etchells, J. P. (2020). A PXY-Mediated Transcriptional Network Integrates Signaling Mechanisms to Control Vascular Development in Arabidopsis. Plant Cell. **32**: 319-335.

Lachowiec, J., Mason, G. A., Schultz, K., and Queitsch, C. (2018). Redundancy, Feedback, and Robustness in the *Arabidopsis thaliana BZR/BEH* Gene Family. Front. Genet. **9**: 523.

Feltz, C. J., and Miller, G. E. (1996). An asymptotic test for the equality of coefficients of variation from k population. Stat. Med. **15**: 647–658.

Krishnamoorthy, K., and Lee, M. (2014). Improved tests for the equality of normal coefficients of variation. Comput. Stat. **29**: 215–232.

Yin, Y., Vafeados, D., Tao, Y., Yoshida, S., Asami, T., and Chory, J. (2005). A new class of transcription factors mediates brassinosteroid-regulated gene expression in *Arabidopsis*. Cell **120**: 249-259.

Chen, L. G., Gao, Z., Zhao, Z., Liu, X., Li, Y., Zhang, Y., Liu, X., Sun, Y., and Tang, W. (2019a). BZR1 Family Transcription Factors Function Redundantly and

Indispensably in BR Signaling but Exhibit BRI1-Independent Function in

Regulating Anther Development in *Arabidopsis*. Mol. Plant **12**: 1408-1415.

Chen W, Lv M, Wang Y, Wang PA, Cui Y, Li M, Wang R, Gou X, and Li J.

(2019b). BES1 is activated by EMS1-TPD1-SERK1/2-mediated signaling to

control tapetum development in *Arabidopsis thaliana*. Nat. Commun. **10**: 4164.

Nolan, T. M., Vukašinović, N., Liu, D., Russinova, E., and Yin, Y. (2020).

Brassinosteroids: Multidimensional Regulators of Plant Growth, Development,

and Stress Responses. Plant Cell **32**: 295-318.

Nurani, A. M., Ozawa, Y., Furuya, T., Sakamoto, Y., Ebine, K., Matsunaga, S.,

Ueda, T., Fukuda, H., and Kondo, Y. (2020). Deep Imaging Analysis in VISUAL

Reveals the Role of YABBY Genes in Vascular Stem Cell Fate Determination.

Plant Cell Physiol. **61**: 255-264.

Yin, Y., Wang, Z. Y., Mora-Garcia, S., Li, J., Yoshida, S., Asami, T., and Chory,

J. (2002). BES1 accumulates in the nucleus in response to brassinosteroids to

regulate gene expression and promote stem elongation. Cell **109**: 181–191.

Wang, Z. Y., Nakano, T., Gendron, J., He, J., Chen, M., Vafeados, D., Yang,

Y., Fujioka, S., Yoshida, S., Asami, T., and Chory, J. (2002). Nuclear-localized

BZR1 mediates brassinosteroid-induced growth and feedback suppression of

brassinosteroid biosynthesis. Dev. Cell **2**: 505–513.

Oh, E., Zhu, J. Y., Ryu, H., Hwang, I., and Wang, Z. Y. (2014). TOPLESS

mediates brassinosteroid-induced transcriptional repression through interaction

with BZR1. Nat. commun. **5**: 4140.

He, J. X., Gendron, J. M., Sun, Y., Gampala, S. S., Gendron, N., Sun, C. Q.,
and Wang, Z. Y. (2005). BZR1 is a transcriptional repressor with dual roles in
brassinosteroid homeostasis and growth responses. Science **307**: 1634-1638.
748

749 **Sun, Y., Fan, X. Y., Cao, D. M., Tang, W., He, K., Zhu, J. Y., He, J. X., Bai, M.**
750 **Y., Zhu, S., Oh, E., Patil, S., Kim, T. W., Ji, H., Wong, W. H., Rhee, S. Y., and**
751 **Wang, Z. Y.** (2010). Integration of brassinosteroid signal transduction with the
752 transcription network for plant growth regulation in *Arabidopsis*. Dev. Cell **19**:
753 765–777.
754

755 **Yu, X., Li, L., Zola, J., Aluru, M., Ye, H., Foudree, A., Guo, H., Anderson, S.,**
756 **Aluru, S., Liu, P., Rodermel, S., and Yin, Y.** (2011). A brassinosteroid
757 transcriptional network revealed by genome-wide identification of BESI target
758 genes in *Arabidopsis thaliana*. Plant J. **65**: 634–646.
759

760 **Chung, Y., Maharjan, P. M., Lee, O., Fujioka, S., Jang, S., Kim, B., Takatsuto,**
761 **S., Tsujimoto, M., Kim, H., Cho, S., Park, T., Cho, H., Hwang, I., and Choe, S.**
762 (2011). Auxin stimulates *DWARF4* expression and brassinosteroid biosynthesis
763 in Arabidopsis. Plant J. **66**: 564–578.
764

765 **Nosaki, S., Miyakawa, T., Xu, Y., Nakamura, A., Hirabayashi, K., Asami, T.,**
766 **Nakano, T., and Tanokura, M.** (2018). Structural basis for brassinosteroid
767 response by BIL1/BZR1. Nat. Plants **4**: 771-776.
768

769 **Aoyama, T., Dong, C. H., Wu, Y., Carabelli, M., Sessa, G., Ruberti, I., Morelli,**
770 **G., and Chua, N. H.** (1995). Ectopic expression of the Arabidopsis transcriptional
771 activator Athb-1 alters leaf cell fate in tobacco. Plant Cell **7**: 1773–1785.
772

Oh, E., Zhu, J. Y,. and Wang, Z. Y. (2012). Interaction between BZR1 and PIF4
integrates brassinosteroid and environmental responses. Nat. Cell Biol. **14**: 802–
809.

Deng, T., and Karin, M. (1993). JunB differs from c-Jun in its DNA-binding and

dimerization domains, and represses c-Jun by formation of inactive heterodimers.

Genes Dev. **7**: 479-490.

Jiang, J., Zhang, C., and Wang, X. (2015). A recently evolved isoform of the

transcription factor BES1 promotes brassinosteroid signaling and development

in *Arabidopsis thaliana*. Plant Cell **27**: 361–374.

Tsutsui, H., and Higashiyama, T. (2017). pKAMA-ITACHI Vectors for Highly

Efficient CRISPR/Cas9-Mediated Gene Knockout in Arabidopsis thaliana. Plant

Cell Physiol. **58**: 46-56.

Ohashi-Ito, K., Oda, Y., and Fukuda, H. (2010). *Arabidopsis* VASCULAR-

RELATED NAC-DOMAIN6 directly regulates the genes that govern programmed

cell death and secondary wall formation during xylem differentiation. Plant Cell

22: 3461-3473.

Clough, S. J., and Bent, A. F. (1998). Floral dip: a simplified method for

Agrobacterium-mediated transformation of *Arabidopsis thaliana*. Plant J. **16**: 735-

743.

Schneider, C. A., Rasband, W. S., and Eliceiri, K. W. (2012). NIH Image to

ImageJ: 25 years of image analysis. Nat. Methods **9**: 671–675.

Langfelder, P., and Horvath, S. (2008). WGCNA: an R package for weighted

correlation network analysis. BMC bioinformatics **9**: 559.

Tamaki, T., Oya, S., Naito, M., Ozawa, Y., Furuya, T., Saito, M., Sato, M.,

Wakazaki, M., Toyooka, K., Fukuda, H., Helariutta, Y., and Kondo Y. (2020).

VISUAL-CC system uncovers the role of GSK3 as an orchestrator of vascular cell
type ratio in plants. Commun. Biol. 3: 184.

Figure 1. Classification of the VISUAL-induced genes based on the gene coexpression network analysis

(A) Schematic showing the transdifferentiation process in Col-0 (WT, left), *bes1* mutants (lower), and *apl* mutants (right).

(B) Expression profiles of the VISUAL-induced genes in the VISUAL time-course

microarray data. The VISUAL-induced genes were defined as >8-fold changed

genes when compared VISUAL at 0 and 72 h. Expression was visualized with a

heat map image according to Z score according to their expression levels. Color

bars in the right side indicates each module in accordance with (C).

(C) Coexpression network of the VISUAL-induced genes using the WGCNA

package. Four distinct modules are highlighted with different colors. A node

represents a gene. An edge indicates high correlation between nodes (TOM >

0.02).

(D) Expression profiles of the VISUAL-induced genes in the multiple VISUAL

microarray datasets. VISUAL microarray datasets were provided from each

comparison between WT and *bes1-1* (Saito et al. 2018), WT and *apl* (Kondo et

al. 2016), and SEOR1-negative and SEOR1-positive cells (Kondo et al. 2016).

(E) Expression profiles of the VISUAL-induced genes in the FACS dataset (Bardy

et al., 2007, Zang et al., 2019).

(F) Expression profiles of the cambium-related genes (Module-II) in the FANS

dataset (Shi et al., 2020). The gene names are shown on the right side.

Figure 2. BEH3 provides the stability for secondary growth in the cambium

(A) An unrooted phylogenetic tree of Arabidopsis BES/BZR family members.

(B) Time-course expression analysis of BES1 homologs genes using VISUAL.

Gene expression levels at each time point were analyzed by RNA-seq.

(C-E) GUS activity in *pBEH3:GUS* transgenic plants. GUS staining was

performed using cotyledons subjected to VISUAL (C) shoot of 20-day-old

seedling (D) and hypocotyl section of 7-day-old seedling (E).

(F) Sections of hypocotyl vasculature of WT and *beh3-1* mutant plants grown

under short-day conditions for 21 days.

(G) Relative areas inside the cambium were calculated in comparison with the

mean value of the WT (**$P < 0.01$; Student's *t*-test, $n = 28$ [WT], 30 [*beh3-1*]).

(H) Size variation inside the cambium of WT and *beh3* plants. Black circles

indicate areas inside the cambium of each individual for the odd number samples

in ascending order of (F).

(I) CV of the areas inside the cambium of WT and *beh3* plants. Statistical analysis

of differences between the WT and *beh3* mutants using Asymptotic and M-

SLART.

Scale bars = 1 mm (C), 2 mm (D), and 100 μm (E,F,I).

Figure 3. BEH3 overexpression promotes secondary growth

(A) Sections of hypocotyl vasculature of 11-day-old WT and *bes1-D* seedlings.

Red and green area indicates xylem cells and phloem cells, respectively.

(B-C) Sections of hypocotyl vasculature of β-estradiol-inducible BEH3-

overexpressing plants treated with (+) or without (-) 5 µM β-estradiol for 12 days.

Red and green area in (B) indicate xylem cells and phloem cells, respectively.

(D) Calculated vascular areas in (C) are shown using box-and-whisker plots.

White circles indicate values of each sample (*$P < 0.05$, **$P < 0.01$; Student's *t*-

test; *n* = 23).

Scale bars = 100 µm (A–C).

Fig. 4

Figure 4. BEH3 antagonistically functions against BES1 and BZR1 in VISUAL

(A) Schematic showing the transdifferentiation process in Col-0 (WT, upper) and *bes1* mutants (lower).

(B, C) Ectopic xylem cell differentiation ratio in a combination of *bes/bzr* mutants. Bright blue signal in (B) indicates xylem cell wall stained with BF-170. Quantified xylem differentiation ratio is shown using box-and-whisker plots in (C). White

circles indicate the xylem differentiation ratio for each sample. Different letters

indicate significant differences ($P < 0.05$; Tukey-Kramer test).

(D) Effects of BES/BZR family members toward vascular cell differentiation in

VISUAL. Red, blue, and grey indicate "promote", "suppress", and "no effect",

respectively.

(E) Time-course expression analysis of xylem-specific (*IRX3*) and phloem-

specific (*SEOR1*) marker genes. Relative fold changes were calculated in

comparison with the WT at 0 h after induction. Data represent mean ± standard

deviation (SD; n = 3). Results of two-way analysis of variance (ANOVA) are

summarized in Supplemental Table 1.

(F, G) Ectopic xylem differentiation in triple mutants. Bright blue signal in (F)

indicates xylem cell wall stained with BF-170. Xylem differentiation ratio is shown

using box-and-whisker plots in (G).

(H) Expression levels of *IRX3* and *SEOR1* in triple mutants at 48 h after induction.

Relative fold changes were calculated in comparison with the WT at 0 h after

induction (*n* = 3). Different letters indicate significant differences ($P < 0.05$; Tukey-

Kramer test).

Scale bars = 1 cm in (B,F).

Figure 5. BEH3 competes with other BES/BZR homologs

(A) Transcriptional repressor activity of BES/BZR proteins in the *Nicotiana benthamiana* transient expression assay. Luciferase (LUC) activities in *pDWF4:ELUC* transgenic plants were calculated as average photon counts per second of six leaf disks and were normalized relative to values of DEX-treated samples harboring an empty effector construct.

(B) DNA-binding ability of BES/BZR proteins in the electrophoretic mobility shift assay (EMSA). Asterisks and arrowheads indicate the positions of free DNA and DNA-binding domain (DBD)-containing protein-bound DNA, respectively.

Figure 6. BEH3 p ... **s the re** ... **y of other BES/BZR homologs**

(A, B) Development of fruits and seeds in *bes1* quintuple (*bes1-q*) and hextuple (*bes1-h*) mutants. Number of seeds in each mature fruit are shown using box-and-whisker plots in (B). Different letters indicate significant differences (*P* < 0.05; Tukey-Kramer test; *n* = 10).

(C, D) Primary root length in 11-day-old *bes1-q* and *bes1-h* mutant seedlings. To obtain homozygous *bes1-h* mutant plants, self-fertilized progenies of the T2 heterozygous *bes1-q* mutants, which containing *beh3*$^{+/-}$ mutation, were used. *n* = 10 (WT), 12 (*bes1-q*), and 13 (*bes1-h*).

(E) Expression levels of BES/BZR target genes in the estradiol-inducible BEH3 overexpression line. Eight-day-old seedlings were treated with 10 μM β-estradiol for 9h. Relative gene expression levels were calculated in comparison with the DMSO-treated sample (**P* < 0.01; Student's *t*-test; *n* = 3).

(F) Expression level of BES/BZR target genes in *bes1-D bzr1-D*. Nine-day-old seedlings were used. Relative gene expression levels were calculated in comparison with the WT (**P* < 0.01; Student's *t*-test; *n* = 3).

Scale bars = 1 mm (A and C).

Figure 7. Competitive relationship among BES/BZR family provides the

robustness for the control of vascular stem cell activity

(A) Schematic representation of the mathematical model that describes

competitive binding between BEH3 and other homologs. BEH3 binds to i of n

binding sites at a certain binding rate (P_{BEH3}). Then, other BES1 homologs (P_o)

bind to *j* of *n-i* binding sites at a certain binding rate. Signaling output (*i, j*) is

defined as $\alpha^i \beta^j$; α and β represent coefficients of transcriptional activity (<1 in

the case of a repressor) of BEH3 and other BES1 homologs, respectively.

(B, C) Simulated signaling output when verifying both P_{BEH3} and P_o (*n* =3; α =

0.95; β = 0.8). Means and c oefficient of variation (CV) (CV = standard

deviation/mean) of output values are shown in (B) and (C), respectively.

(D) Model showing the role of BES/BZR family members in the robust control of

vascular stem cell differentiation and maintenance. Scale bars = 100 µm (H–J).

Supplemental Figure 1. Comprehensive analysis of the VISUAL coexpression network and vascular-related transcriptome datasets. (Supports Figure 1)

(A) Expression profiles of the VISUAL-induced genes in the FANS dataset (Shi et al., 2020).

(B) Venn diagram of genes in each module of the VISUAL-induced genes and the cambium-enriched genes (Zang et al., 2019).

(C) Expression levels of the cambium-related genes (Module-II) in the FANS dataset (Shi et al., 2020). The gene names are shown on the right side.

Supplemental Figure 2. All section images of hypocotyl vasculature of WT and *beh3-1* mutants. (Supports Figure 2)

All section images for Figure 2F. Sections of hypocotyl vasculature of WT and *beh3-1* mutants grown under short day conditions for 21 days. Scale bars = 100 µm.

Supplemental Figure 3. Sections of hypocotyl vasculature of the β-estradiol inducible *BEH3*-overexpressing plants. (Supports Figure 3)

Plants were grown with or without 5 μM β-estradiol for 12 days. Relative areas inside the cambium were calculated in comparison with mean value of DMSO treated samples Quantified values were shown in box-and-whisker plots. White circles indicate the xylem differentiation ratio for each sample. (*$P < 0.05$ and **$P < 0.01$; Student's *t*-test).

Supplemental Figure 4. Semi-automatic calculation of xylem differentiation

ratio by the binarization method. (Supports Figure 4)

(A) The procedure of image thresholding in the WT and *bes1-1* after the VISUAL

induction. The BF-170 signal (BF signal) area and the leaf area was shown by

white and blue colour, respectively. The number written in the lower right

represents the BF signal area per the leaf area (%) calculated by the binarization

method.

(B) Xylem area calculated based on BF signal per leaf area (%) in the WT and

bes1-1. Statistical differences were examined by Student's t-test (**$P < 0.005$).

Error bars indicate SD (n = 12).

(C) Comparison of two calculation methods for xylem differentiation ratio. X axis

indicates the ratio of xylem area calculated from BF signal per the leaf area. Y

axis indicates the ratio of xylem area calculated from bright field images per the

leaf area. There is a strong positive correlation between them.

Supplemental Figure 5. Ectopic xylem differentiation in VISUAL with *beh2*

and *bes1 beh2* mutants. (Supports Figure 4)

(A) A bright blue signal indicates xylem cell wall stained with BF-170.

(B) Quantified xylem differentiation ratio was shown by box-and-whisker plots.

White circles indicate the xylem differentiation ratio for each sample. Different

letters indicate significant differences (*P* < 0.05; Tukey-Kramer test).

Col-0   *bes1-1 bzr1-2*   *bes1-1 beh3-1*

Supplemental Figure 6. Ectopic xylem cells in VISUAL observed using

confocal laser scanning microscope. (Supports Figure 4)

Cotyledons of WT (Col-0), *bes1-1 bzr1-2*, and *bes1-1 beh3-1* were cultured with

xylem indicator, BF170, for 4 days. A bright blue signal indicates BF170-

incorporated xylem cell walls. Bars, 100 μm.

Supplemental Figure 7. Preparation of MBP-fused proteins of BES/BZR

family DBDs for EMSA assay. (Supports Figure 5)

(A) Coomassie-stained SDS-PAGE analysis of the maltose-binding protein

(MBP)-fused BES1, BZR1, BEH3 and BEH4 DBDs after purification.

(B) Oligomeric state analysis of MBP-fused BES1, BZR1, BEH3 and BEH4 DBDs

by size exclusion chromatography (SEC). The peak positions of the marker

proteins are indicated by the black triangles at the top of the chromatogram.

Supplemental Figure 8. Combinatory effects on transcriptional repressor activity in the *Nicotiana benthamiana* transient expression assay. (Supports Figure 5)

When *BZR1* or *BEH3* was co-expressed with *BES1*, luciferase (LUC) activities in *pDWF4:ELUC* transgenic plants were calculated as average photon counts per second of six leaf disks and were normalized relative to values of DEX-treated samples harbouring an empty effector construct.

1004

1005 **Supplemental Figure 9. Expression levels of *BEH3* in *bes1-1* and *BES1* in**

1006 ***beh3-1* at 72 hours after VISUAL induction.** (Supports Figure 4)

1007 The data are indicated as relative fold change to Col-0 (*P<0.05, **P<0.01;

1008 Student's t-test, n = 3, error bar = SD).

A
BEH1 (At3G50750)
BEH2 (At4G36780)
BEH3 (At4G18890)
100 bp

B *bes1 quadruple* mutant
(*beh2-5 / bes1-1 bzr1-2 beh4-1*)

```
                      15                      47
                      |                       |
BEH2 (WT)   GGAGGAGGAGGAGGAGGATCATC-GTCGGGACG
   beh2-5   GGAGGAGGAGGAGGAGGATCATCCGTCGGGACG
```

bes1 quintuple mutant
(*beh1-2 / bes1-1 bzr1-2 beh2-5 beh4-1*)

```
                      13                    45
                      |                     |
BEH1 (WT)   GAGGAGGATCAACGGCGGCGACG-GGGAGGATG
   beh1-2   GAGGAGGATCAACGGCGGCGACGTGGGAGGATG
```

bes1 hextuple mutant
(*beh3-3 / bes1-1 bzr1-2 beh1-2 beh2-5 beh4-1*)

```
                      54                      85
                      |                       |
BEH3 (WT)   GCGAGAGCGGCGAAGACGAGCGA-TTGCGGCTA
   beh3-3   GCGAGAGCGGCGAAGACGAGCGATTTGCGGCTA
```

1009

Supplemental Figure 10. Construction of genome-edited lines for BES/BZR

homologs. (Supports Figure 6)

(A) Genomic structure of the encoding region of BEH1, BEH2, and BEH3. Red

arrowheads indicate location of target site.

(B) The targeted sequence (orange) and a part of sequence in both wild-type line

and mutant lines are shown. Each genome-edited line has a 1-bp insertion (red).

1016

Supplemental Figure 11. Time-course expression analysis of BES/BZR

target genes in the β-estradiol-inducible *BEH3* overexpression line.

(Supports Figure 6)

Eight-day-old seedlings were treated with 10 µM β-estradiol to induce *BEH3*

expression. Relative gene expression levels were calculated in comparison with

the DMSO-treated sample at 3 h after induction (*P < 0.05, **P < 0.01; two-way

ANOVA; n = 3). Values of statistical analysis listed in Supplemental Table 2.

1024

1025 Supplemental table 1. Results of two-way analysis of variance (ANOVA)

1026 for Figure 4E.

IRX3   Method : Tukey

Factor	Level 1	Level 2	Average 1	Average 2	Difference	SE	Statistic	P value	* : P<0.05 ** : P<0.01
0h	Col	bes1	1.0000	1.0822	0.0822	12.0819	0.0068	1.0000	
	Col	beh3	1.0000	0.9405	0.0595	12.0819	0.0049	1.0000	
	Col	bes1 beh3	1.0000	0.9004	0.0996	12.0819	0.0082	1.0000	
	bes1	beh3	1.0822	0.9405	0.1416	12.0819	0.0117	1.0000	
	bes1	bes1 beh3	1.0822	0.9004	0.1818	12.0819	0.0150	1.0000	
	beh3	bes1 beh3	0.9405	0.9004	0.0402	12.0819	0.0033	1.0000	
24h	Col	bes1	6.7949	6.3790	0.4159	12.0819	0.0344	1.0000	
	Col	beh3	6.7949	8.8834	2.0886	12.0819	0.1729	0.9981	
	Col	bes1 beh3	6.7949	5.7313	1.0636	12.0819	0.0880	0.9997	
	bes1	beh3	6.3790	8.8834	2.5045	12.0819	0.2073	0.9968	
	bes1	bes1 beh3	6.3790	5.7313	0.6477	12.0819	0.0536	0.9999	
	beh3	bes1 beh3	8.8834	5.7313	3.1522	12.0819	0.2609	0.9936	
48h	Col	bes1	135.5941	13.7125	121.8816	12.0819	10.0880	P < 0.001	**
	Col	beh3	135.5941	173.1634	37.5692	12.0819	3.1096	0.0231	*
	Col	bes1 beh3	135.5941	132.2273	3.3668	12.0819	0.2787	0.9922	
	bes1	beh3	13.7125	173.1634	159.4509	12.0819	13.1975	P < 0.001	**
	bes1	bes1 beh3	13.7125	132.2273	118.5148	12.0819	9.8093	P < 0.001	**
	beh3	bes1 beh3	173.1634	132.2273	40.9360	12.0819	3.3882	0.0121	*
72h	Col	bes1	91.5413	50.4735	41.0677	12.0819	3.3991	0.0118	*
	Col	beh3	91.5413	74.2216	17.3197	12.0819	1.4335	0.4916	
	Col	bes1 beh3	91.5413	77.9913	13.5500	12.0819	1.1215	0.6801	
	bes1	beh3	50.4735	74.2216	23.7481	12.0819	1.9656	0.2287	
	bes1	bes1 beh3	50.4735	77.9913	27.5177	12.0819	2.2776	0.1316	
	beh3	bes1 beh3	74.2216	77.9913	3.7697	12.0819	0.3120	0.9892	

SEOR1   Method : Tukey

Factor	Level 1	Level 2	Average 1	Average 2	Difference	SE	Statistic	P value	* : P<0.05 ** : P<0.01
0h	Col	bes1	1.0000	1.3088	0.3088	10.1316	0.0305	1.0000	
	Col	beh3	1.0000	1.0357	0.0357	10.1316	0.0035	1.0000	
	Col	bes1 beh3	1.0000	0.8107	0.1893	10.1316	0.0187	1.0000	
	bes1	beh3	1.3088	1.0357	0.2730	10.1316	0.0269	1.0000	
	bes1	bes1 beh3	1.3088	0.8107	0.4981	10.1316	0.0492	1.0000	
	beh3	bes1 beh3	1.0357	0.8107	0.2251	10.1316	0.0222	1.0000	
24h	Col	bes1	9.3700	4.5883	4.7817	10.1316	0.4720	0.9645	
	Col	beh3	9.3700	14.3676	4.9975	10.1316	0.4933	0.9598	
	Col	bes1 beh3	9.3700	6.5691	2.8009	10.1316	0.2765	0.9924	
	bes1	beh3	4.5883	14.3676	9.7793	10.1316	0.9652	0.7701	
	bes1	bes1 beh3	4.5883	6.5691	1.9808	10.1316	0.1955	0.9973	
	beh3	bes1 beh3	14.3676	6.5691	7.7984	10.1316	0.7697	0.8673	
48h	Col	bes1	101.4912	56.0745	45.4167	10.1316	4.4827	P < 0.001	**
	Col	beh3	101.4912	135.2594	33.7682	10.1316	3.3330	0.0138	*
	Col	bes1 beh3	101.4912	94.9092	6.5820	10.1316	0.6497	0.9146	
	bes1	beh3	56.0745	135.2594	79.1849	10.1316	7.8156	P < 0.001	**
	bes1	bes1 beh3	56.0745	94.9092	38.8347	10.1316	3.8330	0.0042	**
	beh3	bes1 beh3	135.2594	94.9092	40.3502	10.1316	3.9826	0.0029	**
72h	Col	bes1	112.0916	97.0006	15.0910	10.1316	1.4895	0.4591	
	Col	beh3	112.0916	133.7105	21.6189	10.1316	2.1338	0.1712	
	Col	bes1 beh3	112.0916	120.2861	8.1945	10.1316	0.8088	0.8497	
	bes1	beh3	97.0006	133.7105	36.7099	10.1316	3.6233	0.0069	**
	bes1	bes1 beh3	97.0006	120.2861	23.2855	10.1316	2.2983	0.1266	
	beh3	bes1 beh3	133.7105	120.2861	13.4244	10.1316	1.3250	0.5566	

1027

1028

 Supplemental table 2. Results of two-way analysis of variance (ANOVA)

 for Supplemental Figure 11.

BEH3　　　Method : Tukey

Factor	Level 1	Level 2	Average 1	Average 2	Difference	SE	Statistic	P value	* : P<0.05 ** : P<0.01
3h	DMSO	Est	1.0000	2.2876	1.2876	0.9152	1.4068	0.1848	
6h	DMSO	Est	1.0340	5.1796	4.1456	0.9152	4.5295	P < 0.001	**
9h	DMSO	Est	1.2027	7.2724	6.0697	0.9152	6.6318	P < 0.001	**

BR6ox2　　　Method : Tukey

Factor	Level 1	Level 2	Average 1	Average 2	Difference	SE	Statistic	P value	* : P<0.05 ** : P<0.01
3h	DMSO	Est	1.0000	1.2068	0.2068	0.1267	1.6316	0.1287	
6h	DMSO	Est	1.0450	1.5146	0.4696	0.1267	3.7054	0.0030	**
9h	DMSO	Est	1.2062	1.7024	0.4961	0.1267	3.9144	0.0021	**

DWF4　　　Method : Tukey

Factor	Level 1	Level 2	Average 1	Average 2	Difference	SE	Statistic	P value	* : P<0.05 ** : P<0.01
3h	DMSO	Est	1.0000	0.9968	0.0032	0.0937	0.0340	0.9734	
6h	DMSO	Est	1.0486	1.2521	0.2035	0.0937	2.1705	0.0507	
9h	DMSO	Est	1.1264	1.4091	0.2827	0.0937	3.0160	0.0107	*

1032 **Supplemental table 3. Sequences of sgRNA for CRISPR/Cas9 system.**

Oligo name	Sequence
BEH1-CRISPR-F	ATTGGATCAACGGCGGCGACGGGG
BEH1-CRISPR-R	AAACCCCCGTCGCCGCCGTTGATC
BEH2-CRISPR-F	ATTGGGAGGAGGAGGATCATCGTC
BEH2-CRISPR-R	AAACGACGATGATCCTCCTCCTCC
BEH3-CRISPR-F	ATTGGCGGCGAAGACGAGCGATTG
BEH3-CRISPR-R	AAACCAATCGCTCGTCTTCGCCGC

1033

1034 **Supplemental table 4. List of qRT-PCR primers.**

UBQ14	Left primer	TCCGGATCAGCAGAGGTT
	Right primer	TCTGGATGTTGTAGTCAGCAAGA
	Probe	#121
IRX3	Left primer	TGACATGAATGGTGACGTAGC
	Right primer	CATCAAATGCTCCTTATCACCTT
	Probe	#25
SEOR1	Left primer	AAGACACCAACGCCTCCA
	Right primer	CGATAGCATAGGAGACACTATCAAGA
	Probe	#136
ACT2	Left primer	CCGCTCTTTCTTTCCAAGC
	Right primer	CCGGTACCATTGTCACACAC
	Probe	#30
BEH3	Left primer	AATGTGGACTCCTGGACAA
	Right primer	CTGATCAACACCAGCAGGAA
	Probe	#108
DWF4	Left primer	TTCTCGTTATGACCAACCTAATCTC
	Right primer	AGGATGACGCTCCGTTGTT
	Probe	#145
BR6ox2	Left primer	CCCATGGAGATGGATGGA
	Right primer	CTTTCCAGGGCAAAGCCTA
	Probe	#10

1035