An estimate of the deepest branches of the tree of life from ancient vertically-evolving genes

Edmund R. R. Moody[1], Tara A. Mahendrarajah[2], Nina Dombrowski[2], James W. Clark[1], Celine Petitjean[1], Pierre Offre[2], Gergely J. Szöllősi[3,4,5], Anja Spang[2,6*], Tom A. Williams[1*]

1. School of Biological Sciences, University of Bristol, Bristol BS8 1TH, UK.
2. NIOZ, Royal Netherlands Institute for Sea Research, Department of Marine Microbiology and Biogeochemistry; AB Den Burg, The Netherlands
3. Dept. of Biological Physics, Eötvös Loránd University, 1117 Budapest, Hungary
4. MTA-ELTE "Lendület" Evolutionary Genomics Research Group, 1117 Budapest, Hungary;
5. Institute of Evolution, Centre for Ecological Research, 1121 Budapest, Hungary
6. Department of Cell- and Molecular Biology, Science for Life Laboratory, Uppsala University, SE-75123, Uppsala, Sweden

*Co-corresponding authors: tom.a.williams@bristol.ac.uk, anja.spang@nioz.nl

Abstract

Core gene phylogenies provide a window into early evolution, but different gene sets and analytical methods have yielded substantially different views of the tree of life. Trees inferred from a small set of universal core genes have typically supported a long branch separating the archaeal and bacterial domains. By contrast, recent analyses of a broader set of non-ribosomal genes have suggested that Archaea may not be very divergent from Bacteria, and that estimates of inter-domain distance are inflated due to accelerated evolution of ribosomal proteins along the inter-domain branch. Resolving this debate is key to determining the diversity of the archaeal and bacterial domains, the shape of the tree of life, and our understanding of the early course of cellular evolution. Here, we investigate the evolutionary history of the marker genes key to the debate. We show that estimates of a reduced Archaea-Bacteria (AB) branch length result from inter-domain gene transfers and hidden paralogy in the expanded marker gene set, which act to artifactually diminish the genetic distance between the two domains. By contrast, analysis of a broad range of manually curated marker gene datasets from a sample of 700 Archaea and Bacteria reveal that current methods likely underestimate the AB branch length due to substitutional saturation and poor model fit; that the best-performing phylogenetic markers tend to support longer inter-domain branch lengths; and that the AB branch lengths of ribosomal and non-ribosomal marker genes are statistically indistinguishable. A phylogeny of prokaryotes inferred from the 27 highest-ranked marker genes, including ribosomal and non-ribosomal markers, supported a long AB branch, recovered a clade of DPANN at the base of the Archaea, and placed CPR within Bacteria as the sister group to the Chloroflexota.

Introduction

Much remains unknown about the earliest period of cellular evolution and the deepest divergences in the tree of life. Phylogenies encompassing both Archaea and Bacteria have been inferred from a "universal core" set of 16-56 genes encoding proteins involved in translation and other aspects of the genetic information processing machinery(Ciccarelli et al., 2006; Fournier and Gogarten, 2010; Harris et al., 2003; Hug et al., 2016; Mukherjee et al., 2017; Petitjean et al., 2014; Ramulu et al., 2014; Raymann et al., 2015; Theobald, 2010; Williams et al., 2020). These genes are thought to predominantly evolve vertically and are thus best-suited for reconstructing the tree of life (Ciccarelli et al., 2006; Creevey et al., 2011; Ramulu et al., 2014; Theobald, 2010). In these analyses, the branch separating Archaea from Bacteria (hereafter, the AB branch) is often the longest internal branch in the tree(Cox et al., 2008; Gogarten et al., 1989; Hug et al., 2016; Iwabe et al., 1989; Pühler et al., 1989; Williams et al., 2020). In molecular phylogenetics, branch lengths are usually measured in expected numbers of substitutions per site, with a long branch corresponding to a greater degree of genetic change. Long branches can therefore result from high evolutionary rates, long periods of absolute time, or a combination of the two. If a sufficient number of fossils are available to calibrate them then molecular clock models can, in principle, disentangle the contributions of these effects. However, only very few fossil calibrations (Sugitani et al., 2015) are currently available that are old enough to calibrate early divergences (Betts et al., 2018; Horita and Berndt, 1999; Lepland et al., 2002; van Zuilen et al., 2002), and as a result, the ages and evolutionary rates of the deepest branches of the tree, and estimates of the true biodiversity of the archaeal and bacterial domains, remain highly uncertain.

Recently, Zhu et al. (Zhu et al., 2019) inferred a phylogeny from 381 genes distributed across Archaea and Bacteria using the supertree method ASTRAL (Mirarab et al., 2014). In addition to a large increase in the number of genes compared to other universal marker sets, the functional profile of these markers comprises not only proteins involved in information processing but also proteins affiliated with most other functional COG categories, including metabolic processes (Table S1). The genetic distance (branch length) between the domains (Zhu et al., 2019) was estimated from a concatenation of the same marker genes, resulting in a much shorter AB branch length than observed with the core universal markers (Hug et al., 2016; Williams et al., 2020). These analyses were consistent with the hypothesis (Petitjean et al., 2014; Zhu et al., 2019) that the apparent deep divergence of Archaea and Bacteria might be the result of an accelerated evolutionary rate of genes encoding translational and in particular ribosomal proteins along the AB branch as compared to other genes. Interestingly, the same observation was made previously using a smaller set of 38 non-ribosomal marker proteins (Petitjean et al., 2014), although the difference in AB branch length between ribosomal and non-ribosomal markers in that analysis was reported to be substantially lower (roughly two-fold, compared to roughly ten-fold for the 381 protein set (Petitjean et al., 2014; Zhu et al., 2019).

A higher evolutionary rate of ribosomal genes might result from the accumulation of compensatory substitutions at the interaction surfaces among the protein subunits of the ribosome (Petitjean et al., 2014; Valas and Bourne, 2011), or as a compensatory response to the addition or removal of ribosomal subunits early in evolution (Petitjean et al., 2014).

Alternatively, differences in the inferred AB branch length might result from varying rates or
patterns of evolution between the traditional core genes (Spang et al., 2015; Williams et al.,
2020) and the expanded set (Zhu et al., 2019). Substitutional saturation (multiple substitutions
at the same site (Jeffroy et al., 2006)) and across-site compositional heterogeneity can both
impact the inference of tree topologies and branch lengths (Foster, 2004; Lartillot et al., 2007;
Lartillot and Philippe, 2004; Quang et al., 2008; Wang et al., 2008). These difficulties are
particularly significant for ancient divergences (Gouy et al., 2015). Failure to model site-
specific amino acid preferences has previously been shown to lead to under-estimation of the
AB branch length due to a failure to detect convergent changes (Tourasse and Gouy, 1999;
Williams et al., 2020), although the published analysis of the 381 marker set did not find
evidence of a substantial impact of these features on the tree as a whole (Zhu et al., 2019).
Those analyses also identified phylogenetic incongruence among the 381 markers, but did
not determine the underlying cause (Zhu et al., 2019).
100
This recent work (Zhu et al., 2019) raises two important issues regarding the inference of the
universal tree: first, that estimates of the genetic distance between Archaea and Bacteria from
classic "core genes" may not be representative of ancient genomes as a whole, and second,
that there may be many more suitable genes to investigate early evolutionary history than
generally recognized, providing an opportunity to improve the precision and accuracy of deep
phylogenies. Here, we investigate these issues in order to determine why different marker sets
support different Archaea-Bacteria branch lengths. First, we examine the evolutionary history
of the 381 gene marker set (hereafter, the expanded marker gene set) and identify several
features of these genes, including instances of inter-domain gene transfers and mixed
paralogy, that may contribute to the inference of a shorter AB branch length in supertree and
concatenation analyses. Then, we re-evaluate the marker gene sets used in a range of
previous analyses to determine how these and other factors, including substitutional saturation
and model fit, contribute to inter-domain branch length estimations and the shape of the
universal tree. Finally, we identify a subset of marker genes least affected by these issues,
and use these to estimate an updated tree of the primary domains of life and the genetic
distance between Archaea and Bacteria.

Results and Discussion

Gene transfers and hidden paralogy obscure the genetic distance between Archaea and Bacteria

Genes from the expanded marker set are not widely distributed in Archaea

The 381 gene set was derived from a larger set of 400 genes used to estimate the phylogenetic placement of new lineages as part of the PhyloPhlAn method (Segata et al., 2013). Perhaps reflecting the focus on bacteria in the original application, the phylogenetic distribution of the 381 marker genes in the expanded set varies substantially (Table S1), with many being poorly represented in Archaea. Indeed 25% of the published gene trees (https://biocore.github.io/wol/ (Zhu et al., 2019)) contain less than 0.5% archaeal homologues, with 21 (5%) and 69 (18%) of these trees including no or less than 10 archaeal homologues, respectively. For the remaining 75% of the gene trees, archaeal homologs comprise 0.5%-13.4% of the dataset. While there are many more sequenced bacteria than archaea, 63% of the gene trees possessed genes from less than half of the 669 archaeal genomes included in the analysis, whereas only 22% of the gene trees possessed fewer than half of the total number of 9906 sampled bacterial genomes. These distributions suggest that many of these genes are not broadly present in both domains, and that some might be specific to Bacteria.

Conflicting evolutionary histories of individual marker genes and the inferred species tree

In the focal analysis of the 381 gene set, the tree topology was inferred using the supertree method ASTRAL (Mirarab et al., 2014), with branch lengths inferred on this fixed tree from a marker gene concatenation (Zhu et al., 2019). The topology inferred from this expanded marker set (Zhu et al., 2019) is similar to published trees (Castelle and Banfield, 2018; Hug et al., 2016) and recovers Archaea and Bacteria as reciprocally monophyletic domains, albeit with a shorter AB branch than in earlier analyses. However, the individual gene trees (Zhu et al., 2019) disagree regarding domain monophyly: Archaea and Bacteria are recovered as reciprocally monophyletic groups in only 24 of the 381 published (Zhu et al., 2019) maximum likelihood (ML) gene trees of the expanded marker set (Table S1).

Since single gene trees often fail to strongly resolve ancient relationships, we used approximately-unbiased (AU) tests (Shimodaira, 2002) to evaluate whether the failure to recover domain monophyly in the published ML trees is statistically supported. For computational tractability, we performed these analyses on a 1000-species subsample of the full 10,575-species dataset that was compiled in the original study (Zhu et al., 2019). For 79 of the 381 genes, we could not perform the test because the gene was not found on any of the 74 archaeal genomes present in the 1000-species subsample. For the remaining 302 genes, domain monophyly was rejected ($p < 0.05$) for 232 out of 302 (76.8%) genes. As a comparison, we performed the same test on several smaller marker sets used previously to infer a tree of life (Coleman et al., 2021; Petitjean et al., 2014; Williams et al., 2020); none of

the markers in those sets rejected reciprocal domain monophyly ($p > 0.05$ for all genes, Figure
1(a)). In what follows, we refer to four published marker gene sets as: the Expanded set (381
genes (Zhu et al., 2019)), the Core set (49 genes (Williams et al., 2020), encoding ribosomal
proteins and other conserved information-processing functions; itself a consensus set of
several earlier studies (Da Cunha et al., 2017; Spang et al., 2015; Williams et al., 2012)), the
Non-ribosomal set (38 genes, broadly distributed and explicitly selected to avoid genes
encoding ribosomal proteins (Petitjean et al., 2014)), and the Bacterial set (29 genes used in
a recent analysis of bacterial phylogeny (Coleman et al., 2021)).

To investigate why 232 of the marker genes rejected the reciprocal monophyly of Archaea and
Bacteria, we returned to the full dataset (Zhu et al., 2019), annotated each sequence in each
marker gene family by assigning proteins to KOs, Pfams, and Interpro domains, among others
(Table S1, see Methods for details) and manually inspected the tree topologies (Table S1).
This revealed that the major cause of domain polyphyly observed in gene trees was inter-
domain gene transfer (in 357 out of 381 gene trees (93.7%)) and mixing of sequences from
distinct paralogous families (in 246 out of 381 gene trees (64.6%)). For instance, marker
genes encoding ABC-type transporters (p0131, p0151, p0159, p0174, p0181, p0287, p0306,
po0364), tRNA synthetases   (i.e. p0000, p0011, p0020, p0091, p0094, p0202),
aminotransferases and dehydratases (i.e. p0073/4-aminobutyrate aminotransferase;
p0093/3-isopropylmalate dehydratase) often comprised a mixture of paralogues.

Together, these analyses indicate that the evolutionary histories of the individual markers of
the expanded set differ from each other and from the species tree. Zhu et al. acknowledged
(Zhu et al., 2019) the varying levels of congruence between the marker phylogenies and the
species tree, but did not investigate the underlying causes. Our analyses establish the basis
for these disagreements in terms of gene transfers and the mixing of orthologues and
paralogues within and between domains. Concatenation is based on the assumption that all
of the genes in the supermatrix evolve on the same underlying tree; genes with different gene
tree topologies violate this assumption and should not be concatenated because the
topological differences among sites are not modelled, and so the impact on inferred branch
lengths is difficult to predict. In practice, it is often difficult to be certain that all of the markers
in a concatenate share the same gene tree topology, and the analysis proceeds on the
hypothesis that a small proportion of discordant genes are not expected to seriously impact
the inferred tree. However, the concatenated tree inferred from the expanded marker set
differs from previous trees in that the genetic distance between Bacteria and Archaea is greatly
reduced, such that the AB branch length appears comparable to distances among bacterial
phyla (Zhu et al., 2019). Because an accurate estimate of the AB branch length has a major
bearing on unanswered questions regarding the root of the universal tree (Gouy et al., 2015),
we next evaluated the impact of the conflicting gene histories within the expanded marker set
on inferred AB branch length.

The inferred branch length between Archaea and Bacteria is artifactually shortened by
inter-domain gene transfer and hidden paralogy

To investigate the impact of gene transfers and mixed paralogy on the AB branch length
inferred by gene concatenations (Zhu et al., 2019), we compared branch lengths estimated
from markers that rejected (AU < 0.05) or did not reject (AU > 0.05) the reciprocal monophyly
of Bacteria and Archaea in the 381 marker set (Figure 1(a)). To estimate AB branch lengths

for genes in which the domains were not monophyletic in the ML tree, we first performed a
constrained ML search to find the best gene tree that was consistent with domain monophyly
for each family under the LG+G4+F model in IQ-TREE 2 (Minh et al., 2020). While it may
seem strained to estimate the length of a branch that does not appear in the ML tree, we
reasoned that this approach would provide insight into the contribution of these genes to the
AB branch length in the concatenation, in which they conflict with the overall topology. AB
branch lengths were significantly ($P = 2.159 \times 10^{-12}$, Wilcoxon rank sum test) shorter for markers
that rejected domain monophyly (Figure 1(a); <0.05: mean AB branch length in expected
substitutions/site 0.0130, >0.05: mean AB branch length 0.559). This result suggests that
inter-domain gene transfers reduce the AB branch length when included in a concatenation.
This behaviour might result from marker gene transfers reducing the number of fixed
differences between the domains, so that the AB branch length in a tree in which Archaea and
Bacteria are constrained to be reciprocally monophyletic will tend towards 0 as the number of
transfers increases. Consistent with this hypothesis, we observed that ∆LL, the difference in
log likelihood between the constrained ML tree and ML gene tree (used here as a proxy for
gene verticality), correlates negatively with AB branch length (Figure 1(b)). Furthermore, AB
branch length decreased as increasing numbers of low-verticality markers were added to the
concatenate (Figure 1(c)). Taken together, these results indicate that the inclusion of genes
that do not support the reciprocal monophyly of Archaea and Bacteria in the universal
concatenate reduces the estimated AB branch length by homogenizing the genetic diversity
of the two domains.

Figure 1: Expanded set genes in which Archaea and Bacteria are not monophyletic support a shorter AB branch. *(a) Expanded set genes that reject domain monophyly (p < 0.05, AU test) support significantly shorter AB branch lengths when constrained to follow a domain monophyletic tree (p = 2.159×10-12, Wilcoxon rank-sum test). None of the marker genes from several other published analyses reject domain monophyly (p > 0.05, AU test) for all genes tested. (b) Marker gene verticality (ΔLL, see below) for the expanded gene set normalized by alignment length correlates negatively with the length of the AB branch between Archaea and Bacteria (R2=0.03998, p = 0.0004731). (c) Concatenations of 20-100 markers of the expanded set markers ranked by marker gene verticality (ΔLL) show the same trend, with a reduction in AB branch length as markers with a greater ΔLL are added to the concatenate. ΔLL is the difference between the log likelihood of the ML gene family tree under a free topology search and the log likelihood of the best tree constrained to obey domain monophyly. The trendline is estimated using LOESS regression.*

The age of the last universal common ancestor (LUCA) inferred from strict clocks
does not predict marker gene quality

Reliable age estimates using molecular clock methods require calibrations, but few calibrations exist for the deeper branches of the tree of life. Zhu et al. (Zhu et al., 2019) argued that the expanded marker set is useful for deep phylogeny because estimates of the age of the last universal common ancestor (LUCA) obtained by fitting molecular clocks to their dataset are in agreement with the geological record: a root (LUCA) age of 3.6-4.2 Ga was inferred from the entire 381-gene dataset, consistent with the earliest fossil evidence for life (Betts et al., 2018; Sugitani et al., 2015), whereas estimates from ribosomal markers alone supported a root age of 7 Ga. This age might be considered implausible because it is much older than the age of the Earth and Solar System (with the moon-forming impact occurring ~4.51 Ga (Barboni et al., 2017; Hanan and Tilton, 1987)). However, the palaeobiological plausibility of the age estimate from the 381 gene set does not, in itself, constitute evidence of marker gene suitability. In the original analyses, the age of LUCA was estimated using a maximum likelihood approach, as well as a Bayesian molecular clock with a strict clock (assuming a constant evolutionary rate) or a relaxed clock with a single calibration. A strict clock model does not permit changes in evolutionary rate through time or across branches, and so a longer AB branch will lead to an older inferred LUCA age. Likewise, a relaxed clock model with a single calibration may fail to distinguish molecular distances and geological time. Given that the short AB branch in the expanded gene set results, in part, from phylogenetic incongruence among markers, we evaluated the age of LUCA inferred from the subset of the expanded gene set least affected by these issues. To do so, we analysed the top 5% of gene families according to their ΔLL score (a set of 20 genes, which includes only 1 ribosomal protein) under the same clock model parameters as the original dataset (Figure 2). This analysis resulted in a significantly more ancient age estimate for LUCA (5.5-6.5 Ga), and trimming the alignment to remove poorly-aligning regions resulted in a still older estimate (6.34-6.89 Ga), approaching that of the ribosomal genes (7.46-8.03 Ga). These analyses suggest that, for these data and calibrations, the inferred age for LUCA is not a reliable indicator of marker quality, because analyses using the subset of the data least affected by incongruence more clearly reveals the underlying limitations of strict clock analyses (and indeed relaxed clocks with few calibrations) for dating ancient divergences. In principle, more reliable estimates of LUCA's age might be obtained by using more calibrations. However, unambiguous calibrations remain elusive, particularly for the root and other deep branches of the tree. Despite advances in molecular clock methodology, such calibrations represent the only way to reliably capture the relationship between genetic distance and divergence time.

Figure 2. The inferred age of LUCA is not a reliable indicator of marker quality. *Posterior node age estimates from Bayesian molecular clock analyses of 1) Conserved sites as estimated previously (Zhu et al., 2019); 2) Random sites (Zhu et al., 2019) 3) Ribosomal genes (Zhu et al., 2019) 4) The top 5% of marker gene families according to their ΔLL score (including only 1 ribosomal protein) and 5) The same top 5% of marker genes trimmed using BMGE(Criscuolo and Gribaldo, 2010) to remove highly variable sites. In each case, a strict molecular clock was applied, with the age of the Cyanobacteria-Melainabacteria split constrained between 2.5 and 2.6 Ga.*

Phylogeny of Archaea and Bacteria using ancient vertically-evolving genes

Finding ancient vertically-evolving genes

To estimate the AB branch length and the phylogeny of prokaryotes using a dataset that resolves some of the issues identified above, we performed a meta-analysis of several previous studies to identify a consensus set of vertically-evolving marker genes. We identified unique markers from these analyses by reference to the COG ontology (Dombrowski et al., 2020; Galperin et al., 2019), extracted homologous sequences from a representative sample of 350 archaeal and 350 bacterial genomes, and performed iterative phylogenetics and manual curation to obtain a set of 54 markers that recovered archaeal and bacterial monophyly (see Methods). Subsequently, we ranked these 54 genes by the extent to which they recovered established within-domain relationships using the split score, a criterion described previously (Dombrowski et al., 2020) (see Methods) yielding a final set of 27 markers that were used for inferring an updated universal species tree (see below). Marker genes that better resolved relationships within each domain also supported a longer AB branch length (Figure 3).

Distributions of AB branch lengths for ribosomal and non-ribosomal marker genes are similar

Traditional universal marker sets include many ribosomal proteins (Ciccarelli et al., 2006; Fournier and Gogarten, 2010; Harris et al., 2003; Hug et al., 2016; Williams et al., 2020). If ribosomal proteins experienced accelerated evolution during the divergence of Archaea and Bacteria, this might lead to the inference of an artifactually long AB branch length (Petitjean et al., 2014; Zhu et al., 2019)**.** To investigate this, we plotted the inter-domain branch lengths for the 38 and 16 ribosomal and non-ribosomal genes, respectively, comprising the 54 marker genes set. We found no evidence that there was a longer AB branch associated with ribosomal markers (Figure 4; mean AB branch length for ribosomal proteins 1.35, mean for non-ribosomal 2.25). Prior to manual curation, non-ribosomal markers had a greater number of HGTs and cases of mixed paralogy. In particular, for the original set of 95 markers, 62% of the non-ribosomal markers and 21% of the ribosomal markers were not monophyletic, respectively. These values were 69% and 29% for the 54 markers, and 50% and 33% for the 27 markers. These results imply that manual curation of marker genes is important for deep phylogenetic analyses, particularly when using non-ribosomal markers.

Figure 3. Better phylogenetic markers have longer AB branches. *The plot shows the relationship between split score (a lower split score denotes better recovery of established within-domain relationships, see Methods) and AB branch length (in expected number of substitutions/site) for the 54 highest-ranked marker genes. Marker genes with higher split scores (that split monophyletic groups into multiple subclades) have shorter AB branch lengths (P = 0.0311, r= 0.294). Split scores of ribosomal and non-ribosomal markers were statistically indistinguishable (P = 0.828, Figure S3).*

Figure 4. Among vertically-evolving marker genes, ribosomal genes do not have a longer AB branch length. *The plot shows functional classification of markers against AB branch length using the 54 most vertically evolving markers. We did not see a significant (P = 0.619, Wilcoxon rank sum test) difference between AB branch lengths for ribosomal and non-ribosomal genes.*

Substitutional saturation and poor model fit contribute to underestimation of AB branch
length

For the top 50% of marker genes as determined by split scores (27 genes), we performed an
additional round of single gene tree inference and manual review to identify and remove
remaining sequences which had evidence of HGT or represented distant paralogs. The
resulting single gene trees are provided in the Data Supplement
(10.6084/m9.figshare.13395470). To evaluate the relationship between site evolutionary rate
and AB branch length, we created two concatenations: fastest sites (comprising sites with
highest probability of being in the fastest Gamma rate category; 868 sites) and slowest sites
(sites with highest probability of being in the slowest Gamma rate category, 1604 sites) and
compared relative branch lengths inferred from the entire concatenate using IQ-TREE 2 to
infer site-specific rates (Figure 5). As expected, total tree length is shorter from the slow-
evolving sites, but the relative AB branch length is longer (1.2 substitutions/site, or ~2% of
total tree length, compared to 2.6 substitutions/site, or ~0.04% total tree length for the fastest-
evolving sites). This result suggests that, at fast-evolving sites, some changes along the AB
branch have been overwritten by later events in evolution --- that is, that substitutional
saturation leads to an underestimate of the AB branch length.

Another factor that has been shown to lead to underestimation of genetic distance on deep
branches is a failure to adequately model the site-specific features of sequence evolution
(Lartillot and Philippe, 2004; Schrempf et al., 2020; Wang et al., 2018; Williams et al., 2020).
Amino acid preferences vary across the sites of a sequence alignment, due to variation in the
underlying functional constraints (Lartillot and Philippe, 2004; Quang et al., 2008; Wang et al.,
2008). The consequence is that, at many alignment sites, only a subset of the twenty possible
amino acids are tolerated by selection. Standard substitution models, such as LG+G4+F, are
site-homogeneous, and approximate the composition of all sites using the average
composition across the entire alignment. Such models underestimate the rate of evolution at
highly constrained sites because they do not account for the high number of multiple
substitutions that occur at such sites. The effect is that site-homogeneous models
underestimate branch lengths when fit to site-heterogeneous data. Site-heterogeneous
models have been developed that account for site-specific amino acid preferences, and these
generally show improved fit to real protein sequence data (reviewed in (Williams et al., 2021)).
To evaluate the impact of substitution model fit for these data, we fit a range of models to the
full concatenation, assessing model fit using the Bayesian information criterion (BIC) in IQ-
TREE 2. The AB branch length inferred under the best-fit model, the site-heterogeneous
LG+C60+G4+F model, was 2.52 substitutions/site, ~1.7-fold greater than the branch length
inferred from the site-homogeneous LG+G4+F model (1.45 substitutions/site). Thus,
substitution model fit has a major effect on the estimated length of the AB branch, with better-
fitting models supporting a longer branch length (Table 1). The same trends are evident when
better-fitting site-heterogeneous models are used to analyse the dataset of Zhu et al.:
considering only the top 5% of genes by ΔLL score, the AB branch length is 1.2 under
LG+G4+F, but increases to 2.4 under the best-fitting LG+C60+G4+F model (Figure S2).

Figure 5. Slow- and fast-evolving sites support different shapes for the universal tree. *(i) Tree of Archaea and Bacteria inferred from a concatenation of 27 core genes; (ii) Tree inferred from the fastest-evolving sites; (iii) Tree inferred from the slowest-evolving sites. To facilitate comparison of relative diversity, scale bars are provided separately for each panel. Slow-evolving sites support a relatively long inter-domain branch and less diversity within the domains (that is, shorter between-taxa branch lengths within domains). This suggests that substitution saturation (overwriting of earlier changes) may reduce the relative length of the AB branch at fast-evolving sites and genes.*

Substitution model	BIC	AB branch length
LG+G4+F	5935950.053	1.449090256
LG+C20+G4+F	5783903.997	2.139350118
LG+C40+G4+F	5756823.360	2.469702112
LG+C60+G4+F	5746886.292	2.517828771

350 ***Table 1.* The inferred AB length from a concatenation of the top 27 markers using a**
351 **simple model versus models which account for site compositional heterogeneity.** Using
352 better fitting models, i.e models which allow for across-site compositional heterogeneity, a
353 longer AB branch is inferred.

354
355 ***A phylogeny of Archaea and Bacteria inferred from 27 vertically-evolving marker genes***

356
The topology of our phylogeny of the primary domains of life (Figure 6) is consistent with recent
single-domain trees inferred for Archaea and Bacteria independently (Coleman et al., 2021;
Dombrowski et al., 2020; Williams et al., 2017), although the deep relationships within Bacteria
are only poorly resolved, with the exception of the monophyly of Gracilicutes (Figure 6). A
recent analysis suggested that, among extant lineages, the metabolisms of Clostridia,
Deltaproteobacteria, Actinobacteria and some Aquificae might best preserve the metabolism
of the last bacterial common ancestor (Xavier et al., 2021). Assuming a universal root between
Archaea and Bacteria (Dagan et al., 2010; Gogarten et al., 1989; Iwabe et al., 1989), none of
these groups branch near the bacterial root in our analysis (Figure 6). This is consistent with
previous work (Castelle and Banfield, 2018; Hug et al., 2016; Parks et al., 2017; Raymann et
al., 2015) including the inference of an updated and rooted bacteria phylogeny (Coleman et
al., 2021). Notably, our analysis placed the Candidate Radiation (CPR) (Brown et al., 2015)
as a sister lineage to Chloroflexi (Chloroflexota) rather than as a deep-branching bacterial
superphylum. While this contrasts with initial trees suggesting that CPR may represent an
early diverging sister lineage of all other Bacteria (Brown et al., 2015; Castelle and Banfield,
2018; Hug et al., 2016), our finding is consistent with a recent analysis that recovered CPR
within the Terrabacteria (Coleman et al., 2021). Together, these analyses suggest that the
deep-branching position of CPR in some trees was a result of long branch attraction, a
possibility that has been raised previously (Hug et al., 2016; Méheust et al., 2019).

376
The deep branches of the archaeal subtree are well-resolved in the ML tree and recover clades
of DPANN (albeit at 51% bootstrap support), Asgard (100% bootstrap support), and TACK
Archaea (75% bootstrap support), in agreement with a range of previous studies (Dombrowski
et al., 2020; Guy and Ettema, 2011; Raymann et al., 2015; Williams et al., 2017). We also find
support for the placement of Methanonatronarchaeia (Sorokin et al., 2017) distant to
Halobacteria within the Methanotecta, in agreement with recent analyses and suggesting their
initial placement with Halobacteria (Sorokin et al., 2017) may be an artifact of compositional
attraction (Aouad et al., 2019; Dombrowski et al., 2020; Martijn et al., 2020). Notably, the
Hadesarchaea (92% bootstrap support) and a clade comprising Theionarchaea,
Methanofastidiosa, and Thermococcales (92% bootstrap support) branch basal to the TACK

and Asgard Archaea, respectively, in our analysis, rather than with other Euryarchaeota.
These positions have been previously reported (Adam et al., 2017; Raymann et al., 2015;
Williams et al., 2017), though the extent of euryarchaeotal paraphyly and the lineages involved
has varied among analyses.
391
A broader observation from our analysis is that the phylogenetic diversity of the archaeal and
bacterial domains, measured as substitutions per site in this consensus set of vertically-
evolving marker genes, appears to be similar (Figure 5(i); the mean root to tip distance for
archaea: 2.38, for bacteria: 2.41, the range of root to tip distances for archaea: 1.79-3.01, for
bacteria: 1.70-3.17). Considering only the slowest-evolving category of sites, branch lengths
within Archaea are actually longer than within Bacteria (Figure 5(iii)). This result differs from
some published trees (Hug et al., 2016; Zhu et al., 2019) in which the phylogenetic diversity
of Bacteria has appeared to be significantly greater than that of Archaea. By contrast to those
earlier studies, we analysed a set of 350 genomes from each domain, an approach which may
tend to reduce the differences between them. While we had to significantly downsample the
sequenced diversity of Bacteria, our sampling nonetheless included representatives from all
known major lineages of both domains, and so might be expected to recover a difference in
diversity, if present. Our analyses and a number of previous studies (Hug et al., 2016; Parks
et al., 2018; Petitjean et al., 2014; Zhu et al., 2019) indicate that the choice of marker genes
has a profound impact on the apparent phylogenetic diversity of prokaryotic groups; for
instance, in the proportion of bacterial diversity composed of CPR (Hug et al., 2016; Parks et
al., 2017). Our results demonstrate that slow and fast-evolving sites from the same set of
marker genes support different tree shapes and branch lengths; it therefore seems possible
that between-dataset differences are due, at least in part, to evolutionary rate variation within
and between marker genes.

Figure 6: A phylogeny of Archaea and Bacteria inferred from a concatenation of 27 marker genes. *Consistent with some recent studies (Dombrowski et al., 2020; Guy and Ettema, 2011; Raymann et al., 2015; Williams et al., 2017), we recovered the DPANN, TACK and Asgard Archaea as monophyletic groups. Although the deep branches within Bacteria are poorly resolved, we recovered a sister group relationship between CPR and Chloroflexota, consistent with a recent report (Coleman et al., 2021). The tree was inferred using the best-fitting LG+C60+G4+F model in IQ-TREE 2 (Minh et al., 2020). Branch lengths are proportional to the expected number of substitutions per site. Support values are ultrafast (UFBoot2) bootstraps (Hoang et al., 2018). Numbers in parenthesis refer to number of taxa within each collapsed clade. Please note that collapsed taxa in the Archaea and Bacteria roughly correspond to order- and phylum-level lineages, respectively.*

Conclusion

Core gene phylogenies provide a window into the earliest period of archaeal and bacterial evolution. Concatenation is useful for pooling signal across individual genes, but topology and branch length estimates from concatenations only reflect the underlying tree of life if the individual genes share the same evolutionary history. Our analysis of published datasets (Coleman et al., 2021; Petitjean et al., 2014; Williams et al., 2020; Zhu et al., 2019) indicates that incongruence among marker genes resulting from inter-domain gene transfer and hidden paralogy can lead to an under-estimate of the inter-domain branch length. We performed a re-analysis of marker genes from a range of published analyses, manually curated datasets to identify and remove transferred genes, and estimated an updated phylogeny of Archaea and Bacteria. Considering only this manually curated consensus marker gene dataset, we found no evidence that ribosomal markers overestimate stem length; since they appear to be transferred less frequently than other genes, our analysis affirms that ribosomal proteins are useful markers for deep phylogeny. In general, better markers, regardless of functional category, support a longer AB branch length. A phylogeny inferred from the 27 best-performing markers was consistent with some recent work on early prokaryotic evolution, resolving the major clades within Archaea and nesting the CPR within Terrabacteria. Our analyses suggest that both the true Archaea-Bacteria branch length (Figure 7), and the phylogenetic diversity of Archaea, may be underestimated by even the best current models, a finding that is consistent with a root for the tree of life between the two prokaryotic domains.

Figure 7. The impact of marker gene choice, phylogenetic congruence, alignment trimming, and substitution model fit on estimates of the Archaea-Bacteria branch length. *Analysis using a site-homogeneous model (LG+G4+F) on the complete 381-gene expanded set results in an AB branch substantially shorter than previous estimates. Removing the genes most seriously affected by inter-domain gene transfer, trimming poorly-aligned sites using BMGE (Criscuolo and Gribaldo, 2010), and using the best-fitting site-heterogeneous model available (LG+C60+G4+F) substantially increase the estimated AB length, such that it is comparable with published estimates from the "core" set (Williams et al., 2020) and the consensus set of 27 markers identified in the present study. Branch lengths measured in expected number of substitutions/site.*

Methods

Data

We downloaded the individual alignments from (Zhu et al., 2019) (https://github.com/biocore/wol/tree/master/data/), along with the genome metadata and the individual newick files. We checked each published tree for domain monophyly, and also performed approximately unbiased (AU) (Shimodaira, 2002) tests to assess support for domain monophyly on the underlying sequence alignments using IQ-TREE 2 (Minh et al., 2020). The phylogenetic analyses were carried out using the 'reduced' subset of 1000 taxa outlined by the authors (Zhu et al., 2019), for computational tractability. These markers were also trimmed according to the protocol in the original paper (Zhu et al., 2019), i.e sites with >90% gaps were removed, followed by removal of sequences with >66% gaps.

We also downloaded the Williams et al. (Williams et al., 2020) ("core"), Petitjean et al. (Petitjean et al., 2014) ("non-ribosomal") and Coleman et al. (Coleman et al., 2021) ("bacterial") datasets from their original publications.

Annotations

Proteins used for phylogenetic analyses by Zhu *et al.* (Zhu et al., 2019), were annotated to investigate the selection of sequences comprising each of the marker gene families. To this end, we downloaded the protein sequences provided by the authors from the following repository: https://github.com/biocore/wol/tree/master/data/alignments/genes. To obtain reliable annotations, we analysed all sequences per gene family using several published databases, including the arCOGs (version from 2014) (Seemann, 2014), KOs from the KEGG Automatic Annotation Server (KAAS; downloaded April 2019) (Aramaki et al., 2020), the Pfam database (Release 31.0)(Bateman et al., 2004), the TIGRFAM database (Release 15.0) (Haft et al., 2003), the Carbohydrate-Active enZymes (CAZy) database (downloaded from dbCAN2 in September 2019)(Cantarel et al., 2009), the MEROPs database (Release 12.0) (Rawlings et al., 2016), (Saier et al., 2006), the hydrogenase database (HydDB; downloaded in November 2018) (Søndergaard et al., 2016), the NCBI- non-redundant (nr) database (downloaded in November 2018), and the NCBI COGs database (version from 2020). Additionally, all proteins were scanned for protein domains using InterProScan (v5.29-68.0; settings: --iprlookup --goterms) (Jones et al., 2014).

Individual database searches were conducted as follows: arCOGs were assigned using PSI-BLAST v2.7.1+ (settings: -evalue 1e-4 -show_gis -outfmt 6 -max_target_seqs 1000 -dbsize 100000000 -comp_based_stats F -seg no) (Altschul et al., 1997). KOs (settings: -E 1e-5), PFAMs (settings: -E 1e-10), TIGRFAMs (settings: -E 1e-20) and CAZymes (settings: -E 1e-20) were identified in all archaeal genomes using hmmsearch v3.1b2(Finn et al., 2011). The MEROPsand HydDB databases were searched using BLASTp v2.7.1 (settings: -outfmt 6, -evalue 1e-20). Protein sequences were searched against the NCBI_nr database using DIAMOND v0.9.22.123 (settings: –more-sensitive –e-value 1e-5 –seq 100 –no-self-hits –taxonmap prot.accession2taxid.gz) (Buchfink et al., 2015). For all database searches the best hit for each protein was selected based on the highest e-value and bitscore and all results are summarized in the Data Supplement Table, Annotation_Tables/0_Annotation_tables_full/All_Zhu_marker_annotations_16-12-

2020.tsv.zip. For InterProScan we report multiple hits corresponding to the individual domains
of a protein using a custom script (parse_IPRdomains_vs2_GO_2.py).

Assigned sequence annotations were summarized and all distinct KOs and Pfams were
collected and counted for each marker gene. KOs and Pfams with their corresponding
descriptions were mapped to the marker gene file downloaded from the repository:
https://github.com/biocore/wol/blob/master/data/markers/metadata.xlsx and used in
summarization of the 381 marker gene protein trees (Table S1).

For manual inspection of single marker gene trees, KO and Pfam annotations were mapped
to the tips of the published marker protein trees, downloaded from the repository:
https://github.com/biocore/wol/tree/master/data/trees/genes. Briefly, the Genome ID, Pfam,
Pfam description, KO, KO description, and NCBI Taxonomy string were collected from each
marker gene annotation table and were used to generate mapping files unique to each marker
gene phylogeny, which links the Genome ID to the annotation information
(GenomeID|Domain|Pfam|Pfam Description|KO|KO Description). An in-house perl script
replace_tree_names.pl
(https://github.com/ndombrowski/Phylogeny_tutorial/tree/main/Input_files/5_required_Scripts
) was used to append the summarized protein annotations to the corresponding tips in each
marker gene tree. Annotated marker gene phylogenies were manually inspected using the
following criteria including: 1) retention of reciprocal domain monophyly (Archaea and
Bacteria) and 2) for the presence or absence of potential paralogous families. Paralogous
groups and misannotated families present in the gene trees were highlighted and violations of
search criteria were recorded in Table S1.

Phylogenetic analyses

COG assignment for the Core, Non-Ribosomal, and Bacterial marker genes

First, all gene sequences in the three published marker sets (core, non-ribosomal, and
bacterial) were annotated using the NCBI COGs database (version from 2020). Sequences
were assigned a COG family using hmmsearch v3.3.2 (Finn et al., 2011) (settings: -E 1e-5)
and the best hit for each protein sequence was selected based on the highest e-value and bit
score. To assign the appropriate COG family for each marker gene, we quantified the
percentage distribution of all unique COGs per gene, and selected the family representing the
majority of sequences in each marker gene.

Accounting for overlap, this resulted in 95 unique COG families from the original 119 total
marker genes across all three published datasets (Table S2). Orthologues corresponding to
these 95 COG families were identified in the 700 genomes (350 Archaea, 350 Bacteria, Table
S3) using hmmsearch v3.3.2 (settings: -E 1e-5). The reported BinID and protein accession
were used to extract the sequences from the 700 genomes, which were used for subsequent
phylogenetic analyses.

Marker gene inspection and analysis

We aligned these 95 sequence sets using MAFFT-linsi (Katoh and Toh, 2008) and removed
poorly-aligned positions with BMGE (Criscuolo and Gribaldo, 2010). We inferred initial
maximum likelihood trees (LG+G4+F) for all 95 markers and mapped the KO and Pfam

domains and descriptions, inferred from annotation of the 700 genomes, to the corresponding
tips (see above). Manual inspection took into consideration monophyly of Archaea and
Bacteria and the presence of paralogs, and other signs of contamination (HGT, LBA).
Accordingly, single gene trees that failed to meet reciprocal domain monophyly were excluded,
and any instances of HGT, paralogous sequences, and LBA artefacts were manually removed
from the remaining trees resulting in 54 markers across the three published datasets that were
subject to subsequent phylogenetic analysis (LG+C20+G4+F) and further refinement (see
below).

528
Ranking markers based on split score

We applied an automated marker gene ranking procedure devised previously, the split score
(Dombrowski et al., 2020), to rank each of the 54 markers that satisfied reciprocal monophyly
based on the extent to which they recovered established phylum-, class- or, order-level
relationships within the archaeal and bacterial domains (Table S4).

The script quantifies the number of splits, or occurrences where a taxon fails to cluster within
its expected taxonomic lineage, across all gene phylogenies. Monophyly of archaeal and
bacterial lineages was assessed based on clades defined in Table S4. Briefly, we used
Cluster1 for Archaea in combination with Cluster0 (phylum) or Cluster3 (i.e. on class-level if
defined and otherwise on phylum-level; Table S4) for Bacteria. We then ranked the marker
genes using the following split-score criteria: the number of splits per taxon and the splits
normalized to the species count. The percentage of split phylogenetic groups was used to
determine the highest ranking (top 50%) markers.

Concatenation

Based on the split score ranking of the 54 marker genes (above), the top 50% (27 markers,
Table S4) marker genes were manually inspected using criteria as defined above, and
contaminating sequences were manually removed from the individual sequence files.
Following inspection, marker protein sequences were aligned using MAFFT-LINSI (Katoh and
Standley, 2013) and trimmed using BMGE (Criscuolo and Gribaldo, 2010). We concatenated
the 27 markers into a supermatrix, which was used to infer a maximum-likelihood tree (Figure
6, under LG+C60+G4+F), evolutionary rates (see below), and rate-category supermatrices
as well as to perform model performance tests (see below).

Constraint analysis

We performed a maximum likelihood free topology search using IQ-TREE 2 (Minh et al., 2020)
under the LG+G4+F model, with 1000 bootstrap replicates on each of the markers from the
expanded, bacterial, core and non-ribosomal sets. We also performed a constrained analysis
with the same model, in order to find the maximum likelihood tree in which Archaea and
Bacteria were reciprocally monophyletic. We then compared both trees using the
approximately unbiased (AU) Shimodaira (2002) test in IQ-TREE 2 (Minh et al., 2020) with
10,000 RELL (Shimodaira, 2002) bootstrap replicates. To evaluate the relationship between
marker gene verticality and AB branch length, we calculated the difference in log-likelihood
between the constrained and unconstrained trees in order to rank the genes from the
expanded marker set, made concatenates comprised of the top 20-100 (intervals of 5) of these

marker genes, and inferred the tree length under LG+C10+G4+F with 1000 bootstrap
replicates.

Site and gene evolutionary rates

We inferred rates using the --rate option in IQ-TREE 2 (Minh et al., 2020) for both the 381
marker concatenation from Zhu (Zhu et al., 2019) and the top 5% of marker genes based on
the results of difference in log-likelihood between the constrained tree and free-tree search in
the constraint analysis (above). We also used this method to explore the differences in rates
for the 27 marker set. We built concatenates for sites in the slowest and fastest rate categories,
and inferred branch lengths from each of these concatenates using the tree inferred from the
corresponding dataset as a fixed topology.

Substitution model fit

Model fit tests were undertaken using the top 5% concatenate described above, with the
alignment being trimmed with BMGE 1.12 (Criscuolo and Gribaldo, 2010) with default settings
(BLOSUM62, entropy 0.5) for all of the analyses except the 'untrimmed' LG+G4+F run, other
models   on   the   trimmed   alignment   were   LG+G4+F,   LG+R4+F   and
LG+C10,20,30,40,50,60+G4+F, with 1000 bootstrap replicates. Model fitting was done using
ModelFinder (Kalyaanamoorthy et al., 2017) in IQ-TREE 2 (Minh et al., 2020). For the model
testing for the 27 concatenation, we performed a model finder analysis (-m MFP) including
additional   complex   models   of   evolution,   (i.e.
LG+C60+G4+F,LG+C50+G4+F,LG+C40+G4+F,LG+C30+G4+F,LG+C20+G4+F,LG+C10+G
4+F,LG+G4+F,LG+R4+F) to the default, to find the best fitting model for the analysis. This
revealed that, according to AIC,BIC and cAIC, LG+C60+G4+F was the best fitting model. For
comparison,   we   also   performed   analyses   using   the   following   models:
LG+G4+F,LG+C20+G4+F,LG+C40+G4+F (Table 1).

Molecular clock analyses

Molecular clock analyses were devised to test the effect of genetic distance on the inferred
age of LUCA. Following the approach of Zhu et al (Zhu et al., 2019), we subsampled the
alignment to 100 species. Five alternative alignments were analysed, representing conserved
sites across the entire alignment, randomly selected sites across the entire alignment, only
ribosomal marker genes, the top 5% of marker genes according to ΔLL and the top 5% of
marker genes further trimmed under default settings in BMGE 1.12 (Criscuolo and Gribaldo,
2010). Divergence time analyses were performed in MCMCTree (Yang, 2007) under a strict
clock model. We used the normal approximation approach, with branch lengths estimated in
codeml under the LG+G4 model. In each case, a fixed tree topology was used alongside a
single calibration on the Cyanobacteria-Melainabacteria split. The calibration was modelled
as a uniform prior distribution between 2.5 and 2.6 Ga, with a 2.5% probability that either
bound could be exceeded. For each alignment, four independent MCMC chains were run for
2,000,000 generations to achieve convergence.

Plotting

601   Statistical analyses were performed using R 4.0.4 (R Core Team, 2021), and data were plotted
602   with ggplot2 (Wickham, 2016).

Data and code availability

All of the data, including sequence alignments, trees, annotation files, and scripts associated with this manuscript have been deposited in the FigShare repository at DOI: 10.6084/m9.figshare.13395470.

Acknowledgements

ERRM was supported by a Royal Society Enhancement Award (RGF\EA\180199) to TAW. CP was supported by NERC grant NE/P00251X/1 to TAW. This work was funded by the Gordon and Betty Moore Foundation through grant GBMF9741 to TAW, AS and GJSz. TAW was supported by a Royal Society University Research Fellowship (URF\R\201024). GJSz received funding from the European Research Council under the European Union's Horizon 2020 research and innovation program under Grant Agreement 714774 and Grant GINOP-2.3.2.-15-2016- 00057. AS is supported by the Swedish Research Council (VR starting grant 2016-03559), the NWO-I foundation of the Netherlands Organisation for Scientific Research (WISE fellowship) and the European Research Council (ERC Starting grant 947317, ASymbEL).

References

Adam PS, Borrel G, Brochier-Armanet C, Gribaldo S. 2017. The growing tree of Archaea: new perspectives on their diversity, evolution and ecology. *ISME J* **11**:2407–2425.

Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research* **25**:3389–3402.

Aouad M, Borrel G, Brochier-Armanet C, Gribaldo S. 2019. Evolutionary placement of Methanonatronarchaeia. *Nature Microbiology* **4**:558-559.

Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, Ogata H. 2020. KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive score threshold. *Bioinformatics* **36**:2251–2252.

Barboni M, Boehnke P, Keller B, Kohl IE, Schoene B, Young ED, McKeegan KD. 2017. Early formation of the Moon 4.51 billion years ago. *Science Advances* **3**:e1602365.

Bateman A, Coin L, Durbin R, Finn RD, Hollich V, Griffiths-Jones S, Khanna A, Marshall M, Moxon S, Sonnhammer ELL, Studholme DJ, Yeats C, Eddy SR. 2004. The Pfam protein families database. *Nucleic Acids Research* **32**:D138–41.

Betts HC, Puttick MN, Clark JW, Williams TA, Donoghue PCJ, Pisani D. 2018. Integrated genomic and fossil evidence illuminates life's early evolution and eukaryote origin. *Nature Ecology and Evolution* **2**:1556–1562.

Brown CT, Hug LA, Thomas BC, Sharon I, Castelle CJ, Singh A, Wilkins MJ, Wrighton KC, Williams KH, Banfield JF. 2015. Unusual biology across a group comprising more than 15% of domain Bacteria. *Nature* **523**:208–211.

Buchfink B, Xie C, Huson DH. 2015. Fast and sensitive protein alignment using DIAMOND. *Nat Methods* **12**:59–60.

Cantarel BL, Coutinho PM, Rancurel C, Bernard T, Lombard V, Henrissat B. 2009. The Carbohydrate-Active EnZymes database (CAZy): an expert resource for Glycogenomics. *Nucleic Acids Research* **37**:D233–8.

Castelle CJ, Banfield JF. 2018. Major New Microbial Groups Expand Diversity and Alter our Understanding of the Tree of Life. *Cell* **172**:1181–1197.

Ciccarelli FD, Doerks T, von Mering C, Creevey CJ, Snel B, Bork P. 2006. Toward automatic reconstruction of a highly resolved tree of life. *Science* **311**:1283–1287.

Coleman GA, Davín AA, Mahendrarajah T, Spang A, Hugenholtz P, Szöllősi GJ, Williams TA. 2021. A rooted phylogeny resolves early bacterial evolution. *Science* **372**. doi:10.1126/science.abe5011

Cox CJ, Foster PG, Hirt RP, Harris SR, Embley TM. 2008. The archaebacterial origin of eukaryotes. *Proceedings of the National Academy of Sciences of the United States of America* **105**:20356–20361.

Creevey CJ, Doerks T, Fitzpatrick DA, Raes J, Bork P. 2011. Universally distributed single-copy genes indicate a constant rate of horizontal transfer. *PLoS One* **6**:e22099.

Criscuolo A, Gribaldo S. 2010. BMGE (Block Mapping and Gathering with Entropy): a new software for selection of phylogenetic informative regions from multiple sequence alignments. *BMC Evolutionary Biology* **10**:210.

Da Cunha V, Gaia M, Gadelle D, Nasir A, Forterre P. 2017. Lokiarchaea are close relatives of Euryarchaeota, not bridging the gap between prokaryotes and eukaryotes. *PLoS Genetics* **13**:e1006810.

Dagan T, Roettger M, Bryant D, Martin W. 2010. Genome networks root the tree of life between prokaryotic domains. *Genome Biology and Evolution* **2**:379–392.

Dombrowski N, Williams TA, Sun J, Woodcroft BJ, Lee J-H, Minh BQ, Rinke C, Spang A.

2020. Undinarchaeota illuminate DPANN phylogeny and the impact of gene transfer on archaeal evolution. *Nature Communications* **11**:1–15.

Finn RD, Clements J, Eddy SR. 2011. HMMER web server: interactive sequence similarity searching. *Nucleic Acids Research* **39**:W29–W37.

Foster PG. 2004. Modeling compositional heterogeneity. *Systematic Biology* **53**:485–495.

Fournier GP, Gogarten JP. 2010. Rooting the ribosomal tree of life. *Molecular Biology and Evolution* **27**:1792–1801.

Galperin MY, Kristensen DM, Makarova KS, Wolf YI, Koonin EV. 2019. Microbial genome analysis: the COG approach. *Briefings in Bioinformatics* **20**:1063–1070.

Gogarten JP, Kibak H, Dittrich P, Taiz L, Bowman EJ, Bowman BJ, Manolson MF, Poole RJ, Date T, Oshima T, Konishi J, Denda K, Yoshida M. 1989. Evolution of the vacuolar H+-ATPase: implications for the origin of eukaryotes. *Proceedings of the National Academy of Sciences of the United States of America* **86**:6661–6665.

Gouy R, Baurain D, Philippe H. 2015. Rooting the tree of life: the phylogenetic jury is still out. *Philosophical Transactions of the Royal Society B Biological Sciences* **370**:20140329.

Guy L, Ettema TJG. 2011. The archaeal "TACK" superphylum and the origin of eukaryotes. *Trends in Microbiology* **19**:580–587.

Haft DH, Selengut JD, White O. 2003. The TIGRFAMs database of protein families. *Nucleic Acids Research* **31**:371–373.

Hanan BB, Tilton GR. 1987. 60025: relict of primitive lunar crust? *Earth and Planetary Science Letters* **84**:15–21.

Harris JK, Kelley ST, Spiegelman GB, Pace NR. 2003. The genetic core of the universal ancestor. *Genome Research* **13**:407–412.

Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: Improving the Ultrafast Bootstrap Approximation. *Molecular Biology and Evolution* **35**:518–522.

Horita J, Berndt ME. 1999. Abiogenic methane formation and isotopic fractionation under hydrothermal conditions. *Science* **285**:1055–1057.

Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ, Butterfield CN, Hernsdorf AW, Amano Y, Ise K, Suzuki Y, Dudek N, Relman DA, Finstad KM, Amundson R, Thomas BC, Banfield JF. 2016. A new view of the tree of life. *Nature Microbiology* **1**:16048 doi:10.1038/nmicrobiol.2016.48

Iwabe N, Kuma K, Hasegawa M, Osawa S, Miyata T. 1989. Evolutionary relationship of archaebacteria, eubacteria, and eukaryotes inferred from phylogenetic trees of duplicated genes. *Proceedings of the National Academy of Sciences of the United States of America* **86**:9355–9359.

Jeffroy O, Brinkmann H, Delsuc F, Philippe H. 2006. Phylogenomics: the beginning of incongruence? *Trends in Genetics* **22**:225–231.

Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell A, Nuka G, Pesseat S, Quinn AF, Sangrador-Vegas A, Scheremetjew M, Yong S-Y, Lopez R, Hunter S. 2014. InterProScan 5: genome-scale protein function classification. *Bioinformatics* **30**:1236–1240.

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods* **14**:587–589.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* **30**:772–780.

Katoh K, Toh H. 2008. Recent developments in the MAFFT multiple sequence alignment program. *Briefings in Bioinformatics*. **9**(4):286-298 doi:10.1093/bib/bbn013

Lartillot N, Brinkmann H, Philippe H. 2007. Suppression of long-branch attraction artefacts in the animal phylogeny using a site-heterogeneous model. *BMC Evolutionary Biology* **7**:S4.

Lartillot N, Philippe H. 2004. A Bayesian mixture model for across-site heterogeneities in the amino-acid replacement process. *Molecular Biology and Evolution* **21**:1095–1109.

Lepland A, Arrhenius G, Cornell D. 2002. Apatite in early Archean Isua supracrustal rocks, southern West Greenland: its origin, association with graphite and potential as a

biomarker. *Precambrian Research* **118**:221–241.

Martijn J, Schön ME, Lind AE, Vosseberg J, Williams TA, Spang A, Ettema TJG. 2020.
Hikarchaeia demonstrate an intermediate stage in the methanogen-to-halophile
transition. *Nature Communications* **11**:5490.

Méheust R, Burstein D, Castelle CJ, Banfield JF. 2019. The distinction of CPR bacteria from
other bacteria based on protein family content. *Nature Communications* **10**:4173.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear
R. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in
the Genomic Era. *Molecular Biology and Evolution* **37**:1530–1534.

Mirarab S, Reaz R, Bayzid MS, Zimmermann T, Swenson MS, Warnow T. 2014. ASTRAL:
genome-scale coalescent-based species tree estimation. *Bioinformatics* **30**:i541–8.

Mukherjee S, Seshadri R, Varghese NJ, Eloe-Fadrosh EA, Meier-Kolthoff JP, Göker M,
Coates RC, Hadjithomas M, Pavlopoulos GA, Paez-Espino D, Yoshikuni Y, Visel A,
Whitman WB, Garrity GM, Eisen JA, Hugenholtz P, Pati A, Ivanova NN, Woyke T, Klenk
H-P, Kyrpides NC. 2017. 1,003 reference genomes of bacterial and archaeal isolates
expand coverage of the tree of life. *Nature Biotechnology* **35**:676–683.

Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil P-A, Hugenholtz
P. 2018. A standardized bacterial taxonomy based on genome phylogeny substantially
revises the tree of life. *Nature Biotechnology* **36**: 996-1004. doi:10.1038/nbt.4229

Parks DH, Rinke C, Chuvochina M, Chaumeil P-A, Woodcroft BJ, Evans PN, Hugenholtz P,
Tyson GW. 2017. Recovery of nearly 8,000 metagenome-assembled genomes
substantially expands the tree of life. *Nature Microbiology* **2**:1533–1542.

Petitjean C, Deschamps P, López-García P, Moreira D. 2014. Rooting the domain archaea
by phylogenomic analysis supports the foundation of the new kingdom Proteoarchaeota.
Genome Biology and Evolution **7**:191–204.

Pühler G, Leffers H, Gropp F, Palm P, Klenk HP, Lottspeich F, Garrett RA, Zillig W. 1989.
Archaebacterial DNA-dependent RNA polymerases testify to the evolution of the
eukaryotic nuclear genome. *Proceedings of the National Academy of Sciences of the
United States of America* **86**:4569–4573.

Quang LS, Gascuel O, Lartillot N. 2008. Empirical profile mixture models for phylogenetic
reconstruction. *Bioinformatics* **24**:2317–2323.

Ramulu HG, Groussin M, Talla E, Planel R, Daubin V, Brochier-Armanet C. 2014. Ribosomal
proteins: toward a next generation standard for prokaryotic systematics? *Molecular
Phylogenetics and Evolution* **75**:103–117.

Rawlings ND, Barrett AJ, Finn R. 2016. Twenty years of the MEROPS database of
proteolytic enzymes, their substrates and inhibitors. *Nucleic Acids Research* **44**:D343–
50.

Raymann K, Brochier-Armanet C, Gribaldo S. 2015. The two-domain tree of life is linked to a
new root for the Archaea. *Proceedings of the National Academy of Sciences* **112**:6670–
6675.

R Core Team. 2021. R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria.

Saier MH Jr, Tran CV, Barabote RD. 2006. TCDB: the Transporter Classification Database
for membrane transport protein analyses and information. *Nucleic Acids Research*
34:D181–6.

Schrempf D, Lartillot N, Szöllösi G. 2020. Scalable empirical mixture models that account for
across-site compositional heterogeneity. *Mol Biol Evol*. doi:10.1093/molbev/msaa145

Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**:2068–
2069.

Segata N, Börnigen D, Morgan XC, Huttenhower C. 2013. PhyloPhlAn is a new method for
improved phylogenetic and taxonomic placement of microbes. *Nature Communications*
4:2304.

Shimodaira H. 2002. An approximately unbiased test of phylogenetic tree selection.
Systematic Biology **51**:492–508.

Søndergaard D, Pedersen CNS, Greening C. 2016. HydDB: A web tool for hydrogenase

classification and analysis. *Scientific Reports* **6**:34212.

Sorokin DY, Makarova KS, Abbas B, Ferrer M, Golyshin PN, Galinski EA, Ciordia S, Mena MC, Merkel AY, Wolf YI, van Loosdrecht MCM, Koonin EV. 2017. Discovery of extremely halophilic, methyl-reducing euryarchaea provides insights into the evolutionary origin of methanogenesis. *Nature Microbiology* **2**:17081.

Spang A, Saw JH, Jørgensen SL, Zaremba-Niedzwiedzka K, Martijn J, Lind AE, van Eijk R, Schleper C, Guy L, Ettema TJG. 2015. Complex archaea that bridge the gap between prokaryotes and eukaryotes. *Nature* **521**:173–179.

Sugitani K, Mimura K, Takeuchi M, Lepot K, Ito S. 2015. Early evolution of large micro-organisms with cytological complexity revealed by microanalyses of 3.4 Ga organic-walled microfossils. *Geobiology* **13**:507–521.

Theobald DL. 2010. A formal test of the theory of universal common ancestry. *Nature* **465**:219–222.

Tourasse NJ, Gouy M. 1999. Accounting for Evolutionary Rate Variation among Sequence Sites Consistently Changes Universal Phylogenies Deduced from rRNA and Protein-Coding Genes. *Molecular Phylogenetics and Evolution* **13**:159–168.

Valas RE, Bourne PE. 2011. The origin of a derived superkingdom: how a gram-positive bacterium crossed the desert to become an archaeon. *Biology Direct* **6**:16.

van Zuilen MA, Lepland A, Arrhenius G. 2002. Reassessing the evidence for the earliest traces of life. *Nature* **418**:627–630.

Wang H-C, Li K, Susko E, Roger AJ. 2008. A class frequency mixture model that adjusts for site-specific amino acid frequencies and improves inference of protein phylogeny. *BMC Evolutionary Biology* **8**:331.

Wang H-C, Minh BQ, Susko E, Roger AJ. 2018. Modeling Site Heterogeneity with Posterior Mean Site Frequency Profiles Accelerates Accurate Phylogenomic Estimation. *Systematic Biology* **67**:216–235.

Wickham H. 2016. ggplot2: Elegant graphics for data analysis. Springer-Verlag New York.

Williams T a., Foster PG, Nye TMW, Cox CJ, Embley TM. 2012. A congruent phylogenomic signal places eukaryotes within the Archaea. *Philosophical Transactions of the Royal Society B Biological Sciences* **279**:4870–4879.

Williams TA, Cox CJ, Foster PG, Szöllősi GJ, Embley TM. 2020. Phylogenomics provides robust support for a two-domains tree of life. *Nature Ecology and Evolution* **4**:138–147.

Williams TA, Schrempf D, Szöllősi GJ, Cox CJ, Foster PG, Embley TM. 2021. Inferring the deep past from molecular data. *Genome Biology and Evolution*. doi:10.1093/gbe/evab067

Williams TA, Szöllősi GJ, Spang A, Foster PG, Heaps SE, Boussau B, Ettema TJG, Embley TM. 2017. Integrative modeling of gene and genome evolution roots the archaeal tree of life. *Proceedings of the National Academy of Sciences of the United States of America* **114**:E4602–E4611.

Xavier JC, Gerhards RE, Wimmer JLE, Brueckner J, Tria FDK, Martin WF. 2021. The metabolic network of the last bacterial common ancestor. *Communications Biology* **4**: 413. doi:10.1038/s42003-021-01918-4

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Molecular Biology and Evolution* **24**:1586–1591.

Zhu Q, Mai U, Pfeiffer W, Janssen S, Asnicar F, Sanders JG, Belda-Ferre P, Al-Ghalith GA, Kopylova E, McDonald D, Kosciolek T, Yin JB, Huang S, Salam N, Jiao J-Y, Wu Z, Xu ZZ, Cantrell K, Yang Y, Sayyari E, Rabiee M, Morton JT, Podell S, Knights D, Li W-J, Huttenhower C, Segata N, Smarr L, Mirarab S, Knight R. 2019. Phylogenomics of 10,575 genomes reveals evolutionary proximity between domains Bacteria and Archaea. *Nature Communications* **10**. doi:10.1038/s41467-019-13443-4