Identification and characterization of a G-quadruplex structure in the pre-core promoter region of hepatitis B virus

Vanessa Meier-Stephenson[1,2,4#], Maulik D Badmalia[1#], Tyler Mrozowich[1], Keith CK Lau[2], Sarah K Schultz[1], Darren L Gemmill[1], Carla Osiowy[3], Guido van Marle[2], Carla S Coffin,[2,4*] and Trushar R Patel[1,2,5*]

[1] Alberta RNA Research and Training Institute, Department of Chemistry and Biochemistry, University of Lethbridge, Lethbridge, Alberta, T1K 3M4, Canada
[2] Department of Microbiology, Immunology and Infectious Diseases, Cumming School of Medicine, University of Calgary, Alberta, T2N 4N1, Canada
[3] Viral Hepatitis and Bloodborne Pathogens, National Microbiology Laboratory, Public Health Agency of Canada, Winnipeg, MB, R3E 3R2, Canada
[4] Department of Medicine, Cumming School of Medicine, Calgary, AB, Canada
[5] DiscoveryLab, Faculty of Medicine & Dentistry, University of Alberta, Edmonton, Alberta, Canada

Equal author contribution
* To whom correspondence should be addressed - Carla S Coffin (cscoffin@ucalgary.ca) and Trushar R Patel (trushar.patel@uleth.ca)

Running Head: A quadruplex motif in key HBV promoter region

Keywords: Hepatitis B virus, pre-core promoter, cccDNA, DNA G-quadruplex, HBV surface antigen, ELISA, microscale thermophoresis, small-angle X-ray scattering

ABSTRACT

Worldwide, ~250 million people are chronically infected with the hepatitis B virus (HBV) and are at increased risk of cirrhosis and hepatocellular carcinoma. The HBV persists as covalently closed circular DNA (cccDNA), which acts as the template for all HBV mRNA transcripts. Nucleos(t)ide analogs do not directly target the HBV cccDNA and cannot eradicate the HBV. We have discovered a unique structural motif, a G-quadruplex in HBV's pre-core promoter region that is conserved amongst nearly all genotypes, and is central to critical steps in the viral life-cycle including the production of pre-genomic RNA, core and polymerase proteins, and encapsidation. Thus, an increased understanding of the HBV pre-core may lead to the development of novel anti-HBV cccDNA targets. We utilized biophysical methods to characterize the presence of the G-quadruplex, employed assays using a known quadruplex-binding protein (DHX36) to pull-down HBV cccDNA, and compared HBV infection in HepG2 cells transfected with wild-type and mutant HBV plasmids. This study provides insights into the presence of a G-quadruplex in the HBV pre-core promoter region essential for HBV replication. The evaluation of this critical host-protein interaction site in the HBV cccDNA may ultimately facilitate the development of novel anti-HBV therapeutics against the resilient cccDNA template.

INTRODUCTION

Approximately 250 million people worldwide are chronic hepatitis B virus (HBV) carriers and are at elevated risk of developing cirrhosis, liver failure, and hepatocellular carcinoma (HCC) (1-3). The virus persists, in part, due to the presence of its stable, compact intranuclear minichromosome— the covalently closed circular DNA (cccDNA) (4-7). Current oral therapies against HBV, including nucleotide/side analogues (NA) target the virus during replication and are effective at decreasing viremia, but rarely induce HBV surface antigen loss, i.e., "functional cure" (8-11). Moreover, NAs do not directly target the cccDNA and can lead to viral relapse once treatment is stopped (10-13). Achieving a sterilizing (or virological) HBV cure requires targeting and clearance of the HBV cccDNA episome. There is limited knowledge of the secondary and tertiary structural intricacies of the HBV genome. The discovery of unique HBV cccDNA structural motifs may highlight new areas for therapeutic drug design and intervention.

HBV is a small, compact 3.2 kb, partially double-stranded DNA virus (14). The HBV genome consists of four overlapping open-reading frames for genes surface (S), core (C), polymerase (P), and X, encoding seven viral proteins (14). Transcriptional regulation of these regions is guided by the corresponding promoter regions that enable the binding of various cellular (and viral) proteins to initiate transcription and produce five RNA transcripts, including the pre-C and pre-genomic RNAs (pgRNA) (each approximately 3.5 kb), and the sub-genomic RNAs S1 (2.4 kb), S2 (2.1 kb) and X (0.7 kb) (14). The pre-C / pre-genomic region is one of the most important viral promoter regions. This region initiates transcription of the mRNA template for the production of both the core protein and polymerase and produces the full-length pre-genomic RNA which contains the 5′-hairpin loop or epsilon (ε) (14-16). The epsilon region enables binding of the viral polymerase and initiation of encapsidation or packaging of the viral

template followed by replication and the creation of viral progeny (14,15). Overall, HBV is

classified into eight main genotypes, A-H, based on a >8% nucleotide variation across the

genome (17,18). The genotypes tend to display different propensities to cause clinical disease

and to respond to therapy (19,20). Despite the number of unique genotypes and viral variants due

to the HBV's error-prone reverse transcriptase, promoter regions tend to be highly conserved

amongst all HBV variants / genotypes (21). This is likely owing to their need to interact with key

host proteins for successful propagation.

G-quadruplexes are highly stable non-canonical DNA (or RNA) structures formed in

guanosine-rich nucleic acid regions, where four guanines form a planar quartet (G-quartet)

stabilized by the presence of Hoogsteen-hydrogen bonds (22). Three or more consecutive G-

quartets can stack on top of each other to form parallel, anti-parallel, or hybrid G-quadruplex

structures (23). The G-quadruplexes are further stabilized by monovalent cations that occupy the

central channel between the G-quartets (22). The typical sequence that can produce a quadruplex

is represented by the formula, $-G_3-N_{1-7}-G_3-N_{1-7}-G_3-N_{1-7}-G_3-$, where groups of three guanosines

are separated by one to seven nucleotides, but there are exceptions to this rule (24), such as the

specific quadruplex highlighted in the current study. Over 700,000 putative quadruplex

sequences have been found throughout the human genome, having a predominance in the

telomeric and gene regulatory regions, which has led to the discovery of their roles in key

transcription, translation, and genomic regulatory processes (25-31). In the field of cancer

research, such findings have facilitated the development of a new class of drugs (*i.e.*, G-

quadruplex stabilizers), including the recent telomeric quadruplex stabilizers, CX-3545

(Quarfloxin®) and CX-5461, currently in early Phase clinical trials (32-35). Based on advances

in oncology, quadruplexes have subsequently been found in several viral genomes, including

human immunodeficiency virus-1 (HIV-1) (36), herpes simplex virus (HSV-1) (37,38), hepatitis

C virus (HCV) (39), human papillomavirus (HPV) (40), Epstein-Barr virus (EBV) (41), Kaposi's

sarcoma-related herpesvirus (KSHV) (42), and Zika virus (43); reviewed in (44,45). More

recently, a G-quadruplex has also been found in the promoter region of the HBV's S-gene (46).

The predominance of G-quadruplexes in key regulatory regions is highlighted, suggesting

interesting targets for anti-viral therapy development (44,45).

In the current work, we demonstrate the central, multi-functioning pre-core promoter

region of the HBV genome as a G-quadruplex forming region, confirm its presence through

biophysical methods, and demonstrate its role in viral replication when compared to non-G-

quadruplex-forming mutants *in vitro*. Collectively, this lays the basis for the study of a critical

host-protein interaction and the potential development of unique therapeutic targets for HBV

cccDNA.

RESULTS

The HBV pre-core promoter region contains a highly conserved G-rich sequence.

In our prior analysis of the HBV promoter region using the HBV genome database

(HBVdB: https://hbvdb.ibcp.fr), we noted a highly G-rich region in the pre-core/core promoter

region in all HBV genomes except the HBV G genotype (21). This region would be overlooked

if the rigid formula were used when searching for putative quadruplex sequences, G_{3-5} N_{1-7} G_{3-5}

N_{1-7} G_{3-5} N_{1-7} G_{3-5}, (i.e., where G's are guanosines present in groups of three to five, and N's are

any nucleotide present in one to seven sequences separating these groups of guanosines), but

exceptions exist (47). This region was of significant interest given the fact that it also binds to

host Specificity protein 1 (Sp1), a host transcription factor already known to bind G-

quadruplexes, further supporting the link with the secondary structure in this region. The analysis of single nucleotide mutation frequencies in this region was computed similarly to our previous study (21), but with a focus on all available basal core promoter sequences in the HBV genome database (9939 sequences, Figure 1A), demonstrating remarkable conservation of the guanosine groups.

A

CTGGGAGGAGCTGGGGGGAGGA
CTGGGAGGAGCTGGGGGGAGGA
CTGGGAGGAGCTGGGGGGAGGA
CTGGGAGGAGCTGGGGGAAGGA
CTGGGAGGAGCTGGGGGGAGGA

G	A	G	T	T	G	G	G	G
0.00%	0.50%	0.30%	89.40%	96.80%	0.50%	0.01%	0.02%	0.10%
99.90%	0.10%	98.80%	0.10%	0.10%	97.40%	100.00%	99.90%	99.80%
0.01%	0.01%	0.02%	10.40%	0.01%	0.01%	0.02%	0.00%	0.01%
0.10%	99.40%	0.90%	3.10%	2.00%	0.02%	0.02%	0.10%	0.10%
			0.05%					

Figure 1. (A) Frequency analysis of the guanosines present in the pre-core promoter region of HBV using the HBV genome database https://hbvdb.ibcp.fr, (accessed April, 2019). HBV Genotype G does not contain this G-rich pre-core region and hence was not included in the analysis. (B) Wild-type (*wt*) and mutants (*G1738A*, *G1748A* and *G1738/1748A*) oligomers used in this study.

The pre-core/core promoter region of HBV forms a quadruplex structure.

To determine the ability of the HBV pre-core/core promoter region to form a quadruplex, we utilized a 23-mer oligomer of the wild-type (*wt*) promoter region for use in multiple

biophysical assays (Figure1B). We also designed a mutant (*G1748A*) oligomer based on prior

studies showing loss of Sp1 binding with the single-nucleotide mutation of G to A substitution at

position 1748 (48). Additional mutants were also included (*i.e.*, *G1738A* and *G1738/1748A*) to

determine the nature and necessity of these putative G-quadruplex-disrupting mutations. HBV

genome frequency analysis using the HBV database (www.hbvdb.ibcp.fr) showed >99.8%

conservation across all genotypes hence supporting the use of these mutants for our proposed

studies (Figure 1). Oligomers were solubilized in an appropriate buffer, followed by a heat-

cooled step to allow for G-quadruplex formation. Samples were purified using SEC prior to

performing all experiments to ensure they were free of aggregation and showed a single peak

eluted for subsequent collection and concentration (Figure 2A). First, we performed circular

dichroism spectropolarimetry (CD) experiments to investigate whether the wildtype (*wt*) and

mutant oligomers form a G-quadruplex structure in solution. We observed a peak in ellipticity

upon CD analysis at $\lambda \approx 263$ nm and a minimum at $\lambda \approx 242$ nm (as detailed in Table S1, Figure

2B), suggesting that the *wt* oligomer adopts a parallel G-quadruplex structure in solution (49-51).

Similarly, the *G1738A* mutant showed a comparable quadruplex CD profile. The spectra for the

G1748A and *G1738/1748A* mutants, however, displayed a less prominent pattern despite the

same concentration, consistent with a partially folded quadruplex structure (Figure 2B). We also

studied the G quadruplex formation in the presence of Li^+. We purified G-quadruplexes under

identical conditions except that the buffers contained Li^+ instead of K^+. The CD spectra were

collected for all the samples, and no considerable changes were observed in CD spectra collected

for G-quadruplexes in Li^+ and K^+, suggesting that the DNA sequence can form G-quadruplex

despite possessing two mutations irrespective of the centrally coordinated monovalent cation.

Therefore, we decided to collect MST and SAXS data for G quadruplexes in the presence of K^+

only.

Figure 2. Purification and primary biophysical studies of the pre-core promoter *wt*, *G1738A*,

G1748A and *G1738/1748A* mutant oligomers. (A) The change in size exclusion chromatography

elution profile suggests change in size of G4s (B) Circular dichroism spectroscopy studies of the

pre-core promoter oligomers. The presence of a negative peak at ~242 nm and a positive peak at

~263 nm indicates that the G4s are present at a parallel quadruplex in the solution, the respective

values are tabulated in Table S1. The plots represent the average of the three independent runs

for each.

To visualize and verify the G-quadruplex formation, we employed the HPLC-SAXS

(52,53). This enabled selection of the dataset from a monodispersed region for further analysis.

The absence of upturned profile at lower momentum transfer (*q*) values in Guinier analysis of

SAXS data for *wt* and mutant oligomers implies that all samples are monodispersed and

aggregation free (Figure 3A, panel i and ii). Next, the dimensionless Kratky analysis (54) of

SAXS data for all samples was performed to assess their flexibility and compactness. This

analysis displays a Gaussian curve suggesting that all samples were folded (Figure 3A, panel iii).

The SAXS data were then converted to the electron pair-distance distribution plots (($P(r)$)

function) (Figure 3A, panel iv) using the GNOM software program(55). The characteristic

Gaussian-shaped pattern observed in $P(r)$ for the *wt* sample indicates that it adopts a compact

structure in solution, whereas the *G1738A, G1748A,* and *G1738/1748A* mutant oligomers,

despite the same nucleotide length (23 nucleotides), has an increasingly extended structure in

solution (Figure 3A, panel iv). The radius of gyration (R_g) for *wt* was calculated to be 18.37 Å,

while the *G1738A, G1748A,* and *G1738/1748A* mutants yields increasing values of 19.10, 20.10

and 20.34 Å, respectively, which are in agreement with those obtained from the Guinier analysis

(Table 1). The $P(r)$ function also allows the determination of a maximum particle dimension

(D_{max}) of macromolecules. Based on this analysis, we obtained the D_{max} of 46.85 Å for the *wt*,

while the *G1738A*, *G1748A,* and *G1738/1748A* mutants were 53.14 Å, 60.00 Å and 60.44 Å

respectively, indicating that each mutation leads to a degree of alteration of the G-quadruplex

structure (Figure 3). To further investigate the structural differences between the samples, we

used the $P(r)$ data in the DAMMIF program that allows low-resolution structure determination.

We calculated ten independent low-resolution structures for each sample that provided *X* values

ranging from 1.2 to 1.8 indicating the good quality of our models (Table 1). Subsequently, we

averaged and filtered ten models using DAMAVER program to obtain a representative low-

resolution structure for each sample. We obtained the normalized spatial discrepancy (NSD)

values of 0.55, 0.58, 0.58 and 0.59 for *wt, G1738A*, *G1748A,* and *G1738/1748A* mutant samples,

respectively, indicating that the ten independent low-resolution structures are highly similar to

each other in all cases. The low-resolution structures presented in Figure 3B demonstrate the

compact quadruplex structure of the *wt* sample, while each of the mutations displays a relatively

extended structure in solution. It is noteworthy that the increase in the size of mutants is also

consistent with observed elution volumes from SEC where we observed that the *wt* displayed the

highest elution volume (Figure 2A) which progressively increased in the same order as the

increase in D_{max} i.e. *wt < G1738A < G1748A < G1738/1748A*.

To further confirm that the *wt* oligomer forms a G-quadruplex structure, we used the

previously described DHX36 protein, (a known G-quadruplex-interacting protein that binds with

high affinity to parallel quadruplexes) (53,56-58), and tested for DHX36 ability to recognize the

wt and/or mutant oligomers via MST. Based on the MST results (Figure 4), we obtained a K_d of

76.81 ± 12.21 nM between DHX36 and the *wt* oligomer further confirming that the *wt* oligomer

adopts a G-quadruplex structure in solution that is recognised by DHX36. The K_d values

obtained for the mutant oligomers were 14.17 ± 12.76, 132.14 ± 14.84, and 65.50 ± 7.74 nM for

G1738A, *G1748A* and *G1738/1748A*, respectively.

Figure 3. Low-resolution structural studies of HBV pre-core G-quadruplex (wt, pink) and its

mutant oligomers (*G1738A*, cyan; *G1748A*, orange and *G1738/1748A*, green). (A, top panel) The

top row represents raw SAXS data, where x-axis and y-axis presents momentum transfer (q) and

the intensity of scattered light, respectively. The second row indicates Guinier plots *(Ln(I,q)) vs*

q^2) suggesting that all samples are pure. Guinier analysis also provides the R_g for each sample

based on the low-q region (See Table1). The third row presenting dimensionless Kratky plots for

each sample demonstrates that all samples are folded and suitable for low-resolution model

building. Finally, the last row represents the electron pair-distance distribution function plots that

provides information about the shape of these molecules, their R_g and maximum particle

dimension. (B, bottom panel) Averaged-filtered models derived using DAMMIN and

DAMAVER calculations for wt (pink), *G1738A* (cyan), *G1748A* (orange) and *G1738/1748A*

(green). The two panels show orthogonal views of G4s, the maximum dimensions (D_{max}) of the

G4 are indicated under each structure, detailed dimensions and parameters are tabulated in Table

1.

Figure 4. Microscale thermophoresis (MST) of FITC-labelled G-quadruplex oligomers—(A) *wt*,

(B) *G1738A*, (C) *G1748A* and (D) *G1738/1748A*—with known G-quadruplex binder, DHX36.

The labelled oligomers were held constant while the DHX36 varied in concentration from 0.006

nM to 20 µM in MST buffer (G-quadruplex buffer supplemented with 0.1% Tween20). MST

was measured in triplicate and background corrected against spectra of buffer alone. The data

above is the average of three independent replicates. Dissociation constants, (K_d), were

computed using K_d fitting and are tabulated in Table 2. The binding affinity was observed in the

order of *G1738A<<wt<G1748A<G1738/1748A*.

The quadruplex-binding protein, DHX36, can pull-down HBV cccDNA derived from HBV

infected human liver tissue.

The biophysical analysis demonstrated that the *wt* oligomer exists as a G-quadruplex *in*

vitro. To address the question of whether the G-quadruplex also exists in HBV genome, we

performed a pull-down assay using DHX36-bound-magnetic beads to study if this host protein

can bind with HBV cccDNA within HBV-positive human liver tissue. We used nested PCR to

amplify HBV DNA in the pull-down products and identified the excess DNA (XS) (*i.e*., the

initial viral DNA input), but none after the washes (Figure 5B). We then noted a strong HBV

DNA signal in the eluted product (EI), demonstrating that DHX36 specifically binds and pull-

down HBV cccDNA (Figure 5B). To determine whether DHX36 recognizes other segments of

the cccDNA outside the region of interest, two HBV gene segments were amplified (i.e., the C

gene with only part of its pre-core promoter region and the X gene, where the 3'-end contains the

quadruplex-forming region of interest; Figure 5A and S2). The pulldown assay, PCR and gel

visualization performed for both of these genes demonstrated an HBV DNA signal (specific PCR

amplicon) in the eluted product for the X which contains the G-quadruplex (Figure 5C, EI), but

not for the C gene segment (Figure 5D, EI). Mutant X genes show binding with DHX36 in this

model as well (Figure 5E), consistent with our findings from MST (Figure 5E). Overall, these

results support the presence of the G-quadruplex in the pre-core promoter region of HBV

cccDNA.

Figure 5: DHX36 pulldown assays. (A) segments of the HBV cccDNA PCR amplified for use in the assay and whole cccDNA extracted from HBV-positive liver tissue; (B) pulldown of whole cccDNA (nested PCR amplified with primers specific to the region of incomplete dsDNA of HBV, nt 1778-1920); (C), pulldown of the X-gene segment; (D), pulldown of the C-gene segment. All PCR were performed on 1:100 dilutions of pull-down washes and elution products. PCR water was used for the negative control; a tandem dimer HBV plasmid was used for positive control. El, Elution; xs, excess DNA. (E) Pulldown study using DHX36 bait protein to bind X gene fragments from the mutant plasmid variants. Sample dilution of 1:800 (left panel) and 1:3200 (right panel), done to account for the slightly differing starting amounts of DNA and the dilution at which the signal is no longer picked up from the nickel bead controls.

The G-quadruplex region influences HBV replication.

Following identification of the G-quadruplex in cccDNA, we investigated the functional impact of this structure on HBV replication. Therefore, we employed a wildtype 1.3mer HBV genome plasmid (kindly gifted from Prof. H. Guo)(59) and designed mutants based on the *G1738A, G1748A,* and *G1738/1748A* sequences used above. The mutant plasmids were transfected into HepG2 cells and a separate GFP control plasmid was simultaneously transfected in all wells to monitor for relative transfection efficiency, inferring relative equivalence (Figure S3). Transfection was consistent across all wells, with an estimated efficiency of about 10% by day 1, and up to 20% for the HBV transfected wells.

The HepG2 transfection studies were performed in triplicate and consistently demonstrated a difference in the viral protein products for *wt* and mutant pre-core promoter regions (Figures 6A-C). The level of HBsAg in the supernatant of HepG2 cells transfected with the *G1738A* and *G1738/1748A* mutant HBV were significantly higher than the *wt* ($p < 0.05$) on days 5 and 7 (Figure 6A, Table S4). Similarly, the levels of HBeAg in supernatant also showed significantly lower production among the wild-type vs. mutant plasmids for nearly all paired sets on days 3 through 7 (Figure 6B, Table S4). In contrast, the high amounts of cellular HBcAg was produced by the *wt* HBV compared to the mutant, with significance noted only between the *wt* and *G1738/1748A* mutant combinations on days 3 through 7 (Figure 6C, S Table S4).

Figure 6: Viral protein markers from 1.3mer HBV transfection studies in HepG2 cells. (A) Average supernatant HBsAg; (B) average supernatant HBeAg; and (C) average cellular HBcAg. Samples were collected at Days 1, 3, 5 and 7 post-transfection and analyzed by various ELISA assays (see Methods). Plots represent the averaged results of the three transfections, with each data point representing at least 8 wells of an ELISA plate. Error bars represent 2x standard error. Significance was calculated using one-way ANOVA with a 2-sided Dunnett t post-hoc analysis. *p-value for significance <0.05. P-values are recorded in the Supplementary Information (Table S4).

Analysis of the nucleic acid levels demonstrated an increase in the amount of cellular DNA detectable above plasmid baseline for all the plasmids at day 7, however, there were no significant differences noted between the wild-type vs. mutant variants after corrections for cells and plasmid (Figure 7, Figure S4-S5). In the supernatant-extracted DNA, a signal was noted on day 7 in one of the three replicate experiments for the *wt* only (Figure 7). The lack of this similar finding in the other replicates likely indicates a lower amount of DNA released into the supernatant relative to the starting plasmid, rather than an experimentally relevant finding. The levels of HBV RNA studies in the supernatant were undetectable when corrected for residual DNA (*i.e.*, reverse transcriptase negative controls) (Figure 8A; Figure S6). There was a

consistent baseline signal in cellular RNA throughout the replicates but no significant difference

between the *wt* and various HBV mutants (Figure 8B). Of note, the amount of detectable plasmid

decreased over time (by ~1 log), consistent with the cellular breakdown of the plasmid (Figure

S5) thus the detected surplus DNA was derived from the newly replicating virus.

Figure 7: DNA from HepG2 transfection studies with wt and mutant HBV plasmids (*G1738A*,

G1748A, and *G1738A/1748A*), harvested at days 1, 3, 5, and 7 post-transfection, isolated from

the cells. DNA was extracted using phenol-chloroform extraction method from both the

supernatant and cells. qPCR was performed using pre-core promoter primers and corrected for

plasmid and internal GAPDH control (See Supplementary Information Table S3 and Figures S4-

S5). Supernatant DNA was also isolated and quantified, but found in only one replicate (see

text).

Figure 8: RNA from HepG2 transfection studies with wt and mutant HBV plasmids (*G1738A*, *G1748A*, and *G1738A/1748A*), harvested at days 1, 3, 5, and 7 post-transfection. RNA was extracted using Trizol extraction followed by DNase digestion. qPCR was performed using primer sets to pick up the core transcripts and total RNA (see Table S3) as noted in the schematic (A). The information is represented as a ratio of core transcripts to total RNA for both the supernatant (B) and cells (C) to account for equivalent detection of residual forms of DNA as detected by qPCR.

DISCUSSION

There is increasing recognition of G-quadruplexes throughout the human genome (25,26,30,31,47). They are frequently found in key regulatory regions of genes and regulate numerous critical cellular processes including transcription, translation and genome stability (25-

28,30,47,60-62). The ubiquity of G-quadruplexes extends to the field of virology where G-

quadruplexes have been found in regulatory regions of various viral genomes (36-46).

In this work, we demonstrate the presence of a G-quadruplex in the HBV pre-core

promoter region, a region critical to the transcription of the HBV C gene and translation of viral

core proteins. The current study describes its evolutionary persistence with the aid of an HBV

sequence repository (https://hbvdb.ibcp.fr), which contains an extensive repository of

replication-competent viruses (63). Across over 10,000 available pre-core/core sequences

worldwide, this region is very G-rich and is highly conserved across nearly all major genotypes

(Figure 1), except for HBV genotype G (excluded in current analysis). HBV G is rarely found

and usually in combination with another genotype (*i.e*., A or H), implying a unique strategy to

maintain replication (64-66). This strong degree of conservation of this G-rich region provides a

rationale for its functional role in virus maintenance, especially in HBV with a highly error-

prone polymerase, capable of producing10^{10-11} point mutations daily (67). If a region of the

genome were to be of high importance to the virus (and not an inconsequential 'DNA knot'), it

would be evolutionarily maintained over time.

Through rigorous biophysical analysis, we demonstrate the ability of this G-rich HBV

pre-core promoter region to form a G-quadruplex, comparing results with those of the HBV

single-nucleotide substitutions, *G1738A, G1748A,* and the double mutation *G1738/1748A*. In

particular, the *G1748A* mutation has been shown to prevent host Sp1 binding and transcript

production (48). We hypothesized that this HBV pre-core mutation affects the quadruplex

structure and disrupts host Sp1 quadruplex binding leading to loss of function noted previously

(48). Our CD studies support the formation of a parallel quadruplex in the *wt* oligomer as well as

the *G1738A* mutant, while the *G1748A* and *G1738/1748A* oligomers form less distinct, likely

altered or distorted quadruplexes (Figure 2). This is consistent with reports that CD spectra

amplitude differs as a function of G-quartet stacks, thus the altered spectra observed in the case

of the mutants support the distortion or alteration of the G-quadruplex (51). More specifically,

the *G1738A* mutation does not appear to result in disruption of the "core" of G-quadruplex,

likely owing to its more 5′ position, while the *G1748A* mutation does. Further, the SAXS data

provide succinct low-resolution structures, indicating the *wt* oligomer indeed forms a quadruplex

(D_{max} = 46.85 Å), while each of the mutants forms an increasingly extended conformation—

G1738A (D_{max} = 53.14 Å) > *G1748A* (D_{max} = 60.00 Å) > *G1738/1748A* (D_{max} = 60.44 Å),

highlighting the impact of these mutations on the secondary structure in this region (Figure 3,

Table 1). When comparing the effects of the mutations on the resultant secondary structure with

these two techniques, the CD and SAXS data agree that the *G1738A* mutant results in a minimal

alteration in the G-quadruplex structure. Conversely, the *G1748A* mutation has a greater impact

on the structure, which when combined with the *G1738A* mutation, appears to be synergistic in

the *G1738/1748A* mutant. Finally, using a known high-affinity parallel quadruplex binder,

DHX36, we demonstrate that the *wt* oligomer can interact with DHX36 (Figures 4 and 5), and

provide a K_d value for this binding of 76.81 ± 12.21 nM through MST studies (Figure 4). This is

a comparable range to other reported K_d values of 0.31 and 0.44 μM, respectively, for DHX36

binding with a human telomerase RNA G-quadruplex (hTR_{1-43}) and basic tetramolecular RNA

G-quadruplex (5′-$A_{15}G_5A_{15}$-3′; 'rAGA') (57,68). This G-quadruplex-binding protein has been

shown to interact at picomolar concentrations, whereby full-length DHX36 binds the RNA G-

quadruplex 'rAGA' with K_d of 39 pM while the DNA equivalent (5′-$A_{15}G_5A_{15}$-3′; 'dAGA') is 77

pM (69). One of the strongest affinities with DHX36 noted in the literature appears to be with a

5′-untranslated region G-quadruplex structure from the Zic1 gene, a gene responsible for

producing a zinc-finger protein critical to early development, with a K_d as low as 3 pM (70).

With regards to our mutant oligomers, binding is also demonstrated with K_d values that fall

within this broad range. The *G1738A* mutant appeared to have stronger binding than the *wt*,

raising the possibility of an alternate binding motif or additional stabilizing interactions with the

DHX36 protein. DHX36 has been shown to bind parallel quadruplexes via its N-terminal domain

in an end-on manner where its first 21 residues are responsible for the quadruplex binding

activity (53,69,71,72). An NMR based study examining the interaction between an 18-amino

acid fragment of DHX36 peptide and a parallel quadruplex showed that there are CH/π- and

CH3/ π-stacking interactions occurring between them. This resulted in DHX36 peptide

effectively covering the tetrad from the top, and subsequently positioning the positively charged

side-chains of three lysines in closer proximity to the negatively charged tetrads further

stabilizing the interaction (72,73). Data from CD and SAXS experiments demonstrate that all the

variants form quadruplexes, hence all of them are capable of binding to DHX36. However, the

altered G4 stacking may have perturbed the aforementioned CH/π- and CH3/π -stacking

interactions which resulted in the varied strength of K_d. It may also be a feature of a smaller

system that could be overcome by employing a longer strand of DNA in which the steric

hindrance from a less organized system would disallow binding. A future direction of this work

will involve developing a higher-resolution structure through molecular modelling (i.e.,

molecular dynamic simulations) to create a model consistent with this experimental data.

In addition to *in vitro* biophysical data of controlled oligomers, we demonstrate its

presence in HBV positive human liver tissue. We employed pull-down assays using cccDNA

extracted from HBV infected liver and demonstrated that the known quadruplex-binder, DHX36

can bind and pull HBV cccDNA from solution (Figure 5B). The inclusion of sub-segments of the

HBV genome serves as controls validating the HBV pre-core region as the major binding site.

Although the *wt* and *G1748A* mutant X-gene show binding, the *G1738A* and *G1738/1748A* do

not bind, and is partially consistent with the structural data analysis. Although it is possible that

the contribution of this binding is shared with the other recently identified G-quadruplex in the

pre-S region, the prior study did not evaluate DHX36 interactions with the HBV pre-S region

(46).

To establish the functional significance of the presence of G-quadruplex in the HBV

genome, we compared the wild-type HBV with *G1738A*, *G1748A* and *G1738/1748A* mutant

plasmids in *in vitro* cell culture studies. It was postulated that mutations which disrupt the

quadruplex region would have reduced binding from host proteins to initiate transcription and

thus impact downstream viral replication. To investigate this, we used a wild-type 1.3-mer HBV

plasmid(59) and created each of the above mutants (*G1738A*, *G1748A* and *G1738/1748A*)

through site-directed mutagenesis. Previous studies with a linearized 1-mer HBV model, in

which the HBV genome was excised from the plasmid and allowed to re-circularize by natural

DNA repair mechanisms once in the cell, did not prove effective at producing HBeAg or HBcAg

at detectable levels (see Supplementary Information —Supplementary Experiments and Figures

S7-10)(74). This may have been due to incomplete circularization inside the cell. Since the

location of the cut site is through the core gene, the production of HBeAg and HBcAg proteins

would be fully reliant on the circularization, whereas the transcripts and open reading frame for

HBsAg production would not be affected. Thus, the 1.3-mer HBV model was adopted to

overcome these deficiencies, allow for the more balanced production of all viral transcripts and

enable an analysis of sequence impacts on viral replication within this cell culture system. We

noted differences between mutant and wild-type transfections in some of the markers of viral

replication typically after ~ 5 days (Figures 6-8). Notably, HBsAg in cell supernatant showed

that the mutant variants, *G1738A* and *G1738/1748A,* produced significantly more HBsAg than

the *wt* on days 5 and 7 (Figure 6A). As the transcript for HBsAg production is independent of the

core promoter, the increase in HBsAg for the mutants compared to *wt* promoter region may be

explained by a shift in cellular resources. It is possible, that the downregulation of one transcript

(*i.e*., the core transcript), leads to increased production of other viral transcripts (*i.e*., the HBV S

transcript). This shift in specific viral protein production could occur if the necessary secondary

structure required for transcription factor binding is missing or distorted, as might be the case for

pre-core promoter region G-quadruplex in the presence of the *G1748A* mutant. This phenomenon

was also observed by Li and Ou (48), where they noted a three-fold increase in S RNA transcript

with a G1736 deletion mutation in the pre-core promoter region. Analysis of the HBeAg showed

greater levels produced by HBV mutants compared to *wt* virus (Figure 6B). In comparison, a

significant increase in levels of HBcAg by the *wt* virus compared to the *G1738/1748A* mutant

was observed (Figure 6C). This suggests subtle differences in viral core protein production.

Although *nt* changes within each mutant may disrupt the quadruplex, they do not abolish HBcAg

production, as would have been expected if the quadruplex was critical to transcription factor

binding. This is reflected in our biophysical data where we show the varied binding affinities

with the selective quadruplex binder, DHX36. Notably, the double mutation *G1738/1748A* had

the weakest binding based on MST analysis and the most unfolded state in the SAXS data and

functional studies show it to have the most significant decrease in HBcAg production. The data

suggests that even partial quadruplex formation is sufficient for the cell's usual transcription

machinery and any possible subtle differences may not be detectable in the less-efficient cell

culture model. Further, the data is consistent with results by Biswas et al., involving the

quadruplex study of HBV's S promoter region in which only a partial decrease in downstream

products rather than complete loss of viral protein due to quadruplex-disrupting mutations(46).

We determined the effect of the *G1738A*, *G1748A* and *G1738/1748A* mutations on the

production of viral nucleic acid products. The total supernatant DNA was detected on day 7 post-

transfection in only one of the three replicates (log 5.7 copies DNA/mL in transfection #2). The

cellular DNA was detectable for all plasmid forms on day 7, but without significant differences

among the *wt* and mutant comparators (Figure 7). Total HBV RNA in the supernatant and in the

cells was at a very low-level when corrected for residual DNA (i.e., starting plasmid or other

HBV DNA products). To overcome potential confounders from larger amounts of DNA, we

compared the RNA transcripts as a ratio of those produced from the core promoter (*i.e.*, pre-core

and pgRNA transcripts) to the total RNA (*i.e.*, as measured by a region just prior to the poly-A

tail signal) (Figure 8A). We assume the relative PCR efficiency for each of the two primer sets is

comparable and equivalent for each comparator thus these values represent a relative impact of

the mutation on core and pgRNA transcript production. For *wt* HBV, the ratio of supernatant

core and pgRNA transcripts appear to decrease over time, whereas in the cells, this amount

appears to be relatively stable (Figure 8B,C). For the mutant HBV plasmids, the ratios of core to

total RNA transcripts follow a similar pattern, without a clear trend for favouring alternate

transcripts.

Overall, using the 1.3-mer HBV plasmid model, the differences in the *wt* and various

mutant pre-core promoter HBV transfections suggest the only minimal influence of the

mutations on overall downstream function. There are several possible explanations for this

finding; (1) transcription factor binding to promoter regions of the 1.3-mer HBV plasmid is

independent of or only partially dependent on the quadruplex under study. It is possible that in

the setting of high levels of plasmid, even a less-optimal binding alternative may still favour

production despite potential structural controls. Therefore, this model may not be representative

of natural processes despite successes with this approach for other pathway studies (59,75,76).

(2) It is also possible that the quadruplex structure is not needed for binding and that the primary

nucleotide sequence is the sole influencer of the subtle finding in this model. Fully delineating

the two possibilities in an experimental model would require conditions that affect the

quadruplex formation *in vivo* via the stabilizing metal ion, and that would still be compatible

with cell growth or function. While known small molecule G-quadruplex stabilizers would be

another possibility to investigate this phenomenon, the off-target effects of these agents on the

cells' normal functioning may limit evaluation.

In summary, our work provides robust biophysical evidence and novel functional data for

the existence of a G-quadruplex structure in the pre-core promoter region of the HBV genome.

The data demonstrates that we can use a known quadruplex-binder to bind a physiologically-

relevant, transcriptional template of HBV—the resilient cccDNA. Furthermore, using a well-

validated HBV culture model, we provide evidence for a functional effect on HBV replication.

This unique structural motif represents a novel HBV therapeutic target the HBV cccDNA

minichromosome and potentially informs the development of a sterilizing or virological cure for

HBV infection.

EXPERIMENTAL PROCEDURE

Analysis of the HBV core promoter region.

The HBV genome database (HBVdB: https://hbvdb.ibcp.fr, accessed April, 2019) was

used to access 9939 HBV core promoter sequences from 7 of the 8 main HBV genotypes (977

genotype A; 2478 genotype B; 3313 genotype C; 1545 genotype D; 389 genotype E; 387

genotype F; and 52 genotype H). Sequences were downloaded as a Clustal W alignment (77),

truncated as the 23-mer region (nt1732-1754) and examined for the presence and frequency of

single nucleotide polymorphisms with the aid of the Geneious Prime program (2019.1.3;

www.geneious.com). HBV genotype G sequences lack the highly G-rich region and were

excluded from analysis (21).

Quadruplex preparation.

All DNA oligomers were ordered from AlphaDNA (Montreal, QC, Canada), with high-

performance liquid chromatography (HPLC) purification by the manufacturer. The pre-core

oligomers were as follows: wild-type (*wt*), 5′-CTGGGA**G**GAGCTGGGG**G**AGGAGA-3′ and

single-nucleotide substitution mutants (*G1738A*) 5′-CTGGGA**A**GAGCTGGGG**G**AGGAGA-3′,

(*G1748A*)   5′-CTGGGA**G**GAGCTGGGG**A**AGGAGA-3′   and   (*G1738/1748A*)   5′-

CTGGGA**A**GAGCTGGGG**A**AGGAGA-3′. Samples were dissolved in G-quadruplex buffer (20

mM HEPES, pH 7.5, 100 mM KCl, 1 mM EDTA) at a concentration of 5 μM, confirmed

spectrophotometrically using extinction coefficients for 260 nm (calculated using IDT

OligoAnalyzer tool) of 237,400 M^{-1} cm^{-1}(*wt*), 239,300 M^{-1} cm^{-1}(*G1738A* and *G1748A*) and

241200 M^{-1} cm^{-1} (*G1738/1748A*). Samples were heated to 95°C for ten minutes, then slowly

cooled to room temperature to allow assembly of the G-quadruplex. Samples were concentrated

to 500 μL using a 3 kDa Vivaspin™ 20 filter column (GE Healthcare, Mississauga, ON,

Canada). The conformational mixtures were purified via size exclusion chromatography (SEC)

using a HiLoad Superdex 75 10/300 in G-quadruplex buffer, loaded with 500 μL sample and

eluted at 0.65 mL/min using G-buffer (20 mM HEPES, pH 7.5, 100 mM KCl, 1 mM EDTA).

Only peak fractions (e.g., for G1738A, only fractions from ~12.5 to 15 mL, see Figure 2A) were

pooled and concentrated using a 3 kDa spin column, for use in subsequent experiments.

Circular Dichroism (CD) spectropolarimetry.

497   All spectra for wt and mutant G-quadruplexes were recorded on a calibrated Jasco J-815

498   spectropolarimeter (Jasco Inc., Easton, MD) from 220–320 nm in a 1.0 mm cell and a 32 s

499   integration time. SEC purified oligomers were concentrated to 20 μM in G-quadruplex buffer.

500   Measurements were performed in triplicate and baseline-corrected by subtraction of the degassed

501   buffer alone.

Molecular cloning and protein preparation of DHX36.

DHX36 was prepared as previously described (78). Briefly, amino acid residues 53-105

of human DHX36 were PCR amplified from a gBlock® (Integrated DNA Technologies, Kanata,

ON, Canada) to add NdeI and SalI restriction sites. The PCR product was inserted into the

pET28a(+) vector using NdeI and SalI restriction enzymes and T4 DNA ligase (Thermo Fisher

Scientific, Waltham, MA) and subsequently transformed into chemically competent DH5α *E.*

coli cells (New England Biolabs (NEB), Ipswich, MA). Positive clones were identified, and the

gene sequence was confirmed by Sanger sequencing (Genewiz, South Plainfield, NJ). The final

pET28a-DHX36$_{53-105}$ expression vector contained a vector-encoded N-terminal hexahistidine tag

and a thrombin cleavage site for tag removal. The pET28a-DHX36$_{53-105}$ expression vector was

transformed into competent *E. coli* cells Lemo21(DE3) (NEB), and expression induced via the

addition of 1mM isopropyl β-D-1-thiogalactopyranoside. Cells were harvested three hours post-

induction by centrifugation at 5000 g for 15 minutes, flash-frozen, and stored at -80°C.

Purification of the protein was performed via a nickel affinity column, followed by SEC as

described previously (52,53).

Microscale Thermophoresis (MST).

MST was used to determine the nature and strength of the interactions between the G-

quadruplex with DHX36, where the G-quadruplex oligomers were labelled with FITC and held

constant, similar to described (79,80). First, DHX36 was diluted from 20 µM to 0.006 nM in

MST buffer (G-quadruplex buffer supplemented with 0.1% Tween20). Next, the folded and

SEC-purified oligomers were diluted to 50 nM in MST buffer and added to each of the tubes of

the DHX36 dilution series. Samples were run using Monolith NT.115 standard capillaries

(Nanotemper Technologies, San Francisco, CA). The data were collected at the high excitation

setting with the laser power set at 20%. Microscale thermophoresis was measured in triplicate

and background corrected against spectra of buffer alone. The data from three independent

replicates were analyzed using MO Affinity Analysis software v2.1.3 and fit to the standard K_d

fit model.

Small angle X-ray scattering (SAXS).

HPLC-SAXS data for the wildtype and mutant variant oligomers were collected using the

B21 BioSAXS beamline at the Diamond Light Source (Didcot, UK) synchrotron facility, under a

previously described protocol (52). Using an Agilent 1200 (Agilent Technologies, Stockport,

UK) in-line HPLC with a specialized flow cell, oligomer data was collected at 100 μM

concentrations in G-quadruplex buffer. Each oligomer sample was injected into a Shodex

KW403-4F (Showa Denko America, Inc.) column pre-equilibrated in G-quadruplex buffer. X-

rays were exposed to each frame for 3s and on average, 10 frames from each individual sample

peak were integrated, followed by buffer subtraction, and then merged using ScÅtter (81) as

previously described (80).

The data was imported into software Primus from the ATSAS suite of software programs

version 2.8 (82), Guinier analysis was performed on the dataset to ascertain the quality of the

data and estimate the Radius of Gyration (R_g) (Figure S1). For calculation of *ab initio* low

resolution structures the maximum, linear dimension (D_{max}) was estimated using the pairwise

distance distribution function *P(r)* plot, along with the R_g, estimated using GNOM (55). Using

the *P(r)* plot information, ten models were calculated using DAMMIN (83), by selecting a

different random seed for each model but keeping identical parameters within each set ensuring

no enforced symmetry (*P1*). The program DAMAVER (84) was used to average and filter the

models to produce a representative model, as described previously (85).

Isolation and preparation of HBV cccDNA.

HBV DNA positive liver tissue was obtained under an approved ethics protocol

(University of Calgary Conjoint Health Research Ethics Board, ID# 16636). Protein-free, nuclear

DNA was extracted from liver by Hirt extraction method, as previously described (86). Samples

were treated with T5 exonuclease (NEB, Ipswich, MA, Cat# M0363) to digest relaxed circular

DNA (rcDNA) or remaining genomic fragments. Products were further cleaned by column

purification (E.Z.N.A. Cycle-Pure Kit, Omega Bio-tek, Norcross, GA) to remove the

exonuclease. HBV cccDNA was amplified by nested PCR, using HBV specific primers spanning

the nick region of the rcDNA genome, as described (86,87).

Amplification of HBV DNA fragments.

A full-length HBV (genotype C) plasmid was used for PCR amplification to produce

HBV DNA fragments of the C gene and promoter region that included the quadruplex forming

region (X fragment, "X") or not (C fragment, "C"), using HBV specific gene primers coding for

nt 1606-1974 (368bp, "X") and *nt* 1825-2286 (461bp, "C"), as described (87). The PCR

amplicons were visualized on a 1.2% agarose gel and purified using the Qiagen gel DNA

extraction kit (Qiagen Inc, Montreal) (See Figure S2, Table S3). The gel-purified products were

heated to 95°C for 5 mins, then cooled slowly to room temperature to enable folding.

HBV genome pull-down assays.

Magnetic beads (HisPur Ni-NTA, ThermoFisher, Cat# 88831) and magnetic tube rack

(Qiagen Inc, Montreal) were used for HBV genome pull-down assays using DHX36 (with an

associated His-tag) as the "bait" protein. The protocol provided with the magnetic beads was

loosely followed, with modifications for our system. Firstly, 40 μL of Nickel bead solution was

added to each tube and equilibrated with washes of equilibration/wash buffer (30 mM imidazole

in phosphate-buffered saline (PBS), 0.05% Tween, pH 7.5), per protocol. Secondly, a 1:1

solution of DHX36 in equilibration/wash buffer (end protein concentration of 10 μM) was

incubated with the beads (400 μL/tube) on an end-on-end rotor for 30 mins at 4$^\text{o}$C. The excess

protein was removed, and the beads were washed with buffer until fluorometry readings showed

undetectable protein for two consecutive readings (four washes). A dilution of cccDNA (whole

or fragments) was made with the buffer to provide 400 μL of the solution for each tube (52, 17

and 26 ng/μL, respectively, for the whole cccDNA, C and X regions). The viral DNA was

incubated with beads (+/- bound DHX36) for 2 hours on a rocker at 4$^\text{o}$C. Following removal of

excess DNA, the beads were washed with buffer until fluorometry readings showed undetectable

DNA for two consecutive readings (four washes). The protein-DNA complexes were eluted with

30 μL elution buffer (250 mM imidazole in PBS, pH 7.5).

Detection of cccDNA and cccDNA fragments.

To confirm the presence of HBV cccDNA in the Hirt-extracted, T5 exonuclease-digested

liver samples, nested PCR was performed to amplify the "nicked" region of the rcDNA, or

incompletely dsDNA, using HBV-specific primers (5′-ACTCCTGGACTCTCAGCAATG-3′

(cccDNA-D-For);   5′-GTATGGTGAGGTGAGCAATG-3′   (cccDNA-D-Rev);   5′-

AGGCTGTAGGCACAAATTGGT-3′   (cccDNA-N-For);   and   5′-

GCTTATACGGGTCAATGTCCA-3′ (cccDNA-N-Rev). The products obtained from the X and

C region pull-downs were PCR-amplified using the same primers used for their production (see

above) and amplicons visualized with agarose gels and SafeView® (Applied Biological

Materials Inc., Vancouver) staining under UV light.

HBV plasmids for transfection of HepG2 cells.

HBV mutants (*G1738A*, *G1748A* and *G1738/1748A*) were created using a 1.3mer HBV

plasmid (gifted by Dr. Haitao Guo)(59) and site-directed mutagenesis kit (QuikChangeII, Agilent

Technologies, Santa Clara, CA) with corresponding primers (University of Calgary DNA

Synthesis Lab, Calgary, AB) (see Table S2). The mutant HBV sequences were confirmed by

Sanger sequencing. The original plasmid (*wt*) and the core-mutants (*G1738A*, *G1748A* and

G1738/1748A) were amplified via an *E. coli* (Top10) system (Thermo Fisher, Cat# C404010)

and isolated using the GenElute Plasmid Miniprep kit (Sigma, Cat# PLN350). In addition, an in-

house green fluorescent protein (GFP)-pcDNA3.1(+) plasmid was used as control.

HepG2 transfections and isolation of total nucleic acid and protein

HepG2 cells (New England Biolabs) were grown to 80-90% confluence in Dulbecco's

Modified Enrichment Media (DMEM) supplemented with 10% fetal bovine serum and 100

U/mL penicillin/streptomycin. Approximately 24 hours prior to transfection, the cells were split

and seeded into 6-well plates at concentrations of 1×10^6 cells/well and allowed to re-attach

overnight. On day 0, cell media was exchanged to remove any unattached cells. The

corresponding wells were transfected with GFP alone, GFP + HBV_{wt} or GFP+$HBV_{mutant\ (G1738A,}$

$_{G1748A\ or\ G1738/1748A)}$ (500 ng GFP plasmid/well and 270 ng HBV plasmid/well), using

Lipofectamine3000 (ThermoFisher) per the manufacturer's protocol. On days 1, 3, 5 and 7, cells

were imaged for GFP to determine % transfection relative to GFP signal, and subsequently, both

the supernatant and cells were harvested for detection of HBV RNA, total HBV DNA, and/or

viral proteins. Total RNA was extracted via TRIzol® (Invitrogen, Carlsbad, CA), as described

(87); total DNA was isolated from cells via phenol-chloroform (supernatant) or TRIzol®. Viral

proteins (HBsAg, HBeAg) were detected in the cell supernatant without further manipulation

and using culture media for controls (see ELISA details below). HBcAg in cells was detected

from the lysate of PBS-washed cells treated with 1% Triton X-100 in TBS buffer. All nucleic

acid extractions were performed with simultaneous "mock" extractions under strict lab protocols

to prevent cross-contamination. Transfection experiments were performed in triplicate with

duplicate samples for each condition per day, per experiment.

Quantification of HBV RNA and DNA.

HBV RNA from the transfected cell supernatant was detected by TaqMan® qPCR

following the protocols previously described (88). Total supernatant DNA, cellular RNA and

DNA were detected using SYBR® Green fluorescence (BioRad, Cat# 1725124). An HBV-

containing plasmid was used as a positive control and to produce the dilution series for the qPCR

standards; water and "mock extracted" template and mock transfection were used as negative

controls. A segment of the pre-core promoter region was amplified using the primers HBV-PCP-

For and HBV-PCP-Rev and subsequently corrected for starting plasmid using primers Plasmid-

Guo-For and Plasmid-Guo-Rev (See Table S2). Samples were prepared in triplicate with

concomitant standards and controls per qPCR run and all transfections were performed in

triplicate. The qPCR amplicons from the *wt*, various mutants and GFP transfected controls were

confirmed by agarose gel visualization to confirm single products. In the GFP control, a qPCR

signal was noted with a different melting curve than the HBV product and this was confirmed to

be a single 800-bp product on a gel that was not seen in the HBV transfected samples, where only the expected HBV fragment size of 340 bp was seen. Primer alignment of the GFP pcDNA3.1(+) plasmid confirmed this amplicon size and poor sequence overlap. HBV nucleic acid detection was corrected for each experimental condition (per the GAPDH internal control).

Detection of HBsAg, HBeAg and HBcAg.

HBsAg was detected by sandwich ELISAs via an in-house developed assay. Briefly, using a 96-well plate (Corning Incorporated, Corning, NY, Cat# 3361), ELISA plates were prepared using HBsAg capture Ab (Fitzgerald Industries International, North Acton, MA, Cat#10-H05H) at 1 ug/mL. An HBsAg dilution series (500 ng/mL - 4 ng/mL) was prepared for use as standards (Fitzgerald, Cat #C-CP2019R). To each corresponding well, 100 μL of supernatant or standards was added in triplicate and incubated overnight. The HBsAg was detected using an HRP-conjugated anti-HBsAg antibody (Fitzgerald, Cat# 60C-CR2100RX, 1:2000 dilution), activated by TMB (LifeTech Novex, Cat#00-2023), quenched by 1 M HCl and read at 450 nm (BioRad iMark, USA). The total protein per sample was determined via the BCA protein assay standards kit (Biorad, Cat# 500-0006) and used to correct for varying protein content within the cell supernatant. HBeAg in the supernatant was analyzed via an electrochemiluminescent immunoanalyzer (Roche Elecsys run on the Cobas e411 immunoanalyzer, Roche Diagnostics, Laval, QC) using the sandwich principle involving a reaction with two HBeAg-specific monoclonal antibodies (one biotinylated and the other labeled with a ruthenium complex). The complex was bound to the solid phase consisting of streptavidin-coated magnetic microparticles (assay detection limit of is ≤ 0.30 Paul☐Ehrlich☐Institute U/mL, corresponding to 1.6 cut-off index (COI) in the assay's detection

units). Additionally, cellular HBcAg was analyzed using a commercial kit (Cell Biolabs Inc, San

Diego, CA, Cat# VPK150), following the manufacturer's instructions. All data shown is an

average of three independent runs performed as duplicates.

Statistical analysis.

Data were analyzed using SPSS statistical software, ver 26 (SPSS Inc., Chicago, IL, USA) and

Microsoft Excel. One-way ANOVA analyses with a 2-sided Dunnett t post-hoc analysis were

employed to determine the significance of the transfection parameters. Statistical significance

was set at *p* values of less than 0.05.

DATA AVAILABILITY

All data has been presented in this article.

ACKNOWLEDGEMENTS

We thank Dr Hans-Joachim Wieden for allowing us to use the circular dichroism spectroscopy

instrument.

ACKNOWLEDGEMENT

TRP & CC: conceptualization of the study, designing of experiments, analysis of data,

preparation of manuscript, VMS: designed experiments, performed molecular and cell biology

experiments analysis of data, preparation of manuscript, MDB: designed experiments, performed

biophysics experiments analysis of data, preparation of manuscript, TM: performed MST

experiments, preparation of manuscript KCKL: assisted in cell biology experiments, preparation

of manuscript

SKS: designing of DHX36$_{53-105}$, DLG: expression and purification of DHX36$_{53-105}$ for cell

biology experiments, CO: performed the quantitative ELISA experiments, GvM: data analysis.

All authors contributed to manuscript preparation.

FUNDING INFORMATION

VMS acknowledges the Canadian Network on Hepatitis C Postdoctoral Fellowship. TRP is a

Canada Research Chair in RNA and Protein Biophysics and acknowledges the Canada

Foundation for Innovation infrastructure grant. TRP and CC acknowledge the Cumming School

of Medicine Seed Grant and Alberta Innovates Strategic Research Program grant. TM and MB

were hired using NSERC Discovery (RGPIN-2017-04004) and Alberta Innovates Strategic

Research Program grant, respectively. TM and MB also acknowledges NSERC PGS-D and

MITACS fellowships. HPLC-SAXS data were collected at the B21 beamline, DIAMOND

synchrotron (BAG proposal: SM16028), United Kingdom.

CONFLICTS OF INTEREST

The authors declare that they have no conflicts of interest with the contents of this article.

SUPPLEMENTARY INFORMATION

Supplementary information is available at JBC online.

ABBREVIATION

HBV = Hepatitis B virus,

R_g = Radius of Gyration,

D_{max} = maximum dimension,

MST = Microscale Thermophoresis,

SAXS = Small-angle X-ray Scattering

REFERENCES

1. WHO. (2018) Global Hepatitis Report, 2017. WHO
2. Beasley, R. P. (1988) Hepatitis B virus. The major etiology of hepatocellular carcinoma. *Cancer* **61**, 1942-1956
3. Ganem, D., and Prince, A. M. (2004) Hepatitis B virus infection--natural history and clinical consequences. *N Engl J Med* **350**, 1118-1129
4. Yang, H. C., and Kao, J. H. (2014) Persistence of hepatitis B virus covalently closed circular DNA in hepatocytes: molecular mechanisms and clinical significance. *Emerg Microbes Infect* **3**, e64
5. Michalak, T. I., Pasquinelli, C., Guilhot, S., and Chisari, F. V. (1994) Hepatitis B virus persistence after recovery from acute viral hepatitis. *J Clin Invest* **94**, 907
6. Nassal, M. (2015) HBV cccDNA: viral persistence reservoir and key obstacle for a cure of chronic hepatitis B. *Gut* **64**, 1972-1984
7. Zoulim, F. (2005) New insight on hepatitis B virus persistence from the study of intrahepatic viral cccDNA. *Journal of hepatology* **42**, 302-308
8. Zoulim, F., Lebosse, F., and Levrero, M. (2016) Current treatments for chronic hepatitis B virus infections. *Curr Opin Virol* **18**, 109-116
9. Lee, H. M., and Banini, B. A. (2019) Updates on Chronic HBV: Current Challenges and Future Goals. *Current treatment options in gastroenterology* **17**, 271-291
10. Terrault, N. A., Lok, A. S. F., McMahon, B. J., Chang, K.-M., Hwang, J. P., Jonas, M. M., Brown, R. S., Jr., Bzowej, N. H., and Wong, J. B. (2018) Update on prevention, diagnosis, and treatment of chronic hepatitis B: AASLD 2018 hepatitis B guidance. *Hepatology (Baltimore, Md.)* **67**, 1560-1599
11. Coffin, C. S., Fung, S. K., Alvarez, F., Cooper, C. L., Doucette, K. E., Fournier, C., Kelly, E., Ko, H. H., Ma, M. M., Martin, S. R., Osiowy, C., Ramji, A., Tam, E., and Villeneuve, J. P. (2018) Management of Hepatitis B Virus Infection: 2018 Guidelines from the Canadian Association for the Study of Liver Disease and Association of Medical Microbiology and Infectious Disease Canada. *Canadian Liver Journal* **1**, 156-217
12. Ward, H., Tang, L., Poonia, B., and Kottilil, S. (2016) Treatment of hepatitis B virus: an update. *Future Microbiol* **11**, 1581-1597
13. Chauhan, R., Lingala, S., Gadiparthi, C., Lahiri, N., Mohanty, S. R., Wu, J., Michalak, T. I., and Satapathy, S. K. (2018) Reactivation of hepatitis B after liver transplantation: Current knowledge, molecular mechanisms and implications in management. *World journal of hepatology* **10**, 352-370
14. Quasdorff, M., and Protzer, U. (2010) Control of hepatitis B virus at the level of transcription. *J Viral Hepat* **17**, 527-536
15. Nassal, M. (2008) Hepatitis B viruses: reverse transcription a different way. *Virus Res* **134**, 235-249
16. Turton, K. L., Meier-Stephenson, V., Badmalia, M. D., Coffin, C. S., and Patel, T. R. (2020) Host Transcription Factors in Hepatitis B Virus RNA Synthesis. *Viruses* **12**, 160
17. Bartholomeusz, A., and Schaefer, S. (2004) Hepatitis B virus genotypes: comparison of genotyping methods. *Reviews in Medical Virology* **14**, 3-16
18. Sunbul, M. (2014) Hepatitis B virus genotypes: global distribution and clinical importance. *World journal of gastroenterology* **20**, 5427-5434

19. Buti, M., Rodriguez-Frias, F., Jardi, R., and Esteban, R. (2005) Hepatitis B virus genome
variability and disease progression: the impact of pre-core mutants and HBV genotypes.
Journal of clinical virology : the official publication of the Pan American Society for
Clinical Virology **34 Suppl 1**, S79-82
20. Lin, C. L., and Kao, J. H. (2017) Natural history of acute and chronic hepatitis B: The
role of HBV genotypes and mutants. *Best practice & research. Clinical gastroenterology*
31, 249-255
21. Meier-Stephenson, V., Bremner, W. T. R., Dalton, C. S., Van Marle, G., Coffin, C. S.,
and Patel, T. R. (2018) Comprehensive Analysis of Hepatitis B Virus Promoter Region
Mutations. *Viruses* **10**, 603
22. Burge, S., Parkinson, G. N., Hazel, P., Todd, A. K., and Neidle, S. (2006) Quadruplex
DNA: sequence, topology and structure. *Nucleic acids research* **34**, 5402-5415
23. Dapic, V., Abdomerovic, V., Marrington, R., Peberdy, J., Rodger, A., Trent, J. O., and
Bates, P. J. (2003) Biophysical and biological properties of quadruplex
oligodeoxyribonucleotides. *Nucleic Acids Res* **31**, 2097-2107
24. Vlasenok, M., Varizhuk, A., Kaluzhny, D., Smirnov, I., and Pozmogova, G. (2017) Data
on secondary structures and ligand interactions of G-rich oligonucleotides that defy the
classical formula for G4 motifs. *Data in Brief* **11**, 258-265
25. Huppert, J. L., and Balasubramanian, S. (2005) Prevalence of quadruplexes in the human
genome. *Nucleic Acids Res* **33**, 2908-2916
26. Lam, E. Y., Beraldi, D., Tannahill, D., and Balasubramanian, S. (2013) G-quadruplex
structures are stable and detectable in human genomic DNA. *Nat Commun* **4**, 1796
27. Hansel-Hertsch, R., Di Antonio, M., and Balasubramanian, S. (2017) DNA G-
quadruplexes in the human genome: detection, functions and therapeutic potential.
Nature reviews. Molecular cell biology **18**, 279-284
28. Rigo, R., Palumbo, M., and Sissi, C. (2017) G-quadruplexes in human promoters: A
challenge for therapeutic applications. *Biochimica et Biophysica Acta (BBA) - General*
Subjects **1861**, 1399-1413
29. König, S. L. B., Evans, A. C., and Huppert, J. L. (2010) Seven essential questions on G-
quadruplexes. in *BioMolecular Concepts*
30. Rhodes, D., and Lipps, H. J. (2015) G-quadruplexes and their regulatory roles in biology.
Nucleic Acids Research **43**, 8627-8637
31. Monsen, R. C., DeLeeuw, L., Dean, W. L., Gray, R. D., Sabo, T M., Chakravarthy, S.,
Chaires, J. B., and Trent, J. O. (2020) The hTERT core promoter forms three parallel G-
quadruplexes. *Nucleic Acids Research* **48**, 5720-5734
32. Asamitsu, S., Obata, S., Yu, Z., Bando, T., and Sugiyama, H. (2019) Recent Progress of
Targeted G-Quadruplex-Preferred Ligands Toward Cancer Therapy. *Molecules (Basel,*
Switzerland) **24**, 429
33. Drygin, D., Siddiqui-Jain, A., O'Brien, S., Schwaebe, M., Lin, A., Bliesath, J., Ho, C. B.,
Proffitt, C., Trent, K., Whitten, J. P., Lim, J. K., Von Hoff, D., Anderes, K., and Rice, W.
G. (2009) Anticancer activity of CX-3543: a direct inhibitor of rRNA biogenesis. *Cancer*
research **69**, 7653-7661
34. Sun, D., Thompson, B., Cathers, B. E., Salazar, M., Kerwin, S. M., Trent, J. O., Jenkins,
T. C., Neidle, S., and Hurley, L. H. (1997) Inhibition of human telomerase by a G-
quadruplex-interactive compound. *Journal of medicinal chemistry* **40**, 2113-2116

35. Xu, H., Di Antonio, M., McKinney, S., Mathew, V., Ho, B., O'Neil, N. J., Santos, N. D.,
Silvester, J., Wei, V., Garcia, J., Kabeer, F., Lai, D., Soriano, P., Banáth, J., Chiu, D. S.,
Yap, D., Le, D. D., Ye, F. B., Zhang, A., Thu, K., Soong, J., Lin, S.-c., Tsai, A. H. C.,
Osako, T., Algara, T., Saunders, D. N., Wong, J., Xian, J., Bally, M. B., Brenton, J. D.,
Brown, G. W., Shah, S. P., Cescon, D., Mak, T. W., Caldas, C., Stirling, P. C., Hieter, P.,
Balasubramanian, S., and Aparicio, S. (2017) CX-5461 is a DNA G-quadruplex stabilizer
with selective lethality in BRCA1/2 deficient tumours. *Nature Communications* **8**, 14432
36. Piekna-Przybylska, D., Sullivan, M. A., Sharma, G., and Bambara, R. A. (2014) U3
region in the HIV-1 genome adopts a G-quadruplex structure in its RNA and DNA
sequence. *Biochemistry* **53**, 2581-2593
37. Artusi, S., Nadai, M., Perrone, R., Biasolo, M. A., Palu, G., Flamand, L., Calistri, A., and
Richter, S. N. (2015) The Herpes Simplex Virus-1 genome contains multiple clusters of
repeated G-quadruplex: Implications for the antiviral activity of a G-quadruplex ligand.
Antiviral research **118**, 123-131
38. Frasson, I., Soldà, P., Nadai, M., Lago, S., and Richter, S. N. (2020) Transcription factor
recruitment by parallel G-quadruplexes to promote transcription: the case of herpes
simplex virus-1 ICP4. *bioRxiv*, 2020.2007.2014.202747
39. Luo, X., Xue, B., Feng, G., Zhang, J., Lin, B., Zeng, P., Li, H., Yi, H., Zhang, X. L., Zhu,
H., and Nie, Z. (2019) Lighting up the Native Viral RNA Genome with a Fluorogenic
Probe for the Live-Cell Visualization of Virus Infection. *Journal of the American*
Chemical Society **141**, 5182-5191
40. Tluckova, K., Marusic, M., Tothova, P., Bauer, L., Sket, P., Plavec, J., and Viglasky, V.
(2013) Human papillomavirus G-quadruplexes. *Biochemistry* **52**, 7207-7216
41. Murat, P., Zhong, J., Lekieffre, L., Cowieson, N. P., Clancy, J. L., Preiss, T.,
Balasubramanian, S., Khanna, R., and Tellam, J. (2014) G-quadruplexes regulate
Epstein-Barr virus-encoded nuclear antigen 1 mRNA translation. *Nat Chem Biol* **10**, 358-
364
42. Madireddy, A., Purushothaman, P., Loosbroock, C. P., Robertson, E. S., Schildkraut, C.
L., and Verma, S. C. (2016) G-quadruplex-interacting compounds alter latent DNA
replication and episomal persistence of KSHV. *Nucleic Acids Res* **44**, 3675-3694
43. Fleming, A. M., Ding, Y., Alenko, A., and Burrows, C. J. (2016) Zika Virus Genomic
RNA Possesses Conserved G-Quadruplexes Characteristic of the Flaviviridae Family.
ACS Infectious Diseases
44. Lavezzo, E., Berselli, M., Frasson, I., Perrone, R., Palu, G., Brazzale, A. R., Richter, S.
N., and Toppo, S. (2018) G-quadruplex forming sequences in the genome of all known
human viruses: A comprehensive guide. *PLoS computational biology* **14**, e1006675
45. Metifiot, M., Amrane, S., Litvak, S., and Andreola, M. L. (2014) G-quadruplexes in
viruses: function and potential therapeutic applications. *Nucleic Acids Res* **42**, 12352-
12366
46. Biswas, B., Kandpal, M., and Vivekanandan, P. (2017) A G-quadruplex motif in an
envelope gene promoter regulates transcription and virion secretion in HBV genotype B.
Nucleic Acids Research **45**, 11268-11280
47. Todd, A. K., Johnston, M., and Neidle, S. (2005) Highly prevalent putative quadruplex
sequence motifs in human DNA. *Nucleic Acids Res* **33**, 2901-2907
48. Li, J., and Ou, J. H. (2001) Differential regulation of hepatitis B virus gene expression by
the Sp1 transcription factor. *Journal of virology* **75**, 8400-8406

49.   Vorlickova, M., Kejnovska, I., Sagi, J., Renciuk, D., Bednarova, K., Motlova, J., and
853   Kypr, J. (2012) Circular dichroism and guanine quadruplexes. *Methods (San Diego,*
854   *Calif.)* **57**, 64-75

855   50.   Carvalho, J., Queiroz, J. A., and Cruz, C. (2017) Circular Dichroism of G-Quadruplex: a
856   Laboratory Experiment for the Study of Topology and Ligand Binding. *Journal of*
857   *Chemical Education* **94**, 1547-1551

858   51.   del Villar-Guerra, R., Trent, J. O., and Chaires, J. B. (2018) G-Quadruplex Secondary
859   Structure Obtained from Circular Dichroism Spectroscopy. *Angewandte Chemie*
860   *International Edition* **57**, 7171-7175

861   52.   Meier, M., Moya-Torres, A., Krahn, N. J., McDougall, M. D., Orriss, G. L., McRae, E.
862   K S., Booy, E. P., McEleney, K., Patel, T. R., McKenna, S. A., and Stetefeld, J. (2018)
863   Structure and hydrodynamics of a DNA G-quadruplex with a cytosine bulge. *Nucleic*
864   *Acids Research* **46**, 5319-5331

865   53.   Meier, M., Patel, T. R., Booy, E. P., Marushchak, O., Okun, N., Deo, S., Howard, R.,
866   McEleney, K., Harding, S. E., Stetefeld, J., and McKenna, S. A. (2013) Binding of G-
867   quadruplexes to the N-terminal recognition domain of the RNA helicase associated with
868   AU-rich element (RHAU). *The Journal of biological chemistry* **288**, 35014-35027

869   54.   Rambo, R. P., and Tainer, J. A. (2011) Characterizing flexible and intrinsically
870   unstructured biological macromolecules by SAS using the Porod-Debye law.
871   *Biopolymers* **95**, 559-571

872   55.   Svergun, D. (1992) Determination of the regularization parameter in indirect-transform
873   methods using perceptual criteria. *J Appl Crystallogr* **25**, 495-503

874   56.   Ariyo, E. O., Booy, E. P., Patel, T. R., Dzananovic, E., McRae, E. K., Meier, M.,
875   McEleney, K., Stetefeld, J., and McKenna, S. A. (2015) Biophysical Characterization of
876   G-Quadruplex Recognition in the PITX1 mRNA by the Specificity Domain of the
877   Helicase RHAU. *PloS one* **10**, e0144510

878   57.   Lattmann, S., Giri, B., Vaughn, J. P., Akman, S. A., and Nagamine, Y. (2010) Role of the
879   amino terminal RHAU-specific motif in the recognition and resolution of guanine
880   quadruplex-RNA by the DEAH-box RNA helicase RHAU. *Nucleic Acids Res* **38**, 6219-
881   6233

882   58.   Tippana, R., Chen, M. C., Demeshkina, N. A., Ferré-D'Amaré, A. R., and Myong, S.
883   (2019) RNA G-quadruplex is resolved by repetitive and ATP-dependent mechanism of
884   DHX36. *Nature Communications* **10**, 1855

885   59.   Guo, H., Zhou, T., Jiang, D., Cuconati, A., Xiao, G.-H., Block, T. M., and Guo, J.-T.
886   (2007) Regulation of hepatitis B virus replication by the phosphatidylinositol 3-kinase-
887   akt signal transduction pathway. *Journal of virology* **81**, 10072-10080

888   60.   Henderson, A., Wu, Y., Huang, Y. C., Chavez, E. A., Platt, J., Johnson, F. B., Brosh, R.
889   M., Jr., Sen, D., and Lansdorp, P. M. (2014) Detection of G-quadruplex DNA in
890   mammalian cells. *Nucleic Acids Res* **42**, 860-869

891   61.   Song, J., Perreault, J.-P., Topisirovic, I., and Richard, S. (2016) RNA G-quadruplexes
892   and their potential regulatory roles in translation. *Translation* **4**, e1244031

893   62.   Balasubramanian, S., Hurley, L. H., and Neidle, S. (2011) Targeting G-quadruplexes in
894   gene promoters: a novel anticancer strategy? *Nature Reviews Drug Discovery* **10**, 261

895   63.   Hayer, J., Jadeau, F., Deléage, G., Kay, A., Zoulim, F., and Combet, C. (2013) HBVdb: a
896   knowledge database for Hepatitis B Virus. *Nucleic Acids Research* **41**, D566-D570

897   64.   van der Kuyl, A. C., Zorgdrager, F., Hogema, B., Bakker, M., Jurriaans, S., Back, N. K.,
898   Berkhout, B., Zaaijer, H. L., and Cornelissen, M. (2013) High prevalence of hepatitis B
899   virus dual infection with genotypes A and G in HIV-1 infected men in Amsterdam, the
900   Netherlands, during 2000-2011. *BMC Infect Dis* **13**, 540
901   65.   Osiowy, C., Gordon, D., Borlang, J., Giles, E., and Villeneuve, J. P. (2008) Hepatitis B
902   virus genotype G epidemiology and co-infection with genotype A in Canada. *J Gen Virol*
903   **89**, 3009-3015
904   66.   Tanaka, Y., Sanchez, L. V., Sugiyama, M., Sakamoto, T., Kurbanov, F., Tatematsu, K.,
905   Roman, S., Takahashi, S., Shirai, T., Panduro, A., and Mizokami, M. (2008)
906   Characteristics of hepatitis B virus genotype G coinfected with genotype H in chimeric
907   mice carrying human hepatocytes. *Virology* **376**, 408-415
908   67.   Nowak, M. A., Bonhoeffer, S., Hill, A. M., Boehme, R., Thomas, H. C., and McDade, H.
909   (1996) Viral dynamics in hepatitis B virus infection. *Proceedings of the National*
910   *Academy of Sciences of the United States of America* **93**, 4398-4402
911   68.   Booy, E. P., Meier, M., Okun, N., Novakowski, S. K., Xiong, S., Stetefeld, J., and
912   McKenna, S. A. (2012) The RNA helicase RHAU (DHX36) unwinds a G4-quadruplex in
913   human telomerase RNA and promotes the formation of the P1 helix template boundary.
914   *Nucleic Acids Res* **40**, 4110-4124
915   69.   Creacy, S. D., Routh, E. D., Iwamoto, F., Nagamine, Y., Akman, S. A., and Vaughn, J. P.
916   (2008) G4 resolvase 1 binds both DNA and RNA tetramolecular quadruplex with high
917   affinity and is the major source of tetramolecular quadruplex G4-DNA and G4-RNA
918   resolving activity in HeLa cell lysates. *The Journal of biological chemistry* **283**, 34626-
919   34634
920   70.   Giri, B., Smaldino, P. J., Thys, R. G., Creacy, S. D., Routh, E. D., Hantgan, R. R.,
921   Lattmann, S., Nagamine, Y., Akman, S. A., and Vaughn, J. P. (2011) G4 resolvase 1
922   tightly binds and unwinds unimolecular G4-DNA. *Nucleic Acids Res* **39**, 7161-7178
923   71.   Sexton, A. N., and Collins, K. (2011) The 5' guanosine tracts of human telomerase RNA
924   are recognized by the G-quadruplex binding domain of the RNA helicase DHX36 and
925   function to increase RNA accumulation. *Molecular and cellular biology* **31**, 736-743
926   72.   Heddi, B., Cheong, V. V., Martadinata, H., and Phan, A. T. (2015) Insights into G-
927   quadruplex specific recognition by the DEAH-box helicase RHAU: Solution structure of
928   a peptide–quadruplex complex. *Proceedings of the National Academy of Sciences* **112**,
929   9608-9613
930   73.   Chen, M. C., Tippana, R., Demeshkina, N. A., Murat, P., Balasubramanian, S., Myong,
931   S., and Ferre-D'Amare, A. R. (2018) Structural basis of G-quadruplex unfolding by the
932   DEAH/RHA helicase DHX36. *Nature* **558**, 465-469
933   74.   Lau, K. C., Joshi, S. S., Gao, S., Giles, E., Swidinsky, K., van Marle, G., Bathe, O. F.,
934   Urbanski, S. J., Terrault, N. A., Burak, K. W., Osiowy, C., and Coffin, C. S. (2020)
935   Oncogenic HBV variants and integration are present in hepatic and lymphoid cells
936   derived from chronic HBV patients. *Cancer Lett* **480**, 39-47
937   75.   Mitra, B., Wang, J., Kim, E. S., Mao, R., Dong, M., Liu, Y., Zhang, J., and Guo, H.
938   (2019) Hepatitis B Virus Precore Protein p22 Inhibits Alpha Interferon Signaling by
939   Blocking STAT Nuclear Translocation. *Journal of virology* **93**, e00196-00119
940   76.   Mao, R., Nie, H., Cai, D., Zhang, J., Liu, H., Yan, R., Cuconati, A., Block, T. M., Guo, J.
941   T., and Guo, H. (2013) Inhibition of hepatitis B virus replication by the host zinc finger
942   antiviral protein. *PLoS Pathog* **9**, e1003494

77. Thompson, J. D., Higgins, D. G., and Gibson, T. J. (1994) CLUSTAL W: improving the
sensitivity of progressive multiple sequence alignment through sequence weighting,
position-specific gap penalties and weight matrix choice. *Nucleic acids research* **22**,
4673-4680
78. Booy, E. P., Howard, R., Marushchak, O., Ariyo, E. O., Meier, M., Novakowski, S. K.,
Deo, S. R., Dzananovic, E., Stetefeld, J., and McKenna, S. A. (2014) The RNA helicase
RHAU (DHX36) suppresses expression of the transcription factor PITX1. *Nucleic Acids*
Res **42**, 3346-3361
79. Mrozowich, T., Meier-Stephenson, V., and Patel, T. (2019) *Microscale thermophoresis:*
warming up to a new biomolecular interaction technique,
80. Reuten, R., Patel, T. R., McDougall, M., Rama, N., Nikodemus, D., Gibert, B., Delcros,
J. G., Prein, C., Meier, M., Metzger, S., Zhou, Z., Kaltenberg, J., McKee, K. K., Bald, T.,
Tuting, T., Zigrino, P., Djonov, V., Bloch, W., Clausen-Schaumann, H., Poschl, E.,
Yurchenco, P. D., Ehrbar, M., Mehlen, P., Stetefeld, J., and Koch, M. (2016) Structural
decoding of netrin-4 reveals a regulatory function towards mature basement membranes.
Nat Commun **7**, 13515
81. Rambo, R. (2017) ScÅtter, a JAVA-based applcation for basic analysis of SAXS
datasets. 3.1R ed Ed.
82. Franke, D., Petoukhov, M. V., Konarev, P. V., Panjkovich, A., Tuukkanen, A., Mertens,
H. D. T., Kikhney, A. G., Hajizadeh, N. R., Franklin, J. M., Jeffries, C. M., and Svergun,
D. I. (2017) ATSAS 2.8: a comprehensive data analysis suite for small-angle scattering
from macromolecular solutions. *Journal of applied crystallography* **50**, 1212-1225
83. Franke, D., and Svergun, D. I. (2009) DAMMIF, a program for rapid ab-initio shape
determination in small-angle scattering. *Journal of applied crystallography* **42**, 342-346
84. Volkov, V. V., and Svergun, D. I. (2003) Uniqueness of ab initio shape determination in
small-angle scattering. *J Appl Crystallogr* **36**, 860-864
85. Patel, T. R., Reuten, R., Xiong, S., Meier, M., Winzor, D. J., Koch, M., and Stetefeld, J.
(2012) Determination of a molecular shape for netrin-4 from hydrodynamic and small
angle X-ray scattering measurements. *Matrix biology : journal of the International*
Society for Matrix Biology **31**, 135-140
86. Gao, S., Duan, Z. P., Chen, Y., van der Meer, F., Lee, S. S., Osiowy, C., van Marle, G.,
and Coffin, C. S. (2017) Compartmental HBV evolution and replication in liver and
extrahepatic sites after nucleos/tide analogue therapy in chronic hepatitis B carriers. *J*
Clin Virol **94**, 8-14
87. Lau, K. C. K., Osiowy, C., Giles, E., Lusina, B., van Marle, G., Burak, K. W., and
Coffin, C. S. (2018) Deep sequencing shows low-level oncogenic hepatitis B virus
variants persists post-liver transplant despite potent anti-HBV prophylaxis. *J Viral Hepat*
25, 724-732
88. van Bommel, F., Bartens, A., Mysickova, A., Hofmann, J., Kruger, D. H., Berg, T., and
Edelmann, A. (2015) Serum hepatitis B virus RNA levels as an early predictor of
hepatitis B envelope antigen seroconversion during treatment with polymerase inhibitors.
Hepatology **61**, 66-76
89. Guinier, A., and Fourner, G. (1955) *Small angle scattering of X-rays*, Wiley, New York
90. Svergun, D. I. (1999) Restoring low resolution structure of biological macromolecules
from solution scattering using simulated annealing. *Biophysical journal* **76**, 2879-2886
988

    TABLES

990   **Table 1**: Analysis of small-angle X-ray scattering data for HBV wild-type (*wt*) and mutant (*mut*)

991   core promoter oligomer samples

	wt	*G1738A*	*G1748A*	*G1738/1748A*
M_w (kDa)	7.30	7.28	7.28	7.26
$I(0)^{\#}$	$0.0035 \pm 1.1 \times 10^{-5}$	$0.0041 \pm 1.3 \times 10^{-5}$	$0.0031 \pm 1.6 \times 10^{-5}$	$0.0037 \pm 2.1 \times 10^{-5}$
q.Rg range	0.24 – 1.30	0.23 – 1.30	0.23 – 1.30	0.23 – 1.29
R_g (Å)$^{\#}$	18.37 ± 0.11	19.10 ± 0.11	20.10 ± 0.19	20.34 ± 0.22
$I(0)^{\#\Delta}$	$0.0034 \pm 0.7 \times 10^{-5}$	$0.0039 \pm 1.0 \times 10^{-5}$	$0.0030 \pm 1.2 \times 10^{-5}$	$0.0036 \pm 1.7 \times 10^{-5}$
R_g (Å)$^{\Delta}$	17.18 ± 0.03	18.17 ± 0.04	19.77 ± 0.09	20.43 ± 0.10
D_{max} (Å) $^{\Delta}$	46.85	53.14	60	60.44
χ^{2*}	~1.80	~1.50	~1.50	~1.20
NSD*	0.55 ± 0.01	0.58 ± 0.03	0.58 ± 0.01	0.59 ± 0.01

The molecular weight (M_w) values were calculated using nucleotide sequences.
[#] - obtained from Guinier analysis (89).
[Δ] - determined using $P(r)$ analysis using the GNOM program (55).
[*] - values derived from DAMMIN (90) and DAMAVER (84) analysis.
The error calculation for each SAXS data analysis step is described by individual packages
implemented in ATSAS suite (55, 84, 89, 90).

Table 2: The dissociation constants calculated for DHX36 53-105 and G-quadruplex

variants

Oligomer	K_d (nM)
wt	76.81 ± 12.21
G1738A	14.17 ± 12.76
G1748A	132.14 ± 14.84
G1738/1748A	65.50 ± 7.74

1000   *The error represents standard deviation of 3 independent experiments.

Figure 1

A

	1732 C	T	G	G	G	A	G	G	A	G	T	T	G	G	G	G	G	1748 A	G	G	A	G	1754 A
A	0.50%	0.04%	0.10%	0.10%	0.10%	99.90%	0.20%	0.10%	99.40%	0.90%	0.10%	0.05%	2.00%	0.02%	0.10%	0.10%	0.10%	99.70%	0.30%	0.10%	99.70%	0.70%	82.70%
C	74.50%	0.10%	0.01%	0.03%	0.00%	0.00%	0.01%	0.01%	0.01%	0.02%	10.40%	3.10%	0.01%	0.02%	0.00%	0.01%	0.03%	0.10%	0.00%	0.02%	0.01%	0.10%	2.60%
G	24.90%	0.03%	99.90%	99.80%	99.90%	0.10%	99.80%	99.90%	0.10%	98.80%	0.10%	0.10%	97.40%	100.00%	99.90%	99.80%	99.80%	0.10%	99.50%	99.90%	0.10%	99.10%	12.90%
T	0.10%	99.80%	0.00%	0.01%	0.00%	0.02%	0.00%	0.00%	0.50%	0.30%	89.40%	96.80%	0.50%	0.01%	0.02%	0.10%	0.02%	0.10%	0.10%	0.00%	0.20%	0.10%	1.90%

B

wt	CTGGGA**G**GAGCTGGGG**G**AGGAGA
G1738A	CTGGGA**A**GAGCTGGGG**G**AGGAGA
G1748A	CTGGGA**G**GAGCTGGGG**A**AGGAGA
G1738/1748A	CTGGGA**A**GAGCTGGGG**A**AGGAGA

Figure 2

Figure 3A

Figure 3B

wt	*G1738A*	*G1748A*	*G1738A/1748A*
46.85 Å	53.14 Å	60.00 Å	60.44 Å

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

A

Core transcripts

(-)-DNA

2.4kb

2.1kb

S

P

C

X

0.7kb

Poly-A

3.5kb

P

Total RNA transcripts

B

Relative RNA

Day post transfection

- wt
- G1738A
- G1748A
- G1738/1748A

C

Relative RNA

Day post transfection

- wt
- G1738A
- G1748A
- G1738/1748A