Polycomb suppresses a female gene regulatory network to ensure testicular differentiation

So Maezawa[1, 2, 3, 4*], Masashi Yukawa[2, 5, 9], Kazuteru Hasegawa[1, 2, 10], Ryo Sugiyama[3], Mengwen Hu[1, 2, 6], Miguel Vidal[7], Haruhiko Koseki[8], Artem Barski[2, 5], Tony DeFalco[1, 2], and Satoshi H. Namekawa[1, 2, 6*]

[1] Division of Reproductive Sciences, Division of Developmental Biology, Perinatal Institute, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio, 45229, USA

[2] Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, Ohio, 45229, USA

[3] Department of Animal Science and Biotechnology, School of Veterinary Medicine, Azabu University, Sagamihara, Kanagawa 252-5201, Japan

[4] Faculty of Science and Technology, Department of Applied Biological Science, Tokyo University of Science, Chiba 278-8510, Japan.

[5] Division of Allergy and Immunology, Division of Human Genetics, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio, 45229, USA

[6] Department of Microbiology and Molecular Genetics, University of California, Davis, California, 95616, USA

[7] Centro de Investigaciones Biológicas Margarita Salas, Department of Cellular and Molecular Biology, Madrid 28040, Spain

[8] Developmental Genetics Laboratory, RIKEN Center for Allergy and Immunology, Yokohama, Kanagawa, Japan

[9] Present address: Department of Chemical Pathology, The Chinese University of Hong Kong, Prince of Wales Hospital, Sha Tin, New Territories, Hong Kong

[10] Present address: Stanford Cancer Institute, Stanford University School of Medicine, Stanford,

CA 94305, USA

*Correspondence: E-mail: s-maezawa@rs.tus.ac.jp; snamekawa@ucdavis.edu

Abstract

Gonadal sex determination is controlled by the support cells of testes and ovaries. In testes, the

epigenetic mechanism that maintains cellular memory to suppress female sexual differentiation

remains unknown. Here, we show that Polycomb suppresses a female gene regulatory network in

Sertoli cells, the specific support cells for postnatal testes. Through genetic ablation, we removed

Polycomb repressive complex 1 (PRC1) from embryonic Sertoli cells after sex determination.

PRC1-depleted postnatal Sertoli cells exhibited defective proliferation and cell death, leading to

the degeneration of adult testes. In adult Sertoli cells, PRC1 suppressed the specific, critical genes

required for granulosa cells, the support cells of ovaries, thereby inactivating the female gene

regulatory network. The underlying chromatin of female genes was coated with Polycomb-

mediated repressive modifications: PRC1-mediated H2AK119ub and PRC2-mediated H3K27me3.

Taken together, we identify a critical mechanism centered on Polycomb that maintains the male

fate in adult testes.

Introduction

In mammals, gonadal sex determination takes place in the bipotential somatic cell

precursors of male Sertoli cells and female granulosa cells in embryos (Capel, 2017; Swain &

Lovell-Badge, 1999; Wilhelm, Palmer, & Koopman, 2007). Sertoli cells are the first somatic cells

to differentiate in the XY gonad. In testes, Sertoli cells function as a regulatory hub for both

differentiation and survival of germ cells, thereby determining male sexual fate (Svingen &

Koopman, 2013). The mechanisms maintaining the male cellular identity of Sertoli cells are

fundamental for adult testicular functions, including spermatogenesis and hormone production.

At the time of sex determination in embryos, the commitment to the male fate is triggered

by the expression of the Y-linked *Sry* gene, and, subsequently, the female fate is suppressed (Capel,

2017; Swain & Lovell-Badge, 1999; Wilhelm et al., 2007). Distinct gene regulatory networks

promote the male or female fate and are regulated by strong feedback loops that antagonize each

other, canalizing one fate from the other (Capel, 2017). Sexual fate is interchangeable even after

the initial commitment to Sertoli cells or granulosa cells with the removal of specific, critical

transcription factors. The loss of *Dmrt1* in Sertoli cells leads to derepression of *Foxl2*, a master

regulator of granulosa cell fate, and transdifferentiation of cell fate from Sertoli to granulosa cells

(Matson et al., 2011; Matson & Zarkower, 2012). On the other hand, the loss of *Foxl2* in granulosa

cells leads to derepression of the male gene network and transdifferentiation of cell fate from

granulosa to Sertoli cells (Uhlenhaut et al., 2009). These findings, together with follow-up studies

(Li et al., 2017; Lindeman et al., 2015; Minkina et al., 2014; Nicol et al., 2018; Zhao, Svingen, Ng,

& Koopman, 2015), suggest that active repression of the alternate sexual fate is important for both

testicular and ovarian function, even in adult life.

Epigenetic silencing mechanisms serve as molecular switches for the sex determination of

bipotential somatic cell precursors. The deletion of a Polycomb gene *Cbx2* results in male to female

reversal (Katoh-Fukui et al., 1998), which is mediated through suppression of genes required for

the female fate (Garcia-Moreno et al., 2019). Additionally, regulation of H3K9 methylation is

important for the male sexual fate (Kuroki et al., 2013). Although these studies highlight the

importance of epigenetic mechanisms for initial sex determination, the epigenetic mechanisms by

which male cellular identity is maintained through cell divisions and the proliferation of Sertoli

cells remain to be determined.

Polycomb proteins suppress non-lineage-specific genes and define cellular identities of

each lineage in stem cells and in development (Aloia, Di Stefano, & Di Croce, 2013; Ringrose &

Paro, 2007; Simon & Kingston, 2013). In this study, we show that Polycomb suppresses the female

gene regulatory network in postnatal Sertoli cells, thereby promoting the male gene regulatory

network to ensure male cell fate. We generated loss-of-function mouse models of Polycomb

repressive complex 1 (PRC1) in Sertoli cells after initial sex determination. We show that PRC1 is

required for the proliferation of Sertoli cells, as well as the suppression of non-lineage-specific

genes and the female gene regulatory network in Sertoli cells. Taken together, we identify a critical

mechanism centered on Polycomb that maintains male fate in adult testes.

Results

PRC1 in Sertoli cells is required for the maintenance of spermatogenesis. In postnatal Sertoli

cells (which are detected by the Sertoli cell marker GATA4), RNF2 is highly expressed and the

RNF2-mediated epigenetic mark H2AK119ub is abundant (Figure 1A), which suggests PRC1

functions in these cells. To determine the function of PRC1, we generated a PRC1 loss-of-function

mouse model by removing two redundant catalytic subunits, RNF2 and RING1 (Endoh et al., 2012).

We generated a conditional deletion of *Rnf2* (*Rnf2*cKO) using *Amh*-Cre, which is expressed

specifically in Sertoli cells after embryonic day 14.5 (E14.5) (Holdcraft & Braun, 2004), in a

background of *Ring1*-knockout (KO) mice (*Amh*-Cre; *Rnf2*cKO; *Ring1*-KO: termed PRC1[Amh-]

^{Cre}cKO: PRC1AcKO). Although RNF2 appears to be the most active component in the

heterodimeric E3 ligases of PRC1, the RNF2 paralog RING1 can partially compensate for the loss

of RNF2 (Endoh et al., 2012). Therefore, we made a conditional deletion of *Rnf2* in a background

of *Ring1*-KO mice, which are viable and do not have fertility defects (del Mar Lorente et al., 2000).

This strategy enabled us to define the function of RNF2 without compensation from RING1, while

also representing a "complete" loss-of-function of PRC1 as shown in testicular germ cells

(Maezawa et al., 2017) and in other biological contexts (Endoh et al., 2012; Posfai et al., 2012;

Yokobayashi et al., 2013). Since PRC1 has various components, including CBX2 (Gao et al., 2012),

this strategy allows us to determine the global function of PRC1. At the same time, the use of *Amh*-

Cre allowed us to test the function of PRC1, specifically after the completion of sex determination

at E12.5.

108   PRC1AcKO males have smaller testes compared with littermate controls that harbored

floxed alleles for *Rnf2* on a *Ring1*-KO background without *Amh*-Cre (termed PRC1 control:

PRC1ctrl: Figure 1B and C). We confirmed efficient *Amh*-Cre-mediated recombination by

observing depletion of the RNF2-mediated mark, H2AK119ub, in GATA4$^+$ Sertoli cells of

PRC1AcKO testes at E15.5 (> 95 % efficiency: Figure 1D) and at postnatal day 7 (P7: Figure 1E).

In 6 week-old PRC1AcKO testes, while the tubules with H2AK119ub$^+$ Sertoli cells (escaped Cre-

mediated deletion) appear to have normal morphology, we frequently observed disorganization of

testicular tubules that contain H2AK119ub$^-$ Sertoli cells (underwent Cre-mediated deletion:

arrowheads, Figure 1- figure supplement 1), suggesting a critical function of PRC1 in Sertoli cells

in the organization of testicular tubules. This mosaic pattern is presumably due to incomplete Cre-

mediated recombination, as the Sertoli cells that escaped Cre-mediated recombination might have

repopulated the testes. Consistent with this interpretation, PRC1AcKO males are subfertile, and 9

out of 15 wild-type females mated with 8-11-week old PRC1AcKO males gave birth (Figure 1F)

at comparable litter sizes (Figure 1G). Interestingly, fecundity decreased in aged PRC1AcKO males

(5 months old), as litter sizes were smaller as compared to controls (Figure 1G). To further examine

the phenotype, we measured the blood levels of three hormones critical for testicular homeostasis:

although the levels of testosterone and estradiol were comparable between cKO and control mice,

follicle-stimulating hormone was increased in mutants (Figure 1H). As follicle stimulating

hormone levels are regulated by a feedback mechanism involving Sertoli cells (Oduwole, Peltoketo,

& Huhtaniemi, 2018), we infer that the dysfunction of Sertoli cells and testicular degeneration

caused increased follicle stimulating hormone levels to recover Sertoli cell function.

To determine the cause of testicular degeneration, we next examined the proliferation of

Sertoli cells. In normal mouse development, Sertoli cells proliferate in fetal and neonatal testes

until approximately 2 weeks after birth, and the number of Sertoli cells in adult testes determines

both testis size and daily sperm production (Sharpe, McKinnell, Kivlin, & Fisher, 2003). In P7

control testes, GATA4$^+$ Sertoli cells occasionally co-expressed an S phase marker, PCNA, and

another marker of active cell cycle, Ki67 (Figure 2A), consistent with the active proliferation of

Sertoli cells. However, in PRC1AcKO testes, GATA4$^+$ Sertoli cells were largely devoid of PCNA

and Ki67 (Figure 2A), suggesting impaired proliferation of PRC1AcKO Sertoli cells. Next, we

independently confirmed the proliferation phenotype by using EdU labeling of actively

proliferating cells. EdU was abdominally administrated to P7 mice, and testicular sections were

examined the following day. While GATA4$^+$ Sertoli cells were occasionally EdU$^+$ (approximately

one-fouth) in control testes , GATA4$^+$ Sertoli cells were devoid of EdU signal in PRC1AcKO testes

(Figure 2B). Additional labeling of H2AK119ub confirmed the loss of PRC1 function in GATA4$^+$

Sertoli cells from PRC1AcKO testes (Figure 2B). From these results, we conclude that the loss of

PRC1 disrupts the proliferation of Sertoli cells. This is in contrast with PRC1's function in testicular

germ cells, in which the loss of PRC1 does not affect proliferation but instead causes apoptotic cell

death (Maezawa et al., 2017). This difference suggests a unique function of PRC1 in Sertoli cells

that is distinct from its function in germ cells.

In Sertoli cells, PRC1 suppresses genes required for granulosa cells. We next sought to

determine the genes regulated by PRC1 in Sertoli cells. In PRC1AcKO testes, some Sertoli cells

escaped *Amh*-Cre-mediated recombination. Thus, it was difficult to specifically isolate Sertoli cells

that underwent PRC1 depletion. To precisely determine the function of PRC1 in gene regulation in

Sertoli cells, we used an alternative strategy: we isolated Sertoli cells from a mouse line in which

conditional deletion of PRC1 can be induced by tamoxifen-inducible Cre-mediated recombination

under the control of the endogenous ROSA26 promoter (*ROSA26*-Cre^{ERT}; *Rnf2*$^{floxed/floxed}$; *Ring1*-

KO: termed PRC1$^{ROSA26-CreERT}$cKO: PRC1RcKO). After isolating Sertoli cells from P7 testes, we

cultured Sertoli cells for 4 days in the presence of 4-hydroxytamoxifen (4-OHT), and performed

RNA-sequencing (RNA-seq: Figure 3A). As a control, we isolated Sertoli cells from control mice

(*Rnf2*$^{floxed/floxed}$; *Ring1*-KO) and cultured them in the same 4-OHT conditions. We performed RNA-

seq for two independent biological replicates and confirmed reproducibility between biological

replicates (Figure 3- figure supplement 1A).

Our RNA-seq analyses demonstrated that 338 genes were upregulated in PRC1RcKO

Sertoli cells as compared to controls, while 307 genes were downregulated in PRC1RcKO Sertoli

cells (Figure 3A and B). Gene ontology (GO) analysis showed that upregulated genes were enriched

for functions in neural differentiation, skeletal system, and cell adhesion (Figure 3D). These

categories suggest that PRC1 suppressed expression of non-lineage-specific genes in Sertoli cells.

On the other hand, GO analysis revealed that downregulated genes were enriched for functions in

the cell cycle and M phase (Figure 3D). This result is in accord with the cell cycle arrest we found

in PRC1AcKO Sertoli cells (Figure 2).

Since we anticipated suppression of genes required for granulosa cells by PRC1 in Sertoli

cells, we next investigated the expression level of key genes required for granulosa cells. In

PRC1RcKO Sertoli cells, genes required for female sexual development were upregulated: these

genes included *Rspo1,* an activator of the Wnt pathway (Chassot et al., 2008), and *Foxl2,* a key

transcription factor for granulosa cells (Schmidt et al., 2004) (Figure 3E). Importantly, these female

genes suppress the male fate, and the loss of these genes leads to female-to-male sex reversal

(Ottolenghi et al., 2007; Parma et al., 2006; Schmidt et al., 2004). Consistent with the antagonistic

function of these female genes with the male pathway, key sex determination genes for the male

pathway were downregulated in PRC1RcKO Sertoli cells: these genes included *Sox9,* an

evolutionarily conserved gene for sex determination, which directs the male pathway downstream

of *Sry* (Vidal, Chaboissier, de Rooij, & Schedl, 2001), and *Dmrt1* (Raymond, Murphy, O'Sullivan,

Bardwell, & Zarkower, 2000; Raymond et al., 1998), male-determining signalling (Figure 3E).

185   To determine whether the suppression of female genes was directly regulated by PRC1, we

performed chromatin immunoprecipitation sequencing (ChIP-seq) of PRC1-mediated H2AK119ub

in isolated wild-type Sertoli cells. We further performed ChIP-seq of H3K27me3 in Sertoli cells

since PRC2-mediated H3K27me3 is regulated by PRC1 and its mediated mark H2A119ub

(Blackledge et al., 2014; Cooper et al., 2014). We performed ChIP-seq for two independent

biological replicates and confirmed the reproducibility between biological replicates (Figure 3-

figure supplement 1B). We confirmed the enrichment of H2AK119ub and H3K27me3 around

transcription start sites (TSSs) of *Rspo1* and *Foxl2* (Figure 3F). Compared to the enrichment of

H2AK119ub on these female genes, enrichment of H2AK119ub was relatively low on the TSSs of

the male genes, *Sox9* and *Dmrt1*. Therefore, we conclude that PRC1 directly binds and suppresses

Rspo1 and *Foxl2*.

197   Track views of these ChIP-seq data show that all three marks were enriched at *Foxl2* and

Rspo1 loci (Figure 4A). Furthermore, enrichment of these three marks was found in other loci such

as *Foxf2*, the mutation of which appears in patients with disorders of sex development (Jochumsen

et al., 2008), and *Hoxd13*, which is involved in female reproductive tract development (Du &

Taylor, 2015). These results suggest that PRC1 works with PRC2 to suppress female genes as well

as developmental regulators such as *Fox* and *Hox* genes, which are the classical targets of

Polycomb-mediated gene repression (Lee et al., 2006).

205   To determine the features of genome-wide gene repression mediated by Polycomb

complexes, we analyzed the enrichment of H2AK119ub, H3K27me3, and RNF2 on the upregulated

genes in PRC1RcKO Sertoli cells. H2AK119ub, H3K27me3, and RNF2 were all significantly

enriched on the TSSs of upregulated genes in PRC1RcKO Sertoli cells as compared to all genes in

the genome (Figure 4B). Additional enrichment analysis confirmed the co-enrichment of

H2AK119ub and H3K27me3 (Figure 4C, upper panels) as well as co-enrichment of H2AK119ub

and RNF2 (Figure 4C, lower panels) on upregulated genes in PRC1RcKO Sertoli cells.

Furthermore, average tag density analysis confirmed that enrichment of H2AK119ub on

upregulated genes in PRC1RcKO Sertoli cells occured on upstream regions, gene bodies, and

downstream regions with the highest enrichment near TSSs (Figure 4D). We found a similar

distribution of H3K27me3 around the gene bodies of upregulated genes in PRC1RcKO Sertoli cells

(Figure 4- figure supplement 1). Together, these results confirmed the genome-wide, global

functions of PRC1 in the direct regulation of gene repression in Sertoli cells.

Polycomb globally inactivates the female gene regulatory network in postnatal Sertoli cells.

Previous studies have suggested that sex determination is canalized by the interconnected,

antagonistic network of genes both in males and in females that are controlled by feedback

mechanisms (Capel, 2017). Since Polycomb is implicated in the maintenance of sex-specific gene

regulatory networks through the PRC2-mediated mark H3K27me3 (Garcia-Moreno et al., 2019),

we hypothesized that PRC1 globally inactivates the female gene regulatory network in postnatal

Sertoli cells. Since male- and female-specific gene networks are regulated immediately after sex

determination during fetal stages (Jameson et al., 2012), we reasoned that female gene network

suppression in fetal Sertoli cells is maintained by PRC1 in postnatal Sertoli cells. To test this

hypothesis, we examined the expression profiles of specifically expressed genes in E13.5 granulosa

cells (Jameson et al., 2012) (Figure 5- figure supplement 1), termed "pre-granulosa genes" in

postnatal Sertoli cells. We found that pre-granulosa genes were upregulated in PRC1RcKO

postnatal Sertoli cells as compared to other genes (Figure 5A). On the other hand, specifically

expressed genes in E13.5 Sertoli cells (Jameson et al., 2012) (Figure 5- figure supplement 1),

termed "pre-Sertoli genes," were downregulated in PRC1RcKO postnatal Sertoli cells as compared

to other genes (Figure 5A). We further examined the correlation of each gene and found that pre-

granulosa genes were highly correlated with upregulated genes in PRC1RcKO postnatal Sertoli

cells, while pre-Sertoli genes were highly correlated with downregulated genes in PRC1RcKO

postnatal Sertoli cells (Figure 5B). These results suggest that PRC1 maintains suppression of the

female gene regulatory network, which is initiated at the time of sex determination and is

maintained throughout the development of postnatal Sertoli cells.

We next sought to determine whether PRC1 directly suppresses the female gene regulatory

network in postnatal Sertoli cells. H2AK119ub is significantly enriched on TSSs of pre-granulosa

genes compared to other genes and pre-Sertoli genes (Figure 5C). Among pre-granulosa genes,

enrichment of H2AK119ub is positively correlated with upregulated genes in PRC1RcKO

postnatal Sertoli cells (Figure 5D). We further identified the enrichment of H3K27me3 on TSSs of

pre-granulosa genes in postnatal Sertoli cells (Figure 5- figure supplement 2A), and the correlation

between H3K27me3 and upregulated genes on the pre-granulosa genes in PRC1RcKO postnatal

Sertoli cells (Figure 5- figure supplement 2B). Together, we conclude that PRC1 globally

inactivates the female gene regulatory network in postnatal Sertoli cells.

Discussion

In this study, we demonstrated that PRC1 is required for proliferation of Sertoli cells and suppression of the non-lineage-specific gene expression program and the female gene regulatory network. Among these functions, we infer that suppression of the female gene regulatory network is the key mechanism to ensure male fate in addition to canonical functions of PRC1, which controls proliferation and suppression of non-lineage-specific gene expression programs. Although we did not observe complete infertility using *Amh*-Cre, presumably due to the repopulation of Sertoli cells that escaped Cre-mediated recombination, smaller testis size and abnormal tubule organization of cKO testes (Figure 1) suggest that PRC1 is critical for physiological functions of Sertoli cells. Below, we discuss two molecular aspects underlying these physiological phenotypes: control of proliferation and suppression of the female gene regulatory network.

The proliferation of Sertoli cells is a critical determinant of testicular functions because the number of Sertoli cells defines testicular functions (Sharpe et al., 2003). Rapid proliferation is a prominent feature of juvenile Sertoli cells. In general, Polycomb proteins are associated with the cell cycle checkpoint by directly suppressing the tumor suppressor locus *Cdkn2a/Ink4a/Arf* (Jacobs, Kieboom, Marino, DePinho, & van Lohuizen, 1999), which functions as a barrier to cancer transformation (Serrano et al., 1996). Therefore, enhanced Polycomb activity is a frequent feature of human tumors. We found that *Cdkn2a* was derepressed in PRC1RcKO Sertoli cells (approximately 2.5-fold upregulation: PRKM value for PRC1RcKO: 32.1 v.s. PRC1ctrl: 13.0). Therefore, our results suggest that PRC1 promotes the rapid proliferation of Sertoli cells by suppressing *Cdkn2a.* The function of PRC1 in Sertoli cell proliferation is distinct from Polycomb functions in testicular germ cells, where PRC1 depletion does not alter the proliferation of germ cells (Maezawa et al., 2017). This may be due to the fact that PRC1 and PRC2 are not required for suppression of *Cdkn2a* in germ cells (Maezawa et al., 2017; Mu, Starmer, Fedoriw, Yee, & Magnuson, 2014)*.* The functional difference in testicular germ cells and Sertoli cells highlights the

277   context-dependent functions of Polycomb proteins. However, a recent study revealed a novel

278   activity by which PcGs can regulate cell proliferation through DNA replication independently of

279   *Cdkn2a* (Piunti et al., 2014). Therefore, determining the detailed molecular mechanisms by which

280   PRC1 controls the proliferation of Sertoli cells will be important for future studies.

282   Another critical function of PRC1 in Sertoli cells is the suppression of the female gene

regulatory network. DMRT1 is a critical transcription factor that suppresses the expression of

female genes (Matson et al., 2011). While more than 10-fold upregulation of the female genes

(*Rspo1* and *Foxl2*) was observed for *Dmrt1* mutants (Matson et al., 2011), the degree of

upregulation of these genes was modest in PRC1RcKO Sertoli cells (Figure 3). This finding,

combined with genetic evidence, indicates that DMRT1 could be the direct regulator of suppression

of female genes, while PRC1-mediated mechanisms could be a maintenance mechanism in

response to the primary silencing mechanisms determined by DMRT1. Another possibility is

compensation by PRC1-independent suppression mechanisms. While a portion of PRC2-mediated

H3K27me3 is regulated by variant PRC1 (Blackledge et al., 2014; Cooper et al., 2014), another

portion of H3K27me3, mediated by canonical PRC1 and PRC2 complexes, is not downstream of

PRC1 (Laugesen, Hojfeldt, & Helin, 2019). These PRC2-mediated mechanisms or other silencing

machinery may be responsible for the suppression of female genes.

296   The notion that Polycomb regulates the female gene regulatory network has been supported

by other recent evidence. In bipotential precursor cells, genes involved in sex determination are

marked with bivalent chromatin domains (Garcia-Moreno et al., 2019) that are prevalent in

pluripotent stem cells and in germ cells (Bernstein et al., 2006; Hammoud et al., 2009; Lesch,

Dokshin, Young, McCarrey, & Page, 2013; Maezawa, Hasegawa, et al., 2018; Maezawa, Yukawa,

Alavattam, Barski, & Namekawa, 2018; Sin, Kartashov, Hasegawa, Barski, & Namekawa, 2015).

Maintenance of the male fate was explained by the persistence of H3K27me3 on silent female

genes in Sertoli cells (Garcia-Moreno et al., 2019). Consistent with PRC1's function found in the

current study, Polycomb-mediated silencing may globally suppress the female gene regulatory

network. We found that H2AK119ub was largely associated with this group of female genes

(Figure 5). Therefore, it would be interesting to speculate that antagonistic male and female

networks can be directly coordinated by Polycomb protein functions, including the strong feedback

mechanism underlying both networks. These possibilities raise several outstanding questions to be

addressed in future studies. What are the functions of another Polycomb complex, PRC2, and of

each Polycomb complex component, including CBX2, in postnatal Sertoli cells? What is the

function of Polycomb in female granulosa cells, especially in the suppression of the male gene

regulatory network? Does Polycomb underlie the feedback regulation of each network to define

sexual identity? Our current study provides a foundation to explore these questions.

Methods

Animals

Generation of mutant *Ring1* and *Rnf2* floxed alleles were previously reported (Cales et al., 2008).

Amh-Cre transgenic mice were purchased from The Jackson Laboratory (Stock No: 007915)

(Holdcraft & Braun, 2004). *Rosa-Cre ERT* mice were purchased from The Jackson Laboratory

(Stock No: 008463) (Ventura et al., 2007). A minimum of three independent mice were analyzed

for each experiment. All of the animals were handled according to approved institutional animal

care and use committee (IACUC) protocols (#IACUC2018-0040) of Cincinnati Children's Hospital

Medical Center. Fertility tests were performed with 6-weeks old CD1 female mice (purchased from

Charles river). At least 2 female mice were bred with a male mouse for 2 weeks, and the fertility

was evaluated by the ratio of pregnant to total female mice and number of pups.

Ligand test

Blood samples were collected from C57BL/6N mice aged 8−11 weeks. Serum was separated immediately and stored at -20°C. Hormone assays, including testosterone, estradiol, and follicle stimulating hormone, were performed by the Center for Research in Reproduction at the University of Virginia.

Sertoli cell isolation

Sertoli cells were isolated as previously described with minor modifications (Chang, Lee-Chang, Panneerdoss, MacLean, & Rao, 2011) and collected from C57BL/6N mice aged 6–8 days. Testes were collected in a 24-well plate in Dulbecco's Modified Eagle Medium (DMEM) supplemented with GlutaMax (Thermo Fisher Scientific), non-essential amino acids (NEAA) (Thermo Fisher Scientific), and penicillin and streptomycin (Thermo Fisher Scientific). After removing the *tunica albuginea* membrane, testes were digested with collagenase (1 mg/ml) at 34°C for 20 min to remove interstitial cells, then centrifuged at 188×*g* for 5 min. Tubules were washed with medium and then digested with trypsin (2.5 mg/ml) at 34°C for 20 min to obtain a single-cell suspension. To remove KIT-positive spermatogonia, cells were washed with magnetic cell-sorting (MACS) buffer (PBS supplemented with 0.5% BSA and 5 mM EDTA) and incubated with CD117 (KIT) MicroBeads (Miltenyi Biotec) on ice for 20 min. Cells were separated by autoMACS Pro Separator (Miltenyi Biotec) with the program "possel." Cells in the flow-through fraction were washed with MACS buffer and incubated with CD90.2 (THY1) MicroBeads (Miltenyi Biotec) on ice for 20 min to remove THY1-positive spermatogonia. Cells were separated by autoMACS Pro Separator (Miltenyi Biotec) with the program "posseld." Cells in the flow-through fraction were washed and plated in a 6-well plate for 1 h in the medium supplemented with 10% fetal bovine serum, which promotes adhesion of Sertoli cells. Purity was confirmed by immunostaining.

For PRC1RcKO, cells were cultured for 4 days with 1 µM 4-OHT in Dulbecco's Modified

Eagle Medium (DMEM) supplemented with GlutaMax (Thermo Fisher Scientific), non-essential

amino acids (NEAA) (Thermo Fisher Scientific), and penicillin and streptomycin (Thermo Fisher

Scientific). The same medium was replaced 2 days after the initiation of the culture.

Histological analysis and germ cell slide preparation

For the preparation of testicular paraffin blocks, testes were fixed with 4% paraformaldehyde (PFA)

overnight at 4°C with gentle inverting. Testes were dehydrated and embedded in paraffin. For

histological analysis, 7 µm-thick paraffin sections were deparaffinized and stained with

hematoxylin and eosin. For immunofluorescence analysis of testicular sections, antigen retrieval

was performed by boiling the slides in target retrieval solution (DAKO) for 10 min and letting the

solution cool for 30 min. Sections were blocked with Blocking One Histo (Nacalai) for 1 h at room

temperature and then incubated with primary antibodies overnight at 4°C. The resulting signals

were detected by incubation with secondary antibodies conjugated to fluorophores (Thermo Fisher

Scientific). Sections were counterstained with DAPI. Images were obtained via a laser scanning

confocal microscope A1R (Nikon) and processed with NIS-Elements (Nikon) and ImageJ

(National Institutes of Health) (Schneider, Rasband, & Eliceiri, 2012).

ChIP-sequencing, RNA-sequencing, and data analysis

RNA-seq analyses were performed in the BioWardrobe Experiment Management System

(Kartashov & Barski, 2015). Briefly, reads were aligned by STAR (version STAR_2.5.3a) with

default arguments except --outFilterMultimapNmax 1 and --outFilterMismatchNmax 2. The --

outFilterMultimapNmax parameter was used to allow unique alignments only, and the --

outFilterMismatchNmax parameter was used to allow a maximum of 2 errors. NCBI RefSeq

annotation from the mm10 UCSC genome browser was used, and canonical TSSs (1 TSS per gene)

were analyzed. All reads from the resulting .bam files were split for related isoforms with respect

to RefSeq annotation. Then, the EM algorithm was used to estimate the number of reads for each

isoform. To detect differentially expressed genes between two biological samples, a read count

output file was input to the DESeq2 package (version 1.16.1); then, the program functions

DESeqDataSetFromMatrix and DESeq were used to compare each gene's expression level between

two biological samples. Differentially expressed genes were identified through binominal tests,

thresholding Benjamini-Hochberg- adjusted P values to <0.01. To perform gene ontology analyses,

the functional annotation clustering tool in DAVID (version 6.8) was used, and a background of all

mouse genes was applied. Biological process term groups with a significance of $P < 0.05$ (modified

Fisher's exact test) were considered significant.

Cross-linking ChIP-seq with the ChIPmetation method (Schmidl, Rendeiro, Sheffield, &

Bock, 2015) was performed for H2AK119ub, H3K27me3, and RNF2 as described previously

(Schmidl et al., 2015). Data analysis for both ChIP-seq and RNA-seq was performed in the

BioWardrobe Experiment Management System (https://github.com/Barski-lab/biowardrobe

(Kartashov & Barski, 2015)). Briefly, reads were aligned to the mouse genome (mm10) with

Bowtie (version 1.0.0 (Langmead, Trapnell, Pop, & Salzberg, 2009)), assigned to RefSeq genes

(which have one annotation per gene) using the BioWardrobe algorithm, and displayed on a local

mirror of the UCSC genome browser as coverage. Peaks of H2AK119ub-, H3K27me3- and RNF2-

enrichment were identified using MACS2 (version 2.0.10.20130712 (Zhang et al., 2008)). Pearson

correlations for the genome-wide enrichment of the peaks among ChIP-seq library replicates were

analyzed using SeqMonk (Babraham Institute). Average tag density profiles were calculated

around gene bodies, including 5-kb upstream and 5-kb downstream of the genes. Resulting graphs

were smoothed in 200-bp windows. Enrichment levels for ChIP-seq experiments were calculated

for 4-kb windows, promoter regions of genes (±2 kb surrounding TSSs), and enhancer regions. To

normalize tag value read counts were multiplied by 1,000,000 and then divided by the total number

of reads in each nucleotide position. The total amount of tag values in promoter or enhancer regions

were calculated as enrichment. Microarray data was analyzed using the processed data (Jameson

et al., 2012). Differentially expressed genes were identified through a p-value cutoff of 0.05, and a

fold change cutoff of 2 for the comparison between E13.5 XX supporting cells and E13.5 XY

supporting cells. Highly expressed genes in E13.5 XY supporting cells and in E13.5 XX supporting

cells were termed as "Pre-sertoli genes" and "Pre-granulosa genes", respectively. RNA-seq and

ChIP-seq data reported in this paper have been deposited in GEO under accession code GSE167516.

Acknowledgments

We thank David Zarkower and Vivian Bardwell for their help and discussion in the initial phase of

this project, Katie Gerhardt for editing the manuscript, and the members of the Namekawa and

Maezawa laboratories for discussion and helpful comments regarding the manuscript. This work

was supported by Japan Society for the Promotion of Science Grant-in-Aid for Research Activity

Start-up (19K21196), the Takeda Science Foundation (2019), and the Uehara Memorial Foundation

Research Incentive Grant (2018) to SM, NIH Grants DP2GM119134 to AB, R35GM119458 to TD,

and R01GM098605 and R01GM122776 to SHN.

Competing Interests

The authors declare no competing interests.

References

Aloia, L., Di Stefano, B., & Di Croce, L. (2013). Polycomb complexes in stem cells and embryonic

development. *Development, 140*(12), 2525-2534. doi:10.1242/dev.091553

Bernstein, B. E., Mikkelsen, T. S., Xie, X., Kamal, M., Huebert, D. J., Cuff, J., . . . Lander, E. S.

(2006). A bivalent chromatin structure marks key developmental genes in embryonic stem

cells. *Cell, 125*(2), 315-326. doi:10.1016/j.cell.2006.02.041

Blackledge, N. P., Farcas, A. M., Kondo, T., King, H. W., McGouran, J. F., Hanssen, L. L., . . .

Klose, R. J. (2014). Variant PRC1 complex-dependent H2A ubiquitylation drives PRC2

recruitment   and   polycomb   domain   formation.   *Cell,   157*(6),   1445-1459.

doi:10.1016/j.cell.2014.05.004

Cales, C., Roman-Trufero, M., Pavon, L., Serrano, I., Melgar, T., Endoh, M., . . . Vidal, M. (2008).

Inactivation of the polycomb group protein Ring1B unveils an antiproliferative role in

hematopoietic cell expansion and cooperation with tumorigenesis associated with Ink4a

deletion. *Mol Cell Biol, 28*(3), 1018-1028. doi:10.1128/MCB.01136-07

Capel, B. (2017). Vertebrate sex determination: evolutionary plasticity of a fundamental switch.

Nat Rev Genet, 18(11), 675-689. doi:10.1038/nrg.2017.60

Chang, Y. F., Lee-Chang, J. S., Panneerdoss, S., MacLean, J. A., 2nd, & Rao, M. K. (2011). Isolation

of Sertoli, Leydig, and spermatogenic cells from the mouse testis. *Biotechniques, 51*(5),

341-342, 344. doi:10.2144/000113764

Chassot, A. A., Ranc, F., Gregoire, E. P., Roepers-Gajadien, H. L., Taketo, M. M., Camerino, G.,

Chaboissier, M. C. (2008). Activation of beta-catenin signaling by Rspo1 controls

differentiation of the mammalian ovary. *Human Molecular Genetics, 17*(9), 1264-1277.

doi:10.1093/hmg/ddn016

Cooper, S., Dienstbier, M., Hassan, R., Schermelleh, L., Sharif, J., Blackledge, N. P., . . .

Brockdorff, N. (2014). Targeting polycomb to pericentric heterochromatin in embryonic

stem cells reveals a role for H2AK119u1 in PRC2 recruitment. *Cell Rep, 7*(5), 1456-1470. doi:10.1016/j.celrep.2014.04.012

del Mar Lorente, M., Marcos-Gutierrez, C., Perez, C., Schoorlemmer, J., Ramirez, A., Magin, T., & Vidal, M. (2000). Loss- and gain-of-function mutations show a polycomb group function for Ring1A in mice. *Development, 127*(23), 5093-5100.

Du, H., & Taylor, H. S. (2015). The Role of Hox Genes in Female Reproductive Tract Development, Adult Function, and Fertility. *Cold Spring Harb Perspect Med, 6*(1), a023002. doi:10.1101/cshperspect.a023002

Endoh, M., Endo, T. A., Endoh, T., Isono, K., Sharif, J., Ohara, O., . . . Koseki, H. (2012). Histone H2A mono-ubiquitination is a crucial step to mediate PRC1-dependent repression of developmental genes to maintain ES cell identity. *PLoS Genet, 8*(7), e1002774. doi:10.1371/journal.pgen.1002774

Gao, Z., Zhang, J., Bonasio, R., Strino, F., Sawai, A., Parisi, F., . . . Reinberg, D. (2012). PCGF homologs, CBX proteins, and RYBP define functionally distinct PRC1 family complexes. *Mol Cell, 45*(3), 344-356. doi:10.1016/j.molcel.2012.01.002

Garcia-Moreno, S. A., Lin, Y. T., Futtner, C. R., Salamone, I. M., Capel, B., & Maatouk, D. M. (2019). CBX2 is required to stabilize the testis pathway by repressing Wnt signaling. *PLoS Genet, 15*(5), e1007895. doi:10.1371/journal.pgen.1007895

Hammoud, S. S., Nix, D. A., Zhang, H., Purwar, J., Carrell, D. T., & Cairns, B. R. (2009). Distinctive chromatin in human sperm packages genes for embryo development. *Nature, 460*(7254), 473-478. doi:10.1038/nature08162

Holdcraft, R. W., & Braun, R. E. (2004). Androgen receptor function is required in Sertoli cells for the terminal differentiation of haploid spermatids. *Development, 131*(2), 459-467. doi:10.1242/dev.00957

Jacobs, J. J., Kieboom, K., Marino, S., DePinho, R. A., & van Lohuizen, M. (1999). The oncogene

and Polycomb-group gene bmi-1 regulates cell proliferation and senescence through the

ink4a locus. *Nature, 397*(6715), 164-168. doi:10.1038/16476

Jameson, S. A., Natarajan, A., Cool, J., DeFalco, T., Maatouk, D. M., Mork, L., . . . Capel, B.

(2012). Temporal transcriptional profiling of somatic and germ cells reveals biased lineage

priming of sexual fate in the fetal mouse gonad. *PLoS Genet, 8*(3), e1002575.

doi:10.1371/journal.pgen.1002575

Jochumsen, U., Werner, R., Miura, N., Richter-Unruh, A., Hiort, O., & Holterhus, P. M. (2008).

Mutation analysis of FOXF2 in patients with disorders of sex development (DSD) in

combination with cleft palate. *Sex Dev, 2*(6), 302-308. doi:10.1159/000195679

Kartashov, A. V., & Barski, A. (2015). BioWardrobe: an integrated platform for analysis of

epigenomics and transcriptomics data. *Genome Biol, 16*, 158. doi:10.1186/s13059-015-

0720-3

Katoh-Fukui, Y., Tsuchiya, R., Shiroishi, T., Nakahara, Y., Hashimoto, N., Noguchi, K., &

Higashinakagawa, T. (1998). Male-to-female sex reversal in M33 mutant mice. *Nature,*

393(6686), 688-692. doi:10.1038/31482

Kuroki, S., Matoba, S., Akiyoshi, M., Matsumura, Y., Miyachi, H., Mise, N., . . . Tachibana, M.

(2013). Epigenetic regulation of mouse sex determination by the histone demethylase

Jmjd1a. *Science, 341*(6150), 1106-1109. doi:10.1126/science.1239864

Langmead, B., Trapnell, C., Pop, M., & Salzberg, S. L. (2009). Ultrafast and memory-efficient

alignment of short DNA sequences to the human genome. *Genome Biol, 10*(3), R25.

doi:10.1186/gb-2009-10-3-r25

Laugesen, A., Hojfeldt, J. W., & Helin, K. (2019). Molecular Mechanisms Directing PRC2

Recruitment and H3K27 Methylation. *Mol Cell, 74*(1), 8-18.

doi:10.1016/j.molcel.2019.03.011

Lee, T. I., Jenner, R. G., Boyer, L. A., Guenther, M. G., Levine, S. S., Kumar, R. M., . . . Young, R.

A. (2006). Control of developmental regulators by Polycomb in human embryonic stem

cells. *Cell, 125*(2), 301-313. doi:10.1016/j.cell.2006.02.043

Lesch, B. J., Dokshin, G. A., Young, R. A., McCarrey, J. R., & Page, D. C. (2013). A set of genes

critical to development is epigenetically poised in mouse germ cells from fetal stages

through completion of meiosis. *Proc Natl Acad Sci U S A, 110*(40), 16061-16066.

doi:10.1073/pnas.1315204110

Li, Y., Zhang, L., Hu, Y., Chen, M., Han, F., Qin, Y., . . . Gao, F. (2017). beta-Catenin directs the

transformation of testis Sertoli cells to ovarian granulosa-like cells by inducing Foxl2

expression. *J Biol Chem, 292*(43), 17577-17586. doi:10.1074/jbc.M117.811349

Lindeman, R. E., Gearhart, M. D., Minkina, A., Krentz, A. D., Bardwell, V. J., & Zarkower, D.

(2015). Sexual cell-fate reprogramming in the ovary by DMRT1. *Current Biology, 25*(6),

764-771. doi:10.1016/j.cub.2015.01.034

Maezawa, S., Hasegawa, K., Yukawa, M., Kubo, N., Sakashita, A., Alavattam, K. G., . . .

Namekawa, S. H. (2018). Polycomb protein SCML2 facilitates H3K27me3 to establish

bivalent domains in the male germline. *Proc Natl Acad Sci U S A, 115*(19), 4957-4962.

doi:10.1073/pnas.1804512115

Maezawa, S., Hasegawa, K., Yukawa, M., Sakashita, A., Alavattam, K. G., Andreassen, P. R., . . .

Namekawa, S. H. (2017). Polycomb directs timely activation of germline genes in

spermatogenesis. *Genes Dev, 31*(16), 1693-1703. doi:10.1101/gad.302000.117

Maezawa, S., Yukawa, M., Alavattam, K. G., Barski, A., & Namekawa, S. H. (2018). Dynamic

reorganization of open chromatin underlies diverse transcriptomes during spermatogenesis.

Nucleic Acids Res, 46(2), 593-608. doi:10.1093/nar/gkx1052

Matson, C. K., Murphy, M. W., Sarver, A. L., Griswold, M. D., Bardwell, V. J., & Zarkower, D.

(2011). DMRT1 prevents female reprogramming in the postnatal mammalian testis.

Nature, 476(7358), 101-104. doi:10.1038/nature10239

Matson, C. K., & Zarkower, D. (2012). Sex and the singular DM domain: insights into sexual

regulation, evolution and plasticity. *Nat Rev Genet, 13*(3), 163-174. doi:10.1038/nrg3161

Minkina, A., Matson, C. K., Lindeman, R. E., Ghyselinck, N. B., Bardwell, V. J., & Zarkower, D.

(2014). DMRT1 protects male gonadal cells from retinoid-dependent sexual

transdifferentiation. *Dev Cell, 29*(5), 511-520. doi:10.1016/j.devcel.2014.04.017

Mu, W., Starmer, J., Fedoriw, A. M., Yee, D., & Magnuson, T. (2014). Repression of the soma-

specific transcriptome by Polycomb-repressive complex 2 promotes male germ cell

development. *Genes Dev, 28*(18), 2056-2069. doi:10.1101/gad.246124.114

Nicol, B., Grimm, S. A., Gruzdev, A., Scott, G. J., Ray, M. K., & Yao, H. H. (2018). Genome-wide

identification of FOXL2 binding and characterization of FOXL2 feminizing action in the

fetal gonads. *Human Molecular Genetics, 27*(24), 4273-4287. doi:10.1093/hmg/ddy312

Oduwole, O. O., Peltoketo, H., & Huhtaniemi, I. T. (2018). Role of Follicle-Stimulating Hormone

in Spermatogenesis. *Front Endocrinol (Lausanne), 9*, 763. doi:10.3389/fendo.2018.00763

Ottolenghi, C., Pelosi, E., Tran, J., Colombino, M., Douglass, E., Nedorezov, T., . . . Schlessinger,

D. (2007). Loss of Wnt4 and Foxl2 leads to female-to-male sex reversal extending to germ

cells. *Hum Mol Genet, 16*(23), 2795-2804. doi:10.1093/hmg/ddm235

Parma, P., Radi, O., Vidal, V., Chaboissier, M. C., Dellambra, E., Valentini, S., . . . Camerino, G.

(2006). R-spondin1 is essential in sex determination, skin differentiation and malignancy.

Nat Genet, 38(11), 1304-1309. doi:10.1038/ng1907

Piunti, A., Rossi, A., Cerutti, A., Albert, M., Jammula, S., Scelfo, A., . . . Pasini, D. (2014).

Polycomb proteins control proliferation and transformation independently of cell cycle

checkpoints by regulating DNA replication. *Nat Commun, 5*, 3649.

doi:10.1038/ncomms4649

Posfai, E., Kunzmann, R., Brochard, V., Salvaing, J., Cabuy, E., Roloff, T. C., . . . Peters, A. H.

(2012). Polycomb function during oogenesis is required for mouse embryonic

development. *Genes Dev, 26*(9), 920-932. doi:10.1101/gad.188094.112

Raymond, C. S., Murphy, M. W., O'Sullivan, M. G., Bardwell, V. J., & Zarkower, D. (2000). Dmrt1,

a gene related to worm and fly sexual regulators, is required for mammalian testis

differentiation. *Genes Dev, 14*(20), 2587-2595. doi:10.1101/gad.834100

Raymond, C. S., Shamu, C. E., Shen, M. M., Seifert, K. J., Hirsch, B., Hodgkin, J., & Zarkower,

D. (1998). Evidence for evolutionary conservation of sex-determining genes. *Nature,*

391(6668), 691-695. doi:10.1038/35618

Ringrose, L., & Paro, R. (2007). Polycomb/Trithorax response elements and epigenetic memory of

cell identity. *Development, 134*(2), 223-232. doi:10.1242/dev.02723

Schmidl, C., Rendeiro, A. F., Sheffield, N. C., & Bock, C. (2015). ChIPmentation: fast, robust, low-

input ChIP-seq for histones and transcription factors. *Nat Methods, 12*(10), 963-965.

doi:10.1038/nmeth.3542

Schmidt, D., Ovitt, C. E., Anlag, K., Fehsenfeld, S., Gredsted, L., Treier, A. C., & Treier, M. (2004).

The murine winged-helix transcription factor Foxl2 is required for granulosa cell

differentiation   and   ovary   maintenance.   *Development,   131*(4),   933-942.

doi:10.1242/dev.00969

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image

analysis.   *Nat   Methods,   9*(7),   671-675.   Retrieved   from

https://www.ncbi.nlm.nih.gov/pubmed/22930834

Serrano, M., Lee, H., Chin, L., Cordon-Cardo, C., Beach, D., & DePinho, R. A. (1996). Role of the

INK4a   locus   in   tumor   suppression   and   cell   mortality.   *Cell,   85*(1),   27-37.

doi:10.1016/s0092-8674(00)81079-x

Sharpe, R. M., McKinnell, C., Kivlin, C., & Fisher, J. S. (2003). Proliferation and functional

maturation of Sertoli cells, and their relevance to disorders of testis function in adulthood.

Reproduction, 125(6), 769-784.

Simon, J. A., & Kingston, R. E. (2013). Occupying chromatin: Polycomb mechanisms for getting

to genomic targets, stopping transcriptional traffic, and staying put. *Mol Cell, 49*(5), 808-

824. doi:10.1016/j.molcel.2013.02.013

Sin, H. S., Kartashov, A. V., Hasegawa, K., Barski, A., & Namekawa, S. H. (2015). Poised

chromatin and bivalent domains facilitate the mitosis-to-meiosis transition in the male

germline. *BMC Biol, 13*, 53. doi:10.1186/s12915-015-0159-8

Svingen, T., & Koopman, P. (2013). Building the mammalian testis: origins, differentiation, and

assembly of the component cell populations. *Genes Dev, 27*(22), 2409-2426.

doi:10.1101/gad.228080.113

Swain, A., & Lovell-Badge, R. (1999). Mammalian sex determination: a molecular drama. *Genes*

Dev, 13(7), 755-767. doi:10.1101/gad.13.7.755

Uhlenhaut, N. H., Jakob, S., Anlag, K., Eisenberger, T., Sekido, R., Kress, J., . . . Treier, M. (2009).

Somatic sex reprogramming of adult ovaries to testes by FOXL2 ablation. *Cell, 139*(6),

1130-1142. doi:10.1016/j.cell.2009.11.021

Ventura, A., Kirsch, D. G., McLaughlin, M. E., Tuveson, D. A., Grimm, J., Lintault, L., . . . Jacks,

T. (2007). Restoration of p53 function leads to tumour regression in vivo. *Nature,*

445(7128), 661-665. doi:10.1038/nature05541

Vidal, V. P., Chaboissier, M. C., de Rooij, D. G., & Schedl, A. (2001). Sox9 induces testis

development in XX transgenic mice. *Nat Genet, 28*(3), 216-217. doi:10.1038/90046

Wilhelm, D., Palmer, S., & Koopman, P. (2007). Sex determination and gonadal development in

mammals. *Physiol Rev, 87*(1), 1-28. doi:10.1152/physrev.00009.2006

Yokobayashi, S., Liang, C. Y., Kohler, H., Nestorov, P., Liu, Z., Vidal, M., . . . Peters, A. H. (2013).

PRC1 coordinates timing of sexual differentiation of female primordial germ cells. *Nature,*

495(7440), 236-240. doi:10.1038/nature11918

Zhang, Y., Liu, T., Meyer, C. A., Eeckhoute, J., Johnson, D. S., Bernstein, B. E., . . . Liu, X. S.

(2008). Model-based analysis of ChIP-Seq (MACS). *Genome Biol, 9*(9), R137.

doi:10.1186/gb-2008-9-9-r137

Zhao, L., Svingen, T., Ng, E. T., & Koopman, P. (2015). Female-to-male sex reversal in mice caused

by transgenic overexpression of Dmrt1. *Development, 142*(6), 1083-1088.

doi:10.1242/dev.122184

Figure 1. Deletion of PRC1 in Sertoli cells.
(**A**) RNF2 and RNF2-mediated H2AK119ub localized at GATA4-positive Sertoli cells (yellow arrows) in a testicular section at 6 weeks of age. Regions bordered by yellow squares are magnified in the right panels. Bars in the large panels: 50 μm. Bars in the magnified panels: 20 μm. (**B**) Genotypes and photographs of testes at 3 weeks of age. Measurement scale in the panel: 2 cm. (**C**) Testicular weight/body weight ratio ($\times 10^{-3}$) at 12 weeks of age. $P < 0.0001$, unpaired t-test. (**D, E**) Localization of H2AK119ub and GATA4 in PRC1ctrl and PRC1AcKO at embryonic day 15.5 (**D**) and 1 week of age (**E**). Regions bordered by yellow squares are magnified in the right panels. Bars in the large panels: 50 μm. Bars in the magnified panels: 20 μm. H2AK119ub⁻ Sertoli cells in mutants are shown with white arrows. (**F**) The fertility of PRC1AcKO males, at 8-11 weeks of age and 5 months of age, were tested via crosses with CD1 wild-type females. Numbers of males tested are shown within the bars, and numbers of females with pups and all females are shown above the bars. *$P < 0.05$, Fisher's exact test. (**G**) Litter sizes of breeding tests at 8-11 weeks of age and 5 months of age. *$P < 0.05$, Welch's t-test. (**H**) Ligand hormone tests at 8-11 weeks of age testes. *$P < 0.05$, Welch's t-test. n.s., not significant.

Figure 2. PRC1 is required for proliferation of Sertoli cells.
(**A**) PCNA and Ki67 were not detected in GATA4-positive Sertoli cells (arrows) in a PRC1AcKO testicular section at 6 weeks of age, while PCNA and Ki67 were present in PRC1ctrl Sertoli cells. (**B**) Testicular sections of the indicated genotypes 1 day following the injection of EdU into males at 1 week of age. The presence of EdU-positive Sertoli cells (arrows) was decreased in PRC1AcKO testes. Regions bordered by yellow squares are magnified in the right panels. Bars in the large panels: 50 μm. Bars in the magnified panels: 20 μm.

Figure 3. In Sertoli cells, PRC1 suppresses genes required for granulosa cells.
(**A**) Genotypes and experiment schematic. Sertoli cells were isolated from P7 testes and cultured for 4 days in the presence of 4-OHT prior to RNA-seq analyses. (**B**) The numbers of differentially expressed genes detected by RNA-seq (≥1.5-fold change, P_{adj} < 0.05) in Sertoli cells (Two biological replicates) between PRC1ctrl and PRC1RcKO. (**C**) Heatmaps showing gene expression patterns for upregulated (left) and down-regulated (right) genes in Sertoli cells. (**D**) GO term analyses. (**E**) Expression levels for representative Sertoli and granulosa genes. (**F, G**) H2AK119ub and H3K27me3 ChIP-seq enrichment around the TSSs of representative Sertoli and granulosa genes.

Figure 4. In Sertoli cells, Polycomb-mediated marks are enriched on genes required for granulosa cells.
(**A**) Genome track views of representative genes in the female gene regulatory network. ChIP-seq enrichment in wild-type Sertoli cells is shown (top); RNA-seq peaks in PRC1ctrl and PRC1RcKO Sertoli cells are shown (bottom). (**B**) Box-and-whisker plots showing distributions of enrichment for ChIP-seq data. Central bars represent medians, the boxes encompass 50% of the data points, and the whiskers indicate 90% of the data points. *P*, Mann-Whitney U tests. (**C**) Scatter plots showing ChIP-seq enrichment (±2 kb around TSSs) of indicated modifications on genes upregulated (left panels) and down-regulated (right panels) in Sertoli cells. The distribution of all genes is shown with gray dots. (**D**) Average tag densities of H2AK119ub ChIP-seq enrichment.

Figure 5. Polycomb inactivates the female gene regulatory network in Sertoli cells.
(**A**) Box-and-whisker plots showing distributions of RNA-seq data. Central bars represent medians, the boxes encompass 50% of the data points, and the whiskers indicate 90% of the data points. *** $P < 0.0001$, Mann-Whitney U tests. (**B**) Scatter plots showing the Pearson correlation between RNA-seq data for genes regulated in E13.5 support cells and P7 Sertoli cells. A linear trendline is shown in blue. (**C**) Box-and-whisker plots showing distributions of enrichment for H2AK119ub ChIP-seq data. Central bars represent medians, the boxes encompass 50% of the data points, and the whiskers indicate 90% of the data points. *** $P < 0.0001$, ** $P < 0.005$, Mann-Whitney U tests. (**D**) Scatter plots showing the Pearson correlation between ChIP-seq enrichment (±2 kb around TSSs) and gene expression in Sertoli cells. A linear trendline is shown in blue.

Figure 1- figure supplement 1. Deletion of PRC1 in Sertoli cells causes degeneration of testes at 6 weeks of age. Localization of H2AK119ub and GATA4 in PRC1ctrl and PRC1AcKO at 6 weeks of age. Regions bordered by yellow squares are magnified in the right panels. Bars in the large panels: 50 μm. Bars in the magnified panels: 20 μm. H2AK119ub⁻ Sertoli cells in mutants are shown with white arrows.

Figure 3- figure supplement 1. Biological replicates for RNA-seq and ChIP-seq data.
(**A**) Scatter plots show the reproducibility of RNA-seq enrichment at individual peaks between biological replicates.
(**B**) Scatter plots show the reproducibility of and ChIP-seq enrichment at individual peaks between biological
replicates. Each peak was identified using MACS (P < 1×10⁻⁵). H3K27ac ChIP-seq enrichment levels are shown in
log₂ RPKM values. The color scale indicates RNA-seq or ChIP-seq peak density. Pearson correlation values (R) are
shown.

Figure 4- figure supplement 1. In Sertoli cells, H3K27me3 is enriched on genes required for granulosa cells.
Average tag densities of H3K27me3 ChIP-seq enrichment on the groups of genes indicated in the panel.

Figure 5- figure supplement 1. Expression profiles of specifically expressed genes in E13.5 granulosa cells.
Microarray analysis of gene expression in E13.5 supporting cells (42). Genes with the criteria of 2-fold higher expression in E13.5 XX supporting cells and $P < 0.05$ were termed as "Pre-granulosa genes" and shown in green. Genes with the criteria of 2-fold higher expression in E13.5 XY supporting cells and $P < 0.05$ were termed as "Pre-Sertoli genes" and shown in red.

Figure 5- figure supplement 2. H3K27me3 is involved in the female gene regulatory network in Sertoli cells.
(**A**) Box-and-whisker plots showing distributions of enrichment for H3K27me3 ChIP-seq data. Central bars represent medians, the boxes encompass 50% of the data points, and the whiskers indicate 90% of the data points. *** $P <$ 0.0001, Mann-Whitney U tests. (**B**) Scatter plots showing the Pearson correlation between ChIP-seq enrichment (±2 kb around TSSs) and gene expression in Sertoli cells. A linear trendline is shown in blue.