BIOLOGICAL SCIENCES: Developmental Biology

Polycomb suppresses a female gene regulatory network

in Sertoli cells.

So Maezawa[a, b, c, d*], Masashi Yukawa[b, e], Kazuteru Hasegawa[a, b], Ryo Sugiyama[c], Mengwen

Hu[a, b, f], Miguel Vidal[g], Haruhiko Koseki[h], Artem Barski[b, e], Tony DeFalco[a, b], and Satoshi H.

Namekawa[a, b, f*]

[a] Division of Reproductive Sciences, Division of Developmental Biology, Perinatal Institute,

Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio, 45229, USA

[b] Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, Ohio, 45229,

USA

[c] Department of Animal Science and Biotechnology, School of Veterinary Medicine, Azabu

University, Sagamihara, Kanagawa 252-5201, Japan

[d] Faculty of Science and Technology, Department of Applied Biological Science, Tokyo University

of Science, Chiba 278-8510, Japan.

[e] Division of Allergy and Immunology, Division of Human Genetics, Cincinnati Children's

Hospital Medical Center, Cincinnati, Ohio, 45229, USA

[f] Department of Microbiology and Molecular Genetics, University of California, Davis,

California, 95616, USA

[g] Centro de Investigaciones Biológicas Margarita Salas, Department of Cellular and Molecular

Biology, Madrid 28040, Spain

[h] Developmental Genetics Laboratory, RIKEN Center for Allergy and Immunology, Yokohama,

Kanagawa, Japan

*Corresponding authors: E-mail: s-maezawa@rs.tus.ac.jp; snamekawa@ucdavis.edu

Keywords: Polycomb, Sertoli cells, Sex determination, Testis, Spermatogenesis

Abstract

Gonadal sex determination is controlled by the support cells of testes and ovaries. Sexual fate

becomes labile and interchangeable with the removal of specific, critical transcription factors

from postnatal gonad support cells. In Sertoli cells, the specific support cells for postnatal

testes, the epigenetic mechanism that maintains cellular memory to suppress female sexual

differentiation remains unknown. Here, we show that, in postnatal Sertoli cells, Polycomb

suppresses a female gene regulatory network. Through genetic ablation, we removed

Polycomb repressive complex 1 (PRC1) from embryonic Sertoli cells after sex determination.

PRC1-depleted postnatal Sertoli cells exhibited defective proliferation and cell death, leading

to the degeneration of adult testes. In adult Sertoli cells, PRC1 suppressed the specific, critical

genes required for granulosa cells, the support cells of ovaries, thereby inactivating the female

gene regulatory network. The underlying chromatin of female genes was coated with

Polycomb-mediated repressive modifications: PRC1-mediated H2AK119ub and PRC2-

mediated H3K27me3. Taken together, we identify a critical mechanism centered on

Polycomb that maintains the male fate in adult testes.

Significance

Sex differences in mammals are defined by the reproductive organs, testes and ovaries. In

testes and ovaries, sexual fate is determined by Sertoli cells and granulosa cells, respectively,

which are supporting cells derived from common somatic progenitors. Critical transcription

factors determine sexual fate of Sertoli and granulosa cells, and, remarkably, removal of these

factors reverses sexual identity. Therefore, sexual fate is surprisingly plastic. Here we address

a long-standing question in developmental biology of how male sexual fate is maintained in

testes throughout life. We show that epigenetic machinery of Polycomb repressive complex 1

(PRC1) suppresses a female gene regulatory network in Sertoli cells. Thus, Polycomb

preserves cellular memory and sexual identity of Sertoli cells, thereby defining the testicular

fate.

Introduction

In mammals, gonadal sex determination takes place in the bipotential somatic cell precursors of male Sertoli cells and female granulosa cells in embryos (1-3). Sertoli cells are the first somatic cells to differentiate in the XY gonad. In testes, Sertoli cells function as a regulatory hub for both differentiation and survival of germ cells, thereby determining male sexual fate (4). The mechanisms maintaining the male cellular identity of Sertoli cells are fundamental for adult testicular functions, including spermatogenesis and hormone production.

At the time of sex determination in embryos, the commitment to the male fate is triggered by the expression of the Y-linked *Sry* gene, and, subsequently, the female fate is suppressed (1-3). Distinct gene regulatory networks promote the male or female fate and are regulated by strong feedback loops that antagonize each other, canalizing one fate from the other (3). Sexual fate is interchangeable even after the initial commitment to Sertoli cells or granulosa cells with the removal of specific, critical transcription factors. The loss of *Dmrt1* in Sertoli cells leads to derepression of *Foxl2*, a master regulator of granulosa cell fate, and transdifferentiation of cell fate from Sertoli to granulosa cells (5, 6). On the other hand, the loss of *Foxl2* in granulosa cells leads to derepression of the male gene network and transdifferentiation of cell fate from granulosa to Sertoli cells (7). These findings, together with follow-up studies (8-12), suggest that active repression of the alternate sexual fate is important for both testicular and ovarian function, even in adult life.

Epigenetic silencing mechanisms serve as molecular switches for the sex determination of bipotential somatic cell precursors. The deletion of a Polycomb gene *Cbx2* results in male to female reversal (13), which is mediated through suppression of genes required for the female fate (14). Additionally, regulation of H3K9 methylation is important for the male sexual fate (15). Although these studies highlight the importance of epigenetic mechanisms for initial sex determination, the epigenetic mechanisms by which male cellular identity is maintained through cell divisions and the proliferation of Sertoli cells remain to be determined.

Polycomb proteins suppress non-lineage-specific genes and define cellular identities of each lineage in stem cells and in development (16-18). In this study, we show that Polycomb suppresses the female gene regulatory network in postnatal Sertoli cells, thereby promoting the male gene regulatory network to ensure male cell fate. We generated loss-of-function mouse models of Polycomb repressive complex 1 (PRC1) in Sertoli cells after initial sex determination.

94 We show that PRC1 is required for the proliferation of Sertoli cells, as well as the suppression of
95 non-lineage-specific genes and the female gene regulatory network in Sertoli cells. Taken together,
96 we identify a critical mechanism centered on Polycomb that maintains male fate in adult testes.
97

Results

PRC1 in Sertoli cells is required for the maintenance of spermatogenesis. In postnatal Sertoli
cells (which are detected by the Sertoli cell marker GATA4), RNF2 is highly expressed and the
RNF2-mediated epigenetic mark H2AK119ub is abundant (Fig.1*A*), which suggests PRC1
functions in these cells. To determine the function of PRC1, we generated a PRC1 loss-of-function
mouse model by removing two redundant catalytic subunits, RNF2 and RING1 (19). We generated
a conditional deletion of *Rnf2* (*Rnf2*cKO) using *Amh*-Cre, which is expressed specifically in Sertoli
cells after embryonic day 14.5 (E14.5) (20), in a background of *Ring1*-knockout (KO) mice (*Amh*-
Cre; *Rnf2*cKO; *Ring1*-KO: termed PRC1[Amh-Cre]cKO: PRC1AcKO). Although RNF2 appears to be
the most active component in the heterodimeric E3 ligases of PRC1, the RNF2 paralog RING1 can
partially compensate for the loss of RNF2 (19). Therefore, we made a conditional deletion of *Rnf2*
in a background of *Ring1*-KO mice, which are viable and do not have fertility defects (21). This
strategy enabled us to define the function of RNF2 without compensation from RING1, while also
representing a "complete" loss-of-function of PRC1 as shown in testicular germ cells (22) and in
other biological contexts (19, 23, 24). Since PRC1 has various components, including CBX2 (25),
this strategy allows us to determine the global function of PRC1. At the same time, the use of *Amh*-
Cre allowed us to test the function of PRC1, specifically after the completion of sex determination
at E12.5.
116

PRC1AcKO males have smaller testes compared with littermate controls that harbored
floxed alleles for *Rnf2* on a *Ring1*-KO background without *Amh*-Cre (termed PRC1 control:
PRC1ctrl: Fig. 1*B* and *C*). We confirmed efficient *Amh*-Cre-mediated recombination by observing
depletion of the RNF2-mediated mark, H2AK119ub, in GATA4[+] Sertoli cells of PRC1AcKO testes
at E15.5 (> 95 % efficiency: Fig. 1*D*) and at postnatal day 7 (P7: Fig. 1*E*). In 6 week-old
PRC1AcKO testes, while the tubules with H2AK119ub[+] Sertoli cells (escaped Cre-mediated
deletion) appear to have normal morphology, we frequently observed disorganization of testicular
tubules that contain H2AK119ub[-] Sertoli cells (underwent Cre-mediated deletion: arrowheads, Fig.
S1), suggesting a critical function of PRC1 in Sertoli cells in the organization of testicular tubules.
This mosaic pattern is presumably due to incomplete Cre-mediated recombination, as the Sertoli
cells that escaped Cre-mediated recombination might have repopulated the testes. Consistent with

this interpretation, PRC1AcKO males are subfertile, and 9 out of 15 wild-type females mated with

8-11-week old PRC1AcKO males gave birth (Fig. 1*F*) at comparable litter sizes (Fig.1*G*).

Interestingly, fecundity decreased in aged PRC1AcKO males (5 months old), as litter sizes were

smaller as compared to controls (Fig. 1*G*). To further examine the phenotype, we measured the

blood levels of three hormones critical for testicular homeostasis: although the levels of testosterone

and estradiol were comparable between cKO and control mice, follicle-stimulating hormone was

increased in mutants (Fig. 1*H*). As follicle stimulating hormone levels are regulated by a feedback

mechanism involving Sertoli cells (26), we infer that the dysfunction of Sertoli cells and testicular

degeneration caused increased follicle stimulating hormone levels to recover Sertoli cell function.

To determine the cause of testicular degeneration, we next examined the proliferation of

Sertoli cells. In normal mouse development, Sertoli cells proliferate in fetal and neonatal testes

until approximately 2 weeks after birth, and the number of Sertoli cells in adult testes determines

both testis size and daily sperm production (27). In P7 control testes, GATA4$^+$ Sertoli cells

occasionally co-expressed an S phase marker, PCNA, and another marker of active cell cycle, Ki67

(Fig. 2*A*), consistent with the active proliferation of Sertoli cells. However, in PRC1AcKO testes,

GATA4$^+$ Sertoli cells were largely devoid of PCNA and Ki67 (Fig. 2*A*), suggesting impaired

proliferation of PRC1AcKO Sertoli cells. Next, we independently confirmed the proliferation

phenotype by using EdU labeling of actively proliferating cells. EdU was abdominally

administrated to P7 mice, and testicular sections were examined the following day. While GATA4$^+$

Sertoli cells were occasionally EdU$^+$ (approximately one-fouth) in control testes , GATA4$^+$ Sertoli

cells were devoid of EdU signal in PRC1AcKO testes (Fig. 2*B*). Additional labeling of H2AK119ub

confirmed the loss of PRC1 function in GATA4$^+$ Sertoli cells from PRC1AcKO testes (Fig. 2*B*).

From these results, we conclude that the loss of PRC1 disrupts the proliferation of Sertoli cells.

This is in contrast with PRC1's function in testicular germ cells, in which the loss of PRC1 does

not affect proliferation but instead causes apoptotic cell death (22). This difference suggests a

unique function of PRC1 in Sertoli cells that is distinct from its function in germ cells.

In Sertoli cells, PRC1 suppresses genes required for granulosa cells. We next sought to

determine the genes regulated by PRC1 in Sertoli cells. In PRC1AcKO testes, some Sertoli cells

escaped *Amh*-Cre-mediated recombination. Thus, it was difficult to specifically isolate Sertoli cells

that underwent PRC1 depletion. To precisely determine the function of PRC1 in gene regulation in

Sertoli cells, we used an alternative strategy: we isolated Sertoli cells from a mouse line in which

conditional deletion of PRC1 can be induced by tamoxifen-inducible Cre-mediated recombination

under the control of the endogenous ROSA26 promoter (*ROSA26*-Cre[ERT]; *Rnf2*[floxed/floxed]; *Ring1*-
KO: termed PRC1[ROSA26-CreERT]cKO: PRC1RcKO). After isolating Sertoli cells from P7 testes, we
cultured Sertoli cells for 4 days in the presence of 4-hydroxytamoxifen (4-OHT), and performed
RNA-sequencing (RNA-seq: Fig.3*A*). As a control, we isolated Sertoli cells from control mice
(*Rnf2*[floxed/floxed]; *Ring1*-KO) and cultured them in the same 4-OHT conditions. We performed RNA-
seq for two independent biological replicates and confirmed reproducibility between biological
replicates (Fig. S1*A*).

Our RNA-seq analyses demonstrated that 338 genes were upregulated in PRC1RcKO
Sertoli cells as compared to controls, while 307 genes were downregulated in PRC1RcKO Sertoli
cells (Fig. 3*A* and *B*). Gene ontology (GO) analysis showed that upregulated genes were enriched
for functions in neural differentiation, skeletal system, and cell adhesion (Fig. 3*D*). These categories
suggest that PRC1 suppressed expression of non-lineage-specific genes in Sertoli cells. On the
other hand, GO analysis revealed that downregulated genes were enriched for functions in the cell
cycle and M phase (Fig. 3*D*). This result is in accord with the cell cycle arrest we found in
PRC1AcKO Sertoli cells (Fig. 2).

Since we anticipated suppression of genes required for granulosa cells by PRC1 in Sertoli
cells, we next investigated the expression level of key genes required for granulosa cells. In
PRC1RcKO Sertoli cells, genes required for female sexual development were upregulated: these
genes included *Rspo1,* an activator of the Wnt pathway (29), and *Foxl2,* a key transcription factor
for granulosa cells (30) (Fig. 3*E*). Importantly, these female genes suppress the male fate, and the
loss of these genes leads to female-to-male sex reversal (30-32). Consistent with the antagonistic
function of these female genes with the male pathway, key sex determination genes for the male
pathway were downregulated in PRC1RcKO Sertoli cells: these genes included *Sox9,* an
evolutionarily conserved gene for sex determination, which directs the male pathway downstream
of *Sry* (33), and *Dmrt1* (34, 35), male-determining signalling (Fig. 3*E*).

To determine whether the suppression of female genes was directly regulated by PRC1, we
performed chromatin immunoprecipitation sequencing (ChIP-seq) of PRC1-mediated H2AK119ub
in isolated wild-type Sertoli cells. We further performed ChIP-seq of H3K27me3 in Sertoli cells
since PRC2-mediated H3K27me3 is regulated by PRC1 and its mediated mark H2A119ub (37, 38).
We performed ChIP-seq for two independent biological replicates and confirmed the
reproducibility between biological replicates (Fig. S1*B*). We confirmed the enrichment of

H2AK119ub and H3K27me3 around transcription start sites (TSSs) of *Rspo1* and *Foxl2* (Fig. 3*F*).

Compared to the enrichment of H2AK119ub on these female genes, enrichment of H2AK119ub

was relatively low on the TSSs of the male genes, *Sox9* and *Dmrt1*. Therefore, we conclude that

PRC1 directly binds and suppresses *Rspo1* and *Foxl2*.

Track views of these ChIP-seq data show that all three marks were enriched at *Foxl2* and

Rspo1 loci (Fig. 4*A*). Furthermore, enrichment of these three marks was found in other loci such as

Foxf2, the mutation of which appears in patients with disorders of sex development (39), and

Hoxd13, which is involved in female reproductive tract development (40). These results suggest

that PRC1 works with PRC2 to suppress female genes as well as developmental regulators such as

Fox and *Hox* genes, which are the classical targets of Polycomb-mediated gene repression (41).

To determine the features of genome-wide gene repression mediated by Polycomb

complexes, we analyzed the enrichment of H2AK119ub, H3K27me3, and RNF2 on the upregulated

genes in PRC1RcKO Sertoli cells. H2AK119ub, H3K27me3, and RNF2 were all significantly

enriched on the TSSs of upregulated genes in PRC1RcKO Sertoli cells as compared to all genes in

the genome (Fig. 4*B*). Additional enrichment analysis confirmed the co-enrichment of H2AK119ub

and H3K27me3 (Fig. 4*C,* upper panels) as well as co-enrichment of H2AK119ub and RNF2 (Fig.

4*C,* lower panels) on upregulated genes in PRC1RcKO Sertoli cells. Furthermore, average tag

density analysis confirmed that enrichment of H2AK119ub on upregulated genes in PRC1RcKO

Sertoli cells occured on upstream regions, gene bodies, and downstream regions with the highest

enrichment near TSSs (Fig. 4*D*). We found a similar distribution of H3K27me3 around the gene

bodies of upregulated genes in PRC1RcKO Sertoli cells (Fig. S2). Together, these results

confirmed the genome-wide, global functions of PRC1 in the direct regulation of gene repression

in Sertoli cells.

Polycomb globally inactivates the female gene regulatory network in postnatal Sertoli cells.

Previous studies have suggested that sex determination is canalized by the interconnected,

antagonistic network of genes both in males and in females that are controlled by feedback

mechanisms (3). Since Polycomb is implicated in the maintenance of sex-specific gene regulatory

networks through the PRC2-mediated mark H3K27me3 (14), we hypothesized that PRC1 globally

inactivates the female gene regulatory network in postnatal Sertoli cells. Since male- and female-

specific gene networks are regulated immediately after sex determination during fetal stages (42),

we reasoned that female gene network suppression in fetal Sertoli cells is maintained by PRC1 in

postnatal Sertoli cells. To test this hypothesis, we examined the expression profiles of specifically

expressed genes in E13.5 granulosa cells (42) (Fig. S3), termed "pre-granulosa genes" in postnatal

Sertoli cells. We found that pre-granulosa genes were upregulated in PRC1RcKO postnatal Sertoli

cells as compared to other genes (Fig. 5*A*). On the other hand, specifically expressed genes in E13.5

Sertoli cells (42) (Fig. S3), termed "pre-Sertoli genes," were downregulated in PRC1RcKO

postnatal Sertoli cells as compared to other genes (Fig. 5*A*). We further examined the correlation

of each gene and found that pre-granulosa genes were highly correlated with upregulated genes in

PRC1RcKO postnatal Sertoli cells, while pre-Sertoli genes were highly correlated with

downregulated genes in PRC1RcKO postnatal Sertoli cells (Fig. 5*B*). These results suggest that

PRC1 maintains suppression of the female gene regulatory network, which is initiated at the time

of sex determination and is maintained throughout the development of postnatal Sertoli cells.

We next sought to determine whether PRC1 directly suppresses the female gene regulatory

network in postnatal Sertoli cells. H2AK119ub is significantly enriched on TSSs of pre-granulosa

genes compared to other genes and pre-Sertoli genes (Fig. 5*C*). Among pre-granulosa genes,

enrichment of H2AK119ub is positively correlated with upregulated genes in PRC1RcKO

postnatal Sertoli cells (Fig. 5*D*). We further identified the enrichment of H3K27me3 on TSSs of

pre-granulosa genes in postnatal Sertoli cells (Fig. S4*A*), and the correlation between H3K27me3

and upregulated genes on the pre-granulosa genes in PRC1RcKO postnatal Sertoli cells (Fig. S4*B*).

Together, we conclude that PRC1 globally inactivates the female gene regulatory network in

postnatal Sertoli cells.

Discussion

In this study, we demonstrated that PRC1 is required for proliferation of Sertoli cells and

suppression of the non-lineage-specific gene expression program and the female gene regulatory

network. Among these functions, we infer that suppression of the female gene regulatory network

is the key mechanism to ensure male fate in addition to canonical functions of PRC1, which controls

proliferation and suppression of non-lineage-specific gene expression programs. Although we did

not observe complete infertility using *Amh*-Cre, presumably due to the repopulation of Sertoli cells

that escaped Cre-mediated recombination, smaller testis size and abnormal tubule organization of

cKO testes (Fig. 1) suggest that PRC1 is critical for physiological functions of Sertoli cells. Below,

we discuss two molecular aspects underlying these physiological phenotypes: control of

proliferation and suppression of the female gene regulatory network.

The proliferation of Sertoli cells is a critical determinant of testicular functions because the

number of Sertoli cells defines testicular functions (27). Rapid proliferation is a prominent feature

of juvenile Sertoli cells. In general, Polycomb proteins are associated with the cell cycle checkpoint

by directly suppressing the tumor suppressor locus *Cdkn2a*/*Ink4a*/*Arf* (43), which functions as a

barrier to cancer transformation (44). Therefore, enhanced Polycomb activity is a frequent feature

of human tumors. We found that *Cdkn2a* was derepressed in PRC1RcKO Sertoli cells

(approximately 2.5-fold upregulation: PRKM value for PRC1RcKO: 32.1 v.s. PRC1ctrl: 13.0).

Therefore, our results suggest that PRC1 promotes the rapid proliferation of Sertoli cells by

suppressing *Cdkn2a.* The function of PRC1 in Sertoli cell proliferation is distinct from Polycomb

functions in testicular germ cells, where PRC1 depletion does not alter the proliferation of germ

cells (22). This may be due to the fact that PRC1 and PRC2 are not required for suppression of

Cdkn2a in germ cells (22, 45)*.* The functional difference in testicular germ cells and Sertoli cells

highlights the context-dependent functions of Polycomb proteins. However, a recent study revealed

a novel activity by which PcGs can regulate cell proliferation through DNA replication

independently of *Cdkn2a* (46). Therefore, determining the detailed molecular mechanisms by

which PRC1 controls the proliferation of Sertoli cells will be important for future studies.

Another critical function of PRC1 in Sertoli cells is the suppression of the female gene

regulatory network. DMRT1 is a critical transcription factor that suppresses the expression of

female genes (5). While more than 10-fold upregulation of the female genes (*Rspo1* and *Foxl2*)

was observed for *Dmrt1* mutants (5), the degree of upregulation of these genes was modest in

PRC1RcKO Sertoli cells (Fig. 3). This finding, combined with genetic evidence, indicates that

DMRT1 could be the direct regulator of suppression of female genes, while PRC1-mediated

mechanisms could be a maintenance mechanism in response to the primary silencing mechanisms

determined by DMRT1. Another possibility is compensation by PRC1-independent suppression

mechanisms. While a portion of PRC2-mediated H3K27me3 is regulated by variant PRC1 (37, 38),

another portion of H3K27me3, mediated by canonical PRC1 and PRC2 complexes, is not

downstream of PRC1 (47). These PRC2-mediated mechanisms or other silencing machinery may

be responsible for the suppression of female genes.

The notion that Polycomb regulates the female gene regulatory network has been supported

by other recent evidence. In bipotential precursor cells, genes involved in sex determination are

marked with bivalent chromatin domains (14) that are prevalent in pluripotent stem cells and in

germ cells (48-53). Maintenance of the male fate was explained by the persistence of H3K27me3

on silent female genes in Sertoli cells (14). Consistent with PRC1's function found in the current

study, Polycomb-mediated silencing may globally suppress the female gene regulatory network.

We found that H2AK119ub was largely associated with this group of female genes (Fig. 5).

Therefore, it would be interesting to speculate that antagonistic male and female networks can be

directly coordinated by Polycomb protein functions, including the strong feedback mechanism

underlying both networks. These possibilities raise several outstanding questions to be addressed

in future studies. What are the functions of another Polycomb complex, PRC2, and of each

Polycomb complex component, including CBX2, in postnatal Sertoli cells? What is the function of

Polycomb in female granulosa cells, especially in the suppression of the male gene regulatory

network? Does Polycomb underlie the feedback regulation of each network to define sexual

identity? Our current study provides a foundation to explore these questions.

Methods

Animals

Generation of mutant *Ring1* and *Rnf2* floxed alleles were previously reported (54). *Amh-Cre*

transgenic mice were purchased from The Jackson Laboratory (Stock No: 007915) (20). *Rosa-Cre*

ERT mice were purchased from The Jackson Laboratory (Stock No: 008463) (55). A minimum of

three independent mice were analyzed for each experiment. Institutional Animal Care and Use

Committee approved this work: protocol no. IACUC2018-0040. Fertility tests were performed with

6-weeks old CD1 female mice (purchased from Charles river). At least 2 female mice were bred

with a male mouse for 2 weeks, and the fertility was evaluated by the ratio of pregnant to total

female mice and number of pups.

Ligand test

Blood samples were collected from C57BL/6N mice aged 8−11 weeks. Serum was separated

immediately and stored at -20°C. Hormone assays, including testosterone, estradiol, and follicle

stimulating hormone, were performed by the Center for Research in Reproduction at the University

of Virginia.

Sertoli cell isolation

Sertoli cells were isolated as previously described with minor modifications (56) and collected from

C57BL/6N mice aged 6–8 days. Testes were collected in a 24-well plate in Dulbecco's Modified

Eagle Medium (DMEM) supplemented with GlutaMax (Thermo Fisher Scientific), non-essential amino acids (NEAA) (Thermo Fisher Scientific), and penicillin and streptomycin (Thermo Fisher Scientific). After removing the *tunica albuginea* membrane, testes were digested with collagenase (1 mg/ml) at 34°C for 20 min to remove interstitial cells, then centrifuged at $188\times g$ for 5 min. Tubules were washed with medium and then digested with trypsin (2.5 mg/ml) at 34°C for 20 min to obtain a single-cell suspension. To remove KIT-positive spermatogonia, cells were washed with magnetic cell-sorting (MACS) buffer (PBS supplemented with 0.5% BSA and 5 mM EDTA) and incubated with CD117 (KIT) MicroBeads (Miltenyi Biotec) on ice for 20 min. Cells were separated by autoMACS Pro Separator (Miltenyi Biotec) with the program "possel." Cells in the flow-through fraction were washed with MACS buffer and incubated with CD90.2 (THY1) MicroBeads (Miltenyi Biotec) on ice for 20 min to remove THY1-positive spermatogonia. Cells were separated by autoMACS Pro Separator (Miltenyi Biotec) with the program "posseld." Cells in the flow-through fraction were washed and plated in a 6-well plate for 1 h in the medium supplemented with 10% fetal bovine serum, which promotes adhesion of Sertoli cells. Purity was confirmed by immunostaining.

For PRC1RcKO, cells were cultured for 4 days with 1 µM 4-OHT in Dulbecco's Modified Eagle Medium (DMEM) supplemented with GlutaMax (Thermo Fisher Scientific), non-essential amino acids (NEAA) (Thermo Fisher Scientific), and penicillin and streptomycin (Thermo Fisher Scientific). The same medium was replaced 2 days after the initiation of the culture.

Histological analysis and germ cell slide preparation

For the preparation of testicular paraffin blocks, testes were fixed with 4% paraformaldehyde (PFA) overnight at 4°C with gentle inverting. Testes were dehydrated and embedded in paraffin. For histological analysis, 7 µm-thick paraffin sections were deparaffinized and stained with hematoxylin and eosin. For immunofluorescence analysis of testicular sections, antigen retrieval was performed by boiling the slides in target retrieval solution (DAKO) for 10 min and letting the solution cool for 30 min. Sections were blocked with Blocking One Histo (Nacalai) for 1 h at room temperature and then incubated with primary antibodies overnight at 4°C. The resulting signals were detected by incubation with secondary antibodies conjugated to fluorophores (Thermo Fisher Scientific). Sections were counterstained with DAPI. Images were obtained via a laser scanning confocal microscope A1R (Nikon) and processed with NIS-Elements (Nikon) and ImageJ (National Institutes of Health) (57).

ChIP-sequencing, RNA-sequencing, and data analysis

RNA-seq analyses were performed in the BioWardrobe Experiment Management System (58).
Briefly, reads were aligned by STAR (version STAR_2.5.3a)75 with default arguments except --
outFilterMultimapNmax 1 and --outFilterMismatchNmax 2. The --outFilterMultimapNmax
parameter was used to allow unique alignments only, and the --outFilterMismatchNmax parameter
was used to allow a maximum of 2 errors. NCBI RefSeq annotation from the mm10 UCSC genome
browser 76 was used, and canonical TSSs (1 TSS per gene) were analyzed. All reads from the
resulting .bam files were split for related isoforms with respect to RefSeq annotation. Then, the EM
algorithm was used to estimate the number of reads for each isoform. To detect differentially
expressed genes between two biological samples, a read count output file was input to the DESeq2
package (version 1.16.1); then, the program functions DESeqDataSetFromMatrix and DESeq were
used to compare each gene's expression level between two biological samples. Differentially
expressed genes were identified through binominal tests, thresholding Benjamini-Hochberg-
adjusted P values to <0.01. To perform gene ontology analyses, the functional annotation clustering
tool in DAVID (version 6.8) was used, and a background of all mouse genes was applied.
Biological process term groups with a significance of $P < 0.05$ (modified Fisher's exact test) were
considered significant.

Cross-linking ChIP-seq with the ChIPmetation method (59) was performed for
H2AK119ub, H3K27me3, and RNF2 as described previously (59). Data analysis for both ChIP-
seq and RNA-seq was performed in the BioWardrobe Experiment Management System
(https://github.com/Barski-lab/biowardrobe (58)). Briefly, reads were aligned to the mouse genome
(mm10) with Bowtie (version 1.0.0 (60)), assigned to RefSeq genes (which have one annotation
per gene) using the BioWardrobe algorithm, and displayed on a local mirror of the UCSC genome
browser as coverage. Peaks of H2AK119ub-, H3K27me3- and RNF2-enrichment were identified
using MACS2 (version 2.0.10.20130712 (61)). Pearson correlations for the genome-wide
enrichment of the peaks among ChIP-seq library replicates were analyzed using SeqMonk
(Babraham Institute). Average tag density profiles were calculated around gene bodies, including
5-kb upstream and 5-kb downstream of the genes. Resulting graphs were smoothed in 200-bp
windows. Enrichment levels for ChIP-seq experiments were calculated for 4-kb windows, promoter
regions of genes (±2 kb surrounding TSSs), and enhancer regions. To normalize tag value read
counts were multiplied by 1,000,000 and then divided by the total number of reads in each
nucleotide position. The total amount of tag values in promoter or enhancer regions were calculated
as enrichment. Microarray data was analyzed using the processed data (42). Differentially
expressed genes were identified through a p-value cutoff of 0.05, and a fold change cutoff of 2 for
the comparison between E13.5 XX supporting cells and E13.5 XY supporting cells. Highly

expressed genes in E13.5 XY supporting cells and in E13.5 XX supporting cells were termed as

"Pre-sertoli genes" and "Pre-granulosa genes", respectively. RNA-seq and ChIP-seq data reported

in this paper will be deposited to GEO.

Acknowledgments

We thank David Zarkower and Vivian Bardwell for their help and discussion in the initial phase of

this project, Katie Gerhardt for editing the manuscript, and the members of the Namekawa and

Maezawa laboratories for discussion and helpful comments regarding the manuscript. This work

was supported by Grant-in-Aid for Research Activity Start-up (19K21196), the Takeda Science

Foundation (2019), and the Uehara Memorial Foundation Research Incentive Grant (2018) to SM,

NIH Grants GM119134 to AB, and GM098605 and GM122776 to SHN.

Figure legends

Figure 1. Deletion of PRC1 in Sertoli cells.

(**A**) RNF2 and RNF2-mediated H2AK119ub localized at GATA4-positive Sertoli cells (yellow

arrows) in a testicular section at 6 weeks of age. Regions bordered by yellow squares are magnified

in the right panels. Bars in the large panels: 50 μm. Bars in the magnified panels: 20 μm. (**B**)

Genotypes and photographs of testes at 6 weeks of age. Measurement scale in the panel: 2 cm. (**C**)

Testicular weight/body weight ratio ($\times 10^{-3}$) at 12 weeks of age. $P < 0.0001$, unpaired t-test. (**D, E**)

Localization of H2AK119ub and GATA4 in PRC1ctrl and PRC1AcKO at embryonic day 15.5 (**D**)

and 1 week of age (**E**). Regions bordered by yellow squares are magnified in the right panels. Bars

in the large panels: 50 μm. Bars in the magnified panels: 20 μm. H2AK119ub⁻ Sertoli cells in

mutants are shown with white arrows. (**F**) The fertility of PRC1AcKO males, at 8-11 weeks of age

and 5 months of age, were tested via crosses with CD1 wild-type females. Numbers of males tested

are shown within the bars, and numbers of females with pups and all females are shown above the

bars. *$P < 0.05$, Fisher's exact test. (**G**) Litter sizes of breeding tests at 8-11 weeks of age and 5

months of age. *$P < 0.05$, Welch's t-test. (**H**) Ligand hormone tests at 8-11 weeks of age testes. *P

< 0.05, Welch's t-test. n.s., not significant.

Figure 2. PRC1 is required for proliferation of Sertoli cells.

(**A**) PCNA and Ki67 were not detected in GATA4-positive Sertoli cells (arrows) in a PRC1AcKO

testicular section at 6 weeks of age, while PCNA and Ki67 were present in PRC1ctrl Sertoli cells.

(**B**) Testicular sections of the indicated genotypes 1 day following the injection of EdU into males

at 1 week of age. The presence of EdU-positive Sertoli cells (arrows) was decreased in PRC1AcKO

testes. Regions bordered by yellow squares are magnified in the right panels. Bars in the large

panels: 50 μm. Bars in the magnified panels: 20 μm.

Figure 3. In Sertoli cells, PRC1 suppresses genes required for granulosa cells.

(**A**) Genotypes and experiment schematic. Sertoli cells were isolated from P7 testes and cultured

for 4 days in the presence of 4-OHT prior to RNA-seq analyses. (**B**) The numbers of differentially

expressed genes detected by RNA-seq (≥1.5-fold change) in Sertoli cells (2 biological replicates)

between PRC1ctrl and PRC1RcKO. (**C**) Heatmaps showing gene expression patterns for

upregulated (left) and down-regulated (right) genes in Sertoli cells. (**D**) GO term analyses. (**E**)

Expression levels for representative Sertoli and granulosa genes. (**F, G**) H2AK119ub and

H3K27me3 ChIP-seq enrichment around the TSSs of representative Sertoli and granulosa genes.

Figure 4. In Sertoli cells, Polycomb-mediated marks are enriched on genes required for

granulosa cells.

(**A**) Genome track views of representative genes in the female gene regulatory network. ChIP-seq

enrichment in wild-type Sertoli cells is shown (top); RNA-seq peaks in PRC1ctrl and PRC1RcKO

Sertoli cells are shown (bottom). (**B**) Box-and-whisker plots showing distributions of enrichment

for ChIP-seq data. Central bars represent medians, the boxes encompass 50% of the data points,

and the whiskers indicate 90% of the data points. *P*, Mann-Whitney U tests. (**C**) Scatter plots

showing ChIP-seq enrichment (±2 kb around TSSs) of indicated modifications on genes

upregulated (left panels) and down-regulated (right panels) in Sertoli cells. The distribution of all

genes is shown with gray dots. (**D**) Average tag densities of H2AK119ub ChIP-seq enrichment.

Figure 5. Polycomb inactivates the female gene regulatory network in Sertoli cells.

(**A**) Box-and-whisker plots showing distributions of RNA-seq data. Central bars represent medians,

the boxes encompass 50% of the data points, and the whiskers indicate 90% of the data points. ***

P < 0.0001, Mann-Whitney U tests. (**B**) Scatter plots showing the Pearson correlation between

RNA-seq data for genes regulated in E13.5 support cells and P7 Sertoli cells. A linear trendline is

shown in blue. (**C**) Box-and-whisker plots showing distributions of enrichment for H2AK119ub

ChIP-seq data. Central bars represent medians, the boxes encompass 50% of the data points, and

the whiskers indicate 90% of the data points. *** *P* < 0.0001, ** *P* < 0.005, Mann-Whitney U tests.

(**D**) Scatter plots showing the Pearson correlation between ChIP-seq enrichment (±2 kb around

TSSs) and gene expression in Sertoli cells. A linear trendline is shown in blue.

Figure S1. Deletion of PRC1 in Sertoli cells causes degeneration of testes at 6 weeks of age.
Localization of H2AK119ub and GATA4 in PRC1ctrl and PRC1AcKO at 6 weeks of age. Regions bordered by yellow squares are magnified in the right panels. Bars in the large panels: 50 μm. Bars in the magnified panels: 20 μm. H2AK119ub⁻ Sertoli cells in mutants are shown with white arrows.

Figure S1. Biological replicates for RNA-seq and ChIP-seq data.
(**A**) Scatter plots show the reproducibility of RNA-seq enrichment at individual peaks between biological replicates. (**B**) Scatter plots show the reproducibility of and ChIP-seq enrichment at individual peaks between biological replicates. Each peak was identified using MACS ($P < 1 \times 10^{-5}$). H3K27ac ChIP-seq enrichment levels are shown in $\log_2$ RPKM values. The color scale indicates RNA-seq or ChIP-seq peak density. Pearson correlation values (R) are shown.

Figure S2. In Sertoli cells, H3K27me3 is enriched on genes required for granulosa cells.
Average tag densities of H3K27me3 ChIP-seq enrichment on the groups of genes indicated in the panel.

Figure S3. Expression profiles of specifically expressed genes in E13.5 granulosa cells.
Microarray analysis of gene expression in E13.5 supporting cells (42). Genes with the criteria of 2-fold higher expression in E13.5 XX supporting cells and $P < 0.05$ were termed as "Pre-granulosa genes" and shown in green. Genes with the criteria of 2-fold higher expression in E13.5 XY supporting cells and $P < 0.05$ were termed as "Pre-sertoli genes" and shown in red.

Figure S4. H3K27me3 is involved in the female gene regulatory network in Sertoli cells. (**A**) Box-and-whisker plots showing distributions of enrichment for H3K27me3 ChIP-seq data. Central bars represent medians, the boxes encompass 50% of the data points, and the whiskers indicate 90% of the data points. *** $P < 0.0001$, Mann-Whitney U tests. (**B**) Scatter plots showing the Pearson correlation between ChIP-seq enrichment (±2 kb around TSSs) and gene expression in Sertoli cells. A linear trendline is shown in blue.

References

1. A. Swain, R. Lovell-Badge, Mammalian sex determination: a molecular drama. *Genes Dev* **13**, 755-767 (1999).

2. D. Wilhelm, S. Palmer, P. Koopman, Sex determination and gonadal development in mammals. *Physiol Rev* **87**, 1-28 (2007).

3. B. Capel, Vertebrate sex determination: evolutionary plasticity of a fundamental switch. *Nature reviews. Genetics* **18**, 675-689 (2017).

4. T. Svingen, P. Koopman, Building the mammalian testis: origins, differentiation, and assembly of the component cell populations. *Genes Dev* **27**, 2409-2426 (2013).

5. C. K. Matson *et al.*, DMRT1 prevents female reprogramming in the postnatal mammalian testis. *Nature* **476**, 101-104 (2011).

6. C. K. Matson, D. Zarkower, Sex and the singular DM domain: insights into sexual regulation, evolution and plasticity. *Nat Rev Genet* **13**, 163-174 (2012).

7. N. H. Uhlenhaut *et al.*, Somatic sex reprogramming of adult ovaries to testes by FOXL2 ablation. *Cell* **139**, 1130-1142 (2009).

8. A. Minkina *et al.*, DMRT1 protects male gonadal cells from retinoid-dependent sexual transdifferentiation. *Dev Cell* **29**, 511-520 (2014).

9. R. E. Lindeman *et al.*, Sexual cell-fate reprogramming in the ovary by DMRT1. *Curr Biol* **25**, 764-771 (2015).

10. L. Zhao, T. Svingen, E. T. Ng, P. Koopman, Female-to-male sex reversal in mice caused by transgenic overexpression of Dmrt1. *Development (Cambridge, England)* **142**, 1083-1088 (2015).

11. Y. Li *et al.*, beta-Catenin directs the transformation of testis Sertoli cells to ovarian granulosa-like cells by inducing Foxl2 expression. *The Journal of biological chemistry* **292**, 17577-17586 (2017).

12.   B. Nicol *et al.*, Genome-wide identification of FOXL2 binding and characterization of
FOXL2 feminizing action in the fetal gonads. *Hum Mol Genet* **27**, 4273-4287 (2018).

13.   Y. Katoh-Fukui *et al.*, Male-to-female sex reversal in M33 mutant mice. *Nature* **393**, 688-
692 (1998).

14.   S. A. Garcia-Moreno *et al.*, CBX2 is required to stabilize the testis pathway by repressing
Wnt signaling. *PLoS genetics* **15**, e1007895 (2019).

15.   S. Kuroki *et al.*, Epigenetic regulation of mouse sex determination by the histone
demethylase Jmjd1a. *Science* **341**, 1106-1109 (2013).

16.   L. Ringrose, R. Paro, Polycomb/Trithorax response elements and epigenetic memory of
cell identity. *Development (Cambridge, England)* **134**, 223-232 (2007).

17.   L. Aloia, B. Di Stefano, L. Di Croce, Polycomb complexes in stem cells and embryonic
development. *Development (Cambridge, England)* **140**, 2525-2534 (2013).

18.   J. A. Simon, R. E. Kingston, Occupying chromatin: Polycomb mechanisms for getting to
genomic targets, stopping transcriptional traffic, and staying put. *Molecular cell* **49**, 808-
824 (2013).

19.   M. Endoh *et al.*, Histone H2A mono-ubiquitination is a crucial step to mediate PRC1-
dependent repression of developmental genes to maintain ES cell identity. *PLoS genetics*
8, e1002774 (2012).

20.   R. W. Holdcraft, R. E. Braun, Androgen receptor function is required in Sertoli cells for
the terminal differentiation of haploid spermatids. *Development (Cambridge, England)*
131, 459-467 (2004).

21.   M. del Mar Lorente *et al.*, Loss- and gain-of-function mutations show a polycomb group
function for Ring1A in mice. *Development (Cambridge, England)* **127**, 5093-5100 (2000).

22.   S. Maezawa *et al.*, Polycomb directs timely activation of germline genes in
spermatogenesis. *Genes & development* **31**, 1693-1703 (2017).

23.   S. Yokobayashi *et al.*, PRC1 coordinates timing of sexual differentiation of female
550   primordial germ cells. *Nature* **495**, 236-240 (2013).

551   24.   E. Posfai *et al.*, Polycomb function during oogenesis is required for mouse embryonic
552   development. *Genes & development* **26**, 920-932 (2012).

553   25.   Z. Gao *et al.*, PCGF homologs, CBX proteins, and RYBP define functionally distinct PRC1
554   family complexes. *Molecular cell* **45**, 344-356 (2012).

555   26.   O. O. Oduwole, H. Peltoketo, I. T. Huhtaniemi, Role of Follicle-Stimulating Hormone in
556   Spermatogenesis. *Front Endocrinol (Lausanne)* **9**, 763 (2018).

557   27.   R. M. Sharpe, C. McKinnell, C. Kivlin, J. S. Fisher, Proliferation and functional maturation
558   of Sertoli cells, and their relevance to disorders of testis function in adulthood.
559   *Reproduction (Cambridge, England)* **125**, 769-784 (2003).

560   28.   S. Vainio, M. Heikkila, A. Kispert, N. Chin, A. P. McMahon, Female development in
561   mammals is regulated by Wnt-4 signalling. *Nature* **397**, 405-409 (1999).

562   29.   A. A. Chassot *et al.*, Activation of beta-catenin signaling by Rspo1 controls differentiation
563   of the mammalian ovary. *Hum Mol Genet* **17**, 1264-1277 (2008).

564   30.   D. Schmidt *et al.*, The murine winged-helix transcription factor Foxl2 is required for
565   granulosa cell differentiation and ovary maintenance. *Development (Cambridge, England)*
566   **131**, 933-942 (2004).

567   31.   P. Parma *et al.*, R-spondin1 is essential in sex determination, skin differentiation and
568   malignancy. *Nature genetics* **38**, 1304-1309 (2006).

569   32.   C. Ottolenghi *et al.*, Loss of Wnt4 and Foxl2 leads to female-to-male sex reversal extending
570   to germ cells. *Human molecular genetics* **16**, 2795-2804 (2007).

571   33.   V. P. Vidal, M. C. Chaboissier, D. G. de Rooij, A. Schedl, Sox9 induces testis development
572   in XX transgenic mice. *Nature genetics* **28**, 216-217 (2001).

573   34.   C. S. Raymond *et al.*, Evidence for evolutionary conservation of sex-determining genes.
574   *Nature* **391**, 691-695 (1998).

35. C. S. Raymond, M. W. Murphy, M. G. O'Sullivan, V. J. Bardwell, D. Zarkower, Dmrt1, a

gene related to worm and fly sexual regulators, is required for mammalian testis

differentiation. *Genes & development* **14**, 2587-2595 (2000).

36. H. H. Yao, W. Whoriskey, B. Capel, Desert Hedgehog/Patched 1 signaling specifies fetal

Leydig cell fate in testis organogenesis. *Genes & development* **16**, 1433-1440 (2002).

37. N. P. Blackledge *et al.*, Variant PRC1 complex-dependent H2A ubiquitylation drives PRC2

recruitment and polycomb domain formation. *Cell* **157**, 1445-1459 (2014).

38. S. Cooper *et al.*, Targeting polycomb to pericentric heterochromatin in embryonic stem

cells reveals a role for H2AK119u1 in PRC2 recruitment. *Cell reports* **7**, 1456-1470 (2014).

39. U. Jochumsen *et al.*, Mutation analysis of FOXF2 in patients with disorders of sex

development (DSD) in combination with cleft palate. *Sex Dev* **2**, 302-308 (2008).

40. H. Du, H. S. Taylor, The Role of Hox Genes in Female Reproductive Tract Development,

Adult Function, and Fertility. *Cold Spring Harb Perspect Med* **6**, a023002 (2015).

41. T. I. Lee *et al.*, Control of developmental regulators by Polycomb in human embryonic

stem cells. *Cell* **125**, 301-313 (2006).

42. S. A. Jameson *et al.*, Temporal transcriptional profiling of somatic and germ cells reveals

biased lineage priming of sexual fate in the fetal mouse gonad. *PLoS genetics* **8**, e1002575

(2012).

43. J. J. Jacobs, K. Kieboom, S. Marino, R. A. DePinho, M. van Lohuizen, The oncogene and

Polycomb-group gene bmi-1 regulates cell proliferation and senescence through the ink4a

locus. *Nature* **397**, 164-168 (1999).

44. M. Serrano *et al.*, Role of the INK4a locus in tumor suppression and cell mortality. *Cell*

85, 27-37 (1996).

45. W. Mu, J. Starmer, A. M. Fedoriw, D. Yee, T. Magnuson, Repression of the soma-specific

transcriptome by Polycomb-repressive complex 2 promotes male germ cell development.

Genes & development **28**, 2056-2069 (2014).

46.   A. Piunti *et al.*, Polycomb proteins control proliferation and transformation independently

of cell cycle checkpoints by regulating DNA replication. *Nature communications* **5**, 3649

(2014).

47.   A. Laugesen, J. W. Hojfeldt, K. Helin, Molecular Mechanisms Directing PRC2

Recruitment and H3K27 Methylation. *Mol Cell* **74**, 8-18 (2019).

48.   B. E. Bernstein *et al.*, A bivalent chromatin structure marks key developmental genes in

embryonic stem cells. *Cell* **125**, 315-326 (2006).

49.   S. S. Hammoud *et al.*, Distinctive chromatin in human sperm packages genes for embryo

development. *Nature* **460**, 473-478 (2009).

50.   B. J. Lesch, G. A. Dokshin, R. A. Young, J. R. McCarrey, D. C. Page, A set of genes critical

to development is epigenetically poised in mouse germ cells from fetal stages through

completion of meiosis. *Proceedings of the National Academy of Sciences of the United

States of America* **110**, 16061-16066 (2013).

51.   H. S. Sin, A. V. Kartashov, K. Hasegawa, A. Barski, S. H. Namekawa, Poised chromatin

and bivalent domains facilitate the mitosis-to-meiosis transition in the male germline. *BMC

biology* **13**, 53 (2015).

52.   S. Maezawa *et al.*, Polycomb protein SCML2 facilitates H3K27me3 to establish bivalent

domains in the male germline. *Proceedings of the National Academy of Sciences of the

United States of America* **115**, 4957-4962 (2018).

53.   S. Maezawa, M. Yukawa, K. G. Alavattam, A. Barski, S. H. Namekawa, Dynamic

reorganization of open chromatin underlies diverse transcriptomes during spermatogenesis.

Nucleic Acids Res **46**, 593-608 (2018).

54.   C. Cales *et al.*, Inactivation of the polycomb group protein Ring1B unveils an

antiproliferative role in hematopoietic cell expansion and cooperation with tumorigenesis

associated with Ink4a deletion. *Mol Cell Biol* **28**, 1018-1028 (2008).

55.   A. Ventura *et al.*, Restoration of p53 function leads to tumour regression in vivo. *Nature*
627   **445**, 661-665 (2007).

628   56.   Y. F. Chang, J. S. Lee-Chang, S. Panneerdoss, J. A. MacLean, 2nd, M. K. Rao, Isolation of
629   Sertoli, Leydig, and spermatogenic cells from the mouse testis. *Biotechniques* **51**, 341-342,
630   344 (2011).

631   57.   C. A. Schneider, W. S. Rasband, K. W. Eliceiri, NIH Image to ImageJ: 25 years of image
632   analysis. *Nat Methods* **9**, 671-675 (2012).

633   58.   A. V. Kartashov, A. Barski, BioWardrobe: an integrated platform for analysis of
634   epigenomics and transcriptomics data. *Genome biology* **16**, 158 (2015).

635   59.   C. Schmidl, A. F. Rendeiro, N. C. Sheffield, C. Bock, ChIPmentation: fast, robust, low-
636   input ChIP-seq for histones and transcription factors. *Nat Methods* **12**, 963-965 (2015).

637   60.   B. Langmead, C. Trapnell, M. Pop, S. L. Salzberg, Ultrafast and memory-efficient
638   alignment of short DNA sequences to the human genome. *Genome biology* **10**, R25 (2009).

639   61.   Y. Zhang *et al.*, Model-based analysis of ChIP-Seq (MACS). *Genome biology* **9**, R137
640   (2008).

A

PRC1ctrl:
Ring1-/-; *Rnf2*f/f

PRC1RcKO:
Ring1-/-; *Rnf2*f/f; *Rosa26-creERT2*Tg/+

Isolation of Sertoli cells at P7 → 4-day culture with 4-OHT → Analysis

B

Numbers of genes
338 — 1.5x upregulated in PRC1RcKO
307 — 1.5x downregulated in PRC1RcKO

C

1.5x upregulated in PRC1RcKO — ctrl dKO — 338 genes
1.5x downregulated in PRC1RcKO — ctrl dKO — 307 genes
RPKM Color Key: 80, 10, 1.25

D

1.5x upregulated in PRC1RcKO
338 genes
-log$_{10}$ (P-value)

Neuron differentiation
Skeletal system
Cell adhesion
Regionalization
Regulation of cell
Prostate gland

1.5x downregulated in PRC1RcKO
307 genes
-log$_{10}$ (P-value)

Cell cycle
M phase
Organelle fission
Chromosome
DNA replication initiation
DNA repair

E

RNA-seq
PRC1ctrl
PRC1RcKO
Log$_2$ RPKM
Sox9 *Dmrt1* *Foxl2* *Rspo1*

F

Wild-type Sertoli cells
ChIP-seq enrichment around TSSs
Normalized enrichment

H2AK1119ub
Sox9 *Dmrt1* *Foxl2* *Rspo1*

H3K27me3
Sox9 *Dmrt1* *Foxl2* *Rspo1*
Sertoli genes | Granulosa genes

A Gene expression

Upregulated
in PRC1RcKO
Sertoli cells

Downregulated
in PRC1RcKO
Sertoli cells

B Correlation of differential gene expression:
E13.5 supporting cells vs postnatal Sertoli cells

Expressed in
pre-granulosa cells

Expressed in
pre-Sertoli cells

Upregulated
in PRC1RcKO
Sertoli cells

Downregulated
in PRC1RcKO
Sertoli cells

C H2AK119ub around TSS

D Repression of pre-granulosa genes
by H2AK119ub in Sertoli cells

Upregulated
in PRC1RcKO
Sertoli cells

Downregulated
in PRC1RcKO
Sertoli cells

Gene expression in E13.5 supporting cells
by microarray (Jameson *et al*, 2012, *PLoS Gen.*)

- Pre-Granulosa genes (575 genes)
- Pre-Sertoli genes (679 genes)
- Other genes (9242 genes)

A H3K27me3 around TSS

B Repression of pre-granulosa gene
by H3K27me3 in Sertoli cells