Sahul's megafauna were vulnerable to plant-community changes due to their position in the trophic network

John Llewelyn[1] (johnsllewelyn@gmail.com), Giovanni Strona[2] (goblinshrimp@gmail.com), Matthew C. McDowell[3] (matthew.mcdowell@utas.edu.au), Christopher N. Johnson[4] (c.n.johnson@utas.edu.au), Katharina J. Peters[1] (katharina.peters@flinders.edu.au), Daniel B. Stouffer[5] (Daniel.stouffer@canterbury.ac.nz), Sara N. de Visser[6] (sndevisser@hotmail.com), Frédérik Saltré[1] (frederik.saltre@flinders.edu.au), Corey J. A. Bradshaw[1] (Corey.Bradshaw@flinders.edu.au)

[1]Global Ecology, College of Science and Engineering and ARC Centre of Excellence for Australian Biodiversity and Heritage, Flinders University, GPO Box 2100, Adelaide, South Australia 5001, Australia
[2]Research Centre for Ecological Change, University of Helsinki, Viikinkaari 1, Biocentre 3, 00790, Helsinki, Finland
[3]Dynamics of Eco-Evolutionary Pattern and ARC Centre of Excellence for Australian Biodiversity and Heritage, University of Tasmania, Tasmania 7001, Australia
[4]School of Natural Sciences and Australian Research Council Centre of Excellence for Australian Biodiversity and Heritage, University of Tasmania, Private Bag 55, Hobart, Tasmania 7001, Australia
[5]Centre for Integrative Ecology, School of Biological Sciences, University of Canterbury, Christchurch, New Zealand
[6]Community and Conservation Ecology, Centre for Ecological and Evolutionary Studies, University of Groningen, PO Box 11103, 9700 Groningen, The Netherlands

Short running title: Trophic cascades and megafauna

Key words: ecological network, coextinction, biotic interactions, extinction event, Late Pleistocene, food web

Author contributions

Project conceptualization by J.L., G.S., C.J.A.B., and F.S. Data curation by J.L., G.S., K.J.P., C.N.J., M.C.M., and S.N.d-V. Investigation and formal analysis by J.L., G.S., and C.J.A.B. Methodology by J.L., G.S., C.J.A.B., and D.B.S. Original draft preparation by J.L., G.S., C.J.A.B., and F.S. All authors contributed to reviewing and editing the draft.

Data and code accessibility

All relevant data and code are available at Github.com/JohnLlewelyn/Inferring-networks-and-modelling-trophic-cascades (DOI: 10.5281/zenodo.3756224)

Corresponding author
John Llewelyn
Email: johnsllewelyn@gmail.com
Phone: +61 429 506 956
Fax: +8 8201 3399
Address: Global Ecology Lab, College of Science & Engineering, Flinders University

Abstract

Extinctions stemming from environmental change often trigger trophic cascades and

coextinctions. However, it remains unclear whether trophic cascades were a large contributor

to the megafauna extinctions that swept across several continents in the Late Pleistocene. The

pathways to megafauna extinctions are particularly unclear for Sahul (landmass comprising

Australia and New Guinea), where extinctions happened earlier than on other continents. We

investigated the role of bottom-up trophic cascades in Late Pleistocene Sahul by constructing

pre-extinction (~ 80 ka) trophic network models of the vertebrate community of Naracoorte,

south-eastern Australia. These models allowed us to predict vertebrate species' vulnerability

to cascading extinctions based on their position in the network. We tested whether the

observed extinctions could be explained by bottom-up cascades, or if they should be

attributed to other external causes. Species that disappeared from the community were more

vulnerable, overall, to bottom-up cascades than were species that survived. The position of

extinct species in the network – having few or no predators – also suggests they might have

been particularly vulnerable to a new predator. These results provide quantitative evidence

that trophic cascades and naivety to predators could have contributed to the megafauna

extinction event in Sahul.

Introduction

Of all the extinctions that have ever occurred on Earth, many — potentially most — have

been coextinctions [1]. In some cases, these coextinctions involved host-specific parasites

that were doomed by the extinction of their host species, or flowering plants imperilled by the

extinction of their pollinators [2]. Coextinctions have also been mediated through trophic

interactions between herbivores and vegetation, and between predators and their prey [3,4].

Therefore, if we are to understand past extinction events and predict future extinctions, we

need to be able to infer coextinction cascades accurately.

Changes in the primary producer component of a community can trigger bottom-up

cascades and profoundly alter ecological communities [5]. However, it is unclear which

species are most vulnerable to bottom-up cascades. On the one hand, it has been argued that

top predators and species from high trophic levels are particularly sensitive to food-web

perturbations and reductions in habitat area/primary productivity [6–9]. However, others have

concluded that changes in the diversity of primary producers or nutrient content most

strongly affect herbivores, and the cascading effects on higher trophic levels are dampened by

trophic distance [10–13]. Similarly, extinction-risk assessments by the International Union

for Conservation of Nature (IUCN) and recorded recent extinctions suggest that herbivorous

terrestrial vertebrates are particularly vulnerable to extinction [14], a pattern that might partly

be explained by the sensitivity of lower trophic levels to bottom-up cascades. The uncertainty

regarding how vulnerability to bottom-up cascades varies with species traits (such as trophic

level) has limited our ability to assess the importance of bottom-up cascades in past

extinction events, and to predict how these cascades might unfold in the future.

Identifying the vulnerability of species to bottom-up (or top-down) coextinction cascades

relies heavily on understanding species interactions within an ecological community. To this

end, ecological network modelling is an invaluable tool for representing ecological

communities from the perspectives of species interactions and for studying the consequences

of changes in these interactions [15]. In ecological network models, organismal groups (e.g.,

species, age groups, populations, or individuals) are represented by nodes, and interactions —

which can be weighted or unweighted — are represented by links (edges). The interaction

type most frequently used to build ecological network models are trophic interactions (i.e.,

food webs). For contemporary communities, there is a growing number of studies that use

detailed information on species interactions to build network models and to study trophic

cascades [16–18]. Unfortunately, similar approaches are challenging to apply to palaeo-

communities because of the lack of data on ancient trophic interactions. However, by

combining contemporary and palaeo-data to infer trophic interactions, this limitation can be

overcome and network models of palaeo-communities can be constructed [19,20]. For

example, Pires et al. (2015) used this approach to model Late Pleistocene mammal

communities in the Americas, concluding that (1) pre-existing American mammal networks

were not especially unstable (i.e., they were similar to modern networks in Africa in terms of

population densities re-establishing after simulated perturbations), and (2) the arrival of

humans destabilized the networks because this new predator increased network connectance

(i.e., the proportion of potential links that are realized). Investigations of paleo mass

extinctions are fortified through the use of network modelling, as these methods provide

insights into the causes and consequences of extinction events. Such events can also be used

as a means of validating modelling methods because the outcomes (i.e., extinctions) are

known. Despite these opportunities, the application of network modelling to investigate

palaeo-extinction events remains under-utilised.

Megafauna (animals > 44 kg) extinctions swept across several continents during the Late

Pleistocene (126,000 – 12,000 years ago), with the highest proportions of genera lost from

Sahul (landmass including Australia and New Guinea) and the Americas [21,22]. While the causes of these extinctions are still debated, most evidence points toward the arrival of anatomically modern humans and/or climate change [23–25]. Irrespective of the root causes, large extinction events such as these always involve both primary and secondary (or co-) extinctions [3]. Indeed, it has been argued that the loss of prey species led to large predators going extinct in the Late Pleistocene [26,27]. Although the arrival of modern humans and/or climate change have been identified as the most likely ultimate causes of megafauna extinctions in the Late Pleistocene, vegetation change associated with human arrival and/or climate change has been identified as a pathway through which these ultimate causes could have triggered extinctions (i.e., via bottom-up trophic cascades) [28–31].

The Late Pleistocene megafauna assemblage of Sahul was distinct from that of other continents in that all the large mammals were marsupials or monotremes [32]. Giant reptiles and birds were also a prominent component of the continent's megafauna [33]. While Sahul's megafauna included many species over the standard body-mass threshold of 44 kg, the term 'megafauna' is often extended to include species with a body mass above that of their surviving relatives [32] — a definition we have adopted here. Identifying the pathways by which Sahul's unique megafauna were lost is challenging because their extinctions happened much earlier in Sahul than elsewhere [34]. To characterize such ancient extinction events, a sufficient number of dated fossilized remains is necessary [35]. The most detailed and well-studied fossil record spanning the megafauna extinction event in Sahul comes from the Naracoorte region in south-eastern Australia (Figure 1). This fossil record offers an exceptional picture of the species living in the region over the past 500,000 years, including the ecological community at the time of the main megafauna extinction event that occurred approximately 44,000 years ago in Naracoorte [25]. Thus, the Naracoorte fossil record is the best platform available from which to model the ecological and environmental processes potentially involved in megafauna extinctions in Sahul.

We assessed how vulnerability to bottom-up cascades varies with network-position attributes and whether bottom-up coextinction cascades stemming from the loss of basal resources (i.e., primary producers/plants) could have played a role in the megafauna extinctions of Sahul. First, we built small, synthetic networks (3 to 20 nodes) varying in topology (i.e., structure of connections), and calculated each node's coextinction vulnerability using two methods: (*i*) simulation, and (*ii*) Bayesian networks. This allowed us to test the general influence of trophic level, diet breadth, and number of connected basal resources on vulnerability to bottom-up coextinction cascades.

Next, we used Naracoorte as a model system to assess whether bottom-up coextinction

cascades could explain which species went extinct during the Late Pleistocene event. We

constructed an entire terrestrial, palaeo-vertebrate assemblage (including all terrestrial

vertebrate classes), and combined this assemblage with palaeo and contemporary data to infer

trophic interactions and build network models (Figure 2). These network models consisted of

nodes (species) with directed, unweighted links. We then computed each species'

vulnerability to coextinction via bottom-up cascades using the simulation method we

validated with the synthetic networks, and we compared the coextinction vulnerabilities (and

the traits influencing them) between extinct and extant (surviving into the Holocene) species.

In addition to vulnerability to bottom-up coextinction cascades, we also tested for differences

in the network positions of extant and extinct species to determine if relative position could

have made extinct species more vulnerable in other ways (e.g., more vulnerable to a new

predator). By incorporating interactions involving all vertebrate species, we adopted a holistic

approach to studying megafauna extinctions of the Late Pleistocene.

We found that vulnerability to bottom-up cascades decreases with increasing trophic

level, diet breadth, and basal connections, and that extinct species were more vulnerable to

bottom-up coextinction cascades than were extant species. This suggests that bottom-up

trophic cascades possibly contributed to the megafauna extinction event in Sahul. Our results

also indicate that extinct species had fewer predators than did surviving species, suggesting

that extinct species might have been particularly sensitive to the arrival of the new predator

Homo sapiens.

Methods

We aimed to: (1) identify how vulnerability to bottom-up cascades varies with trophic level,

diet breadth, and basal connections using synthetic networks, and (2) develop and interrogate

ecological network models representing the Naracoorte ecosystem prior to the main

megafauna-extinction pulse and the arrival of humans in the region (~ 44,000 years ago) [25].

Below, we describe how we generated the synthetic networks, measured node vulnerability to

bottom-up cascades, built ecologically realistic models of the Naracoorte network, and tested

whether bottom-up coextinction cascades could explain which species were lost from the

Naracoorte network (see Supplementary Figure S1 for a flowchart of methods for the

construction and analysis of the Naracoorte network).

Synthetic networks

We randomly generated 196 'synthetic' networks that differed in topology. The networks

varied in size (number of nodes: 3 to 20), number of links (2 to 92), and connectance (0.07 to

0.33). Within these networks, nodes varied in trophic level (1 to 4), number of 'in' links (1 to

8), and the number of basal nodes to which non-basal nodes were directly or indirectly

connected through 'in' links (1 to 8). We generated these networks to test how a node's

vulnerability to the removal/extinction of basal nodes (analogous to plants/primary

producers) varied depending on the node's trophic level, diet breadth (number of 'in' links),

and the number of basal resources to which it was connected.

Coextinction vulnerability in synthetic networks

We inferred vulnerability of nodes to bottom-up cascades via: (*i*) simulations, and (*ii*) a

Bayesian network method [36]. We applied two different approaches because there is

uncertainty regarding the best methods to infer vulnerability to bottom-up cascades [3], so

consensus using both approaches would provide more robust results than relying on only one.

In the simulation method, primary extinctions occur by randomly removing basal nodes from

the network, after which coextinctions are simulated by removing nodes that had lost all their

'in' links. Based on 1000 iterations of each network, we calculated the average coextinction

vulnerability of each node as the proportion of total basal resources remaining when

coextinction occurred. Applying the Bayesian-network method to the same networks [36],

baseline extinction probabilities are assigned to each node and then each node's accumulative

extinction vulnerability is calculated using the network structure (taking into account each

node's dependencies on other nodes). In our case, we adjusted baseline probabilities of

extinction so that primary extinctions were restricted to basal nodes.

Analysis of synthetic networks

We fit mixed-effects models to the results from the simulation and Bayesian network

approaches to assess if the two methods yielded similar results in terms of the effects of

trophic level, diet breadth, and basal connections on vulnerability to bottom-up coextinction

cascades. Prior to fitting, we scaled the independent variables so that the units of the

regression coefficients were the same for all variables. The full/global mixed-effects model

had coextinction vulnerability as the dependent variable (response), trophic level, diet

breadth, number of connected basal nodes, and the interactions between these traits as

independent variables (fixed effects), and network identity as a random effect. We compared

support for the full models with every combination of nested/reduced model using Akaike's

information criterion weights ($w\text{AIC}_c$). If including trophic level, diet breadth, and number of

basal connections resulted in models with higher $w\text{AIC}_c$, this suggests these variables affect

vulnerability. We calculated model-averaged (via $w\text{AIC}_c$) coefficients for the independent

variables to identify how each variable influenced vulnerability. We also extracted marginal

R^2 from three mixed effects models that had either trophic level, diet breadth, or basal

connections as the only independent variable to estimate how much variation in vulnerability

each of these variables explained.

Naracoorte study region

The World Heritage-listed (from 1994) Naracoorte Caves in south-eastern South Australia

(37° 02′ 24″ S, 140° 48′ 00″ E) encompass a series of limestone caves that opened and closed

to the surface at different locations, and at different times, over the last 500,000 years [37].

These openings acted as natural pitfall traps, capturing snapshots of Naracoorte's biodiversity

at different periods from at least 500,000 years ago to the present. In addition to the fossils of

the many animals that fell into these natural traps, there are remains of species that lived in

the caves, such as owls and bats, and their prey [38]. Consequently, the Naracoorte Caves

provide an ideal platform from which to build palaeo-ecological network models to gain

insight into how these long-lost ecosystems functioned and changed over time.

Species data

To build a species assemblage list (i.e., to identify the nodes to include in the network

models), we used two data sources: *FosSahul* 2.0 [35,39] and the Atlas of Living Australia.

FosSahul 2.0 is a database of dated fossil records from Sahul, including an automated

quality-rating method for date reliability [40]. We extracted and vetted records from the

Naracoorte region (defined as the region between 35° 32′ 48″ S and 38° 6′ 50″ S, and

between 139° 10′ 42″ E and 141° 0′ 21″ E) from *FosSahul* 2.0 that had high- or intermediate-

quality dates (A*, A or B) [35], and whose age was younger than 200,000 years before

present. We chose this cut-off age to provide a large enough period to include dated fossils

from all/most megafauna species that lived in the region immediately prior to the main

extinction event; if we had made the period too narrow, few megafauna species would have

been captured despite their likely presence in the region at the time. Fossil records suffer

from taphonomic biases (biases in the accumulation and preservation of different organisms),

and, consequently, some species that were present in Late Pleistocene Naracoorte are unlikely

to be represented in the fossil record. Furthermore, there are biases for studying and dating

particular groups of species due to academic and amateur interests [41]. These biases, along

with the fact that *FosSahul* was primarily designed to document megafauna remains rather

than smaller species, means that *FosSahul* 2.0 does not include all vertebrate species present

in Late Pleistocene Naracoorte. To account for this gap in the species list, we supplemented

FosSahul 2.0 data with contemporary and historical species records from the Naracoorte

Coastal Plains bioregion from the Atlas of Living Australia online repository (ala.org.au;

accessed 3 January 2019). The Atlas of Living Australia has detailed species records of

vertebrates from this bioregion, and so it captures most of the diversity of extant and recently

extinct vertebrate species. We extracted data pertaining to all terrestrial vertebrates from the

region, and removed species that were introduced since European arrival, as well as vagrants

and erroneous records (species well-outside their normal distribution and not present in the

fossil record), and strictly coastal species such as marine birds that do not use inland

waterways. Our final species list, built using *FosSahul* 2.0 and the Atlas of Living Australia,

included 280 birds, 81 mammals, 50 reptiles, and 12 amphibians (Supplementary Table S1).

The fossil record and phylogeography of extant mammal species (i.e., the vertebrate class

most intensively studied) from the Naracoorte region suggest that, at a fine spatial scale, the

distributions of some species expanded and contracted with climatic fluctuations during the

Late Pleistocene and Holocene, but that during times of contraction species persisted at a

regional scale (as in our study) in refugia [42–44]. Furthermore, most extant mammals

recorded in the Late Pleistocene fossil record at Naracoorte were present (living) in the

Naracoorte bioregion when Europeans arrived, suggesting little species turnover (apart from

megafauna) between the Late Pleistocene and European arrival [42–44]. Together, these

results suggest that our approach of using the fossil record and modern presence data

provides a reasonable estimate of the species likely present in the region in the Late

Pleistocene.

To infer trophic links, we required information on each species' body mass and broad

trophic category (whether it ate plants and/or fungi, invertebrates, terrestrial vertebrates, or

fish). For extant species, we extracted much of this information from large databases: snake

database [45]; Australian bird database [46]; *PanTHERIA* (mammal database) [47]; lizard

database [48]; *AmphiBIO* (amphibian database) [49]; tropical bird database [50]; Amphibian

database [51]; reptile database [52]; and Elton traits databases (mammals and birds) [53].

However, information on body mass was not available for some extant reptiles and

amphibians. For these species, we estimated body mass using their body length (from field

guides) and validated allometric relationships [45,48,51,54,55]. Extinct megafaunal species were not included in any of the aforementioned databases, so for these species we obtained body mass and diet data from the literature (see Supplementary Table S2). The 423 vertebrate species in the Naracoorte assemblage included 125 that consumed vertebrates, 249 that consumed plants, 362 that consumed invertebrates, and 48 that consumed fish (Supplementary Table S1). Of the 423 species, 273 consumed more than one of these resource groups. Mean species body mass ranged from 0.4 g to 2700 kg (Supplementary Table S1).

Inferring trophic links

For almost all extant species, information on trophic interactions is incomplete, but the lack of knowledge regarding trophic interactions is even greater for long-extinct species. To overcome this limitation and build realistic ecological network models, various methods have been developed for inferring trophic links using species' traits such as body size (i.e., larger predators tend to eat larger prey, and predators are usually bigger than their prey) [56]. Others have built on this approach to improve prediction accuracy [57–60] by adding phylogenetic and physiological information, prohibiting impossible or unlikely links, incorporating specific morphological features such as biting force and cuticular thickness, and taking into account abundance. However, most of this research focusses on fish or invertebrates, with the effectiveness of these methods rarely applied or validated for terrestrial vertebrates, but see [61,62].

   We therefore developed and validated a new method, based on the body-size trophic-niche model [56], to infer trophic links between terrestrial vertebrates. The body-size trophic-niche model that we adapted consists of two quantile regressions: (*i*) one defining the upper prey-size limit given predator mass, and (*ii*) the other defining the lower prey-size limit given predator mass. If a species falls within the upper and lower limits for a particular predator, it is inferred as potential prey for that predator. We used a large predator-prey interaction dataset to identify these body-size relationships between terrestrial predators and their prey, and tested whether these relationships varied depending on the predator's taxonomic class (i.e., did including predator class as an independent variable improve the fit of the body size quantile regressions?).

   We extracted the interaction dataset from *GloBI*, an online repository of biotic interactions [63]. The dataset consisted of 3893 records: 958 records of predation by non-marine mammals, 2711 by birds, 199 by reptiles, and 25 by amphibians (Supplementary Table S3)

[64]. We extracted data on mean body mass for the species from the same databases we used
to add this information to the Naracoorte species list (see above). Once we identified the best
trophic-niche model (i.e., the combination of quantile regressions that best fit the upper and
lower prey-size limits) using the entire *GloBI* dataset and Bayesian information criterion
(BIC; Supplementary Methods S1), we validated this method of assigning trophic links by
comparing model performance when applied first to the *GloBI* data divided into training and
validation datasets, and then to a well-resolved trophic network from the Serengeti [65] (see
also S. de Visser unpublished data; Supplementary Methods S1). We used the true skill
statistic to evaluate model performance and found that the top-ranked model (according to
BIC) also performed best at assigning links in the validation step (Supplementary Methods
S1, Supplementary Table S4, Supplementary Figure S2, script available at [64]). We used the
best trophic-niche model to identify potential prey for each predator of vertebrates (see
methods below; Supplementary Methods S1; script available [64]).
While including additional species traits could improve the accuracy of inferred predator-
prey interactions, we used only three readily available traits (broad diet, body size, and
predator taxonomic class — with reptiles and amphibians grouped together). We used only
these three traits because: (1) using few traits is compatible with the quantile regression
framework, (2) they can easily be extracted for vertebrate species in most assemblages (and
therefore the method can be widely applied), and (3) our validation steps demonstrated the
resulting performance of the trophic-niche model is sufficient (true skill statistic [TSS] = 0.6
when applied to the Serengeti assemblage; TSS varies from -1 to 1, with a score of 0
indicating no better than random; Supplementary Table S5).
340
Naracoorte networks
To build realistic ecological networks for the Naracoorte assemblage, we applied the trophic-
niche model to the species list, removed excess links between vertebrates (to account for
overestimating the number of predator-prey links), and added links to non-terrestrial
vertebrate food resources. However, there is uncertainty regarding which vertebrate
predator/prey links to delete as well as how many links to add from non-terrestrial vertebrate
food resources to vertebrates. To address this uncertainty, we used a randomization approach
in the link-removal and -addition steps described below, and generated 1000 versions of the
network. That is, we randomly removed (for the vertebrate predator-prey links) or added (for
the herbivores, invertivores, and piscivores) links in the range indicated as realistic based on
contemporary species' diet breadths. Because we do not know exactly where to add or

remove these links, we generated 1000 versions of the network so our results were not

skewed by the particular links selected.

We used the trophic-niche model to assign potential predator-prey links in the Naracoorte

species assemblage. Although trophic-niche models are good at identifying potential links,

they almost always overestimate the number of realized links [66]. This is because predators

are unlikely to consume all prey within their size range — some species are not palatable, are

dangerous, too rare, difficult to capture, use different microhabitats, or have other ecological

characteristics that make them unsuitable for regular consumption [66]. To build a network

with a more realistic structure, we assigned a probability to each interaction based on the

prey's position in the predator's prey-size range and a Gaussian distribution centred on this

range (with a standard deviation equal to one quarter of the predator's prey-size range); the

highest probability was for prey close to the centre of the prey-size range (i.e., centre of the

distribution) and decreased the closer the prey was to the predator's limits. For each predator,

we randomly sampled from a density kernel fit to published carnivore diet breadths ($n = 12$;

sampled between 1 and twice the maximum diet breadth in Supplementary Table S6a;

Supplementary Methods S1) to select the number of potential prey that were 'realized' prey.

In assigning the sampled diet breadths to individual predators, predators with more potential

prey (indicated by the trophic-niche model) were assigned larger diet breadths than were

those with fewer potential prey. To account for different degrees of dietary specialisation, we

Poisson-resampled the number of potential prey for each predator before ranking predators

according to their number of potential prey, slightly shuffling relative diet breadths between

network models. Then, using the assigned diet breadths, we randomly selected from the

potential prey, taking into account the probability of the predator-prey interaction. This

method resulted in a vertebrate network with realistic connectance (proportion of potential

links that are realized), and with most of each predator's prey closer to the centre, rather than

the limits, of their prey-size range.

Terrestrial vertebrates not only consume other terrestrial vertebrates, they also consume

invertebrates, plants, fungi, and fish. In addition to inferring trophic links among terrestrial

vertebrates, we therefore needed to add links to vertebrates from these other food resources.

However, we did not include detail on individual species within these resource groups

because: (*i*) our study focusses on terrestrial vertebrate species; (*ii*) invertebrate, plant, and

fungal diversities are not well resolved for most ecosystems (including for Late Pleistocene

Naracoorte); and (*iii*) fish only constitute a small part of the Naracoorte community in terms

of biodiversity and biomass. We therefore generated a pool of *n* species for each of these

groups (n = 1300 for plants, 6000 for invertebrates, and 23 for freshwater fish), with the

number of invertebrate and plant species calculated based on the described diversity in these

groups relative to terrestrial vertebrate diversity in Australia [67], and the number of fish

determined by the diversity of freshwater fish recorded in the Naracoorte Coastal Plains

bioregion in Atlas of Living Australia (ala.org.au; accessed 6 April 2021). To assign links to

vertebrates from species in these groups, we used published records of vertebrate diet breadth

for 20 herbivores, 6 invertivores, and 9 piscivores (Supplementary Tables S6a and 6b). We fit

kernel densities to the invertivore, herbivore, and piscivore diet breadth data, and randomly

sampled (within a diet breadth range of 1 to twice the maximum diet breadth recorded for that

trophic guild in Supplementary Table S6a and S6b) from these distributions to assign the

number of links between each resource group and vertebrate consumer in each of the 1000

network models. However, because the published piscivore diet breadths (Supplementary

Table S6b) came from terrestrial Australian predators that are not exclusively piscivorous, we

multiplied the number of fish in the diets of these predators by the number of resource groups

from which the predator fed before fitting and sampling from the kernel densities. This

modification was made to offset the adjustment accounting for inflated diet breadths in

omnivores (described below).

For vertebrates that fed from more than one group (i.e., omnivores, which represent over

half the vertebrates in this network), we proportionally adjusted the number of 'in' links

(food resources) depending on from how many food groups they fed. For example, if a

species consumed from two groups (e.g., vertebrates and invertebrates), we randomly deleted

half of the links from each group; if they fed from three groups, we randomly removed two-

thirds of the links from each food group (and so on). We made these deletions to prevent

omnivores from having inflated diet breadths.

In some cases involving water birds, we used empirical evidence to avoid assigning

unrealistic trophic links. For example, pelicans are large and, consequently, the trophic-niche

model predicts that pelicans take large vertebrate prey. However, we know that pelicans are

not birds of prey and do not consume large terrestrial animals. Thus, we restricted the

allocation of trophic links for such birds to prevent them from feeding on unrealistically large

terrestrial vertebrates (they were prevented from consuming prey that weighed over a third of

their body mass).

We completed the networks by generating links from plants to invertebrates and from

plants to fish. To determine diet breadth for these herbivores, we sampled from a Pareto

distribution (alpha = 1.02, truncated at 52) following ref [68]. The alpha and truncation

values were based on a temperate woodland system (i.e., similar to Naracoorte) [68–70].

Analysis of Naracoorte models

We estimated the coextinction vulnerability of Naracoorte vertebrates using the same

simulation approach as we applied to the synthetic networks i.e., plant nodes were iteratively

(and randomly) removed to simulate primary extinctions, and coextinctions were triggered

when animals lost all their food resources. We measured each species' coextinction

vulnerability as the proportion of plant nodes remaining when coextinction occurred. We

repeated the simulations 1000 times for each of the 1000 network models to test whether: (1)

the effects of trophic level, diet breadth, and basal connections (direct and indirect) on

coextinction vulnerability were consistent with results obtained from the synthetic models,

and (2) vulnerability to bottom-up cascades differed between extinct and extant species.

To test whether the coextinction vulnerability results were consistent with those from the

synthetic models, we fit linear-regression models to the data with vulnerability to

coextinction cascades as the response, and trophic level, diet breadth, basal connections, and

the interactions between these variables as independent variables (fixed effects). Rather than

using the raw data from the 1000 network models, we used each species mean value for each

trait and vulnerability for these analyses. We then compared relative model probabilities

(wAIC$_c$) for the full model to all possible reduced models, and examined the coefficients of

the main effects to determine if they were similar to those obtained from the synthetic

networks.

We compared susceptibility of Naracoorte's extinct *versus* extant species to bottom-up

coextinction cascades in three steps. First, we compared the groups in terms of their trophic

level, diet breadth, and number of basal connections to determine if differences in these

variables suggest one group would be more vulnerable than the other. Second, to test for an

association between coextinction vulnerability and extinction status, we compared wAIC$_c$

support for two models: a null model with vulnerability to coextinction as the dependent

variable and no fixed effects *versus* a model that was identical to the first model, except with

extinction status as an independent variable. Third, we ran a randomization test to assess the

probability that extinct species were more vulnerable to coextinction than were extant

species. Here, we sampled the coextinction vulnerabilities of 10 extinct and 10 extant species

from each of the 1000 models 20 times (i.e., using the raw data rather than species means),

and each time calculated the mean difference in coextinction vulnerability between the two

groups.

We also assessed the position of extinct *versus* extant species in the network using 12

different network metrics: *trophic level, pageRank*, *betweenness centrality*, *eigenvector*

centrality, *closeness centrality (in)*, *coreness (in)*, *degree (in)*, *eccentricity (in)*, *closeness*

centrality (out), *coreness (out)*, *degree (out)*, and *eccentricity (out)* (Supplementary Table S7

for metric descriptions). We calculated the metrics followed by an 'in' or 'out' for each node

using links pointing towards (in) or away from (out) the focal node. We chose these metrics

because they are commonly used, node-level metrics describing position in the network. For

each metric, we calculated the species' mean value across the 1000 network models. After

checking for highly correlated metrics and removing those identified as redundant (i.e.,

metrics that had a Spearman's $\rho > 0.8$), we ran a principal component analysis and visually

inspected for grouping of extinct and extant species. The reduced list of metrics included

closeness centrality (out), *eccentricity (out)*, *degree (in)*, *coreness (in)*, *betweenness*, and

PageRank.

Results

Synthetic networks

The two methods calculating bottom-up coextinction vulnerability (simulation and Bayesian

network) yielded similar results in terms of the effects of trophic level, diet breadth, and basal

connections on node vulnerability to bottom-up coextinction cascades. $w\text{AIC}_c$ indicated that

the full models (i.e., that had all three independent variables and their interactions) were more

strongly supported than reduced models using both approaches, with the full models having

$w\text{AIC}_c > 0.999$ (Supplementary Tables S8 and S9). The weighted model-averaged

coefficients describing the relationships between the three independent variables and

coextinction vulnerability were negative, irrespective of which of the two methods we used to

calculate vulnerability (Figure 3A; Supplementary Tables S8 and S9). These negative

correlations indicate that vulnerability to bottom-up cascades decreased with increasing

number of basal connections, diet breadth, and trophic level. Marginal R^2 of the three reduced

models, each of which had one of the three variables as a main effect, suggest that the

number of basal connections explained most of the variation in coextinction vulnerability in

the synthetic networks (Figure 3B). We restricted our analyses of the Naracoorte network

models to the simulation method because both approaches yielded similar results, and

because the Bayesian network method was prohibitively time consuming for networks of the

size of the Naracoorte models.

Naracoorte network

In terms of the effects of trophic level, diet breadth, and basal connections on vulnerability,

the patterns in the Naracoorte network were similar to those from the synthetic networks

(Supplementary Table S10). wAIC$_c$ strongly supported the full model over reduced models

(wAIC$_c \approx 1$ for the full model; Supplementary Table S10), and the three main effects were

negatively correlated with coextinction vulnerability (Figure 4A; Supplementary Table S10).

Reduced models that had either trophic level, diet breadth, or basal connections as the only

independent variable had $R^2 > 0.23$, indicating that each of these variables were associated

with a substantial proportion of variation in vulnerability (Figure 4B).

Extinct species had fewer basal connections, narrower diet breadths, and came from lower

trophic levels, on average, than those species that survived into the Holocene (Figure 5A-C;

mean basal connections: 298 *vs.* 515; diet breadth: 34 *vs.* 79; and trophic level: 2.3 *vs.* 2.9 for

extinct *versus* extant species). Extinct species also had higher coextinction vulnerability than

did surviving species (Figure 5D; mean coextinction vulnerability ± 95% confidence interval:

0.045 ± 0.015 *vs.* 0.008 ± 0.002 for extinct and extant species, respectively), a result

consistent with coextinction vulnerability being higher for species with fewer basal

connections, narrower diet breadth, and from lower trophic levels.

To test for an association between species' extinction status and vulnerability to bottom-up

cascades, we compared support for a model with coextinction vulnerability as the dependent

variable and extinction status as the only fixed effect *versus* a null model (a random-intercept

model with no fixed effects), and we also did a randomization test comparing the

coextinction vulnerability of extinct *versus* extant species. Both approaches indicated that

extinction status was associated with vulnerability to bottom-up cascades, with extinct

species more vulnerable than those that survived into the Holocene. The wAIC$_c$ for the model

with extinction status as a fixed effect was ~ 1, indicating that this model was strongly

supported over the null model. The randomization test showed that extinct species had a

probability of 0.91 of being more vulnerable to bottom-up coextinctions than extant species

(Figure 6).

Extinct species differed from extant species in terms of their position in the network.

Principal component analysis of six network-position metrics showed that extinct and extant

species fell into two distinct groups according to the second principal component (dimension

2 in Figure 7A). The main contributors to this principal component are metrics describing a
node's connection to the network through its 'out' links, including the closeness centrality
(out) and eccentricity (out) metrics (Figure 7A; Supplementary Figures S3 and S4). Closer
examination of the out links (i.e., number of predators) showed that, on average, extinct
species had < 1 predator, whereas extant species had > 3 (Figure 7B; mean number of
predators: 0.2 *vs.* 3.3 for extinct *vs.* extant species). Indeed, the average number of predators
was lower for extinct than extant species in all 1000 models of the Naracoorte network.
526
Discussion
Our analyses demonstrate that a species' vulnerability to bottom-up coextinction cascades
varies depending on its trophic level, diet breadth, and number of basal connections. We also
found that the position of extinct species in the Naracoorte network — being primarily
herbivorous (low trophic level), and therefore having relatively narrow diet breadths and few
pathways to basal resources — might have made them more vulnerable to bottom-up
coextinction cascades than were co-occurring species that survived into the Holocene. The
Naracoorte results suggest that trophic cascades potentially contributed to the megafauna
extinction event in south-eastern Sahul. However, the extinct species from Naracoorte also
had fewer predators than did extant species, a network position attribute that would likely
have made them more vulnerable to the arrival of the new 'super predator' — humans [71].
The Naracoorte and synthetic network models revealed that vulnerability to bottom-up
coextinction cascades precipitated by plant extinctions decreased with increasing trophic
level, diet breadth, and number of basal connections (Figures 3 and 4; Supplementary Tables
S8, S9 and S10). Our results therefore support previous findings that species with narrower
diet breadths/fewer pathways to basal resources are more vulnerable to bottom-up
coextinction cascades [72–74]. However, previous research does not provide a clear
expectation regarding the relationship between trophic level and extinction vulnerability. It is
often assumed that higher trophic levels are more vulnerable to extinction than are lower
levels due to the cumulative effects of disturbances on lower trophic levels (on which higher
trophic levels depend), and direct persecution by humans [8,9]. However, our analyses that
specifically tested for sensitivity of species to primary extinctions in the plant community
imply that vulnerability to these bottom-up cascades in fact *decreases* with trophic level.
Consistent with these results, several manipulative experiments of ecological communities

have revealed that changes in the plant component of the community most strongly affect

herbivores, and impacts on higher trophic levels diminish with trophic distance [10,13,75].

This pattern has also been identified in theoretical studies. Applying Rosenzweig–

MacArthur models and synthetic (but ecologically feasible) networks, the loss of primary

producers triggered extinctions in herbivores before doing so in carnivores, and herbivores

were more vulnerable to these cascades than were carnivores [74]. However, our

vulnerability scores were based on coextinction being triggered when a consumer lost all

food resources. Coextinctions could be triggered at lower thresholds and/or vary between

species. While the congruence between our results and those from previous studies support

the methods and threshold we used, further investigation into how coextinction threshold

influences the effect of network position on node vulnerability, as well as how coextinction

thresholds covary with species/community traits, is needed to refine methods for predicting

the probability and magnitude of bottom-up cascades.

The extinction pattern observed in the Naracoorte assemblage could have been the result

of bottom-up cascades triggered by changes in the plant community, as demonstrated by our

vulnerability modelling. This leads to the question: did vegetation change at the same time as

the megafauna disappeared? Studies in other regions of Sahul have detected shifts in

vegetation roughly coinciding with, or immediately preceding, megafauna extinction.

Hypothesised drivers of these vegetation shifts include land-use changes associated with

human arrival (i.e., fire-stick farming) [28,76] and/or climate change [77]. However, there are

no detailed studies on the vegetation of Naracoorte spanning the Late Pleistocene extinction

event (but see [78] for a review of broad proxies of vegetative change over this period). The

megafauna's disappearance from Naracoorte did, nonetheless, coincide with an unusually

cool period (Supplementary Figure S5) and the arrival of humans (~ 44,000 years ago) [25],

offering the intriguing possibility that changes in climate and/or land use triggered shifts in

vegetation that had consequences for higher trophic levels in the network. To evaluate this

possibility, more studies are required to model vegetation changes in south-eastern Sahul

(including the Naracoorte region), and these must be validated using the pollen record and/or

other fossil data.

By considering the network position of all vertebrate species in the assemblage, a clear

difference between extinct and extant species emerged — extinct species had fewer predators

than did species that survived (mean number of predators: 0.2 *versus* 3.3 for extinct *versus*

extant species, respectively; Figure 4B). This predator naivety, coupled with the species'

slow life histories, likely made megafauna especially vulnerable to new predators [34,79–81]

and suggests that hunting by humans could have adversely affected megafauna. Thus, a

network modelling approach to assessing extinction vulnerability suggests that bottom-up

and/or top-down processes could have selectively removed the now-extinct species from the

Naracoorte community. However, there remains scope to address uncertainties regarding the

structure of the Naracoorte network and the methods for estimating vulnerability to

ecological cascades. As palaeo-vegetation, invertebrate, trophic (including detailed

information on the diets of extinct species), and climate data improve, network models can

incorporate this information to build more refined networks, and include more detailed top-

down and bottom-up forcings to assess the plausibility of the different potential causes and

pathways to extinction — including what (if any) role humans played in the megafauna's

demise.

In summary, our network modelling of Late Pleistocene Naracoorte suggests bottom-up

coextinction cascades and/or predator naivety and the arrival of humans could have

contributed to the megafauna extinction event in Sahul. Indeed, that our network models

showed that extinct species were vulnerable to bottom-up cascades *and* new predation

pressures lends support to recent research suggesting that climate change (that can shift

vegetation and lead to bottom-up cascades) and human arrival together drove the megafauna

extinction trajectories in much of south-eastern Sahul [25]. Our research, along with other

recent studies [19,62,82], demonstrates that network modelling is a powerful tool for

investigating and understanding ancient extinction events. By developing methods to model

whole-community responses to change and validating these methods using ancient extinction

events, we can also provide better estimates of the fates of contemporary communities as the

sixth mass-extinction event unfolds [83].

Acknowledgements

We thank E. Reed, C. Carbone, M. Tucker, C. Dickman and V.K. Llewelyn for their

constructive input.

Funding

This work was supported by the Australian Research Council Centre of Excellence for

Australian Biodiversity and Heritage (grant number CE170100015).

References

1. Dunn RR, Harris NC, Colwell RK, Koh LP, Sodhi NS. 2009 The sixth mass coextinction: are most endangered species parasites and mutualists? *Proc. R. Soc. B* **276**, 3037–3045. (doi:10.1098/rspb.2009.0413)

2. Koh LP, Dunn RR, Sodhi NS, Colwell RK, Proctor HC, Smith VS. 2004 Species coextinctions and the biodiversity crisis. *Science* **305**, 4. (doi:10.1126/science.1101101)

3. Colwell RK, Dunn RR, Harris NC. 2012 Coextinction and persistence of dependent species in a changing world. *Annu. Rev. Ecol. Evol. Syst.* **43**, 183–203. (doi:10.1146/annurev-ecolsys-110411-160304)

4. Estes JA *et al.* 2011 Trophic downgrading of Planet Earth. *Science* **333**, 301–306. (doi:10.1126/science.1205106)

5. Kagata H, Ohgushi T. 2006 Bottom-up trophic cascades and material transfer in terrestrial food webs. *Ecol. Res.* **21**, 26–34. (doi:10.1007/s11284-005-0124-z)

6. Schoener TW. 1989 Food webs from the small to the large: the Robert H. MacArthur award lecture. *Ecology* **70**, 1559–1589. (doi:10.2307/1938088)

7. Didham RK, Lawton JH, Hammond PM, Eggleton P. 1998 Trophic structure stability and extinction dynamics of beetles (Coleoptera) in tropical forest fragments. *Phil. Trans. R. Soc. Lond. B* **353**, 437–451. (doi:10.1098/rstb.1998.0221)

8. Duffy JE. 2003 Biodiversity loss, trophic skew and ecosystem functioning. *Ecol. Lett.* **6**, 680–687. (doi:10.1046/j.1461-0248.2003.00494.x)

9. Purvis A, Gittleman JL, Cowlishaw G, Mace GM. 2000 Predicting extinction risk in declining species. *Proc. R. Soc. B* **267**, 1947–1952. (doi:10.1098/rspb.2000.1234)

10. Scherber C *et al.* 2010 Bottom-up effects of plant diversity on multitrophic interactions in a biodiversity experiment. *Nature* **468**, 553–556. (doi:10.1038/nature09492)

11. Dyer LA, Letourneau D. 2002 Top-down and bottom-up diversity cascades in detrital vs. living food webs: Diversity cascades in a tropical system. *Ecol. Lett.* **6**, 60–68. (doi:10.1046/j.1461-0248.2003.00398.x)

12. Gruner DS. 2004 Attenuation of top-down and bottom-up forces in a complex terrestrial community. *Ecology* **85**, 3010–3022. (doi:10.1890/04-0020)

13. Kagata H, Nakamura M, Ohgushi T. 2005 Bottom-up cascade in a tri-trophic system: different impacts of host-plant regeneration on performance of a willow leaf beetle and its natural enemy. *Ecol. Entomol.* **30**, 58–62. (doi:10.1111/j.0307-6946.2005.00667.x)

14. Atwood TB, Valentine SA, Hammill E, McCauley DJ, Madin EMP, Beard KH, Pearse WD. 2020 Herbivores at the highest risk of extinction among mammals, birds, and reptiles. *Sci. Adv.* **6**, eabb8458. (doi:10.1126/sciadv.abb8458)

15. Säterberg T, Sellman S, Ebenman B. 2013 High frequency of functional extinctions in
ecological networks. *Nature* **499**, 468–470. (doi:10.1038/nature12277)

16. Boit A, Martinez ND, Williams RJ, Gaedke U. 2012 Mechanistic theory and modelling of
complex food-web dynamics in Lake Constance. *Ecol. Lett.* **15**. (doi:10.1111/j.1461-
0248.2012.01777.x)

17. Coll M, Lotze HK, Romanuk TN. 2008 Structural degradation in Mediterranean sea food
webs: testing ecological hypotheses using stochastic and mass-balance modelling.
Ecosystems **11**, 939–960. (doi:10.1007/s10021-008-9171-y)

18. Pocock MJO, Evans DM, Memmott J. 2012 The robustness and restoration of a network
of ecological networks. *Science* **335**, 973–977. (doi:10.1126/science.1214915)

19. Dunne JA, Labandeira CC, Williams RJ. 2014 Highly resolved early Eocene food webs
show development of modern trophic structure after the end-Cretaceous extinction.
Proc. R. Soc. B **281**, 20133280–20133280. (doi:10.1098/rspb.2013.3280)

20. Roopnarine PD. 2006 Extinction cascades and catastrophe in ancient food webs.
Paleobiology **32**, 1–19. (doi:10.1666/05008.1)

21. Barnosky AD. 2004 Assessing the causes of Late Pleistocene extinctions on the
continents. *Science* **306**, 70–75. (doi:10.1126/science.1101476)

22. Koch PL, Barnosky AD. 2006 Late Quaternary extinctions: state of the debate. *Annu. Rev.*
Ecol. Evol. Syst. **37**, 215–250. (doi:10.1146/annurev.ecolsys.34.011802.132415)

23. Lorenzen ED *et al.* 2011 Species-specific responses of Late Quaternary megafauna to
climate and humans. *Nature* **479**, 359–364. (doi:10.1038/nature10574)

24. Metcalf JL *et al.* 2016 Synergistic roles of climate warming and human occupation in
Patagonian megafaunal extinctions during the Last Deglaciation. *Sci. Adv.* **2**, e1501682.
(doi:10.1126/sciadv.1501682)

25. Saltré F, Chadoeuf J, Peters KJ, McDowell MC, Friedrich T, Timmermann A, Ulm S,
Bradshaw CJA. 2019 Climate-human interaction associated with southeast Australian
megafauna extinction patterns. *Nat. Commun.* **10**, 5311. (doi:10.1038/s41467-019-
13277-0)

26. Galetti M *et al.* 2018 Ecological and evolutionary legacy of megafauna extinctions:
anachronisms and megafauna interactions. *Biol. Rev.* **93**, 845–862.
(doi:10.1111/brv.12374)

27. Ripple WJ, Van Valkenburgh B. 2010 Linking top-down forces to the Pleistocene
megafaunal extinctions. *BioScience* **60**, 516–526. (doi:10.1525/bio.2010.60.7.7)

28. Miller GH. 2005 Ecosystem Collapse in Pleistocene Australia and a Human Role in
Megafaunal Extinction. *Science* **309**, 287–290. (doi:10.1126/science.1111288)

29. Villavicencio NA, Lindsey EL, Martin FM, Borrero LA, Moreno PI, Marshall CR, Barnosky
AD. 2016 Combination of humans, climate, and vegetation change triggered Late
Quaternary megafauna extinction in the Última Esperanza region, southern Patagonia,
Chile. *Ecography* **39**, 125–140. (doi:10.1111/ecog.01606)

30. Faith JT. 2011 Late Pleistocene climate change, nutrient cycling, and the megafaunal
extinctions in North America. *Quaternary Science Reviews* **30**, 1675–1680.
(doi:10.1016/j.quascirev.2011.03.011)

31. Araújo T, Machado H, Mothé D, dos Santos Avila L. 2021 Species distribution modeling
reveals the ecological niche of extinct megafauna from South America. *Quat. res.* , 1–8.
(doi:10.1017/qua.2021.24)

32. Johnson C. 2006 *Australia's mammal extinctions: a 50,000-year history*. Cambridge
University Press.

33. Stuart AJ. 2015 Late Quaternary megafaunal extinctions on the continents: a short
review. *Geol. J.* **50**, 338–363. (doi:10.1002/gj.2633)

34. Johnson CN *et al.* 2016 What caused extinction of the Pleistocene megafauna of Sahul?
Proc. R. Soc. B **283**, 20152399. (doi:10.1098/rspb.2015.2399)

35. Peters KJ, Saltré F, Friedrich T, Jacobs Z, Wood R, McDowell M, Ulm S, Bradshaw CJA.
2019 FosSahul 2.0, an updated database for the Late Quaternary fossil records of Sahul.
Sci. Data **6**, 272. (doi:10.1038/s41597-019-0267-3)

36. Eklöf A, Tang S, Allesina S. 2013 Secondary extinctions in food webs: a Bayesian network
approach. *Methods Ecol. Evol.* **4**, 760–770. (doi:10.1111/2041-210X.12062)

37. Reed EH, Bourne SJ. 2009 Pleistocene fossil vertebrate sites of the south east region of
South Australia II. *T. Roy. Soc. South Aust.* **133**, 30–40.
(doi:10.1080/03721426.2009.10887108)

38. Macken AC, Reed EH. 2013 Late Quaternary small mammal faunas of the Naracoorte
Caves World Heritage Area. *T. Roy. Soc. South Aust.* **137**, 53–67.
(doi:10.1080/3721426.2013.10887171)

39. Rodríguez-Rey M *et al.* 2016 Data descriptor: a comprehensive database of quality-rated
fossil ages for Sahul's Quaternary vertebrates. *Sci. Data* **3**, 1–7.
(doi:10.1038/sdata.2016.53)

40. Rodríguez-Rey M *et al.* 2015 Criteria for assessing the quality of Middle Pleistocene to
Holocene vertebrate fossil ages. *Quat. Geochronol.* **30**, 69–79.
(doi:10.1016/j.quageo.2015.08.002)

41. Starrfelt J, Liow LH. 2016 How many dinosaur species were there? Fossil bias and true
richness estimated using a Poisson sampling model. *Phil. Trans. R. Soc. B* **371**, 20150219.
(doi:10.1098/rstb.2015.0219)

42. Macken AC, Reed EH. 2014 Postglacial reorganization of a small-mammal
paleocommunity in southern Australia reveals thresholds of change. *Ecol. Monogr.* **84**,
563–577. (doi:10.1890/13-0713.1)

43. Macken AC, Prideaux GJ, Reed EH. 2012 Variation and pattern in the responses of
mammal faunas to Late Pleistocene climatic change in southeastern South Australia. *J.*
Quat. Sci. **27**, 415–424. (doi:10.1002/jqs.1563)

44. Prideaux GJ, Roberts RG, Megirian D, Westaway KE, Hellstrom JC, Olley JM. 2007
Mammalian responses to Pleistocene climate change in southeastern Australia. *Geology*
35, 33–36. (doi:10.1130/G23070A.1)

45. Feldman A, Meiri S. 2013 Length-mass allometry in snakes: snake length-mass allometry.
Biol. J. Linnean. Soc. **108**, 161–172. (doi:10.1111/j.1095-8312.2012.02001.x)

46. Garnett ST *et al.* 2015 Biological, ecological, conservation and legal information for all
species and subspecies of Australian bird. *Sci. Data* **2**, 150061.
(doi:10.1038/sdata.2015.61)

47. Jones KE *et al.* 2009 PanTHERIA: a species-level database of life history, ecology, and
geography of extant and recently extinct mammals: *Ecological Archives* E090-184.
Ecology **90**, 2648–2648. (doi:10.1890/08-1494.1)

48. Meiri S. 2010 Length-weight allometries in lizards. *J. Zool.* (doi:10.1111/j.1469-
7998.2010.00696.x)

49. Oliveira BF, São-Pedro VA, Santos-Barrera G, Penone C, Costa GC. 2017 AmphiBIO, a
global database for amphibian ecological traits. *Sci. Data* **4**, 170123.
(doi:10.1038/sdata.2017.123)

50. Read QD, Baiser B, Grady JM, Zarnetske PL, Record S, Belmaker J. 2018 Tropical bird
species have less variable body sizes. *Biol. Lett.* **14**, 20170453.
(doi:10.1098/rsbl.2017.0453)

51. Santini L, Benítez-López A, Ficetola GF, Huijbregts MAJ. 2018 Length-mass allometries in
amphibians. *Integr. Zool.* **13**, 36–45. (doi:10.1111/1749-4877.12268)

52. Slavenko A, Tallowin OJS, Itescu Y, Raia P, Meiri S. 2016 Late Quaternary reptile
extinctions: size matters, insularity dominates. *Global Ecol. Biogeogr.* **25**, 1308–1320.
(doi:10.1111/geb.12491)

53. Wilman H, Belmaker J, Simpson J, de la Rosa C, Rivadeneira MM, Jetz W. 2014
EltonTraits 1.0: Species-level foraging attributes of the world's birds and mammals:
Ecological Archives E095-178. *Ecology* **95**, 2027–2027. (doi:10.1890/13-1917.1)

54. Cogger H. 2014 *Reptiles and amphibians of Australia*. CSIRO publishing.

55. Wilson SK, Swan G. 2008 *A complete guide to reptiles of Australia*. 2nd Ed. New Holland
Publishers.

56. Gravel D, Poisot T, Albouy C, Velez L, Mouillot D. 2013 Inferring food web structure from
predator-prey body size relationships. *Methods Ecol. Evol.* **4**, 1083–1090.
(doi:10.1111/2041-210X.12103)

57. Brose U *et al.* 2019 Predator traits determine food-web architecture across ecosystems.
Nat. Ecol. Evol. **3**, 919–927. (doi:10.1038/s41559-019-0899-x)

58. Brousseau P-M, Gravel D, Handa IT. 2018 Trait matching and phylogeny as predictors of
predator-prey interactions involving ground beetles. *Funct. Ecol.* **32**, 192–202.
(doi:10.1111/1365-2435.12943)

59. Eklof A, Helmus MR, Moore M, Allesina S. 2012 Relevance of evolutionary history for
food web structure. *Proc. R. Soc. B* **279**, 1588–1596. (doi:10.1098/rspb.2011.2149)

60. McLeod AM, Leroux SJ, Chu C. 2020 Effects of species traits, motif profiles, and
environment on spatial variation in multi-trophic antagonistic networks. *Ecosphere* **11**.
(doi:10.1002/ecs2.3018)

61. Morales-Castilla I, Matias MG, Gravel D, Araújo MB. 2015 Inferring biotic interactions
from proxies. *Trends Ecol. Evol.* **30**, 347–356. (doi:10.1016/j.tree.2015.03.014)

62. Pires MM, Koch PL, Fariña RA, de Aguiar MAM, dos Reis SF, Guimarães PR. 2015
Pleistocene megafaunal interaction networks became more vulnerable after human
arrival. *Proc. R. Soc. B* **282**, 20151367. (doi:10.1098/rspb.2015.1367)

63. Poelen JH, Simons JD, Mungall CJ. 2014 Global Biotic Interactions: an open infrastructure
to share and analyze species-interaction datasets. *Ecol. Inform.*
(doi:10.1016/j.ecoinf.2014.08.005)

64. Llewelyn J, Strona G, Bradshaw CJA. 2020 Inferring networks and modelling trophic
cascades. *GitHub and Zenodo* (doi:10.5281/zenodo.3756224)

65. de Visser SN, Freymann BP, Olff H. 2011 The Serengeti food web: empirical
quantification and analysis of topological changes under increasing human impact:
Topological changes under human impact. *J. Anim. Ecol.* **80**, 484–494.
(doi:10.1111/j.1365-2656.2010.01787.x)

66. Marples NM, Speed MP, Thomas RJ. 2018 An individual-based profitability spectrum for
understanding interactions between predators and their prey. *Biological Journal of the
Linnean Society* **125**, 1–13. (doi:10.1093/biolinnean/bly088)

67. Chapman AD. 2009 Numbers of living species in Australia and the world. , 1–78.

68. Forister ML *et al.* 2015 The global distribution of diet breadth in insect herbivores. *Proc.
Natl. Acad. Sci. USA* **112**, 442–447. (doi:10.1073/pnas.1423042112)

69. Fraser RA, Wells RT. 2006 Palaeontological excavation and taphonomic investigation of
the late Pleistocene fossil deposit in Grant Hall, Victoria Fossil Cave, Naracoorte, South
Australia. *Alcheringa* **30**, 147–161. (doi:10.1080/03115510609506860)

70. Reed L. 2012 Of mice and megafauna: new insights into Naracoorte's fossil deposits.
journal of the Australasian Cave and Karst Management Association **86**, 7–14.
(doi:hdl.handle.net/2440/93551)

71. Darimont CT, Fox CH, Bryan HM, Reimchen TE. 2015 The unique ecology of human
predators. *Science* **349**, 858–860. (doi:10.1126/science.aac4249)

72. Chichorro F, Juslén A, Cardoso P. 2019 A review of the relation between species traits
and extinction risk. *Biological Conservation* **237**, 220–229.
(doi:10.1016/j.biocon.2019.07.001)

73. Purvis A, Jones KE, Mace GM. 2000 Extinction. *BioEssays* **22**, 1123–1133.
(doi:10.1002/1521-1878(200012)22:12<1123::AID-BIES10>3.0.CO;2-C)

74. Kaneryd L *et al.* 2012 Species-rich ecosystems are vulnerable to cascading extinctions in
an increasingly variable world: Extinction Cascades in a Variable World. *Ecol. Evol.* **2**,
858–874. (doi:10.1002/ece3.218)

75. Schädler M, Brandl R, Kempel A. 2010 Host plant genotype determines bottom-up
effects in an aphid-parasitoid-predator system. *Entomol. Exp. Appl.* **135**, 162–169.
(doi:10.1111/j.1570-7458.2010.00976.x)

76. Turney CSM *et al.* 2001 Redating the onset of burning at Lynch's Crater (North
Queensland): implications for human settlement in Australia. *J. Quaternary Sci.* **16**, 767–
771. (doi:10.1002/jqs.643)

77. DeSantis LRG, Field JH, Wroe S, Dodson JR. 2017 Dietary responses of Sahul (Pleistocene
Australia–New Guinea) megafauna to climate and environmental change. *Paleobiology*
43, 181–195. (doi:10.1017/pab.2016.50)

78. Bampton T. 2021 Palaeoenvironmental proxies used to reconstruct the Quaternary of
Australia: a case study from Naracoorte Caves, South Australia. *Helicite* **46**, 1–13.

79. Flannery TF. 1990 Pleistocene faunal loss: implications of the aftershock for Australia's
past and future. *Archaeol. Oceania* **25**, 45–55. (doi:10.1002/j.1834-
4453.1990.tb00232.x)

80. Johnson CN. 2002 Determinants of loss of mammal species during the Late Quaternary
'megafauna' extinctions: life history and ecology, but not body size. *Proc. R. Soc. B* **269**,
2221–2227. (doi:10.1098/rspb.2002.2130)

81. Bradshaw CJ, Johnson CN, Llewelyn J, Weisbecker V, Strona G, Saltré F. 2021 Relative
demographic susceptibility does not explain the extinction chronology of Sahul's
megafauna. *eLife* **10**, e63870. (doi:10.7554/eLife.63870)

82. Roopnarine PD, Angielczyk KD, Wang SC, Hertog R. 2007 Trophic network models
explain instability of Early Triassic terrestrial communities. *Proc. R. Soc. B* **274**, 2077–
2086. (doi:10.1098/rspb.2007.0515)

83. Ceballos G, Ehrlich PR, Dirzo R. 2017 Biological annihilation via the ongoing sixth mass
extinction signaled by vertebrate population losses and declines. *Proc. Natl. Acad. Sci.*
USA **114**, E6089–E6096. (doi:10.1073/pnas.1704949114)

Figure 1

Figure 1. Sahul (top right insert) and the Naracoorte bioregion/Naracoorte Caves in south-eastern Sahul (main figure). These maps show coastline/sea level as they were during the Last Glacial Maximum (LGM; approximately 19,000 to 26,500 years ago). The grey area and thick outline indicate the land area during the LGM, the thinner lines show present-day coastlines and borders between countries and Australian states and territories, and the green area highlights the Naracoorte bioregion.

Figure 2

Figure 2. Example of an inferred ecological network model of the Late Pleistocene Naracoorte assemblage. Small points represent vertebrate species (nodes) and lines represent trophic interactions (links). Point colour shows trophic group (e.g., herbivores, carnivores etc), and extinct nodes are black. Due to uncertainty regarding trophic interactions, 1000 versions/models of the Naracoorte network were inferred and analysed. Plants, invertebrates and fish are shown as single large points in this figure.

Figure 3

Figure 3. The effects of basal connections, diet breadth, and trophic level on vulnerability to bottom-up cascades in synthetic networks. Vulnerabilities were calculated using either a simulation or Bayesian network approach. Plot **A** shows the weighted, model-averaged coefficients of the main effects (both methods), and plot **B** shows the marginal R^2 from mixed-effects models that had either basal connections, diet breadth, or trophic level as the only fixed effect (both methods). Error bars in plot A indicate 95% confidence intervals.

Figure 4

Figure 4. The effects of basal connections, diet breadth, and trophic level on vulnerability (calculated using the simulation approach) to bottom-up cascades in the Naracoorte network models. Each species' mean number of basal connections, diet breadth, trophic level, and vulnerability to bottom-up cascades was calculated, and linear regression models were fit to the mean data. Plot **A** shows the weighted, model-averaged coefficients of the main effects, and plot **B** shows the adjusted R^2 from linear regression models that had either basal connections, diet breadth, or trophic level as the only fixed effect. Error bars in plot A indicate 95% confidence intervals.

Figure 5. Comparison of species that survived into the Holocene (extant, shown in green) to those that went extinct in the Late Pleistocene (extinct, shown in red) from the Naracoorte network. Panel **A** shows the number of basal nodes (plants) connected directly and indirectly to each node via 'in' links; panel **B** shows diet breadth of each node; panel **C** indicates trophic level; and panel **D** shows the calculated coextinction vulnerability. The panels, which all include density violin plots and scatterplots, are based on each species' mean score across the 1000 Naracoorte network models.

Figure 6

Figure 6. Density plot showing results from a randomization test (20,000 replicates) comparing bottom-up coextinction vulnerability of extinct *versus* extant species from the Naracoorte network models. Each species' coextinction vulnerability was calculated using the simulation method (removing plant nodes and then removing animal nodes that no longer had connections to plant nodes). From these results, the coextinction vulnerabilities of 10 extinct and 10 extant species were sampled (with replacement), and the mean differences in coextinction between the groups calculated. This process was repeated 20 times for the results from each of the 1000 network models and used to build the density plot. The red area of the density plot indicates higher vulnerability for extinct species, and the green area indicates higher vulnerability for extant species.

Figure 7

Figure 7. Comparison of extinct *versus* extant species in terms of their position in the Naracoorte trophic network. Panel **A** is a biplot of the first two principal components from a principal component analysis of six variables describing species' positions in the network. Panel **B** is a violin and scatter plot showing the number of species that preyed on each extant *versus* extinct species. We inferred 1000 models of the Naracoorte network, and from these models calculated each species' mean value of each metric for use in the principal-components biplot and predator-diversity plot.