Sahul's megafauna were vulnerable to extinction due to their position in the trophic network

John Llewelyn[1] (johnsllewelyn@gmail.com), Giovanni Strona[2] (goblinshrimp@gmail.com), Matthew C. McDowell[3] (matthew.mcdowell@utas.edu.au), Christopher N. Johnson[4] (c.n.johnson@utas.edu.au), Katharina J. Peters[1] (katharina.peters@flinders.edu.au), Daniel B. Stouffer[5] (Daniel.stouffer@canterbury.ac.nz), Sara N. de Visser[6] (sndevisser@hotmail.com), Frédérik Saltré[1] (frederik.saltre@flinders.edu.au), Corey J. A. Bradshaw[1] (Corey.Bradshaw@flinders.edu.au)

[1]Global Ecology, College of Science and Engineering and ARC Centre of Excellence for Australian Biodiversity and Heritage, Flinders University, GPO Box 2100, Adelaide, South Australia 5001, Australia
[2]Research Centre for Ecological Change, University of Helsinki, Viikinkaari 1, Biocentre 3, 00790, Helsinki, Finland
[3]Dynamics of Eco-Evolutionary Pattern and ARC Centre of Excellence for Australian Biodiversity and Heritage, University of Tasmania, Tasmania 7001, Australia
[4]School of Natural Sciences and Australian Research Council Centre of Excellence for Australian Biodiversity and Heritage, University of Tasmania, Private Bag 55, Hobart, Tasmania 7001, Australia
[5]Centre for Integrative Ecology, School of Biological Sciences, University of Canterbury, Christchurch, New Zealand
[6]Community and Conservation Ecology, Centre for Ecological and Evolutionary Studies, University of Groningen, PO Box 11103, 9700 Groningen, The Netherlands

Short running title: Trophic cascades and megafauna

Key words: ecological network, coextinction, biotic interactions, extinction event, Late Pleistocene, food web

Author contributions
Project conceptualization by J.L., G.S., C.J.A.B., and F.S. Data curation by J.L., G.S., K.J.P., C.N.J., M.C.M., and S.N.d-V. Investigation and formal analysis by J.L., G.S., and C.J.A.B. Methodology by J.L., G.S., C.J.A.B., and D.B.S. Original draft preparation by J.L., G.S., C.J.A.B., and F.S. All authors contributed to reviewing and editing the draft.

Data and code accessibility
All relevant data and code are available at Github.com/JohnLlewelyn/Inferring-networks-and-modelling-trophic-cascades (DOI: 10.5281/zenodo.3756224)

Corresponding author
John Llewelyn
Email: johnsllewelyn@gmail.com
Phone: +61 429 506 956
Fax: +8 8201 3399
Address: Global Ecology Lab, College of Science & Engineering, Flinders University

Sahul's megafauna were vulnerable to extinction due to their

position in the trophic network

Abstract

Extinctions stemming from environmental change often trigger trophic cascades and

coextinctions. However, it remains unclear whether trophic cascades were a large

contributor to the megafauna extinctions that swept across several continents in the Late

Pleistocene. The pathways to megafauna extinctions are particularly unclear for Sahul

(landmass comprising Australia and New Guinea), where extinctions happened earlier than

on other continents. We investigated the role of bottom-up trophic cascades in Late

Pleistocene Sahul by constructing pre-extinction (~ 80 ka) trophic network models of the

vertebrate community of Naracoorte, south-eastern Australia. These models allowed us to

predict vertebrate species' vulnerability to cascading extinctions based on their position in

the network. We tested whether the observed extinctions could be explained by bottom-up

cascades, or if they should be attributed to other external causes. Species that disappeared

from the community were more vulnerable, overall, to bottom-up cascades than were

species that survived. The position of extinct species in the network – having few or no

predators – also suggests they might have been particularly vulnerable to a new predator.

These results provide quantitative evidence that trophic cascades and naivety to predators

could have contributed to the megafauna extinction event in Sahul.

Introduction

Megafauna (animals > 44 kg) extinctions swept across several continents during the Late Pleistocene (126,000 – 12,000 years ago), with the highest proportions of genera lost from Sahul (the landmass including Australia and New Guinea) and the Americas [1,2]. While the causes of these extinctions are still debated, most evidence points toward the arrival of anatomically modern humans and/or climate change [3–5]. Irrespective of the root causes, large extinction events always involve both primary and secondary (or co-) extinctions [6]. That is, some extinctions result from the direct impact of environmental change or over exploitation (i.e., primary extinctions), and then coextinctions occur in species that depend directly or indirectly on species lost to primary extinction [7–9].

Of all the extinctions that have ever occurred on Earth, many — potentially the majority — have been coextinctions [10]. In some cases, these coextinctions involved host-specific parasites that were doomed by the extinction of their host species, or flowering plants imperilled by the extinction of their pollinators [11]. Coextinctions have also been mediated through trophic interactions between herbivores and vegetation, and between predators and their prey [6,12]. Indeed, coextinctions via trophic cascades have previously been implicated in Late Pleistocene megafauna extinctions [13,14].

Bottom-up trophic cascades, stemming from changes in the plant/primary producer community, can profoundly alter ecological communities [15]. A recent study found that the trophic group most strongly affected by changes in the plant community are herbivores [16]. Extinction risk ratings from the International Union for Conservation of Nature (IUCN) and recent extinctions suggest that herbivorous terrestrial vertebrates are indeed

particularly vulnerable to extinction [17], a pattern that might in part be explained by the

sensitivity of herbivores to bottom-up cascades. However, it remains unclear whether

bottom-up cascades generally impact lower trophic levels more strongly [18], and whether

bottom-up cascades and coextinctions played a role in the megafauna extinction events of

the Late Pleistocene.

Identifying the vulnerability of species to bottom-up (or top-down) coextinction cascades

relies heavily on understanding species interactions within an ecological community. To this

end, ecological network modelling is an invaluable tool for representing ecological

communities from a species-interaction perspective and studying the consequences of

changes in these interactions [19]. In ecological network models, organismal groups (e.g.,

species, age groups, populations, or individuals) are represented by nodes, and interactions

– which can be weighted or un-weighted – are represented by links. The interaction type

most frequently used to build ecological network models are trophic interactions, creating

networks analogous to food webs. For contemporary communities, there is a growing

number of studies that use detailed information on species interactions to build network

models and study trophic cascades [20–22]. Unfortunately, similar approaches are

challenging to apply to palaeo communities because of the lack of data on ancient trophic

interactions. However, by combining contemporary and palaeo data to infer trophic

interactions, this limitation can be overcome and network models of palaeo communities

can be constructed [23,24]. For example, Pires et al. (2015) used this approach to model

Late Pleistocene mammal communities in the Americas, concluding that (1) pre-existing

American mammal networks were not especially unstable (i.e., they were similar to modern

networks in Africa in terms of population densities re-establishing after simulated

perturbations), and (2) the arrival of humans destabilized the networks because this new

predator increased network connectance (i.e., the proportion of potential links that are

realized). However, the vulnerability of extinct *versus* extant species to coextinction was not

tested, nor whether their conclusions would hold true in other continents presenting vastly

different species assemblages.

The Late Pleistocene megafauna assemblage of Sahul was distinct from that of other

continents in that all the large mammals were marsupials or monotremes [25]. Giant

reptiles and birds were also a prominent component of this continent's megafauna [26].

While Sahul's megafauna included many species over the standard body-mass threshold for

megafauna of 44 kg, the term 'megafauna' is often extended in Sahul to include species with

a body mass above that of their surviving relatives [25] — a definition we have adopted

here. Identifying the pathways by which Sahul's unique megafauna were lost is challenging

because the Late Pleistocene megafauna extinctions happened much earlier in Sahul than

elsewhere [27]. To understand such ancient extinction events, a sufficient number of dated

fossilized remains is necessary [28]. The most detailed and well-studied fossil record

spanning the megafauna extinction event in Sahul comes from the Naracoorte region in

south-eastern Australia. This fossil record offers an exceptional picture of the species living

in the region over the past 500,000 years, including the ecological community at the time of

the main megafauna extinction event that occurred approximately 44,000 years ago in

Naracoorte [5]. Thus, the Naracoorte fossil record is the best platform available from which

to model the ecological and environmental processes potentially involved in the megafauna

extinction event in Sahul.

Using Naracoorte as a model system to assess whether bottom-up coextinction cascades

were an important component of the overall extinction event in Late Pleistocene Sahul, we

constructed an entire terrestrial, palaeo-vertebrate assemblage (including all terrestrial

vertebrate classes), and combined this assemblage with palaeo and contemporary data to

infer trophic interactions and build network models (Figure 1). These network models

consisted of nodes (species) with directed, unweighted links. We then computed each

species' vulnerability to coextinction via bottom-up cascades, by iteratively taking the

product of the vulnerabilities of the focal species' resource nodes using an algorithm similar

to PageRank. The resultant vulnerabilities were assessed in three steps: (1) we tested for a

relationship between trophic level and vulnerability to coextinction, (2) we compared

trophic composition of the extant (defined as species that survived the Late Pleistocene

extinction event) and extinct species groups to see whether vulnerable trophic levels were

over-represented in the extinct group; and (3) we focused on carnivores (the group we have

the most information on regarding trophic interactions) to test whether extinct carnivores

were more vulnerable to bottom-up cascades than were extant carnivores. In addition to

bottom-up coextinction cascade vulnerability, we also tested for differences in the network

positions of extant and extinct species to see if relative position could have made extinct

species more vulnerable in other ways (e.g., more vulnerable to a new predator).

By incorporating interactions involving all vertebrate species, we adopted a holistic

approach to studying megafauna extinctions of the Late Pleistocene. Overall, we found that

extinct species were more vulnerable to coextinction than were extant species, and

conclude that bottom-up trophic cascades possibly contributed to this megafauna extinction

event. Our results also indicate that extinct species had fewer predators than did surviving

species, suggesting extinct species might have been particularly sensitive to the arrival of

the new predator (humans). By using network modelling, our study therefore demonstrates

that Sahul's megafauna were vulnerable to extinction due to their relative positions in their

ecological community.

Methods

Our aim was to develop and interrogate ecological network models representing the

Naracoorte ecosystem prior to the main megafauna-extinction pulse and the arrival of

humans in the region (~ 44,00 years ago) [5]. Below, we describe how we built ecologically

realistic models of the Naracoorte network and simulated bottom-up

extinction/coextinction cascades (see Supplementary Figure S1 for a simplified flowchart of

our methods).

Study region

The World Heritage-listed (from 1994) Naracoorte Caves in south-eastern South Australia

(37° 02' 24" S, 140° 48' 00" E) encompass a series of limestone caves that opened and closed

to the surface at different locations, and at different times, over the last 500,000 years [29].

These openings acted as natural pitfall traps, capturing snapshots of Naracoorte's

biodiversity at different periods from at least 500,000 years ago to the present. In addition

to the fossils of the many animals that fell into these natural traps, there are remains of

species that lived in the caves, such as owls and bats, and their prey [30]. Consequently, the

Naracoorte Caves provide an ideal platform from which to build palaeo-ecological network

models to gain insight into how these long-lost ecosystems functioned and changed over

time.

Data

To build a species assemblage list (i.e., to identify the nodes to include in the network

models), we used two data sources: *FosSahul* 2.0 [28,31] and the Atlas of Living Australia.

FosSahul 2.0 is a database of dated fossil records from Sahul, including an automated

quality-rating method for date reliability [32]. We extracted and vetted records from the

Naracoorte region (defined as the region between 35° 32′ 48″ S and 38° 6′ 50″ S, and

between 139° 10′ 42″ E and 141° 0′ 21″ E) from *FosSahul* 2.0 that had high- or intermediate-

quality dates (A*, A or B) [28], and whose age was younger than 200,000 years before

present. We chose this cut-off age to provide a large enough time period to include dated

fossils from all/most megafauna species that lived in the region immediately prior to the

main extinction event; if we had made the time period too narrow, few megafauna species

would have been captured despite their likely presence in the region at the time. Fossil

records suffer from taphonomic biases (biases in the accumulation and preservation of

different organisms), and consequently some species that were present in Late Pleistocene

Naracoorte are unlikely to be represented in the fossil record. Furthermore, there are biases

for studying and dating particular groups of species due to academic and amateur interests

[33]. These biases, along with the fact that *FosSahul* was primarily designed to document

megafauna remains rather than smaller species, means that *FosSahul* 2.0 does not include

all vertebrate species present in Late Pleistocene Naracoorte. To account for this gap in the

species list, we supplemented *FosSahul* 2.0 data with contemporary and historical species

records from the Naracoorte Coastal Plains bioregion from the Atlas of Living Australia

online repository (ala.org.au; accessed 3 January 2019). The Atlas of Living Australia has

detailed species records of vertebrates from this bioregion, and so it captures most of the

diversity of extant and recently extinct vertebrate species. We extracted data pertaining to

all terrestrial vertebrates from the region, and removed species that were introduced since

European arrival, as well as vagrants and erroneous records (species well-outside their

normal distribution and not present in the fossil record), and strictly coastal species such as

marine birds that do not use inland waterways. Our final species list, built using *FosSahul* 2.0

and the Atlas of Living Australia, included 423 vertebrate species (Supplementary Table S1).

To infer trophic links, we required information on each species' body mass and broad

trophic category (whether it ate plants and/or fungi, terrestrial invertebrates, terrestrial

vertebrates, or aquatic animals). For extant species, we extracted much of this information

from large databases: snake database [34]; Australian bird database [35]; PanTHERIA

(mammal database) [36]; lizard database [37]; AmphiBIO (amphibian database) [38];

tropical bird database [39]; Amphibian database [40]; reptile database [41]; and Elton traits

databases (mammals and birds) [42]. However, information on body mass was not available

for some extant reptiles and amphibians. For these species, we estimated body mass using

their body length (from field guides) and validated allometric relationships [34,37,40,43,44].

Extinct megafaunal species were not included in any of the aforementioned databases, so

for these species we obtained data on body mass and diet from the literature (see

Supplementary Table S2). The 423 vertebrate species in the Naracoorte assemblage

included 109 that consumed vertebrates, 221 that consumed plants, 318 that consumed

invertebrates, and 75 that consumed aquatic animals (Supplementary Table S1). Of the 423

species, 238 consumed more than one trophic group. Mean body mass ranged from 0.4 g to

2700 kg (Supplementary Table S1).

Inferring trophic links

For almost all extant species, information on trophic interactions is incomplete, but the lack

of knowledge regarding trophic interactions is even greater for long-extinct species. To

overcome this limitation and build realistic ecological network models, various methods

have been developed for inferring trophic links using species' traits such as body size (i.e.,

larger predators tend to eat larger prey, and predators are usually bigger than their prey

[45]). Others have built on this approach to improve prediction accuracy [46–49] by adding

phylogenetic and physiological information, prohibiting impossible or unlikely links,

incorporating specific morphological features such as biting force and cuticular thickness,

and taking into account abundance. However, most of this research focusses on fish or

invertebrates, with the effectiveness of these methods rarely applied or validated for

terrestrial vertebrates (but see [50,51]).

We therefore developed and validated a new method, based on the body-size trophic niche

model [45], to infer trophic links between terrestrial vertebrates. The body-size trophic

niche model [45], that we adapted, consists of two quantile regressions: one defines the

upper prey-size limit given predator mass, and the other defines the lower prey-size limit

given predator mass. If a species falls within the upper and lower limits for a particular

predator, it is inferred as potential prey for that predator. We used a large predator-prey

interaction dataset to identify these body-size relationships between terrestrial predators

and their prey, and tested whether these relationships varied depending on taxonomic class

of the predator (i.e., did including predator class as an independent variable improve the fit

of the body size quantile regressions?).

We extracted the interaction dataset from GloBI, an online repository of biotic interactions

[52]. The dataset consisted of 3893 records: 958 records of predation by non-marine

mammals, 2711 by birds, 199 by reptiles, and 25 by amphibians (Supplementary Table S3)

[53]. We extracted data on mean body mass for the species from the same databases we

used to add this information to the Naracoorte species list (see above). Once we identified

the best trophic niche model (i.e., the combination of quantile regressions that best fit the

upper and lower prey-size limits) using the entire GloBI data set and Bayesian information

criterion (BIC; Supplementary Methods S1), we validated this method of assigning trophic

links by comparing model performance when applied first to the GloBI data divided into

training and validation datasets, and then to a well-resolved trophic network from the

Serengeti [54] (plus S. de Visser unpublished data; Supplementary Methods S1). We used

the true skill statistic to evaluate model performance and found that the top-ranked model

(according to BIC) also performed best at assigning links in the validation step

(Supplementary Methods S1, Supplementary Table S4, Supplementary Figure S2, script

available at [53]). We used the Serengeti network to quantify how much our best trophic

niche model overestimated the number of interactions, so we could adjust for this

overestimation when building networks (see methods below, Supplementary Methods S1,

script available at [53]).

While the inclusion of additional species traits could improve accuracy of predator-prey

interaction inferences, we used only three readily available traits (broad diet, body size, and

predator taxonomic class with reptiles and amphibians grouped together). We restricted our

method to using these three traits because: (1) using few traits is compatible with the

quantile regression framework, (2) they can easily be extracted for vertebrate species in

most assemblages (and therefore the method can be widely applied), and (3) our validation

steps demonstrated the resulting trophic niche model's performance is sufficient (true skill

statistic = 0.6 when applied to the Serengeti assemblage).

Networks

To build realistic ecological networks for the Naracoorte assemblage, we applied the trophic

niche model to the species list, removed excess links between vertebrates (to account for

overestimating the number of predator-prey links), and added links to non-terrestrial

vertebrate food resources. However, there is uncertainty regarding which vertebrate

predator/prey links to delete as well as how many links to add from non-terrestrial

vertebrate food resources to vertebrates. To address this uncertainty, we used a

randomization approach in the link-removal and -addition steps described below, and

generated 1000 versions of the network. That is, we randomly removed (for the vertebrate

predator-prey links) or added (for the herbivores, invertivores, and predators of aquatic

animals) links in the range indicated as realistic based on contemporary ecological networks

and species' diet breadths. Because we do not know exactly where to add or remove these

links, we generated 1000 versions of the network so our results were not skewed by the

particular links selected.

We used our trophic niche model to assign potential predator-prey links in the Naracoorte

species assemblage. Then we assigned a probability to each potential predator-prey

interaction based on the prey's position in the predator's prey-size range and a Gaussian

distribution centred on this range (with a standard deviation equal to a quarter of the

predator's prey-size range); the highest probability was for prey close to the centre of the

prey-size range (i.e., centre of the distribution) and decreased the closer the prey was to the

predator's limits. For each predator, we randomly sampled from the Serengeti

overprediction density kernel (Supplementary Methods S1) to determine the proportion of

links to keep, and then randomly selected from the potential links taking into account the

probability of the predator-prey interaction. This procedure resulted in a vertebrate

network with realistic connectance (proportion of potential links that are realized), and with

most of each predator's prey closer to the centre, rather than the limits, of their prey-size

range.

Terrestrial vertebrates not only consume other terrestrial vertebrates, they also consume

invertebrates, plants and fungi, and aquatic animals. In addition to inferring trophic links

among terrestrial vertebrates, we therefore needed to add links to vertebrates from these

other food resources. However, because our study focusses on vertebrate species, and

because invertebrate, plant, and fungi diversities are not well resolved for most ecosystems

(including Late Pleistocene Naracoorte), we did not include detail on individual species

within these groups. We therefore generated a pool of n species for each group (n = 1440

for plants, 5620 for invertebrates, and 104 for aquatic animals), with these pool numbers

calculated based on the described diversity in these groups relative to terrestrial vertebrate

diversity in Australia [55,56]. To assign links from vertebrates to species from these groups,

we used published records of vertebrate diet breadth (number of species consumed) for 34

herbivores, 15 invertivores, and 6 predators of aquatic species (Supplementary Table 5). To

these data, we fit kernel density estimates from which we randomly sampled (within a diet

breadth range of 1 to twice the maximum diet breadth recorded for that trophic guild in

Supplementary Table S5) to assign the number of links between each resource group and

vertebrate consumer in each of the 1000 network models.

We completed the networks by generating links from plants to invertebrates, and from

plants to aquatic animals. For these subnetworks, we randomly selected a value of

connectance between 0 and 0.1 for each of the 1000 replicated models. That value defined

the density of plant-invertebrate and plant-aquatic animal links (we did not generate

invertebrate-invertebrate links). Each plant, invertebrate, and aquatic animal node was

randomly assigned a probability from a log-Normal distribution with a mean = 0. We then

allocated links between nodes by sampling them according to these probabilities. In each

simulation, we attributed the plant, invertebrate, and aquatic animal groups a σ value

defining the shape of the log-Normal distribution, randomly sampled between 0.5 and 1.5

(which varied diet overlap and competition for resources in these subnetworks). We

allocated plant-invertebrate and plant-aquatic animal links until we reached the

predetermined connectance.

For vertebrates that fed from more than one group (i.e., omnivores, which represent over

half the vertebrates in this network), we made a proportional adjustment to the number of

'in' links (food resources) depending on how many food groups they fed from. For example,

if a species consumed from two groups, such as vertebrates and invertebrates, we randomly

deleted half of the links from each group; if they fed from three groups, we randomly

removed two-thirds of the links from each food group (and so on). We made these deletions

to prevent omnivores from having inflated diet breadths.

In some cases involving water birds, we used empirical evidence to avoid assigning

unrealistic trophic links. For example, pelicans are large and, consequently, the trophic niche

model predicts that pelicans take large vertebrate prey. However, we know that pelicans are

not birds of prey and do not consume large terrestrial animals. Thus, we restricted the

allocation of trophic links for such birds to prevent them from feeding on unrealistically

large terrestrial vertebrates (they were prevented from consuming prey that weighed over a

third of their body mass).

Vulnerability to coextinction

We computed a measure of relative risk to bottom-up coextinction cascades for each node

in the network based on the node's probability of losing a certain percentage (t%) of its

most vulnerable resources. The resource loss threshold, t%, was assigned for each of the

1000 replicated network models by randomly sampling between 0 and 1. Since the problem

of calculating a node's probability of losing a percentage of its resources is iterative (with

the extinction probability of one node depending on the unknown probability of extinction

of other nodes, and so on), we used an algorithmic approach similar to the popular

PageRank [57]. For this, we first attributed an arbitrary extinction risk of 0.9 to all nodes in

the network, and then reiterated the procedure of computing extinction risk for each node

in the network as the product of the extinction risk of the t% most vulnerable nodes having

outgoing links pointing towards the focal node [53]. We continued the reiteration process

until the risk for all nodes stopped changing.

Analysis

We compared susceptibility of extinct *versus* extant species to bottom-up coextinction

cascades in two ways: (1) by assessing the vulnerability of different trophic levels and

comparing their prevalence in the extinct *versus* extant group, and (2) by assessing the

vulnerability of extinct *versus* extant predators (at the species level). We used a trophic-

rather than species-level comparison for the entire assemblage because, for most trophic

groups (herbivores, invertivores, and aquatic feeders), we allocated resource-consumer

links by sampling from diet-breadth density kernels without using species traits to

differentiate species within the trophic group. For the trophic-level comparison, we first

assessed whether vulnerability to bottom-up coextinction cascades (according to the

vulnerability algorithm) differed between trophic levels, and then compared the proportion

of each trophic level in the extinct *versus* extant groups to see whether vulnerable trophic

levels were over-represented in either group. We applied a randomization test to compare

the proportion of each trophic group among extinct and extant species by sampling (with

replacement) 10 extant and 10 extinct species from the Naracoorte assemblage, and

subtracting the number of species from the trophic level of interest in the extinct sample

from that number in the extant sample. We repeated this process 10,000 times and used

the ensuing results to make density plots to assess whether any of the trophic groups were

over-represented among extinct or extant species.

In contrast to herbivores, invertivores, and aquatic feeders, resource links differed among

predators depending on the species' traits. Therefore, in addition to the trophic-level

comparison, we compared individual predators to see if extinct species were at a higher risk of

coextinction from bottom-up cascades than were extant predators. To test this, we compared

the weight of evidence (according to AICc) for two generalized mixed-effects models fit to the

predator coextinction vulnerability data. In both models, vulnerability was the dependent

variable and species was a random effect. We compared one model (random-intercept) with no

fixed effects to another model where the species' extinction status was the only fixed effect. If

the latter model had a higher weight of evidence, it is implied that extinct and extant predators

differed in their vulnerability.

We assessed the position of extinct *versus* extant species in the network using 12 different

network metrics: *trophic level, pageRank, betweenness centrality, eigenvector centrality,*

closeness centrality (in), coreness (in), degree (in), eccentricity (in), closeness centrality (out),

coreness (out), degree (out), and *eccentricity (out)* (Supplementary Table S6 for metric

descriptions). Metrics followed by an 'in' or 'out' were calculated for each node using links

pointing towards (in) or away from (out) the focal node. We chose these metrics because

they are commonly used, node-level metrics describing position in the network. For each

metric, we calculated the species' mean value across the 1000 network models. After

checking for highly correlated metrics and removing those identified as redundant (i.e.,

metrics that had a Spearman's ρ > 0.8), we ran a principal component analysis and visually

inspected for grouping of extinct and extant species. The reduced list of metrics included

closeness (out), *eccentricity* (out), *degree* (in and out), and *PageRank*.

Results

Our coextinction-vulnerability measure found that herbivores were more vulnerable to

bottom-up cascades than were species from mid- or high-trophic levels (Figure 2a).

Herbivores were over-represented among extinct species (67% of extinct species vs. 11% of

extant species; Figure 2b), whereas the less-vulnerable species occupying mid-trophic levels

were over-represented among extant species (11% of extinct species vs. 63% of extant

species; Figure 2c). Carnivores made up a similar proportion of species in the extinct and

extant species groups (22% of extinct species vs. 25% of extant species; Figure 2d).

For predatory species (the trophic group for which a species' in links differed depending on

the species' traits), extinct predators were more vulnerable to bottom-up cascades than

were extant predators (mean coextinction vulnerability of 0.15 and 0.09 for extinct and

extant predators, respectively; Figure 3). The predator vulnerability mixed-effects model

with extinction status as a fixed effect had an AICc weight (wAICc) of 0.986, meaning that

this model was supported 68.9 times more than the intercept-only model (information-

theoretic evidence ratio = 0.986/0.014 = 68.9).

Extinct species differed from extant species in terms of their position in the network.

Principal component analysis of six network-position metrics shows that extinct and extant

species fall into two distinct groups according to the first principal component (dimension 1

in Figure 4a). The main contributors to this principal component are metrics describing a

node's connection to the network through its 'out' links (Figure 4a; Supplementary Figure

S3). Closer examination of out links (i.e., number of predators) shows that, on average,

extinct species had < 1 predator, whereas extant species had > 8 (Figure 4b). Indeed, the

average number of predators was lower for extinct species than for extant species in all

1000 models of the Naracoorte network. Extinct species also had fewer in links (number of

trophic resources), on average, than did extant species (Figure 4c).

Discussion

Our analyses demonstrate the position of extinct species in the Late Pleistocene Naracoorte

network made them more vulnerable to bottom-up coextinction cascades than co-occurring

extant species, suggesting that trophic cascades might have contributed to the megafauna

extinction event in south-eastern Sahul. However, extinct species also had fewer predators

than did extant species, a network position attribute that would likely have made them

more vulnerable to the arrival of the new 'super predator' – humans [58].

Herbivores, the trophic group most vulnerable to bottom-up cascades in our analyses (as

well as in field experiments involving plant and invertebrate communities) [16], were over-

represented among the extinct species from Naracoorte (Figure 2b). Furthermore,

Naracoorte's extinct predators were more vulnerable to bottom-up cascades than were

predatory species that survived into the Holocene (Figure 3). Together, these results suggest

the species lost from Naracoorte in the Late Pleistocene tended to be those more

vulnerable to bottom-up trophic cascades. This leads to the question: did vegetation change

and initiate trophic cascades in Naracoorte at the same time as the megafauna

disappeared? Studies in other regions of Sahul have detected shifts in vegetation roughly

coinciding with or preceding megafauna extinction. Hypothesised drivers of these shifts

include land-use changes associated with human arrival (i.e., fire-stick farming) [59,60]

and/or climate change [61]. However, there are no studies on the vegetation of Naracoorte

spanning the Late Pleistocene extinction event. Nonetheless, the megafauna's

disappearance from Naracoorte coincided with an unusually cool period (Supplementary

Figure S3) and the arrival of humans (~ 44,000 years ago) [5], offering the intriguing

possibility that changes in climate or land use triggered shifts in vegetation that had

consequences for higher trophic levels in the network. To evaluate this possibility, more

studies are required to model vegetation changes in south-eastern Sahul (including the

Naracoorte region), and these must be validated using the pollen record and/or other fossil

data.

By considering the network position of all vertebrate species in the assemblage, a clear

difference between extinct and extant species emerges: extinct species had far fewer

predators than did species that survived (mean number of predators: 0.7 *versus* 8.1 for

extinct *versus* extant species, respectively; Figure 4b). This predator naivety, coupled with

the species' slow life histories, likely made megafauna especially vulnerable to new

predators [27,62,63] and suggests hunting by humans could have adversely affected

megafauna. Megafauna also had fewer trophic resources, on average, than did surviving

species (mean number of food species: 38.4 *versus* 72.7 for extinct *versus* extant species,

respectively; Figure 4c), but this difference reflected the over-representation of herbivores

(which have narrower diet breadth than other trophic groups in our models; Supplementary

Table S5) among the extinct species. As palaeo-vegetation, invertebrate, trophic and climate

data improve, network models can eventually incorporate this information to build more

refined network models, and include more detailed top-down and bottom-up forcings to

assess the plausibility of the different potential causes and pathways to extinction —

including what (if any) role humans played in the megafauna's demise.

In summary, our network modelling of Late Pleistocene Naracoorte suggests bottom-up

coextinction cascades and/or predator naivety and the arrival of humans could have driven

the megafauna extinction event in Sahul. Indeed, that our network modelling showed that

extinct species were vulnerable to bottom-up cascades *and* new predation pressures lends

support to recent research suggesting that climate change (that can shift vegetation and

lead to bottom-up cascades) and human arrival drove the megafauna extinction trajectories

in south-eastern Sahul [5]. Our research, along with other recent studies [23,51,65],

demonstrates that network modelling is a powerful tool for investigating and understanding

ancient extinction events. By developing methods to model whole-community responses to

change and validating these methods using ancient extinction events, we can also provide

better estimates of the fates of contemporary communities as the sixth mass-extinction

event unfolds [66].

Acknowledgements

We thank E. Reed, C. Carbone, M. Tucker, C. Dickman and V.K. Llewelyn for their

constructive input.

Funding

This work was supported by the Australian Research Council Centre of Excellence for

Australian Biodiversity and Heritage (grant number CE170100015).

References

1.   Barnosky AD. 2004 Assessing the causes of Late Pleistocene extinctions on the
continents. *Science* **306**, 70–75. (doi:10.1126/science.1101476)

2.   Koch PL, Barnosky AD. 2006 Late Quaternary extinctions: state of the debate. *Annu. Rev.*
Ecol. Evol. Syst. **37**, 215–250. (doi:10.1146/annurev.ecolsys.34.011802.132415)

3.   Lorenzen ED *et al.* 2011 Species-specific responses of Late Quaternary megafauna to
climate and humans. *Nature* **479**, 359–364. (doi:10.1038/nature10574)

4. Metcalf JL *et al.* 2016 Synergistic roles of climate warming and human occupation in
Patagonian megafaunal extinctions during the Last Deglaciation. *Sci. Adv.* **2**, e1501682.
(doi:10.1126/sciadv.1501682)

5. Saltré F, Chadoeuf J, Peters KJ, McDowell MC, Friedrich T, Timmermann A, Ulm S,
Bradshaw CJA. 2019 Climate-human interaction associated with southeast Australian
megafauna extinction patterns. *Nat. Commun.* **10**, 5311. (doi:10.1038/s41467-019-
13277-0)

6. Colwell RK, Dunn RR, Harris NC. 2012 Coextinction and persistence of dependent species
in a changing world. *Annu. Rev. Ecol. Evol. Syst.* **43**, 183–203. (doi:10.1146/annurev-
ecolsys-110411-160304)

7. Dunne JA, Williams RJ, Martinez ND. 2002 Network structure and biodiversity loss in
food webs: robustness increases with connectance. *Ecol. Lett.* **5**, 558–567.
(doi:10.1046/j.1461-0248.2002.00354.x)

8. Pimm SL. 1980 Food web design and the effect of species deletion. *Oikos* **35**, 139.
(doi:10.2307/3544422)

9. Strona G, Bradshaw CJA. 2018 Co-extinctions annihilate planetary life during extreme
environmental change. *Sci Rep* **8**, 16724. (doi:10.1038/s41598-018-35068-1)

10. Dunn RR, Harris NC, Colwell RK, Koh LP, Sodhi NS. 2009 The sixth mass coextinction: are
most endangered species parasites and mutualists? *Proc. R. Soc. B* **276**, 3037–3045.
(doi:10.1098/rspb.2009.0413)

11. Koh LP, Dunn RR, Sodhi NS, Colwell RK, Proctor HC, Smith VS. 2004 Species coextinctions
and the biodiversity crisis. *Science* **305**, 4.

12. Estes JA *et al.* 2011 Trophic downgrading of Planet Earth. *Science* **333**, 301–306.
(doi:10.1126/science.1205106)

13. Galetti M *et al.* 2018 Ecological and evolutionary legacy of megafauna extinctions:
anachronisms and megafauna interactions. *Biol. Rev.* **93**, 845–862.
(doi:10.1111/brv.12374)

14. Ripple WJ, Van Valkenburgh B. 2010 Linking top-down forces to the Pleistocene
megafaunal extinctions. *BioScience* **60**, 516–526. (doi:10.1525/bio.2010.60.7.7)

15. Aberhan M, Kiessling W. 2015 Persistent ecological shifts in marine molluscan
assemblages across the end-Cretaceous mass extinction. *Proc Natl Acad Sci USA* **112**,
7207–7212. (doi:10.1073/pnas.1422248112)

16. Scherber C *et al.* 2010 Bottom-up effects of plant diversity on multitrophic interactions
in a biodiversity experiment. *Nature* **468**, 553–556. (doi:10.1038/nature09492)

17. Atwood TB, Valentine SA, Hammill E, McCauley DJ, Madin EMP, Beard KH, Pearse WD.
2020 Herbivores at the highest risk of extinction among mammals, birds, and reptiles.
Sci. Adv. **6**, eabb8458. (doi:10.1126/sciadv.abb8458)

18. Haddad NM, Crutsinger GM, Gross K, Haarstad J, Knops JMH, Tilman D. 2009 Plant
species loss decreases arthropod diversity and shifts trophic structure. *Ecology Letters*
12, 1029–1039. (doi:10.1111/j.1461-0248.2009.01356.x)

19. Säterberg T, Sellman S, Ebenman B. 2013 High frequency of functional extinctions in
ecological networks. *Nature* **499**, 468–470. (doi:10.1038/nature12277)

20. Boit A, Martinez ND, Williams RJ, Gaedke U. 2012 Mechanistic theory and modelling of
complex food-web dynamics in Lake Constance. *Ecol. Lett.* **15**. (doi:10.1111/j.1461-
0248.2012.01777.x)

21. Coll M, Lotze HK, Romanuk TN. 2008 Structural degradation in Mediterranean sea food
webs: testing ecological hypotheses using stochastic and mass-balance modelling.
Ecosystems **11**, 939–960. (doi:10.1007/s10021-008-9171-y)

22. Pocock MJO, Evans DM, Memmott J. 2012 The robustness and restoration of a network
of ecological networks. *Science* **335**, 973–977. (doi:10.1126/science.1214915)

23. Dunne JA, Labandeira CC, Williams RJ. 2014 Highly resolved early Eocene food webs
show development of modern trophic structure after the end-Cretaceous extinction.
Proc. R. Soc. B **281**, 20133280–20133280. (doi:10.1098/rspb.2013.3280)

24. Roopnarine PD. 2006 Extinction cascades and catastrophe in ancient food webs.
Paleobiology **32**, 1–19. (doi:10.1666/05008.1)

25. Johnson C. 2006 *Australia's mammal extinctions: a 50,000-year history*. Cambridge
University Press.

26. Stuart AJ. 2015 Late Quaternary megafaunal extinctions on the continents: a short
review. *Geol. J.* **50**, 338–363. (doi:10.1002/gj.2633)

27. Johnson CN *et al.* 2016 What caused extinction of the Pleistocene megafauna of Sahul?
Proc. R. Soc. B **283**, 20152399. (doi:10.1098/rspb.2015.2399)

28. Peters KJ, Saltré F, Friedrich T, Jacobs Z, Wood R, McDowell M, Ulm S, Bradshaw CJA.
2019 FosSahul 2.0, an updated database for the Late Quaternary fossil records of Sahul.
Sci Data **6**, 272. (doi:10.1038/s41597-019-0267-3)

29. Reed EH, Bourne SJ. 2009 Pleistocene fossil vertebrate sites of the south east region of
South Australia II. *T. Roy. Soc. South Aust.* **133**, 30–40.
(doi:10.1080/03721426.2009.10887108)

30. Macken AC, Reed EH. 2013 Late Quaternary small mammal faunas of the Naracoorte
Caves World Heritage Area. *T. Roy. Soc. South Aust.* **137**, 53–67.
(doi:10.1080/3721426.2013.10887171)

31. Rodríguez-Rey M *et al.* 2016 Data descriptor: a comprehensive database of quality-rated
fossil ages for Sahul's Quaternary vertebrates. *Sci. Data* , 7.

32. Rodríguez-Rey M *et al.* 2015 Criteria for assessing the quality of Middle Pleistocene to
Holocene vertebrate fossil ages. *Quat. Geochronol.* **30**, 69–79.
(doi:10.1016/j.quageo.2015.08.002)

33. Starrfelt J, Liow LH. 2016 How many dinosaur species were there? Fossil bias and true
richness estimated using a Poisson sampling model. *Phil. Trans. R. Soc. B* **371**, 20150219.
(doi:10.1098/rstb.2015.0219)

34. Feldman A, Meiri S. 2013 Length-mass allometry in snakes: snake length-mass allometry.
Biol. J. Linnean. Soc. **108**, 161–172. (doi:10.1111/j.1095-8312.2012.02001.x)

35. Garnett ST *et al.* 2015 Biological, ecological, conservation and legal information for all
species and subspecies of Australian bird. *Sci. Data* **2**, 150061.
(doi:10.1038/sdata.2015.61)

36. Jones KE *et al.* 2009 PanTHERIA: a species-level database of life history, ecology, and
geography of extant and recently extinct mammals: *Ecological Archives* E090-184.
Ecology **90**, 2648–2648. (doi:10.1890/08-1494.1)

37. Meiri S. 2010 Length-weight allometries in lizards. *J. Zool.* (doi:10.1111/j.1469-
7998.2010.00696.x)

38. Oliveira BF, São-Pedro VA, Santos-Barrera G, Penone C, Costa GC. 2017 AmphiBIO, a
global database for amphibian ecological traits. *Sci. Data* **4**, 170123.

39. Read QD, Baiser B, Grady JM, Zarnetske PL, Record S, Belmaker J. 2018 Tropical bird
species have less variable body sizes. *Biol. Lett.* **14**, 20170453.
(doi:10.1098/rsbl.2017.0453)

40. Santini L, Benítez-López A, Ficetola GF, Huijbregts MAJ. 2018 Length-mass allometries in
amphibians. *Integr. Zool.* **13**, 36–45. (doi:10.1111/1749-4877.12268)

41. Slavenko A, Tallowin OJS, Itescu Y, Raia P, Meiri S. 2016 Late Quaternary reptile
extinctions: size matters, insularity dominates. *Global Ecol. Biogeogr.* **25**, 1308–1320.
(doi:10.1111/geb.12491)

42. Wilman H, Belmaker J, Simpson J, de la Rosa C, Rivadeneira MM, Jetz W. 2014
EltonTraits 1.0: Species-level foraging attributes of the world's birds and mammals:
Ecological Archives E095-178. *Ecology* **95**, 2027–2027. (doi:10.1890/13-1917.1)

43. Cogger H. 2014 *Reptiles and amphibians of Australia*. CSIRO publishing.

44. Wilson SK, Swan G. 2008 *A complete guide to reptiles of Australia*. 2nd Ed. New Holland
Publishers.

45. Gravel D, Poisot T, Albouy C, Velez L, Mouillot D. 2013 Inferring food web structure from
predator-prey body size relationships. *Methods Ecol. Evol.* **4**, 1083–1090.
(doi:10.1111/2041-210X.12103)

46. Brose U *et al.* 2019 Predator traits determine food-web architecture across ecosystems.
Nat. Ecol. Evol. **3**, 919–927. (doi:10.1038/s41559-019-0899-x)

47. Brousseau P-M, Gravel D, Handa IT. 2018 Trait matching and phylogeny as predictors of
predator-prey interactions involving ground beetles. *Funct. Ecol.* **32**, 192–202.
(doi:10.1111/1365-2435.12943)

48. Eklof A, Helmus MR, Moore M, Allesina S. 2012 Relevance of evolutionary history for
food web structure. *Proc. R. Soc. B* **279**, 1588–1596. (doi:10.1098/rspb.2011.2149)

49. McLeod AM, Leroux SJ, Chu C. 2020 Effects of species traits, motif profiles, and
environment on spatial variation in multi-trophic antagonistic networks. *Ecosphere* **11**.
(doi:10.1002/ecs2.3018)

50. Morales-Castilla I, Matias MG, Gravel D, Araújo MB. 2015 Inferring biotic interactions
from proxies. *Trends Ecol. Evol.* **30**, 347–356. (doi:10.1016/j.tree.2015.03.014)

51. Pires MM, Koch PL, Fariña RA, de Aguiar MAM, dos Reis SF, Guimarães PR. 2015
Pleistocene megafaunal interaction networks became more vulnerable after human
arrival. *Proc. R. Soc. B* **282**, 20151367. (doi:10.1098/rspb.2015.1367)

52. Poelen JH, Simons JD, Mungall CJ. 2014 Global Biotic Interactions: an open infrastructure
to share and analyze species-interaction datasets. *Ecol. Inform.*
(doi:https://doi.org/10.1016/j.ecoinf.2014.08.005 accessed on 17/04/2019)

53. Llewelyn J, Strona G, Bradshaw CJA. 2020 Inferring networks and modelling trophic
cascades. *GitHub and Zenodo* (doi:10.5281/zenodo.3756224)

54. de Visser SN, Freymann BP, Olff H. 2011 The Serengeti food web: empirical
quantification and analysis of topological changes under increasing human impact:
Topological changes under human impact. *J. Anim. Ecol.* **80**, 484–494.
(doi:10.1111/j.1365-2656.2010.01787.x)

55. Chapman AD. 2009 Numbers of living species in Australia and the world.

56. Grosberg RK, Vermeij GJ, Wainwright PC. 2012 Biodiversity in water and on land. *Current
Biology* **22**, R900–R903. (doi:10.1016/j.cub.2012.09.050)

57. Page L, Brin S, Motwani R, Winograd T. 1999 The PageRank citation ranking: Bringing
order to the web.

58. Darimont CT, Fox CH, Bryan HM, Reimchen TE. 2015 The unique ecology of human
predators. *Science* **349**, 858–860. (doi:10.1126/science.aac4249)

59. Miller GH. 2005 Ecosystem Collapse in Pleistocene Australia and a Human Role in
Megafaunal Extinction. *Science* **309**, 287–290. (doi:10.1126/science.1111288)

60. Turney CSM *et al.* 2001 Redating the onset of burning at Lynch's Crater (North
Queensland): implications for human settlement in Australia. *J. Quaternary Sci.* **16**, 767–
771. (doi:10.1002/jqs.643)

61. DeSantis LRG, Field JH, Wroe S, Dodson JR. 2017 Dietary responses of Sahul (Pleistocene
Australia–New Guinea) megafauna to climate and environmental change. *Paleobiology*
43, 181–195. (doi:10.1017/pab.2016.50)

62. Flannery TF. 1990 Pleistocene faunal loss: implications of the aftershock for Australia's
past and future. *Archaeol. Oceania* **25**, 45–55. (doi:10.1002/j.1834-
4453.1990.tb00232.x)

63. Johnson CN. 2002 Determinants of loss of mammal species during the Late Quaternary
'megafauna' extinctions: life history and ecology, but not body size. *Proc. R. Soc. B* **269**,
2221–2227. (doi:10.1098/rspb.2002.2130)

64. Wood JR, Wilmshurst JM, Rawlence NJ, Bonner KI, Worthy TH, Kinsella JM, Cooper A.
2013 A megafauna's microfauna: gastrointestinal parasites of New Zealand's extinct
Moa (Aves: Dinornithiformes). *PLoS One* **8**, e57315. (doi:10.1371/journal.pone.0057315)

65. Roopnarine PD, Angielczyk KD, Wang SC, Hertog R. 2007 Trophic network models
explain instability of Early Triassic terrestrial communities. *Proc. R. Soc. B* **274**, 2077–
2086. (doi:10.1098/rspb.2007.0515)

66. Ceballos G, Ehrlich PR, Dirzo R. 2017 Biological annihilation via the ongoing sixth mass
extinction signaled by vertebrate population losses and declines. *Proc. Natl. Acad. Sci.*
USA **114**, E6089–E6096. (doi:10.1073/pnas.1704949114)

639
640

Figure captions

Figure 1. Example of an inferred ecological network model of the Late Pleistocene Naracoorte assemblage. Small points represent vertebrate species (nodes) and lines represent trophic interactions (links). Point colour shows trophic group (e.g., herbivores, carnivores etc), and extinct nodes are black. Due to uncertainty regarding trophic interactions, 1000 versions/models of the Naracoorte network were inferred and analysed. Plants, invertebrates and aquatic animals are shown as single large points in this figure because of the large number of nodes in these groups (see methods).

Figure 2. Vulnerability of Naracoorte trophic groups to coextinction from bottom-up coextinction cascades, and proportion of each trophic group among the extinct *versus* extant species groups. Panel (a) shows vulnerability to bottom-up coextinction cascades of three trophic groups: herbivores, mid-trophic level species (between trophic levels 2 and 3), and carnivores. A node's vulnerability to coextinction was determined by iteratively taking the product of the vulnerabilities of the focal node's resource nodes using an algorithm similar to PageRank. The horizontal lines inside the boxes correspond to the medians, the boxes indicate the upper and lower quartiles, whiskers show 1.5× the interquartile range, and points are outliers. Panels (b), (c), and (d) are density plots showing results from randomization tests comparing the proportion of each trophic group among the extinct *versus* extant species groups. A density plot centred on 0 indicates that the proportion of species belonging to that trophic group was similar/identical in the extinct and extant species groups.

Figure 3. Comparison of extinct *versus* extant predators from the Naracoorte network in terms of their vulnerability to bottom-up coextinction cascades. A node's vulnerability to coextinction was determined by iteratively taking the product of the vulnerabilities of the focal node's resource nodes using an algorithm similar to PageRank. Plot points and error bars show mean and 95% confidence intervals, respectively.

Figure 4. Comparison of extinct *versus* extant species in terms of their position in the Naracoorte trophic network. Panel a) is a biplot of the first two principal components from a principal component analysis of six variables describing species' position in the network. Panel b) is a boxplot showing the number of species that preyed on each extant *versus* extinct species. Panel c) is a boxplot of the number of resource nodes for extinct *versus* extant species. In the boxplots, the centre line indicates the median, the box shows the upper and lower quartiles, the whiskers indicate 1.5× the interquartile range, and the points are outliers. We inferred 1000 models of the Naracoorte network, and from these models calculated each species' mean value of each metric for use in the principal-components biplot, predator-diversity boxplot, and resource-diversity boxplot.