Genome streamlining, proteorhodopsin, and organic nitrogen metabolism in freshwater nitrifiers

Justin C. Podowski[a], Sara F. Paver[a], Ryan J. Newton[b], Maureen L. Coleman[a#]

[a] *Department of the Geophysical Sciences, University of Chicago, Chicago, Illinois, USA*

[b] *School of Freshwater Sciences, University of Wisconsin Milwaukee, Milwaukee, Wisconsin, USA*

Corresponding author:
   Maureen Coleman
   University of Chicago
   Department of the Geophysical Sciences
   Hinds Laboratory
   5734 S. Ellis Ave.
   Chicago, IL 60637

   Telephone: (773) 702-8352
   Fax: (773) 702-0207
   E-mail: mlcoleman@uchicago.edu

Running title: Nitrifiers across the Laurentian Great Lakes

Abstract

Microbial nitrification is a critical process governing nitrogen availability in aquatic systems. Freshwater nitrifiers have received little attention, leaving many unanswered questions about their taxonomic distribution, functional potential, and ecological interactions. Here, we reconstructed genomes to infer the metabolism and ecology of free-living picoplanktonic nitrifiers across the Laurentian Great Lakes, a connected series of five of Earth's largest lakes. Surprisingly, ammonia oxidizing Bacteria (AOB) related to *Nitrosospira* dominated over ammonia oxidizing Archaea (AOA) at nearly all stations, with distinct ecotypes prevailing in the transparent, oligotrophic upper lakes compared to Lakes Erie and Ontario. Unexpectedly, one ecotype of *Nitrosospira* encodes proteorhodopsin, which could enhance survival in conditions where ammonia oxidation is inhibited or substrate limited. Nitrite oxidizing Bacteria (NOB) *Ca. Nitrotoga* and *Nitrospira* fluctuated in dominance, with the latter prevailing in deeper, less productive basins. Genome reconstructions reveal highly reduced genomes and features consistent with genome streamlining, along with diverse adaptations to sunlight and oxidative stress and widespread capacity for organic nitrogen use. Our findings expand the known functional diversity of nitrifiers and establish their ecological genomics in large lake ecosystems. By elucidating links between microbial biodiversity and biogeochemical cycling, our work also informs ecosystem models of the Laurentian Great Lakes, a critical freshwater resource experiencing rapid environmental change.

Importance (150 words)

Microorganisms play critical roles in Earth's nitrogen cycle. In lakes, microorganisms called nitrifiers derive energy from reduced nitrogen compounds. In doing so, they transform nitrogen into a form that can ultimately be lost to the atmosphere by a process called denitrification, which helps mitigate nitrogen pollution from fertilizer runoff and sewage. Despite their importance, freshwater nitrifiers are virtually unexplored. To understand their diversity and

function, we reconstructed genomes of freshwater nitrifiers across some of Earth's largest

freshwater lakes, the Laurentian Great Lakes. We discovered several new species of nitrifiers

specialized for clear low nutrient waters, and distinct species in comparatively turbid Lake Erie.

Surprisingly, one species may be able to harness light energy using a protein called

proteorhodopsin, despite the fact that nitrifiers typically live in deep dark water. Our work reveals

unique biodiversity of the Great Lakes and fills key gaps in our knowledge of an important

microbial group, the nitrifiers.

Introduction

The oxidation of ammonia to nitrate powers the growth of nitrifying microorganisms and represents a critical flux in the global nitrogen cycle. Microbial nitrification of ammonia released from organic matter degradation produces nitrate, which can then be removed from the system by denitrification (1). As chemolithoautotrophs, nitrifiers are also a major source of dark carbon fixation (2), which may contribute significant organic carbon to the microbial food web of the ocean's interior (3–5) and of deep freshwater lakes (6).

Microbial nitrifiers are found in Archaea and several phyla of Bacteria, spanning diverse physiology and ecology; understanding the drivers and consequences of this diversity across ecosystems is a fundamental challenge. Ammonia-oxidizing Archaea (AOA) in the phylum *Thaumarchaeota* dominate the mesopelagic oceans (7), likely due to their high affinity for ammonia (8) and streamlined genomes (9). In freshwater systems, AOA are abundant in some oligotrophic lakes, while ammonia-oxidizing Bacteria (AOB) affiliated with the *Nitrosomonadaceae* (Betaproteobacteria) are highly variable but tend to dominate more eutrophic systems (10–16). Complicating this picture, however, there is considerable physiological variation within each group, such as low-nutrient-adapted clades of AOB (17, 18) and the ability of some strains to use alternative substrates like urea (18, 19). Within the AOA, there are also distinct ecotypes that appear to segregate with depth in the water column, in both marine (7) and freshwater systems (10). In freshwaters especially — which are poorly characterized compared to the oceans — it remains difficult to predict which AOA and AOB taxa are likely to dominate in a given system (16).

For aquatic nitrite oxidizers, which span the phyla *Nitrospira, Nitrospinae,* and *Proteobacteria,* niche differentiation is even less clear. The oceans are dominated by exclusively marine lineages (2, 20), consistent with ancient salinity-associated divergence. Cultivated strains of NOB show variation in substrate affinity and physiology (20–22), but the phylogenetic conservation of these traits, and their influence on environmental distributions, are

poorly understood. Moreover, recent studies have discovered that NOB are capable of

alternative energy metabolisms (23, 24) and can access nitrogen from cyanate and urea (25,

26), expanding their ecological potential. In freshwater systems, the NOB *Ca.* Nitrotoga

(Betaproteobacteria) was only recently discovered to be widespread (27), and the diversity of

this genus and factors favoring its success are unknown.

Here, we use the Laurentian Great Lakes as a model system to examine niche

partitioning among planktonic freshwater nitrifiers. The Great Lakes hold 20% of Earth's surface

freshwater and more than half of this volume receives little to no light (< 1% surface irradiance).

This system, while hydrologically connected, spans strong trophic and chemical gradients:

ultraoligotrophic Lake Superior supports rates of primary production and nitrification comparable

to the ocean gyres (28, 29), while Lake Erie supports greater production (30) and more than 70-

fold higher nitrification rate (31). Between these extremes, Lake Ontario has low ambient

ammonium concentrations like Lake Superior (32) but nitrification rates up to four times higher

(33). While previous studies reported that AOA and AOB dominate Lakes Superior and Erie,

respectively (14, 29), recent community profiling has revealed broader diversity in both

ammonia-oxidizing and nitrite-oxidizing lineages (34–36). We sought to link taxonomic,

genomic, and metabolic diversity of nitrifiers with the varied biogeochemistry of the Great Lakes,

using genome reconstructions and abundance profiling. Our results uncover novel lineages and

metabolic capabilities, and provide the first large-scale assessment of freshwater nitrifier

genomics.

Results and Discussion

Niche partitioning of nitrifiers across the Great Lakes

To map free-living picoplanktonic (here defined as cells that pass through a 1.6µm filter)

nitrifiers across the Great Lakes, we searched our recent 16S rRNA datasets for known

nitrifying taxa (34). We detected putative AOB in the genus *Nitrosospira* (Betaproteobacteria,

family *Nitrosomonadaceae*) and AOA similar to *Nitrosarchaeum* (family *Nitrosopumilaceae*),

along with putative NOB in the genera *Ca.* Nitrotoga (Betaproteobacteria, family

Gallionellaceae) and *Nitrospira* (family *Nitrospiraceae*). We did not detect 16S rRNA amplicons

from *Nitrosococcus*, *Nitrococcus*, *Nitrospina* or *Nitrobacter*. The highest relative abundances of

picoplanktonic nitrifiers were observed in deep samples from eastern Erie and Ontario (9-24%

of total amplicons), compared to 2-14% in Michigan, Huron and Superior; Erie and Ontario also

have higher cell concentrations and higher surface chlorophyll (Dataset S1). The relative

abundance of nitrifiers was negatively correlated with photosynthetically active radiation (PAR;

Spearman's rho = -0.89, $p < 2.2e-16$) and reached a maximum below the depth of 1% PAR in

each lake, up to 20% of amplicon sequences (Fig S1a). The relative abundances of ammonia-

and nitrite-oxidizing taxa were strongly correlated (Spearman's rho = 0.918, $p < 2.2e-16$; Fig

S1b). Free-living nitrifiers were rare (<0.1% relative abundance) in bottom water samples from

the southern basin of Huron (HU15M) and western basin of Erie (HU91M); these two stations

are the shallowest in our dataset and have relatively high light penetration to the bottom (~1%

PAR). Chlorophyll-*a* concentration was also negatively correlated with the relative abundance of

nitrifiers (Spearman's rho = -0.677, $p < 1.7e-7$; Fig S1c). These findings are consistent with

previous work demonstrating photoinhibition of nitrification (37–40), as well as potential

competition with phototrophs for ammonium (41).

The taxonomic assemblage of nitrifiers differed across lakes and even among stations

within a lake (Fig 1, Dataset S1), in association with variable productivity and nitrogen

availability. Surface ammonium is typically below 300 nM except in Erie, where it is several-fold

higher and spatially variable; nitrate, on the other hand, is very high across the lakes but lowest

in Erie due to biological uptake (42, 43). Few measurements of urea exist but it can exceed

ammonium (44) (Dataset S2). AOB (*Nitrosomonadaceae*) were observed across all lakes. In

contrast, AOA (*Nitrosopumilaceae*) sequences only exceeded 0.5% relative abundance at the

three deepest stations (SU08M, MI41M, ON55M) where the ratio of AOB:AOA ranged from 10:1

to 1:3. We found pronounced shifts in the dominant NOB across stations (Fig 1), and all stations

except those in Lake Ontario showed strong dominance (greater than ten-fold) of either *Ca.*

Nitrotoga (family *Gallionellaceae*) or *Nitrospira*. *Nitrospira* was the only nitrite oxidizer detected

in Superior and the dominant nitrite oxidizer in parts of Michigan (MI41M, MI18M). By contrast,

Ca. Nitrotoga was the only nitrite oxidizer observed in Erie and Huron and the dominant nitrite

oxidizer at the shallowest station in Michigan (MI27M). Within each taxon, a single 16S rRNA

oligotype dominated the AOA, *Ca*. Nitrotoga, and *Nitrospira,* while several oligotypes of

Nitrosomonadaceae shifted abundance across samples (Fig S2), consistent with ecotypic

diversity as discussed below.

***Ecotypic variation in abundant streamlined* Nitrosospira**

We reconstructed 15 genomes of the AOB *Nitrosospira*, substantially expanding genome

descriptions for this genus (45–47). Based on a phylogenomic tree, free-living Great Lakes

Nitrosospira fall into two major clades, both of which are distinct from published species; each of

these clades also includes MAGs recovered from lakes Biwa and Baikal, suggesting novel

globally distributed freshwater lineages (Fig S3). One clade, which we call NspGL1, has a highly

reduced genome (median 1.42 Mb) and low G+C content (40.7%) (Fig 2, Dataset S3). The

second clade was resolved into three subclades (denoted NspGL2a, 2b, and 3; Fig S3) based

on phylogeny and average nucleotide identity (ANI), all with small genome sizes of 1.45-1.68

Mb and 50% G+C content (Fig 2, Dataset S3). Compared to 86 reference *Nitrosomonadaceae*

genomes, Great Lakes *Nitrosospira* genomes are not only smaller (median estimated complete

genome size: reference = 3.21 Mb, GL = 1.45 Mb; Table 1), but also have shorter intergenic

spacers, fewer paralogs, fewer pseudogenes, and fewer sigma factors (Table 1, Fig S4, Dataset

S4), consistent with genome streamlining to reduce resource demands (48). Based on short

read mapping, these subclades are ecologically distinct: NspGL1 and NspGL2b — with the

smallest genomes — are the dominant AOB in the upper oligotrophic lakes, while NspGL2a is

only abundant in Ontario and NspGL3 is only abundant in Erie (Fig 2). Hereafter we refer to

these subclades as ecotypes due to their phylogenetic and ecological divergence.

We next compared gene content between our Great Lakes *Nitrosospira* and 86

Nitrosomonadaceae reference genomes. On average, Great Lakes *Nitrosospira* encode far

fewer two-component signal transduction systems (NspGL = 5-8, mean reference = 19),

transposases (NspGL = 0-7, mean reference = 39), motility genes (NspGL = 0-4, mean

reference = 52), pilus and secretion genes (NspGL = 2-9, mean reference = 27), and defense-

related genes (NspGL = 4-11, mean reference = 39) (Dataset S5). They also lack functions

related to biofilm formation such as polysaccharide matrix production (e.g. *pel* genes) and

extracellular protein targeting (exosortase and PEP-CTERM motifs). We note that while our 15

new *Nitrosospira* MAGs have high estimated completion (median 98.6%; Dataset S3), some

gene absences may reflect incomplete assemblies. Nevertheless, this overall picture of gene

content in Great Lakes AOB contrasts with *Nitrosospira* isolates from soil (45, 47) and even to

oligotrophic *Nitrosomonas* isolates (49), and is consistent with a passive planktonic lifestyle in

extremely low-nutrient systems.

We next compared metabolic potential among Great Lakes AOB ecotypes to understand

their ecological preferences for upper lakes (NspGL1, NspGL2b), Ontario (NspGL2a), and Erie

(NspGL3). Surprisingly, all seven NspGL1 MAGs encode proteorhodopsin, a light-driven proton

pump that supports bacterial energy production (50, 51). They also carry the genes necessary

to synthesize its chromophore retinal, including 15,15'-Beta-carotene dioxygenase (*blh*),

lycopene cyclase (*crtY*), phytoene dehydrogenase (*crtI*), phytoene synthase (*crtB*), and GGPP

synthase (*crtE*) ((52, 53); Fig 3a). We also identified proteorhodopsin in a single-cell amplified

genome representing NspGL1 from Lake Michigan (Fig 3a; 99.8% ANI with NspGL1 MAGs),

demonstrating that it is not an artifact of metagenome assembly. To our knowledge, this is the

first example of a nitrifier with proteorhodopsin. All NspGL1 proteorhodopsins share residues

H95, D127 and E138 along with a short beta-turn (G111-P116) between helices B and C, which

are characteristic features of proteorhodopsin as distinct from sensory and other rhodopsins

(54), and the presence of leucine at position 135 suggests green light tuning (55) (Fig 3b). All of

the genes in this module have highest similarity to homologs from *Polynucleobacter*, but are

flanked by Nitrosomonadaceae-like genes, suggesting recent horizontal gene transfer (Fig 3a).

The predicted NspGL1 proteorhodopsins cluster with *Polynucleobacter*, *Methylopumilus*, and

other freshwater Betaproteobacteria in Supercluster III as defined by MicRhoDE ((56); Fig 3c).

We compared the homologous genome region in two highly similar MAGs from Lakes Biwa and

Baikal (Fig S5); these contigs lack the proteorhodopsin module but appear to flank a variable

region where the contig assembly ends. A proteorhodopsin photosystem could support survival

of NspGL1 in the presence of sunlight, which has been shown to inhibit ammonia oxidation (37,

57). In the upper lakes where NspGL1 is abundant, light penetration is high well below the

thermocline in stratified periods (58), and deep water taxa are seasonally advected to the

surface by water column mixing (34). In addition to proteorhodopsin, NspGL1 — but not the

other three ecotypes of Great Lakes *Nitrosospira* — encode a class I cyclopyrimidine dimer

photolyase, which uses light energy to repair UV-induced DNA damage, and the catalase-

peroxidase *katG*, suggesting that the NspGL1 ecotype is adapted to relatively shallow depths in

the water column (Fig 2).

Great Lakes *Nitrosospira* carry a reduced, ecotype-specific complement of nitrogen

metabolism genes compared to reference AOB (Fig 2, Dataset S5; gene absences were verified

as described in Materials & Methods). All are presumed to have the core ammonia oxidation

enzymes ammonia monooxygenase and hydroxylamine dehydrogenase; these genes were

assembled and binned as expected in some MAGs, and were manually identified on short

unbinned contigs in other cases (Dataset S6; see Materials & Methods). Surprisingly, all Great

Lakes *Nitrosospira* MAGs lack the copper protein nitrosocyanin, whose precise function is

unknown but so far has been found in all described AOB except one member of the *N.*

oligotropha clade (49). Based on the expanded set of genomes analyzed here, the lack of

nitrosocyanin likely extends beyond the Great Lakes MAGs to closely related freshwater and

marine strains, along with five more members of the *N. oligotropha* clade (Fig S3); its absence

may be related to the divergence of these clades. Only NspGL1 and NspGL2b encode NO-

forming nitrite reductase (NirK), which confers nitrite tolerance (59); this result is surprising

given that these two clades dominate the upper lakes where productivity and reduced N are

lowest. None of the Great Lakes ecotypes encode NO reductase (NorCBQD) and NspGL1 lacks

cytochrome P460 family proteins, both of which are common in AOB and implicated in nitrogen

oxide metabolism (18, 49). Nitrogen acquisition is also distinct among Great Lakes AOB:

NspGL1 lacks an apparent ammonium transporter, but encodes urease structural and

accessory genes (*ureABCEFG*) and a high-affinity urea transporter (*urtABCDE).* Further, all

Great Lakes ecotypes encode a high-affinity amino acid transporter (*livFGHM*); these genes are

rare (<5%) in reference genomes and could supply reduced nitrogen and/or organic carbon.

Finally, NspGL1 and NspGL3 have genes for producing cyanophycin, an intracellular storage

compound for nitrogen (47, 60). Together, the distinctive gene complements present in Great

Lakes *Nitrosospira* illustrate the variability and adaptability of AOB gene content, even across a

connected freshwater habitat.

As with nitrogen metabolism, carbon metabolism is also distinct between Great Lakes

and reference AOB, and among Great Lakes ecotypes (Fig 2, Dataset S5). Unlike most

reference AOB, Great Lakes *Nitrosospira* lack two key enzymes of the oxidative pentose

phosphate pathway, glucose-6-phosphate dehydrogenase and 6-phosphogluconate

dehydrogenase. All ecotypes except Erie-specific NspGL3 also lack genes for glycogen

synthesis and degradation, suggesting that they are unable to store and access this carbon

reserve. The key enzyme for carbon fixation, RuBisCO, has evolved several kinetically distinct

forms whose distribution likely reflects ecological pressures (61). NspGL1 and NspGL3 both

contain Form IA RuBisCO, while NspGL2a and NspGL2b contain Form IC RuBisCO (Fig 2; (61,

62)). NspGL1 genomes also possess an alpha carboxysome-like *cso* operon, similar to

Nitrosomonas eutropha C91 (62), though our draft assembly lacks the expected carbonic

anhydrase (*csoS3/csoSCA*). Carboxysome-associated RuBisCO may allow NspGL1, the

ecotype most strongly adapted to energy and nutrient limitation, to more efficiently fix CO_2 by

minimizing the wasteful oxygenation reaction and reducing the cellular nitrogen allocation to

RuBisCO (61). The ranges of kinetic properties observed in other autotrophs for Form IAq

(found in NspGL3) and Form IC (found in NspGL2a and NspGL2b) overlap, and therefore more

work is needed to understand the fitness advantages, if any, that this RuBisCO diversity confers

on Great Lakes nitrifiers.

***Streamlined freshwater* Thaumarchaeota**

We reconstructed three similar genomes (>99% ANI) of *Nitrosarchaeum* (NarchGL; Fig

S6, Dataset S3) from three separate samples (two from Superior, one from Ontario), consistent

with our low observed 16S rRNA diversity for *Thaumarchaeota*. These NarchGL genomes are

very similar (~99% ANI) to two genomes reconstructed from Lake Baikal, located thousands of

kilometers away (63). Their next closest relatives are also from freshwater environments, and

phylogenetic clustering suggests that salinity is an important driver of divergence throughout the

Nitrosopumilaceae (Fig S6). As a group, the *Thaumarchaeota* tend to have smaller genomes,

lower G+C content, higher coding density, and fewer paralogs and pseudogenes than nitrifying

bacterial taxa; even within this group, NarchGL genomes fall below the 30th percentile in size

and have significantly fewer paralogs than average (Table1, Fig S4, Dataset S4). Using our

reconstructed genomes as probes for metagenomic read recruitment, NarchGL were detected

in Superior, Michigan, and Ontario; they represented roughly one-third of ammonia oxidizers in

the mid-hypolimnion of station SU08M (Fig 2).

NarchGL share nearly 90% of their predicted proteins with close relatives including *Ca.*

Nitrosarchaeum limnia; at the same time, they show distinctive patterns in gene content that

pinpoint the key selective pressures of deep lakes (Fig 2, Dataset S5). All three NarchGL

genomes encode urease and a urea transporter, implicating urea as a vital source of nitrogen

for energy and/or biosynthesis. Consistent with phosphorus scarcity in much of the Great Lakes

(64), NarchGL encode high affinity transport systems for phosphate and potentially

phosphonates, though we did not identify a phosphonate lyase. In addition to both CRISPR/Cas

enzymes cas1 and cas4, NarchGL genomes contain several phage proteins, suggesting viral

infection and integration events may be common. DNA photolyases, which have been found in

epipelagic clades of marine *Thaumarchaeota* (7), are present in all low salinity *Nitrosarchaeum*

including NarchGL, suggesting NarchGL are exposed to sunlight due to high water clarity (58)

and/or annual mixing in the Great Lakes. NarchGL also lack the common tRNA modification 4-

thiouridylation (indicated by K04487 and PF02568-PF18297 (65)); we propose that the absence

of this modification, which is susceptible to near-UV radiation (65), is also related to sunlight

exposure.

Genomes of NarchGL reveal striking reduction in environmental sensing, response, and

regulatory functions, relative to most other *Nitrosarchaeum* and *Nitrosopumilaceae* (Dataset

S5). NarchGL encode 9-12 domains representing the general archaeal transcription factors,

TATA binding protein (TBPs, PF00352) and transcription factor B (TFBs, PF00382 and

PF08271), compared to 21 in *Ca.* N. limnia. NarchGL lack common domains found in two-

component systems that transmit environmental signals to control gene expression or protein

activity (domains PF02743, PF00672, PF00512, PF00072; NarchGL = 0, *Ca.* N. limnia = 53-54

copies per genome). Further, they are depleted in ArsR family transcription factors (PF01022; 0

copies in NarchGL vs. 2-3 in *Ca.* N. limnia), P-II proteins for regulation of nitrogen metabolism

(PF00543; 1 copy per NarchGL genome vs. 5 in *Ca.* N. limnia), and other potential regulatory

domains (CBS PF00571: 5 in NarchGL vs. 18-19 in *Ca.* N. limnia; USP PF00582: 1 in NarchGL

vs. 15 in *Ca.* N. limnia). This extremely limited regulatory capacity in NarchGL stands in sharp

contrast to closely related *Ca.* N. limnia, but instead parallels the oceanic minimalist *Ca.*

Nitrosopelagicus brevis (9).

Expanded diversity of Ca.* Nitrotoga *with reduced genomes

Despite the broad distribution of *Ca.* Nitrotoga in freshwater systems and beyond, only

six genomes are available, derived from rivers heavily impacted by urban and agricultural

influence, a wastewater treatment plant, and coastal sediment (27, 66, 67). Hence the metabolic

and phylogenetic diversity of this group is virtually unexplored. We reconstructed six new MAGs

of *Ca.* Nitrotoga, which form two clusters with >99% ANI within each cluster and ~97% ANI

between clusters (NtogaGL1a and NtogaGL1b; Fig S3). These new *Ca.* Nitrotoga MAGs are far

smaller than published genomes (median GL = 1.44 Mb, reference = 2.61-2.98 Mb), have

shorter intergenic regions, fewer sigma factors, and fewer paralogs (Table 1, Fig S4, Dataset

S4), consistent with genome streamlining (48). They lack functions such as motility and

chemotaxis, pilus biogenesis, and DNA repair (*mutLS*) (Dataset S5). Great Lakes *Ca.* Nitrotoga

also encode markedly fewer two-component systems for sensing and responding to

environmental cues than four river-derived genomes (NtogaGL = 2-6 per genome vs. 30-35 in

four reference genomes; reference strain KNB has 7). Compared to reference genomes,

NtogaGL have fewer defense-related genes (restriction-modification, toxin-antitoxin, and

CRISPR-Cas systems; mean NtogaGL = 11 vs. 39 for references), and transposases (mean

NtogaGL = 3 vs. 19 for references) (Dataset S5). While incomplete assembly of hypervariable

genome regions may explain some of these absences, the overall genome properties are

consistent with a relatively stable low-nutrient environment and planktonic lifestyle.

The reduced genomes of NtogaGL1a/b help clarify core features of the genus *Nitrotoga*,

along with accessory functions that may enable local adaptation in specific populations. To date,

sequenced *Nitrotoga* including NtogaGL1a/b encode similar electron transport pathways,

including NADH dehydrogenase-Complex I, succinate dehydrogenase-Complex II, and

Alternative Complex III, along with high-affinity cbb3-type cytochrome oxidases suggesting

adaptation to low oxygen conditions. They also share the Calvin cycle for carbon fixation, a

complete TCA cycle, and an evolutionarily distinct nitrite oxidoreductase (NXR) from other NOB

(27, 66, 67). All *Nitrotoga* to date also share transporters for amino acids and peptides, potential

sources of C and/or N. *Nitrotoga* can also potentially access reduced sulfur compounds for

energy via sulfite dehydrogenase, suggesting metabolic flexibility beyond nitrite oxidation.

Beyond these similarities, the small genomes of NtogaGL1a/b are distinct from

previously described *Nitrotoga* in many ways. NtogaGL1a/b lack NiFe hydrogenase to use

hydrogen as an energy source. They also lack nitrogen metabolism functions including

assimilatory nitrite reductase (*nirBD*) and nitrite reductase to NO (*nirK*). Based on gene content,

NtogaGL1a/b appear unable to use hexoses like glucose, since they lack the glycolytic enzyme

phosphofructokinase and the Entner-Doudoroff pathway, similar to *Nitrobacter winogradskyi*

(68). Consistent with this, they also lack genes for storage and breakdown of glycogen (Dataset

S5). All but one of the NtogaGL1a/1b genomes encode cyanate lyase (*cynS*), which is found in

other NOB but no *Nitrotoga* to date (25, 69, 70). The *cynS* gene, adjacent to *glnK-amtB* for

ammonium sensing and transport, likely functions in N assimilation, as recently described for

Nitrospinae (71). While cyanase has been shown to mediate reciprocal feeding between some

NOB and ammonia oxidizers (25), it remains to be seen whether such an interaction occurs in

the free-living (<1.6um) size-fraction and dilute environment sampled here. Notably, cyanase

from NtogaGL1a/1b, along with predicted *Nitrospirae* proteins from Lake Baikal and soil, form a

distinct phylogenetic cluster from most nitrifier cyanase proteins observed to date (Fig S7).

The two ANI-based clusters we detected, NtogaGL1a and NtogaGL1b, appear to be

phylogenetically and ecologically distinct ecotypes. Based on short-read mapping, NtogaGL1b

dominates Erie, while NtogaGL1a dominates all other *Ca.* Nitrotoga-containing samples (Fig 4).

We found several metabolic genes that differentiate the two ecotypes. ER-specific NtogaGL1b

genomes share a region encoding thiosulfate dehydrogenase (*tsdA*), cytochromes, transport of

sulfur-containing compounds, lactate dehydrogenase (ldh*), a two-component system, and a

Crp-family transcription factor (Fig S8). This region may be involved in oxidizing thiosulfate as

an energy source, and sensing and responding to redox changes that accompany seasonal

hypoxia in Lake Erie. The corresponding region in NtogaGL1a encodes an integrase and

photolyase, consistent with greater DNA photodamage in the more transparent waters of

Michigan, Huron, and Ontario where NtogaGL1a is abundant.

Great Lakes* Nitrospira *reveal adaptations to sunlit oxic environment

We reconstructed six closely related genomes of *Nitrospira* (~99% ANI; Fig S9, Dataset

S3), representing the predominant NOB throughout Lake Superior and in parts of Michigan,

Ontario, and Huron (Fig. 4; Dataset S1). These genomes, which we refer to as NspiraGL, fall

within lineage II (Fig S9), which is broadly distributed across soil, freshwater, and engineered

habitats (20); however, genome analyses to date have focused on strains from wastewater and

engineered systems, leaving major blind spots. NspiraGL share core features of *Nitrospira*

metabolism, including a periplasmic-facing NXR that is advantageous under substrate-limiting

conditions, multiple cytochrome *bd*-like oxidases, and the reverse TCA cycle for carbon fixation

(69). However, as with *Ca.* Nitrotoga, the *Nitrospira* genomes we reconstructed in the Great

Lakes are markedly smaller than published reference genomes (median NspiraGL = 1.83 Mb,

median reference = 3.72 Mb), with higher coding density and fewer paralogs, sigma factors and

pseudogenes (Fig. S4; Dataset S4), consistent with genome streamlining theory (48).

Compared to 75 lineage II *Nitrospira* reference genomes, NspiraGL have reduced capacity for

environmental sensing (two-component systems: NspiraGL = 7, mean reference = 26), transport

(NspiraGL = 76-83, mean reference = 140), defense (NspiraGL = 7-8, mean reference = 26),

and transposition (NspiraGL = 0-2, mean reference = 15), and lack pilus or flagellar motility

(Dataset S5). NspiraGL encode just five sigma factors, compared to 18 in *N. moscoviensis.*

Further, NspiraGL genomes encode a single NXR, while *N. moscoviensis* carries five copies

that are differentially regulated (26, 72). NspiraGL also lack the *glnE* gene for glutamine

synthetase (GS) adenylyltransferase, suggesting that GS activity is not repressed by this

mechanism. Together, these features suggest limited regulatory and ecological flexibility,

consistent with a relatively constant, oligotrophic environment.

Compared to other *Nitrospira,* NspiraGL exhibit limited energetic flexibility, but can

access diverse nitrogen sources (Fig. 4, Dataset S5). We predict that NspiraGL are unable to

grow on hydrogen or formate as alternative energy sources (23, 26), as they lack NiFe-

hydogenase and formate dehydrogenase. The glycolysis and oxidative TCA cycles appear to be

incomplete, lacking phosphofructokinase and citrate synthase, respectively; this suggests a

limited capacity for organic carbon utilization. NspiraGL lack *nirK,* encoding NO-forming nitrite

reductase, which is found in a majority of reference genomes. To obtain nitrogen for

biosynthesis, NspiraGL encode a high-affinity nitrate/nitrite/cyanate transporter (*nrtABC*),

assimilatory nitrite reductase (*nirA*), and cyanase (*cynS*), along with *amt* family ammonium

transporter. Although none of the NspiraGL MAGs include urease (*ureCBA*), one does contain

urease accessory proteins (*ureEFGD*) and two contain a urea transporter (*urtABCD*),

suggesting incomplete assembly of the urea utilization pathway. As with *Ca.* Nitrotoga, we

suggest that cyanase, along with urease where present, functions in nitrogen assimilation rather

than cross-feeding, given the dilute environment and free-living planktonic cells.

Beyond energy, carbon, and nitrogen metabolism, we discovered striking differences

between NspiraGL and reference *Nitrospira* related to DNA repair. NspiraGL encode two

additional photolyase-related proteins, along with a class I cyclopyrimidine dimer (CPD)

photolyase found in most reference *Nitrospira* (Fig 5). Photolyases use blue light energy to

repair DNA lesions caused by UV radiation (73). The two additional genes in NspiraGL are

adjacent and share best hits with *Betaproteobacteria*, suggesting recent horizontal transfer (Fig

S10). One likely encodes an FeS-BCP photolyase, which repairs (6-4) dipyrimidine lesions (74,

75). The other shares an FAD-binding domain with photolyases but the C-terminal region has

no recognizable domains (Fig 5). This protein is widespread in aquatic bacteria and has not

been functionally characterized, though an actinobacterial homolog was suggested to be

involved in light sensing and regulation (76). Beyond photolyases, NspiraGL also encode uracil-

DNA glycosylase (UNG), which removes misincorporated uracil from DNA. Uracil results from

deamination of cytosine, which can occur spontaneously or be induced by NO (77). In addition

to the photolyases and UNG that repair DNA lesions, NspiraGL encode translesion DNA

polymerase V (*umuCD*) which enables replication to proceed past lesions. Together, these

genes indicate that *Nitrospira* in the Great Lakes experience significant DNA damage, including

UV-induced damage that also requires light for the repair process, in hypolimnion waters with

high transparency (58) and/or during seasonal mixing.

Other major differences between NspiraGL and reference *Nitrospira* are related to

reactive oxygen species (ROS) (Dataset S5). Surprisingly, despite their oxic habitat, NspiraGL

lack superoxide dismutase (SOD), monofunctional catalase (*katE*), and bacterioferritin, which

limits the Fenton reaction by sequestering free iron. However, all six NspiraGL MAGs, but few

reference genomes (7% of 75), have recently acquired bifunctional catalase-peroxidase *katG*;

interestingly we also observed *katG* in Great Lakes *Ca.* Nitrotoga and *Nitrosospira* (Fig 2, Fig 4).

The absence of SOD suggests that NspiraGL does not produce damaging levels of endogenous

superoxide, perhaps because NspiraGL lack the major respiratory and non-respiratory

flavoproteins that produce ROS in other SOD-containing *Nitrospira* (78). Unlike superoxide,

H_2O_2 can cross membranes, and is known to be produced by both photooxidation of dissolved

organic matter and dark heterotrophic activity (79). The lakes where NspiraGL dominate have

high water clarity (58) and low productivity and are fully oxic, consistent with abiotic

photochemistry as the primary source of exogenous ROS; this stress may have selected for

katG as a defense. NspiraGL also lack cytochrome *c* peroxidase, which is found in 70 of 75

reference genomes; this protein is proposed to function in anaerobic respiration of H_2O_2 (80)

and therefore its absence in NspiraGL is consistent with a constant oxic environment. Together,

these results indicate that *Nitrospira* in the Great Lakes face distinct ROS pressures that have

shaped their gene content.

Conclusions

The Laurentian Great Lakes harbor nitrifiers that are phylogenetically related, but

markedly different in genome size and functional capacity, from their well-studied relatives

inhabiting wastewater systems, soils, and even other freshwater systems. By examining the

entire nitrifier assemblage at once, we detected common features across taxa that illuminate the

selective pressures faced by microbes in deep lakes. All the lineages we describe show small

genome sizes (1.3-1.7 Mb), reduced capacity for environmental sensing and response, and

adaptation to a passive (i.e. non-motile) planktonic lifestyle, features which have not been

previously associated with AOB, *Nitrospira*, and *Ca.* Nitrotoga. Within the AOB *Nitrosospira*, we

found ecotypes with a gradient of genome reduction that maps onto their habitats' trophic

gradient: from NspGL1 (1.4 Mb, low GC, upper lakes) to NspGL2b (1.5 Mb, upper lakes) to

NspGL2a (1.6 Mb, Ontario) to NspGL3 (1.7 Mb, Erie) (Fig 2). The thaumarchaeal NarchGL have

markedly reduced regulatory capacity like the open ocean strain *Nitrosopelagicus brevis* (9).

The NOB NspiraGL have genomes 50-60% smaller than described *Nitrospira* and dominate,

along with the AOB NspGL1 and the AOA NarchGL, the deeper more oligotrophic basins, while

Ca. Nitrotoga favor shallower, more productive basins. The emergence of Erie-specific ecotypes

of both *Nitrosospira* (NspGL3) and *Ca.* Nitrotoga (NtogaGL1b) demonstrates how distinct this

habitat is compared to the other lakes. Importantly, our findings here represent planktonic cells

in the smallest size fraction (<1.6 µm); it is likely, especially in Erie, that particle-associated

nitrifiers may be abundant and genetically distinct.

Nitrifiers inhabiting the transparent waters of the upper Great Lakes show distinctive

adaptations to light including diverse photolyases, ROS detoxification, and even

proteorhodopsin. This discovery is surprising, given that nitrifiers are rare in the surface mixed

layer of the Great Lakes (Fig. S1) and that photoinhibition of ammonia oxidation and nitrifier

growth is well documented (37, 40, 57). We propose that proteorhodopsin could be used to

augment energy metabolism when ammonia oxidation is photoinhibited and/or ammonia is

substrate limited. Water clarity has increased over the past several decades in Lakes Michigan

and Huron, now surpassing that of Lake Superior (58). High light penetration along with

seasonal mixing likely exposes deep water cells to damaging levels of light and oxidative stress.

Future cultivation and physiological studies should examine photoinhibition and potential

phototrophy in Great Lakes nitriifiers.

The capacity for nitrification is found across multiple phyla, and our work unveils new

clues to understanding the ecological and evolutionary potential of these diverse lineages. This

collective nitrifier diversity undoubtedly influences the cycling of carbon and nitrogen across this

ecosystem, and future work will explore the differential contributions to nitrification by the distinct

lineages we described here. Understanding what controls the diversity of nitrifiers and other key

functional groups, and the consequences of this diversity for biogeochemistry, are essential for

forecasting the effects of rapid environmental change across the large lakes of the world

(e.g.(81)) and predicting impacts on the critical ecosystem services they provide (82).

Materials & Methods

Sample collection

Water samples were collected from the Laurentian Great Lakes aboard the R/V *Lake

Guardian*, during the biannual Water Quality Surveys conducted by the U.S. EPA Great Lakes

National Program Office (83). Station information is provided in Supplemental Dataset S8. Data

presented here were collected in April and August 2012. Samples were collected using a CTD

rosette sampler (Sea-Bird Scientific) at the surface (2 m), deep chlorophyll maximum (if

present), the mid-hypolimnion (depths ranging from 19 m in Erie to 200 m in Superior; Dataset

S1), and near the bottom of the water column (10 m above the lake bottom at most stations, 1 m

above bottom at shallow stations). For each sample, 5-8L of water was pre-filtered through a

GF/A glass fiber filter (Whatman 1820-047; nominal pore size 1.6 µm) to exclude eukaryotic

phytoplankton and particle associated microbes, and cells were collected on 0.22 µm Sterivex

filters (Millipore SVGP01050). Filters were stored at -80°C. For dissolved nutrient analysis, 0.22

µm filtrate was collected in 125 ml acid-clean HDPE bottles (Nalgene) and stored at -20°C.

Samples for single cell amplified genomes (SAG) were collected in August 2014. For each

sample, 1 ml of raw water was incubated with 100 µl of glycerol-TE buffer (20 ml 100X TE pH 8

+ 100 ml glycerol + 60 ml water; final concentration after sample addition is 10mM Tris, 1mM

EDTA, 5% glycerol) for 10 minutes in the dark, then flash frozen in liquid nitrogen and stored at -

80°C until processing.

Physicochemical data

CTD profiles, water chemistry and chlorophyll-*a* data were collected by the U.S. EPA

according to standard protocols (84) and retrieved from the Great Lakes Environmental

Database (https://cdx.epa.gov/) for 2012 and 2013. In addition, we measured dissolved nitrogen

species from August 2013 samples. Ammonium concentrations were measured using the OPA

method in a 96-well plate (85). Nitrate and nitrite concentrations were measured using the

Greiss reaction method in a 96-well plate (86). Urea concentrations were measured in a 24 well

plate using a colorimetric reaction (87).

16S rRNA analysis

The full 16S rRNA amplicon dataset was described by Paver and colleagues (34). Here

we focus on data from the V4☐V5 region (primers: 515F☐Y, 926R (88)), collected in 2012 in

tandem with metagenome samples from select stations. We classified sequences using the

Silva v. 132 database (89) and the wang method (90) as implemented by mothur (91).

Sequences classified to each detected family of nitrifiers (ammonia oxidizer families

Nitrosomonadaceae and *Nitrosopumilaceae*; nitrite oxidizer families *Gallionellaceae* and

Nitrospiraceae) with a mothur-assigned confidence score above 90 were delineated into

taxonomic units using minimum entropy decomposition with a minimum substantive abundance

of 10 (92).

Metagenome and single-cell genome sequencing

One station per lake in Superior, Michigan, Huron, and Ontario, and two stations in Erie,

were selected for metagenome sequencing. Spring 2012 metagenome samples were collected

from the surface, and Summer 2012 metagenome samples were collected from the mid-

hypolimnion (depths listed in Dataset S1). DNA was extracted using a modified

phenol:chloroform extraction protocol (34) and libraries prepared according to the Illumina

TruSeq protocol. Samples from spring 2012 were sequenced at the Joint Genome Institute

using Illumina HiSeq (2x150bp). Samples from summer 2012 were sequenced at the University

of Chicago Functional Genomics Core Facility using Illumina HiSeq 2500 (2x250bp).

To confirm the presence of proteorhodopsin, we analyzed a single cell amplified genome

from *Nitrosospira* collected from Lake Michigan and sequenced by the Joint Genome Institute.

Quality filtered reads were downloaded from JGI IMG/ER and normalized using bbnorm.sh with

target=100 and mindepth=2. Normalized reads were assembled using SPAdes 3.1.11 in single

cell mode (93) with flags --sc and --careful. Resulting scaffolds were annotated identically to

MAGs as described below.

Obtaining metagenome-assembled genomes

Raw reads for spring surface samples were quality controlled at the Joint Genome

Institute, using bbduk.sh for adapter trimming (ktrim=r, minlen=40, minlenfraction=0.6,mink=11,

tbo, tpe, k=23, hdist=1, hdist2=1, ftm=5) and quality filtering (maq=8, maxns=1, minlen=40,

minlenfraction=0.6, k=27, hdist=1, trimq=12, qtrim=rl). Raw reads for summer hypolimnion

samples were adapter trimmed, quality filtered, and interleaved using bbduk (parameters:

ktrim=r, mink=8, hdist=2, k=21, forcetrimleft=10, forcetrimright=199, minlen=150) using BBTools

suite version 35.74 (https://sourceforge.net/projects/bbmap/). Separate assemblies of quality

filtered reads were carried out for each metagenome using metaSPAdes 3.1.11 --meta mode

using default k sizes of 21, 33, 55 (94). To enable binning based on sequence coverage,

forward and reverse reads were merged using bbmerge in BBtools, using qtrim2=r

trimq=10,13,16 and adapter=default. Merged short reads were then mapped onto each

assembly using bowtie2 2.2.9 in --sensitive mode (95), and this coverage information was used

to bin assembled contigs. Binning was performed using MetaBAT2 2.12.1 (96), Binsanity 0.2.6.3

(97) and CONCOCT 1.0.0 (98) using default parameters. The resulting bins were scored,

aggregated, and de-replicated using DAS_Tool 1.1.1 (99) followed by manual curation using

Anvi'o 4.0 (100). We assessed genome completion and contamination of manually curated bins

using CheckM 1.1.0 lineage_wf (101), and all new MAGs presented here are greater than 70%

complete with less than 10% contamination (Dataset S3). Potential nitrifiers were screened by

searching for ammonia monooxygenase, hydroxylamine oxidoreductase and nitrite

oxidoreductase within reconstructed genomes using blastp 2.5.0 (102). For bins where any of

these genes were detected, we identified bacterial single copy core genes (103) or archaeal

single copy core genes (104) using HMMER (105), as implemented in Anvi'o. Single copy core

genes were queried against proteins predicted from bacterial and archaeal genomes in RefSeq

(NCBI) (106), and taxonomic identity of these core genes was ascertained based on a least

common ancestor approach using a 0.1% window around the bit score of the best hit using

KronaTools 2.7.1 (107). Taxonomic assignment was further validated using GTDB-tk 1.0.0

(108). Grouping of MAGs into clades and subclades based on ANI was carried out using

fastANI 1.1.0 (109). Genome characteristics for each genome group were calculated as the

median of those values for the group. Estimated complete genome size was calculated for

MAGs and for references in the pangenome analysis using CheckM (101) completion and

contamination, as follows: Estimated = Actual*(1-Contamination) / Completion. To quantify the

abundance of each clade/ecotype across samples, we used competitive mapping of merged

short reads using bowtie2 in sensitive mode against all nitrifier MAGs, summing up mapped

read count across all MAGs in a given clade/ecotype, and dividing by total mapped nitrifier

reads in a sample; these values are shown in bubble plots (Fig 2, 4).

Annotation and gene cluster analysis
553
Reference genomes were obtained from GenBank (accession numbers listed in

Supplemental Dataset S4). The full pangenome analyses included all the genomes listed in

Supplemental Dataset S4, but we only report results from the subset of genomes most closely

related to our MAGs. This subset consists of 86 *Nitrosomonadaceae*, 5 *Ca.* Nitrotoga, 78

Nitrosopumilaceae within Thaumarchaeota, and 75 *Nitrospira* that fall within Lineage II.

Reference genomes were treated consistently with GL MAGs, with *de novo* gene calling by

prodigal 2.6.3 (110) via Anvi'o. Unless otherwise noted, default settings were used for all

software. Genes were annotated using InterProScan 5.30-69.0 (111), GhostKOALA (112) and

eggnog-mapper 1.0.3 against the bactNOG database (113). Gene cluster analysis was carried

out using the Anvi'o pangenome pipeline (114), using blastp to determine sequence similarity,

ITEP to eliminate weak similarity (115) and MCL to cluster, using a minbit of 0.5, MCL inflation

of 2 and minimum gene occurrence of 1 (116). Sigma factors were tallied by identifying gene

clusters annotated with the following PFAMs: PF00309, PF03979, PF00140, PF04542,

PF04539, PF04545, PF08281. Pseudogene counts were retrieved where available from NCBI

PGAP annotated genomes (117). Paralog counts are reported as the number of gene clusters

with more than one gene per genome. Intergenic spacers were calculated using bedtools

complementBed function (118). Prokka 1.14.5 (119) was used to generate GenBank-format files

from MAGs and SAGs, and genoPlotR 0.8.9 (120) was used to generate initial gene

neighborhood maps.

Gene tree construction

The NspGL1 proteorhodopsin sequence was inserted into the MicRhoDE rhodopsin tree

using pplacer (121) through the MicRhoDE Galaxy pipeline (56). We then constructed a more

targeted phylogenetic tree using aligned reference sequences of Supercluster III from

MicRhoDE, filtered to exclude fragments shorter than 220 amino acids. To this alignment, we

added NspGL1 sequences using MAFFT 7.310 (122) along with high similarity sequences from

NCBI nr that were not present in MicRhoDE. The tree was inferred using RaxML 8.2.12 with

model PROTGAMMALG (123). The tree was visualized in iTOL (124) and more distant clusters

were collapsed for clarity.

A cyanase phylogenetic tree was created using sequences drawn from querying

NtogaGL cyanase against NCBI nr using blastp, as well as sequences from references (2, 25,

125). Sequences were aligned using MAFFT (122) and the tree was inferred using RaxML

8.2.12 with model PROTGAMMALG (123). Tree was visualized in iTOL (124) and branches

were colored based on the taxonomy of the parent genome.

Photolyase-related proteins in GL MAGs were identified by searching for the following

features: K01669, COG0415, PF03441, PF00875, PF04244, SSF48173, SSF52425. Reference

proteins (n=56) spanning the previously defined families of photolyases and cryptochromes

(126) were obtained from UniProt, along with aquatic bacterial sequences described by

Maresca and colleagues (76). The reference sequences were aligned using MAFFT (122), and

sequences from GL MAGs were added using the MAFFT --addfragments option. The tree was

estimated using IQ-TREE 2 1.6.11 (127) and visualized using iTOL (124).

Phylogenomic tree construction

Nitrospirae, *Thaumarchaeota*, *Gallionellaceae* and *Nitrosomonadaceae* genomes were

downloaded from Genbank (NCBI) (128) and included in phylogenomic trees for their respective

family. Phylogenomic analyses were carried out within Anvi'o. Briefly, single copy core genes

were extracted as described above, individually aligned at the protein level using muscle (129),

and concatenated for each genome. Concatenated alignments were trimmed using Gblocks

0.91b (130) and analyzed by RAxML 8.2.12 (123) to create a phylogenetic tree using the

PROTGAMMALG model and 50 bootstraps. Trees were visualized in iTOL (124).

<u>Proteorhodopsin assembly verification</u>

We used several approaches to validate the presence of proteorhodopsin in assembled

Nitrosospira genomes, to rule out the possibility of chimeric assemblies from different species.

We note that proteorhodopsin-containing contigs were independently assembled and binned

together with core *Nitrosospira* contigs from seven different samples (i.e. each sample was

assembled and binned separately, rather than co-assembled). In five of seven cases,

proteorhodopsin and retinal biosynthesis genes were assembled together with core *Nitrosospira*

genes on the same contig. To rule out a systematic reproducible error in assembly and/or

binning, we compared these seven MAGs to a single cell amplified *Nitrosospira* genome (SAG)

from Lake Michigan, obtained as part of another project with the JGI. This SAG was processed

through JGI's standard decontamination pipeline and manually investigated to ensure lack of

contamination. We found no evidence of contaminating core genes, as all core genes had best

hits to either *Nitrosospira* or more generally *Nitrosomonadaceae* in nr. SAG contigs were

matched to homologous contigs from NspGL1 MAGs to determine if any SAG contigs were

unique using FastANI 1.1.0 (109) with --visualize flag. All contigs from this *Nitrosospira* SAG

were found within an NspGL1 MAG. Bandage 0.8.1 (131) was used to manually inspect the

assembly graph around the contig that contained the NspGL1 *Nitrosospira* proteorhodopsin to

ensure that the assembled contig did not represent a chimeric contig or inappropriate

scaffolding. We verified that a single, unique path exists from the beginning to the end of the

NspGL1 contig containing proteorhodopsin (Fig 3). Further, we verified that consistent coverage

across this contig existed by mapping short reads from the original sample using bowtie2 (95)

and viewing results using Integrated Genomics Viewer 2.7.0 (132) A closely related assembly

of the same genomic region from Lake Biwa did not show evidence of proteorhodopsin; to

confirm this difference between the Biwa and Great Lakes MAGs, we mapped reads from Lake

Biwa (133) (BioProject PRJDB6644) onto the assembled contig described above using bowtie2

(95)**.** This analysis demonstrated that while a large fraction of the NspGL1 contig in question

recruited reads from Lake Biwa at high identity (98-99%), starting upstream of proteorhodopsin

and retinal biosynthesis, this contig no longer recruited reads from Lake Biwa.

<u>Manual identification of key nitrification genes</u>

Despite recovery of 15 high completion MAGs in NspGL1/2a/2b/3, many of these MAGs

lacked key nitrification genes in *amo* and *hao* operons. This was largely due to the fact that *amo*

and *hao* operons were often assembled on small contigs below the minimum size cutoff we

imposed for binning contigs. Difficulty in assembling these contigs was likely in part due to the

several *amo* and *hao* operons with extremely high identity to one another in each genome, a

phenomenon which has been observed in other *Nitrosospira* (18). Manual assembly graph

inspection with Bandage (131) supported this hypothesis, as did assessment of abundance of

short reads associated with *amo* operons from NspGL and comparison of abundance of short

reads associated with core gene *rpoB* from NspGL, using ROCker (134). Still, an exemplar

MAG from at least one representative of each ecotype (NspGL1/2a/2b/3) was found with both

amo and *hao* operons. Further, manual inspection of unbinned contigs confirmed that *amo* and

hao operons existed on contigs in every sample from which a MAG for a particular ecotype was

recovered. That is, for every time that an NspGL1 MAG was recovered from a sample, we were

able to determine that an *amo* and *hao* operon which could be affiliated with NspGL1 existed,

even if it was not correctly binned. Affiliation for these unbinned key nitrification genes was

carried out by alignment of *amoAB* and *haoAB* sequences to *amoAB* and *haoAB* sequences

correctly binned in NspGL ecotypes. This process was also carried out for two NtogaGL1a

MAGs for *nxrAB*, which were poorly assembled in those two samples. Dataset S6 summarizes

the presence of genes related to nitrification and nitrogen metabolism across all our MAGs.

<u>Verification of gene absences</u>

Metagenome-assembled genomes typically comprise tens or even hundreds of contigs,

and this fragmented nature makes it impossible to say with certainty whether a particular gene

is truly absent. To substantiate our claims of gene absence based on MAGs, we used several

lines of evidence. First, we note that our MAGs have high estimated completion (median 96.4%,

mean 94.3%), based on the presence of universal core gene markers. Second, for all new

lineages described here except NspGL3, we assembled multiple similar MAGs independently

from different samples, and we inferred gene absences only if the absence was replicated in

multiple assemblies. Together these two factors provide strong support for cases where a

missing gene would be expected to occur in a region of predominantly core genes; however

these factors are less informative for cases where a missing gene might occur in a genomic

island, because we have no way of assessing the completion of regions lacking core genes, and

islands tend to have systematic poor assemblies across samples. A third line of evidence that

we considered is chromosome organization: if a single gene is deleted from an otherwise

conserved region of synteny, then this deletion should be apparent in a gene neighborhood

diagram (e.g. Figures S5, S7, S8, S10). Unfortunately in many cases, our MAGs are too

dissimilar from reference genomes and share little synteny with them, so this approach is not

always informative.

We used a fourth approach based on quantitative analysis of short reads to verify gene

absences. If a suspected missing gene were actually present in the population, but failed to

assemble and/or bin with the rest of the genome, then it should be detectable in the

unassembled short reads. The frequency of a gene in the population can be estimated from its

abundance in the short reads, compared to the abundance of core marker genes in the short

reads. We implemented this approach as follows. We searched unassembled short reads for

each gene of interest that we identified as absent from MAGs (e.g. nitrosocyanin) using tblastn.

Short reads with significant similarity were then filtered by best-hit taxonomy to the appropriate

nitrifier group (i.e. *Nitrosomonadaceae*, *Ca.* Nitrotoga, *Nitrospira*, *Thaumarchaea*). These filtered

short reads were enumerated and length-normalized (1000 * number of short reads / length of

the target gene of interest). The same procedure was repeated for genes expected to be

present in every cell (e.g. *amoAB*, *hao*, *nxrAB*, ribosomal protein genes) for comparison. If a

putative missing gene (based on MAGs) has near-zero detection in the short reads, we can be

confident that the gene is truly missing (or has undetectable sequence similarity, or was so

recently acquired from another lineage that its best hit points to a different taxon). By contrast, if

a putative missing gene (based on MAGs) does recover short reads, then the gene may be

present in genomes related to our MAGs but was unassembled/unbinned, or the gene may be

present in another lineage of nitrifiers that is not represented by our MAGs. Short read-based

quantification of select genes is presented in Supplemental Dataset S7 and described in

Supplemental Text.

Statistical analysis and plots

All statistical comparisons were carried out in R version 3.5.3 (135) and plots were generated

using ggplot2 3.2.0 (136). Code and data files are available at

bitbucket.org/greatlakes/gl_nitrifiers.

Data Availability

The metagenome-assembled genomes presented here are available via NCBI BioProject

PRJNA636190. 16S rRNA data is available at NCBI BioProject PRJNA591360. Metagenomes

sequenced by JGI are available at http://genome.jgi.doe.gov, Project IDs 1045056, 1045059,

1045062, 1045065, 1045068, 1045071. Single-cell amplified genome is available at

img.jgi.doe.gov, IMG Genome ID 3300033241. Raw reads are available in NCBI SRA

(SRR14240538-SRR14240543) or through JGI with project IDs listed above.

Acknowledgments

The work conducted by the U.S. Department of Energy Joint Genome Institute, a DOE Office of

Science User Facility, is supported under Contract No. DE-AC02-05CH11231. Sequencing

support was provided by the DOE JGI Community Sequencing Program (CSP #1565,

#503460). Computational resources were provided by the UChicago Research Computing

Center. Funding for this work was provided by Illinois-Indiana Sea Grant (Grant #

NA14OAR170095), UChicago Women's Board, and the National Science Foundation (OCE-

1830011 to MLC). We thank the science staff in the Great Lakes National Program Office of the

US EPA, and the captain and crew of the R/V *Lake Guardian*, for facilitating sample collection.

We thank members of the Coleman and Waldbauer labs for assistance with sample collection

and processing, and for discussion and comments on the manuscript.

Competing Interests

The authors declare no competing interests.

Figure & Table Legends

Table 1. Evidence for genome streamlining in nitrifiers from the Laurentian Great Lakes.

Genome features were compared between Great Lakes MAGs and reference genomes using a

two-sided Wilcoxon/Mann-Whitney test. n.s., not significant at 0.05 level. Only genomes with

>70% completion and <10% contamination are included.

Figure 1. Dissolved inorganic nitrogen availability and distribution of nitrifiers across the Great

Lakes. **(a)** Oxidized nitrogen concentrations. Values include NOx concentrations from published

studies (n=128; (14, 33, 35, 137, 138)), US EPA Water Quality Surveys in 2012 and 2013

(n=1626 from GLENDA database), and this study (n=20). **(b)** Ammonium concentrations.

Values are derived from the literature as in (a) (n=118) and from this study (n=20). **(c)**

Distribution of nitrifiers across the Great Lakes. Top panel: map of sampling stations; stars

indicate stations chosen for metagenome analysis. Bottom panel: relative abundance of

ammonia-oxidizing (green) and nitrite-oxidizing (pink) families based on 16S rRNA V4-V5

amplicon sequencing, sampled in the mid-hypolimnion (except western Erie, sampled 1m from

bottom). Data is plotted roughly west to east as indicated on the map.

Figure 2. Genome properties and cross-lake distribution of ammonia oxidizing organisms,

showing both Archaea (top) and Betaproteobacteria (bottom). Rows highlighted in green

represent clusters of genomes reconstructed from the Great Lakes and median values are

shown for genome size, GC content, and coding density. For catalase, "E" indicates

monofunctional catalase *katE*; "G" indicates bifunctional catalase-peroxidase *katG*. For carbon

fixation, RuBisCO type is shown in parentheses (61); HP/HB, 3-Hydroxypropionate/4-

Hydroxybutyrate cycle; CBB, Calvin-Benson-Bassham cycle. Bubble plot shows composition of

ammonia oxidizers in hypolimnion samples, using MAGs as probes to recruit metagenomic

reads (values sum to 100% for each lake column). Genes identified in only a subset of genomes

are shown as (+).

Figure 3. Evidence for proteorhodopsin (PR) in *Nitrosospira* from the Great Lakes. (a) Gene

neighborhood surrounding PR in *Nitrosospira* MAG MC17_S15_bin_110 and SAG 207399.

Genes are colored according to the best BLAST hit taxonomy in the NCBI nr database. (b)

Alignment of predicted *Nitrosospira* PR with reference sequences. Diagnostic features are

highlighted (54, 55): blue boxes, diagnostic residues for PR; pink box, residue indicative of blue

or green tuning; green underline, shorter beta-sheet region in PR. Sequence accession

numbers: bacteriorhodopsin P02945, actinorhodopsin A0A1D9E0H1, sensory rhodopsin

P42196, *Dokdonia* PR EAQ40507.1, SAR86 blue-tuned PR Q4PP54, SAR86 green-tuned PR

Q9F7P4, SAR11 PR A6YQL7. (c) Phylogenetic tree showing close relatives of *Nitrosospira* PR

within Supercluster III, as defined by MicRhoDE database (56). Neighboring clusters have been

collapsed for clarity.

Figure 4. Genome properties and cross-lake distribution of nitrite oxidizing taxa *Nitrospira* (top)

and *Ca.* Nitrotoga (Betaproteobacteria*;* bottom). Rows highlighted in pink represent clusters of

genomes reconstructed from the Great Lakes and median values are shown for genome size,

GC content, and coding density. rTCA, reductive tricarboxylic acid cycle; CBB, Calvin-Benson-

Bassham cycle; ONR, octaheme nitrite reductase. Values in parentheses indicate RuBisCO

type (61). Bubble plot shows composition of NOB per lake based on metagenomic read

mapping. Genes identified in only a subset of genomes are indicated by (+). The asterisk (*)

indicates for *Ca.* Nitrotoga, one *nxrAB* copy was recovered in genome assemblies, but short

read analysis suggests two copies per genome (Supplemental Text).

Figure 5. Distinct photolyase proteins in NspiraGL. **a)** Phylogenetic tree showing families of

photolyases. Three families are found in NspiraGL: CPD class I photolyase, FeS-BCP / CRYPro

family, and an uncharacterized CPF-related family found in diverse Bacteria. CPD Class I

photolyases are also found in other nitrifiers including *Ca.* Nitrotoga NtogaGL1a, *Nitrosospira*

NspGL1, and *Nitrosarchaeum* NarchGL. **b)** Domain structure of the three photolyase families

present in NspiraGL.

Supplemental Material

Supplemental Text. Detailed description of short-read analyses to verify gene absences.

10.6084/m9.figshare.15130350

Figure S1. Relationships between environmental factors and relative abundance of nitrifiers in

the Great Lakes. Relative abundance based on 16S rRNA V4-V5 region amplicon sequencing

from Summer 2012. Nitrifier abundance includes sequences assigned to the families

Nitrosopumilaceae (AOA), *Nitrosomonadaceae* (AOB), *Nitrospiraceae* (NOB), and

Gallionellaceae (NOB). (a) Relative abundance of nitrifiers (% of total community) compared to

photosynthetically active radiation (PAR). (b) Correlation between the relative abundances of

nitrite oxidizers and ammonia oxidizers in each sample. A 1:1 line (solid) and linear regression

fit (dashed) are shown for reference. (c) Relative abundance of nitrifiers (% of total community)

compared to chlorophyll *a* concentration. PAR and chlorophyll data from the EPA Great Lakes

Environmental Database (https://cdx.epa.gov/). Values are presented in Supplemental Dataset

S1.

Figure S2. Oligotype composition of nitrifiers across the Great Lakes. Each color represents a

different 16S rRNA oligotype.

Figure S3. Phylogenomic tree of the *Nitrosomonadales* order of *Betaproteobacteria*, focusing

on the *Nitrosomonadaceae* and *Gallionellaceae* families. Tree was constructed using 139

concatenated single-copy core genes (B.J. Campbell, L. Yu, J.F. Heidelberg, and D.L.

Kirchman, PNAS 108:12776-12781, 2011). Colored squares indicate the environment of origin;

colored circles denote the lake of origin for genomes from the Laurentian Great Lakes; yellow

stars indicate clades containing strains that lack nitrosocyanin.

Figure S4. Histograms of nitrifier genome properties. Rows show each taxonomic group and

columns show genome properties. Each circle represents a single genome/MAG. Great Lakes

MAGs shown as filled circles.

Figure S5. Gene neighborhood surrounding proteorhodopsin in *Nitrosospira* draft genomes

from deep lakes. Light blue, shared genes on reverse strand; dark blue, shared genes on

forward strand; pink, tRNA genes; orange, PR module in NspGL1 MAG; gray, genes unique to

one MAG.

Figure S6. Phylogenomic tree of *Nitrosopumilaceae* and *Cenarchaeum*. Tree was constructed

using concatenated single-copy core genes as defined in (C. Rinke *et al.,* Nature 499:431-437,

2013)*.* Colored boxes correspond to the environment of origin; genomes from the Laurentian

Great Lakes are indicated by colored circles. Names in bold indicate cultured isolates or

enrichments.

Figure S7. Cyanase diversity and distribution among nitrite oxidizing bacteria. a) Phylogenetic

tree showing distinct clades of cyanase, highlighting Great Lakes *Nitrospira* and *Ca.* Nitrotoga.

b) Gene synteny surrounding the cyanase gene in NspiraGL and reference *Nitrospira*, and a

cyanase insertion in NtogaGL compared to reference *Ca.* Nitrotoga.

Figure S8. Gene content distinguishing NtogaGL1a and NtogaGL1b. Genes colored in blue are

shared between both genomes; orange colored genes are unique to either genome.

Figure S9. Phylogenomic tree of Lineage I and II *Nitrospira*. Tree was constructed using

concatenated single-copy core genes (B.J. Campbell, L. Yu, J.F. Heidelberg, and D.L.

Kirchman, PNAS 108:12776-12781, 2011). Colored boxes correspond to the environment of

origin; genomes from the Laurentian Great Lakes are indicated by circles color coded by lake of

origin.

Figure S10. Gene neighborhood surrounding photolyase genes in NspiraGL, compared to

reference *N. moscoviensis*. Genes shared between the two genomes are colored light blue

(protein coding) or magenta (rRNA, tRNA); non-shared genes are gray; photolyase genes are

colored to match the tree in Figure 5.

Dataset S1. Relative abundance of nitrifiers and associated environmental parameters. Relative

abundance based on 16S rRNA amplicons is reported as proportion of total community reads.

The full 16S rRNA dataset and flow cytometry cell counts are presented in Paver et al. (2020)

Environ Microbiol 22(1): 433-446. Chlorophyll a concentrations, irradiance values and dissolved

oxygen measurements are from the US EPA Water Quality Survey (available from

https://cdx.epa.gov/). Relative abundance based on metagenome read recruitment to MAGs is

reported as the proportion of total sequenced reads for that sample.

10.6084/m9.figshare.15130350

Dataset S2. Nitrate (NOx), ammonium, and urea concentrations compiled from literature, EPA

Water Quality Survey (available from https://cdx.epa.gov/) or measured in this study. NA, not

available; n.d., not detected 10.6084/m9.figshare.15130350

Dataset S3. Assembly statistics and taxonomic information for metagenome-assembled

genomes (MAGs) of nitrifiers in the Great Lakes. 10.6084/m9.figshare.15130350

Dataset S4. Genome properties for newly assembled MAGs and reference genomes included

in this study. 10.6084/m9.figshare.15130350

Dataset S5. Protein functions differentiating genomes of Great Lakes nitrifiers from previously

published genomes. Genome averages are calculated for reference genomes and GL MAGs.

Genome-level counts for each individual GL MAG and for selected reference genomes are also

shown. 10.6084/m9.figshare.15130350

Dataset S6. Detailed list of presence/absence of key nitrification genes in MAGs presented in

this study. NCBI Protein IDs listed when a MAG contains the target gene. 0 indicates the gene

was not recovered in the MAG. 10.6084/m9.figshare.15130350

Dataset S7. Short read analysis for verifying gene loss. Each target gene was searched in

unassembled short reads as described in Materials & Methods. Values indicate the number of

mapped reads, normalized to gene length. n.d., not detected. 10.6084/m9.figshare.15130350

Dataset S8. Stations are from the US EPA Water Quality Survey (available from

https://cdx.epa.gov/). 10.6084/m9.figshare.15130350

References

1. Canfield DE, Glazer AN, Falkowski PG. 2010. The evolution and future of Earth's nitrogen

cycle. Science 330:192–196.

2. Pachiadaki MG, Sintes E, Bergauer K, Brown JM, Record NR, Swan BK, Mathyer ME,

Hallam SJ, Lopez-Garcia P, Takaki Y, Nunoura T, Woyke T, Herndl GJ, Stepanauskas R.

2017. Major role of nitrite-oxidizing bacteria in dark ocean carbon fixation. Science

358:1046–1051.

3. Reinthaler T, van Aken HM, Herndl GJ. 2010. Major contribution of autotrophy to microbial

carbon cycling in the deep North Atlantic's interior. Deep Sea Res II 57:1572–1580.

4. Swan BK, Martinez-Garcia M, Preston CM, Sczyrba A, Woyke T, Lamy D, Reinthaler T,

Poulton NJ, Masland EDP, Gomez ML, Sieracki ME, DeLong EF, Herndl GJ,

Stepanauskas R. 2011. Potential for chemolithoautotrophy among ubiquitous bacteria

lineages in the dark ccean. Science 333:1296–1300.

5. Baltar F, Herndl GJ. 2019. Ideas and perspectives: Is dark carbon fixation relevant for

oceanic primary production estimates? Biogeosciences 16:3793–3799.

6. Callieri C, Coci M, Eckert EM, Salcher MM, Bertoni R. 2014. Archaea and Bacteria in deep

lake hypolimnion: in situ dark inorganic carbon uptake. J Limnol 73:31-38.

7. Santoro AE, Richter RA, Dupont CL. 2019. Planktonic marine Archaea. Ann Rev Mar Sci

11:131–158.

8. Schleper C. 2010. Ammonia oxidation: different niches for bacteria and archaea? ISME J

4:1092–1094.

9. Santoro AE, Dupont CL, Richter RA, Craig MT, Carini P, McIlvin MR, Yang Y, Orsi WD,

Moran DM, Saito MA. 2015. Genomic and proteomic characterization of "*Candidatus*

Nitrosopelagicus brevis": An ammonia-oxidizing archaeon from the open ocean. Proc Natl

Acad Sci USA 112:1173–1178.

10. Auguet J-C, Triadó-Margarit X, Nomokonova N, Camarero L, Casamayor EO. 2012.

Vertical segregation and phylogenetic characterization of ammonia-oxidizing Archaea in a

deep oligotrophic lake. ISME J 6:1786–1797.

11. Herber J, Klotz F, Frommeyer B, Weis S, Straile D, Kolar A, Sikorski J, Egert M,

Dannenmann M, Pester M. 2020. A single Thaumarchaeon drives nitrification in deep

oligotrophic Lake Constance. Environ Microbiol 22:212–228.

12. Urbach E, Vergin KL, Young L, Morse A, Larson GL, Giovannoni SJ. 2001. Unusual

bacterioplankton community structure in ultra-oligotrophic Crater Lake. Limnol Oceanogr

46:557–572.

13. Okazaki Y, Fujinaga S, Tanaka A, Kohzu A, Oyagi H, Nakano S. 2017. Ubiquity and

quantitative significance of bacterioplankton lineages inhabiting the oxygenated

hypolimnion of deep freshwater lakes. ISME J 11:2279–2293.

14. Mukherjee M, Ray A, Post AF, McKay RM, Bullerjahn GS. 2016. Identification,

enumeration and diversity of nitrifying planktonic archaea and bacteria in trophic end

members of the Laurentian Great Lakes. J Great Lakes Res 42:39–49.

15. Hugoni M, Etien S, Bourges A, Lepère C, Domaizon I, Mallet C, Bronner G, Debroas D,

Mary I. 2013. Dynamics of ammonia-oxidizing Archaea and Bacteria in contrasted

freshwater ecosystems. Res Microbiol 164:360–370.

16. Hayden CJ, Beman JM. 2014. High abundances of potentially active ammonia-oxidizing

Bacteria and Archaea in oligotrophic, high-altitude lakes of the Sierra Nevada, California,

USA. PLoS ONE 9:e111560.

17. Bollmann A, Bar-Gilissen M-J, Laanbroek HJ. 2002. Growth at low ammonium

concentrations and starvation response as potential factors involved in niche differentiation

among ammonia-oxidizing bacteria. Appl Environ Microbiol 68:4751–4757.

18. Sedlacek CJ, McGowan B, Suwa Y, Sayavedra-Soto L, Laanbroek HJ, Stein LY, Norton

JM, Klotz MG, Bollmann A. 2019. A physiological and genomic comparison of

Nitrosomonas cluster 6a and 7 ammonia-oxidizing bacteria. Microb Ecol 78:985–994.

19.   Alonso-Sáez L, Waller AS, Mende DR, Bakker K, Farnelid H, Yager PL, Lovejoy C,

905   Tremblay J-É, Potvin M, Heinrich F, Estrada M, Riemann L, Bork P, Pedrós-Alió C,

906   Bertilsson S. 2012. Role for urea in nitrification by polar marine Archaea. Proc Natl Acad

907   Sci USA 109:17989–17994.

908   20.   Daims H, Lücker S, Wagner M. 2016. A new perspective on microbes formerly known as

909   nitrite-oxidizing bacteria. Trends Microbiol 24:699–712.

910   21.   Nowka B, Daims H, Spieck E. 2015. Comparison of oxidation kinetics of nitrite-oxidizing

911   bacteria: nitrite availability as a key factor in niche differentiation. Appl Environ Microbiol

912   81:745–753.

913   22.   Wegen S, Nowka B, Spieck E. 2019. Low temperature and neutral pH define "*Candidatus*

914   Nitrotoga sp.*" as a competitive nitrite oxidizer in coculture with *Nitrospira defluvii*. Appl

915   Environ Microbiol 85:e02569-18.

916   23.   Koch H, Galushko A, Albertsen M, Schintlmeister A, Gruber-Dorninger C, Lucker S,

917   Pelletier E, Le Paslier D, Spieck E, Richter A, Nielsen PH, Wagner M, Daims H. 2014.

918   Growth of nitrite-oxidizing bacteria by aerobic hydrogen oxidation. Science 345:1052–

919   1054.

920   24.   Füssel J, Lücker S, Yilmaz P, Nowka B, van Kessel MAHJ, Bourceau P, Hach PF,

921   Littmann S, Berg J, Spieck E, Daims H, Kuypers MMM, Lam P. 2017. Adaptability as the

922   key to success for the ubiquitous marine nitrite oxidizer *Nitrococcus*. Science Adv

923   3:e1700807.

924   25.   Palatinszky M, Herbold C, Jehmlich N, Pogoda M, Han P, von Bergen M, Lagkouvardos I,

925   Karst SM, Galushko A, Koch H, Berry D, Daims H, Wagner M. 2015. Cyanate as an

926   energy source for nitrifiers. Nature 524:105–108.

927   26.   Koch H, Lücker S, Albertsen M, Kitzinger K, Herbold C, Spieck E, Nielsen PH, Wagner M,

928   Daims H. 2015. Expanded metabolic versatility of ubiquitous nitrite-oxidizing bacteria from

929   the genus *Nitrospira*. Proc Natl Acad Sci USA 112:11371–11376.

930   27.   Boddicker AM, Mosier AC. 2018. Genomic profiling of four cultivated *Candidatus* Nitrotoga

931   spp. predicts broad metabolic potential and environmental distribution. ISME J 12:2864–

932   2882.

933   28.   Sterner RW. 2010. In situ-measured primary production in Lake Superior. J Great Lakes

934   Res 36:139–149.

935   29.   Small GE, Bullerjahn GS, Sterner RW, Beall BFN, Brovold S, Finlay JC, McKay RML,

936   Mukherjee M. 2013. Rates and controls of nitrification in a large oligotrophic lake. Limnol

937   Oceanogr 58:276–286.

938   30.   Vollenweider RA, Munawar M, Stadelmann P. 1974. A comparative review of

939   phytoplankton and primary production in the Laurentian Great Lakes. J Fish Res Bd Can

940   31:739–762.

941   31.   Clevinger CC, Heath RT, Bade DL. 2014. Oxygen use by nitrification in the hypolimnion

942   and sediments of Lake Erie. J Great Lakes Res 40:202–207.

943   32.   Neilson MA, Stevens RJJ. 1987. Spatial heterogeneity of nutrients and organic matter in

944   Lake Ontario. Can J Fish Aquat Sci 44:2192–2203.

945   33.   Lean DRS, Knowles R. 1987. Nitrogen transformations in Lake Ontario. Can J Fish Aquat

946   Sci 44:2133–2143.

947   34.   Paver SF, Newton RJ, Coleman ML. 2020. Microbial communities of the Laurentian Great

948   Lakes reflect connectivity and local biogeochemistry. Environ Microbiol 22:433–446.

949   35.   Rozmarynowycz MJ, Beall BFN, Bullerjahn GS, Small GE, Sterner RW, Brovold SS,

950   D'souza NA, Watson SB, McKay RML. 2019. Transitions in microbial communities along a

951   1600 km freshwater trophic gradient. J Great Lakes Res 45:263–276.

952   36.   Fujimoto M, Cavaletto J, Liebig JR, McCarthy A, Vanderploeg HA, Denef VJ. 2016.

953   Spatiotemporal distribution of bacterioplankton functional groups along a freshwater

954   estuary to pelagic gradient in Lake Michigan. J Great Lakes Res 42:1036–1048.

37.   Hooper AB, Terry KR. 1974. Photoinactivation of ammonia oxidation in *Nitrosomonas*. J

Bacteriol 119:899-906.

38. Horrigan SG, Springer AL. 1990. Oceanic and estuarine ammonium oxidation: effects of

light. Limnol Oceanogr 35:479–482.

39. Guerrero M, Jones R. 1996. Photoinhibition of marine nitrifying bacteria. I. Wavelength-

dependent response. Mar Ecol Prog Ser 141:183–192.

40. Merbt SN, Stahl DA, Casamayor EO, Martí E, Nicol GW, Prosser JI. 2012. Differential

photoinhibition of bacterial and archaeal ammonia oxidation. FEMS Microbiol Lett 327:41–

46.

41. Smith JM, Chavez FP, Francis CA. 2014. Ammonium uptake by phytoplankton regulates

nitrification in the sunlit ocean. PLoS ONE 9:e108173.

42. Kumar S, Sterner RW, Finlay JC, Brovold S. 2007. Spatial and temporal variation of

ammonium in Lake Superior. J Great Lakes Res 33:581–591.

43. Dove A, Chapra SC. 2015. Long-term trends of nutrients and trophic response variables

for the Great Lakes. Limnol Oceanogr 60:696–721.

44. Belisle BS, Steffen MM, Pound HL, Watson SB, DeBruyn JM, Bourbonniere RA, Boyer GL,

Wilhelm SW. 2016. Urea in Lake Erie: Organic nutrient sources as potentially important

drivers of phytoplankton biomass. J Great Lakes Res 42:599–607.

45. Rice MC, Norton JM, Valois F, Bollmann A, Bottomley PJ, Klotz MG, Laanbroek HJ, Suwa

Y, Stein LY, Sayavedra-Soto L, Woyke T, Shapiro N, Goodwin LA, Huntemann M, Clum A,

Pillay M, Kyrpides N, Varghese N, Mikhailova N, Markowitz V, Palaniappan K, Ivanova N,

Stamatis D, Reddy TBK, Ngan CY, Daum C. 2016. Complete genome of *Nitrosospira*

briensis C-128, an ammonia-oxidizing bacterium from agricultural soil. Stand Genomic Sci

11:46.

46. Garcia JC, Urakawa H, Le VQ, Stein LY, Klotz MG, Nielsen JL. 2013. Draft genome

sequence of *Nitrosospira* sp. strain APG3, a psychrotolerant ammonia-oxidizing bacterium

isolated from sandy lake sediment. Genome Announcements 1:e00930-13.

47.   Norton JM, Klotz MG, Stein LY, Arp DJ, Bottomley PJ, Chain PSG, Hauser LJ, Land ML,

983   Larimer FW, Shin MW, Starkenburg SR. 2008. Complete genome sequence of

984   *Nitrosospira multiformis*, an ammonia-oxidizing bacterium from the soil environment. Appl

985   Environ Microbiol 74:3559–3572.

986   48.   Giovannoni SJ, Thrash JC, Temperton B. 2014. Implications of streamlining theory for

987   microbial ecology. ISME J 8:1553–1565.

988   49.   Bollmann A, Sedlacek CJ, Norton J, Laanbroek HJ, Suwa Y, Stein LY, Klotz MG, Arp D,

989   Sayavedra-Soto L, Lu M, Bruce D, Detter C, Tapia R, Han J, Woyke T, Lucas SM, Pitluck

990   S, Pennacchio L, Nolan M, Land ML, Huntemann M, Deshpande S, Han C, Chen A,

991   Kyrpides N, Mavromatis K, Markowitz V, Szeto E, Ivanova N, Mikhailova N, Pagani I, Pati

992   A, Peters L, Ovchinnikova G, Goodwin LA. 2013. Complete genome sequence of

993   *Nitrosomonas* sp. Is79, an ammonia oxidizing bacterium adapted to low ammonium

994   concentrations. Stand Genomic Sci 7:469–482.

995   50.   Beja O. 2000. Bacterial Rhodopsin: Evidence for a new type of phototrophy in the sea.

996   Science 289:1902–1906.

997   51.   Béjà O, Spudich EN, Spudich JL, Leclerc M, DeLong EF. 2001. Proteorhodopsin

998   phototrophy in the ocean. Nature 411:786–789.

999   52.   Sabehi G, Loy A, Jung K-H, Partha R, Spudich JL, Isaacson T, Hirschberg J, Wagner M,

1000   Béjà O. 2005. New insights into metabolic properties of marine bacteria encoding

1001   proteorhodopsins. PLoS Biol 3:e273.

1002   53.   Martinez A, Bradley AS, Waldbauer JR, Summons RE, DeLong EF. 2007. Proteorhodopsin

1003   photosystem gene expression enables photophosphorylation in a heterologous host. Proc

1004   Natl Acad Sci USA 104:5590–5595.

1005   54.   Reckel S, Gottstein D, Stehle J, Löhr F, Verhoefen M-K, Takeda M, Silvers R, Kainosho M,

1006   Glaubitz C, Wachtveitl J, Bernhard F, Schwalbe H, Güntert P, Dötsch V. 2011. Solution

1007   NMR structure of proteorhodopsin. Angewandte Chemie International Edition 50:11942–

11946.

55. Kralj JM, Bergo VB, Amsden JJ, Spudich EN, Spudich JL, Rothschild KJ. 2008.

Protonation state of Glu142 differs in the green- and blue-absorbing variants of

proteorhodopsin. Biochemistry 47:3447–3453.

56. Boeuf D, Audic S, Brillet-Guéguen L, Caron C, Jeanthon C. 2015. MicRhoDE: a curated

database for the analysis of microbial rhodopsin diversity and evolution. Database (Oxford)

bav080.

57. Hyman MR, Arp DJ. 1992. 14C2H2- and 14CO2-labeling studies of the de novo synthesis

of polypeptides by *Nitrosomonas europaea* during recovery from acetylene and light

inactivation of ammonia monooxygenase. J Biol Chem 267:1534–1545.

58. Yousef F, Shuchman R, Sayers M, Fahnenstiel G, Henareh A. 2017. Water clarity of the

Upper Great Lakes: Tracking changes between 1998–2012. J Great Lakes Res 43:239–

247.

59. Beaumont HJE, Hommes NG, Sayavedra-Soto LA, Arp DJ, Arciero DM, Hooper AB,

Westerhoff HV, van Spanning RJM. 2002. Nitrite reductase of *Nitrosomonas europaea* is

not essential for production of gaseous nitrogen oxides and confers tolerance to nitrite. J

Bacteriol 184:2557–2560.

60. Watzer B, Forchhammer K. 2018. Cyanophycin synthesis optimizes nitrogen utilization in

the unicellular cyanobacterium *Synechocystis* sp. strain PCC 6803. Appl Environ Microbiol

84:e01298-18.

61. Badger MR, Bek EJ. 2008. Multiple Rubisco forms in proteobacteria: their functional

significance in relation to CO2 acquisition by the CBB cycle. J Exp Bot 59:1525–1541.

62. Rae BD, Long BM, Badger MR, Price GD. 2013. Functions, compositions, and evolution of

the two types of carboxysomes: polyhedral microcompartments that facilitate CO2 fixation

in cyanobacteria and some proteobacteria. Microbiol Mol Biol Rev 77:357–379.

63. Cabello-Yeves PJ, Zemskaya TI, Rosselli R, Coutinho FH, Zakharenko AS, Blinov VV,

Rodriguez-Valera F. 2018. Genomes of novel microbial lineages assembled from the sub-

ice waters of Lake Baikal. Appl Environ Microbiol 84:e02132-17.

64. Sterner RW, Smutka TM, McKay RML, Xiaoming Q, Brown ET, Sherrell RM. 2004.

Phosphorus and trace metal limitation of algae and bacteria in Lake Superior. Limnol

Oceanogr 49:495–507.

65. Ryals J, Hsu RY, Lipsett MN, Bremer H. 1982. Isolation of single-site Escherichia coli

mutants deficient in thiamine and 4-thiouridine syntheses: identification of a nuvC mutant. J

Bacteriol 151:899–904.

66. Kitzinger K, Koch H, Lücker S, Sedlacek CJ, Herbold C, Schwarz J, Daebeler A, Mueller

AJ, Lukumbuzya M, Romano S, Leisch N, Karst SM, Kirkegaard R, Albertsen M, Nielsen

PH, Wagner M, Daims H. 2018. Characterization of the first "*Candidatus* Nitrotoga" isolate

reveals metabolic versatility and separate evolution of widespread nitrite-oxidizing bacteria.

mBio 9:e01186-18.

67. Ishii K, Fujitani H, Sekiguchi Y, Tsuneda S. 2020. Physiological and genomic

characterization of a new "*Candidatus* Nitrotoga" isolate. Environ Microbiol 22:2365–2382.

68. Starkenburg SR, Chain PSG, Sayavedra-Soto LA, Hauser L, Land ML, Larimer FW,

Malfatti SA, Klotz MG, Bottomley PJ, Arp DJ, Hickey WJ. 2006. Genome sequence of the

chemolithoautotrophic nitrite-oxidizing bacterium *Nitrobacter winogradskyi* Nb-255. Appl

Environ Microbiol 72:2050–2063.

69. Lücker S, Wagner M, Maixner F, Pelletier E, Koch H, Vacherie B, Rattei T, Damste JSS,

Spieck E, Le Paslier D, Daims H. 2010. A *Nitrospira* metagenome illuminates the

physiology and evolution of globally important nitrite-oxidizing bacteria. Proc Natl Acad Sci

USA 107:13479–13484.

70. Lücker S, Nowka B, Rattei T, Spieck E, Daims H. 2013. The genome of *Nitrospina gracilis*

illuminates the metabolism and evolution of the major marine nitrite oxidizer. Front

Microbiol 4:27.

71.   Kitzinger K, Marchant HK, Bristow LA, Herbold CW, Padilla CC, Kidane AT, Littmann S,

Daims H, Pjevac P, Stewart FJ, Wagner M, Kuypers MMM. 2020. Single cell analyses

reveal contrasting life strategies of the two main nitrifiers in the ocean. Nat Commun

11:767.

72.   Mundinger AB, Lawson CE, Jetten MSM, Koch H, Lücker S. 2019. Cultivation and

transcriptional analysis of a canonical *Nitrospira* under stable growth conditions. Front

Microbiol 10:1325.

73.   Sancar A. 2003. Structure and function of DNA photolyase and cryptochrome blue-light

photoreceptors. Chem Rev 103:2203–2238.

74.   Zadow A von, Ignatz E, Pokorny R, Essen L-O, Klug G. 2016. *Rhodobacter sphaeroides*

CryB is a bacterial cryptochrome with (6–4) photolyase activity. FEBS J 283:4291–4309.

75.   Zhang F, Scheerer P, Oberpichler I, Lamparter T, Krauß N. 2013. Crystal structure of a

prokaryotic (6-4) photolyase with an Fe-S cluster and a 6,7-dimethyl-8-ribityllumazine

antenna chromophore. Proc Natl Acad Sci USA 110:7217–7222.

76.   Maresca JA, Keffer JL, Hempel PP, Polson SW, Shevchenko O, Bhavsar J, Powell D,

Miller KJ, Singh A, Hahn MW. 2019. Light modulates the physiology of nonphototrophic

Actinobacteria. J Bacteriol 201:e00740-18.

77.   Wink DA, Kasprzak KS, Maragos CM, Elespuru RK, Misra M, Dunams TM, Cebula TA,

Koch WH, Andrews AW, Allen JS, Et A. 1991. DNA deaminating ability and genotoxicity of

nitric oxide and its progenitors. Science 254:1001–1003.

78.   Imlay JA. 2013. The molecular mechanisms and physiological consequences of oxidative

stress: lessons from a model bacterium. Nat Rev Microbiol 11:443–454.

79.   Zhang T, Hansel CM, Voelker BM, Lamborg CH. 2016. Extensive dark biological

production of reactive oxygen species in brackish and freshwater ponds. Environ Sci

Technol 50:2983–2993.

80.   Khademian M, Imlay JA. 2017. *Escherichia coli* cytochrome *c* peroxidase is a respiratory

oxidase that enables the use of hydrogen peroxide as a terminal electron acceptor. Proc

Natl Acad Sci USA 114:E6922–E6931.

81. O'Reilly CM, Sharma S, Gray DK, Hampton SE, Read JS, Rowley RJ, Schneider P,

Lenters JD, McIntyre PB, Kraemer BM, Weyhenmeyer GA, Straile D, Dong B, Adrian R,

Allan MG, Anneville O, Arvola L, Austin J, Bailey JL, Baron JS, Brookes JD, Eyto E de,

Dokulil MT, Hamilton DP, Havens K, Hetherington AL, Higgins SN, Hook S, Izmest'eva LR,

Joehnk KD, Kangur K, Kasprzak P, Kumagai M, Kuusisto E, Leshkevich G, Livingstone

DM, MacIntyre S, May L, Melack JM, Mueller☐Navarra DC, Naumenko M, Noges P,

Noges T, North RP, Plisnier P-D, Rigosi A, Rimmer A, Rogora M, Rudstam LG, Rusak JA,

Salmaso N, Samal NR, Schindler DE, Schladow SG, Schmid M, Schmidt SR, Silow E,

Soylu ME, Teubner K, Verburg P, Voutilainen A, Watkinson A, Williamson CE, Zhang G.

2015. Rapid and highly variable warming of lake surface waters around the globe.

Geophys Res Lett 42:10773-10781.

82. Sterner RW, Keeler B, Polasky S, Poudel R, Rhude K, Rogers M. 2020. Ecosystem

services of Earth's largest freshwater lakes. Ecosystem Services 41:101046.

83. Barbiero RP, Lesht BM, Hinchey EK, Nettesheim TG. 2018. A brief history of the U.S. EPA

Great Lakes National Program Office's water quality survey. J Great Lakes Res 44:539–

546.

84. US Environmental Protection Agency. 2003. Sampling and analytical procedures for

GLNPO's open lake water quality survey of the Great Lakes. Great Lakes National

Program Office Illinois, USA.

85. Holmes RM, Aminot A, Kérouel R, Hooker BA, Peterson BJ. 1999. A simple and precise

method for measuring ammonium in marine and freshwater ecosystems. Can J Fish Aquat

Sci 56:1801–1808.

86. Miranda KM, Espey MG, Wink DA. 2001. A rapid, simple spectrophotometric method for

simultaneous detection of nitrate and nitrite. Nitric Oxide 5:62–71.

87. Revilla M, Alexander J, Glibert PM. 2005. Urea analysis in coastal waters: comparison of

enzymatic and direct methods. Limnol Oceanogr Methods 3:290–299.

88. Parada AE, Needham DM, Fuhrman JA. 2016. Every base matters: assessing small

subunit rRNA primers for marine microbiomes with mock communities, time series and

global field samples. Environ Microbiol 18:1403–1414.

89. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO.

2013. The SILVA ribosomal RNA gene database project: improved data processing and

web-based tools. Nucleic Acids Res 41:D590–D596.

90. Wang Q, Garrity GM, Tiedje JM, Cole JR. 2007. Naïve Bayesian classifier for rapid

assignment of rRNA sequences into the new bacterial taxonomy. Appl Environ Microbiol

73:5261–5267.

91. Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. 2013. Development of a

dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence

data on the MiSeq Illumina sequencing platform. Appl Environ Microbiol 79:5112–5120.

92. Eren AM, Maignien L, Sul WJ, Murphy LG, Grim SL, Morrison HG, Sogin ML. 2013.

Oligotyping: differentiating between closely related microbial taxa using 16S rRNA gene

data. Methods Ecol Evol 4:1111–1119.

93. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,

Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G,

Alekseyev MA, Pevzner PA. 2012. SPAdes: A new genome assembly algorithm and its

applications to single-cell sequencing. J Comput Biol 19:455–477.

94. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. 2017. metaSPAdes: a new versatile

metagenomic assembler. Genome Res 27:824–834.

95. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods

9:357–359.

96. Kang DD, Froula J, Egan R, Wang Z. 2015. MetaBAT, an efficient tool for accurately

reconstructing single genomes from complex microbial communities. PeerJ 3:e1165.

97. Graham ED, Heidelberg JF, Tully BJ. 2017. BinSanity: unsupervised clustering of

environmental microbial assemblies using coverage and affinity propagation. PeerJ

5:e3035.

98. Alneberg J, Bjarnason BS, de Bruijn I, Schirmer M, Quick J, Ijaz UZ, Lahti L, Loman NJ,

Andersson AF, Quince C. 2014. Binning metagenomic contigs by coverage and

composition. Nat Methods 11:1144–1146.

99. Sieber CMK, Probst AJ, Sharrar A, Thomas BC, Hess M, Tringe SG, Banfield JF. 2018.

Recovery of genomes from metagenomes via a dereplication, aggregation and scoring

strategy. Nat Microbiol 3:836–843.

100. Eren AM, Esen ÖC, Quince C, Vineis JH, Morrison HG, Sogin ML, Delmont TO. 2015.

Anvi'o: an advanced analysis and visualization platform for 'omics data. PeerJ 3:e1319.

101. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM:

assessing the quality of microbial genomes recovered from isolates, single cells, and

metagenomes. Genome Res 25:1043–1055.

102. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.

BLAST+: architecture and applications. BMC Bioinform 10:421.

103. Campbell BJ, Yu L, Heidelberg JF, Kirchman DL. 2011. Activity of abundant and rare

bacteria in a coastal ocean. Proc Natl Acad Sci USA 108:12776–12781.

104. Rinke C, Schwientek P, Sczyrba A, Ivanova NN, Anderson IJ, Cheng J-F, Darling A,

Malfatti S, Swan BK, Gies EA, Dodsworth JA, Hedlund BP, Tsiamis G, Sievert SM, Liu W-

T, Eisen JA, Hallam SJ, Kyrpides NC, Stepanauskas R, Rubin EM, Hugenholtz P, Woyke

T. 2013. Insights into the phylogeny and coding potential of microbial dark matter. Nature

499:431–437.

105. Mistry J, Finn RD, Eddy SR, Bateman A, Punta M. 2013. Challenges in homology search:

HMMER3 and convergent evolution of coiled-coil regions. Nucleic Acids Res 41:e121–

e121.

106. O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, Rajput B, Robbertse

B, Smith-White B, Ako-Adjei D, Astashyn A, Badretdin A, Bao Y, Blinkova O, Brover V,

Chetvernin V, Choi J, Cox E, Ermolaeva O, Farrell CM, Goldfarb T, Gupta T, Haft D,

Hatcher E, Hlavina W, Joardar VS, Kodali VK, Li W, Maglott D, Masterson P, McGarvey

KM, Murphy MR, O'Neill K, Pujar S, Rangwala SH, Rausch D, Riddick LD, Schoch C,

Shkeda A, Storz SS, Sun H, Thibaud-Nissen F, Tolstoy I, Tully RE, Vatsan AR, Wallin C,

Webb D, Wu W, Landrum MJ, Kimchi A, Tatusova T, DiCuccio M, Kitts P, Murphy TD,

Pruitt KD. 2016. Reference sequence (RefSeq) database at NCBI: current status,

taxonomic expansion, and functional annotation. Nucleic Acids Res 44:D733–D745.

107. Ondov BD, Bergman NH, Phillippy AM. 2011. Interactive metagenomic visualization in a

Web browser. BMC Bioinform 12.

108. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. 2020. GTDB-Tk: a toolkit to classify

genomes with the Genome Taxonomy Database. Bioinformatics 36:1925-1927.

109. Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. 2018. High throughput

ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. Nat Commun

9:1–8.

110. Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal:

prokaryotic gene recognition and translation initiation site identification. BMC Bioinform 11.

111. Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J,

Mitchell A, Nuka G, Pesseat S, Quinn AF, Sangrador-Vegas A, Scheremetjew M, Yong S-

Y, Lopez R, Hunter S. 2014. InterProScan 5: genome-scale protein function classification.

Bioinformatics 30:1236–1240.

112. Kanehisa M, Sato Y, Morishima K. 2016. BlastKOALA and GhostKOALA: KEGG tools for

functional characterization of genome and metagenome sequences. J Mol Biol 428:726–

731.

113. Huerta-Cepas J, Forslund K, Coelho LP, Szklarczyk D, Jensen LJ, von Mering C, Bork P.

2017. Fast genome-wide functional annotation through orthology assignment by eggNOG-

Mapper. Mol Biol Evol 34:2115–2122.

114. Delmont TO, Eren AM. 2018. Linking pangenomes and metagenomes: the

Prochlorococcus metapangenome. PeerJ 6:e4320.

115. Benedict MN, Henriksen JR, Metcalf WW, Whitaker RJ, Price ND. 2014. ITEP: An

integrated toolkit for exploration of microbial pan-genomes. BMC Genomics 15:8.

116. van Dongen S, Abreu-Goodger C. 2012. Using MCL to extract clusters from networks.

Methods Mol Biol 804:281–295.

117. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze

A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI prokaryotic genome annotation pipeline.

Nucleic Acids Res 44:6614–6624.

118. Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic

features. Bioinformatics 26:841–842.

119. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068–

2069.

120. Guy L, Roat Kultima J, Andersson SGE. 2010. genoPlotR: comparative gene and genome

visualization in R. Bioinformatics 26:2334–2335.

121. Matsen FA, Kodner RB, Armbrust EV. 2010. pplacer: linear time maximum-likelihood and

Bayesian phylogenetic placement of sequences onto a fixed reference tree. BMC

Bioinform 11:538.

122. Katoh K, Misawa K, Kuma K, Miyata T. 2002. MAFFT: a novel method for rapid multiple

sequence alignment based on fast Fourier transform. Nucleic Acids Res 30:3059–3066.

123. Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of

large phylogenies. Bioinformatics 30:1312–1313.

124. Letunic I, Bork P. 2016. Interactive tree of life (iTOL) v3: an online tool for the display and

annotation of phylogenetic and other trees. Nucleic Acids Res 44:W242–W245.

125. Spang A, Poehlein A, Offre P, Zumbrägel S, Haider S, Rychlik N, Nowka B, Schmeisser C,

Lebedeva EV, Rattei T, Böhm C, Schmid M, Galushko A, Hatzenpichler R, Weinmaier T,

Daniel R, Schleper C, Spieck E, Streit W, Wagner M. 2012. The genome of the ammonia-

oxidizing *Candidatus* Nitrososphaera gargensis: insights into metabolic versatility and

environmental adaptations. Environ Microbiol 14:3122–3145.

126. Vechtomova YuL, Telegina TA, Kritsky MS. 2020. Evolution of proteins of the DNA

photolyase/cryptochrome family. Biochemistry Moscow 85:131–153.

127. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A,

Lanfear R. 2020. IQ-TREE 2: New models and efficient methods for phylogenetic inference

in the genomic era. Mol Biol Evol 37:1530–1534.

128. Benson DA, Karsch-Mizrachi I, Lipman DJ, Ostell J, Wheeler DL. 2005. GenBank. Nucleic

Acids Res 33:D34–D38.

129. Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Res 32:1792–1797.

130. Castresana J. 2000. Selection of conserved blocks from multiple alignments for their use in

phylogenetic analysis. Mol Biol Evol 17:540–552.

131. Wick RR, Schultz MB, Zobel J, Holt KE. 2015. Bandage: interactive visualization of de

novo genome assemblies. Bioinformatics 31:3350–3352.

132. Thorvaldsdóttir H, Robinson JT, Mesirov JP. 2013. Integrative Genomics Viewer (IGV):

high-performance genomics data visualization and exploration. Brief Bioinform 14:178–

192.

133. Okazaki Y, Nishimura Y, Yoshida T, Ogata H, Nakano S-I. 2019. Genome-resolved viral

and cellular metagenomes revealed potential key virus-host interactions in a deep

freshwater lake. Environ Microbiol 21:4740-4754.

134. Orellana LH, Rodriguez-R LM, Konstantinidis KT. 2017. ROCker: accurate detection and

quantification of target genes in short-read metagenomic data sets by modeling sliding-

window bitscores. Nucleic Acids Res 45:e14.

135. Team RC. 2018. R: A language and environment for statistical computing. R Foundation

for Statistical Computing V, Austria. 2016.

136. Wickham H. 2009. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag, New

York.

137. Murphy TP. 1980. Ammonia and nitrate uptake in the lower Great Lakes. Can J Fish Aquat

Sci 37:1365–1372.

138. Gardner WS, Lavrentyev PJ, Cavaletto JF, McCarthy MJ, Eadie BJ, Johengen TH, Cotner

JB. 2004. Distribution and dynamics of nitrogen and microbial plankton in southern Lake

Michigan during spring transition 1999-2000. J Geophys Res: Oceans 109:C03007.

1253

Figure 1

Figure 2

	% GC	Genome size (Mb)	Coding density (%)	Catalase	Photophysiology Photolyase	Proteorhodopsin	Retinal synthesis	*nirK*	*norCBQD*	Cytochrome P460	Nitrosocyanin	Urease	Carbon fixation pathway
Thaumarchaea													
NarchGL *(n=3)*	31.4	1.24	90	-	+	-	-	+	-	-	-	+	HP/HB
Nitrosarchaeum limnia SFB1	32	1.7	89	-	+	-	-	+	-	-	-	-	HP/HB
Nitrosopumilus maritimus SCM1	34	1.6	92	-	-	-	-	+	-	-	-	-	HP/HB
Nitrosopelagicus brevis CN25	33	1.2	94	-	+	-	-	+	-	-	-	-	HP/HB
Betaproteo.													
NspGL1 *(n=7)*	40.7	1.42	89	G	+	+	+	+	-	-	-	+	CBB (IAc)
NspGL3 *(n=1)*	50.3	1.66	90	-	-	-	-	-	-	+	-	-	CBB (IAq)
NspGL2a *(n=2)*	50.1	1.57	90	-	-	-	-	-	-	+	-	-	CBB (IC)
NspGL2b *(n=5)*	50.0	1.53	89	-	-	-	-	(+)	-	+	-	-	CBB (IC)
Nitrosospira briensis C-128	54	3.2	83	E	-	-	-	+	-	-	+	+	CBB (IC)
Nitrosospira multiformis ATCC 25196	54	3.2	86	E	-	-	-	+	+	-	+	+	CBB (IC)
Nitrosomonas ureae Nm10	44	3.3	86	E	+	-	-	+	-	+	+	+	CBB (IC/IAq)
Nitrosomonas europaea ATCC 19718	50	3.8	88	EG	-	-	-	+	+	+	+	-	CBB (IAq)

N physiology

Distribution across lakes

Relative abundance
25 50 75

Figure 3

Tree scale: 0.1

Figure 4

| | | % GC | Genome size (Mb) | Coding density (%) | # Sigma factors | Photolyase | Catalase | Formate dehydrogenase | NiFe hydrogenase | Thiosulfate dehydrogenase (*tsdA*) | N physiology | | | | | Carbon fixation pathway | Distribution across lakes |
											Assim. nitrite reduction	*nirK*	# *nxrAB* copies	Cyanase	Urease		
Nitrospira	NspiraGL *(n=6)*	57.8	1.83	89	5	+++	G	-	-	-	*nirA*	-	1	+	(+)	rTCA	
	Nitrospira japonica NJ1	58	4.1	90	27	+	E	-	-	-	ONR	+	3	+	+	rTCA	
	Nitrospira moscoviensis NSP M-1	61	4.5	89	18	+	EG	+	+	+	ONR	+	5	+	+	rTCA	
	Nitrospira defluvii	59	4.3	89	17	+	-	+	-	+	*nirA*	+	2	+	-	rTCA	
Betaproteo.	NtogaGL1a *(n=4)*	51.7	1.42	91	4	(+)	G	-	-	-	-	-	1*	+	-	CBB (IC)	
	NtogaGL1b *(n=2)*	51.8	1.46	91	4	-	G	-	-	+	-	-	1*	+	-	CBB (IC)	
	Ca. Nitrotoga CP45	49	2.8	85	8	+	EG	-	+	-	*nirBD*	+	1	-	-	CBB (IC, ID/IA)	
	Ca. Nitrotoga KNB	50	2.6	85	6	+	G	-	+	-	*nirBD*	+	2	-	-	CBB (IC, ID/IA)	

Relative abundance
25 50 75

Figure 5

a

b

Table 1. Evidence for genome streamlining in nitrifiers from the Laurentian Great Lakes. W, two-sided Wilcoxon/Mann-Whitney test statistic. n.s., not significant at 0.05 level.

taxonomic group	genome feature	median (reference)	median (GL)	W	p value	n (ref)	n (GL)
Nitrosomonadaceae	coding density	0.856	0.892	34	5.5E-09	86	15
	est complete size	3210560	1450843	1285	1.0E-09	86	15
	median intergenic	114	80	1254	6.3E-09	86	15
	paralogs	123	29	1189	2.1E-07	86	15
	pseudogenes	101	11	1215	9.0E-10	81	15
	sigma factors	8	4	1275	1.2E-09	86	15
Nitrospira	coding density	0.876	0.894	64	0.0074	64	6
	est complete size	3790956	1828031	373	1.5E-04	64	6
	median intergenic	90	78	299	0.026	64	6
	paralogs	212	49	376	1.2E-04	64	6
	pseudogenes	69	9	182	2.9E-04	31	6
	sigma factors	13	5	379	8.3E-05	64	6
Ca. Nitrotoga	coding density	0.857	0.910	0	0.0080	5	6
	est complete size	2858108	1441179	30	0.0081	5	6
	median intergenic	122	72	30	0.0080	5	6
	paralogs	93	23	30	0.0081	5	6
	pseudogenes	18	8	6	n.s.	1	6
	sigma factors	8	4	30	0.0054	5	6
Thaumarchaeota (Nitrosopumilaceae)	coding density	0.900	0.898	102	n.s.	62	3
	est complete size	1398741	1242579	153	n.s.	62	3
	median intergenic	61	66	60	n.s.	62	3
	paralogs	85	38	175	0.011	62	3
	pseudogenes	22	13	82	n.s.	34	3