1

Large lakes harbor streamlined free-living nitrifiers

Justin C. Podowski[a], Sara F. Paver[a], Ryan J. Newton[b], Maureen L. Coleman[a#]

[a] *Department of the Geophysical Sciences, University of Chicago, Chicago, Illinois, USA*

[b] *School of Freshwater Sciences, University of Wisconsin Milwaukee, Milwaukee, Wisconsin, USA*

Corresponding author:

Maureen Coleman

University of Chicago

Department of the Geophysical Sciences

Hinds Laboratory

5734 S. Ellis Ave.

Chicago, IL 60637

Telephone: (773) 702-8352

Fax: (773) 702-0207

E-mail: mlcoleman@uchicago.edu

Running title: Nitrifiers across the Laurentian Great Lakes

Abstract

Microbial nitrification is a critical process governing nitrogen availability in aquatic systems and provides energy for aphotic carbon fixation. Nitrifiers are phylogenetically and physiologically diverse, but the factors controlling this diversity, and its biogeochemical implications, are poorly understood, particularly in freshwater systems. Here, we characterized the taxonomic, genomic, and metabolic diversity of free-living nitrifiers across Earth's largest freshwater system, the Laurentian Great Lakes. Surprisingly, ammonia oxidizing Bacteria (AOB) related to *Nitrosospira* dominated over ammonia oxidizing Archaea (AOA) at most stations, even in oligotrophic lakes with extremely low substrate concentrations. Nitrite oxidizing Bacteria (NOB) *Ca.* Nitrotoga and *Nitrospira* each dominated a subset of stations, even switching within a lake. Among four ecotypes of *Nitrosospira* we identified, one encodes proteorhodopsin, which could enhance survival of this AOB in transparent, low nutrient lakes. Genome reconstructions for free-living AOA, AOB, and NOB indicate small genomes across all taxa and diverse adaptations to sunlight and oxidative stress. Our findings expand the known functional diversity of nitrifiers and provide the first integrated picture of freshwater nitrifier genomics and ecology. As lakes worldwide undergo rapid environmental change, understanding the relationship between microbial biodiversity and nitrogen cycling is essential for predicting future ecosystem health and services.

Introduction

The oxidation of ammonia to nitrate powers the growth of nitrifying microorganisms and represents a critical flux in the global nitrogen cycle. In aquatic systems, microbial nitrification of ammonia released from sinking organic matter produces nitrate, which can then be removed from the system by denitrification (1). As chemolithoautotrophs, nitrifiers are also a major source of dark carbon fixation (2), which may contribute significant organic carbon to the microbial food web of the ocean's interior (3–5) and of deep freshwater lakes (6).

Microbial nitrifiers encompass wide diversity in phylogeny, physiology, and ecology, and understanding the drivers and consequences of this diversity across ecosystems is a fundamental challenge. Ammonia-oxidizing Archaea (AOA) in the phylum *Thaumarchaeota* dominate the mesopelagic oceans (7,8), likely due to their high affinity for ammonia (9,10) and streamlined genomes (11). In freshwater systems, AOA have been found to dominate oligotrophic lakes, while ammonia-oxidizing Bacteria (AOB) affiliated with the *Nitrosomonadaceae* (Betaproteobacteria) dominate more eutrophic systems (12–17). Complicating this picture, however, there is considerable genomic and physiological variation within each group. Some lineages, such as the thaumarchaeal *Nitrosopelagicus* (11) and bacterial *Nitrosomonas* cluster 6a (18,19), seem to be more specialized for nutrient scarcity than other related lineages. Within the AOA, there are also distinct ecotypes that appear to segregate with depth in the water column, in both marine (8,20) and freshwater systems (12). In freshwaters especially – which are poorly characterized compared to the oceans – it remains difficult to predict which ammonia-oxidizing taxa are likely to dominate in a given system.

For aquatic nitrite oxidizers, which span the phyla *Nitrospira, Nitrospinae,* and *Proteobacteria,* niche differentiation is even less clear. The oceans are dominated by exclusively marine lineages (2,21,22), consistent with ancient salinity-driven divergence. Cultivated strains of NOB show variation in substrate affinity and physiology (21,23,24), but the phylogenetic conservation of these traits, and their influence on environmental distributions, are

poorly understood. Moreover, recent studies have discovered that NOB are capable of

alternative energy metabolisms (25,26) and can access nitrogen from cyanate and urea (27–

29), expanding their ecological potential. In freshwater systems, the NOB *Ca.* Nitrotoga

(Betaproteobacteria) was only recently discovered to be widespread (30), and the diversity of

this genus and factors favoring its success are unknown.

Here, we use the Laurentian Great Lakes as a model system to examine niche

partitioning among planktonic freshwater nitrifiers. The Great Lakes hold 20% of Earth's surface

freshwater and more than half of this volume receives little to no light (< 1% surface irradiance).

This system, while hydrologically connected, spans strong trophic and chemical gradients:

ultraoligotrophic Lake Superior supports rates of primary production and nitrification comparable

to the ocean gyres (31,32), while Lake Erie supports greater production (33) and more than 70-

fold higher nitrification rate (34). Between these extremes, Lake Ontario has low ambient

ammonium concentrations like Lake Superior (35) but nitrification rates up to four times higher

(36). While previous studies reported that AOA and AOB dominate Lakes Superior and Erie,

respectively (16,32), recent community profiling has revealed broader diversity in both

ammonia-oxidizing and nitrite-oxidizing lineages (37–39). We sought to link taxonomic,

genomic, and metabolic diversity of nitrifiers with the varied biogeochemistry of the Great Lakes,

using genome reconstructions and abundance profiling. Our results uncover novel lineages and

metabolic capabilities and provide the first large-scale assessment of freshwater nitrifier

genomics.

Materials and Methods

Sample collection

Water samples were collected from the Laurentian Great Lakes aboard the R/V *Lake*

Guardian, during the biannual Water Quality Surveys conducted by the U.S. EPA Great Lakes

National Program Office (40). Data presented here were collected in April and August 2012.

Samples were collected using a CTD rosette sampler (Sea-Bird Scientific) at the surface (2 m),

deep chlorophyll maximum (if present), the mid-hypolimnion, and near the bottom of the water

column (10 m above the lake bottom at most stations, 1 m above bottom at shallow stations).

For each sample, 5-8L of water was pre-filtered through a GF/A glass fiber filter (Whatman

1820-047; nominal pore size 1.6 µm) to exclude eukaryotic phytoplankton and particle

associated microbes, and cells were collected on 0.22 µm Sterivex filters (Millipore

SVGP01050). Filters were stored at -80°C. For dissolved nutrient analysis, 0.22 µm filtrate was

collected in 125 ml acid-clean HDPE bottles (Nalgene) and stored at -20°C. Samples for single

cell amplified genomes (SAG) were collected in August 2014. For each sample, 1 ml of raw

water was incubated with 100 µl of glycerol-TE buffer (20 ml 100X TE pH 8 + 100 ml glycerol +

60 ml water; final concentration after sample addition is 10mM Tris, 1mM EDTA, 5% glycerol)

for 10 minutes in the dark, then flash frozen in liquid nitrogen and stored at -80°C until

processing.

Physicochemical data

CTD profiles, water chemistry and chlorophyll-*a* data were collected by the U.S. EPA

according to standard protocols (41) and retrieved from the Great Lakes Environmental

Database (https://cdx.epa.gov/) for 2012 and 2013. In addition, we measured dissolved nitrogen

species from August 2013 samples. Ammonium concentrations were measured using the OPA

method in a 96-well plate (42). Nitrate and nitrite concentrations were measured using the

Greiss reaction method in a 96-well plate (43).

16S rRNA analysis

The full 16S rRNA amplicon dataset was described by Paver and colleagues (37). Here

we focus on data from the V4-V5 region (primers 515F-Y, 926R; ref. 44), collected in 2012 in

tandem with metagenome samples from select stations. We classified sequences using the

Silva v. 132 database (45) and the wang method (46) as implemented by mothur (47).

Sequences classified to each detected family of nitrifiers (ammonia oxidizer families

Nitrosomonadaceae and *Nitrosopumilaceae*; nitrite oxidizer families *Gallionellaceae* and

Nitrospiraceae) with a mothur-assigned confidence score above 90 were delineated into

taxonomic units using minimum entropy decomposition with a minimum substantive abundance

of 10 (48).

Metagenome and single-cell genome analysis

One station per lake in Superior, Michigan, Huron, and Ontario, and two stations in Erie,

were selected for metagenome sequencing. DNA was extracted using a modified

phenol:chloroform extraction protocol (37) and libraries prepared according to the Illumina

TruSeq protocol. Samples from spring 2012 were sequenced at the Joint Genome Institute

using Illumina HiSeq (2x150bp). Samples from summer 2012 were sequenced at the University

of Chicago Functional Genomics Core Facility using Illumina HiSeq 2500 (2x250bp). Separate

assemblies of quality filtered reads were carried out for each metagenome using metaSPAdes

3.1.11 (49). Detailed QC, assembly, contig binning, and annotation are described in

Supplementary Methods.

To confirm the presence of proteorhodopsin, we analyzed a single cell amplified genome

from *Nitrosospira* collected from Lake Michigan and sequenced by the Joint Genome Institute.

Quality filtered reads were downloaded from JGI IMG/ER and normalized using bbnorm.sh with

target=100 and mindepth=2. Normalized reads were assembled using SPAdes in single cell

mode (50) with flags --sc and --careful. Resulting scaffolds were annotated identically to MAGs

as described in Supplementary Methods.

Phylogenomic trees and individual gene trees for proteorhodopsin, cyanase, and

photolyase were constructed as described in Supplementary Methods.

Results

Niche partitioning of nitrifiers across the Great Lakes

To map free-living (here defined as cells that pass through a 1.6μm filter) nitrifiers across

the Great Lakes, we searched our recent 16S rRNA datasets for known nitrifying taxa (37). We

detected putative AOB in the genus *Nitrosospira* (Betaproteobacteria) and AOA similar to

Nitrosoarchaeum, along with putative NOB in the genera *Ca.* Nitrotoga (Betaproteobacteria) and

Nitrospira (Nitrospirae). We did not detect 16S rRNA amplicons from *Nitrosococcus*,

Nitrococcus, *Nitrospina* or *Nitrobacter*. The relative abundance of nitrifiers was negatively

correlated with photosynthetically active radiation (PAR; Spearman's rho = -0.89, $p < 2.2e{-}16$)

and reached a maximum below the depth of 1% PAR in each lake, up to 20% of amplicon

sequences (Fig S1a). The relative abundances of ammonia- and nitrite-oxidizing taxa were

strongly correlated (Spearman's rho = 0.918, $p < 2.2e{-}16$; Fig S1b). Free-living nitrifiers were

rare (<0.1% relative abundance) in bottom water samples from the southern basin of Huron

(HU15M) and western basin of Erie (HU91M); these two stations are the shallowest in our

dataset and have relatively high light penetration to the bottom (~1% PAR). Chlorophyll-*a*

concentration was also negatively correlated with the relative abundance of nitrifiers

(Spearman's rho = -0.677, $p < 1.7e{-}7$; Fig S1c). These findings are consistent with previous

work demonstrating photoinhibition of nitrification (51–54), as well as potential competition with

phototrophs for ammonium (55).

The taxonomic assemblage of nitrifiers, along with nitrogen availability, differed across

lakes and even across stations within a lake (Fig 1, Tables S1-S3). AOB (*Nitrosomonadaceae*)

were observed across all lakes. In contrast, AOA (*Nitrosoarchaeum*) sequences only exceeded

0.5% relative abundance at the three deepest stations (SU08M, MI41M, ON55M) where the

ratio of AOB:AOA ranged from 10:1 to 1:3 (Table S1). A single dominant oligotype accounted

for more than 85% of *Nitrosoarchaeum* sequences in all samples, while four major oligotypes of

Nitrosomonadaceae were observed (data not shown). We found pronounced shifts in the

dominant NOB across stations (Fig 1), and all stations except those in Lake Ontario showed

strong dominance (greater than ten-fold) of either *Ca.* Nitrotoga or *Nitrospira*. *Nitrospira* was the

only nitrite oxidizer detected in Superior and the dominant nitrite oxidizer in parts of Michigan

(MI41M, MI18M). By contrast, *Ca.* Nitrotoga was the only nitrite oxidizer observed in Erie and

Huron and the dominant nitrite oxidizer at the shallowest station in Michigan (MI27M). In all but

one sample, a single dominant oligotype of *Ca*. Nitrotoga and *Nitrospira* accounted for >74%

and >84% of sequences from each taxon, respectively.

***Ecotypic variation in abundant streamlined* Nitrosospira**

We reconstructed 15 genomes of the AOB *Nitrosospira*, substantially expanding genome

descriptions for this genus (56–58). Based on a phylogenomic tree, free-living Great Lakes

Nitrosospira fall into two major clades, both of which are distinct from published species; each of

these clades also includes MAGs recovered from lakes Biwa and Baikal, suggesting novel

globally distributed freshwater lineages (Fig S2). One clade, which we call NspGL1, has a

reduced genome (average 1.4 Mb) and low G+C content (41%) (Fig 2, Table S4), typical of

streamlined genomes (59). The second clade was resolved into three subclades (denoted

NspGL2a, 2b, and 3; Fig S2) based on phylogeny and average nucleotide identity (ANI), all with

small genome sizes of 1.5-1.8 Mb and 50% G+C content (Fig 2, Table S4). Based on short read

mapping, these subclades are ecologically distinct: NspGL1 and NspGL2b -- with the smallest

genomes -- are the dominant AOB in the upper oligotrophic lakes, while NspGL2a is only

abundant in Ontario and NspGL3 is only abundant in Erie (Fig 2). Hereafter we refer to these

subclades as ecotypes due to their phylogenetic and ecological separation.

We compared the genomes of our Great Lakes *Nitrosospira* to 87 reference AOB

genomes belonging to the family *Nitrosomonadaceae,* revealing distinctive patterns of gene

content. Great Lakes *Nitrosospira* genomes encode fewer two-component signal transduction

systems, transposases, motility genes, pilus and secretion genes, and defense-related genes,

and their complement of transporters is distinct from most reference genomes (Table S5). They

also lack functions related to biofilm formation such as polysaccharide matrix production (e.g.

pel genes) and extracellular protein targeting (exosortase and PEP-CTERM motifs). These

features are in sharp contrast to *Nitrosospira* isolates from soil (56,58) and to oligotrophic

Nitrosomonas isolates (60) and are consistent with a passive planktonic lifestyle in extremely

low-nutrient systems.

We next compared metabolic potential among Great Lakes AOB ecotypes to understand

their ecological preferences for upper lakes (NspGL1, NspGL2b), Ontario (NspGL2a), and Erie

(NspGL3). Surprisingly, all seven NspGL1 MAGs contain a cluster of genes encoding

proteorhodopsin and retinal biosynthesis, including 15,15'-Beta-carotene dioxygenase (*blh*),

lycopene cyclase (*crtY*), phytoene dehydrogenase (*crtI*), phytoene synthase (*crtB*), and GGPP

synthase (*crtE*) (Fig 3a). We also identified proteorhodopsin in a single cell amplified genome

(SAG) representing NspGL1 from Lake Michigan (Fig 3a), demonstrating that it is not an artifact

of metagenome assembly. To our knowledge, this is the first example of a nitrifier with

proteorhodopsin. All NspGL1 proteorhodopsins share residues H95, D127 and E138 along with

a short beta-turn (G111-P116) between helices B and C, which are characteristic features of

proteorhodopsin as distinct from sensory and other rhodopsins (61), and the presence of

leucine at position 135 suggests green light tuning (62) (Fig 3b). All of the genes in this module

have highest similarity to homologs from *Polynucleobacter*, but are flanked by

Nitrosomonadaceae-like genes, suggesting recent horizontal gene transfer (Fig 3a). The

predicted NspGL1 proteorhodopsins cluster with *Polynucleobacter*, *Methylopumilus*, and other

freshwater Betaproteobacteria in Supercluster III as defined by MicRhoDE (63) (Fig S3). We

compared the homologous genome region in two highly similar MAGs from Lakes Biwa and

Baikal (Fig S2); these contigs lack the proteorhodopsin module but appear to flank a variable

region where the contig assembly ends (Fig S4). A proteorhodopsin photosystem could support

survival of NspGL1 in the presence of sunlight, which has been shown to inhibit ammonia

oxidation (51,64). In the upper lakes where NspGL1 is abundant, light penetration is high well

below the thermocline in stratified periods (65), and deep water taxa are seasonally advected to

the surface by water column mixing (37). In addition to proteorhodopsin, NspGL1 – but not the

other three ecotypes of Great Lakes *Nitrosospira* – encode a class I cyclopyrimidine dimer

photolyase, which uses light energy to repair UV-induced DNA damage, and the catalase-

peroxidase *katG*, suggesting that the NspGL1 ecotype is adapted to relatively shallow depths in

the water column (Fig 2).

Great Lakes *Nitrosospira* carry a reduced, ecotype-specific complement of nitrogen

metabolism genes compared to reference AOB (Fig 2; Table S5). All are presumed to have the

core ammonia oxidation enzymes ammonia monooxygenase and hydroxylamine

dehydrogenase; these genes were assembled and binned as expected in some MAGs and

were manually identified on short unbinned contigs in other cases. Surprisingly, all Great Lakes

Nitrosospira lack the copper protein nitrosocyanin, whose precise function is unknown but so far

has been found in all AOB except one member of the *N. oligotropha* clade (60). The lack of

nitrosocyanin extends beyond the Great Lakes MAGs to closely related freshwater and marine

strains, along with five more members of the *N. oligotropha* clade (Fig S2); its absence may be

related to the divergence of these clades. Only NspGL1 and NspGL2b encode NO-forming

nitrite reductase (NirK), which confers nitrite tolerance (66); this result is surprising given that

these two clades dominate the upper lakes where productivity and reduced N are lowest. None

of the Great Lakes MAGs encode NO reductase (NorCBQD), and NspGL1 lacks cytochrome

P460 family proteins; these genomic patterns may influence N_2O emissions across the Great

Lakes. Nitrogen acquisition is also distinct among Great Lakes AOB: NspGL1 lacks an apparent

ammonium transporter, but encodes urease structural and accessory genes (*ureABCEFG*) and

a high-affinity urea transporter (*urtABCDE)*. Further, all Great Lakes ecotypes encode a high-

affinity amino acid transporter (*livFGHM*) and a secreted non-specific nuclease; these genes are

rare (<5%) in reference genomes and could supply reduced nitrogen and/or organic carbon.

Finally, NspGL1 and NspGL3 have genes for producing cyanophycin, an intracellular storage

compound for nitrogen (58,67). Together, the minimal and distinctive gene complements

present in Great Lakes *Nitrosospira* illustrate the variability and adaptability of AOB gene

content across habitats.

As with nitrogen metabolism, carbon metabolism is also distinct between Great Lakes

and reference AOB, and among Great Lakes ecotypes (Fig 2; Table S5). Unlike most reference

AOB, Great Lakes *Nitrosospira* lack two key enzymes of the oxidative pentose phosphate

pathway, glucose-6-phosphate dehydrogenase and 6-phosphogluconate dehydrogenase. All

ecotypes except Erie-specific NspGL3 also lack genes for glycogen synthesis and degradation.

These findings suggest continuous autotrophy, rather than "feast or famine" dynamics of carbon

storage and utilization, or diel periodicity as in photoautotrophs. The key enzyme for carbon

fixation, RuBisCO, has evolved several kinetically distinct forms whose distribution likely reflects

ecological pressures (68). NspGL1 and NspGL3 both contain Form IA RuBisCO, while

NspGL2a and NspGL2b contain Form IC RuBisCO (68,69) (Fig 2). NspGL1 genomes also

possess an alpha carboxysome-like *cso* operon, similar to *Nitrosomonas eutropha* C91 (69),

though our draft assembly lacks the expected carbonic anhydrase (*csoS3/csoSCA*).

Carboxysome-associated RuBisCO may allow NspGL1, the ecotype most strongly adapted to

energy and nutrient limitation, to more efficiently fix CO_2 by minimizing the wasteful oxygenation

reaction and reducing the cellular nitrogen allocation to RuBisCO (68). The ranges of kinetic

properties observed in other autotrophs for Form IAq (found in NspGL3) and Form IC (found in

NspGL2a and NspGL2b) overlap, and therefore more work is needed to understand the fitness

advantages, if any, that this RuBisCO diversity confers on Great Lakes nitrifiers.

Streamlined freshwater Thaumarchaeota

We reconstructed three similar genomes (>99% ANI) of *Nitrosoarchaeum* (NarchGL; Fig

S5, Fig S6, Table S4) from three separate samples (two from Superior, one from Ontario),

consistent with our low observed 16S rRNA diversity for *Thaumarchaeota*. These NarchGL

genomes are very similar (~99% ANI) to two genomes reconstructed from Lake Baikal, located

thousands of kilometers away (70). Their next closest relatives are also from freshwater

environments, and phylogenetic clustering suggests that salinity is an important driver of

divergence throughout the *Nitrosopumilaceae* (Fig S5). Using our reconstructed genomes as

probes for metagenomic read recruitment, NarchGL were detected in Superior, Michigan, and

Ontario; they represented roughly one-third of ammonia oxidizers in the mid-hypolimnion of

station SU08M, reaching relative abundance of approximately 0.5% (Fig 2).

NarchGL share nearly 90% of their predicted proteins with close relatives including *Ca.*

Nitrosoarchaeum limnia; at the same time, they show distinctive patterns in gene content that

pinpoint the key selective pressures of deep lakes (Fig 2, Table S6). All three NarchGL

genomes encode urease and a urea transporter, implicating urea as a vital source of nitrogen

for energy and/or biosynthesis. Consistent with phosphorus scarcity in much of the Great Lakes

(71), NarchGL encode high affinity transport systems for phosphate and potentially

phosphonates, though we did not identify a phosphonate lyase. In addition to both CRISPR/Cas

enzymes cas1 and cas4, NarchGL genomes contain several phage proteins, suggesting viral

infection and integration events may be common. DNA photolyases, which have been found in

epipelagic clades of marine *Thaumarchaeota* (72–74), are present in all low salinity

Nitrosoarchaeum including NarchGL, suggesting NarchGL are exposed to sunlight due to high

water clarity (65) and/or annual mixing in the Great Lakes. NarchGL also lack the common

tRNA modification 4-thiouridylation (indicated by IscS K04487 and ThiI-NFACT PF02568-

PF18297; ref. 75); we propose that the absence of this modification, which is susceptible to

near-UV radiation (75), is also related to sunlight exposure.

While aquatic AOA are typically adapted to oligotrophic conditions, NarchGL appear

especially streamlined. Their genomes reveal striking reduction in environmental sensing,

response, and regulatory functions, relative to other *Nitrosoarchaeum* and *Nitrosopumilaceae*

(Table S6). NarchGL encode fewer variants of the general archaeal transcription factors, TATA

binding protein (TBPs, PF00352) and transcription factor B (TFBs, PF00382 and PF08271),

combinations of which influence differential gene expression (76). NarchGL also encode fewer

two-component systems that transmit environmental signals to control gene expression or

protein activity (domains: cache PF02743, HAMP PF00672, His kinase A PF00512, receiver

PF00072). Further, they are depleted in ArsR family transcription factors (PF01022), P-II

proteins for regulation of nitrogen metabolism (PF00543), and other potential regulatory

domains (CBS PF00571, USP PF00582). This extremely limited regulatory capacity in NarchGL

stands in sharp contrast to closely related *Ca.* N. limnia, but instead parallels the oceanic

minimalist *Ca.* Nitrosopelagicus brevis (11) (Table S6).

Expanded diversity of Ca.* Nitrotoga *with reduced genomes

Despite the broad distribution of *Ca.* Nitrotoga in freshwater systems and beyond, only

six genomes are available, derived from rivers heavily impacted by urban and agricultural

influence, a wastewater treatment plant, and coastal sediment (30,77,78). Hence the metabolic

and phylogenetic diversity of this group is virtually unexplored. We reconstructed six new MAGs

of *Ca*. Nitrotoga, which form two clusters with >99% ANI within each cluster and ~97% ANI

between clusters (NtogaGL1a and NtogaGL1b; Fig S2). These new *Ca*. Nitrotoga MAGs are far

smaller than published genomes (1.5 Mb compared to 2.6-3.0 Mb; Fig 4, Table S4), lacking

functions such as motility and chemotaxis, pilus biogenesis, and DNA repair (*mutLS*) (Table

S7). Great Lakes *Ca.* Nitrotoga also encode markedly fewer two-component systems for

sensing and responding to environmental cues, defense-related genes (restriction-modification,

toxin-antitoxin, and Crispr-Cas systems), transposases, and efflux transporters (Table S7).

While incomplete assembly of hypervariable genome regions may explain some of these

absences, the overall genome properties are consistent with a relatively stable low-nutrient

environment and planktonic lifestyle.

The reduced genomes of NtogaGL1a/b help clarify core features of the genus *Nitrotoga*,

along with accessory functions that may enable local adaptation in specific populations. To date,

sequenced *Ca.* Nitrotoga including NtogaGL1a/b encode similar electron transport pathways,

including NADH dehydrogenase-Complex I, succinate dehydrogenase-Complex II, and

Alternative Complex III, along with high-affinity cbb3-type cytochrome oxidases suggesting

adaptation to low oxygen conditions. They also share the Calvin cycle for carbon fixation, a

complete TCA cycle, and an evolutionarily distinct nitrite oxidoreductase (NXR) from other NOB

(30,77,78). All *Ca.* Nitrotoga to date also share transporters for amino acids and peptides,

potential sources of C and/or N. *Ca.* Nitrotoga can also potentially access reduced sulfur

compounds for energy via sulfite dehydrogenase, suggesting metabolic flexibility beyond nitrite

oxidation.

Beyond these similarities, the small genomes of NtogaGL1a/b are distinct from

previously described *Ca.* Nitrotoga in many ways. NtogaGL1a/b lack NiFe hydrogenase to use

hydrogen as an energy source. They also lack nitrogen metabolism functions including

assimilatory nitrite reductase (*nirBD*), nitrite reductase to NO (*nirK*), and nitric oxide reductase

(*norQD*). Based on gene content, NtogaGL1a/b appear unable to use hexoses like glucose,

since they lack the glycolytic enzyme phosphofructokinase and the Entner-Doudoroff pathway,

similar to *Nitrobacter winogradskyi* (79). Consistent with this, they also lack genes for storage

and breakdown of glycogen (Table S7). All but one of the NtogaGL1a/1b genomes encode

cyanate lyase (*cynS*), which is found in other NOB but no *Ca.* Nitrotoga to date (27,80,81). The

cynS gene, adjacent to *glnK-amtB* for ammonium sensing and transport, likely functions in N

assimilation, as recently described for *Nitrospinae* (22). While cyanase has been shown to

mediate reciprocal feeding between some NOB and ammonia oxidizers (27), this interaction is

unlikely in the free-living (<1.6um) size-fraction and dilute environment sampled here. Notably,

cyanase from NtogaGL1a/1b, along with predicted *Nitrospirae* proteins from Lake Baikal and

soil, form a distinct phylogenetic cluster from most nitrifier cyanase proteins observed to date

(Fig S7).

The two ANI-based clusters we detected, NtogaGL1a and NtogaGL1b, appear to be

phylogenetically and ecologically distinct ecotypes. Based on short-read mapping, NtogaGL1b

dominates Erie, while NtogaGL1a dominates all other *Ca.* Nitrotoga-containing samples (Fig 4).

We found several metabolic genes that differentiate the two ecotypes. ER-specific NtogaGL1b

genomes share a region encoding thiosulfate dehydrogenase (*tsdA*), cytochromes, transport of

sulfur-containing compounds, lactate dehydrogenase (ldh*),* a two-component system, and a

Crp-family transcription factor (Fig S8). This region may be involved in oxidizing thiosulfate as

an energy source, and sensing and responding to redox changes that accompany seasonal

hypoxia in Lake Erie. The corresponding region in NtogaGL1a encodes an integrase and

photolyase, consistent with greater DNA photodamage in the more transparent waters of

Michigan, Huron, and Ontario where NtogaGL1a is abundant. NtogaGL1a also lacks superoxide

oxidase (*cybB*) found in NtogaGL1b and reference strains, suggesting these two ecotypes

experience differential oxidative stress.

Great Lakes* Nitrospira *reveal adaptations to sunlit oxic environment

We reconstructed six closely related genomes of *Nitrospira* (~99% ANI; Fig S9, Table

S4), representing the predominant NOB throughout Lake Superior and in parts of Michigan,

Ontario, and Huron (Fig. 4; Table S1). These genomes, which we refer to as NspiraGL, fall

within lineage II (Fig S9), which is broadly distributed across soil, freshwater, and engineered

habitats (21); however, genome analyses to date have focused on strains from wastewater and

engineered systems, leaving major blind spots. NspiraGL share core features of *Nitrospira*

metabolism, including a periplasmic-facing NXR that is advantageous under substrate-limiting

conditions, multiple cytochrome *bd*-like oxidases, and the reverse TCA cycle for carbon fixation

(80). However, as with *Ca.* Nitrotoga, the *Nitrospira* genomes we reconstructed in the Great

Lakes are markedly smaller (~1.8 Mb) than published reference genomes (Fig. 4; Table S4),

with reduced capacity for environmental sensing, transport, defense, and transposition, and a

lack of pilus or flagellar motility (Table S8). NspiraGL encode just five sigma factors, compared

to 18 in *Nitrospira moscoviensis,* consistent with genome streamlining theory (59). Further,

NspiraGL genomes encode a single NXR, while *N. moscoviensis* carries five copies that are

differentially regulated (29,82). NspiraGL also lack the *glnE* gene for glutamine synthetase (GS)

adenylyltransferase, suggesting that GS activity is not repressed by this mechanism. Together,

these features suggest limited regulatory and ecological flexibility, consistent with a relatively

constant, oligotrophic environment.

Compared to other *Nitrospira,* NspiraGL exhibit limited energetic flexibility, but can

access diverse nitrogen sources (Fig. 4, Table S8). We predict that NspiraGL are unable to

grow on hydrogen or formate as alternative energy sources (25,29), as they lack NiFe-

hydogenase and formate dehydrogenase. The glycolysis and oxidative TCA cycles appear to be

incomplete, lacking phosphofructokinase and citrate synthase, respectively; this suggests a

limited capacity for organic carbon utilization. Regarding nitrogen metabolism, NspiraGL lack

nirK, encoding NO-forming nitrite reductase, but maintain nitric oxide reductase (*norQD*),

implying that cellular NO is formed by other mechanisms. To obtain nitrogen for biosynthesis,

NspiraGL encode a high-affinity nitrate/nitrite/cyanate transporter (*nrtABC*), assimilatory nitrite

reductase (*nirA*), and cyanase (*cynS*), along with two *amt* family ammonium transporters.

Although none of the NspiraGL MAGs include urease (*ureCBA*), one does contain urease

accessory proteins (*ureEFGD*) and two contain a urea transporter (*urtABCD*), suggesting

incomplete assembly of the urea utilization pathway. As with *Ca.* Nitrotoga, we suggest that

cyanase, along with urease where present, functions in nitrogen assimilation rather than cross-

feeding, given the dilute environment and free-living planktonic cells.

Beyond energy, carbon, and nitrogen metabolism, we discovered striking differences

between NspiraGL and reference *Nitrospira* related to DNA repair. NspiraGL encode two

additional photolyase-related proteins, along with a class I cyclopyrimidine dimer (CPD)

photolyase found in most reference *Nitrospira* (Fig 5, Fig S10). Photolyases use blue light

energy to repair DNA lesions caused by UV radiation (83). The two additional genes in

NspiraGL are adjacent and share best hits with Betaproteobacteria, suggesting recent

horizontal transfer (Fig 5). One likely encodes an FeS-BCP photolyase, which repairs (6-4)

dipyrimidine lesions (84,85). The other shares an FAD-binding domain with photolyases but the

C-terminal region has no recognizable domains (Fig 5b). This protein is widespread in aquatic

bacteria and has not been functionally characterized, though an actinobacterial homolog was

suggested to be involved in light sensing and regulation (86). Beyond photolyases, NspiraGL

also encode uracil-DNA glycosylase (UNG), which removes misincorporated uracil from DNA.

Uracil results from deamination of cytosine, which can occur spontaneously or be induced by

NO (87). In addition to the photolyases and UNG that repair DNA lesions, NspiraGL encode

translesion DNA polymerase V (*umuCD*) which enables replication to proceed past lesions.

Together, these genes indicate that *Nitrospira* in the Great Lakes experience significant DNA

damage, including UV-induced damage that also requires light for the repair process, in

hypolimnion waters with high transparency (65) and/or during seasonal mixing.

Other major differences between NspiraGL and reference *Nitrospira* are related to

reactive oxygen species (ROS) (Table S8). Surprisingly, despite their oxic habitat, NspiraGL

lack superoxide dismutase (SOD), monofunctional catalase (*katE*), classic regulators of

oxidative stress response (*oxyR* and *perR*), and bacterioferritin, which limits the Fenton reaction

by sequestering free iron. However, all six NspiraGL MAGs, but few reference genomes (7% of

75), have recently acquired bifunctional catalase-peroxidase *katG*; interestingly we also

observed *katG* in Great Lakes *Ca.* Nitrotoga and *Nitrosospira* (Fig 2, Fig 4). The absence of

SOD suggests that NspiraGL does not produce damaging levels of endogenous superoxide,

perhaps because NspiraGL lack the major respiratory and non-respiratory flavoproteins that

produce ROS in other SOD-containing *Nitrospira* (88). Unlike superoxide, H_2O_2 can cross

membranes, and is known to be produced by both photooxidation of dissolved organic matter

and dark heterotrophic activity (89). The lakes where NspiraGL dominate have high water clarity

(65) and low productivity, consistent with abiotic photochemistry as the primary source of

exogenous ROS; this stress may have selected for *katG* as a defense, though without the

canonical *oxyR* regulation. The absence of bacterioferritin could be related to iron limitation (71).

NspiraGL also lack cytochrome *c* peroxidase, which is found in 70 of 75 reference genomes;

this protein is proposed to function in anaerobic respiration of H_2O_2 (90) and therefore its

absence in NspiraGL is consistent with a constant oxic environment. Together, these results

indicate that *Nitrospira* in the Great Lakes face distinct ROS pressures that have shaped their

gene content.

Discussion

The Laurentian Great Lakes harbor nitrifiers that are phylogenetically related, but

markedly different in genome size and functional capacity, from their well-studied relatives

inhabiting wastewater systems, soils, and even other freshwater systems. By examining the

entire nitrifier assemblage at once, we detected common features across taxa that illuminate the

selective pressures faced by microbes in deep lakes. All the lineages we describe show small

genome sizes (1.3-1.8 Mb), reduced capacity for environmental sensing and response, and

adaptation to a passive (i.e. non-motile) planktonic lifestyle, expanding the paradigm of

streamlining observed for oceanic AOA (11) to AOB, *Nitrospira*, and *Ca.* Nitrotoga in

oligotrophic freshwaters. Within the cosmopolitan *Nitrosospira*, we found ecotypes with a

gradient of genome reduction that maps onto their habitats' trophic gradient: from NspGL1 (1.4

Mb, low GC, upper lakes) to NspGL2b (1.5 Mb, upper lakes) to NspGL2a (1.6 Mb, Ontario) to

NspGL3 (1.8 Mb, Erie) (Fig 2). The thaumarchaeal NarchGL have markedly reduced regulatory

capacity like the open ocean strain *Nitrosopelagicus brevis* (11). The NOB NspiraGL have

genomes 50-60% smaller than described *Nitrospira* and dominate, along with the AOB NspGL1

and the AOA NarchGL, the deeper more oligotrophic basins, while *Ca.* Nitrotoga favor shallower

more productive basins. The emergence of Erie-specific ecotypes of both *Nitrosospira*

(NspGL3) and *Ca.* Nitrotoga (NtogaGL1b) demonstrates how distinct this habitat is compared to

the other lakes.

Nitrifiers inhabiting the transparent waters of the upper Great Lakes show distinctive

adaptations to light including diverse photolyases, ROS detoxification, and even

proteorhodopsin. This discovery is surprising, given that nitrifiers are rare in the surface mixed

layer of the Great Lakes (Fig. S1) and that photoinhibition of ammonia oxidation and nitrifier

growth is well documented (51-54,64). We propose that proteorhodopsin could be used to

augment energy metabolism when ammonia oxidation is photoinhibited. Water clarity has

increased over the past several decades in Lakes Michigan and Huron, now surpassing that of

Lake Superior (65). High light penetration along with seasonal mixing likely exposes deep water

cells to damaging levels of light and oxidative stress. Future cultivation and physiological studies

will document photoinhibition and potential phototrophy in Great Lakes nitriifiers.

The capacity for nitrification is found across multiple phyla, and our work unveils new

clues to understanding the ecological and evolutionary potential of these diverse lineages. This

collective nitrifier diversity undoubtedly influences the cycling of carbon and nitrogen across this

ecosystem, and future work will explore the differential contributions to nitrification by the distinct

lineages we described here. Understanding what controls the diversity of nitrifiers and other key

functional groups, and the consequences of this diversity for biogeochemistry, are essential for

forecasting the effects of rapid environmental change across the large lakes of the world (e.g.

ref. 91) and predicting impacts on the critical ecosystem services they provide (92).

References

1.   Canfield DE, Glazer AN, Falkowski PG. The evolution and future of Earth's nitrogen cycle.
Science 2010; 330:192–6.
2.   Pachiadaki MG, Sintes E, Bergauer K, Brown JM, Record NR, Swan BK, et al. Major role
of nitrite-oxidizing bacteria in dark ocean carbon fixation. *Science* 2017; 358:1046–51.
3.   Reinthaler T, van Aken HM, Herndl GJ. Major contribution of autotrophy to microbial
carbon cycling in the deep North Atlantic's interior. *Deep Sea Res Part II: Topical Studies*
in Oceanography 2010; 57:1572–80.
4.   Swan BK, Martinez-Garcia M, Preston CM, Sczyrba A, Woyke T, Lamy D, et al. Potential
for chemolithoautotrophy among ubiquitous bacteria lineages in the dark ocean. *Science*
2011; 333:1296–300.
5.   Baltar F, Herndl GJ. Ideas and perspectives: Is dark carbon fixation relevant for oceanic
primary production estimates? *Biogeosciences* 2019; 16:3793–9.
6.   Callieri C, Coci M, Eckert EM, Salcher MM, Bertoni R. Archaea and Bacteria in deep lake
hypolimnion: in situ dark inorganic carbon uptake. *J Limnol* 2014; 73:31-38.
7.   Wuchter C, Abbas B, Coolen MJL, Herfort L, van Bleijswijk J, Timmers P, et al. Archaeal
nitrification in the ocean. *Proc Natl Acad Sci USA* 2006; 103:12317–22.
8.   Francis CA, Roberts KJ, Beman JM, Santoro AE, Oakley BB. Ubiquity and diversity of
ammonia-oxidizing archaea in water columns and sediments of the ocean. *Proc Natl Acad*
Sci USA 2005; 102:14683–8.
9.   Schleper C. Ammonia oxidation: different niches for bacteria and archaea? *ISME J* 2010;
4:1092–4.
10. Martens-Habbena W, Berube PM, Urakawa H, de la Torre JR, Stahl DA. Ammonia
oxidation kinetics determine niche separation of nitrifying Archaea and Bacteria. *Nature*
2009; 461:976–9.
11. Santoro AE, Dupont CL, Richter RA, Craig MT, Carini P, McIlvin MR, et al. Genomic and
proteomic characterization of "*Candidatus* Nitrosopelagicus brevis": An ammonia-oxidizing
archaeon from the open ocean. *Proc Natl Acad Sci USA* 2015; 112:1173–8.
12. Auguet J-C, Triadó-Margarit X, Nomokonova N, Camarero L, Casamayor EO. Vertical
segregation and phylogenetic characterization of ammonia-oxidizing Archaea in a deep
oligotrophic lake. *ISME J* 2012; 6:1786–97.
13. Herber J, Klotz F, Frommeyer B, Weis S, Straile D, Kolar A, et al. A single Thaumarchaeon
drives nitrification in deep oligotrophic Lake Constance. *Environ Microbiol* 2020; 22:212–
28.
14. Urbach E, Vergin KL, Young L, Morse A, Larson GL, Giovannoni SJ. Unusual
bacterioplankton community structure in ultra-oligotrophic Crater Lake. *Limnol Oceanogr*
2001; 46:557–72.
15. Okazaki Y, Fujinaga S, Tanaka A, Kohzu A, Oyagi H, Nakano S. Ubiquity and quantitative
significance of bacterioplankton lineages inhabiting the oxygenated hypolimnion of deep
freshwater lakes. *ISME J* 2017; 11:2279–93.
16. Mukherjee M, Ray A, Post AF, McKay RM, Bullerjahn GS. Identification, enumeration and
diversity of nitrifying planktonic archaea and bacteria in trophic end members of the
Laurentian Great Lakes. *J Great Lakes Res* 2016; 42:39–49.
17. Hugoni M, Etien S, Bourges A, Lepère C, Domaizon I, Mallet C, et al. Dynamics of
ammonia-oxidizing Archaea and Bacteria in contrasted freshwater ecosystems. *Res*
Microbiol 2013; 164:360–70.
18. Bollmann A, Bar-Gilissen M-J, Laanbroek HJ. Growth at low ammonium concentrations
and starvation response as potential factors involved in niche differentiation among
ammonia-oxidizing bacteria. *Appl Environ Microbiol* 2002; 68:4751–7.
19. Sedlacek CJ, McGowan B, Suwa Y, Sayavedra-Soto L, Laanbroek HJ, Stein LY, et al. A

physiological and genomic comparison of *Nitrosomonas* cluster 6a and 7 ammonia-
oxidizing bacteria. *Microb Ecol* 2019; 78:985–94.
20. Beman JM, Popp BN, Francis CA. Molecular and biogeochemical evidence for ammonia
oxidation by marine Crenarchaeota in the Gulf of California. *ISME J* 2008; 2:429–41.
21. Daims H, Lücker S, Wagner M. A new perspective on microbes formerly known as nitrite-
oxidizing bacteria. *Trends Microbiol* 2016; 24:699–712.
22. Kitzinger K, Marchant HK, Bristow LA, Herbold CW, Padilla CC, Kidane AT, et al. Single
cell analyses reveal contrasting life strategies of the two main nitrifiers in the ocean. *Nature
Comm* 2020; 11:767.
23. Nowka B, Daims H, Spieck E. Comparison of oxidation kinetics of nitrite-oxidizing bacteria:
nitrite availability as a key factor in niche differentiation. *Appl Environ Microbiol* 2015;
81:745–53.
24. Wegen S, Nowka B, Spieck E. Low temperature and neutral pH define "*Candidatus*
Nitrotoga sp." as a competitive nitrite oxidizer in coculture with *Nitrospira defluvii. Appl
Environ Microbiol* 2019; 85:e02569-18.
25. Koch H, Galushko A, Albertsen M, Schintlmeister A, Gruber-Dorninger C, Lucker S, et al.
Growth of nitrite-oxidizing bacteria by aerobic hydrogen oxidation. *Science* 2014;
345:1052–4.
26. Füssel J, Lücker S, Yilmaz P, Nowka B, van Kessel MAHJ, Bourceau P, et al. Adaptability
as the key to success for the ubiquitous marine nitrite oxidizer *Nitrococcus. Science
Advances* 2017; 3:e1700807.
27. Palatinszky M, Herbold C, Jehmlich N, Pogoda M, Han P, von Bergen M, et al. Cyanate as
an energy source for nitrifiers. *Nature* 2015; 524:105–8.
28. Kitzinger K, Padilla CC, Marchant HK, Hach PF, Herbold CW, Kidane AT, et al. Cyanate
and urea are substrates for nitrification by Thaumarchaeota in the marine environment. *Nat
Microbiol* 2019; 4:234.
29. Koch H, Lücker S, Albertsen M, Kitzinger K, Herbold C, Spieck E, et al. Expanded
metabolic versatility of ubiquitous nitrite-oxidizing bacteria from the genus *Nitrospira. Proc
Natl Acad Sci USA* 2015; 112:11371–6.
30. Boddicker AM, Mosier AC. Genomic profiling of four cultivated *Candidatus* Nitrotoga spp.
predicts broad metabolic potential and environmental distribution. *ISME J* 2018; 12:2864–
82.
31. Sterner RW. In situ-measured primary production in Lake Superior. *J Great Lakes Res*
2010; 36:139–49.
32. Small GE, Bullerjahn GS, Sterner RW, Beall BFN, Brovold S, Finlay JC, et al. Rates and
controls of nitrification in a large oligotrophic lake. *Limnol Oceanogr* 2013; 58:276–86.
33. Vollenweider RA, Munawar M, Stadelmann P. A comparative review of phytoplankton and
primary production in the Laurentian Great Lakes. *J Fish Res Bd Can* 1974; 31:739–62.
34. Clevinger CC, Heath RT, Bade DL. Oxygen use by nitrification in the hypolimnion and
sediments of Lake Erie. *J Great Lakes Res* 2014; 40:202–7.
35. Neilson MA, Stevens RJJ. Spatial heterogeneity of nutrients and organic matter in Lake
Ontario. *Can J Fish Aquat Sci* 1987; 44:2192–203.
36. Lean DRS, Knowles R. Nitrogen transformations in Lake Ontario. *Can J Fish Aquat Sci*
1987; 44:2133–43.
37. Paver SF, Newton RJ, Coleman ML. Microbial communities of the Laurentian Great Lakes
reflect connectivity and local biogeochemistry. *Environ Microbiol* 2020; 22:433–46.
38. Rozmarynowycz MJ, Beall BFN, Bullerjahn GS, Small GE, Sterner RW, Brovold SS, et al.
Transitions in microbial communities along a 1600 km freshwater trophic gradient. *J Great
Lakes Res* 2019; 45:263–76.
39. Fujimoto M, Cavaletto J, Liebig JR, McCarthy A, Vanderploeg HA, Denef VJ.
Spatiotemporal distribution of bacterioplankton functional groups along a freshwater

estuary to pelagic gradient in Lake Michigan. *J Great Lakes Res* 2016; 42:1036–48.

40. Barbiero RP, Lesht BM, Hinchey EK, Nettesheim TG. A brief history of the U.S. EPA Great
Lakes National Program Office's water quality survey. *J Great Lakes Res* 2018; 44:539–
46.

41. US Environmental Protection Agency. Sampling and analytical procedures for GLNPO's
open lake water quality survey of the Great Lakes. Great Lakes National Program Office,
Illinois, USA; 2003.

42. Holmes RM, Aminot A, Kérouel R, Hooker BA, Peterson BJ. A simple and precise method
for measuring ammonium in marine and freshwater ecosystems. *Can J Fish Aquat Sci*
1999; 56:1801–8.

43. Miranda KM, Espey MG, Wink DA. A rapid, simple spectrophotometric method for
simultaneous detection of nitrate and nitrite. *Nitric Oxide* 2001; 5:62–71.

44. Parada AE, Needham DM, Fuhrman JA. Every base matters: assessing small subunit
rRNA primers for marine microbiomes with mock communities, time series and global field
samples. *Environ Microbiol* 2016; 18:1403–14.

45. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal
RNA gene database project: improved data processing and web-based tools. *Nucleic*
Acids Res 2013; 41:D590–6.

46. Wang Q, Garrity GM, Tiedje JM, Cole JR. Naïve Bayesian classifier for rapid assignment
of rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol* 2007;
73:5261–7.

47. Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. Development of a dual-
index sequencing strategy and curation pipeline for analyzing amplicon sequence data on
the MiSeq Illumina sequencing platform. *Appl Environ Microbiol* 2013; 79:5112–20.

48. Eren AM, Maignien L, Sul WJ, Murphy LG, Grim SL, Morrison HG, et al. Oligotyping:
differentiating between closely related microbial taxa using 16S rRNA gene data. *Methods*
Ecol Evol 2013; 4:1111–9.

49. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. metaSPAdes: a new versatile
metagenomic assembler. *Genome Res* 2017; 27:824–34.

50. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a
new genome assembly algorithm and its applications to single-cell sequencing. *J Comput*
Biol 2012; 19:455–77.

51. Hooper AB, Terry KR. Photoinactivation of ammonia oxidation in *Nitrosomonas*. *J Bacteriol*
1974; 119:899-906.

52. Horrigan SG, Springer AL. Oceanic and estuarine ammonium oxidation: Effects of light.
Limnol Oceanogr 1990; 35:479–82.

53. Guerrero M, Jones R. Photoinhibition of marine nitrifying bacteria. I. Wavelength-
dependent response. *Mar Ecol Prog Ser* 1996; 141:183–92.

54. Merbt SN, Stahl DA, Casamayor EO, Martí E, Nicol GW, Prosser JI. Differential
photoinhibition of bacterial and archaeal ammonia oxidation. *FEMS Microbiol Lett* 2012;
327:41–6.

55. Smith JM, Chavez FP, Francis CA. Ammonium uptake by phytoplankton regulates
nitrification in the sunlit ocean. *PLoS ONE* 2014; 9:e108173.

56. Rice MC, Norton JM, Valois F, Bollmann A, Bottomley PJ, Klotz MG, et al. Complete
genome of *Nitrosospira briensis* C-128, an ammonia-oxidizing bacterium from agricultural
soil. *Stand Genomic Sci* 2016; 11:46.

57. Garcia JC, Urakawa H, Le VQ, Stein LY, Klotz MG, Nielsen JL. Draft genome sequence of
Nitrosospira sp. Strain APG3, a psychrotolerant ammonia-oxidizing bacterium isolated
from sandy lake sediment. *Genome Announcements* 2013; 1:e00930-13.

58. Norton JM, Klotz MG, Stein LY, Arp DJ, Bottomley PJ, Chain PSG, et al. Complete
genome sequence of *Nitrosospira multiformis*, an ammonia-oxidizing bacterium from the

soil environment. *Appl Environ Microbiol* 2008; 74:3559–72.

59. Giovannoni SJ, Cameron Thrash J, Temperton B. Implications of streamlining theory for
microbial ecology. *ISME J* 2014; 8:1553–65.

60. Bollmann A, Sedlacek CJ, Norton J, Laanbroek HJ, Suwa Y, Stein LY, et al. Complete
genome sequence of *Nitrosomonas sp.* Is79, an ammonia oxidizing bacterium adapted to
low ammonium concentrations. *Stand Genomic Sci* 2013; 7:469–82.

61. Reckel S, Gottstein D, Stehle J, Löhr F, Verhoefen M-K, Takeda M, et al. Solution NMR
structure of proteorhodopsin. *Angewandte Chemie* 2011; 50:11942–6.

62. Kralj JM, Bergo VB, Amsden JJ, Spudich EN, Spudich JL, Rothschild KJ. Protonation state
of Glu142 differs in the green- and blue-absorbing variants of proteorhodopsin. *Biochem*
2008; 47:3447–53.

63. Boeuf D, Audic S, Brillet-Guéguen L, Caron C, Jeanthon C. MicRhoDE: a curated
database for the analysis of microbial rhodopsin diversity and evolution. *Database* 2015;
bav080.

64. Hyman MR, Arp DJ. $^{14}C_2H_2$- and $^{14}CO_2$-labeling studies of the de novo synthesis of
polypeptides by *Nitrosomonas europaea* during recovery from acetylene and light
inactivation of ammonia monooxygenase. *J Biol Chem* 1992; 267:1534–45.

65. Yousef F, Shuchman R, Sayers M, Fahnenstiel G, Henareh A. Water clarity of the Upper
Great Lakes: Tracking changes between 1998–2012. *J Great Lakes Res* 2017; 43:239–47.

66. Beaumont HJE, Hommes NG, Sayavedra-Soto LA, Arp DJ, Arciero DM, Hooper AB, et al.
Nitrite reductase of *Nitrosomonas europaea* is not essential for production of gaseous
nitrogen oxides and confers tolerance to nitrite. *J Bacteriol* 2002; 184:2557–60.

67. Watzer B, Forchhammer K. Cyanophycin synthesis optimizes nitrogen utilization in the
unicellular cyanobacterium Synechocystis sp. strain PCC 6803. *Appl Environ Microbiol*
2018; 84:e01298-18.

68. Badger MR, Bek EJ. Multiple Rubisco forms in proteobacteria: their functional significance
in relation to CO_2 acquisition by the CBB cycle. *J Exp Bot* 2008; 59:1525–41.

69. Rae BD, Long BM, Badger MR, Price GD. Functions, compositions, and evolution of the
two types of carboxysomes: polyhedral microcompartments that facilitate CO_2 fixation in
cyanobacteria and some Proteobacteria. *Microbiol Mol Biol Rev* 2013; 77:357–79.

70. Cabello-Yeves PJ, Zemskaya TI, Rosselli R, Coutinho FH, Zakharenko AS, Blinov VV, et
al. Genomes of novel microbial lineages assembled from the sub-ice waters of Lake
Baikal. *Appl Environ Microbiol* 2018; 84: e02132-17.

71. Sterner RW, Smutka TM, McKay RML, Xiaoming Q, Brown ET, Sherrell RM. Phosphorus
and trace metal limitation of algae and bacteria in Lake Superior. *Limnol Oceanogr* 2004;
49:495–507.

72. Bayer B, Vojvoda J, Offre P, Alves RJE, Elisabeth NH, Garcia JA, et al. Physiological and
genomic characterization of two novel marine thaumarchaeal strains indicates niche
differentiation. *ISME J* 2016; 10:1051–63.

73. Luo H, Tolar BB, Swan BK, Zhang CL, Stepanauskas R, Ann Moran M, et al. Single-cell
genomics shedding light on marine Thaumarchaeota diversification. *ISME J* 2014; 8:732–
6.

74. Ahlgren NA, Chen Y, Needham DM, Parada AE, Sachdeva R, Trinh V, et al. Genome and
epigenome of a novel marine Thaumarchaeota strain suggest viral infection,
phosphorothioation DNA modification and multiple restriction systems. *Environ Microbiol*
2017; 19:2434–52.

75. Ryals J, Hsu RY, Lipsett MN, Bremer H. Isolation of single-site *Escherichia coli* mutants
deficient in thiamine and 4-thiouridine syntheses: identification of a nuvC mutant. *J*
Bacteriol 1982; 151:899–904.

76. Karr EA. Transcription Regulation in the Third Domain. In: Sariaslani S, Gadd GM (eds).
Advances in Applied Microbiology. Academic Press, 2014, pp. 101–33.

77. Kitzinger K, Koch H, Lücker S, Sedlacek CJ, Herbold C, Schwarz J, et al. Characterization
of the first "*Candidatus* Nitrotoga" isolate reveals metabolic versatility and separate
evolution of widespread nitrite-oxidizing bacteria. *mBio* 2018; 9:e01186-18.
78. Ishii K, Fujitani H, Sekiguchi Y, Tsuneda S. Physiological and genomic characterization of
a new "*Candidatus* Nitrotoga" isolate. *Environ Microbiol* 2020; 22:2365–82.
79. Starkenburg SR, Chain PSG, Sayavedra-Soto LA, Hauser L, Land ML, Larimer FW, et al.
Genome sequence of the chemolithoautotrophic nitrite-oxidizing bacterium *Nitrobacter*
winogradskyi Nb-255. *Appl Environ Microbiol* 2006; 72:2050–63.
80. Lücker S, Wagner M, Maixner F, Pelletier E, Koch H, Vacherie B, et al. A *Nitrospira*
metagenome illuminates the physiology and evolution of globally important nitrite-oxidizing
bacteria. *Proc Natl Acad Sci USA* 2010; 107:13479–84.
81. Lücker S, Nowka B, Rattei T, Spieck E, Daims H. The genome of *Nitrospina gracilis*
illuminates the metabolism and evolution of the major marine nitrite oxidizer. *Front*
Microbiol 2013; 4:27.
82. Mundinger AB, Lawson CE, Jetten MSM, Koch H, Lücker S. Cultivation and transcriptional
analysis of a canonical *Nitrospira* under stable growth conditions. *Front Microbiol* 2019;
10:1325.
83. Sancar A. Structure and function of DNA photolyase and cryptochrome blue-light
photoreceptors. *Chem Rev* 2003; 103:2203–38.
84. Zadow A von, Ignatz E, Pokorny R, Essen L-O, Klug G. *Rhodobacter sphaeroides* CryB is
a bacterial cryptochrome with (6–4) photolyase activity. *FEBS J* 2016; 283:4291–309.
85. Zhang F, Scheerer P, Oberpichler I, Lamparter T, Krauß N. Crystal structure of a
prokaryotic (6-4) photolyase with an Fe-S cluster and a 6,7-dimethyl-8-ribityllumazine
antenna chromophore. *Proc Natl Acad Sci USA* 2013; 110:7217–22.
86. Maresca JA, Keffer JL, Hempel PP, Polson SW, Shevchenko O, Bhavsar J, et al. Light
modulates the physiology of nonphototrophic Actinobacteria. *J Bacteriol* 2019; 201:
e00740-18.
87. Wink DA, Kasprzak KS, Maragos CM, Elespuru RK, Misra M, Dunams TM, et al. DNA
deaminating ability and genotoxicity of nitric oxide and its progenitors. *Science* 1991;
254:1001–3.
88. Imlay JA. The molecular mechanisms and physiological consequences of oxidative stress:
lessons from a model bacterium. *Nat Rev Microbiol* 2013; 11:443–54.
89. Zhang T, Hansel CM, Voelker BM, Lamborg CH. Extensive dark biological production of
reactive oxygen species in brackish and freshwater ponds. *Environ Sci Technol* 2016;
50:2983–93.
90. Khademian M, Imlay JA. *Escherichia coli* cytochrome *c* peroxidase is a respiratory oxidase
that enables the use of hydrogen peroxide as a terminal electron acceptor. *Proc Natl Acad*
Sci USA 2017; 114:E6922–31.
91. O'Reilly CM, Sharma S, Gray DK, Hampton SE, Read JS, Rowley RJ, et al. Rapid and
highly variable warming of lake surface waters around the globe. *Geophys Res Lett* 2015;
42:10773-10781.
92. Sterner RW, Keeler B, Polasky S, Poudel R, Rhude K, Rogers M. Ecosystem services of
Earth's largest freshwater lakes. *Ecosystem Services* 2020; 41:101046.
93. Murphy TP. Ammonia and nitrate uptake in the lower Great Lakes. *Can J Fish Aq Sci*
1980; 37:1365–72.
94. Gardner WS, Lavrentyev PJ, Cavaletto JF, McCarthy MJ, Eadie BJ, Johengen TH, et al.
Distribution and dynamics of nitrogen and microbial plankton in southern Lake Michigan
during spring transition 1999-2000. *J Geophys Res* 2004; 109: C03007.
725

Figure Legends

Figure 1. Dissolved inorganic nitrogen availability and distribution of nitrifiers across the Great Lakes. **(a)** Oxidized nitrogen concentrations. Values include NOx concentrations from published studies (n=128; ref. 16,36,38,93,94), US EPA Water Quality Surveys in 2012 and 2013 (n=1626 from GLENDA database), and this study (n=20) (Table S2). **(b)** Ammonium concentrations. Values are derived from the literature as in (a) (n=118) and from this study (n=20) (Table S3). **(c)** Distribution of nitrifiers across the Great Lakes. Top panel: map of sampling stations. Stars indicate stations chosen for metagenome analysis. Bottom panel: relative abundance of ammonia-oxidizing (green) and nitrite-oxidizing (pink) families based on 16S rRNA V4-V5 amplicon sequencing, sampled in the mid-hypolimnion (except western Erie, sampled 1m from bottom). Data is plotted roughly west to east as indicated on the map.

Figure 2. Genome properties and cross-lake distribution of ammonia oxidizing organisms, showing both Archaea (top) and Betaproteobacteria (bottom). Rows highlighted in green represent clusters of genomes reconstructed from the Great Lakes; expanded phylogenetic trees are shown in Figures S2 and S5. For catalase, "E" indicates monofunctional catalase *katE*; "G" indicates bifunctional catalase-peroxidase *katG*. For carbon fixation, RuBisCO type is shown in parentheses (68); HP/HB, 3-Hydroxypropionate/4-Hydroxybutyrate cycle; CBB, Calvin-Benson-Bassham cycle. Bubble plot shows composition of ammonia oxidizers in hypolimnion samples, using MAGs as probes to recruit metagenomic reads (values sum to 100% for each lake column). Genes identified in only a subset of genomes are shown as (+).

Figure 3. Evidence for proteorhodopsin (PR) in *Nitrosospira* from the Great Lakes. (a) Gene neighborhood surrounding PR in *Nitrosospira* MAG MC17_S15_bin_110 and SAG 207399. Genes are colored according to the best BLAST hit taxonomy in the NCBI nr database. (b) Alignment of predicted *Nitrosospira* PR with reference sequences. Diagnostic features are

highlighted (61,62): blue boxes, diagnostic residues for PR; pink box, residue indicative of blue

or green tuning; green underline, shorter beta-sheet region in PR. Sequence accession

numbers: bacteriorhodopsin P02945, actinorhodopsin A0A1D9E0H1, sensory rhodopsin

P42196, *Dokdonia* PR EAQ40507.1, SAR86 blue-tuned PR Q4PP54, SAR86 green-tuned PR

Q9F7P4, SAR11 PR A6YQL7.

Figure 4. Genome properties and cross-lake distribution of nitrite oxidizing taxa *Nitrospira* (top)

and *Ca.* Nitrotoga (Betaproteobacteria*;* bottom). Rows highlighted in pink represent clusters of

genomes reconstructed from the Great Lakes. Expanded phylogenetic trees are shown in

Figures S2 and S6. rTCA, reductive tricarboxylic acid cycle; CBB, Calvin-Benson-Bassham

cycle; ONR, octaheme nitrite reductase. Values in parentheses indicate RuBisCO type (68).

Bubble plot shows composition of NOB per lake based on metagenomic read mapping. Genes

identified in only a subset of genomes are indicated by (+).

Figure 5. Distinct photolyase proteins in NspiraGL. **a)** Phylogenetic tree showing families of

photolyases. Three families are found in NspiraGL: CPD class I photolyase, FeS-BCP / CRYPro

family, and an uncharacterized CPF-related family found in diverse Bacteria. CPD Class I

photolyases are also found in other nitrifiers including *Ca*. Nitrotoga NtogaGL1a, *Nitrosospira*

NspGL1, and *Nitrosoarchaeum* NarchGL. **b)** Domain structure of the three photolyase families

present in NspiraGL.

Acknowledgments

Sequencing support for this project was provided by the DOE Joint Genome Institute

Community Sequencing Program (CSP #1565, #503460). Computational resources were

provided by the UChicago Research Computing Center. Funding for this work was provided by

Illinois-Indiana Sea Grant (Grant # NA14OAR170095), UChicago Women's Board, and the

National Science Foundation (OCE-1830011 to MLC). We thank Glenn Warren and the science

staff in the Great Lakes National Program Office of the US EPA, and the captain and crew of the

R/V *Lake Guardian*, for facilitating sample collection. We thank members of the Coleman and

Waldbauer labs for assistance with sample collection and processing, and for discussion and

comments on the manuscript.

Data Availability

The metagenome-assembled genomes presented here are available via NCBI BioProject

PRJNA636190. 16S rRNA data is available at NCBI BioProject PRJNA591360. Metagenomes

sequenced by JGI are available at http://genome.jgi.doe.gov, Project IDs 1045056, 1045059,

1045062, 1045065, 1045068, 1045071. Single cell amplified genome is available at

img.jgi.doe.gov, IMG Genome ID 3300033241.

Competing Interests

The authors declare no competing interests.

Figure 1

Figure 2

Figure 3

a

b

	90	100	110	120	130	140
Bacteriorhodopsin	A I A F T M Y L S M L L G Y G L T M V P F G G E Q N P I Y W A - - - R Y A D W L F T T P L L L D - - -
Actinorhodopsin	A I A A Y H Y F R M F D S F N H - - A F A G A E G N P D L Y N V G Y R Y V D W F L T V P L L L V E T V A
Sensory rhodopsin	G I A A V A Y V V M A L G V G W - - V P V A E - - - R T V F A P - - R Y I D W I L T T P L I V - - - - Y
Dokdonia PR	F I A A V H Y W Y M R D Y - - W - - Q A N A E - - S P T F F - - - - R Y V D W V L T V P L M C V E F F L
SAR86 PR (blue-tuned)	G V A F W H Y L Y M R G V - - W - - I Y A G E - - T P T V F - - - - R Y I D W L I T V P L Q I I E F Y L
SAR86 PR (green-tuned)	G I A F W H Y M Y M R G V - - W - - I E T G D - - S P T V F - - - - R Y I D W L L T V P L L I C E F Y L
SAR11 PR	G I A F I H Y M Y M R D V - - W - - V M T G E - - S P T V Y - - - - R Y I D W L I T V P L L M L E F Y F
Type 1 *Nitrosospira* PR	L V A A V H Y F Y M R D V - - W - - V A T G S - - T P T V Y - - - - R Y I D W L I T V P L L M I E F Y L

Figure 4

		% GC	Genome size (Mb)	Coding density (%)	# Sigma factors	Photolyase	Catalase	Formate dehydrogenase	NiFe hydrogenase	Thiosulfate dehydrogenase (*tsdA*)	Assim. nitrite reduction	*nirK*	*norQD*	# *nxrAB* copies	Cyanase	Urease	Carbon fixation pathway	Distribution across lakes
Nitrospira	NspiraGL	58	1.8	89	5	+++	G	-	-	-	*nirA*	-	+	1	+	(+)	rTCA	
	Nitrospira japonica NJ1	58	4.1	90	27	+	E	-	-	-	ONR	+	+	3	+	+	rTCA	
	Nitrospira moscoviensis NSP M-1	61	4.5	89	18	+	EG	+	+	+	ONR	+	+	5	+	+	rTCA	
	Nitrospira defluvii	59	4.3	89	17	+	-	+	-	+	*nirA*	+	+	2	+	-	rTCA	
Betaproteo.	NtogaGL1a	51	1.4	91	4	(+)	G	-	-	-	-	-	-	1	+	-	CBB (IC)	
	NtogaGL1b	52	1.5	91	4	-	G	-	-	+	-	-	-	1	+	-	CBB (IC)	
	Ca. Nitrotoga CP45	49	2.8	85	8	+	EG	-	+	-	*nirBD*	+	+	1	-	-	CBB (IC, ID/IA)	
	Ca. Nitrotoga KNB	50	2.6	85	6	+	G	-	+	-	*nirBD*	+	+	2	-	-	CBB (IC, ID/IA)	

N physiology

Relative abundance
25 50 75

Figure 5

a

b

Supplementary Information

Extended Methods

<u>Metagenome QC and Assembly</u>

Raw reads for spring surface samples were quality controlled at the Joint Genome Institute,
using bbduk.sh for adapter trimming (ktrim=r, minlen=40, minlenfraction=0.6,mink=11, tbo, tpe,
k=23, hdist=1, hdist2=1, ftm=5) and quality filtering (maq=8, maxns=1, minlen=40,
minlenfraction=0.6, k=27, hdist=1, trimq=12, qtrim=rl). Raw reads for summer hypolimnion
samples were adapter trimmed, quality filtered, and interleaved using bbduk (parameters:
ktrim=r, mink=8, hdist=2, k=21, forcetrimleft=10, forcetrimright=199, minlen=150) using BBTools
suite version 35.74 (https://sourceforge.net/projects/bbmap/). Separate assemblies of quality
filtered reads were carried out for each metagenome using metaSPAdes 3.1.11 --meta mode
using default k sizes of 21, 33, 55 (1). For mapping, forward and reverse reads were merged
using bbmerge in BBtools, using qtrim2=r trimq=10,13,16 and adapter=default. Merged short
reads from all 12 metagenomes were mapped onto each assembly using bowtie2 in --sensitive
mode (2), and together this information was used to bin assembled contigs. Binning was
performed using MetaBAT2 (3), Binsanity (4) and CONCOCT (5) using default parameters. The
resulting bins were scored, aggregated, and de-replicated using DAS_Tool (6) followed by
manual curation using Anvi'o (7). We assessed genome completion and contamination of
manually curated bins using CheckM lineage_wf (8), and all genomes presented here are
greater than 70% complete and have less than 10% contamination (Table S1). Potential
nitrifiers were screened by searching for ammonia monooxygenase, hydroxylamine
oxidoreductase and nitrite oxidoreductase within reconstructed genomes using blastp (9). For
bins where any of these genes were detected, we identified bacterial single copy core genes
(10) or archaeal single copy core genes (11) using HMMER (12), as implemented in Anvi'o.
Single copy core genes were queried against proteins predicted from bacterial and archaeal

genomes in RefSeq (13), and taxonomic identity of these core genes was ascertained based on a least common ancestor approach using a 0.1% window around the bit score of the best hit using KronaTools (14). Taxonomic assignment was further validated using GTDB-tk (15). Grouping of MAGs into clades and subclades based on ANI was carried out using fastANI (16).

Annotation and gene clustering

Genes from putative nitrifier genomes were annotated using InterProScan 5.30-69.0 (17) and eggnog-mapper against the bactNOG database (18). Gene cluster analysis was carried out using the Anvi'o pangenome pipeline (19), using blastp to determine sequence similarity, ITEP to eliminate weak similarity (20) and MCL to cluster, using a minbit of 0.5, MCL inflation of 2 and minimum gene occurrence of 1 (21). Assignment of KEGG Orthology to gene clusters was carried out using GhostKOALA (22). Sigma factors were tallied by identifying gene clusters annotated with the following PFAMs: PF00309, PF03979, PF00140, PF04542, PF04539, PF04545, PF08281. Prokka (23) was used to generate GenBank file format files from MAGs and SAGs, and genoPlotR (24) was used to generate gene neighborhood maps

Gene tree construction

The NspGL1 proteorhodopsin sequence was inserted into the MicRhoDE rhodopsin tree using pplacer (25) through the MicRhoDE Galaxy pipeline (26). We then constructed a more targeted phylogenetic tree using aligned reference sequences of Supercluster III from MicRhoDE, filtered to exclude fragments shorter than 220 amino acids. To this alignment, we added NspGL1 sequences using MAFFT (27) along with high similarity sequences from NCBI nr that were not present in MicRhoDE. The tree was inferred using RaxML with model PROTGAMMALG (28). The tree was visualized in iTOL (29) and more distant clusters were collapsed for clarity.

A cyanase phylogenetic tree was created using sequences drawn from querying NtogaGL cyanase against NCBI nr using blastp, as well as sequences from references (30–32).

Sequences were aligned using MAFFT (27) and the tree was inferred using RaxML with model PROTGAMMALG (28). Tree was visualized in iTOL (29) and branches were colored based on the taxonomy of the parent genome.

Photolyase-related proteins in GL MAGs were identified by searching for the following features: K01669, COG0415, PF03441, PF00875, PF04244, SSF48173, SSF52425. Reference proteins (n=56) spanning the previously defined families of photolyases and cryptochromes (33) were obtained from UniProt, along with aquatic bacterial sequences described by Maresca and colleagues (34). The reference sequences were aligned using MAFFT (27), and sequences from GL MAGs were added using the MAFFT --addfragments option. The tree was estimated using IQ-TREE 2 (35) and visualized using iTOL (29).

Phylogenomic tree construction

All *Nitrospirae*, *Thaumarchaeota*, *Gallionellaceae* and *Nitrosomonadaceae* genomes were downloaded from Genbank (36) and included in phylogenetic trees for their respective family. Single copy core genes were extracted as described above and individually aligned at the protein level using muscle (37). Core proteins were then concatenated for each genome. Concatenated alignments were trimmed using Gblocks (38) and analyzed by RAxML 8.2.12 (28) to create a phylogenetic tree using the PROTGAMMALG model and 50 bootstraps. Trees were visualized in iTOL (29).

Proteorhodopsin assembly verification

We used several approaches to validate the presence of proteorhodopsin in assembled *Nitrosospira* genomes, to rule out the possibility of chimeric assemblies from different species. We note that proteorhodopsin-containing contigs were independently assembled and binned together with core *Nitrosospira* contigs from seven different samples (i.e. each sample was

assembled and binned separately, rather than co-assembled). In five of seven cases,

proteorhodopsin and retinal biosynthesis genes were assembled together with core *Nitrosospira*

genes on the same contig. To rule out a systematic reproducible error in assembly and/or

binning, we compared these seven MAGs to a single cell amplified *Nitrosospira* genome (SAG)

from Lake Michigan, obtained as part of another project with the JGI. This SAG was processed

through JGI's standard decontamination pipeline and manually investigated to ensure lack of

contamination. We found no evidence of contaminating core genes, as all core genes had best

hits to either *Nitrosospira* or more generally *Nitrosomonadaceae* in nr. SAG contigs were

matched to homologous contigs from NspGL1 MAGs to determine if any SAG contigs were

unique using FastANI (16) with --visualize flag. All contigs from this *Nitrosospira* SAG were

found within an NspGL1 MAG. Bandage (39) was used to manually inspect the assembly graph

around the contig that contained the NspGL1 *Nitrosospira* proteorhodopsin to ensure that the

assembled contig did not represent a chimeric contig or inappropriate scaffolding. We verified

that a single, unique path exists from the beginning to the end of the NspGL1 contig containing

proteorhodopsin (Fig 3, Fig S4). Further, we verified that consistent coverage across this contig

existed by mapping short reads from the original sample using bowtie2 (2) and viewing results

using Integrated Genomics Viewer (40) A closely related assembly of the same genomic region

from Lake Biwa did not show evidence of proteorhodopsin; to confirm this difference between

the Biwa and Great Lakes MAGs, we mapped reads from Lake Biwa (41) (BioProject

PRJDB6644) onto the assembled contig described above (Fig 3, Fig S4) using bowtie2 (2)**.** This

analysis demonstrated that while a large fraction of the NspGL1 contig in question recruited

reads from Lake Biwa at high identity (98-99%), starting upstream of proteorhodopsin and

retinal biosynthesis, this contig no longer recruited reads from Lake Biwa.

<u>Manual identification of key nitrification genes</u>

Despite recovery of 15 high completion MAGs in NspGL1/2a/2b/3, many of these MAGs

lacked key nitrification genes in *amo* and *hao* operons. This was largely due to the fact that *amo* and *hao* operons were often assembled on small contigs below the minimum size cutoff we imposed for binning contigs. Difficulty in assembling these contigs was likely in part due to the several *amo* and *hao* operons with extremely high identity to one another in each genome, a phenomenon which has been observed in other *Nitrosospira* (42). Manual assembly graph inspection with Bandage (39) supported this hypothesis, as did assessment of abundance of short reads associated with *amo* operons from NspGL and comparison of abundance of short reads associated with core gene *rpoB* from NspGL, using ROCker (43). Still, an exemplar MAG from at least one representative of each ecotype (NspGL1/2a/2b/3) was found with both *amo* and *hao* operons. Further, manual inspection of unbinned contigs confirmed that *amo* and *hao* operons existed on contigs in every sample from which a MAG for a particular ecotype was recovered. That is, for every time that an NspGL1 MAG was recovered from a sample, we were able to determine that an *amo* and *hao* operon which could be affiliated with NspGL1 existed, even if it was not correctly binned. Affiliation for these unbinned key nitrification genes was carried out by alignment of *amoAB* and *haoAB* sequences to *amoAB* and *haoAB* sequences correctly binned in NspGL ecotypes. This process was also carried out for two NtogaGL1a MAGs for *nxrAB*, which were poorly assembled in those two samples.

Data analysis and plots

All data were analyzed in R version 3.5.3 (44) and plots were generated using ggplot2 (45).

Figure S1. Relationships between environmental factors and relative abundance of nitrifiers in the Great Lakes. Relative abundance based on 16S rRNA V4-V5 region amplicon sequencing from Summer 2012. Nitrifier abundance includes sequences assigned to the familes *Nitrosopumilaceae* (AOA), *Nitrosomonadaceae* (AOB), *Nitrospiraceae* (NOB), and *Gallionellaceae* (NOB). (a) Relative abundance of nitrifiers (% of total community) compared to photosynthetically active radiation (PAR). (b) Correlation between the relative abundances of nitrite oxidizers and ammonia oxidizers in each sample. A 1:1 line (solid) and linear regression fit (dashed) are shown for reference. (c) Relative abundance of nitrifiers (% of total community) compared to chlorophyll *a* concentration. PAR and chlorophyll data from the EPA Great Lakes Environmental Database (https://cdx.epa.gov/).

Figure S2. Phylogenomic tree of the *Nitrosomonadales* order of *Betaproteobacteria*, focusing on the *Nitrosomonadaceae* and *Gallionellaceae* families. Tree was constructed using 139 concatenated single-copy core genes (10). Colored squares indicate the environment of origin; colored circles denote the lake of origin for genomes from the Laurentian Great Lakes; yellow stars indicate clades containing strains that lack nitrosocyanin.

Figure S3. Proteorhodopsin phylogeny. Top: Phylogenetic placement of *Nitrosospira* PR. Rhodopsin sequences and classifications from MicRhoDE database (26); star shows NspGL1 PR. Bottom: Close relatives of *Nitrosospira* PR within Supercluster III. Neighboring clusters have been collapsed for clarity.

Figure S4. Shared synteny surrounding the proteorhodopsin module in *Nitrosospira* draft genomes from deep lakes. Light blue, shared genes on reverse strand; dark blue, shared genes on forward strand; pink, tRNA genes; orange, PR module in NspGL1 MAG; gray, genes unique to one MAG.

Ca. Cenarchaeum sp. (NP-γ-1)

Nitrosopumilaceae PNGco C binSS26
Nitrosopumilaceae IBTHAUMO2
Nitrosopumilus sp H8
Nitrosopumilus sp H13
Nitrosopumilus sp B06
Nitrosopumilus sp D6
Nitrosopumilaceae archaeon bin 79
Nitrosopumilus sp SZUA 465
Nitrosopumilales archaeon CG15 37_12
Ca. Nitrosoarchaeum limnia SFB1
Ca. Nitrosoarchaeum limnia BG20
Nitrosoarchaeum koreense MY1
Nitrosopumilus sp CG10 33_7
Nitrosopumilales archaeon CG15 33_23
Nitrosopumilales archaeon CG 34_10
Nitrosopumilales archaeon CG 34_8
Nitrosopumilales archaeon CG11 33_24
Nitrosopumilaceae F1-20-MAGs118
Nitrosopumilaceae F1-60-MAGs143
Nitrosopumilaceae F1-20-MAGs117
Nitrosopumilaceae F1-80-MAGs016
Nitrosopumilaceae F1-120-MAGs239
Nitrosopumilaceae F1-60-MAGs144
Nitrosopumilaceae F1-140-MAGs129
Nitrosoarchaeum sp GW199 bin 3
Nitrosoarchaeum sp Baikal deep G180
Nitrosoarchaeum sp BD3
Nitrosopumilaceae PowLak16 MAG35
Nitrosoarchaeum sp Baikal deep G181
Nitrosoarchaeum sp Baikal G1
MC17 S15 bin 49
ON33M 5M MetaBAT2 044   NarchGL
SU08M 6M MetaBAT2 048
Nitrosopumilus sp PNGco_C_binVS21
Nitrosopumilus sp PNGco_S_binVS9
Nitrosopumilus sp 32_1
Nitrosopumilus sp AR
Ca. Nitrosopumilus koreensis AR1
Nitrosopumilus sp SJ
Nitrosopumilus piranensis D3C
Marine Group I thaumarchaeote SCGC_AAA799_P11
Marine Group I thaumarchaeote SCGC_AAA799_B03
Marine Group I thaumarchaeote SCGC_AAA799_D11
Marine Group I thaumarchaeote SCGC_AAA799_N04
Marine Group I thaumarchaeote SCGC_AAA799_E16
Marine Group I thaumarchaeote SCGC_RSA3
Ca. Nitrosopumilus sp SW
Nitrosopumilus maritimus SCM1
Nitrosopumilus sp GoM MAG6
Nitrosopumilus sp GoM MAG5
Nitrosopumilus sp GoM MAG3
Nitrosopumilus sp 33_1
Nitrosopumilus sp GoM MAG4
Thaumarchaeota archaeon casp thauma3
Thaumarchaeota archaeon SCGC AAA282_K18
Thaumarchaeota archaeon NP142
Nitrosopumilus sp SZUA 238
Nitrosopumilus sp LS_AOA
Thaumarchaeota archaeon NORP47
Ca. Nitrosopumilus sp NM25
Nitrosopumilus adriaticus NF5
Ca. Nitrosopumilus sp DDS1
Ca. Nitrosopumilus salaria BD31
Nitrosopumilus salaria SZUA 335
Nitrosopumilales archaeon CG 34_16
Nitrosopumilus sp das_tool_concoct_135
Ca. Nitrosopumilus sediminis AR2
Nitrosopumilus sp UBA526
Nitrosopumilus sp UBA527
Thaumarchaeota archaeon Guaymas 69
uncultured marine group I AG_319_N20
Nitrosopumilus sp BACL13 MAG-121220-bin23
Nitrosopumilus sp UBA241
Nitrosopumilus sp BACL13 MAG-120910-bin56
uncultured marine group I AG_319_O22
Ca. Nitrosomarinus catalina SPOT01
Nitrosopumilus sp Nsub
Nitrosomarinus sp GoM MAG2
Nitrosopumilus sp MED740
Nitrosopumilus sp MED_G94
Thaumarchaeota archaeon Baikal G2
Nitrosopumilus sp Baikal deep G182
Thaumarchaeota archaeon casp thauma1
Nitrosopumilaceae F1-20-MAGs116
Nitrosopumilaceae F1-140-MAGs130
Nitrosopumilaceae F1-120-MAGs240
Nitrosopumilaceae F1-80-MAGs017
Nitrosopumilaceae F1-60-MAGs142

Ca. Nitrosoarchaeum sp. (NP-γ-2.2)

Nitrosopumilus sp. (NP-γ-2.1)

low salinity

low salinity

Environment
- freshwater
- brackish
- brackish sediment
- marine
- marine sediment
- fish tank
- groundwater
- soil
- sponge symbiont

Great Lakes
- Superior
- Michigan
- Huron
- Erie
- Ontario

Tree scale: 0.1   10

Figure S5. Phylogenomic tree of *Nitrosopumilaceae* and *Cenarchaeum*. Tree was constructed using concatenated single-copy core genes as defined in (11). This tree represents a zoomed in view of *Nitrosopumilaceae* and *Cenarchaeum* from larger tree of all *Thaumarchaeota* and *Aigarchaeota*. Colored boxes correspond to the environment of origin; genomes from the Laurentian Great Lakes are indicated by colored circles. Names in bold indicate cultured isolates or enrichments.

Figure S6. Phylogenomic tree of *Thaumarchaeota* and *Aigarchaeota*. Tree was constructed using concatenated single-copy core genes from (11), and all available genomes in Genbank as of February 6 2020. Colors denote clades described in (46). Star indicates placement of NarchGL.

Figure S7. Cyanase diversity and distribution among nitrite oxidizing bacteria. a) Phylogenetic tree showing distinct clades of cyanase, highlighting Great Lakes *Nitrospira* and *Ca.* Nitrotoga. b) Gene synteny surrounding the cyanase gene in NspiraGL and reference *Nitrospira*, and a cyanase insertion in NtogaGL compared to reference *Ca.* Nitrotoga.

Figure S8. Gene content distinguishing NtogaGL1a and NtogaGL1b. Genes colored in blue are shared between both genomes; orange colored genes are unique to either genome.

Figure S9. Phylogenomic tree of Lineage I and II *Nitrospira*. Tree was constructed using concatenated single-copy core genes described in (10). Colored boxes correspond to the environment of origin; genomes from the Laurentian Great Lakes are indicated by circles color coded by lake of origin.

Figure S10. Gene neighborhood surrounding photolyase genes in NspiraGL, compared to reference *N. moscoviensis*. Genes shared between the two genomes are colored light blue (protein coding) or magenta (rRNA, tRNA); non-shared genes are gray; photolyase genes are colored to match the tree in Figure 5.

References

1. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. metaSPAdes: a new versatile metagenomic assembler. *Genome Res* 2017; 27:824–34.
2. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 2012; 9:357–9.
3. Kang DD, Froula J, Egan R, Wang Z. MetaBAT, an efficient tool for accurately reconstructing single genomes from complex microbial communities. *PeerJ* 2015; 3:e1165.
4. Graham ED, Heidelberg JF, Tully BJ. BinSanity: unsupervised clustering of environmental microbial assemblies using coverage and affinity propagation. *PeerJ* 2017; 5:e3035.
5. Alneberg J, Bjarnason BS, de Bruijn I, Schirmer M, Quick J, Ijaz UZ, et al. Binning metagenomic contigs by coverage and composition. *Nat Methods* 2014; 11:1144–6.
6. Sieber CMK, Probst AJ, Sharrar A, Thomas BC, Hess M, Tringe SG, et al. Recovery of genomes from metagenomes via a dereplication, aggregation and scoring strategy. *Nat Microbiol* 2018; 3:836–43.
7. Eren AM, Esen ÖC, Quince C, Vineis JH, Morrison HG, Sogin ML, et al. Anvi'o: an advanced analysis and visualization platform for 'omics data. *PeerJ* 2015; 3:e1319.
8. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 2015; 25:1043–55.
9. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+: architecture and applications. *BMC Bioinform* 2009; 10:421.
10. Campbell BJ, Yu L, Heidelberg JF, Kirchman DL. Activity of abundant and rare bacteria in a coastal ocean. *Proc Natl Acad Sci USA* 2011; 108:12776–81.
11. Rinke C, Schwientek P, Sczyrba A, Ivanova NN, Anderson IJ, Cheng J-F, et al. Insights into the phylogeny and coding potential of microbial dark matter. *Nature* 2013; 499:431–7.
12. Mistry J, Finn RD, Eddy SR, Bateman A, Punta M. Challenges in homology search: HMMER3 and convergent evolution of coiled-coil regions. *Nucleic Acids Res* 2013; 41:e121–e121.
13. O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, et al. Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res* 2016; 44:D733–45.
14. Ondov BD, Bergman NH, Phillippy AM. Interactive metagenomic visualization in a Web browser. *BMC Bioinform* 2011; 12:385.
15. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database. *Bioinformatics* 2020; 36:1925-1927.
16. Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun* 2018; 9:1–8.
17. Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, et al. InterProScan 5: genome-scale protein function classification. *Bioinformatics* 2014; 30:1236–40.
18. Huerta-Cepas J, Forslund K, Coelho LP, Szklarczyk D, Jensen LJ, von Mering C, et al. Fast genome-wide functional annotation through orthology assignment by eggNOG-Mapper. *Mol Biol Evol* 2017; 34:2115–22.
19. Delmont TO, Eren AM. Linking pangenomes and metagenomes: the *Prochlorococcus* metapangenome. *PeerJ* 2018; 6:e4320.
20. Benedict MN, Henriksen JR, Metcalf WW, Whitaker RJ, Price ND. ITEP: An integrated toolkit for exploration of microbial pan-genomes. *BMC Genomics* 2014; 15:8.
21. van Dongen S, Abreu-Goodger C. Using MCL to extract clusters from networks. *Methods Mol Biol* 2012; 804:281–95.
22. Kanehisa M, Sato Y, Morishima K. BlastKOALA and GhostKOALA: KEGG tools for

functional characterization of genome and metagenome sequences. *J Mol Biol* 2016; 428:726–31.

23. Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 2014; 30:2068–9.

24. Guy L, Roat Kultima J, Andersson SGE. genoPlotR: comparative gene and genome visualization in R. *Bioinformatics* 2010; 26:2334–5.

25. Matsen FA, Kodner RB, Armbrust EV. pplacer: linear time maximum-likelihood and Bayesian phylogenetic placement of sequences onto a fixed reference tree. *BMC Bioinform* 2010; 11:538.

26. Boeuf D, Audic S, Brillet-Guéguen L, Caron C, Jeanthon C. MicRhoDE: a curated database for the analysis of microbial rhodopsin diversity and evolution. *Database (Oxford)* 2015; bav080.

27. Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res* 2002; 30:3059–66.

28. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 2014; 30:1312–3.

29. Letunic I, Bork P. Interactive tree of life (iTOL) v3: an online tool for the display and annotation of phylogenetic and other trees. *Nucleic Acids Res* 2016; 44:W242–5.

30. Pachiadaki MG, Sintes E, Bergauer K, Brown JM, Record NR, Swan BK, et al. Major role of nitrite-oxidizing bacteria in dark ocean carbon fixation. *Science* 2017; 358:1046–51.

31. Spang A, Poehlein A, Offre P, Zumbrägel S, Haider S, Rychlik N, et al. The genome of the ammonia-oxidizing *Candidatus* Nitrososphaera gargensis: insights into metabolic versatility and environmental adaptations. *Environ Microbiol* 2012; 14:3122–45.

32. Palatinszky M, Herbold C, Jehmlich N, Pogoda M, Han P, von Bergen M, et al. Cyanate as an energy source for nitrifiers. *Nature* 2015;524:105–8.

33. Vechtomova YuL, Telegina TA, Kritsky MS. Evolution of proteins of the DNA photolyase/cryptochrome family. *Biochemistry Moscow* 2020; 85:131–53.

34. Maresca JA, Keffer JL, Hempel PP, Polson SW, Shevchenko O, Bhavsar J, et al. Light modulates the physiology of nonphototrophic Actinobacteria. *J Bacteriol* 2019; 201: e00740-18.

35. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, et al. IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era. *Mol Biol Evol* 2020; 37:1530–4.

36. Benson DA, Karsch-Mizrachi I, Lipman DJ, Ostell J, Wheeler DL. GenBank. *Nucleic Acids Res* 2005; 33:D34–8.

37. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 2004; 32:1792–7.

38. Castresana J. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol Biol Evol* 2000; 17:540–52.

39. Wick RR, Schultz MB, Zobel J, Holt KE. Bandage: interactive visualization of de novo genome assemblies. *Bioinformatics* 2015; 31:3350–2.

40. Thorvaldsdóttir H, Robinson JT, Mesirov JP. Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. *Brief Bioinform* 2013; 14:178–92.

41. Okazaki Y, Nishimura Y, Yoshida T, Ogata H, Nakano S-I. Genome-resolved viral and cellular metagenomes revealed potential key virus-host interactions in a deep freshwater lake. *Environ Microbiol* 2019; 21:4740-4754.

42. Sedlacek CJ, McGowan B, Suwa Y, Sayavedra-Soto L, Laanbroek HJ, Stein LY, et al. A physiological and genomic comparison of *Nitrosomonas* cluster 6a and 7 ammonia-oxidizing bacteria. *Microb Ecol* 2019; 78:985–94.

43. Orellana LH, Rodriguez-R LM, Konstantinidis KT. ROCker: accurate detection and quantification of target genes in short-read metagenomic data sets by modeling sliding-window bitscores. *Nucleic Acids Res* 2017; 45:e14.
44. Team RC. R: A language and environment for statistical computing. R Foundation for Statistical Computing V, Austria, 2018.
45. Wickham H. *ggplot2: Elegant graphics for data analysis*. New York: Springer-Verlag; 2009.
46. Alves RJE, Minh BQ, Urich T, von Haeseler A, Schleper C. Unifying the global phylogeny and environmental distribution of ammonia-oxidising archaea based on *amoA* genes. *Nature Commun* 2018; 9:1517.