1 **An integrated comparative physiology and molecular approach pinpoints mediators of**
2 **breath-hold capacity in dolphins**
3

4 Ashley M. Blawas[1], Kathryn E. Ware[2], Emma Schmaltz[1], Larry Zheng[1], Jake Spruance[2], Austin
5 S. Allen[1], Nicole West[3], Nicolas Devos[4], David L. Corcoran[4], Douglas P. Nowacek[1,5], William
6 C. Eward[6,7], Andreas Fahlman[8,9], Jason A. Somarelli[2,7]
7
8 [1]Nicholas School of the Environment, Duke University Marine Laboratory, Beaufort, NC,
9 28516, USA.
10
11 [2]Department of Medicine, Duke University Medical Center, Durham, NC, 27710, USA.
12
13 [3]Dolphin Quest, Oahu, 5000 Kahala Ave, Honolulu, HI, 96816, USA.
14
15 [4]Duke Center for Genomic and Computational Biology, Duke University, Durham, NC, 27708,
16 USA.
17
18 [5]Pratt School of Engineering, Duke University, Durham, NC, 27708, USA.
19
20 [6]Department of Orthopaedic Surgery, Duke University Medical Center, Durham, NC, 27710,
21 USA.
22
23 [7]Duke University Medical Center, Duke Cancer Institute, Durham, NC, 27710, USA.
24
25 [8]Global Diving Research, Inc., Ottawa, ON, K2J 5E8, Canada.
26
27 [9]Research Department, Fundación Oceanogràfic de la Comunitat Valenciana, Gran Vía Marqués
28 del Turia 19, 46005 Valencia, Spain.
29
Address correspondence to: jason.somarelli@duke.edu; andreas.fahlman@duke.edu

Keywords: ischemic stress tolerance; cetaceans; diving physiology; oceans and human health

Abstract

Ischemic events, such as ischemic heart disease and ischemic stroke, are the number one cause of

death globally. Ischemia prevents blood, carrying essential nutrients and oxygen, from reaching

tissues and organ systems, leading to cell and tissue death, and eventual organ failure. While

humans are relatively intolerant to these ischemic events, other species, such as marine

mammals, have evolved remarkable tolerance to chronic ischemia/reperfusion during diving.

Here we capitalized on the unique adaptations of bottlenose dolphins (*Tursiops truncatus*) as a

comparative model of ischemic stress and hypoxia tolerance to identify molecular features

associated with breath-holding. Using RNA-Seq we observed time-dependent upregulation of the

arachidonate 5-lipoxygenase (ALOX5) gene during breath-holding. Consistent with the RNA-

Seq data, we also observed increased ALOX5 enzymatic activity in the serum of dolphins

undergoing breath holds. ALOX5 has previously been shown to be activated during hypoxia in

rodent models, and its metabolites, leukotrienes, induce vasoconstriction. The upregulation of

ALOX5 occurred within the estimated aerobic dive limit of the species, suggesting that ALOX5

enzymatic activity may promote tolerance to ischemic stress through sustained vasoconstriction

in dolphins during diving. These observations pinpoint a potential molecular mechanism by

which dolphins, and perhaps other marine mammals, have adapted to the prolonged breath-holds

associated with diving.

Introduction

<u>Ischemic stress and hypoxia are associated with negative clinical outcomes in humans.</u>

Maintenance of homeostatic function in mammalian tissues is directly dependent on a continuous

supply of oxygenated blood. Interruption of this blood supply, known as ischemia, results in

reduction in local oxygenation compared to normal physiologic levels, or hypoxia, and can lead

to inflammation and cell/tissue death (Bona et al., 1999; Choi, 1996; Eltzschig and Carmeliet,

2011; Gottlieb and Engler, 1999; Murdoch et al., 2005). Ischemia is the causative factor in

multiple clinical settings, and ischemic heart disease is the number one cause of death globally,

accounting for over 9 million deaths each year (Nowbar et al., 2019; World Health Organization,

2018).

<u>Marine mammals have evolved tolerance to ischemic stress.</u> While humans have little tolerance

for ischemic stress and hypoxia, a number of species have evolved unique physiologies that

allow them to seemingly thrive despite regular tissue-level ischemia and low-oxygen

environments. One group of animals that undergo repeated daily ischemic events is marine

mammals. During a dive, a marine mammal experiences a suite of cardiovascular changes that

aid in reducing aerobic metabolism (Irving et al., 1941; Scholander, 1940). As part of this

response, both heart rate (f_H) and stroke volume decrease, resulting in reduced cardiac output

(Fahlman et al., 2020b, 2019b). Increased peripheral resistance, through selective

vasoconstriction, helps assure that mean arterial blood pressure is maintained, at least in studies

on forced diving in seals (Blix et al., 1976; Zapol et al., 1979). Ultimately, this response

conserves oxygen in the blood and lungs for oxygen-sensitive tissues like the brain and the heart,

while the skeletal muscles rely on endogenous myoglobin-bound oxygen for aerobic metabolism

Figure 1. Dolphins as a model of ischemia. A. Dolphins and other cetaceans have increased oxygen stores that are reapportioned compared to humans. Oxygen store data were reported in Ponganis et al., 2011 (human) and Kooyman and Ponganis, 2018 (dolphin). B. The enhanced oxygen stores and diving capacity of dolphins makes them a unique model to study ischemic stress tolerance. Dive data and calculated aerobic dive limit (cADL) were reported by AIDA and Foster and Sheel, 2005 (human) and Fahlman et al., 2018 (dolphin).

(Davis and Kanatous, 1999; Fahlman et al., 2009). While these responses to submersion in water

are largely conserved across all vertebrates, many of the physiological adaptations that support

diving are exaggerated in marine mammals compared to other taxa (Kooyman and Ponganis,

1998; Panneton, 2013) (**Figure 1**). For example, maintenance of

increased peripheral resistance does not appear to occur in human breath-hold divers, as mean

arterial blood pressure increases with dive duration (Breskovic et al., 2011; Gooden, 1994;

Taboni et al., 2019). These physiological differences highlight the tremendous potential to study

marine mammals as model organisms for the investigation of adaptations to ischemic and

hypoxic stress tolerance, and the cardiorespiratory plasticity that helps prevent hypertension

(Blawas et al., 2021; Fahlman et al., 2019b, 2020b).

<u>Marine mammals have evolved molecular adaptations to ischemic stress tolerance.</u> Increasing

attention has been paid to the defenses marine mammals possess against the oxidant by-products

and inflammation associated with ischemic, hypoxia, and reperfusion at the molecular level

(Allen and Vázquez-Medina, 2019; Hindle, 2020; Zhu et al., 2018). Using phylogenetic and

evolutionary convergence approaches, several gene families have been identified to contribute to

the increased ischemic stress tolerance of marine mammals, including hypoxia-inducible factor 1

(HIF-1) (Bi et al., 2015; Johnson et al., 2005, 2004), genes relating to the glutathione system and

peroxiredoxins (Bagchi et al., 2018; Tift et al., 2014; Yim et al., 2014; Zhou et al., 2018), and

several genes linked to oxygen storage, particularly hemoglobin and myoglobin (Mirceta et al.,

2013; Nery et al., 2013; Tian et al., 2017, 2016). Yet, few studies have examined differential

gene expression in marine mammals under conditions of ischemia and hypoxia (i.e. diving

conditions).

98   Here, we investigate the dynamic molecular changes that occur during an apnea in

bottlenose dolphins using genomic analysis of peripheral blood mononuclear cells (PBMCs) and

serum sampled at regular intervals during breath-holds. We couple these analyses with

previously-published f_H measurements from the same dolphins to understand how the timing of

molecular changes relates to the physiologic dive response (Blawas et al., 2021; Fahlman et al.,

2019b, 2020b). Our integrated analyses pinpoint a gene regulatory network centered around the

arachidonate 5-lipoxygenase (ALOX5) gene and its downstream metabolites, leukotrienes, as

differentially activated during breath-holding. This activation of ALOX5 is consistent with

cardiovascular control through reduction in f_H and peripheral vasoconstriction to efficiently

manage O_2 use during diving. Based on our collective results we propose a model in which the

ALOX5 pathway is upregulated during extended breath-holds as a mechanism to sustain

vasoconstriction and maintain oxygen stores for critical organs while diving.

Materials and Methods

Data collection and animal information. Four adult male bottlenose dolphins (*Tursiops*

truncatus) housed at Dolphin Quest Oahu (Honolulu, HI, USA) with an average (± S.D.) age of

22.8±9.9 years (range = 11 – 35 years) and body mass of 198.1±42.9 kg (range = 147.0 – 251.7

kg, **Table 1**) participated in this study. All data were collected under voluntary participation and

the animals could end a trial at any time. Routine veterinary assessments include venous blood

sampling, and the dolphins that participated in this study had previously been desensitized to the

blood sampling protocol. The study protocols were accepted by Dolphin Quest and the Animal

Care and Welfare Committee at the Oceanogràfic (OCE-17-16, amendments OCE-29-18 and

OCE-3-19i).

Table 1. Animal ID, age (years), body mass (kg), and included analyses for all dolphins in the study.

Animal ID	Age (years)	Body Mass (kg)	RNA-Seq	Lipoxygenase assay
6JK5	24	200.9	x	x
9FL3	35	251.7	x	
9ON6	21	192.8	x	x
83H1	11	147.0		x
Mean±S.D.	22.8±9.9	198.1±42.9		

Experimental trials. Serum was isolated from whole blood samples at baseline, 3 minutes, and 4

½ - 5 minutes of breath-holding on fasted dolphins at Dolphin Quest, Oahu, March 2018 and

May 2019. All trials were performed in the morning, when the animals were fasted with at least

12 hours having passed since the last meal on the previous day to minimize the potential

confounding effect of nutritional state. To ensure that the samples were collected during resting

behavior each breath-hold was proceeded by 2 minutes of rest or slow swimming at the surface.

A trial was initiated when the dolphin rolled into dorsal recumbency with its blowhole

submerged and continued for approximately 5 minutes. The breath-hold ended when the animal

rolled into ventral recumbency and took a breath. Prior to this study the animals had previously

participated in breath-hold experiments of durations up to 5 minutes (Fahlman et al., 2019a,

2020b).

<u>Blood collection and processing</u>. Whole blood was collected from tail flukes at baseline (0-30

seconds into the breath-hold) and during breath-holding for 3 minutes and 4 ½ (2018) or 5

(2019) minutes while the animal was in dorsal recumbency with its blowhole submerged (**Figure**

2A). Blood was collected into PAXgene tubes and RNA-Seq was performed subsequent to

shipping, red blood cell lysis, RNA extraction (**Figure 2A**). All samples were shipped the same

day via overnight courier to Duke University for downstream processing. For RNA extraction,

tubes were equilibrated to room temperature for 2 hours to achieve complete lysis of blood cells.

Subsequently, tubes were centrifuged at 4,000 x g for 10 minutes. Pellets were resuspended in 4

mL of RNase-free water and RNA was extracted according to the PAXgene Blood RNA kit

(PreAnalytiX #762164). Prior to library prep, RNA quality was evaluated on a Bioanalyzer 2100

(Agilent). Stranded mRNA-seq libraries were prepared using the Nugen Universal Plus mRNA-

seq Library preparation kit with Globin AnyDeplete (Tecan #9147-A01). Libraires were

sequenced at 150bp paired-end on one lane of an Illumina NovaSeq 6000 instrument S-Prime

flow cell.   Library preparation and sequencing was performed in conjunction with the Duke

University Sequencing and Genomic Technologies Shared Resource. Samples collected in 2018

were used to conduct RNA-Seq analysis and samples collected in 2019 were used for the

lipoxygenase assays.

RNA-Seq data analysis. RNA-seq data was processed using the TrimGalore toolkit (Krueger,

2020) which employs Cutadapt (Martin, 2011) to trim low-quality bases and Illumina sequencing

adapters from the 3' end of the reads. Only reads that were 20nt or longer after trimming were

kept for further analysis. Reads were mapped to the turTru1v92 version of the dolphin genome

and transcriptome (Kersey et al., 2012) using the STAR RNA-seq alignment tool (Dobin et al.,

2013). Reads were kept for subsequent analysis if they mapped to a single genomic location.

Gene counts were compiled using the HTSeq tool (Anders et al., 2015). Only genes that had at

least 10 reads in any given library were used in subsequent analysis. Normalization and

differential expression across the time points were carried out using the DESeq2 (Love et al.,

2014) Bioconductor (Huber et al., 2015) package with the R statistical programming

environment (R Core Team, 2020). The false discovery rate was calculated to control for

multiple hypothesis testing. To identify relevant molecular features of dolphin breath-holding we

first analyzed the RNA-Seq data from all individuals at baseline using gene set enrichment

analysis (GSEA) (Mootha et al., 2003; Subramanian et al., 2005). GSEA is a standard pathway

analysis tool that calculates enrichment scores for annotated pathways based on the rank order of

genes present in the data for each pathway. Pathways with genes that are more up- or down-

regulated are more likely to be enriched in a data set than pathways whose genes are randomly

distributed throughout the data. Pathway enrichments in dolphin PBMCs at baseline, with genes

ranked on total expression value, were compared with human whole blood pathway enrichments

from the Genotype-Tissue Expression (GTEx) project.

Construction of gene regulatory networks. Gene expression networks were created using

GeneMANIA (Franz et al., 2018), implemented within the Cytoscape platform (Shannon et al.,

2003). For time-dependent gene network construction all nodes with 0 or 1 connection were

trimmed out of the networks. Two additional non-coding RNA genes were eliminated (RF00016

and RF00026). To quantify network connectivity, all genes in the network were individually

ranked by the following network parameters: degree, clustering coefficient, closeness,

betweenness, neighborhood connectivity, and stress. These rankings were summed to generate a

sum rank score for each gene. Pathway enrichments were performed in STRING using the

trimmed network of 123 genes. Human whole blood transcriptomics data used for the analyses

described in this manuscript were obtained from the Genotype-Tissue Expression (GTEx)

Program Portal (https://gtexportal.org/home/) accessed on 9/20/2020.

Lipoxygenase assays. Briefly, 5 ml of blood was collected directly into BD Vacutainer® SST™

Tubes (SST) using a 21 g, ¾ in. winged infusion set with a BD Vacutainer adapter and holder.

Tubes were gently inverted 5 times to activate clotting reagent and allowed to clot at room

temperature for 30 minutes in an upright position. Tubes were centrifuged at 1,500 x g for 15

minutes to separate serum fractions, and serum was transferred to 15 ml conical tubes, frozen on

dry ice, and shipped to Duke University for downstream analyses. Sera were stored at -80°C

until use. Lipoxygenase activity was quantified from 1 µg of total protein using a Fluorometric

Lipoxygenase Activity Assay Kit (BioVision Inc; cat. #K978).

Results

RNA-Seq from dolphins at baseline pinpoints enriched gene regulatory networks. All samples produced between 30 and 40 million reads, with no time-dependent changes in read counts across samples (**Supplementary Figure 1A**). Principle components analysis and hierarchical clustering of all samples (three individual dolphins x three time points) revealed both individual- and within-individual time-dependent grouping of the data (**Supplementary Figure 1B, C**). GSEA identified multiple pathways enriched in dolphin PBMCs at baseline when ranked by total expression, including hedgehog signaling and several pathways relevant to blood cell metabolism, including heme metabolism, coagulation, IL6/JAK/STAT3 activation, apical junctions, and allograft rejection (**Figure 2B, C**). GSEA also identified enrichment of pathways related to apical junctions, interferon alpha response, estrogen response, complement activity, and heme metabolism in RNA-Seq data from GTEx human whole blood transcriptomes (**Figure 2D**). Comparison of dolphin baseline RNA-Seq data ranked by total expression with the top 100 and 500 most highly-expressed genes in human whole blood showed significant enrichment (FDR<0.0001) (**Figure 2E**). Together these analyses suggest that significant overlap exists in mRNA expression at both the gene-level and pathway-level between dolphin and human blood.

Breath-holding induces upregulation of multiple regulatory pathways. We next reasoned that patterns of step-wise increases in mRNA expression may pinpoint molecular responses to breath-holding common across individuals. We constructed gene regulatory networks for 136 genes with step-wise increases in mRNA expression from baseline to 3 minutes and again from 3 minutes to 4 ½ minutes (**Figure 3A**). We performed network analysis to identify genes that are upregulated and have the most network interactions. To do this we analyzed the time-dependent

216   gene regulatory network for the following parameters: degree, clustering coefficient, closeness,

217   betweenness, neighborhood connectivity, and stress. We then plotted the sum rank score of these

218   network parameters with gene expression for each gene in the network. These analyses

219   pinpointed arachidonate 5-lipoxygenase (ALOX5) as among the most connected genes with a

220   time-dependent increase in expression (**Figure 3B, C**). Additional genes, including EPX,

Figure 2. RNA-Seq from dolphin peripheral blood mononuclear cells reveals enrichment of pathways similar to humans. A. Whole blood from dolphins undergoing fasted breath-holds at baseline (0-30 seconds), 3 minutes, and 4 ½ minutes was collected from tail flukes and stored in PAXgene tubes for RNA extraction of peripheral blood mononuclear cells and RNA-Seq. B. Gene set enrichment analysis of baseline RNA-Seq data ranked by total expression pinpoints highly expressed relevant pathways. C. Enrichment plots for heme metabolism, coagulation, and IL6/JAK/STAT3 signaling from baseline dolphin RNA-Seq data. D. GSEA-based pathway enrichment from GTEx human whole blood RNA-Seq data ranked by total expression. E. GSEA enrichment plots comparing dolphin RNA-Seq data ranked by total expression with top 100 and top 500 expressed genes in human whole blood.

PTGDR2, SIX5, DCN, ADAMTS12, and GLRX2 demonstrated upregulation and/or high

network connectivity (**Figure 3B, C**). We used GeneMANIA to infer transcription factor and

microRNA targets from this time-dependent network. The gene regulatory network produced

from these genes displayed enrichment in targets from several transcription factor families,

including GATA and the <u>s</u>mall, <u>m</u>others <u>a</u>gainst <u>d</u>ecapentaplegic (SMAD) families (**Figure 3D**),

both of which have been implicated in hematopoietic development and regulation (Blank and

Karlsson, 2011; Lentjes et al., 2016). Network inference also pinpointed enrichment of targets of

multiple microRNAs, including the miR148A/B/152 family, miR492, miR186, miR518A-2, the

Figure 3. Time-dependent upregulation of gene regulatory pathways during dolphin breath holding. A. Gene regulatory network formed by the time-dependent increases in mRNAs from baseline to 3 minutes and 4 ½ minutes. Fold changes for each gene over time are indicated by darker red and larger nodes. B. Network analysis of genes within a co-expression network with increased expression over time. C. Top genes with increased expression sorted by their network analysis parameters. D. GeneMANIA-based transcription factor inference pinpoints GATA and SMAD transcription factor targets within the time-dependent network. E. MicroRNA enrichment inference based on the time-dependent network. F. Functional pathway enrichments for the time-dependent gene regulatory network.

miR130A/B/301 family, and miR205 (**Figure 3E**). Consistent with the identification of ALOX5

as a core network node, the network was functionally enriched the synthesis of 5-

eicosatetraenoic acid pathway, which is an initial step in the production of arachidonic acid by

ALOX5 (**Figure 3F**).

Arachidonate 5-Lipoxygenase (ALOX5) and subsequent lipoxygenase activity enhanced in

breath-holding dolphins. Based on the network analyses we focused on ALOX5 for further

validation. At the mRNA level, ALOX5 was one of just two genes, along with IL5RA, that was

significantly upregulated in all three individuals during breath-holding (**Figure 4A, B**).

Lipoxygenase assays from serum of three individual dolphins collected in 2019 revealed time-

dependent increases in lipoxygenase activity during breath-holding in all three individuals,

consistent with the RNA-Seq analyses (**Figure 4C**). Comparison of the timing of these molecular

changes with previously-published f_H measurements from the same dolphins demonstrated that

changes in gene expression and enzymatic activity were coincident with the expected timing of

bradycardia based on the heart rate data (**Figure 4D**). Overlay of the RNA-Seq data for ALOX5

expression with the heart rate data shows the upregulation of ALOX5 is concomitant with lower

heart rate (**Figure 4E**).

Figure 4. Dolphins induce ALOX5 activity during breath-holding. A. ALOX5 and B. IL5RA mRNA expression is significantly increased over time during breath-holding. C. Individual dolphin lipoxygenase activity in whole blood collected at an independent sampling date. D. Physiological measurements of heart rate for three individual dolphins (black lines from ECG data previously published in Blawas et al., 2020, dashed lines from echocardiogram data previously published in Fahlman et al., 2020) over time. Inset shows heart rate for humans performing breath-holds with facial immersion in water (dark gray in inset) overlaid on dolphin heart rate. Human heart rate traces were digitally extracted from Arnold, 1985; Andersson et al., 2004; and Shattock and Tipton, 2012. E. Overlay of heart rate data with ALOX activity in three individual dolphins. F. Hypothesized mechanism through which ALOX5 improves dive performance and mitigates ischemic stress tolerance.

Discussion

Dolphins and other cetaceans have evolved exquisite physiological adaptations to deal with the

challenges of a fully aquatic lifestyle including having a hydrodynamic shape to reduce drag

(Fish, 1993), counter-current heat exchangers for thermoregulation (Favilla and Costa, 2020;

Pabst et al., 1999), and cardiorespiratory plasticity for exquisite management of circulation and

respiratory gases (Blawas et al., 2021; Fahlman et al., 2020b, 2020a, 2019b; Noren et al., 2012).

The well-known dive response, a suite of adaptations that support reduced aerobic metabolism

during diving, involves apnea, bradycardia, and peripheral vasoconstriction that assures

maintained mean arterial blood pressure as blood flow to peripheral tissues is reduced and allows

regulation of perfusion to conserve oxygen-rich blood for the brain and heart. To maintain a

constant mean arterial blood pressure and prevent hypertension, these adaptations must work in

concert to ensure efficient autoregulation; however, extended dives also result in frequent events

of ischemia and hypoxia (Fahlman et al., 2019a; McKnight et al., 2019; Ridgway et al., 1969).

While these cardiorespiratory adaptations have been studied from the perspective of the changes

in f_H associated with diving, we are not aware of any study that has measured blood pressure in

voluntarily diving cetaceans. Thus, little is known about whether dolphins are able to maintain

constant mean arterial blood pressure throughout the breath-hold. In addition, knowledge of the

molecular adaptations that contribute to enhanced tolerance to hypoxia and ischemic stress, and

that prevent reperfusion injury during and following a dive, is rudimentary at best. To address

this lack of understanding, we combined an integrated genomics and systems-level analysis of

breath-hold responses at the molecular level with existing physiological measurements to define

the molecular responses to breath-holding in dolphins.

While this study is limited by a small sample size and relatively short breath-hold

durations, our analyses identified candidate genes and pathways with time-dependent changes in

expression throughout the breath-holds that were validated in functional studies using

independently-collected samples and assays. Consequently, these results provide evidence for

fine-scale cardiovascular control in bottlenose dolphins at the genetic level, and suggest that

dolphins may manage blood pressure changes during diving using both autonomic and molecular

pathways to regulate peripheral vasomotor control. Notably, these molecular changes occurred

within the calculated aerobic dive limit (cADL) of bottlenose dolphins - the duration of a dive

that can be sustained without requiring anaerobic respiration at the cellular level which has been

estimated to be 6.5 minutes (Fahlman et al., 2018). This suggests that changes in gene

expression may operate on a short enough time-scale that they could play a role in driving the

physiological changes that are observed during the breath-hold and recovery. It is also worth

considering the possibility that changes in gene expression could occur to support specific

physiological responses to diving during a dive, and that this gene expression differs when the

animal is at the surface. Future studies will be focused on using novel technologies, such as

GRO-Seq (Lopes et al., 2017) and others to measure nascent mRNAs, as well as measuring later

time points to understand the changes that occur upon recovery from breath-holds.

To provide physiological context for these molecular alterations on the time scales

observed, we compared molecular changes to changes in previously published f_H patterns in the

same individual dolphins during submerged breath-holds (Blawas et al., 2021; Fahlman et al.,

2020b). If we assume that the appearance of vasoconstriction is coincident with bradycardia, our

data provide evidence of an increase in the expression of a gene, ALOX5, known to promote

vasoconstriction coincident with the onset of vasoconstriction. Vasoconstriction, or a narrowing

of the blood vessels, has been suggested as a mechanism by which marine mammals during

forced dives have been observed to optimize the use of onboard oxygen stores in the blood and

muscle (Davis and Kanatous, 1999; Scholander and Grinnell, 1942; Zapol et al., 1979). Given

the long assumed link between vasoconstriction and bradycardia in marine mammals, the rapid

bradycardia we observed suggests that vasoconstriction was occurring in the dolphins in this

study during breath-holds (Hochachka, 1981; Van Citters et al., 1965). We found that changes in

gene expression occurred in all animals during the 5-minute breath-hold trials and that the same

gene families that were upregulated in the dolphins during breath-holds help manage

vasoconstriction in mice (Ichinose et al., 2001) and humans (Friedman et al., 1984).

Our integrated approach reveals possible molecular underpinnings that may support and

act synergistically with the cardiac response to breath-holding in bottlenose dolphins.

Specifically, we identified a suite of candidate genes that may support peripheral

vasoconstriction and provide defense against ischemic and hypoxic stress in dolphins, including

the GATA and SMAD transcription factors, several microRNAs, a disintegrin and

metalloproteinase with thrombospondin motifs 12 (ADAMTS12), mitochondrial glutaredoxin-2

(Glrx2), and ALOX5. Interestingly, many of these factors play known roles in regulating

hypoxia, hematopoiesis, and ischemic stress responses. For example, the GATA transcription

factor family is an important modulator of hematopoietic development of T lymphocytes, mast

cells, and erythrocytes (Lentjes et al., 2016). Likewise, the SMAD family regulates

hematopoietic stem cells (Blank and Karlsson, 2011). Of the microRNAs identified from our

analysis of target enrichments, nearly all have been shown to be protective against ischemia-

induced cell death, including miR148A (Zheng et al., 2018), miR492 (Guo et al., 2020), miR186

(Bostjancic et al., 2009; Li et al., 2013; Wang et al., 2018), miR130 (Lu et al., 2015), and

miR205 (Chen et al., 2019). At the protein-coding gene level, ADAMTS12 genetic variation is

associated with pediatric stroke (Witten et al., 2020), GLRX2 is implicated in neuroprotection

during hypoxia and ischemia (Romero et al., 2015), and ALOX5 is known to be induced by

hypoxia (Porter et al., 2014) and mediates the production of leukotrienes, which induce

bronchoconstriction and vasoconstriction (Poeckel and Funk, 2010). In addition, both ALOX5

and IL5RA have been identified as susceptibility genes associated with asthma and asthmatic

inflammation in humans (Cheong et al., 2005; Mougey et al., 2013), and a monoclonal antibody

to the IL5RA ligand, IL5, is FDA-approved for the treatment of severe eosinophilic asthma

(Fala, 2016; Pavord et al., 2012). Given the intricate connection between molecular control and

physiologic function to manage ischemia, hypoxia, and inflammatory responses in humans and

rodent models, (Bartels et al., 2013) it is intriguing to speculate as to how dolphins and other

marine mammals may uncouple or leverage these interconnected processes for improved

tolerance to ischemic/hypoxic stress without the pathological consequences associated with

hyper-stimulation of these processes.

These results demonstrate that the ALOX5 pathway is upregulated in bottlenose dolphins

during breath-holds and offer a potential mechanism for maintaining elevated peripheral

resistance through vasoconstriction, which helps manage blood distribution and the available

oxygen for critical organs. We suggest that the upregulation of ALOX5 could support a genetic

response that is secondary to the autonomic response during diving to prolong vasoconstriction

and maintain mean arterial blood pressure during extended periods of submersion (**Figure 4F**).

Interestingly, the changes we observed occurred within the cADL of the species, indicating that

fluctuations in gene expression could be occurring during regular dives. These fast-acting

changes in gene expression that support vasoconstriction provide evidence for fine-scale control

of perfusion in dolphins and an ability to maintain constant blood pressure, as is observed during

forced dives of pinnipeds (Blix et al., 1976; Zapol et al., 1979). Additionally, the data show that

during the breath-holds a large suite of candidate genes are upregulated that may support an

increased tolerance to the hypoxia and ischemic conditions that are expected to arise in some

peripheral tissues during diving. By interpreting these molecular data in the context of the

physiological changes known to occur in bottlenose dolphins during a breath-hold, we have

identified several genes that are upregulated during apnea in dolphins and may be an additional

mechanism to reinforce vasoconstriction while also providing defense against the hypoxia and

ischemia resulting from this response.

By examining molecular data through a physiological lens, these data connect the cellular

and tissue-level responses of dolphins to apnea to understand whether the bottlenose dolphin

may be genetically tuned to dive and withstand the hypoxia and the potential implications of this

to translational medicine. Our results uncover potential candidates at the intersection of ischemia,

hypoxia, and vasoconstriction that may contribute to the exquisite adaptation of dolphins and

other marine mammals to life in the ocean.

Abbreviations List

cADL, calculated aerobic dive limit

ALOX5, Arachidonate 5-Lipoxygenase

GSEA, Gene Set Enrichment Analysis

f_H, heart rate

IL5RA, Interleukin 5 receptor, alpha

PBMC, peripheral blood mononuclear cells

Acknowledgments

The authors wish to thank the marine mammal specialists, veterinarians, and dolphins at Dolphin

Quest, Oahu, The Duke Sequencing and Genomics Technologies Shared Resource, and the Duke

Genomics Analysis and Bioinformatics Shared Resource. The authors also thank the Laboratory

of Dr. Peter Hoffman at The University of Hawai'i for providing a centrifuge and other

equipment for sample processing. This work was supported through Dolphin Quest, Oahu (JAS)

and the Triangle Center for Evolutionary Medicine (JAS). AMB was supported by a Bureau of

Ocean Energy Management (BOEM) Environmental Study and the E. Bayard Halsted

Scholarship in Science, History, and Journalism. The authors would like to thank Giselle Vargas

and Mallissa Vuong for their role in data collection.

References

Allen, K.N., Vázquez-Medina, J.P., 2019. Natural Tolerance to Ischemia and Hypoxemia in
Diving Mammals: A Review. Front. Physiol. 10.
https://doi.org/10.3389/fphys.2019.01199
Anders, S., Pyl, P.T., Huber, W., 2015. HTSeq—a Python framework to work with high-
throughput sequencing data. Bioinformatics 31, 166–169.
https://doi.org/10.1093/bioinformatics/btu638
Bagchi, A., Batten, A.J., Levin, M., Allen, K.N., Fitzgerald, M.L., Hückstädt, L.A., Costa, D.P.,
Buys, E.S., Hindle, A.G., 2018. Intrinsic anti-inflammatory properties in the serum of
two species of deep-diving seal. Journal of Experimental Biology 221.
https://doi.org/10.1242/jeb.178491
Bartels, K., Grenz, A., Eltzschig, H.K., 2013. Hypoxia and inflammation are two sides of the
same coin. Proc Natl Acad Sci U S A 110, 18351–18352.
https://doi.org/10.1073/pnas.1318345110
Bi, J., Hu, B., Zheng, J., Wang, J., Xiao, W., Wang, D., 2015. Characterization of the hypoxia-
inducible factor 1 alpha gene in the sperm whale, beluga whale, and Yangtze finless
porpoise. Mar Biol 162, 1201–1213. https://doi.org/10.1007/s00227-015-2662-4
Blank, U., Karlsson, S., 2011. The role of Smad signaling in hematopoiesis and translational
hematology. Leukemia 25, 1379–1388. https://doi.org/10.1038/leu.2011.95
Blawas, A.M., Nowacek, D.P., Allen, A.S., Rocho-Levine, J., Fahlman, A., 2021. Respiratory
sinus arrhythmia and submersion bradycardia in bottlenose dolphins (Tursiops truncatus).
Journal of Experimental Biology. https://doi.org/10.1242/jeb.234096
Blix, A.S., Kjekshus, J.K., Enge, I., Bergan, A., 1976. Myocardial Blood Flow in the Diving
Seal. Acta Physiologica Scandinavica 96, 277–280. https://doi.org/10.1111/j.1748-
1716.1976.tb10196.x
Bona, E., Andersson, A.-L., Blomgren, K., Gilland, E., Puka-Sundvall, M., Gustafson, K.,
Hagberg, H., 1999. Chemokine and Inflammatory Cell Response to Hypoxia-Ischemia in
Immature Rats. Pediatric Research 45, 500–509. https://doi.org/10.1203/00006450-
199904010-00008
Bostjancic, E., Zidar, N., Glavac, D., 2009. MicroRNA microarray expression profiling in human
myocardial infarction. Dis Markers 27, 255–268. https://doi.org/10.3233/DMA-2009-
0671

Breskovic, T., Uglesic, L., Zubin, P., Kuch, B., Kraljevic, J., Zanchi, J., Ljubkovic, M., Sieber,
A., Dujic, Z., 2011. Cardiovascular changes during underwater static and dynamic
breath-hold dives in trained divers. J Appl Physiol (1985) 111, 673–678.
https://doi.org/10.1152/japplphysiol.00209.2011
Chen, W., Ruan, Y., Zhao, S., Ning, J., Rao, T., Yu, W., Zhou, X., Liu, C., Qi, Y., Cheng, F.,
2019. MicroRNA-205 inhibits the apoptosis of renal tubular epithelial cells via the
PTEN/Akt pathway in renal ischemia-reperfusion injury. Am J Transl Res 11, 7364–
7375.
Cheong, H.S., Kim, L.H., Park, B.L., Choi, Y.H., Park, H.-S., Hong, S.-J., Choi, B.W., Park, C.-
S., Shin, H.D., 2005. Association analysis of interleukin 5 receptor alpha subunit
(IL5RA) polymorphisms and asthma. J Hum Genet 50, 628–634.
https://doi.org/10.1007/s10038-005-0304-2
Choi, D.W., 1996. Ischemia-induced neuronal apoptosis. Current Opinion in Neurobiology 6,
667–672. https://doi.org/10.1016/S0959-4388(96)80101-2
Davis, R.W., Kanatous, S.B., 1999. Convective oxygen transport and tissue oxygen consumption
in Weddell seals during aerobic dives. Journal of Experimental Biology 202, 1091–1113.
Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,
M., Gingeras, T.R., 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29,
15–21. https://doi.org/10.1093/bioinformatics/bts635
Eltzschig, H.K., Carmeliet, P., 2011. Hypoxia and Inflammation. New England Journal of
Medicine 364, 656–665. https://doi.org/10.1056/NEJMra0910283
Fahlman, A., Brodsky, M., Miedler, S., Dennison, S., Ivančić, M., Levine, G., Rocho-Levine, J.,
Manley, M., Rocabert, J., Borque-Espinosa, A., 2019a. Ventilation and gas exchange
before and after voluntary static surface breath-holds in clinically healthy bottlenose
dolphins, Tursiops truncatus. Journal of Experimental Biology 222.
https://doi.org/10.1242/jeb.192211
Fahlman, A., Cozzi, B., Manley, M., Jabas, S., Malik, M., Blawas, A., Janik, V., 2020a.
Conditioned variation in heart rate during static breath-holds in the bottlenose dolphin
(Tursiops truncatus). Front. Physiol. 11. https://doi.org/10.3389/fphys.2020.604018
Fahlman, A., Hooker, S.K., Olszowka, A., Bostrom, B.L., Jones, D.R., 2009. Estimating the
effect of lung collapse and pulmonary shunt on gas exchange during breath-hold diving:
the Scholander and Kooyman legacy. Respir Physiol Neurobiol 165, 28–39.
https://doi.org/10.1016/j.resp.2008.09.013
Fahlman, A., Jensen, F.H., Tyack, P.L., Wells, R.S., 2018. Modeling Tissue and Blood Gas
Kinetics in Coastal and Offshore Common Bottlenose Dolphins, Tursiops truncatus.
Front. Physiol. 9. https://doi.org/10.3389/fphys.2018.00838
Fahlman, A., Miedler, S., Marti-Bonmati, L., Fernandez, D.F., Caballero, P.M., Arenarez, J.,
Rocho-Levine, J., Robeck, T., Blawas, A., 2020b. Cardiorespiratory coupling in
cetaceans; a physiological strategy to improve gas exchange? Journal of Experimental
Biology. https://doi.org/10.1242/jeb.226365
Fahlman, A., Miedler, S., Rocho-Levine, J., Jabois, A., Arenarez, J., Marti-Bonmati, L., García-
Párraga, D., Cauture, F., 2019b. Re-evaluating the significance of the dive response
during voluntary surface apneas in the bottlenose dolphin, Tursiops truncatus. Scientific
Reports 9. https://doi.org/10.1038/s41598-019-45064-8

Fala, L., 2016. Nucala (Mepolizumab): First IL-5 Antagonist Monoclonal Antibody FDA
Approved for Maintenance Treatment of Patients with Severe Asthma. Am Health Drug
Benefits 9, 106–110.
Favilla, A.B., Costa, D.P., 2020. Thermoregulatory Strategies of Diving Air-Breathing Marine
Vertebrates: A Review. Front. Ecol. Evol. 8. https://doi.org/10.3389/fevo.2020.555509
Fish, F.E., 1993. Power output and propulsive efficiency of swimming bottlenose dolphins
(Tursiops truncatus. J. Exp. Biol 179–193.
Franz, M., Rodriguez, H., Lopes, C., Zuberi, K., Montojo, J., Bader, G.D., Morris, Q., 2018.
GeneMANIA update 2018. Nucleic Acids Res 46, W60–W64.
https://doi.org/10.1093/nar/gky311
Friedman, Z., Lunyong, V.E., Courtney, J., Smith, H., Berkowitz, P., Sun, F., 1984.
Prostaglandin formation in the isolated human ductus arteriosus, aorta, pulmonary and
umbilical arteries. Prostaglandins Leukot Med 14, 279–286.
https://doi.org/10.1016/0262-1746(84)90211-7
Gooden, B.A., 1994. Mechanism of the human diving response. Integr Physiol Behav Sci 29, 6–
16. https://doi.org/10.1007/BF02691277
Gottlieb, R.A., Engler, R.L., 1999. Apoptosis in Myocardial Ischemia-Reperfusion. Annals of
the New York Academy of Sciences 874, 412–426. https://doi.org/10.1111/j.1749-
6632.1999.tb09255.x
Guo, L.-L., Chen, H.-H., Qu, F.-C., Lu, Q.-H., 2020. Clinical significance of miR-492 in
peripheral blood of acute myocardial infarction. Eur Rev Med Pharmacol Sci 24, 9041–
9045. https://doi.org/10.26355/eurrev_202009_22849
Hindle, A.G., 2020. Diving deep: Understanding the genetic components of hypoxia tolerance in
marine mammals. Journal of Applied Physiology.
https://doi.org/10.1152/japplphysiol.00846.2019
Hochachka, P.W., 1981. Brain, Lung, and Heart Functions during Diving and Recovery. Science
212, 509–514.
Huber, W., Carey, V.J., Gentleman, R., Anders, S., Carlson, M., Carvalho, B.S., Bravo, H.C.,
Davis, S., Gatto, L., Girke, T., Gottardo, R., Hahne, F., Hansen, K.D., Irizarry, R.A.,
Lawrence, M., Love, M.I., MacDonald, J., Obenchain, V., Oleś, A.K., Pagès, H., Reyes,
A., Shannon, P., Smyth, G.K., Tenenbaum, D., Waldron, L., Morgan, M., 2015.
Orchestrating high-throughput genomic analysis with Bioconductor. Nat Methods 12,
115–121. https://doi.org/10.1038/nmeth.3252
Ichinose, F., Zapol, W.M., Sapirstein, A., Ullrich, R., Tager, A.M., Coggins, K., Jones, R.,
Bloch, K.D., 2001. Attenuation of hypoxic pulmonary vasoconstriction by endotoxemia
requires 5-lipoxygenase in mice. Circ Res 88, 832–838.
https://doi.org/10.1161/hh0801.089177
Irving, L., Scholander, P.F., Grinnell, S.W., 1941. The respiration of the porpoise, tursiops
truncatus. Journal of Cellular and Comparative Physiology 17, 145–168.
https://doi.org/10.1002/jcp.1030170203
Johnson, P., Elsner, R., Zenteno-Savín, T., 2005. Hypoxia-inducible factor 1 proteomics and
diving adaptations in ringed seal. Free Radical Biology and Medicine 39, 205–212.
https://doi.org/10.1016/j.freeradbiomed.2005.03.008
Johnson, P., Elsner, R., Zenteno-Savín, T., 2004. Hypoxia-Inducible Factor in Ringed Seal
(Phoca hispida) Tissues. Free Radical Research 38, 847–854.
https://doi.org/10.1080/10715760410001725526

Kersey, P.J., Staines, D.M., Lawson, D., Kulesha, E., Derwent, P., Humphrey, J.C., Hughes,
D.S.T., Keenan, S., Kerhornou, A., Koscielny, G., Langridge, N., McDowall, M.D.,
Megy, K., Maheswari, U., Nuhn, M., Paulini, M., Pedro, H., Toneva, I., Wilson, D.,
Yates, A., Birney, E., 2012. Ensembl Genomes: an integrative resource for genome-scale
data from non-vertebrate species. Nucleic Acids Res. 40, D91-97.
https://doi.org/10.1093/nar/gkr895
Kooyman, G.L., Ponganis, P.J., 1998. The physiological basis of diving to depth: birds and
mammals. Annu. Rev. Physiol. 60, 19–32.
https://doi.org/10.1146/annurev.physiol.60.1.19
Krueger, F., 2020. FelixKrueger/TrimGalore.
Lentjes, M.H.F.M., Niessen, H.E.C., Akiyama, Y., de Bruïne, A.P., Melotte, V., van Engeland,
M., 2016. The emerging role of GATA transcription factors in development and disease.
Expert Rev Mol Med 18, e3. https://doi.org/10.1017/erm.2016.2
Li, C., Fang, Z., Jiang, T., Zhang, Q., Liu, C., Zhang, C., Xiang, Y., 2013. Serum microRNAs
profile from genome-wide serves as a fingerprint for diagnosis of acute myocardial
infarction and angina pectoris. BMC Med Genomics 6, 16. https://doi.org/10.1186/1755-
8794-6-16
Lopes, R., Agami, R., Korkmaz, G., 2017. GRO-seq, A Tool for Identification of Transcripts
Regulating Gene Expression. Methods Mol Biol 1543, 45–55.
https://doi.org/10.1007/978-1-4939-6716-2_3
Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biology 15, 550. https://doi.org/10.1186/s13059-
014-0550-8
Lu, C., Wang, X., Ha, T., Hu, Y., Liu, L., Zhang, X., Yu, H., Miao, J., Kao, R., Kalbfleisch, J.,
Williams, D., Li, C., 2015. Attenuation of cardiac dysfunction and remodeling of
myocardial infarction by microRNA-130a are mediated by suppression of PTEN and
activation of PI3K dependent signaling. J Mol Cell Cardiol 89, 87–97.
https://doi.org/10.1016/j.yjmcc.2015.10.011
Martin, M., 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal 17, 10–12. https://doi.org/10.14806/ej.17.1.200
McKnight, J.C., Bennett, K.A., Bronkhorst, M., Russell, D.J.F., Balfour, S., Milne, R., Bivins,
M., Moss, S.E.W., Colier, W., Hall, A.J., Thompson, D., 2019. Shining new light on
mammalian diving physiology using wearable near-infrared spectroscopy. PLOS Biology
17, e3000306. https://doi.org/10.1371/journal.pbio.3000306
Mirceta, S., Signore, A.V., Burns, J.M., Cossins, A.R., Campbell, K.L., Berenbrink, M., 2013.
Evolution of Mammalian Diving Capacity Traced by Myoglobin Net Surface Charge.
Science 340, 1234192–1234192. https://doi.org/10.1126/science.1234192
Mootha, V.K., Lindgren, C.M., Eriksson, K.-F., Subramanian, A., Sihag, S., Lehar, J.,
Puigserver, P., Carlsson, E., Ridderstråle, M., Laurila, E., Houstis, N., Daly, M.J.,
Patterson, N., Mesirov, J.P., Golub, T.R., Tamayo, P., Spiegelman, B., Lander, E.S.,
Hirschhorn, J.N., Altshuler, D., Groop, L.C., 2003. PGC-1alpha-responsive genes
involved in oxidative phosphorylation are coordinately downregulated in human diabetes.
Nat. Genet. 34, 267–273. https://doi.org/10.1038/ng1180
Mougey, E., Lang, J.E., Allayee, H., Teague, W.G., Dozor, A.J., Wise, R.A., Lima, J.J., 2013.
ALOX5 polymorphism associates with increased leukotriene production and reduced

lung function and asthma control in children with poorly controlled asthma. Clin Exp
Allergy 43, 512–520. https://doi.org/10.1111/cea.12076

Murdoch, C., Muthana, M., Lewis, C.E., 2005. Hypoxia Regulates Macrophage Functions in
Inflammation. The Journal of Immunology 175, 6257–6263.
https://doi.org/10.4049/jimmunol.175.10.6257

Nery, M.F., Arroyo, J.I., Opazo, J.C., 2013. Accelerated evolutionary rate of the myoglobin gene
in long-diving whales. J. Mol. Evol. 76, 380–387. https://doi.org/10.1007/s00239-013-
9572-1

Noren, S.R., Kendall, T., Cuccurullo, V., Williams, T.M., 2012. The dive response redefined:
underwater behavior influences cardiac variability in freely diving dolphins. Journal of
Experimental Biology 215, 2735–2741. https://doi.org/10.1242/jeb.069583

Nowbar, A.N., Gitto, M., Howard, J.P., Francis, D.P., Al-Lamee, R., 2019. Mortality From
Ischemic Heart Disease. Circulation: Cardiovascular Quality and Outcomes 12, e005375.
https://doi.org/10.1161/CIRCOUTCOMES.118.005375

Pabst, D.A., Rommel, S.A., McLellan, W.A., 1999. Functional anatomy of marine mammals, in:
Biology of Marine Mammals.

Panneton, M.W., 2013. The Mammalian Diving Response: An Enigmatic Reflex to Preserve
Life? Physiology 28, 284–297. https://doi.org/10.1152/physiol.00020.2013

Pavord, I.D., Korn, S., Howarth, P., Bleecker, E.R., Buhl, R., Keene, O.N., Ortega, H., Chanez,
P., 2012. Mepolizumab for severe eosinophilic asthma (DREAM): a multicentre, double-
blind, placebo-controlled trial. Lancet 380, 651–659. https://doi.org/10.1016/S0140-
6736(12)60988-X

Poeckel, D., Funk, C.D., 2010. The 5-lipoxygenase/leukotriene pathway in preclinical models of
cardiovascular disease. Cardiovasc Res 86, 243–253. https://doi.org/10.1093/cvr/cvq016

Porter, K.M., Kang, B.-Y., Adesina, S.E., Murphy, T.C., Hart, C.M., Sutliff, R.L., 2014. Chronic
hypoxia promotes pulmonary artery endothelial cell proliferation through H_2O_2-induced
5-lipoxygenase. PLoS One 9, e98532. https://doi.org/10.1371/journal.pone.0098532

R Core Team, 2020. R: A Language and Environment for Statistical Computing. R Foundation
for Statistical Computing, Vienna, Austria.

Ridgway, S.H., Scronce, B.L., Kanwisher, J., 1969. Respiration and Deep Diving in the
Bottlenose Porpoise. Science 166, 1651–1654.

Romero, J.I., Hanschmann, E.-M., Gellert, M., Eitner, S., Holubiec, M.I., Blanco-Calvo, E.,
Lillig, C.H., Capani, F., 2015. Thioredoxin 1 and glutaredoxin 2 contribute to maintain
the phenotype and integrity of neurons following perinatal asphyxia. Biochim Biophys
Acta 1850, 1274–1285. https://doi.org/10.1016/j.bbagen.2015.02.015

Scholander, P.F., 1940. Experimental Investigations on the Respiratory Function in Diving
Mammals and Birds. Hvalradets Skrifter.

Scholander, P.F., Grinnell, S.W., 1942. Aerobic and anaerobic changes in seal muscles during
diving 11.

Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N.,
Schwikowski, B., Ideker, T., 2003. Cytoscape: a software environment for integrated
models of biomolecular interaction networks. Genome Res 13, 2498–2504.
https://doi.org/10.1101/gr.1239303

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,
Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., Mesirov, J.P., 2005. Gene set
enrichment analysis: a knowledge-based approach for interpreting genome-wide

expression profiles. Proc Natl Acad Sci U S A 102, 15545–15550.
https://doi.org/10.1073/pnas.0506580102

Taboni, A., Fagoni, N., Moia, C., Vinetti, G., Ferretti, G., 2019. Gas exchange and
cardiovascular responses during breath-holding in divers. Respir Physiol Neurobiol 267,
27–34. https://doi.org/10.1016/j.resp.2019.06.002

Tian, R., Wang, Z., Niu, X., Zhou, K., Xu, S., Yang, G., 2016. Evolutionary Genetics of Hypoxia
Tolerance in Cetaceans during Diving. Genome Biol Evol 8, 827–839.
https://doi.org/10.1093/gbe/evw037

Tian, R., Yin, D., Liu, Y., Seim, I., Xu, S., Yang, G., 2017. Adaptive Evolution of Energy
Metabolism-Related Genes in Hypoxia-Tolerant Mammals. Front. Genet. 8.
https://doi.org/10.3389/fgene.2017.00205

Tift, M.S., Ponganis, P.J., Crocker, D.E., 2014. Elevated carboxyhemoglobin in a marine
mammal, the northern elephant seal. Journal of Experimental Biology 217, 1752–1757.
https://doi.org/10.1242/jeb.100677

Van Citters, R.L., Franklin, D.L., Smith, O.A., Watson, N.W., Elsner, R.W., 1965.
Cardiovascular adaptations to diving in the northern elephant seal Mirounga
angustirostris. Comparative Biochemistry and Physiology 16, 267–276.
https://doi.org/10.1016/0010-406X(65)90324-5

Wang, R., Bao, H., Zhang, S., Li, R., Chen, L., Zhu, Y., 2018. miR-186-5p Promotes Apoptosis
by Targeting IGF-1 in SH-SY5Y OGD/R Model. Int J Biol Sci 14, 1791–1799.
https://doi.org/10.7150/ijbs.25352

Witten, A., Rühle, F., de Witt, M., Barysenka, A., Stach, M., Junker, R., Nowak-Göttl, U., Stoll,
M., 2020. ADAMTS12, a new candidate gene for pediatric stroke. PLoS One 15,
e0237928. https://doi.org/10.1371/journal.pone.0237928

World Health Organization, 2018. Global Health Estimates 2016: Deaths by Cause, Age, Sex, by
Country and by Region, 2000- 2016. Geneva.

Yim, H.-S., Cho, Y.S., Guang, X., Kang, S.G., Jeong, J.-Y., Cha, S.-S., Oh, H.-M., Lee, Jae-Hak,
Yang, E.C., Kwon, K.K., Kim, Y.J., Kim, T.W., Kim, W., Jeon, J.H., Kim, S.-J., Choi,
D.H., Jho, S., Kim, H.-M., Ko, J., Kim, H., Shin, Y.-A., Jung, H.-J., Zheng, Y., Wang, Z.,
Chen, Y., Chen, M., Jiang, A., Li, E., Zhang, S., Hou, H., Kim, T.H., Yu, L., Liu, S.,
Ahn, K., Cooper, J., Park, S.-G., Hong, C.P., Jin, W., Kim, H.-S., Park, C., Lee, K.,
Chun, S., Morin, P.A., O'Brien, S.J., Lee, H., Kimura, J., Moon, D.Y., Manica, A.,
Edwards, J., Kim, B.C., Kim, S., Wang, J., Bhak, J., Lee, H.S., Lee, Jung-Hyun, 2014.
Minke whale genome and aquatic adaptation in cetaceans. Nature Genetics 46, 88–92.
https://doi.org/10.1038/ng.2835

Zapol, W.M., Liggins, G.C., Schneider, R.C., Qvist, J., Snider, M.T., Creasy, R.K., Hochachka,
P.W., 1979. Regional blood flow during simulated diving in the conscious Weddell seal.
Journal of Applied Physiology 47, 968–973. https://doi.org/10.1152/jappl.1979.47.5.968

Zheng, D., Li, Z., Wei, X., Liu, R., Shen, A., He, D., Tang, C., Wu, Z., 2018. Role of miR-148a
in Mitigating Hepatic Ischemia-Reperfusion Injury by Repressing the TLR4 Signaling
Pathway via Targeting CaMKIIα in Vivo and in Vitro. Cell Physiol Biochem 49, 2060–
2072. https://doi.org/10.1159/000493716

Zhou, X., Sun, D., Guang, X., Ma, S., Fang, X., Mariotti, M., Nielsen, R., Gladyshev, V.N.,
Yang, G., 2018. Molecular Footprints of Aquatic Adaptation Including Bone Mass
Changes in Cetaceans. Genome Biol Evol 10, 967–975.
https://doi.org/10.1093/gbe/evy062

Zhu, K., Ge, D., Wen, Z., Xia, L., Yang, Q., 2018. Evolutionary Genetics of Hypoxia and Cold
Tolerance in Mammals. J Mol Evol 86, 618–634. https://doi.org/10.1007/s00239-018-
9870-8
634

	🧑	🐬
Average dive duration (min)	0.5-0.67	0.6
Max dive duration (min)	11.6	13
cADL (min)	—	6.5

Animal ID	Age (years)	Body Mass (kg)	RNA-Seq	Lipoxygenase assay
6JK5	24	200.9	x	x
9FL3	35	251.7	x	
9ON6	21	192.8	x	x
83H1	11	147.0		x
Mean±S.D.	22.8±9.9	198.1±42.9		