Itaconate and derivatives reduce interferon responses and inflammation in influenza A virus infection

Aaqib Sohail[1,9*], Azeem A. Iqbal[1,9*], Nishika Sahini[1,9*], Mohamed Tantawy[1,9], Moritz Winterhoff[1,9], Thomas Ebensen[2], Robert Geffers[3], Klaus Schughart[4], Fangfang Chen[1,9], Matthias Preusse[1,2], Marina C. Pils[5], Carlos A. Guzman[2], Ahmed Mostafa[6], Stephan Pleschka[6], Christine Falk[7], Alessandro Michelucci[8], Frank Pessler[1,9,**]

[1]Biomarkers for Infectious Diseases, [2]Vaccinology and Applied Microbiology, [3]Genome Analytics, [4]Infection Genetics, [5]Mouse Pathology Platform, Helmholtz Centre for Infection Research, 31824 Braunschweig, Germany

[6]Institute for Medical Virology, Justus-Liebig-University, 35392 Giessen, Germany

[7]Department of Transplantation Immunology, Hannover Medical School, 30625 Hannover, Germany

[8]Dept. of Oncology, Luxembourg Institute of Health, 1526 Luxembourg

[9]TWINCORE Centre for Experimental and Clinical Infection Research, 30625 Hannover, Germany

* Joint first authors

** Corresponding author

Frank Pessler, MD, PhD

Tel. +49 511 220027-167, Mobile +49 160 9728 0364

frank.pessler@twincore.de

Abstract

Itaconate has recently emerged as a metabolite with immunomodulatory properties. We evaluated effects of endogenous itaconate and exogenous itaconate, dimethyl-, and 4-octyl-itaconate on host responses to influenza A virus infection. Infection induced *ACOD1* (the enzyme catalyzing itaconate synthesis) mRNA in monocytes and macrophages, which correlated with viral replication and was abrogated by itaconate treatment. Pulmonary inflammation and weight loss were greater in *Acod1*$^{-/-}$ than wild-type mice, and ectopic synthesis of itaconate in human epithelial cells reduced infection-induced inflammation. The compounds induced different recruitment programs in infected human macrophages, and transcriptome profiling revealed that they reversed infection-triggered interferon responses and modulated inflammation in cell lines, PBMC, and lung tissue. Single-cell RNA sequencing of PBMC revealed that infection induced *ACOD1* exclusively in monocytes, whereas treatment silenced IFN-responses in monocytes, lymphocytes, and NK cells. Viral replication did not increase under treatment despite the dramatically repressed IFN responses, but 4-octyl itaconate inhibited viral transcription in PBMC. The results reveal dramatic reprogramming of host responses by itaconate and derivatives and their potential as adjunct treatments for hyperinflammation in viral infection.

Key words: innate immunity - influenza virus – pneumonia – single cell RNA sequencing - treatment.

Introduction

As exemplified by influenza viruses (Ramos & Fernandez-Sesma, 2015) and pathogenic coronaviruses (Ragab, Salah Eldin et al., 2020), overshooting host inflammation, oxidative stress and cell damage/death are major determinants of morbidity and mortality in severe viral infections of humans. Consequently, there is a great need for adjunct treatments that can be given in conjunction with direct-acting anti-virals, with the aim to modulate the host response in such a way that end-organ damage is reduced and clinical outcome is improved.

Itaconic acid initially attracted great interest in the field of biotechnology due to its highly reactive double bonds, which make it an efficient intermediate in the biosynthesis of industrial polymers (reviewed in (Cordes, Michelucci et al., 2015)). The discoveries that its synthesis is highly induced during macrophage activation (Strelko, Lu et al., 2011) and that *immune response gene 1* (*Irg1,* since then renamed *aconitate decarboxylase*) encodes the enzyme cis-aconate decarboxylase (ACOD1 in humans, CAD in other organisms), which converts cis-aconitate to itaconate (Chen, Lukat et al., 2019, Michelucci, Cordes et al., 2013), have triggered intense efforts to elucidate functions of the endogenous ACOD1/itaconate axis in inflammation and infection. The early recognition that itaconate possesses anti-inflammatory and immunomodulatory properties also suggested that itaconate and prodrugs derived from it may have potential as immunomodulatory and cytoprotective interventions for inflammation-driven diseases (reviewed in (Hooftman & O'Neill, 2019, O'Neill & Artyomov, 2019)).

Most evidence of anti-inflammatory and immunomodulatory effects of itaconate has actually been obtained by treating cells or small animals with chemically modified variants that are presumably taken up more efficiently. 4-Octyl-itaconate (4OI) contains an added 8-carbon chain and acts primarily through activation of the NRF2 pathway (Mills, Ryan et al., 2018). It has, for instance, shown cytoprotective and anti-inflammatory effects in a variety of models, including lipopolysaccharide (LPS)-induced macrophage activation and LPS-induced sepsis in mice (Mills et al., 2018), in a neuronal model of oxidative stress (Liu, Feng et al., 2018), and in peripheral blood mononuclear cells (PBMC) from patients with systemic lupus erythematosus (Tang, Wang et al., 2018). Dimethyl-itaconate (DI) has also been shown to exert NRF2 independent anti-inflammatory effects by inhibiting IκBζ activity (Bambouskova, Gorvel et al., 2018). Furthermore, DI possesses cytoprotective effects, which was, for instance, shown by reduced cell damage in mouse models of cardiac ischemia (Lampropoulou, Sergushichev et al., 2016).

However, cytoprotective properties have also been demonstrated for unmodified itaconate,
notably in a rat model of cerebral ischemic-reperfusion injury (Cordes, Lucas et al., 2020). Both
compounds likely exert cytoprotective effects by reducing reactive oxygen species (ROS)
generation via inhibition of succinate dehydrogenase (Cordes et al., 2020, Cordes, Wallace et al.,
2016, Lampropoulou et al., 2016). Considering these varied effects of itaconate and derivatives, it
is not surprising that administration of DI has demonstrated beneficial effects in a variety of
mouse models of infection-associated inflammation such as LPS-induced mastitis (Zhao, Jiang et
al., 2019) and endometritis (Xu, Jiang et al., 2020b), and fungal keratitis (Gu, Lin et al., 2020).
Following early evidence that *Acod1* transcription is strongly induced in severe influenza A virus
(IAV) infection in mice (Preusse, Tantawy et al., 2013), its impact on host susceptibility to viral
infections has also been studied, and protective effects of itaconate on neurons exposed to RNA
viruses have been reported (Cho, Proll et al., 2013, Daniels, Kofman et al., 2019, Passalacqua,
Purdy et al., 2019) . On the other hand, in a recent study using a mouse model of IAV infection,
deleting the *Acod1* locus did not affect weight loss or survival, whereas it clearly worsened the
outcome of *Mycobacterium tuberculosis* infection in the same mouse strain (Nair, Huynh et al.,
2018). This lack of effect on IAV infection was surprising because abnormally increased
inflammation is usually associated with deleterious effects on the host during viral infections. For
instance, it has been shown that overactive inflammatory responses are associated with decreased
survival and increased weight loss in IAV infected inbred mouse strains (Alberts, Srivastava et
al., 2010). While the role of endogenous itaconate in protection from viral infections thus remains
incompletely understood, the above results do suggest a strong potential of external application of
itaconate or its derivatives to ameliorate inflammation and other deleterious consequences of
viral infections.
Focusing on IAV as one of the most important human respiratory viral pathogens (Collaborators,
2018), we therefore examined role and regulation of the endogenous ACOD1/itaconate axis in a
mouse model of IAV infection and subsequently performed a detailed analysis of the modulatory
effects of exogenous addition of itaconate and derivatives on inflammatory responses in IAV-
infected human cell lines, explanted lung tissue, and PBMC, both at the bulk and single-cell
levels. We find that loss of ACOD1 and itaconate synthesis led to increased inflammation during
infection and that exogenous itaconate, DI, and 4OI substantially reduced inflammation, most

notably IFN responses, without enhancing viral replication. 4OI was unique in that it also exerted

a direct anti-viral effect.

Results

110

111 ***Acod1* expression and itaconate synthesis are induced in mouse lung during IAV-infection**

112 **and limit pulmonary inflammation and disease severity.** Compared with C57BL/6J, DBA/2J

113 mice are more susceptible to IAV infection in that weight loss is more rapid, mortality is higher,

114 and inflammation is more pronounced (Preusse, Schughart et al., 2017, Srivastava, Blazejewska

115 et al., 2009). We have shown previously that *Acod1* transcription in the first 48 h of infection is

116 higher in IAV-infected lung from DBA/2J than from C57BL/6J mice (Preusse et al., 2013). A

117 reanalysis of our RNAseq data from a longer time course (Wilk, Pandey et al., 2015) showed that

118 this difference persisted at least through day 3 post infection (p.i.) and that *Acod1* expression

119 nearly normalized by day 14 in the surviving strain (**Fig. 1A**). This analysis also revealed a

120 remarkable correlation with expression of *Tnfaip3* (encoding an anti-inflammatory

121 deubiquitinase) and a weaker positive correlation with *Hmox1* and *2* (encoding inducible anti-

122 oxidant enzymes)*,* all of which are known to be co-regulated with *Acod1* (Li, Zhang et al., 2013,

123 Uddin, Joe et al., 2016). *Acod1* was the 4th most highly upregulated pulmonary mRNA 48 h p.i.

124 in C57BL/6J mice, where it appeared in a cluster of transcripts pertaining to IFN responses and

125 innate immunity (**Fig. 1B**). These results underscored the close association of *Acod1* expression

126 with systemic inflammation and oxidative stress in IAV-infected mice. We therefore tested

127 whether targeted deletion of *Acod1* would affect inflammatory responses and host susceptibility

128 to IAV infection. Itaconate was not detectable in lungs from IAV-infected *Acod1$^{-/-}$* mice; it was

129 highly elevated on days 7/8 p.i. in the *Acod1$^{+/+}$* and, significantly less, in *Acod1$^{+/-}$* mice, and

130 returned to normal by day 14 (**Fig. 1C**). Both weight loss and mortality were higher in *Acod1$^{-/-}$*

131 than *Acod$^{+/+}$* female C57BL/6J mice (**Fig. 1D,E**), with greatest changes occurring around day 8,

132 the time of maximal itaconate accumulation. Histologically assessed pulmonary inflammation

133 was greatest in *Acod1$^{-/-}$* mice and intermediate in *Acod1$^{+/-}$* mice (**Fig. 1F**). Thus, itaconate

134 synthesis correlated with disease activity, and loss of *Acod1* was associated with more severe

135 disease and higher pulmonary inflammation.

136 **IAV infection induces *ACOD1* expression in human PBMC and macrophages.** We then

137 tested whether IAV infection increased *ACOD1* mRNA levels in human PBMC and M1/M2

138 macrophages. Phorbol myristate acetate (PMA)-differentiated THP-1 (dTHP-1) cells were

139 analyzed for comparison. Viral transcription was highest in M2 cells (**Fig. 2A**). The strongest

ACOD1 mRNA induction was seen in M2 cells and undifferentiated PBMC, and the weakest in

M1 cells and dTHP-1 cells (**Fig. 2B-E**), even though the latter two cell types did support a

significant degree of viral transcription (**Fig. 2A,D**). As in IAV-infected mouse lung (**Fig. 1**),

TNFAIP3 and *ACOD1* transcriptional correlated with each other (**Fig. 2C**). Infection of dTHP-1

cells with three IAV (H1N1) strains of differential replication efficiency (Petersen, Mostafa et al.,

2018b) showed the strongest *ACOD1* and *TNFAIP3* induction by the two strains with the highest

and the weakest by the strain with the lowest replication efficiency (**Fig. 2D,E**). Thus, the extent

of *ACOD1* induction after IAV infection differed significantly among the cell types tested, was

strongest in M2 primary cells, was accompanied in all cases by an increase in *TNFAIP3* mRNA,

and correlated with viral replication.

Itaconate and DI markedly reduce IFN responses in dTHP-1 cells without affecting viral

RNA replication. We then tested whether these compounds could modulate cellular

inflammation due to IAV infection. In infected dTHP-1 cells, treatments with itaconate or DI did

not affect viral hemagglutinin (*HA*) mRNA expression, but both reduced induction of *CXCL10*, a

key pro-inflammatory chemokine during influenza virus infection, with DI being several-fold

more potent (**Fig. 3B,C, Fig. S1**). The 12 h time point was chosen because our previous work had

identified it as the peak of IFN-I responses at the mRNA level in IAV-infected dTHP-1 cells

(Petersen, Mostafa et al., 2018a). We then assessed global effects of itaconate and DI on cellular

gene mRNA expression. A comparison of differentially expressed genes (DEGs) suggested that

the two compounds had both shared and unique effects on gene expression (**Fig. 3D**). Both

compounds had considerable effects on gene expression in uninfected cells (**Fig. 3E,F**). As

expected, IAV infection of untreated cells caused a major induction of IFN-regulated transcripts

by 12 h p.i. (see mRNAs labeled in the volcano plot in **Fig. 3G**). Strikingly, addition of both

itaconate and DI led to global down-regulation of IFN-regulated genes. This was accompanied by

upregulation of a substantial number of other transcripts, indicating that the compounds also

modulated other cellular processes (**Fig. 3H,I**). A principle component analysis (PCA) indicated

that effects of the compounds on the cells were so pronounced at these concentrations that treated

infected and treated uninfected cells clustered closely together, but far away from untreated

infected or uninfected cells (**Fig. S2**). Indeed, a hierarchical clustering analysis revealed an

across-the-board normalization of a clade of IFN- and inflammation-related transcripts by the

compounds, but also clusters of DEGs that were upregulated by either or both compounds with

respect to both uninfected and infected cells, further indicating effects upon cell homeostasis in

general (**Fig. S3**). mRNA and protein expression may be differentially affected by IAV infection

due to virus-induced "host-cell shut-off". We therefore assessed release of inflammation-related

polypeptides in supernatants from the same experiment (**Fig. 3J**). This analysis confirmed the

downregulation of pro-inflammatory factors (e.g., IP-10, MCP-1) by itaconate and DI, but also

showed increased IL-8 (DI only) and minor increases of IL1B (both IA and DI) in uninfected

cells. Of note, both compounds also increased levels of the anti-inflammatory IL1 receptor

antagonist (IL1RA) in supernatants from both infected and uninfected cells. A comprehensive

view of the protein targets that were detected by this assay revealed that the two compounds

effected substantially different changes in the target cells, as several factors were induced even in

uninfected cells specifically by DI or itaconate (**Fig. S4**). Thus, itaconate and DI apparently act

on macrophages to release different recruitment programs that modify local inflammatory cell

populations.

Analysis of enriched GO terms revealed that IFN-I signaling was strongly induced in the infected

cells, which was prevented by treatment with itaconate and DI (**Fig. 4**). This analysis also

revealed that both compounds activated metabolic processes, whereas itaconate also modulated

processes relating to differentiation and membrane signaling. A KEGG pathway analysis

additionally revealed induction of classic pro-inflammatory pathways such as *TNF signaling*,

TLR receptor signaling, and *chemokine signaling* by infection, all of which were depleted by

itaconate and DI treatment (**Fig. S5**). Taken together, these results revealed strong anti-IFN and

anti-inflammatory effects of itaconate and DI, but also suggested that they exert other important

effects on cell metabolism and differentiation, some of which differ between the compounds.

Anti-inflammatory effects of ectopically expressed itaconate, and exogenous itaconate and
193
DI on IAV-infected respiratory epithelial cells. We then tested whether the effects of itaconate
194

and DI could also be seen in cells not endogenously expressing *ACOD1*. For this purpose, we

studied the respiratory epithelial cell line A549, which supports productive IAV infection and

does not express *ACOD1*. As expected, infection of A549 cells did not result in induction of

ACOD1 mRNA, but transient transfection with an *ACOD1*-expressing plasmid led to significant

ACOD1 mRNA expression (**Fig. 5A**), which was accompanied by appearance of substantial

concentrations of IA (**Fig. 5B**). While there was no effect on viral *HA* mRNA expression,

transfection with the *ACOD1*-expressing vector led to a marked reduction in infection-associated

CXCL10 induction (**Fig. 5C,D**). A preliminary treatment experiment showed that (as opposed to

THP-1 cells) DI was cytotoxic at 1.0 mM in A549 cells, and the 0.5 mM concentration was

therefore used. Treatment with DI resulted in a modest reduction of *HA* expression early in

infection and a marked reduction of *CXCL10*, *IL6*, and *IFNB* expression (**Fig. 5E-H**). Exogenous

itaconate, too, reduced *CXCL10* mRNA, but less efficiently (**Fig. S6**). Taken together, these

results obtained with A549 cells suggest that immunomodulatory effects of itaconate and DI are

independent of the ability of the cell type to express endogenous *ACOD1*, and that similar effects

on IFN-directed gene expression can be obtained with ectopic, endogenous synthesis of itaconate

and exogenously added itaconate and DI.

Next, we assessed effects of itaconate and DI on cellular transcriptomes in IAV-infected A549

cells. PCA revealed the expected normalization of gene expression due to DI treatment, but also

increasing effects of the compounds on overall cell homeostasis (**Fig. S7**). As in THP-1 cells,

hierarchical clustering analysis revealed clear clustering of each of the four groups and the close

relationship of the uninfected and DI-treated infected cells (**Fig. S8**). However, the anti-

inflammatory effect of itaconate was weaker than that of DI, and at the higher (40 mM)

concentration, the itaconate signature became so dominant that the itaconate-treated infected

group now formed its own clade (**Fig. S9**). IAV infection enriched GO terms related to IFN-I and

–II responses as well as other antiviral and pro-inflammatory terms, which could all be reduced to

near baseline by treatment with DI or itaconate (**Fig. 6**). Visualization of differentially expressed

IFN-related mRNA also showed that, in spite of its overall lower effect on pro-inflammatory

signaling than DI, itaconate treatment at the lower (20 mM) concentration did have a

considerable impact on IFN-I expression (**Fig 6B**). In addition, itaconate uniquely inhibited

NFKB signaling and *fatty acid metabolism*. KEGG-pathway analysis additionally revealed

depletion of other pro-inflammatory pathways, including *TNF signaling* and *necroptosis,* by the

treatments (**Fig. S10**).

Anti-inflammatory effect of itaconate and DI in a human lung tissue explant (HLTE) model

of IAV infection. To test whether the *ACOD1*/itaconate axis and the anti-inflammatory effects of

itaconate and DI are functional in the end-organ of IAV infection in humans, we then used a

human lung tissue culture model. Indeed, infection with IAV led to brisk transcription of viral *HA*

and host *CXCL10* mRNA in a 72 h time course (**Fig. S11**), whereas *ACOD1* mRNA levels did

not change significantly. Adding itaconate and DI reduced expression of *CXCL10* mRNA in

tissue and IP10 protein in culture supernatant, and itaconate additionally reduced ISG15 mRNA

in tissue, whereas viral titers in the supernatant were unaffected (**Fig. 7**). Re-analysis of a

published dataset from a model using a different IAV strain (H3N2) and histologically normal

lung tissue (Matos, Wunderlich et al., 2019) revealed a pronounced induction of *ACOD1* and

TNFAIP3 expression, but (as opposed to *in vivo* infection of mouse lung) not induction of

HMOX1 or *2* (**Fig. S12).**

Anti-inflammatory effects of itaconate, DI, and 4OI on IAV-infected PBMC. PBMC play

important roles in influenza infection as they contain cells of innate and adaptive immunity, and

because circulating monocytes (which can sustain a nonproductive influenza virus infection) are

considered an important source of cytokines that support systemic spread of inflammation (Duan,

Hibbs et al., 2016). We therefore investigated effects of the compounds on IAV infected PBMCs

both at the bulk and single cell levels. Analysis with targeted PCR- and immuno-assays

demonstrated vigorous transcription of viral RNA, induction of *ACOD1* mRNA, and a brisk pro-

inflammatory response both at the mRNA and protein level (**Fig. 8A,B**). The induction of

ACOD1 mRNA expression in peripheral blood in IAV infection was also corroborated by

reanalysis of a published dataset from whole blood from patients with moderate and severe

influenza (**Fig. S13**). The effects of itaconate were mixed in that there was no significant change

in *ACOD1*, *IFNB1*, and *TNF* mRNA expression in cells or of IL1B protein in supernatants. In

contrast, DI and 4OI led to a marked reduction of IFN and pro-inflammatory cytokine expression

at the mRNA and protein level, as well as of *ACOD1* mRNA. As exemplified by the effect of DI

and 4OI on *TNF* and *ACOD1* expression, the treatments reduced expression of some mRNA

targets even in uninfected cells, suggesting that they also reduced some baseline inflammation.

Of note, 4OI in addition led to a pronounced (95%) reduction in *HA* transcription (p=<0.0001),

indicating inhibition of viral RNA replication. PCA of bulk PBMC transcriptomes revealed

strong effects of the compounds on cell transcriptomes in general, which led to one outlying DI-

treated sample and two itaconate-treated samples, whereas all four 4OI-treated samples formed a

clearly discernible group (**Fig. S14**). The most significant transcriptome changes were driven by

4OI, which had major effects on many genes that were not regulated by IAV infection or

itaconate or DI treatments (**Fig. S15**). Nonetheless, induction of *HMOX1* by DI in IAV-infected

samples was evident. Functional enrichment analysis of GO terms based on the transcriptomes

revealed that infection induced the expected antiviral responses, including *response to IFN-I*, as

well as several other inflammation-related terms (**Fig. 8C**). At least one of the three compounds

prevented activation of each of the infection-associated terms. However, consistent with its lack

of effect on *IFNB* expression (**Fig. 8**), itaconate did not affect *response to IFN-I* or *influenza A*.

Several terms that were depleted by the compounds in infected PBMC were not enriched by

infection alone, suggesting that the compounds diminished also a baseline activation of the

PBMC. Interestingly, 4OI treatment strongly enriched terms relating to chromatin conformation,

which turned out to be driven by upregulation of histone deacetylase epression. A KEGG

pathway analysis essentially confirmed the findings of the GO term analysis (**Fig. S16**), including

that 4OI enriched pathways relating to chromatin structure such as *Alcoholism*. Taken together,

these results consolidated our above findings that itaconate and derivatives can modulate

inflammation due to IAV infection, but also suggested that DI and 4OI may have more robust

anti-IFN and anti-inflammatory effects than itaconate. Considering the variable anti-

inflammatory effects of itaconate and the miscellaneous effects of 4OI on gene expression in this

model, we subsequently focused on DI to decipher effects at the single-cell level.

Single-cell RNAseq of PBMC identifies monocytes as the main PBMC host cell of IAV

infection, sole source of *ACOD1* expression, and predominant target of immunosuppressive

effects. Thirteen cell types could be identified in the PBMC scRNAseq data set, but we focused

the analyses on the overarching five cell types: CD4+ and CD8+ T cells, B cells, NK cells, and

monocytes (**Fig. 9A,B**). Monocytes constituted the least numerous cell type, and DI treatment of

control and infected cells led to a further reduction (**Fig. S17**). It was not possible to discern

whether this was due to decreased survival or a technical artefact. Monocytes were essentially the

only cell type expressing viral RNA, as only negligible amounts of viral transcripts were detected

in lymphocytes and NK cells. DI treatment appeared to result in a slight increase in viral RNA

expression in monocytes (**Fig. 9C**). IAV infection upregulated expression of *ACOD1* exclusively,

and *CXCL10* predominantly, in monocytes, whereas *TNFAIP3*, *IFIT1,* and *ISG15* were

upregulated also in the NK, T and B cell compartments (**Fig. 9D**). DI treatment markedly reduced

expression of *ACOD1*, *IFIT1,* and *CXCL10*, but –interestingly– it reduced *TNFAIP3* and *ISG15*

(albeit weakly) expression only in monocytes. Consistent with the observation that they

constituted the main PBMC host cell, IAV infection triggered the most vigorous transcriptomic

host response in monocytes, which was driven by the expected IFN responses (**Fig. 9E**).

However, discernable differential expression (mostly of IFN-related RNAs) was also observed in

the other cell types. CD8+ T cells are shown as a representative example in **Fig. 9F**, the other cell

types in **Fig. S18**. Monocytes were the only cell type that mounted a transcriptomic response to

DI treatment of uninfected cells. In infection, treatment with DI led to a global downregulation of

IFN-related transcripts in monocytes and also downregulation of a subpopulation of IFN-related

transcripts in lymphocytes and NK cells. When considering all DEGs irrespective of magnitude

of differential expression, DI exerted comparable effects on monocytes and T cells, and only

somewhat less on B cells. However, when considering only transcripts with FC $>|2|$, a

preferential effect on monocytes became apparent (**Fig. 9G,H**).

GO term enrichment analysis of the monocyte scRNAseq data confirmed the major enrichment of

IFN-related processes by infection and their depletion by DI treatment, which had been seen in

all models in this study. In addition, infection broadly dampened RNA metabolism, ribosome

assembly, and protein synthesis and export, which was consistent with the cytostatic effects of

type I IFN. DI treatment reversed all these effects and also stimulated some in uninfected cells

(**Fig. 10A**). A KEGG enrichment analysis of these monocyte data confirmed the depletion of

IFN-related pathways (*influenza, measles, cytosolic DNA sensing pathway*) by DI treatment, but

also of other pro-inflammatory pathways (*TLR signaling, cytokine-cytokine receptor interaction,*

NF-kB signaling) in infected and uninfected cells (**Fig. S19**). Of note, DI treatment apparently

stimulated ribosome function in both infected and control monocytes.

The Venn diagrams in **Fig. 10B** illustrate that monocytes are the cell type in which the most

biological processes were activated by infection and depleted by DI treatment, but reduction of

antiviral and pro-inflammatory responses was a common theme in the other cell types as well

(**Fig. 10C**). Indeed, GO term analysis identified four major IFN-related processes as a central

functional network that was upregulated by infection and downregulated by DI treatment during

infection in all five cell types (**Fig. 10B**). A marked stimulation of protein synthesis and export

by DI was also apparent in the uninfected lymphocytes and NK cells. Finally, in order to identify

a common transcriptomic network we performed GO and KEGG pathway enrichment analyses of

the 81 transcripts that were commonly differentially expressed in IAV infection and regulated in

the opposite direction by DI treatment in all five cell types (center of the Venn diagram in **Fig.**

9G). While the GO analysis by and large confirmed the findings shown in **Fig. 10B,** the KEGG

pathway analysis revealed a major effect of DI treatment on *allograft rejection*, *graft-vs.-host*

disease, and *antigen processing and presentation,* suggesting a broader interference with immune

cell activation by DI (**Fig. S20**).

Discussion

In this first dedicated study on the impact of itaconate and derivatives on host responses to IAV infection, we found that (i) the endogenous ACOD1/itaconate axis limits pulmonary inflammation due to the infection in a murine model, (ii) ectopically expressed ACOD1 possesses anti-IFN activity in an epithelial ell line that normally does not express this enzyme, and (iii) that exogenously applied itaconate and/or derivatives restrict IFN-responses and modulate classic pro-inflammatory cytokines across all models tested.

Anti-inflammatory effects of itaconate and derivatives. A major suppression of IFN and some pro-inflammatory cytokine responses was the recurrent theme in all models, i.e. immortalized cells, primary cells, and human lung tissue. This is fully consistent with the early reports that DI has strong anti-inflammatory properties (Lampropoulou et al., 2016), which was subsequently corroborated in diverse models of inflammation and infection, also using 4OI (e.g., (Gu et al., 2020, Xu, Jiang et al., 2020a, Zhao et al., 2019)). We used itaconate and DI in parallel in several models, which enabled comparing their relative effectiveness. Overall, results with DI were most consistent, and in spite of its higher toxicity on A549 cells than on dTHP-1 cells, the lower dose that had to be used on A549 cells still possessed strong anti-inflammatory effects. Itaconate effects were less consistent and much higher doses had to be used to achieve anti-IFN effects similar to DI. Itaconate did not reduce expression of *IFNB1*, *TNF* mRNA or IL1B protein in PBMC. The lack of *TFN* repression by itaconate in that model is consistent with previous reports that it usually does not affect TNF-α level (Bambouskova et al., 2018). Recent work has also revealed the potential of itaconate to actually enhance IL1-β release (Muri, Wolleb et al., 2020). Of note, at higher doses, itaconate and strong electrophilic compounds can actually increase inflammation due to CASP-8 dependent inflammasome activation, and this effect is magnified when the cells are not pre-incubated with the compound before adding the pro-inflammatory stimulus (Muri et al., 2020), as we did in the PBMC infections. However, these authors did not investigate effects on IFN suppression. 4OI demonstrated the expected potent anti-inflammatory effects, but also broad effects on processes relating to, e.g., chromatin structure, the potential effects of which on cell homeostasis and cell-pathogen interactions will require further study.

Anti-inflammatory effect of ectopic synthesis of itaconate in A549 cells. Despite the multitude of

studies on effects of exogenous administration of itaconate derivatives on cellular inflammation,

it had not been tested whether ectopic synthesis of itaconate in a cell type that does not naturally

express *ACOD1* has similar effects. Our transfection study (Fig. 5) demonstrated that ectopic,

endogenously synthesized itaconate exerts very similar anti-IFN effects as exogenously added

itaconate or DI. Of note, the achieved itaconate concentration is very close to that measured in

LPS/IFNG stimulated dTHP-1 cells (Chen et al, unpublished data), suggesting that it is

physiological.

Cell-type specific and -independent anti-inflammatory effects of itaconate and DI. A particular

strength of our study lies in the use of scRNAseq to identify major target cells in peripheral

blood. Firstly, this analysis identified monocytes as the major PBMC type that is infected by

IAV. Secondly, we found that on one hand DI can attenuate IFN-responses in lymphocytes and

NK cell as well, but that the immunosuppressive effect is greatest on monocytes. Taken together,

these results suggest that even though pharmacological use of itaconate and derivatives would

modulate host responses in a diverse array of cells, monocytes/macrophages likely constitute a

major functional hub in the genesis of inflammation in IAV infection and subsequent treatment

responses to itaconate-based adjunct treatments.

Antiviral effect of 4OI. Olagnier et al. recently reported that Nrf2 activators including 4OI and

dimethyl fumarate limit inflammation in cellular models of SARS-CoV-2 infection (Olagnier,

Farahani et al., 2020) in an NRF2-dependent manner. Of note, as opposed to influenza viruses,

infection with SARS-CoV-2 suppresses the NRF2 pathway, which can be reversed in part by

treatment with Nrf2 activators. These authors in addition observed an antiviral effect of 4OI,

which proved to be independent of NRF2- and type I IFN-signaling. Our finding of a remarkable

suppression of IAV RNA replication in PBMC corroborates the notion that 4OI has strong

antiviral effects in addition to its well-documented anti-inflammatory properties (Olagnier et al.,

2020). Further research is needed to validate our finding in other cell types, as we tested it in

PBMC only. Nonetheless, the gene expression analysis of itaconate, DI- and 4OI-infected PBMC

identified several processes/pathways uniquely regulated by 4OI. Further research should focus

on assessing whether any of them could contribute to the antiviral effect of 4OI.

Infection-independent effects on target cells. By using relatively high doses of itaconate and DI
and assessing effects of the compounds on uninfected cells or their effects on processes that were
not affected by infection alone, our study also provides a first comprehensive look at global
effects of the compounds on target cells. In particular, itaconate and 4OI exerted pronounced,
broad effects on gene expression in A549 cells and PBMCs, respectively. Clearly, it will be of
great interest to investigate how these effects relate to differences in toxicities on different cell
types, or to differences in their effects on virus-host interactions.

Importance of endogenous ACOD1 expression. In the mouse model, we observed a more severe
clinical phenotype and higher inflammation in *Acod1$^{-/-}$* mice. This agrees with work in a mouse
model of *M. tuberculosis* infection, where inflammation and mortality were higher in the *Acod1$^{-/-}$*
animals (Nair et al., 2018). However, IAV infection was also assessed in that study, and deleting
the *Acod1* gene did not affect weight loss or survival. Those authors used equal numbers of male
and female mice, whereas we used female mice only. Further research is necessary into possible
sex-specific effects of the *Acod1* gene in influenza infection. Of note, we found upregulation of
ACOD1 by IAV infection and its prevention by DI treatment in myeloid cells, i.e. dTHP-1 and
peripheral blood monocytes, and upregulation of *ACOD1* mRNA in blood from patients with
moderate and severe influenza infection as compared with healthy donors. Thus, induction of
ACOD1 expression (and thereby itaconate synthesis) likely is a common feature in human
influenza. We have recently identified the active center of ACOD1 enzyme and found that
naturally occurring loss-of-function mutations are extremely rare, suggesting that physiologic
itaconate responses have been important in human evolution (Chen et al., 2019). It is therefore
tempting to speculate that one beneficial role of an active ACOD1/itaconate axis is to reduce the
risk of death or organ damage from infection-associated inflammation and that loss-of-function
genotypes have been selected against during times of high infectious disease burdens.

Clinical implications. The current COVID-19 pandemic has, once again, focused the research
limelight on the deleterious effects of overshooting innate immune responses in acute viral
infections. In a current report, Lee et al. found that intense IFN responses in PBMC from patients
are a hallmark of both severe COVID-19 and influenza (Lee, Park et al., 2020). Of note, a recent
systematic review of 45 studies on corticosteroid use for COVID-19 found a significant
beneficial effect (van Paassen, Vos et al., 2020), whereas to date there are no clinically effective

antivirals against SARS-CoV-2. Therefore, the concept of adjunct treatments to control

deleterious host immune responses in viral infections is more pertinent than ever. Considering (i)

the well documented cytoprotective properties of itaconate and derivatives and (ii) our findings of

remarkable anti-inflammatory and anti-IFN effects in the absence of increased viral replication,

itaconate derivatives would constitute a highly promising class of compounds for such

applications. Considering the more reliable results obtained with DI as compared to unmodified

itaconate, the lower required concentrations of DI and 4OI, but also the striking anti-viral effects

of 4OI, it appears that chemical variants of itaconate, rather than its native form, will take the

lead in further translational development to the bedside.

Acknowledgements. We thank V. Kaever, H. Bähre and staff of the RCU Metabolomics at

Hannover Medical School for support with the itaconate measurements, Elena Reinhard for

assistance with the animal experiments, the HZI Animal Facility for expert animal care, the HZI

Genome Analytics Platform for microarray and RNA sequencing, L. Gröbe for cell sorting, D.

Jonigk and P. Braubach (Institute of Pathology, Hannover Medical School) for providing human

lung tissue. We thank Hans Jörg Hauser (HZI) for helpful comments on the manuscript.

Conflict of interest. The authors declare that none of them have a conflict of interest relating to

conduct of the study or publication of the manuscript.

Authors' contributions. AS, AAI, CF, FC,NS, KS, MP, MT, and TE performed experiments

and analyzed data, MW performed statistical analyses and prepared figures, RG performed the

gene expression studies, MCP performed histological analyses, CAG, AM, SP and AM provided

reagents and concepts to study design. All authors edited the manuscript and consented to its

publication. FP oversaw the study and wrote the final draft of the manuscript.

Data Availability

The gene expression data are available through GEO at accession number GSE162210,

GSE162260, GSE162261 and GSE164922. These data are currently private and access can be

given to reviewers. The data previously published in (Wilk et al., 2015) are available at

GSE66040.

Methods

Cell culture. Cell lines were obtained from German Collection of Microorganisms and Cell Cultures GmbH (DSMZ), Braunschweig, Germany. The human myelomonocytic leukemia cell line THP-1 was propagated in RPMI 1640 medium (GIBCO$^®$ Life Technologies™) supplemented with 10% fetal calf serum (FCS) and 2 mM L-glutamine. Cells (2.5×10^5 cells/ml) were differentiated with 200 nM phorbol-12-myristate-13-acetate (PMA, Sigma-Aldrich, product no.P-8139) for 48 h, followed by incubation in fresh RPMI medium for another 24 hours (h). Human adenocarcinoma cells resembling type II alveolar epithelial cells (A549) were propagated in DMEM medium supplemented with 10% FCS and 2 mM L-glutamine.

Primary cell isolation and differentiation. Primary human monocytes were isolated from buffy coats from healthy blood donors by density gradient centrifugation (Histopaque™, Sigma-Aldrich). For studies of monocytes and M1/M2-type macrophages, CD14+ cells were isolated by magnetic activated cell sorting (CD14+ Cell Isolation Kit; Miltenyi Biotec). 2×10^5 monocytes were then differentiated into M1-type macrophages cells by incubation with 1000 U/ml GM-CSF (granulocyte macrophage-colony stimulating factor, CellGenix) and into M2-type macrophages by incubation with 100 ng/ml M-CSF (macrophage-colony stimulating factor, Miltenyi Biotec) in serum-free DC medium (CellGenix) for 5 days.

Virus strains. A common reference strain of influenza A virus (PR8M, A/PuertoRico/8/34 H1N1, kindly provided by Stefan Ludwig (University of Münster, Germany) (Blazejewska, Koscinski et al., 2011) was used throughout. In the experiment shown in Fig. 2, we additionally used the clinical isolate Gi-WT (A/Giessen/6/2009 H1N1-WT) and the reassortant Gi-NS-PR8 containing the NS segment of the PR8 strain on the backbone of SOIV-WT, which replicates faster than the WT (Petersen et al., 2018b). Viruses were propagated in the chorio-allantoic cavity of 10-day-old embryonated eggs for 48 h at 37°C. Fluid from the chorio-allantoic cavity was collected and the virus was titrated by focus-forming unit (FFU) assay and stored in aliquots at -80°C until use.

Viral infections and treatment with itaconate, DI, and 4OI. 2.5×10^5 dTHP-1 or A549 cells were infected with the indicated IAV strain at an MOI of 1, cells were centrifuged at 300 g for 15 min and incubated at 37°C for 2 h. Cells were subsequently washed twice with PBS and fresh

post-infection medium was added to the cells. Cells were re-incubated at 37°C for the indicated

lengths of time. The respective amount of IA, DI, or 4OI was added to RPMI/DMEM complete

medium and pH was adjusted to 7.5 with 1 M KOH solution. 4OI (100 mM) was dissolved in

DMSO, and the final concentration of 4OI (25 μM) thus resulted in a 0.025% concentration of

DMSO in the media. Media were filtered by using vacuum filters with 0.22 μm pore size

(Millipore). Cells were incubated overnight with the compounds, and virus was then added for 2

h. Unbound virus was then removed by replacing the medium with fresh medium containing the

compounds at the same concentrations. For PBMC infections, cells were isolated from 7 donors

and seeded as 4×10^6 cells/well in a 6 well plate and simultaneously infected for 12 h with strain

PR8M (MOI=1) with or without IA (10 mM), DI (0.5 mM), or 4OI (25 μM). Thus, PBMC were

not preincubated with the compounds before infection. Strain A/California/7/2009 (reagent

15/252, obtained from National Institute for Biological Standards and Control, London, UK) was

used for the mouse infections, applying 5×10^5 FFU (20 μL) by the intranasal route.

Mouse model. All animal procedures were performed following guidelines from the Federation

of European Laboratory Animal Science Associations. The study was approved by the regulatory

authority of the German Federal Sate of Lower Saxony (Niedersächsisches Landesamt für

Verbraucherschutz und Umwelt, LAVES). *ACOD1$^{-/-}$*, *ACOD1$^{+/-}$*, and *ACOD1$^{+/+}$* mice with

C57BL/6J background had originally been generated by Dr. Haruhiko Koseki in the RIKEN

Institute (Yokohama, Japan) by the use of stem cells which were purchased from the Repository

of Knockout Mouse Project under strain ID Irg1tm1a (KOMP) Wtsi. Age-matched groups of

female *ACOD1$^{-/-}$*, *ACOD1$^{+/-}$*, and *ACOD1$^{+/+}$* mice were derived from the same heterozygous

breeding pair. Animals were bred at the University of Luxembourg and then transferred to the

Animal Experimental Unit of the Helmholtz-Centre for Infection Research, where they were kept

under Animal Biosafety Level 2 and non-specific pathogen-free conditions. Animals were housed

seperated according to genotype, up to 5 mice per cage. For IAV infections, mice were sedated

by intraperitoneal injection of ketamine (10 mg/mL) and xylazine (1 mg/mL) in 0.9% NaCl,

infected intranasally and then observed for weight loss and survival for up to 15 days. Mice with

weight loss >20% had to be sacrificed (CO_2 asphyxiation) and were counted as non-survivors. In

a separate experiment, mice were sacrificed on days 8/9 and 14 p.i. (one mouse that died on day

11 was also included) and lungs examined for histopathological changes. Formalin-fixed,

paraffin-embedded sections of mouse lungs were stained with hematoxylin/eosin (H&E). To

assess the degree of inflammation and histopathological lesions, 10 parameters of inflammation

and tissue damage were scored by an expert in mouse pathology (MP), who was not aware of the

identity of the slides, according to the scale below.

Table 1. Scoring system to assess inflammation in IAV-infected mouse lung				
Parameter	0	1	2	3
Severity of infiltration of inflammatory cells	Not present	Mild	Moderate	Severe
Area involved	No changes	1-30%	30-70%	>70%

Human lung tissue explant model. The study was approved by the Ethics Committee of

Hannover Medical School (MHH), and all donors gave informed consent for experimental use of

their tissue. Tissue was obtained at the time of medically indicated lung transplantation from

patients with end-stage lung disease due to emphysema or pulmonary arterial hypertension. The

overall tissue quality and gross pathological changes were assessed by a board-certified

pathologist (Dept. of Pathology, Hannover Medical School). Healthy appearing tissue was then

dissected and sectioned into pieces of approx. 30–50 mm^3 (typical weight, 30-50 mg). Tissue

pieces were cultured overnight in individual wells in RPMI medium, containing DI (1 mM) or

itaconate (25 mM) as indicated. After this pre-treatment + washing step in medium, tissue pieces

were infected with IAV ($2x10^5$ FFU/ml) for 24 h at 37°C in medium containing DI or IA.

RT-qPCR. RNA was purified using the Nucleospin RNA purification kit (Machery Nagel) and

on-column removal of DNA by digestion with rDNase (Machery Nagel) for 15 min. at RT.

cDNA was synthesized with the PrimeScript™ kit (TaKaRa, Shiga, Japan) using 400 ng RNA in

a 10 µl reaction. RT-qPCR reactions were set up in a final volume of 20 µl, using the SensiFast™

SYBR® No-ROX Kit (Bioline, Taunton, MA) and the primers listed in **Table 2**. RT-qPCR was

performed in a LightCycler® 2.0 instrument (Roche, Mannheim, Germany), using 45 cycles of

the following program: 95°C for 15 sec., 60°C for 15 sec., and 72°C for 15 sec. To exclude

artefacts resulting from primer dimer formation, melting curve analysis was performed using the

sequence 95°C for 15 sec., 60°C for 15 sec., 95°C for 1 min. and 37°C for 30 sec. Relative

expression of the host mRNA targets was calculated using the $2^{-\Delta\Delta CT}$ method(Paijo, Döring et al.,

2016), and expression of IAV *HA* mRNA by normalizing against expression of HPRT as internal

control.

Table 2. List of RT-qPCR primers

Gene name	Primer name	Sequence
ISG15	ISG15-F	TGTCGGTGTCAGAGCTGAAG
	ISG15-R	AGAGGTTCGTCGCATTTGTC
IL6	IL6-F	CTACATTTGCCGAAGAGCCC
	IL6-R	CCCTGACCCAACCACAAATG
HPRT	HPRT-F	GAACGTCTTGCTCGAGATGTG
	HPRT-R	CCAGCAGGTCAGCAAAGAATT
IL1B	IL-1β-F	TACCCAAAGAAGAAGATGGAA
	IL-1β-R	GAGGTGCTGATGTACCAGTTG
CXCL10	CXCL10_F	CTGCTTTGGGGGTTTATCAGA
	CXCL10_R	CCACTGAAAGAATTTGGGC
A20	A20_F	ATGCACCGATACACACTGGA
	A20_R	CACAAGCTTCCGGACTTCTC
IL10	IL10-F	TACCTGGGTTGCCAAGCCT
	IL10-R	AGAAATCGATGACAGCGCC
HA	HA-F	CTCGTGCTATGGGGCATTCA
	HA-R	TTGCAATCGTGGACTGGTGT
IFNB1	IFNβ_F	CAGCAATTTTCAGTGTCAGAAGC
	IFNβ_R	TCATCCTGTCCTTGAGGCAGT
ACOD1	ACOD1_F	ATGCTGCTTTTGTGAACGGTG
	ACOD1-R	CTACCACGGAAGGGGGATGGA

Microarray analyses. RNA was extracted using the RNeasy kit (Qiagen), RNA quality was

checked with a Bioanalyzer 2100 (Agilent Technologies) and samples with RNA integrity

number >7 were used for microarray analysis. Total RNA/Poly-A RNA Control Mixture was

prepared by adding poly-A RNA controls (Affymetrix). This poly-A RNA was then used to

prepare double standard cDNA. Antisense RNA (complimentary RNA or cRNA) was then

synthesized and amplified by *in vitro* transcription (IVT) of the ds-cDNA template using T7

RNA polymerase. Enzymes, salts, inorganic phosphates, and unincorporated nucleotides were

removed to prepare the cRNA for 2nd-cycle ds-cDNA synthesis. After verifying quality and yield

of cRNA, sense-strand cDNA was synthesized by the reverse transcription of cRNA. cRNA

template was hydrolyzed leaving ds-cDNA, which was then purified. cDNA was fragmented and

labeled by terminal deoxynucleotidyl transferase (TdT) using the Affymetrix proprietary DNA

Labeling Reagent that is covalently linked to biotin. Cartridge Array Hybridization was

conducted on the GeneChip® Instrument (Affymetrix) using Thermo Fisher Scientific

microarrays (Clariom™ S Assay, human**).** Raw .CEL files from Clariom S Pico Assay

microarrays for hg.38 were imported into the Transcriptome Analysis Console (TAC4.0.2)

Software (ThermoFisher Scientific). Array QC and data normalization was performed using the

robust multiarray average (RMA) method(Irizarry, Hobbs et al., 2003). Differentially expressed

genes (DEGs) were identified using limma:Linear Models for microarray and RNA-Seq Data

(www.Bioconductor.org) and one-way ANOVA.

Differential gene expression analysis of bulk RNAseq data. Differential gene expression for

Fig. 1B was calculated using generalized linear models contained in the edgeR package

(doi:10.1093/bioinformatics/btp616). P values were adjusted using false discovery rate (FDR),

using FDR <0.05 to define significance of differential gene expression.

Single cell RNA sequencing

Barcoding and cDNA synthesis. The single cell suspension was loaded onto a well on a 10x

Chromium Single Cell instrument (10x Genomics). Barcoding and cDNA synthesis were

performed according to the manufacturer's instructions. Briefly, the 10x™ GemCode™

Technology partitions thousands of cells into nanoliter-scale Gel Bead-In-EMulsions (GEMs),

where all the cDNA generated from an individual cell share a common 10x Barcode. In order to

identify the PCR duplicates, Unique Molecular Identifier (UMI) was also added. The GEMs were

incubated with enzymes to produce full length cDNA, which was then amplified by PCR to

generate enough quantity for library construction. Quality was checked using the Agilent

Bioanalyzer High Sensitivity Assay.

Library construction and quality control. The cDNA libraries were constructed using the 10x

ChromiumTM Single cell 3' Library Kit according to the manufacturer's original protocol.

Briefly, after the cDNA amplification, enzymatic fragmentation and size selection were

performed using SPRI select reagent (Beckman Coulter, Cat# B23317) to optimize the

cDNA size. P5, P7, a sample index and read 2 (R2) primer sequence were added by end

repair, A-tailing, adaptor ligation and sample-index PCR. The final single cell 3' library

contains a standard Illumina paired-end constructs (P5 and P7), Read 1 (R1) primer sequence,

16 bp 10x barcode, 10 bp randomer, 89 bp cDNA fragments, R2 primer sequence and 8

bp sample index. For post library construction QC, 1 µl of sample was diluted 1:10

and ran on the Agilent Bioanalyzer High Sensitivity chip for qualitative analysis.

Single-cell RNA sequencing and generation of data matrix. PBMC were infected and treated as

for the microarray analyses. Cells were then stained with propidium iodide (Apoptosis Detection

Kit, eBioscience, cat. no. 88-8005-74), and dead cells and doublets were removed by FACS. Live

cells (including those undergoing apoptosis) were used to make single-cell suspensions using the

10x Genomics platform. Libraries were sequenced on an Illumina NovaSeq 6000 2x50 paired-

end kits using the following read length: 26 bp Read1 for cell barcode and UMI, 8 bp I7 index for

sample index and 89 bp Read2 for transcript. Cell Ranger 1.3 (http://10xgenomics.com) was used

to process Chromium single-cell 3' RNA-seq output. First, "cellranger mkfastq"

demultiplexed the sequencing samples based on the 8bp sample index read to generate fastq files

for the Read1 and Read2, followed by extraction of 16bp cell barcode and 10bp UMI. Second,

"cellranger count" aligned the Read2 to the pre-built human reference genome (hg19, GRCh38)

using STAR. Then, aligned reads were used to generate data matrix only when they had valid

barcodes and UMI, and mapped to exons (Ensembl GTFs GRCm38.p4) without PCR duplicates.

Valid cell barcodes were defined based on UMI distribution.

Single-cell gene expression data analysis including filtering, normalization, and clustering was

processed   using   Seurat   V3.1   (https://satijalab.org/seurat/)   in   the   R   environment

(www.bioconductor). Cells were removed when they had < 500 genes and > 25% reads mapped
to mitochondrial expression genome. After filtering and doublet removal, 6456 single cells
(control = 1870, DI =1474, IAV=1701, IAV & DI= 1411) remained. Raw counts were
normalized by the global-scaling normalization method "LogNormalize". Principal component
analysis score was used to determine the 'dimensionality' of all cells. Scatter plots were obtained
using the UMAP method. Cell clusters were identified as different cell populations based on the
expression of recently published canonical markers (**Table 3**). Differential gene expression was
determined for each cell type using DESeq2 (Love, Huber et al., 2014). IAV infected cells were
identified by the presence of IAV encoded transcripts, as mapped against NIAID Influenza
Research   Database   (Zhang,   Aevermann   et   al.,   2017),
www.fludb.org/brc/fluStrainDetails.spg?strainName=A/Giessen/6/2009(H1N1)&decorator=influ
enza. mRNA encoding the viral M protein could not be identified, likely due to a sequence
mismatch.

Table 3. mRNA markers used to define PBMC cell subpopulations in scRNAseq					
Cell type	**Sub-type**	**Marker(s)**	**Sub-sub type**	**Marker(s)**	**Ref.**
T cells	CD4	*CD3D, IL7R*	CD4 naïve	*LEF1, ATM, SELL, KLF2, ITGA6*	(Butler, Hoffman et al., 2018, Cano-Gamez, Soskic et al., 2020, Szabo, Levitin et al., 2019)
	CD4		Effector memory	*KLRB1 IL7R*	(Butler et al., 2018, Cano-Gamez et al., 2020)

	CD4		CD4 memory	*PASK*	(Butler et al., 2018, Cano-Gamez et al., 2020)
	CD4		Treg	*FOXP3*	(Li, Li et al., 2015)
	CD8	*CD3D, CD8A*	Effector memory	*GZMK*	(Butler et al., 2018, Chu, Berner et al., 2020)
	CD8		Cytotoxic	*GNLY*	(Butler et al., 2018, Szabo et al., 2019)
B cells		*MS4A1*	Activated B cells	*MIR155HG*	(Butler et al., 2018)
NK cells		*NKG7 GNLY*			(Butler et al., 2018)
Monocytes		*CD14, LYZ*			(Butler et al., 2018)

Enzyme-linked immunoassay. Supernatants from the same PBMC as used for RNA analyses
were collected 12 h p.i. Protein concentrations were measured using ELISA MAX Deluxe Sets
(Biolegend; IL1B cat. no. 437004; IL6, 430501; TNFa, 430201; IFNG, 430101) as per the
manufacturer's instructions. Briefly, capture antibody was used to pre-coat the wells O/N. After
removing the capture antibody and washing the plates, block buffer was applied for 1 h at room
temperature at 500 rpm. Again, plates were washed at least 4 times before adding the samples
(100 μL, diluted as needed) and standards for 2 h at RT. The plates were then washed again 4

times, and detection antibody was added to each well. The plate was further incubated for 1 h at

RT and then washed 4 times. Avidin-HRP conjugate was added and incubated for 30 min.

Substrate was added until color developed, and the samples were measured after 15-20 min at

450 and 570 nm.

Multiplex assay for inflammation-related polypeptides. Concentration of 27 inflammation-

related human proteins were measured in cell culture supernatants by using Human 27-plex

BIORAD cytokine panel (Cat. No.171-A1112, BIORAD). Standard curves were generated for all

27 cytokine standards with eight 2-fold serial dilutions starting from 32 ng/ml. 100 μl of assay

buffer was added to each well of a microfiltration plate, followed by addition of 50 μl of beads

suspension. After washing the beads with assay buffer, 50 μl of standard or the sample

(supernatant) was added to each well and incubated for 30 min at room temperature with gentle

shaking. 25 μl of antibody premix for detection was added into each well followed by incubation

for 30 min at room temperature with gentle shaking. Three washing steps were performed using

50 μl of assay buffer in each well, followed by washing with streptavidin solution for 10 min at

room temperature with shaking. Final washing with 125 μl of assay buffer was performed before

quantification by using BIO-PLEX Manager software version 4.

Immunofluorescense. THP-1 cells were seeded and differentiated with 80% confluency in a 24

well plate with coverslips. Cells were fixed with 2.5% formaldehyde for 20 min and the reaction

stopped by adding 0.1 M glycine for 5 minutes (min). Cells were then permeabilized with 0.2%

Triton X100 in PBS for 5 min. Cells were incubated for 1 h with primary antibody (rabbit-anti

p65 mAB, Cell Signaling, cat. no. 8242S) 1:200 diluted in BSA/PBS 1 mg/ml. Cells were then

washed three times with PBS and incubated in the dark for 45 min with Alexafluor594 secondary

antibody (1:100) (Cell Signaling, cat.# 8889S) in BSA/PBS 1 mg/ml and incubated in the dark for

45 min. Cells were washed three times with PBS. To stain nuclei, DAPI was added at 1:10,000 in

PBS for 10 min. Cells were washed three times with PBS and mounted with 20 μl of DAKO

mounting medium per coverslip on glass slides and stored in the dark until imaged. The images

were analysed using Image J s/w (imagej.net).

Quantification of itaconate. Itaconate concentrations were measured by liquid-chromatography

tandem mass-spectrometry (HPLC-MS/MS) using a liquid chromatography system (NEXERA

LC, Shimadzu, Japan) coupled to a triple quadrupole - ion trap mass spectrometer (QTRAP 5500,

Sciex, Framingham, MA) and Analyst 1.7 software (Sciex, Framingham, MA, USA), as

described in detail in (Kuhn, 2017).

Statistics. Data are shown mean ± standard error of the mean (SEM) unless stated otherwise.

Significance of between-group and across-group differences was assessed with the statistical tests

indicated in the figure legends, using Prism 5.02 and 8.0 (GraphPad Software Inc.). Wherever

exact p values are not stated, significance is graded as $*p<0.05$, $**p<0.01$, $***p<0.001$.

Principle component analysis was performed with the R package "PCAtools"

(https/github.com/kevinblighe/PCAtools), and **hierarchical clustering analysis** with the R tool

pheatmap (cran.r-project.org/web/packages/pheatmap/pheatmap.pdf).

Reanalysis of published data sets. Reanalyses of publically available published data sets were

performed in the R environment (version 3.6.3) (R_Core_Team, 2014), using package beeswarm

(version 0.2.3.) (Eklund, 2016) for visualization as indicated.

Gene set enrichment analysis. Features with linear fold change (FC) $\geq|1.5|$ and unadjusted P-

value ≤ 0.05 were used for enrichment analysis (Subramanian, Tamayo et al., 2005) based on

gene ontology (GO Biological Process) and KEGG pathways, using GEne SeT AnaLysis Toolkit

(Liao, Wang et al., 2019). Enrichment/depletion strip charts were made using ggplot2 (Wickham,

2016). Volcano plots to shows DEGs were made using R package "EnhancedVolcano"

(https://github.com/kevinblighe/EnhancedVolcano). Molecular Signatures Database (MSigDB)

v7.2 (Liberzon, Birger et al., 2015) was used to collect the information of genes involved in gene

set enrichment analyses.

Figure legends

Figure 1. Reciprocal interactions between *Acod1* expression and inflammation during
influenza A virus infection in mice. A. Higher pulmonary expression of *Acod1, Tnfaip3,* and
Hmox1 and *2* mRNA in DBA/2J (D2) than in C57BL/6J (B6) mice following IAV infection
(reanalysis of published RNAseq data (Wilk et al., 2015)) (n=3 each strain and time point).
*p<0.05; **p<0.01; ***p<0.001 (pairwise t-test with Bonferroni-Holm multiple comparisons
correction). **B.** *Acod1* mRNA is among the most highly upregulated transcripts in mouse lung
within 48 h of IAV infection (RNAseq, n=3 per time point). **C.** Lower pulmonary levels of
itaconate in IAV-infected *Acod1*$^{+/-}$ than in *Acod1*$^{+/+}$ mice. Itaconate was not detected in the
Acod1$^{-/-}$ mice (n=36 *Acod1*$^{+/+}$, 27 *Acod1*$^{+/-}$, 4 *Acod1*$^{-/-}$ mice). *p<0.05; **p<0.01; ***p<0.001
(unpaired t-test). **D.** Greater weight loss of female *Acod1*$^{-/-}$ C57BL/6J mice during infection with
IAV. Mice had to be euthanized when weight loss exceeded 20%; n=9 *Acod1*$^{+/+}$, 10 *Acod1*$^{+/-}$, 9
Acod1$^{-/-}$ mice. *p<0.05; **p<0.01; ***p<0.001 (unpaired t-test) **E.** Survival data of the same
experiment as in C. Median survival after day 7 was significantly lower in *Acod1*$^{-/-}$ mice
(p=0.018, Mann-Whitney-U test). **F.** Semi-quantitative histopathological scoring of lungs on days
8/9 (mid), 11 and 14 (late) reveals highest inflammation in *Acod1*$^{-/-}$ mice; n=9 *Acod1*$^{+/+}$, 10
Acod1$^{+/-}$, 9 *Acod1*$^{-/-}$ mice (all female). *p<0.05; **p<0.01; ***p<0.001 (two-way ANOVA,
Tukey's multiple comparisons test) **G.** H&E stained lung sections showing inflammation severity
scores of 0 (left) and 3 (right).

Figure 2. Differences among human myeloid cells in *ACOD1* mRNA induction during IAV
infection. A. Induction of *ACOD1* and *TNFAIP3* mRNA (RT-qPCR) in M1 and M2
macrophages and PBMCs during IAV infection. **B.** Induction of *ACOD1* and *TNFAIP3* (RT-
qPCR) in dTHP-1 cells correlates with virulence of three IAV strains. Mean ±SEM (n=4). *HA* =
IAV hemagglutinin. *p<0.05; **p<0.01; ***p<0.001 (two-way ANOVA, Tukey's multiple
comparisons test, with reference to M1 macrophages in A, and to Gi-WT in B).

Figure 3. Effects of itaconate and DI on cellular inflammation due to IAV infection of
dTHP-1 cells. A. Outline of experiment. dTHP-1 cells were treated with itaconate or DI
overnight, or left untreated, and then infected with IAV. Dose-response curves of effects of a
range of itaconate and DI and concentrations on *HA* and *CXCL10* mRNA expression are shown

in **Fig. S1. B**-**C**. 48 h time course, DI = 0.25 mM (n=3). mRNA expression (RT-qPCR) of *HA*

(**B**) and *CXCL10* mRNA (**C**) *p<0.05; **p<0.01; ***p<0.001 (unpaired t-test). **D-I.** Global

transcriptomic changes (microarray analysis) due to itaconate (25 mM) and DI (1 mM) treatment

of uninfected and IAV-infected dTHP-1 cells (n=3). **D.** Venn diagram showing unique and

common differentially expressed genes by itaconate and DI treatment of infected cells. **E-I.**

Volcano plots showing differential mRNA expression in uninfected dTHP-1 cells under itaconate

and DI treatment (**E-F**) and infected dTHP-1 cells with or without itaconate or DI treatment (**G-**

I). The 2-3 most significantly differentially expressed mRNAs are identified by labels. **J.** Effects

of itaconate (25 mM) and DI (1 mM) treatment on levels of inflammation-related polypeptides in

dTHP-1 cell supernatants 12 h after IAV infection or mock infection. Mean ± SEM (n=3).

*p<0.05; **p<0.01; ***p<0.001 (One-way ANOVA, Tukey's multiple comparison test).

Figure 4. Itaconate and DI treatment leads to a global downregulation of IFN-I signaling in

IAV-infected dTHP-1 cells. Enrichment analysis of gene ontology (GO Biological Process)

terms based on the microarray data shown in Fig. 3, using DEGs (p<0.05, FC >|1.5|) as input. GO

terms (FDR ≤0.05) reveal major induction of antiviral and inflammatory responses by infection,

which are decreased by both itaconate and DI treatment.

Figure 5. Effect of overexpression of ACOD1 and exogenous treatment with DI on IAV-

infected A549 cells. A. A549 cells were transfected with pCMV6-Hu-ACOD1, pCMV6 empty

vector, or mock transfected for 48 h and then infected with IAV for 48 h (n=3). Expression of the

indicated mRNAs was measured by RT-qPCR, itaconate concentrations by LC-MS/MS. *p<0.05;

p<0.01; ***p<0.001 (2-way ANOVA, Tukey's multiple comparisons test) **B. A549 cells were

treated with 0.25 mM DI overnight, infected with IAV for 48 h (n=3), and expression of the

indicated mRNAs measured by RT-qPCR. Mean ± SEM (n = 3). *p<0.05; **p<0.01; ***p<0.001

(2-way ANOVA, Sidak's multiple comparisons test).

Figure 6. GO term enrichment analysis of itaconate and DI reveals global effects on IFN-

signaling and cellular inflammation in A549. A549 cells were, in the case of treatment,

incubated overnight with DI (0.5 mM) or itaconate (20 mM), and then infected with strain PR8M.

Gene expression was determined by microarray analysis 24 h p.i. (n=4). **A**. GO enrichment (FDR

<0.05) analysis using DEGs (p<0.05, FC >|1.5|) as input. **B**. Hierarchical cluster analysis

illustrating downregulation of a subset of genes involved in IFNA/B signaling.

Fig. 7. Anti-inflammatory effect of itaconate and DI on IAV-infected human lung tissue

explants. Human primary lung tissue from patients with emphysema and pulmonary arterial

hypertension was incubated overnight with itaconate (25 mM), DI (1 mM), or medium only and

was then infected with IAV for 24 h (n=5 donors, 3 tissue pieces per donor). **A**. Viral titers in

supernatant (focus forming assay). **B-D**. *CXCL10* mRNA (**B**), and ISG15 (**C**) mRNA in tissue

(RT-qPCR); IP10 concentration in supernatant (EIA) (**D**). Mean ± SEM (n=6 donors, 3 tissue

pieces per replicate). *p<0.05; **p<0.01; ***p<0.001 (Mann-Whitney U test).

Fig. 8. Anti-inflammatory effects of itaconate, DI, and 4OI on IAV-infected human PBMCs.

PBMCs were isolated from freshly donated human blood and then infected with IAV for 12 h in

the presence or absence of itaconate (10 mM), DI (0.5 mM), or 4OI (25 μM) without

preincubation (7 donors, 3 replicates). **A.** Expression of the indicated mRNAs in PBMC (RT-

qPCR). **B.** Concentrations of the indicated proteins in PBMC supernatants (ELISA). Mean +/-

SEM (6 donors, 3 replicates per donor). *p<0.05; ***p<0.01; ***p<0.001 (1-way ANOVA). **C.**

GO enrichment analysis of effects of IA, DI, and 4OI on IAV-infected PBMC. Microarray

analysis of PBMC RNA from four of the donors featured in **A-B**. RNA was pooled from the three

replicates of each donor, resulting in 4 samples per group. GO term analysis was performed on

DEGs (p<0.05, FC>|1.5|) and terms with an FDR <0.02 in at least one condition are shown.

Fig. 9. Single-cell transcriptomic responses of PBMC to IAV infection and DI treatment.

PBMCs were isolated from freshly donated human blood (one donor), and then infected with

IAV for 12 h with or without DI (0.5 mM) in the medium. **A-B**. UMAP plots identifying 13 cell

types (**A**) that make up the five major cell types (**B**) studied. The RNA cell markers used are

listed in **Table 3**. **C-D.** Differential mRNA expression in uninfected, infected and infected/DI-

treated PBMCs. **E.** Differential mRNA expression in monocytes in the indicated paired

comparisons. Major transcriptional reprogramming due to IAV infection is evident, which is

largely prevented by treatment with DI. **F.** Differential mRNA expression in CD8+ T cells in the

indicated paired comparisons. Reprogramming in the presence of IAV is much less, but a DI

treatment effect is evident. **G-H.** Venn diagrams based on RNAs differentially regulated (FDR

<0.05) by both viral infection and DI treatment in each of the 5 major cell type, either irrespective

of FC (**G**) or FC >|2| (**H**).

Figure 10. GO enrichment analysis at the single cell level of DI effects on uninfected and

IAV-infected PBMC. Analysis based on the scRNAseq data shown in Fig. 9. Go terms that are

enriched/depleted in all cell types by infection and DI treatment are marked with a red asterix in

A and C. **A.** GO enrichment analysis of monocytes. **B.** Venn diagrams illustrating GO terms that

are commonly or uniquely regulated (FDR <0.05) in monocytes, CD4 T cells, CD8 T cells, B

cells, and NK cells; due to infection alone, DI treatment of infected cells, or DI treatment of

uninfected cells. Go terms that are enriched/depleted in all cell types by infection and DI

treatment are printed in bold. **C**. Go term analysis of CD4 T cells, CD8 T cells, B cells, and NK

cells.

Supplemental figure legends

Figure S1. Dose response curves of itaconate and DI treatment of IAV-infected dTHP-1

cells. dTHP-1 cells were pretreated overnight with the indicated concentrations of IA or DI,

infected with IAV, and expression of *HA* and *CXCL10* mRNA was measured after 12 h (RT-

qPCR) (n=3).

Figure S2. The impact of itaconate and DI treatments on cell transcriptomes is greater than

that of IAV infection. PCA based on the microarray analysis shown in Fig. 3. Untreated and

treated IAV-infected cells cluster together, whereas untreated infected cells are clearly separated

from all other groups in both principle components.

Figure S3. A. Hierarchical clustering analysis of effects of itaconate and DI on IAV-infected

dTHP-1 cells. Analysis based on the same microarray data as used for Fig. 3 and EV1B, using

the top 100 DEGs (FDR F-test <1.89E-09). Uninfected cells and untreated IAV-infected cells

form one clade and itaconate and DI treated IAV-infected cells the other clade. The brackets

identify (i) a large pro-inflammatory subclade in IAV infection, which is globally downregulated

by the treatments; (ii) a large cluster of DEGs specific for itaconate treatment; (iii) a small cluster

of DEGs specific for DI treatment, and (iv) a subclade comprising DEGs differentially expressed

due to both compounds. **B. Enrichment plots based on the 99 genes contained in GO term**
Response to type I interferon. Enrichment scores are plotted on the y-axis, the mRNAs
(identified by vertical lines) along the x-axis, ranked by fold change. The plots illustrate
pronounced enrichment in a large number of genes due to infection, which is nearly
quantitatively depleted by treatment with both itaconate and DI.

Figure S4. Itaconate and DI induce different recruitment programs in dTHP1 cells.
Supernatants of uninfected and IAV-infected dTHP1 cells, with our without itaconate or DI
treatment, were analyzed for concentrations of 27 cytokines/chemokines by multiplex microbead
array 12 h after infection or mock treatment. A hierarchical clustering analysis was carried out
with those targets that were detected above limit of detection of the assay in most samples. The
microarray analysis of cellular gene expression from the same experiment is shown in Fig. 4
(n=3).

Figure S5. KEGG-pathway analysis of effects of itaconate and DI treatment on IAV-
infected dTHP-1 cells. Analysis based on the microarray data set used for Figure 4. A broad
dampening of IFN-related and pro-inflammatory pathways by both compounds is evident.

Figure S6. Effects of increasing doses of itaconate and DI on *CXCL10* expression in IAV-
infected A549 cells. RT-qPCR using *HPRT1* as reference. A549 cells were infected with IAV
and itaconate and DI treatments were given at the indicated concentrations. Expression of *HA* and
CXCL10 mRNA was measured by RT-qPCR 24 h p.i. (n=3). DI concentrations $\geq$ 1 mM could not
be evaluated due to widespread cytotoxicity seen by light microscopy.

Figure S7. Global effects of itaconate and DI on transcriptomic responses in IAV-infected
A549 cells. PCA based on the microarray analysis used for Fig. 6, but additionally including
treatments with 40 mM itaconate and 0.25 mM DI. PCA showing normalization of IAV-driven
reprogramming of gene expression, but increasing impact of itaconate on cellular responses with
increasing doses.

Figure S8. Effects of itaconate (20 mM) and DI (0.5 mM) on transcriptomic responses in
IAV-infected A549 cells. Analysis based on the microarray data used for the GO analysis in Fig.
6. Unsupervised hierarchical clustering analysis of the 100 most significant DEGs (FDR F-test

<1.46E-10). DI-treated infected and uninfected cells cluster in one clade, and infected and

itaconate-treated infected cells in the other. There is a large clade mostly containing

proinflammatory genes, which are downregulated by DI, whereas effects of itaconate are much

weaker. *DUSP95* is exclusively downregulated by itaconate (arrowhead). There are two smaller

clades of genes that form an "itaconate signature", comprising genes that are, albeit to a lesser

extent, also upregulated by IAV infection, or uniquely by itaconate.

Figure S9. Effects of itaconate (40 mM) and DI (0.5 mM) on transcriptomic responses in

IAV-infected A549 cells. Hierarchical clustering analysis ased on the microarray analysis used

for Fig. S5. Compared to the 20 mM itaconate concentration (**Fig. S8**), itaconate-treated IAV

infection is now in a separate clade, indicating that the impact of itaconate on the cells dominates

that of the infection. Downregulation of the inflammation-driven clade is now more pronounced,

the itaconate-unique signature is stronger, and there also is a clade comprised of genes

downregulated by IAV.

Figure S10. KEGG pathway enrichment analysis of itaconate (20 mM) and DI (0.5 mM)

treatment on IAV infection in A549 cells. Analysis performed based on the microarray data

shown in **Fig. 6** and **S8.**

Figure S11. Primary human lung tissue explants support IAV RNA transcription and

upregulation of CXCL10 mRNA. Human primary lung tissue from patients with emphysema or

pulmonary arterial hypertension (n=7 donors, 3 tissue pieces per donor per treatment) was

infected with IAV for 72 h and *HA* and *CXCL10* mRNA expression measured by RT-qPCR,

using *HPRT1* as internal reference. *CXCL10* levels are expressed as fold change with reference to

uninfected tissue at the same time point. *p<0.05; **p<0.01; ***p<0.001 (Mann-Whitney U

test).

Figure S12. Induction of *ACOD1* and *TNFAIP3* mRNAs during IAV (H3N2) infection of

human lung explants. Reanalysis of a published dataset of gene expression (RNAseq) in IAV

(H3N2) infection of human lung tissue derived from tumor-free margins obtained during

lobectomy for lung carcinoma(Matos et al., 2019). A strong induction of *ACOD1* expression and

a tendency towards increased *TNFAIP3* expression are seen. As opposed to the strong induction

of both genes in the mouse model (**Fig. 1A**), *HMOX1* and *HMOX2* expression is unchanged.

*p<0.05; **p<0.01; ***p<0.001 (pairwise t-tests with pooled SD).

Figure S13. Increased levels of *ACOD1* and *TNFAIP3* mRNA expression in whole blood

from patients with moderate and severe influenza. Reanalysis of a published data set of gene

expression in whole blood from patients with moderate and severe influenza and healthy

controls(Tang, Shojaei et al., 2017). *p<0.05; **p<0.01; ***p<0.001 (pairwise t-tests with

pooled SD).

Figure S14. Transcriptome changes in itaconate, DI, and 4OI treated PBMC are driven

more by the treatments than by infection. Analysis of mRNA expression in a subgroup (n=4)

of the PBMC samples that were used for the targeted assays shown in Fig. 8. A PCA was

performed on the same oligonucleotide microarray data as used for Fig. 8C and J. There is a less

pronounced effect of IAV infection (control and IAV infected samples are not clearly separated)

than on dTHP-1 and A549 cells, but pronounced additional broad changes in gene expression due

to treatment with 4OI, itaconate treatment (note two outliers), and DI (one outlier).

Figure S15. Hierarchical clustering analysis of transcriptomic changes in IAV infected

PBMC and responses to itaconate, DI, and 4OI treatment. The 100 most significant DEGs

were selected (FDR F-test <3.72E-05). Transcriptome changes are mostly due to marked effects

of 4OI (more down- than upregulation) on genes that are not affected by IAV infection,

indicating general effects on cell homeostasis. However, induction of *HMOX1* by DI in IAV

infection is evident (red arrowhead). The blue arrowhead points to the extreme outlier under DI

treatment seen in the PCA.

Figure S16. KEGG pathway analysis of transcriptomic changes in IAV infected PBMC and

responses to itaconate, DI, and 4OI treatment. KEGG terms with an FDR <0.05 in at least one

group were selected. The threshold for inclusion in the chart was raised to FDR <0.01 if a

pathway was enriched/depleted only in a single treatment and not by IAV infection alone (e.g.,

Alcoholism in 4OI treatment).

Figure S17. Effects of IAV infection and DI treatment on relative cell type distribution in

PBMC. A reduction of CD14+ monocytes is apparent under DI treatment of both uninfected and

infected PBMC. **A.** Percentages. **B.** Absolute numbers.

Figure S18. Effects of IAV infection and DI treatment on transcriptomes in the PBMC cell

types not shown in Fig. 9. Analysis based on scRNAseq. Volcano plots comparing effects of DI

treatment, with and without IAV infection, on CD4+ cells, NK cells, and B cells.

Figure S19. Cell-type specific KEGG pathway analysis of effects of DI on infected and

uninfected monocytes, CD4 T cells, CD8 T cells, B cells, and NK cells. Analyses based on the

scRNAseq data used for Fig. 9 and 10. Pathways that are enriched (FDR <0.05) in at least one

cell type are shown.

Figure S20. Identification of central pathways commonly regulated in monocytes, CD4 and

CD8 T cells, B cells, and NK cells in response to IAV infection and DI treatment. GO

Biological Process (A) and KEGG (B) functional enrichment analyses of all genes commonly

differentially expressed in monocytes, CD4 and CD8 T cells, B cells, and NK cells in response to

IAV infection and DI treatment (center of Venn diagram shown in **Fig. 9H,** based on scRNAseq

analysis of PBMC)**.**

References

Alberts R, Srivastava B, Wu H, Viegas N, Geffers R, Klawonn F, Novoselova N, do Valle TZ, Panthier JJ,
Schughart K (2010) Gene expression changes in the host response between resistant and susceptible
inbred mouse strains after influenza A infection. *Microbes Infect* 12: 309-18

Bambouskova M, Gorvel L, Lampropoulou V, Sergushichev A, Loginicheva E, Johnson K, Korenfeld D,
Mathyer ME, Kim H, Huang LH, Duncan D, Bregman H, Keskin A, Santeford A, Apte RS, Sehgal R, Johnson
B, Amarasinghe GK, Soares MP, Satoh T et al. (2018) Electrophilic properties of itaconate and derivatives
regulate the IkappaBzeta-ATF3 inflammatory axis. *Nature* 556: 501-504

Blazejewska P, Koscinski L, Viegas N, Anhlan D, Ludwig S, Schughart K (2011) Pathogenicity of different
PR8 influenza A virus variants in mice is determined by both viral and host factors. *Virology* 412: 36-45

Butler A, Hoffman P, Smibert P, Papalexi E, Satija R (2018) Integrating single-cell transcriptomic data
across different conditions, technologies, and species. *Nat Biotechnol* 36: 411-420

Cano-Gamez E, Soskic B, Roumeliotis TI, So E, Smyth DJ, Baldrighi M, Wille D, Nakic N, Esparza-Gordillo J,
Larminie CGC, Bronson PG, Tough DF, Rowan WC, Choudhary JS, Trynka G (2020) Single-cell
transcriptomics identifies an effectorness gradient shaping the response of CD4(+) T cells to cytokines.
Nat Commun 11: 1801

Chen F, Lukat P, Iqbal AA, Saile K, Kaever V, van den Heuvel J, Blankenfeldt W, Bussow K, Pessler F (2019)
Crystal structure of cis-aconitate decarboxylase reveals the impact of naturally occurring human
mutations on itaconate synthesis. *Proc Natl Acad Sci U S A* 116: 20644-20654

Cho H, Proll SC, Szretter KJ, Katze MG, Gale M, Jr., Diamond MS (2013) Differential innate immune
response programs in neuronal subtypes determine susceptibility to infection in the brain by positive-
stranded RNA viruses. *Nat Med* 19: 458-64

Chu T, Berner J, Zehn D (2020) Two parallel worlds of memory T cells. *Nat Immunol* 21: 1484-1485

Collaborators GI (2018) Mortality, morbidity, and hospitalisations due to influenza lower respiratory tract
infections, 2017: an analysis for the Global Burden of Disease Study 2017. *Lancet Resp Med* 7: 69-89

Cordes T, Lucas A, Divakaruni AS, Murphy AN, Cabrales P, Metallo CM (2020) Itaconate modulates
tricarboxylic acid and redox metabolism to mitigate reperfusion injury. *Mol Metab* 32: 122-135

Cordes T, Michelucci A, Hiller K (2015) Itaconic Acid: The Surprising Role of an Industrial Compound as a
Mammalian Antimicrobial Metabolite. *Annu Rev Nutr* 35: 451-73

Cordes T, Wallace M, Michelucci A, Divakaruni AS, Sapcariu SC, Sousa C, Koseki H, Cabrales P, Murphy
AN, Hiller K, Metallo CM (2016) Immunoresponsive Gene 1 and Itaconate Inhibit Succinate
Dehydrogenase to Modulate Intracellular Succinate Levels. *J Biol Chem* 291: 14274-84

Daniels BP, Kofman SB, Smith JR, Norris GT, Snyder AG, Kolb JP, Gao X, Locasale JW, Martinez J, Gale M,
Jr., Loo YM, Oberst A (2019) The Nucleotide Sensor ZBP1 and Kinase RIPK3 Induce the Enzyme IRG1 to
Promote an Antiviral Metabolic State in Neurons. *Immunity* 50: 64-76 e4

Duan M, Hibbs ML, Chen W (2016) The contributions of lung macrophage and monocyte heterogeneity
to influenza pathogenesis. *Immunol Cell Biol* 95: 225-235

Gu L, Lin J, Wang Q, Li C, Peng X, Fan Y, Lu C, Lin H, Niu Y, Zhu G, Zhao G (2020) Dimethyl itaconate
protects against fungal keratitis by activating the Nrf2/HO-1 signaling pathway. *Immunol Cell Biol* 98:
229-241

Hooftman A, O'Neill LAJ (2019) The Immunomodulatory Potential of the Metabolite Itaconate. *Trends*
Immunol

Irizarry RA, Hobbs B, Collin F, Beazer-Barclay YD, Antonellis KJ, Scherf U, Speed TP (2003) Exploration,
normalization, and summaries of high density oligonucleotide array probe level data. *Biostatistics* 4: 249-
64

Kuhn M (2017) PhD dissertation. Hannover Medical School

Lampropoulou V, Sergushichev A, Bambouskova M, Nair S, Vincent EE, Loginicheva E, Cervantes-
Barragan L, Ma X, Huang SC, Griss T, Weinheimer CJ, Khader S, Randolph GJ, Pearce EJ, Jones RG, Diwan
A, Diamond MS, Artyomov MN (2016) Itaconate Links Inhibition of Succinate Dehydrogenase with
Macrophage Metabolic Remodeling and Regulation of Inflammation. *Cell Metab* 24: 158-66

Lee JS, Park S, Jeong HW, Ahn JY, Choi SJ, Lee H, Choi B, Nam SK, Sa M, Kwon JS, Jeong SJ, Lee HK, Park
SH, Park SH, Choi JY, Kim SH, Jung I, Shin EC (2020) Immunophenotyping of COVID-19 and influenza
highlights the role of type I interferons in development of severe COVID-19. *Sci Immunol* 5

Li Y, Zhang P, Wang C, Han C, Meng J, Liu X, Xu S, Li N, Wang Q, Shi X, Cao X (2013) Immune responsive
gene 1 (IRG1) promotes endotoxin tolerance by increasing A20 expression in macrophages through
reactive oxygen species. *J Biol Chem* 288: 16225-34

Li Z, Li D, Tsun A, Li B (2015) FOXP3+ regulatory T cells and their functional regulation. *Cell Mol Immunol*
12: 558-65

Liao Y, Wang J, Jaehnig EJ, Shi Z, Zhang B (2019) WebGestalt 2019: gene set analysis toolkit with
revamped UIs and APIs. *Nucleic Acids Res* 47: W199-W205

Liberzon A, Birger C, Thorvaldsdottir H, Ghandi M, Mesirov JP, Tamayo P (2015) The Molecular Signatures
Database (MSigDB) hallmark gene set collection. *Cell Syst* 1: 417-425

Liu H, Feng Y, Xu M, Yang J, Wang Z, Di G (2018) Four-octyl itaconate activates Keap1-Nrf2 signaling to
protect neuronal cells from hydrogen peroxide. *Cell Commun Signal* 16: 81

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. *Genome Biol* 15: 550

Matos ADR, Wunderlich K, Schloer S, Schughart K, Geffers R, Seders M, Witt M, Christersson A, Wiewrodt
R, Wiebe K, Barth P, Hocke A, Hippenstiel S, Honzke K, Dittmer U, Sutter K, Rescher U, Rodionycheva S,
Matera N, Ludwig S et al. (2019) Antiviral potential of human IFN-alpha subtypes against influenza A
H3N2 infection in human lung explants reveals subtype-specific activities. *Emerg Microbes Infect* 8: 1763-
1776

Michelucci A, Cordes T, Ghelfi J, Pailot A, Reiling N, Goldmann O, Binz T, Wegner A, Tallam A, Rausell A,
Buttini M, Linster CL, Medina E, Balling R, Hiller K (2013) Immune-responsive gene 1 protein links
metabolism to immunity by catalyzing itaconic acid production. *Proc Natl Acad Sci U S A* 110: 7820-5

Mills EL, Ryan DG, Prag HA, Dikovskaya D, Menon D, Zaslona Z, Jedrychowski MP, Costa ASH, Higgins M,
Hams E, Szpyt J, Runtsch MC, King MS, McGouran JF, Fischer R, Kessler BM, McGettrick AF, Hughes MM,
Carroll RG, Booty LM et al. (2018) Itaconate is an anti-inflammatory metabolite that activates Nrf2 via
alkylation of KEAP1. *Nature* 556: 113-117

Muri J, Wolleb H, Broz P, Carreira EM, Kopf M (2020) Electrophilic Nrf2 activators and itaconate inhibit
inflammation at low dose and promote IL-1beta production and inflammatory apoptosis at high dose.
Redox Biol 36: 101647

Nair S, Huynh JP, Lampropoulou V, Loginicheva E, Esaulova E, Gounder AP, Boon ACM, Schwarzkopf EA,
Bradstreet TR, Edelson BT, Artyomov MN, Stallings CL, Diamond MS (2018) Irg1 expression in myeloid
cells prevents immunopathology during M. tuberculosis infection. *J Exp Med* 215: 1035-1045

O'Neill LAJ, Artyomov MN (2019) Itaconate: the poster child of metabolic reprogramming in macrophage
function. *Nat Rev Immunol* 19: 273-281

Olagnier D, Farahani E, Thyrsted J, Blay-Cadanet J, Herengt A, Idorn M, Hait A, Hernaez B, Knudsen A,
Iversen MB, Schilling M, Jorgensen SE, Thomsen M, Reinert LS, Lappe M, Hoang HD, Gilchrist VH, Hansen
AL, Ottosen R, Nielsen CG et al. (2020) SARS-CoV2-mediated suppression of NRF2-signaling reveals
potent antiviral and anti-inflammatory activity of 4-octyl-itaconate and dimethyl fumarate. *Nat Commun*
11: 4938

Paijo J, Döring M, Spanier J, Grabski E, Nooruzzaman M, Schmidt T, Witte G, Messerle M, Hornung V,
Kaever V (2016) cGAS senses human cytomegalovirus and induces type I interferon responses in human
monocyte-derived cells. *PLoS Pathog* 12: e1005546

Passalacqua KD, Purdy JG, Wobus CE (2019) The inert meets the living: The expanding view of metabolic
alterations during viral pathogenesis. *PLoS Pathog* 15: e1007830

Petersen H, Mostafa A, Tantawy MA, Iqbal AA, Hoffmann D, Tallam A, Selvakumar B, Pessler F, Beer M,
Rautenschlein S (2018a) NS Segment of a 1918 Influenza A Virus-Descendent Enhances Replication of
H1N1pdm09 and Virus-Induced Cellular Immune Response in Mammalian and Avian Systems. *Front*
Microbiol 9: 526

Preusse M, Schughart K, Pessler F (2017) Host Genetic Background Strongly Affects Pulmonary microRNA
Expression before and during Influenza A Virus Infection. *Front Immunol* 8: 246

Preusse M, Tantawy MA, Klawonn F, Schughart K, Pessler F (2013) Infection- and procedure-dependent
effects on pulmonary gene expression in the early phase of influenza A virus infection in mice. *BMC*
Microbiol 13: 293

Ragab D, Salah Eldin H, Taeimah M, Khattab R, Salem R (2020) The COVID-19 Cytokine Storm; What We
Know So Far. *Front Immunol* 11: 1446

Ramos I, Fernandez-Sesma A (2015) Modulating the Innate Immune Response to Influenza A Virus:
Potential Therapeutic Use of Anti-Inflammatory Drugs. *Front Immunol* 6: 361

Srivastava B, Blazejewska P, Hessmann M, Bruder D, Geffers R, Mauel S, Gruber AD, Schughart K (2009)
Host genetic background strongly influences the response to influenza A virus infections. *PLoS ONE* 4:
e4857

Strelko CL, Lu W, Dufort FJ, Seyfried TN, Chiles TC, Rabinowitz JD, Roberts MF (2011) Itaconic acid is a
mammalian metabolite induced during macrophage activation. *J Am Chem Soc* 133: 16386-16389

Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL,
Golub TR, Lander ES, Mesirov JP (2005) Gene set enrichment analysis: a knowledge-based approach for
interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* 102: 15545-50

Szabo PA, Levitin HM, Miron M, Snyder ME, Senda T, Yuan J, Cheng YL, Bush EC, Dogra P, Thapa P, Farber
DL, Sims PA (2019) Single-cell transcriptomics of human T cells reveals tissue and activation signatures in
health and disease. *Nat Commun* 10: 4706

Tang BM, Shojaei M, Parnell GP, Huang S, Nalos M, Teoh S, O'Connor K, Schibeci S, Phu AL, Kumar A, Ho
J, Meyers AFA, Keynan Y, Ball T, Pisipati A, Kumar A, Moore E, Eisen D, Lai K, Gillett M et al. (2017) A

novel immune biomarker IFI27 discriminates between influenza and bacteria in patients with suspected
respiratory infection. *Eur Respir J* 49

Tang C, Wang X, Xie Y, Cai X, Yu N, Hu Y, Zheng Z (2018) 4-Octyl Itaconate Activates Nrf2 Signaling to
Inhibit Pro-Inflammatory Cytokine Production in Peripheral Blood Mononuclear Cells of Systemic Lupus
Erythematosus Patients. *Cell Physiol Biochem* 51: 979-990

Uddin MJ, Joe Y, Kim S-K, Jeong SO, Ryter SW, Pae H-O, Chung HT (2016) IRG1 induced by heme
oxygenase-1/carbon monoxide inhibits LPS-mediated sepsis and pro-inflammatory cytokine production.
Cell Mol Immunol 13: 170

van Paassen J, Vos JS, Hoekstra EM, Neumann KMI, Boot PC, Arbous SM (2020) Corticosteroid use in
COVID-19 patients: a systematic review and meta-analysis on clinical outcomes. *Crit Care* 24: 696

Wickham H (2016) *Elegant Graphics for Data Analysis*. Springer Verlag, New York

Wilk E, Pandey AK, Leist SR, Hatesuer B, Preusse M, Pommerenke C, Wang J, Schughart K (2015) RNAseq
expression analysis of resistant and susceptible mice after influenza A virus infection identifies novel
genes associated with virus replication and important for host resistance to infection. *BMC Genomics* 16:
655

Xu M, Jiang P, Sun H, Yuan X, Gao S, Guo J, Zhao C, Hu X, Liu X, Fu Y (2020a) Dimethyl itaconate protects
against lipopolysaccharide-induced endometritis by inhibition of TLR4/NF-kappaB and activation of
Nrf2/HO-1 signaling pathway in mice. *Iran J Basic Med Sci* 23: 1239-1244Zhang Y, Aevermann BD,
Anderson TK, Burke DF, Dauphin G, Gu Z, He S, Kumar S, Larsen CN, Lee AJ, Li X, Macken C, Mahaffey C,
Pickett BE, Reardon B, Smith T, Stewart L, Suloway C, Sun G, Tong L et al. (2017) Influenza Research
Database: An integrated bioinformatics resource for influenza virus research. *Nucleic Acids Res* 45: D466-
d474

Zhao C, Jiang P, He Z, Yuan X, Guo J, Li Y, Hu X, Cao Y, Fu Y, Zhang N (2019) Dimethyl itaconate protects
against lippolysacchride-induced mastitis in mice by activating MAPKs and Nrf2 and inhibiting NF-kappaB
signaling pathways. *Microb Pathog* 133: 103541

1028

Figure 1

Figure 2

Figure 3

dTHP-1 cells: GO Biological Process (FDR < 0.05)

Figure 5

Figure 6

A

A549 cells: GO Biological Process (FDR < 0.05)

B

Figure 7

Figure 8

Figure 9

Figure 10

Figure S1

Figure S2

dTHP-1 cells

- 🔴 1 mM DI / Uninfected
- 🟡 1 mM DI / Infection
- 🟢 25 mM Ita / Uninfected
- 🟦 25 mM Ita / Infection
- 🔵 Uninfected
- 🟣 Infection

Untreated / Infection

25 mM Ita / Infection

25 mM Ita / Uninfected

Uninfected

1 mM DI / Infection

1 mM DI / Uninfected

PC1, 44.7% variation

PC2, 19.81% variation

Figure S3

Figure S4

dTHP-1 supernatants

Figure S5

dTHP-1 cells, KEGG (FDR<0.05)

Figure S7

A549 cells

- 🔴 0.25 mM DI / Uninfected
- 🟠 0.25 mM DI / Infection
- 🟡 0.5 mM DI / Uninfected
- 🟢 0.5 mM DI / Infection
- 🟢 20 mM Ita / Uninfected
- 🔵 20 mM Ita / Infection
- 🔵 40 mM Ita / Uninfected
- 🟣 40 mM Ita / Infection
- 🟣 Uninfected
- 🔴 Infection

Infection

20 mM Ita / Infection

0.25 mM DI / Infection

Uninfected

40 mM Ita / Infection

0.5 mM DI / Infection

20 mM Ita / Uninfected

0.5 mM DI / Uninfected

40 mM Ita / Uninfected

0.25 mM DI / Uninfected

PC1, 33.51% variation

PC2, 20.96% variation

Figure S8

Figure S9

Figure S10

A549 cells, KEGG (FDR<0.05)

Figure S12

Figure S13

Figure S14

PBMC

PC2, 18.7% variation

DI / Infection

Infection

Uninfected

Ita / Infection

4OI / Infection

-25   0   25   100

PC1, 30.33% variation

Figure S15

Figure S16

PBMC, KEGG (FDR<0.05)

A Effects of IAV infection and DI treatment on relative cell type distribution in PBMCs

IAV	-	-	+	+
DI	-	+	-	+

B Absolute numbers of analyzed cells

	IAV	-	-	+	+
	DI	-	+	-	+
CD4+ T		823	664	743	686
NK		283	205	216	229
CD8+ T		530	439	525	537
B		160	136	139	114
Monocytes		74	30	78	25
Sum		1870	1474	1701	1591

Figure S18

PBMC, KEGG (FDR < 0.05)

Figure S20

A
PBMC, GO Biological Process

B
PBMC, KEGG