Itaconate and derivatives reduce interferon response and inflammation

in influenza A virus infection

Aaqib Sohail[1,9*], Azeem A. Iqbal[1,9*], Nishika Sahini[1,9*], Mohamed Tantawy[1,9], Moritz

Winterhoff[1,9], Thomas Ebensen[2] , Robert Geffers[3], Klaus Schughart[4], Fangfang Chen[1,9], Matthias

Preusse[1,2], Marina C. Pils[5], Carlos A. Guzman[2], Ahmed Mostafa[6], Stephan Pleschka[6], Christine

Falk[7], Alessandro Michelucci[8], Frank Pessler[1,9,**]

[1]Biomarkers for Infectious Diseases, [2]Vaccinology and Applied Microbiology, [3]Genome

Analytics, [4]Infection Genetics, [5]Mouse Pathology Platform, Helmholtz Centre for Infection

Research, 31824 Braunschweig, Germany

[6]Institute for Medical Virology, Justus-Liebig-University, 35392 Giessen, Germany

[7]Department of Transplantation Immunology, Hannover Medical School, 30625 Hannover,

Germany

[8]Dept. of Oncology, Luxembourg Institute of Health, 1526 Luxembourg

[9]TWINCORE Centre for Experimental and Clinical Infection Research, 30625 Hannover,

Germany

*Joint first authors

**Corresponding author

Frank Pessler, MD, PhD

Tel. +49 511 220027-167, Mobile +49 160 9728 0364

frank.pessler@twincore.de

Total character count 59381

Abstract

Itaconate has recently emerged as a metabolite with immunomodulatory properties. We evaluated effects of endogenous itaconate and exogenous itaconate, dimethyl-, and 4-octyl-itaconate on host responses to influenza A virus infection. Infection induced *ACOD1* (the enzyme that catalyzes itaconate synthesis) mRNA in monocytes and macrophages, which correlated with viral replication and was abrogated by itaconate treatment. Weight loss and pulmonary inflammation were greater in *Acod1*$^{-/-}$ than wild-type mice. Investigations in cell lines, human lung tissue, and peripheral blood mononuclear cells revealed that the compounds reversed infection-triggered interferon responses and pro-inflammatory gene expression. Ectopic synthesis of itaconate in epithelial cells also reduced inflammation. Single-cell RNA sequencing of PBMC showed that infection induced *ACOD1* exclusively in monocytes, whereas dimethyl-itaconate treatment silenced IFN-responses in monocytes, lymphocytes, and natural killer cells. Viral replication did not increase under itaconate and dimethyl-itaconate treatment despite the dramatically repressed IFN responses, but 4-octyl itaconate inhibited viral transcription in PBMC. The results reveal the potential of itaconate derivatives as adjunct treatments for deleterious host responses in infection by influenza viruses and potentially other viral pathogens.

Key words: innate immunity - influenza virus – pneumonia – single cell RNA sequencing - treatment.

Introduction

As exemplified by influenza viruses (Ramos & Fernandez-Sesma, 2015) and pathogenic coronaviruses (Ragab et al., 2020), overshooting host inflammation, oxidative stress and cell damage/death are major determinants of morbidity and mortality in severe viral infections of humans. Consequently, there is a great need for adjunct treatments that can be given in conjunction with direct-acting anti-virals, with the aim to modulate the host response in such a way that end-organ damage is reduced and clinical outcome is improved.

Itaconic acid initially attracted great interest in the field of biotechnology due to its highly reactive double bonds, which make it an efficient intermediate in the biosynthesis of industrial polymers (reviewed in (Cordes et al., 2015)). The discoveries that its synthesis is highly induced during macrophage activation (Strelko et al., 2011) and that *immune response gene 1* (*Irg1,* since then renamed *aconitate decarboxylase*) encodes the enzyme cis-aconate decarboxylase (ACOD1 in humans, CAD in other organisms), which converts cis-aconitate to itaconate (Chen et al., 2019, Michelucci et al., 2013), have triggered intense efforts to elucidate functions of the endogenous ACOD1/itaconate axis in inflammation and infection. The early recognition that itaconate possesses anti-inflammatory and immunomodulatory properties also suggested that itaconate and prodrugs derived from it may have potential as immunomodulatory and cytoprotective interventions for inflammation-driven diseases (reviewed in (Hooftman & O'Neill, 2019, O'Neill & Artyomov, 2019)).

Most evidence of anti-inflammatory and immunomodulatory effects of itaconate has actually been obtained by treating cells or small animals with chemically modified variants that are presumably taken up more efficiently. 4-Octyl-itaconate (4OI) contains an added 8-carbon chain and acts primarily through activation of the NRF2 pathway (Mills 2018). It has, for instance, shown cytoprotective and anti-inflammatory effects in a variety of models, including lipopolysaccharide (LPS)-induced macrophage activation and LPS-induced sepsis in mice (Mills 2018), in a neuronal model of oxidative stress (Liu 2018), and in peripheral blood mononuclear cells (PBMC) from patients with systemic lupus erythematosus (Tang 2018). Dimethyl-itaconate (DI) has also been shown to exert NRF2 independent anti-inflammatory effects by inhibiting IκBζ activity (Bambouskova 2018). Furthermore, DI possesses cytoprotective effects, which was, for instance, shown by reduced cell damage in mouse models of cardiac ischemia

(Lampropoulou 2016). However, cytoprotective properties have also been demonstrated for
unmodified itaconate, notably in a rat model of cerebral ischemic-reperfusion injury (Cordes
2020). Both compounds likely exert cytoprotective effects by reducing reactive oxygen species
(ROS) generation via inhibition of succinate dehydrogenase (Cordes 2020, Cordes 2016,
Lampropoulou 2016). Considering these varied effects of itaconate and derivatives, it is not
surprising that administration of DI has demonstrated beneficial effects in a variety of mouse
models of infection-associated inflammation such as LPS-induced mastitis (Zhao 2019) and
endometritis (Xu 2020b), and fungal keratitis (Gu 2020). Following early evidence that *Acod1*
transcription is strongly induced in severe influenza A virus (IAV) infection in mice (Preusse
2013), its impact on host susceptibility to viral infections has also been studied, and protective
effects of itaconate on neurons exposed to RNA viruses have been reported (Cho 2013, Daniels
2019). On the other hand, in a recent study using a mouse model of IAV infection, deleting the
Acod1 locus did not affect weight loss or survival, whereas it clearly worsened the outcome of
Mycobacterium tuberculosis infection in the same mouse strain (Nair 2018). This lack of effect
on IAV infection was surprising because abnormally increased inflammation is usually asociated
with deleterious effects on the host during viral infections. For instance, it has been shown that
overactive inflammatory responses are associated with decreased survival and increased weight
loss in IAV infected inbred mouse strains (Alberts, Srivastava et al., 2010). While the role of
endogenous itaconate in protection from viral infections thus remains incompletely understood,
the above results do suggest a strong potential of external application of itaconate or its
derivatives to ameliorate inflammation and other deleterious consequences of viral infections.
Focusing on IAV as one of the most important human respiratory viral pathogens (Collaborators,
2018), we therefore examined role and regulation of the endogenous ACOD1/itaconate axis in a
mouse model of IAV infection and subsequently performed a detailed analysis of the modulatory
effects of exogenous addition of itaconate and derivatives on inflammatory responses in IAV-
infected human cell lines, explanted lung tissue, and PBMC, both at the bulk and single-cell
levels. We find that loss of ACOD1 and itaconate synthesis led to increased inflammation during
infection and that exogenous itaconate, DI, and 4OI substantially reduced inflammation, most
notably IFN responses, without enhancing viral replication. 4OI was unique in that it also exerted
a direct anti-viral effect.

Results

Acod1 **expression and itaconate synthesis are induced in mouse lung during IAV-infection**

and limit pulmonary inflammation and disease severity. Compared with C57BL/6J, DBA/2J

mice are more susceptible to IAV infection in that weight loss is more rapid, mortality is higher,

and inflammation is more pronounced (Preusse 2017, Srivastava 2009). We have shown

previously that *Acod1* transcription in the first 48 h of infection is higher in IAV-infected lung

from DBA/2J than from C57BL/6J mice (Preusse 2013). A reanalysis of our RNAseq data from

a longer time course (Wilk 2015) showed that this difference persisted at least through day 3 post

infection (p.i.) and that *Acod1* expression nearly normalized by day 14 in the surviving strain

(**Fig. 1A**). This analysis also revealed a remarkable correlation with expression of *Tnfaip3*

(encoding an anti-inflammatory deubiquitinase) and a weaker positive correlation with *Hmox1*

and *2* (encoding inducible anti-oxidant enzymes)*,* all of which are known to be co-regulated with

Acod1 (Li, Zhang et al., 2013, Uddin, Joe et al., 2016). *Acod1* was the 4[th] most highly

upregulated pulmonary mRNA 48 h p.i. in C57BL/6J mice, where it appeared in a cluster of

transcripts pertaining to IFN responses and innate immunity (**Fig. 1B**). These results underscored

the close association of *Acod1* expression with systemic inflammation and oxidative stress in

IAV-infected mice. We therefore tested whether targeted deletion of *Acod1* would affect

inflammatory responses and host susceptibility to IAV infection. Itaconate was not detectable in

lungs from IAV-infected $Acod1^{-/-}$ mice; it was highly elevated on days 7/8 p.i. in the $Acod1^{+/+}$

and, significantly less, in $Acod1^{+/-}$ mice, and returned to normal by day 14 (**Fig. 1C**). Both

weight loss and mortality were higher in $Acod1^{-/-}$ than $Acod^{+/+}$ female C57BL/6J mice (**Fig.**

1D,E), with greatest changes occurring around day 8, the time of maximal itaconate

accumulation. Histologically assessed pulmonary inflammation was greatest in $Acod1^{-/-}$ mice and

intermediate in $Acod1^{+/-}$ mice (**Fig. 1F**). Thus, itaconate synthesis correlated with disease

activity, and loss of *Acod1* was associated with more severe disease and higher pulmonary

inflammation.

IAV infection induces *ACOD1* **expression in human PBMC and macrophages.** Unless stated

otherwise, all infections human cells and tissue were done using the strain PR8M

(A/PuertoRico/8/34 H1N1). We first tested whether IAV infection increased *ACOD1* mRNA

levels in human PBMC and M1/M2 macrophages. Phorbol myristate acetate (PMA)-

differentiated THP-1 (dTHP-1) cells were analyzed for comparison. Viral transcription, as

measured by hemagglutinin (*HA*) mRNA levels, was highest in M2 cells (**Fig. 2A**). The strongest

ACOD1 induction was seen in M2 cells and undifferentiated PBMC, and the weakest in M1 cells

and dTHP-1 cells (**Fig. 2B-E**), even though the latter two cell types did support a significant

degree of viral transcription (**Fig. 2A,D**). As in IAV-infected mouse lung (**Fig. 1**), *TNFAIP3* and

ACOD1 transcriptional correlated with each other (**Fig. 2C**). Infection of dTHP-1 cells with three

IAV (H1N1) strains of differential replication efficiency (Petersen, Mostafa et al., 2018b)

showed the strongest *ACOD1* and *TNFAIP3* induction by the two strains with the highest and the

weakest by the strain with the lowest replication efficiency (**Fig. 2D,E**). Thus, the extent of

ACOD1 induction after IAV infection differed significantly among the cell types tested, was

strongest in M2 primary cells, was accompanied in all cases by an increase in *TNFAIP3* mRNA,

and correlated with viral replication.

Itaconate and DI markedly reduce IFN responses in dTHP-1 cells without affecting viral

RNA replication. We then tested whether these compounds could modulate cellular

inflammation due to IAV infection. In infected dTHP-1 cells, treatments with itaconate or DI did

not affect viral hemagglutinin (*HA)* mRNA expression, but both reduced induction of *CXCL10*, a

key pro-inflammatory chemokine during influenza virus infection, with DI being several-fold

more potent (**Fig. 3B,C, Fig. S1**). The 12 h time point was chosen because our previous work

had identified it as the peak of IFN-I responses at the mRNA level in IAV-infected dTHP-1 cells

(Petersen, Mostafa et al., 2018a). We then assessed global effects of itaconate and DI on cellular

gene mRNA expression. A comparison of differentially expressed genes (DEGs) suggested that

the two compounds had both shared and unique effects on gene expression (**Fig. 3D**). Both

compounds had considerable effects on gene expression in uninfected cells (**Fig. 3E,F**). As

expected, IAV infection of untreated cells caused a major induction of IFN-regulated transcripts

by 12 h p.i. (see mRNAs labeled in the volcano plot in **Fig. 3G**). Strikingly, addition of both

itaconate and DI led to global down-regulation of IFN-regulated genes. This was accompanied

by upregulation of a substantial number of other transcripts, indicating that the compounds also

modulated other cellular processes (**Fig. 3H,I**). A principle component analysis (PCA) indicated

that effects of the compounds on the cells were so pronounced at these concentrations that

treated infected and treated uninfected cells clustered closely together, but far away from

untreated infected or uninfected cells (**Fig. S2**). Indeed, a hierarchical clustering analysis

revealed an across-the-board normalization of a clade of IFN- and inflammation-related

transcripts by the compounds, but also clusters of DEGs that were upregulated by either or both

compounds with respect to both uninfected and infected cells, further indicating effects upon cell

homeostasis in general (**EV1A,B**). mRNA and protein expression may be differentially affected

by IAV infection due to virus-induced "host-cell shut-off". We therefore assessed release of

inflammation-related polypeptides in supernatants from the same experiment (**Fig. 3J**). This

analysis confirmed the downregulation of pro-inflammatory factors (e.g., IP-10, MCP-1) by

itaconate and DI, but also showed increased IL-8 (DI only) and minor increases of IL1B (both IA

and DI) in uninfected cells. Of note, both compounds also increased levels of the anti-

inflammatory IL1 receptor antagonist (IL1RA) in supernatants from both infected and uninfected

cells. A comprehensive view of the protein targets that were detected by this assay revealed that

the two compounds effected substantially different changes in the target cells, as several factors

were induced even in uninfected cells specifically by DI or itaconate (**EV2**). Thus, itaconate and

DI apparently act on macrophages to release different recruitment programs that modify local

inflammatory cell populations.

Analysis of enriched GO terms revealed that IFN-I signaling was strongly induced in the infected

cells, which was prevented by treatment with itaconate and DI (**Fig. 4**). This analysis also

revealed that both compounds activated metabolic processes, whereas itaconate also modulated

processes relating to differentiation and membrane signaling. A KEGG pathway analysis

additionally revealed induction of classic pro-inflammatory pathways such as *TNF signaling*,

TLR receptor signaling, and *chemokine signaling* by infection, all of which were depleted by

itaconate and DI treatment (**Fig. S3**). However, in a separate experiment, there were only

tendencies of IA and DI to reduce nuclear localization of NF-kB during infection, whereas

reduction by 4OI was significant. (**EV1C**). Taken together, these results revealed strong anti-IFN

and anti-inflammatory effects of itaconate and DI, but also suggested that they exert other

important effects on cell metabolism and differentiation, some of which differ between the

compounds.

Anti-inflammatory effects of ectopically expressed itaconate, and exogenous itaconate and

DI on IAV-infected respiratory epithelial cells. We then tested whether the effects of itaconate

and DI could also be seen in cells not endogenously expressing *ACOD1*. For this purpose, we

studied the respiratory epithelial cell line A549, which supports productive IAV infection and

does not express *ACOD1*. As expected, infection of A549 cells did not result in induction of

ACOD1 mRNA, but transient transfection with an *ACOD1*-expressing plasmid led to significant

ACOD1 mRNA expression (**Fig. 5A**), which was accompanied by appearance of substantial

concentrations of IA (**Fig. 5B**). While there was no effect on viral *HA* mRNA expression,

transfection with the *ACOD1*-expressing vector led to a marked reduction in infection-associated

CXCL10 induction (**Fig. 5C,D**). A preliminary treatment experiment showed that (as opposed to

THP-1 cells) DI was cytotoxic at 1.0 mM in A549 cells, and the 0.5 mM concentration was

therefore used. Treatment with DI resulted in a modest reduction of *HA* expression early in

infection and a marked reduction of *CXCL10*, *IL6*, and *IFNB* expression (**Fig. 5E-H**). Exogenous

itaconate, too, reduced *CXCL10* mRNA, but less efficiently (**Fig. S4**). Taken together, these

results obtained with A549 cells suggest that immunomodulatory effects of itaconate and DI are

independent of the ability of the cell type to express endogenous *ACOD1*, and that similar effects

on IFN-directed gene expression can be obtained with ectopic, endogenous synthesis of itaconate

and exogenously added itaconate and DI.

Next, we assessed effects of itaconate and DI on cellular transcriptomes in IAV-infected A549

cells. PCA revealed the expected normalization of gene expression due to DI treatment, but also

increasing effects of the compounds on overall cell homeostasis (**Fig. S5**). As in THP-1 cells,

hierarchical clustering analysis revealed clear clustering of each of the four groups and the close

relationship of the uninfected and DI-treated infected cells (**EV3**). However, the anti-

inflammatory effect of itaconate was weaker than that of DI, and at the higher (40 mM)

concentration, the itaconate signature became so dominant that the itaconate-treated infected

group now formed its own clade (**Fig. S6**). IAV infection enriched GO terms related to IFN-I and

–II responses as well as other antiviral and pro-inflammatory terms, which could all be reduced

to near baseline by treatment with DI or itaconate (**Fig. 6**). Visualization of differentially

expressed IFN-related mRNA also showed that, in spite of its overall lower effect on pro-

inflammatory signaling than DI, itaconate treatment at the lower (20 mM) concentration did have

a considerable impact on IFN-I expression (**Fig 6B**). In addition, itaconate uniquely inhibited

NFKB signaling and *fatty acid metabolism*. KEGG-pathway analysis additionally revealed

depletion of other pro-inflammatory pathways, including *TNF signaling* and *necroptosis,* by the

treatments (**Fig. S7**).

Anti-inflammatory effect of itaconate and DI in a human lung tissue explant (HLTE) model

of IAV infection. To test whether the *ACOD1*/itaconate axis and the anti-inflammatory effects of

itaconate and DI are functional in the end-organ of IAV infection in humans, we then used a

human lung tissue culture model. Indeed, infection with IAV led to brisk transcription of viral

HA and host *CXCL10* mRNA in a 72 h time course (**Fig. S8**), whereas *ACOD1* mRNA levels did

not change significantly. Adding itaconate and DI reduced expression of *CXCL10* mRNA in

tissue and IP10 protein in culture supernatant, and itaconate additionally reduced ISG15 mRNA

in tissue, whereas viral titers in the supernatant were unaffected (**Fig. 7**). Re-analysis of a

published dataset from a model using a different IAV strain (H3N2) and histologically normal

lung tissue (Matos 2019) revealed a pronounced induction of *ACOD1* and *TNFAIP3* expression,

but (as opposed to *in vivo* infection of mouse lung) not induction of *HMOX1* or *2* (**Fig. S9).**

Anti-inflammatory effects of itaconate, DI, and 4OI on IAV-infected PBMC. PBMC play

important roles in influenza infection as they contain cells of innate and adaptive immunity, and

because circulating monocytes (which can sustain a nonproductive influenza virus infection) are

considered an important source of cytokines that support systemic spread of inflammation

(Duan, Hibbs et al., 2016). We therefore investigated effects of the compounds on IAV infected

PBMCs both at the bulk and single cell levels. Analysis with targeted PCR- and immuno-assays

demonstrated vigorous transcription of viral RNA, induction of *ACOD1* mRNA, and a brisk pro-

inflammatory response both at the mRNA and protein level (**Fig. 8A,B**). The induction of

ACOD1 mRNA expression in peripheral blood in IAV infection was also corroborated by

reanalysis of a published dataset from whole blood from patients with moderate and severe

influenza (**Fig. S10**). The effects of itaconate were mixed in that there was no significant change

in *ACOD1*, *IFNB1*, and *TNF* mRNA expression in cells or of IL1B protein in supernatants. In

contrast, DI and 4OI led to a marked reduction of IFN and pro-inflammatory cytokine expression

at the mRNA and protein level, as well as of *ACOD1* mRNA. As exemplified by the effect of DI

and 4OI on *TNF* and *ACOD1* expression, the treatments reduced expression of some mRNA

targets even in uninfected cells, suggesting that they also reduced some baseline inflammation.

Of note, 4OI in addition led to a pronounced (95%) reduction in *HA* transcription (p=<0.0001),

indicating inhibition of viral RNA replication. PCA of bulk PBMC transcriptomes revealed

strong effects of the compounds on cell transcriptomes in general, which led to one outlying DI-

treated sample and two itaconate-treated samples, whereas all four 4OI-treated samples formed a

clearly discernible group (**Fig. S11**). The most significant transcriptome changes were driven by

4OI, which had major effects on many genes that were not regulated by IAV infection or

itaconate or DI treatments (**EV4**). Nonetheless, induction of *HMOX1* by DI in IAV-infected

samples was evident. Functional enrichment analysis of GO terms based on the transcriptomes

revealed that infection induced the expected antiviral responses, including *response to IFN-I*, as

well as several other inflammation-related terms (**Fig. 8C**). At least one of the three compounds

prevented activation of each of the infection-associated terms. However, consistent with its lack

of effect on *IFNB* expression (**Fig. 8**), itaconate did not affect *response to IFN-I* or *influenza A*.

Several terms that were depleted by the compounds in infected PBMC were not enriched by

infection alone, suggesting that the compounds diminished also a baseline activation of the

PBMC. Interestingly, 4OI treatment strongly enriched terms relating to chromatin conformation,

which turned out to be driven by upregulation of histone deacetylase epression. A KEGG

pathway analysis essentially confirmed the findings of the GO term analysis (**Fig. S12**),

including that 4OI enriched pathways relating to chromatin structure such as *Alcoholism*. Taken

together, these results consolidated our above findings that itaconate and derivatives can

modulate inflammation due to IAV infection, but also suggested that DI and 4OI may have more

robust anti-IFN and anti-inflammatory effects than itaconate. Considering the variable anti-

inflammatory effects of itaconate and the miscellaneous effects of 4OI on gene expression in this

model, we subsequently focused on DI to decipher effects at the single-cell level.

Single-cell RNAseq of PBMC identifies monocytes as the main PBMC host cell of IAV

infection, sole source of *ACOD1* expression, and predominant target of immunosuppressive

effects. Thirteen cell types could be identified in the PBMC scRNAseq data set, but we focused

the analyses on the overarching five cell types: CD4+ and CD8+ T cells, B cells, NK cells, and

monocytes (**Fig. 9A,B**). Monocytes constituted the least numerous cell type, and DI treatment of

control and infected cells led to a further reduction (**Fig. S13**). It was not possible to discern
whether this was due to decreased survival or a technical artefact. Monocytes were essentially
the only cell type expressing viral RNA, as only negligible amounts of viral transcripts were
detected in lymphocytes and NK cells. DI treatment appeared to result in a slight increase in viral
RNA expression in monocytes (**Fig. 9C**). IAV infection upregulated expression of *ACOD1*
exclusively, and *CXCL10* predominantly, in monocytes, whereas *TNFAIP3*, *IFIT1,* and *ISG15*
were upregulated also in the NK, T and B cell compartments (**Fig. 9D**). DI treatment markedly
reduced expression of *ACOD1*, *IFIT1,* and *CXCL10*, but –interestingly– it reduced *TNFAIP3* and
ISG15 (albeit weakly) expression only in monocytes. Consistent with the observation that they
constituted the main PBMC host cell, IAV infection triggered the most vigorous transcriptomic
host response in monocytes, which was driven by the expected IFN responses (**Fig. 9E**).
However, discernable differential expression (mostly of IFN-related RNAs) was also observed in
the other cell types. CD8+ T cells are shown as a representative example in **Fig. 9F**, the other
cell types in **Fig. S14**. Monocytes were the only cell type that mounted a transcriptomic response
to DI treatment of uninfected cells. In infection, treatment with DI led to a global downregulation
of IFN-related transcripts in monocytes and also downregulation of a subpopulation of IFN-
related transcripts in lymphocytes and NK cells. When considering all DEGs irrespective of
magnitude of differential expression, DI exerted comparable effects on monocytes and T cells,
and only somewhat less on B cells. However, when considering only transcripts with FC >|2|, a
preferential effect on monocytes became apparent (**Fig. 9G,H**).

GO term enrichment analysis of the monocyte scRNAseq data confirmed the major enrichment
of IFN-related processes by infection and their depletion by DI treatment, which had been seen
in all models in this study. In addition, infection broadly dampened RNA metabolism, ribosome
assembly, and protein synthesis and export, which was consistent with the cytostatic effects of
type I IFN. DI treatment reversed all these effects and also stimulated some in uninfected cells
(**Fig. 10A**). A KEGG enrichment analysis of these monocyte data confirmed the depletion of
IFN-related pathways (*influenza, measles, cytosolic DNA sensing pathway*) by DI treatment, but
also of other pro-inflammatory pathways (*TLR signaling, cytokine-cytokine receptor interaction,*
NF-kB signaling) in infected and uninfected cells (**EV5).** Of note, DI treatment apparently
stimulated ribosome function in both infected and control monocytes.

311 The Venn diagrams in **Fig. 10B** illustrate that monocytes are the cell type in which the most

312 biological processes were activated by infection and depleted by DI treatment, but reduction of

313 antiviral and pro-inflammatory responses was a common theme in the other cell types as well

314 (**Fig. 10C**). Indeed, GO term analysis identified four major IFN-related processes as a central

315 functional network that was upregulated by infection and downregulated by DI treatment during

316 infection in all five cell types (**Fig. 10B**). A marked stimulation of protein synthesis and export

317 by DI was also apparent in the uninfected lymphocytes and NK cells. Finally, in order to identify

318 a common transcriptomic network we performed GO and KEGG pathway enrichment analyses

319 of the 81 transcripts that were commonly differentially expressed in IAV infection and regulated

320 in the opposite direction by DI treatment in all five cell types (center of the Venn diagram in **Fig.**

321 **9G**). While the GO analysis by and large confirmed the findings shown in **Fig. 10B,** the KEGG

322 pathway analysis revealed a major effect of DI treatment on *allograft rejection*, *graft-vs.-host*

323 *disease*, and *antigen processing and presentation,* suggesting a broader interference with

324 immune cell activation by DI (**Fig. S15**).

Discussion

In this first dedicated study on the impact of itaconate and derivatives on host responses to IAV

In this first dedicated study on the impact of itaconate and derivatives on host responses to IAV

infection, we found that (i) the endogenous ACOD1/itaconate axis limits infection-induced

pulmonary inflammation in a murine model, (ii) ectopically expressed ACOD1 possesses anti-

IFN activity in an epithelial cell line that normally does not express this enzyme, and (iii) that

exogenously applied itaconate and/or derivatives restrict IFN-responses and modulate classic

pro-inflammatory cytokines across all models tested.

Anti-inflammatory effects of itaconate and derivatives. A major suppression of IFN and some

pro-inflammatory cytokine responses was the recurrent theme in all models, i.e. immortalized

cells, primary cells, and human lung tissue. This is fully consistent with the early reports that DI

has strong anti-inflammatory properties (Lampropoulou 2016), which was subsequently

corroborated in diverse models of inflammation and infection, also using 4OI (e.g., (Gu 2020,

Xu, Jiang 2020a, Zhao 2019). We used itaconate and DI in parallel in several models, which

enabled comparing their relative effectiveness. Overall, results with DI were most consistent, and

in spite of its higher toxicity on A549 cells than on dTHP-1 cells, the lower dose still possessed

strong anti-inflammatory effects. Itaconate effects were less consistent and much higher doses

had to be used to achieve anti-IFN effects similar to DI. Itaconate did not reduce expression of

IFNB1, *TNF* mRNA or IL1B protein in PBMC. The lack of *TFN* repression by itaconate in that

model is consistent with previous reports that it usually does not affect TNF-α levels

(Bambouskova et al., 2018). Recent work has also revealed the potential of itaconate to actually

enhance IL1B release (Muri 2020). Of note, at higher doses, itaconate and strong electrophilic

compounds can actually increase inflammation due to CASP-8 dependent inflammasome

activation, and this effect is magnified when the cells are not pre-incubated with the compound

before adding the pro-inflammatory stimulus (Muri 2020), as we did in PBMC infections.

However, these authors did not investigate effects on IFN suppression. 4OI demonstrated the

expected potent anti-inflammatory effects, but also broad effects on processes relating to, e.g.,

chromatin structure, the potential effects of which on cell homeostasis and cell-pathogen

interactions will require further study.

Anti-inflammatory effect of ectopic synthesis of itaconate in A549 cells. Despite the multitude of
studies on effects of exogenous administration of itaconate derivatives on cellular inflammation,
it had not been tested whether ectopic synthesis of itaconate in a cell type that does not naturally
express *ACOD1* has similar effects. Our transfection study (Fig. 5) demonstrated that ectopic,
endogenously synthesized itaconate exerts very similar anti-IFN effects as exogenously added
itaconate or DI. Of note, the achieved itaconate concentration is very close to that measured in
LPS/IFNγ stimulated dTHP-1 cells (Chen et al, unpublished data), suggesting that it is
physiological.

Cell-type specific and -independent anti-inflammatory effects of IA and DI. A particular strength
of our study lies in the use of scRNAseq to identify major target cells in peripheral blood. Firstly,
this analysis identified monocytes as the major PBMC type that is infected by IAV. Secondly,
we found that on one hand DI can attenuate IFN-responses in lymphocytes and NK cell as well,
but that the immunosuppressive effect is greatest on monocytes. Taken together, these results
suggest that even though pharmacological use of itaconate and derivatives would modulate host
responses in a diverse array of cells, monocytes/macrophages likely constitute a major functional
hub in the genesis of inflammation in IAV infection and subsequent treatment responses to
itaconate-based adjunct treatments.

Antiviral effect of 4OI. Olagnier et al. (2020) recently reported that Nrf2 activators including 4OI
and dimethyl fumarate limit inflammation in cellular models of SARS-CoV-2 infection in an
NRF2-dependent manner. Of note, as opposed to influenza viruses, infection with SARS-CoV-2
suppresses the NRF2 pathway, which can be reversed in part by treatment with Nrf2 activators.
These authors in addition observed an antiviral effect of 4OI, which proved to be independent of
NRF2- and type I IFN-signaling. Our finding of a remarkable suppression of IAV RNA
expression in PBMC corroborates the notion that 4OI has strong antiviral effects in addition to
its well-documented anti-inflammatory properties. Further research is needed to validate our
finding in other cell types, as we tested it in PBMC only. Nonetheless, the gene expression
analysis of itaconate, DI- and 4OI-infected PBMC identified several processes/pathways
uniquely regulated by 4OI. Further research should focus on assessing whether any of them
could contribute to the antiviral effect of 4OI.

Infection-independent effects on target cells. By using relatively high doses of itaconate and DI

and assessing effects of the compounds on uninfected cells or their effects on processes that were

not changed during infection, our study also provides a first comprehensive look at global effects

of the compounds on specific target cells. In particular, itaconate and 4OI exerted pronounced

effects on gene expression in A549 cells and PBMCs, respectively. Clearly, it will be of great

interest to investigate how these effects relate to differences in toxicities on different cell types,

or to differences in their effects upon virus-host interactions.

Importance of endogenous ACOD1 expression. In the mouse model, we observed a more severe

clinical phenotype and higher inflammation in *Acod1*$^{-/-}$ mice. This agrees with work in a mouse

model of *M. tuberculosis* infection, where inflammation and mortality were higher in the *Acod1*$^{-/-}$

animals (Nair 2018). However, IAV infection was also assessed in that study, and deleting the

Acod1 gene did not affect weight loss or survival. Those authors used equal numbers of male and

female mice, whereas we used female mice only. Further research is required into possible sex-

specific effects of the *Acod1* gene in influenza infection. Of note, we found upregulation of

ACOD1 by IAV infection and its prevention by DI treatment in myeloid cells, i.e. dTHP-1 and

peripheral blood monocytes, and upregulation of *ACOD1* mRNA in blood from patients with

moderate and severe influenza infection as compared with healthy donors. Thus, induction of

ACOD1 expression (and thereby itaconate synthesis) likely is a common feature in human

influenza. We have recently identified the active center of ACOD1 enzyme and found that

naturally occurring loss-of-function mutations are extremely rare, suggesting that physiologic

itaconate responses have been important in human evolution (Chen 2019). It is therefore

tempting to speculate that one beneficial role of an active ACOD1/itaconate axis is to reduce the

risk of death or organ damage from infection-associated hyper-inflammation and that loss-of-

function genotypes have been selected against during times of high infectious disease burdens.

Clinical implications. The current COVID-19 pandemic has, once again, focused the research

limelight on the deleterious effects of overshooting innate immune responses in acute viral

infections. In a current report, Lee et al. found that intense IFN responses in PBMC from patients

are a hallmark of both severe COVID-19 and influenza (Lee 2020). Of note, a recent systematic

review of 45 studies on corticosteroid use for COVID-19 found a significant beneficial effect

(van Paassen 2020), whereas to date there are no clinically effective antivirals against SARS-
CoV-2. Therefore, the concept of adjunct treatments to control deleterious host immune
responses in viral infections is more pertinent than ever. Considering (i) the well documented
cytoprotective properties of itaconate and derivatives and (ii) our findings of remarkable anti-
inflammatory and anti-IFN effects in the absence of increased viral replication, itaconate
derivatives would constitute a highly promising class of compounds for such applications.
Considering the more reliable results obtained with DI as compared to unmodified itaconate, the
lower required concentrations of DI and 4OI, but also the striking anti-viral effects of 4OI, it
appears that chemical variants of itaconate, rather than its native form, will take the lead in
further translational development to the bedside.

Detailed methods are found in the Appendix

Acknowledgements. We thank V. Kaever, H. Bähre and staff of the RCU Metabolomics at
Hannover Medical School for support with the itaconate measurements, Elena Reinhard for
assistance with the animal experiments, the HZI Animal Facility for expert animal care, the HZI
Genome Analytics Platform for microarray and RNA sequencing, L. Gröbe for cell sorting, D.
Jonigk and P. Braubach (Institute of Pathology, Hannover Medical School) for providing human
lung tissue. We thank Hans Jörg Hauser (HZI) for helpful comments on the manuscript.

Conflict of interest. The authors declare that none of them have a conflict of interest relating to
conduct of the study or publication of the manuscript.

Data Availability

The gene expression data are available through GEO at accession number GSE162210,
GSE162260, GSE162261 and GSE164922. These data are currently private and access can be
given to reviewers. The data previously published in (Wilk et al., 2015) are available at
GSE66040.

The Paper Explained

PROBLEM. Interferon responses are part of the primary host defenses against infections. However, excessive inflammation is often a major factor in severe disease or even death due to respiratory infections such as influenza, as it can lead to acute respiratory distress syndrome and sepsis-like multiorgan involvement.

RESULTS. We applied itaconate and derivatives to influenza A virus infected human cells and lung tissue and found that these compounds markedly repress interferon responses and some pro-inflammatory processes without increasing viral replication. In fact, 4-octyl itaconate greatly decreased viral RNA replication in peripheral blood mononuclear cells. Single cell RNA sequencing identified monocytes as the predominant blood cell infected with the virus and also the major target of itaconate activity.

IMPACT. The results suggest that novel medications based on itaconate may be effective in dampening deleterious inflammation in influenza infection without the risk of favoring viral spread in the patient.

Figure legends

Figure 1. Reciprocal interactions between *Acod1* expression and inflammation during influenza A virus infection in mice. A. Higher pulmonary expression of *Acod1, Tnfaip3,* and *Hmox1* and *2* mRNA in DBA/2J (D2) than in C57BL/6J (B6) mice following IAV infection (reanalysis of published RNAseq data (Wilk et al., 2015)) (n=3 each strain and time point). *p<0.05; **p<0.01; ***p<0.001 (pairwise t-test with Bonferroni-Holm multiple comparisons correction). **B.** *Acod1* mRNA is among the most highly upregulated transcripts in mouse lung within 48 h of IAV infection (RNAseq, n=8 per time point). **C.** Lower pulmonary levels of itaconate in IAV-infected *Acod1$^{+/-}$* than in *Acod1$^{+/+}$* mice. Itaconate was not detected in the *Acod1$^{-/-}$* mice. (n=36 *Acod1$^{+/+}$*, 27 *Acod1$^{+/-}$*, 4 *Acod1$^{-/-}$* mice). *p<0.05; **p<0.01; ***p<0.001 (unpaired t-test). **D.** Greater weight loss of female *Acod1$^{-/-}$* C57BL/6J mice during infection with IAV. Mice had to be euthanized when weight loss exceeded 20%; n=9 *Acod1$^{+/+}$*, 10 *Acod1$^{+/-}$*, 9 *Acod1$^{-/-}$* mice. *p<0.05; **p<0.01; ***p<0.001 (unpaired t-test) **E.** Survival data of the same experiment as in C. Median survival after day 7 was significantly lower in *Acod1$^{-/-}$* mice (p=0.018, Mann-Whitney-U test). **F.** Semi-quantitative histopathological scoring of lungs on days 8/9 (mid), 11 and 14 (late) reveals highest inflammation in *Acod1$^{-/-}$* mice; n=9 *Acod1$^{+/+}$*, 10 *Acod1$^{+/-}$*, 9 *Acod1$^{-/-}$* mice (all female). *p<0.05; **p<0.01; ***p<0.001 (two-way ANOVA, Tukey's multiple comparisons test) **G.** H&E stained lung sections showing inflammation severity scores of 0 (left) and 3 (right).

Figure 2. Differences among human myeloid cells in *ACOD1* mRNA induction during IAV infection. A. Induction of *ACOD1* and *TNFAIP3* mRNA (RT-qPCR) in M1 and M2 macrophages and PBMCs during IAV infection. **B.** Induction of *ACOD1* and *TNFAIP3* (RT-qPCR) in dTHP-1 cells correlates with virulence of three IAV strains. Mean ±SEM (n=4). *HA* = IAV hemagglutinin. *p<0.05; **p<0.01; ***p<0.001 (two-way ANOVA, Tukey's multiple comparisons test, with reference to M1 macrophages in A, and to Gi-WT in B).

Figure 3. Effects of itaconate and DI on cellular inflammation due to IAV infection of dTHP-1 cells. A. Outline of experiment. dTHP-1 cells were treated with itaconate or DI overnight, or left untreated, and then infected with IAV. Dose-response curves of effects of a range of itaconate and DI and concentrations on *HA* and *CXCL10* mRNA expression are shown

in **Fig. S1. B**-**C**. 48 h time course, DI = 0.25 mM (n=3). mRNA expression (RT-qPCR) of *HA*

(**B**) and *CXCL10* mRNA (**C**) *p<0.05; **p<0.01; ***p<0.001 (unpaired t-test). **D-I.** Global

transcriptomic changes (microarray analysis) due to itaconate (25 mM) and DI (1 mM) treatment

of uninfected and IAV-infected dTHP-1 cells (n=4). **D.** Venn diagram showing unique and

common differentially expressed genes by itaconate and DI treatment of infected cells. **E-I.**

Volcano plots showing differential mRNA expression in uninfected dTHP-1 cells under

itaconate and DI treatment (**E-F**) and infected dTHP-1 cells with or without itaconate or DI

treatment (**G-I**). **J.** Effects of itaconate (25 mM) and DI (1 mM) treatment on levels of

inflammation-related polypeptides in dTHP-1 cell supernatants 12 h after IAV infection or mock

infection. Mean ± SEM (n = 3). *p<0.05; **p<0.01; ***p<0.001 (One-way ANOVA, Tukey's

multiple comparison test).

Figure 4. Itaconate and DI treatment leads to a global downregulation of IFN-I signaling in

IAV-infected dTHP-1 cells. Enrichment analysis of gene ontology (GO Biological Process)

terms based on the microarray data shown in Fig. 3, using DEGs (p<0.05, FC >|1.5|) as input.

GO terms (FDR ≤0.05) reveal major induction of antiviral and inflammatory responses by

infection, which are decreased by both itaconate and DI treatment.

Figure 5. Effect of overexpression of ACOD1 and exogenous treatment with DI on IAV-

infected A549 cells. A. A549 cells were transfected with pCMV6-Hu-ACOD1, pCMV6 empty

vector, or mock transfected for 48 h and then infected with IAV for 48 h (n=3). Expression of the

indicated mRNAs was measured by RT-qPCR, itaconate concentrations by LC-MS/MS.

*p<0.05; **p<0.01; ***p<0.001 (2-way ANOVA, Tukey's multiple comparisons test) **B.** A549

cells were treated with 0.25 mM DI overnight, infected with IAV for 48 h (n=3), and expression

of the indicated mRNAs measured by RT-qPCR. Mean ± SEM (n = 3). *p<0.05; **p<0.01;

***p<0.001 (2-way ANOVA, Sidak's multiple comparisons test).

Figure 6. GO term enrichment analysis of itaconate and DI reveals global effects on IFN-

signaling and cellular inflammation in A549. A549 cells were, in the case of treatment,

incubated overnight with DI (0.5 mM) or itaconate (20 mM), and then infected with strain

PR8M. Gene expression was determined by microarray analysis 24 h p.i. (n=3). **A**. GO

enrichment (FDR <0.05) analysis using DEGs (p<0.05, FC >|1.5|) as input. **B**. Hierarchical

cluster analysis illustrating itaconate treatment downregulating a subset of genes involved in

IFNA/B signaling.

Fig. 7. Anti-inflammatory effect of itaconate and DI on IAV-infected human lung tissue

explants. Human primary lung tissue from patients with emphysema and pulmonary arterial

hypertension was incubated overnight with itaconate (25 mM), DI (1 mM), or medium only and

was then infected with IAV for 24 h (5 donors, 3 tissue pieces per donor). **A**. Viral titers in

supernatant (focus forming assay). **B-D**. *CXCL10* mRNA (**B**), and ISG15 (**C**) mRNA in tissue

(RT-qPCR); IP10 concentration in supernatant (EIA) (**D**). Mean ± SEM (n=6). *p<0.05;

**p<0.01; ***p<0.001 (Mann-Whitney U test).

Fig. 8. Anti-inflammatory effects of DI and 4OI on IAV-infected human PBMCs. PBMCs

were isolated from freshly donated human blood and then infected with IAV for 12 h in the

presence or absence of itaconate (10 mM), DI (0.5 mM), or 4OI (25 μM) without preincubation

(7 donors, 3 replicates). **A.** Expression of the indicated mRNAs in PBMC (RT-qPCR). **B.**

Concentrations of the indicated proteins in PBMC supernatants (ELISA). Mean +/- SEM (6

donors, 3 replicates per donor). *p<0.05; ***p<0.01; ***p<0.001 (1-way ANOVA). **C.** GO

enrichment analysis of effects of IA, DI, and 4OI on IAV-infected PBMC. Microarray analysis

of PBMC RNA from four of the donors featured in **A-B**. RNA was pooled from the three

replicates of each donor, resulting in 4 samples per group. GO term analysis was performed on

DEGs (p<0.05, FC>|1.5|) and terms with an FDR <0.02 in at least one condition are shown.

Fig. 9. Single-cell transcriptomic responses of PBMC to IAV infection and DI treatment.

PBMCs were isolated from freshly donated human blood (one donor), and then infected with

IAV for 12 h with or without DI (0.5 mM) in the medium. **A-B**. UMAP plots identifying 13 cell

types (**A**) that make up the five major cell types (**B**) studied. The RNA cell markers used are

listed in Table S1. **C-D.** Differential mRNA expression in uninfected, infected and infected/DI-

treated PBMCs. **E.** Differential mRNA expression in monocytes in the indicated paired

comparisons. Major transcriptional reprogramming due to IAV infection is evident, which is

largely prevented by treatment with DI. **F.** Differential mRNA expression in CD8+ T cells in the

indicated paired comparisons. Reprogramming in the presence of IAV is much less, but a DI

treatment effect is evident. **G-H.** Venn diagrams based on RNAs differentially regulated (FDR

<0.05) by both viral infection and DI treatment in each of the 5 major cell type, either

irrespective of FC (**G**) or FC >|2| (**H**).

Figure 10. GO enrichment analysis at the single cell level of DI effects on uninfected and

IAV-infected PBMC. Analysis based on the scRNAseq data shown in Fig. 9. Go terms that are

enriched/depleted in all cell types by infection and DI treatment are marked with a red asterix. **A.**

GO enrichment analysis of monocytes. **B.** Venn diagrams illustrating GO terms that are

commonly or uniquely regulated (FDR <0.05) in monocytes, CD4 T cells, CD8 T cells, B cells,

and NK cells; due to infection alone, DI treatment of infected cells, or DI treatment of uninfected

cells. **C**. Go term analysis of CD4 T cells, CD8 T cells, B cells, and NK cells.

EV1 A. Hierarchical clustering analysis of effects of itaconate and DI on IAV-infected

dTHP-1 cells. Analysis based on the same microarray data as used for Fig. 3 and EV1B, using

the top 100 DEGs (FDR F-test <1.89E-09). Uninfected cells and untreated IAV-infected cells

form one clade and itaconate and DI treated IAV-infected cells the other clade. The brackets

identify (i) a large pro-inflammatory subclade in IAV infection, which is globally downregulated

by the treatments; (ii) a large cluster of DEGs specific for itaconate treatment; (iii) a small cluster

of DEGs specific for DI treatment, and (iv) a subclade comprising DEGs differentially expressed

due to both compounds. **B. Enrichment plots based on the 99 genes contained in GO term**

Response to type I interferon. Enrichment scores are plotted on the y-axis, the mRNAs

(identified by vertical lines) along the x-axis, ranked by fold change. The plots illustrate

pronounced enrichment in a large number of genes due to infection, which is nearly

quantitatively depleted by treatment with both itaconate and DI. **C. 4OI prevents increased**

NFκB nuclear localization in IAV infection. IAV infection of dTHP-1 cells, with or without

treatment with IA (25 mM), DI (0.5 mM) and 4OI (25 μM). Data are based on fluorescent

intensities resulting from colocalization of NFκB (red) in nuclei (DAPI, blue) and represent the

mean of at least 5 representative images per replicate sample (n=3). *p<0.05; **p<0.01;

***p<0.001 (unpaired t-test with Welch's correction).

EV2. Effects of itaconate (20 mM) and DI (0.5 mM) on transcriptomic responses in IAV-

infected A549 cells. Analysis based on the microarray data used for the GO analysis in Fig. 6.

Unsupervised hierarchical clustering analysis of the 100 most significant DEGs (FDR F-test

<1.46E-10). DI-treated infected and uninfected cells cluster in one clade, and infected and

itaconate-treated infected cells in the other. There is a large clade mostly containing

proinflammatory genes, which are downregulated by DI, whereas effects of itaconate are much

weaker. *DUSP95* is exclusively downregulated by itaconate (arrowhead). There are two smaller

clades of genes that form an "itaconate signature", comprising genes that are, albeit to a lesser

extent, also upregulated by IAV infection, or uniquely by itaconate.

EV3. Hierarchical clustering analysis of transcriptomic changes in IAV infected PBMC

and responses to itaconate, DI, and 4OI treatment. The 100 most significant DEGs were

selected (FDR F-test <3.72E-05). Transcriptome changes are mostly due to marked effects of

4OI (more down- than upregulation) on genes which are not affected by IAV infection,

indicating general effects on cell homeostasis. However, induction of *HMOX1* by DI in IAV

infection is evident (arrow). The arrow points to the extreme outlier under DI treatment seen in

the PCA.

EV4. Cell-type specific KEGG pathway analysis of effects of DI on infected and uninfected

monocytes, CD4 T cells, CD8 T cells, B cells, and NK cells. Analyses based on the scRNAseq

data used for Fig. 9 and 10.

References

Alberts R, Srivastava B, Wu H, Viegas N, Geffers R, Klawonn F, Novoselova N, do Valle TZ, Panthier JJ,

Schughart K (2010) Gene expression changes in the host response between resistant and susceptible

inbred mouse strains after influenza A infection. *Microbes Infect* 12: 309-18

Bambouskova M, Gorvel L, Lampropoulou V, Sergushichev A, Loginicheva E, Johnson K, Korenfeld D,

Mathyer ME, Kim H, Huang LH, Duncan D, Bregman H, Keskin A, Santeford A, Apte RS, Sehgal R, Johnson

B, Amarasinghe GK, Soares MP, Satoh T et al. (2018) Electrophilic properties of itaconate and derivatives

regulate the IkappaBzeta-ATF3 inflammatory axis. *Nature* 556: 501-504

Chen F, Lukat P, Iqbal AA, Saile K, Kaever V, van den Heuvel J, Blankenfeldt W, Bussow K, Pessler F (2019)

Crystal structure of cis-aconitate decarboxylase reveals the impact of naturally occurring human

mutations on itaconate synthesis. *Proc Natl Acad Sci U S A* 116: 20644-20654

Cho H, Proll SC, Szretter KJ, Katze MG, Gale Jr M, Diamond MS (2013) Differential innate immune

response programs in neuronal subtypes determine susceptibility to infection in the brain by positive-

stranded RNA viruses. *Nature medicine* 19: 458-464

Collaborators GI (2018) Mortality, morbidity, and hospitalisations due to influenza lower respiratory

tract infections, 2017: an analysis for the Global Burden of Disease Study 2017. *Lancet Resp Med* 7: 69-

89

Cordes T, Lucas A, Divakaruni AS, Murphy AN, Cabrales P, Metallo CM (2020) Itaconate modulates

tricarboxylic acid and redox metabolism to mitigate reperfusion injury. *Mol Metab* 32: 122-135

Cordes T, Michelucci A, Hiller K (2015) Itaconic Acid: The Surprising Role of an Industrial Compound as a

Mammalian Antimicrobial Metabolite. *Annu Rev Nutr* 35: 451-73

Cordes T, Wallace M, Michelucci A, Divakaruni AS, Sapcariu SC, Sousa C, Koseki H, Cabrales P, Murphy

AN, Hiller K, Metallo CM (2016) Immunoresponsive Gene 1 and Itaconate Inhibit Succinate

Dehydrogenase to Modulate Intracellular Succinate Levels. *J Biol Chem* 291: 14274-84

Daniels BP, Kofman SB, Smith JR, Norris GT, Snyder AG, Kolb JP, Gao X, Locasale JW, Martinez J, Gale M,

Jr., Loo YM, Oberst A (2019) The Nucleotide Sensor ZBP1 and Kinase RIPK3 Induce the Enzyme IRG1 to

Promote an Antiviral Metabolic State in Neurons. *Immunity* 50: 64-76 e4

Duan M, Hibbs ML, Chen W (2016) The contributions of lung macrophage and monocyte heterogeneity

to influenza pathogenesis. *Immunol Cell Biol* 95: 225-235

Eklund A (2016) beeswarm: The BeeSwarm Plot, an Alternative to Stripchart. *https://CRANR-

projectorg/package=beeswarm*

Gu L, Lin J, Wang Q, Li C, Peng X, Fan Y, Lu C, Lin H, Niu Y, Zhu G, Zhao G (2020) Dimethyl itaconate

protects against fungal keratitis by activating the Nrf2/HO-1 signaling pathway. *Immunol Cell Biol* 98:

229-241

Hooftman A, O'Neill LAJ (2019) The Immunomodulatory Potential of the Metabolite Itaconate. *Trends*

Immunol

Kuhn M (2017) PhD dissertation. Hannover Medical School. Hannover, Germany.

Lampropoulou V, Sergushichev A, Bambouskova M, Nair S, Vincent EE, Loginicheva E, Cervantes-

Barragan L, Ma X, Huang SC, Griss T, Weinheimer CJ, Khader S, Randolph GJ, Pearce EJ, Jones RG, Diwan

A, Diamond MS, Artyomov MN (2016) Itaconate Links Inhibition of Succinate Dehydrogenase with

Macrophage Metabolic Remodeling and Regulation of Inflammation. *Cell Metab* 24: 158-66

Lee JS, Park S, Jeong HW, Ahn JY, Choi SJ, Lee H, Choi B, Nam SK, Sa M, Kwon JS, Jeong SJ, Lee HK, Park

SH, Park SH, Choi JY, Kim SH, Jung I, Shin EC (2020) Immunophenotyping of COVID-19 and influenza

highlights the role of type I interferons in development of severe COVID-19. *Sci Immunol* 5

Li Y, Zhang P, Wang C, Han C, Meng J, Liu X, Xu S, Li N, Wang Q, Shi X, Cao X (2013) Immune responsive

gene 1 (IRG1) promotes endotoxin tolerance by increasing A20 expression in macrophages through

reactive oxygen species. *J Biol Chem* 288: 16225-34

Liu H, Feng Y, Xu M, Yang J, Wang Z, Di G (2018) Four-octyl itaconate activates Keap1-Nrf2 signaling to

protect neuronal cells from hydrogen peroxide. *Cell Commun Signal* 16: 81

Matos ADR, Wunderlich K, Schloer S, Schughart K, Geffers R, Seders M, Witt M, Christersson A,

Wiewrodt R, Wiebe K, Barth P, Hocke A, Hippenstiel S, Honzke K, Dittmer U, Sutter K, Rescher U,

Rodionycheva S, Matera N, Ludwig S et al. (2019) Antiviral potential of human IFN-alpha subtypes

against influenza A H3N2 infection in human lung explants reveals subtype-specific activities. *Emerg*

Microbes Infect 8: 1763-1776

Michelucci A, Cordes T, Ghelfi J, Pailot A, Reiling N, Goldmann O, Binz T, Wegner A, Tallam A, Rausell A

(2013) Immune-responsive gene 1 protein links metabolism to immunity by catalyzing itaconic acid

production. *Proceedings of the National Academy of Sciences* 110: 7820-7825

Mills EL, Ryan DG, Prag HA, Dikovskaya D, Menon D, Zaslona Z, Jedrychowski MP, Costa ASH, Higgins M,

Hams E, Szpyt J, Runtsch MC, King MS, McGouran JF, Fischer R, Kessler BM, McGettrick AF, Hughes MM,

Carroll RG, Booty LM et al. (2018) Itaconate is an anti-inflammatory metabolite that activates Nrf2 via

alkylation of KEAP1. *Nature* 556: 113-117

Muri J, Wolleb H, Broz P, Carreira EM, Kopf M (2020) Electrophilic Nrf2 activators and itaconate inhibit

inflammation at low dose and promote IL-1beta production and inflammatory apoptosis at high dose.

Redox Biol 36: 101647

Nair S, Huynh JP, Lampropoulou V, Loginicheva E, Esaulova E, Gounder AP, Boon ACM, Schwarzkopf EA,

Bradstreet TR, Edelson BT, Artyomov MN, Stallings CL, Diamond MS (2018) Irg1 expression in myeloid

cells prevents immunopathology during M. tuberculosis infection. *J Exp Med* 215: 1035-1045

O'Neill LAJ, Artyomov MN (2019) Itaconate: the poster child of metabolic reprogramming in macrophage

function. *Nat Rev Immunol*

Olagnier D, Farahani E, Thyrsted J, Blay-Cadanet J, Herengt A, Idorn M, Hait A, Hernaez B, Knudsen A,

Iversen MB, Schilling M, Jorgensen SE, Thomsen M, Reinert LS, Lappe M, Hoang HD, Gilchrist VH, Hansen

AL, Ottosen R, Nielsen CG et al. (2020) SARS-CoV2-mediated suppression of NRF2-signaling reveals

potent antiviral and anti-inflammatory activity of 4-octyl-itaconate and dimethyl fumarate. *Nat Commun*

11: 4938

Petersen H, Mostafa A, Tantawy MA, Iqbal AA, Hoffmann D, Tallam A, Selvakumar B, Pessler F, Beer M,

Rautenschlein S (2018a) NS Segment of a 1918 Influenza A Virus-Descendent Enhances Replication of

H1N1pdm09 and Virus-Induced Cellular Immune Response in Mammalian and Avian Systems. *Frontiers*

in Microbiology 9: 526

Petersen H, Mostafa A, Tantawy MA, Iqbal AA, Hoffmann D, Tallam A, Selvakumar B, Pessler F, Beer M,

Rautenschlein S, Pleschka S (2018b) NS Segment of a 1918 Influenza A Virus-Descendent Enhances

Replication of H1N1pdm09 and Virus-Induced Cellular Immune Response in Mammalian and Avian

Systems. *Front Microbiol* 9: 526

Preusse M, Schughart K, Pessler F (2017) Host Genetic Background Strongly Affects Pulmonary

microRNA Expression before and during Influenza A Virus Infection. *Front Immunol* 8: 246

Preusse M, Tantawy MA, Klawonn F, Schughart K, Pessler F (2013) Infection- and procedure-dependent

effects on pulmonary gene expression in the early phase of influenza A virus infection in mice. *BMC*

Microbiol 13: 293

R_Core_Team (2014) R: A language and environment for statistical computing. *R Foundation for*

Statistical Computing, Vienna, Austria URL http://www.R-project.org/

Ragab D, Salah Eldin H, Taeimah M, Khattab R, Salem R (2020) The COVID-19 Cytokine Storm; What We

Know So Far. *Front Immunol* 11: 1446

Ramos I, Fernandez-Sesma A (2015) Modulating the Innate Immune Response to Influenza A Virus:

Potential Therapeutic Use of Anti-Inflammatory Drugs. *Front Immunol* 6: 361

Srivastava B, Blazejewska P, Hessmann M, Bruder D, Geffers R, Mauel S, Gruber AD, Schughart K (2009)

Host genetic background strongly influences the response to influenza a virus infections. *PLOS One* 4:

e4857

Strelko CL, Lu W, Dufort FJ, Seyfried TN, Chiles TC, Rabinowitz JD, Roberts MF (2011) Itaconic acid is a

mammalian metabolite induced during macrophage activation. *J Am Chem Soc* 133: 16386-16389

Tang C, Wang X, Xie Y, Cai X, Yu N, Hu Y, Zheng Z (2018) 4-Octyl Itaconate Activates Nrf2 Signaling to

Inhibit Pro-Inflammatory Cytokine Production in Peripheral Blood Mononuclear Cells of Systemic Lupus

Erythematosus Patients. *Cell Physiol Biochem* 51: 979-990

Uddin MJ, Joe Y, Kim S-K, Jeong SO, Ryter SW, Pae H-O, Chung HT (2016) IRG1 induced by heme

oxygenase-1/carbon monoxide inhibits LPS-mediated sepsis and pro-inflammatory cytokine production.

Cellular and Molecular Immunology 13: 170

van Paassen J, Vos JS, Hoekstra EM, Neumann KMI, Boot PC, Arbous SM (2020) Corticosteroid use in

COVID-19 patients: a systematic review and meta-analysis on clinical outcomes. *Crit Care* 24: 696

Wilk E, Pandey AK, Leist SR, Hatesuer B, Preusse M, Pommerenke C, Wang J, Schughart K (2015) RNAseq

expression analysis of resistant and susceptible mice after influenza A virus infection identifies novel

genes associated with virus replication and important for host resistance to infection. *BMC Genomics* 16:

655

Xu M, Jiang P, Sun H, Yuan X, Gao S, Guo J, Zhao C, Hu X, Liu X, Fu Y (2020a) Dimethyl itaconate protects

against lipopolysaccharide-induced endometritis by inhibition of TLR4/NF-kappaB and activation of

Nrf2/HO-1 signaling pathway in mice. *Iran J Basic Med Sci* 23: 1239-1244

Xu M, Jiang P, Sun H, Yuan X, Gao S, Guo J, Zhao J, Hu X, Liu X, Fu Y (2020b) Dimethyl itaconate protects

against lipopolysaccharide-induced endometritis by inhibition of TLR4/NF-κB and activation of Nrf2/HO-

1 signaling pathway in mice. *Iran J Basic Med Sci* 23: 1239-1244

Zhao C, Jiang P, He Z, Yuan X, Guo J, Li Y, Hu X, Cao Y, Fu Y, Zhang N (2019) Dimethyl itaconate protects

against lippolysacchride-induced mastitis in mice by activating MAPKs and Nrf2 and inhibiting NF-kappaB

signaling pathways. *Microb Pathog* 133: 103541

Figure 1

Figure 2

Figure 3

dTHP-1 cells: GO Biological Process (FDR < 0.05)

Figure 5

Figure 6

A

A549 cells: GO Biological Process (FDR < 0.05)

B

Figure 7

Figure 8

Figure 9

Figure 10