Ocular elongation and retraction in foveated reptiles

Ashley M. Rasys[1], Shana H. Pau[2], Katherine E. Irwin[1], Sherry Luo[2], Paul A. Trainor[3,4], Douglas B. Menke[2], and James D. Lauderdale[1,5]

[1]Department of Cellular Biology, The University of Georgia, Athens, GA 30602, USA; [2]Department of Genetics, The University of Georgia, Athens, GA 30602, USA; [3]Stowers Institute for Medical Research, Kansas City, MO 64110 USA; [4]Department of Anatomy & Cell Biology, The University of Kansas School of Medicine, Kansas City, KS 66160; [5]Neuroscience Division of the Biomedical and Health Sciences Institute, The University of Georgia, Athens, GA 30602, USA

Running title: Eye morphogenesis in foveated lizards

Key words: Anole, chameleon, lizard, eye, development, fovea, ocular morphogenesis, myopia

Corresponding author and person to whom request should be addressed:
James D. Lauderdale
Department of Cellular Biology
University of Georgia
Athens, GA 30602, USA
Ph. 706-542-7433
Fax: 706-542-4271
e-mail: jdlauder@uga.edu

Douglas B. Menke
Department of Genetics
University of Georgia
Athens, GA 30602, USA
Ph. 706-542-9557
Fax: 706-542-3910
e-mail: dmenke@uga.edu

Funding:
This work was funded by National Science Foundation awards 1149453 to D.B.M. and 1827647 to D.B.M. and J.D.L. and a Society for Developmental Biology Emerging Models grant to A.M.R. A.M.R. was supported by NIH training grant T32GM007103 and by an ARCS Foundation Scholarship. P.A.T is supported by the Stowers Institute for Medical Research

Disclosure Statement: The authors have nothing to disclose. The funders of this research had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Abstract

Background: Pronounced asymmetric changes in ocular globe size during eye

development have been observed in a number of species ranging from humans to

lizards. In contrast, largely symmetric changes in globe size have been described for

other species like rodents. We propose that asymmetric changes in the three-

dimensional structure of the developing eye correlate with the types of retinal

remodeling needed to produce areas of high photoreceptor density. To test this idea, we

systematically examined three-dimensional aspects of globe size as a function of eye

development in the bifoveated brown anole, *Anolis sagrei*.

Results: During embryonic development, the anole eye undergoes dynamic changes in

ocular shape. Initially spherical, the eye elongates in the presumptive foveal regions of

the retina and then proceeds through a period of retraction that returns the eye to its

spherical shape. During this period of retraction, pit formation and photoreceptor cell

packing are observed. We found a similar pattern of elongation and retraction

associated with the single fovea of the veiled chameleon, *Chamaeleo calyptratus*.

Conclusions: These results, together with those reported for other foveated species,

support the idea that areas of high photoreceptor packing occur in regions where the

ocular globe asymmetrically elongates and retracts during development.

Key Findings

• The eyes of the brown anole, *Anolis sagrei*, and veiled chameleon, *Chamaeleo*

calyptratus undergo dynamic asymmetrical changes in ocular shape during

development.

• In both species, asymmetric elongation and retraction of the ocular globe is

associated with fovea morphogenesis.

• Pit formation and photoreceptor cell packing in the foveal area occur when the

corresponding region of the ocular globe is retracting relative to adjacent regions.

Introduction

Decades of experimental work have revealed a great deal about the developmental

mechanisms that govern the patterning, differentiation, and growth of the vertebrate

eye. Much of this understanding has come from functional studies in mouse, rat,

chicken, *Xenopus,* zebrafish, and medaka.[1,2] Additional insights have come through

genetic studies of human syndromes that feature eye defects.[3] Consequently, we know

a great deal about the genes and signaling pathways that regulate development of the

core structures common to all vertebrate eyes, including the cornea, lens and retina.[4-7]

Missing from our current understanding of vertebrate eye development, however, is

detailed knowledge about the developmental pathways that regulate the formation of

specialized structures that are only present in the eyes of certain vertebrate species.

These structures include the conus papillaris, areas or streaks, and foveae. Studies of

eye formation in diverse vertebrate groups are needed to determine how these

specialized structures form and to achieve a more complete understanding of vertebrate

eye development and evolution. Notably, modern investigations of eye development

have almost completely excluded reptiles, a tremendously successful amniote group

represented by over 10,000 extant species.[8]

Although eye development in reptiles remains poorly studied, other aspects of

reptilian biology have been actively explored. For instance, *Anolis*, a lizard genus with

approximately 400 recognized species, has served as an important model system for

studies of evolution, ecology, physiology, behavior, and neuroendocrinology for many

years.[9] More recently, *Anolis* has also emerged as a system to investigate reptile

development and the mechanisms that contribute to morphological evolution.[10-16] The

brown anole lizard, *Anolis sagrei*, is particularly well-suited for developmental studies

due its small size, ease of husbandry, continuous egg production, high fertility, and low

cost. In addition, *ex ovo* culture systems and gene-editing have been established for

this species, which presents the opportunity for pharmacological and genetic

manipulation of *Anolis* embryos during development.[12,17,18] Of interest for studies of eye

development, *Anolis* lizards possess specialized structures that include a bifoveated

retina and a highly vascularized conus papillaris.[19,20]

Here we describe morphological and histological aspects of eye development in *A.*

sagrei. We pay particular attention to alterations in ocular globe shape, which is an

interesting, but poorly understood, aspect of eye development. Although typically only

studied postnatally in the context of myopia in humans,[21-23] changes in ocular shape

during embryonic development have been observed in a number of foveated species,

including humans,[24-30] and non-human primates,[26,31] as well as geckos,[32,33] suggesting

the presence of a conserved morphogenetic mechanism. The bifoveated brown anole is

a good model system in which to study the mechanisms underlying fovea development

in a vertebrate eye. In this study, we provide the first systematic three-dimensional

assessment of the dynamic changes in ocular shape with an emphasis on ocular

elongation and retraction and its association with fovea formation.

Material and methods

Animals

All experimental procedures were conducted in accordance with the National

Institutes of Health Guide for the Care and Use of Laboratory Animals under protocols

approved and overseen by the University of Georgia (anoles) and Stowers Institute for

Medical Research (chameleons) Institutional Animal Care and Use Committees. *Anolis*

sagrei lizards were maintained in a breeding colony at the University of Georgia

following guidelines described by *Sanger et al.,* 2008.[34] Eggs were collected weekly

from natural matings and placed in 100 X 15 mm lidded petri dishes containing moist

vermiculite and incubated at 27-28°C and 70% humidity. Adults and hatchlings were

euthanized using methods consistent with the American Veterinary Medical Association

(AVMA) Guidelines for the Euthanasia of Animals.[35,36] *Chamaeleo calyptratus* were

maintained in a breeding colony at the Stowers Institute for Medical Research (Kansas

City, Missouri) following guidelines described by Diaz et al., 2015[37] and Diaz et al.,

2019.[38] Eggs were collected at the time of oviposition and incubated at 26-28˚C and

50% humidity on damp vermiculite. Male and female embryos of both species were

used for these studies.

Staging

Embryonic development of *Anolis* lizards typically takes place over a 30-33 day

period, starting with fertilization, which takes place internally.[39] Early embryogenesis

proceeds within the oviduct. *A. sagrei* embryos obtained from eggs that were collected

after egg-laying were staged as described by Sanger et al.*,* 2008.[39] Embryos younger

than those captured by the Sanger staging series were denoted with the prefix "PL" for

pre-laying followed by a number. We describe here 5 PL timepoints, which includes the

first few embryos of the Sanger staging series (Sanger St 1-3 correspond to PL 3-5). PL

stage embryos were collected from gravid adult females following euthanasia. *C.*

calyptratus embryos were staged following criteria described by Diaz et al., 2017 and

Diaz et al., 2019 and stage matched to the anole using Sanger's morphological

criteria.[38-40]

Dissection

Lizard embryos were removed from their shells using a blunt pair of forceps and iris

scissors in 1x phosphate-buffered saline (PBS; 137 mM NaCl, 2.7 mM KCl, 10 mM

Na_2HPO_4, 1.8 mM KH_2PO_4, pH 7.4). Upon removal of yolk and amniotic sac with fine

forceps, embryos were placed into 60 ml of fresh 1x PBS solution with 1 ml of 0.4%

pharmaceutical grade, neutrally buffered tricaine (TRICAINE-S; Western Chemical Inc)

to anesthetize for imaging. Eyes were enucleated from embryo stages >4 with fine

forceps and placed in Bouin's fixative at 4˚C overnight on a rocker. Following fixation,

eyes were washed five times at 15 min per wash in 1x PBS. Specimens were stored in

70% ethanol solution (EtOH) at 4˚C until processed for histology. Whole chameleon

embryos were dissected from eggs in a similar manner, fixed in Bouin's at 4˚C

overnight, washed in 1x PBS, and stored in 70% EtOH prior to shipment from the

Stowers Institute for Medical Research. Upon arrival at the University of Georgia,

embryos were slowly rehydrated in a series of graded EtOH/PBS solutions. Once fully

rehydrated, eyes were carefully removed from embryos.

Whole Eye Measurements

Prior to fixing, anole eyes were positioned in both a lateral and dorsal orientation and

imaged with a ZEISS Discovery V12 SteREO microscope and AxioCam while in 1x PBS

solution. AxioVision 4 software (Carl Zeiss MicroImaging) was used to take axial

measurements along the dorsoventral (y), nasotemporal (x), and lateromedial (z)

aspects of imaged eyes from embryos at Sanger stages 4-18, the hatchling (Hch), and

the adult (Adt). The lateral view, which encompassed the whole cornea, was used to

measure the y- and x-axes, whereas the dorsal view was imaged to obtain x- and z-

axes. Because x-axial measurements can be acquired from both views, we included

only the dorsal x-axial value in our dataset and used the other value as a control for

proper orientation. Measurements from eyes that were not correctly positioned were

excluded from the dataset. We normalized the x- and z-axial measurements from each

lizard to that same individual's y-axis and then multiplied this number to the mean (μ) of

that individual's stage group y-axis (μ_y), $\frac{x \, or \, z}{y} * \mu_y$ = x- or z-axis$_N$. To identify trends in

ocular growth, a ratio was then calculated for each axis by taking the raw y-axis dataset

and the normalized datasets (x- and z-axis$_N$) from every lizard and dividing these values

with the corresponding μ of the raw y-, x-, and z-axial lengths of the hatchling,

$\frac{y-axis, \; x-axisN, \; or \; z-axisN \; (St \, 4-18; \, Hch; \, Adt)}{\mu y, \; \mu x, \; or \; \mu z \, (Hch)}$. Chameleon eyes were processed as reported

above except imaging was performed post-fixation and rehydration. Prism 7 (GraphPad

Software) and JMP V14.1 (JMP SAS) were used for graph generation and data

analyses. As a few of the sample groups did not have a normal distribution, we used a

nonparametric one-way ANOVA (Kruskal-Wallis) and Mann-Whitney for our statistical

analyses.

Paraffin Sectioning

Eyes were dehydrated in a series of graded ethanol solutions 70%, 80%, 90%, 96%

and 100% (twice) for a minimum of 15 min each and then soaked in xylene for a total of

30 min for all embryonic stages. Tissue specimens were incubated in a series of 3

paraffin wax jars for 30 min at 65˚C, embedded in paraffin, and serially sectioned

horizontally at 10 µm. In adult specimens, the dorsal aspect of the eye was punctured

with a 0.15 mm minutien pin prior to processing in xylenes and paraffin waxes to

facilitate wax entry. Processing time in xylene in the adult was extended up to a total of

2 hrs. Eyes were serial sectioned on a horizontal plane. Sections were stained with

hematoxylin & eosin following standard protocols and mounted in Cytoseal (Thermo

Scientific™ Richard-Allan Scientific™). Photomosaic images were generated using a

KEYENCE BZ-700 microscope with Keyence image stitching software. Adobe

Photoshop CC (2017.01 release) was used to digitally enhance contrast and adjust

white balance of images.

Results

Anatomy of the adult anole eye

Laterally positioned in the skull, the adult eye externally is oblate spheroid in shape

with a prominent convex cornea slightly biased toward the nasal region (Figure 1a,b).

Peripheral to the cornea is the sclera sulcus, whose curvature is supported by 14 sclera

ossicles (Figure 1b,c). The sclera ossicles are uniformly orientated in a patterned ring

except for a slight extension in the temporal region of the eye (Figure 1c). The anole's

radial pupil is positioned centrally, fashioned by a heavily pigmented iris with an array of

iridophores and melanophores that extend into the circumferential sclera sulcus (Figure

1b,c). The iris is asymmetric with dorsal and ventral notches defining the boundary

between the larger temporal and smaller nasal region (Figure 1c, arrowheads). Dorsally,

a protruding blood vessel is present that extends from the optic nerve, wraps around the

region of the center fovea (a small bulge in the medial region), and dissipates towards

the dorsal nasal area of the eye (Figure 1b, narrow arrowheads). The optic nerve (not

shown) exits the eye ventrally and temporally to the central fovea.

Internally, the adult anole eye is very similar to other vertebrates, possessing a

cornea, iris/ciliary body, lens, and retina (Figure 2a). Anteriorly, the transparent cornea,

which is exterior to the lens and iris, is composed of epithelial, stromal, and endothelial

layers (Figure 2a, Rasys et al.*, in prep*). These layers thicken towards the limbal area

where the cornea and anterior margin of the sclera ossicles meet. The sclera ossicles

extend from the limbal area outward towards the sclera proper in overlapping sheets

making up the sclera sulcus. Underlying this sclera sulcus is the long thin ciliary body

(Figure 2a). The ciliary body lacks a ciliary process and comprises an inner non-

pigmented layer and an outer pigmented epithelial layer (Rasys et al.*, in prep*). These

layers extend from the neural retina to the iris-ciliary boundary. Beyond this boundary is

the iris proper which is closely associated with the lens (Figure 2a). Both the inner and

outer iris epithelium are pigmented. The lens is oval shaped with a central nucleus,

inner cortex, and outer annular pad (Figure 2a). Anchoring the lens are zonular fibers

that extend from the ciliary inner non-pigmented epithelium (Figure 2a). Posteriorly, the

eye includes the retina, retinal pigmented epithelium, choroidal, and sclera layers. The

neural retina is avascular and composed of a ganglion cell layer (GCL), inner and outer

plexiform layers (IPL and OPL), inner and outer nuclear layers (INL and ONL) (Figure

2b,c).

Each eye has two foveae with one in the central retina and a second in the temporal

retina (Figure 2). The central fovea, located slightly temporal (4-5 degrees) to the optical

axis (the central or mid-point of the eye), exhibits a distinct pit accompanied by a higher

density of photoreceptor cells compared to the peripheral retina, as has been previously

described for other anoles.[19,20] The retina at the central foveola is devoid of the GCL,

INL and ONL (Figure 2a,c). The retina in the parafoveal region has a larger number of

cell bodies in the INL and ONL compared to the peripheral retina (Figure 2a,c). A

second, shallower fovea is located in the temporal retina roughly 45 degrees from the

optical axis or 40 degrees from the visual axis (central fovea) (Figure 2). This temporal

fovea exhibits a shallow pit accompanied by an increase in photoreceptor cell density.

At its center, all the retinal cell layers are retained, but layers are thinner than the

surrounding peripheral retina (Figure 2b). The temporal fovea is approximately one-third

the area of the central fovea, with fewer photoreceptor cells.

Eye formation

In anoles, fertilization takes place internally and embryonic development occurs over

a 30-33 day period.[39] As a consequence, the earliest stages of eye development take

place while the eggs are in the oviduct. Embryonic stages pre-egg lay are denoted with

the prefix "PL". Stages post-egg lay use the Sanger staging series[39] and are denoted

using the prefix "St". Stages PL 3-5 correspond to Sanger stages St 1-3.[39] For these

earliest timepoints we use the PL nomenclature to help distinguish pre-lay stages (PL 1-

5) from post-lay stages (Sanger St 4-19).

The initial stages of eye development occur between PL 1 and PL 5. Optic vesicle

formation is evident in embryos at PL 1 (Figure 3a), which is about 1 day after

fertilization and 3 days before egg lay. The lens placode is visible at PL 2 (see

developmental poster in supplementary data). By PL 3, the optic cup and lens pit are

present (Figure 3b). Rapidly following this period, the cornea separates (PL 5) from the

newly minted lens vesicle (PL 4) and the optic fissure begins to close (PL 5)

(developmental poster). At the time of egg lay (~St 4), the anole eye is spherical, trace

amounts of pigmentation are evident in the temporal region of the eye (Figure 3c) and

present nasally by St 5 (Figure 3d). At this time (St 5), the prospective central and

temporal foveal regions are evident as thickenings in the retina, also defined as retinal

mounding (Figure 3g, *Rasys et. al.,* in prep). St 5 also marks the first appearance of the

sclera sulcus, evidenced by a slight depression in the temporal area, which extends to

the nasal side by St 6 (Figure 3d). Eventually, the sclera sulcus forms a complete ring

encircling the cornea (developmental poster, *whole eye stages 6-8*). Shortly following

this period, pigment begins to increase - initially in the iris between St 6-10 and then

throughout the rest of the eye. In the iris, pigment is deposited first as a narrow band

along the horizontal axis (St 6-7) before radiating outward throughout the dorsal and

ventral regions (St 8-9) (Poster, *Whole eye*). By St 10, pigment in the iris is black and

evenly distributed. Granules are also just becoming obvious throughout the whole eye,

but more so in the temporal region. At St 13, retinal mounding is no longer present in

the central and reduced in the temporal foveal regions (Figure 3h). The eye is a light

brown color which darkens between St 15-17 and is completely black by the time of

hatching (Figure 4b; developmental poster). During this period the sclera ossicles that

shape the sulcus and provide support to the underlying ocular structures, are starting to

form. The sclera anlagen first manifests as a ring of pale conjunctiva papillae around the

cornea between St 11-12. By St 13, scleral sheets are present although they are small

and by St 15-16 these expand, radiating outward and eventually overlapping (St 17)

with the neighboring plates (developmental poster). Iridophores (reflective pigments)

scattered throughout the iris and sclera sulcus region, are also apparent during this

time.

Eye morphogenesis

The anole eye exhibits conspicuous asymmetric changes in morphology during

development (Fig 4). To assess potentially complex changes in the three-dimensional

shape of the globe, measurements were made along the three anatomical axes of the

eye at different stages of development. For this study, the dorsoventral axis was defined

as the y-axis, the nasotemporal axis was defined as the x-axis, and the lateromedial

axis was defined as the z-axis (Figure 4a). The lateromedial (z) axis is also the optical

axis and passes through anterior (iris and lens) and posterior (retina) structures of the

eye.

At St 5, the ocular globe is mostly spherical in shape (Figure 4b), with similar lengths

along the three anatomical axes. At late mid-gestation (St 13-14), the ocular globe has a

triangular appearance with increased lengths along the nasotemporal (x) and

lateromedial (z) axes compared to the length of the dorsoventral axis (Figure 3e,h;

Figure 4b). Interestingly, whereas the nasal surface of the globe has a rounded

appearance, both the temporal and medial surfaces have angular shapes. These

differences in surface geometry suggest that elongation in the nasotemporal and

lateromedial planes occurred largely by changes in the temporal and medial regions of

the globe, respectively. The medial region of the globe corresponds to the area of the

retina that gives rise to the central fovea, and the temporal region corresponds to that of

the temporal fovea (Figure 3g-i). The eye at this stage also exhibits considerably less

pigmentation at the medial surface compared to the lateral surface (Figure 4b). At

hatching, the ocular globe has a spherical shape with uniform pigmentation (Figure 3i;

Figure 4b). Remarkably, the globe at this stage has a smaller surface area than that of

mid-gestational embryos (Figure 4b). The change in morphology and size of the globe

in a hatchling compared to that of a mid-gestational embryo suggests that the eye

undergoes asymmetric retraction in the areas encompassing the developing central and

temporal foveae during the period between mid-gestation and hatching.

To better understand the dynamics of globe morphogenesis in the brown anole, axial

measurements along the three anatomical planes were made of eyes at embryonic

stages St 4-18, hatchlings (Hch), and adults (Adt) (Figure S1; Table S1). To facilitate a

quantitative comparison of changes in the three-dimensional shape of the eye, the three

measurements of each eye were converted to a standardized metric (see *Whole Eye*

Measurements in Methods). Briefly, in this metric the axial length of the y-axis, which

did not exhibit appreciable elongation and retraction during development, was used as a

normalization factor for measurements along the x- and z-axes. Variance due to

differences in embryo body and eye size within an individual's stage group was handled

by multiplying the normalized measurements for each individual by the group mean

value (μ) for the y-axis (Figure S1; Table S1). This operation resulted in an overall

reduction in variance seen at each stage in both x- and z-axis$_N$ datasets, which

suggests that the degree to which an eye elongates and retracts is proportional to the

embryo's eye size (Figure S2).

Using the hatchling eye as a reference, regional differences in globe morphogenesis

as a function of developmental stage were assessed by taking the raw y-axis dataset

and the normalized datasets (x- and z-axis$_N$) from every lizard and dividing these values

with the corresponding mean of the raw y-, x-, and z-axial lengths of the hatchling (see

Whole Eye Measurements in Methods). This analysis revealed four distinct phases in

ocular morphogenesis (Figure 4c).

Phase 1, which occurs between embryonic St 4-8, is characterized by rapid growth

of the eye. In embryos at St 4-5, the globe appears to expand uniformly along all 3

axes, which suggests that ocular growth during this period is equally distributed across

the eye. Although the St 5 eye is 50% smaller than the hatchling eye, the eye at these

two stages is quite similar in overall shape (Figure 4b,c; Table S1). At St 6, the globe

begins to exhibit asymmetric expansion, with more growth along the nasotemporal and

lateromedial axes compared the dorsoventral axis. By St 7, expansion along the

lateromedial axis is greater than along the nasotemporal axis.

The start of the phase 2 is defined as the developmental timepoint when

morphological asymmetry of the globe is clearly visible. This condition is met at St 8. It

is also at this point that the overall globe size is similar to that of the hatchling. Between

St 8 and the close of the second phase at St 14, the globe continues to expand

asymmetrically along the nasotemporal and lateromedial axes, with the more

pronounced expansion along the lateromedial axis (Figure 4c,d). The globe reaches

maximum lengths along the nasotemporal and lateromedial axes by St 14. At this stage

the length of the lateromedial axis is 1.4x that of the hatchling, and the length of the

nasotemporal axis is 1.2x that of the hatchling (Figure 4b,c).

Phase 3 is the epoch during which the globe begins to shorten in length along the

nasotemporal and lateromedial axes to regain a spherical shape. This phase

encompasses St 15-18 (Figure 4c,d), and corresponds to the time during which the

fovea acquires its distinctive morphological characteristics (Figure 3h,i). Upon

completion of retraction, the eyes are slightly smaller than those of hatchlings (St 17-18;

Figure 4c,d). The fourth, and final, phase is characterized by a uniform expansion of the

globe, which begins at the close of St 18 and continues into adulthood (Figure 4c). From

hatching, the eye doubles in size by the time the lizard reaches adulthood (Figure 4c).

7   To determine the magnitude of the asymmetric shape changes during ocular

morphogenesis, measurements capturing the maximum extent of elongation (St 13-14,

Figure 4c) were compared to those capturing the maximum extent of retraction (St 18-

Hch, Figure 4c). Axial measurements of globes from lizards at St 13, 14 and 18, and

hatching were compared using one-way ANOVA analyses (nonparametric Kruskal-

Wallis test); normalized values were used for the nasotemporal and lateromedial axes

(for mean and standard deviation, see Table. S1). Although measurements along the

dorsoventral axis were not statistically different across these four developmental time

periods (p-value 0.1585; alpha = 0.05), significant differences were observed for

measurements along both the nasotemporal and lateromedial axes (p-value <0.0001 for

both; alpha = 0.05).

Differences between the groups were compared using the Mann-Whitney test

(Figure 4d). Among the normalized nasotemporal (x-axis$_N$) datasets, St 13 and 14

measurements were significantly different from the St 18 and the hatchling (p-value

<0.0001 for both; alpha = 0.05) but not between St 13 and 14 (p-value 0.1244). Similar

results were observed among the normalized lateromedial (z-axis$_N$) measurements. For

St 18 compared to hatchling, x-axis$_N$ and z-axis$_N$ lengths were significantly different (p-

value <0.0001, and p-value of 0.0017, respectively). This suggests that by St 18 ocular

retraction is finished and the expansion phase is already well underway by the time of

hatching. The large difference in mean x- and z- axis$_N$ lengths between St 14 and 18 (z-

axis$_N$ 2647 to 1807 μm; x-axis$_N$, 2663 to 2151 μm) equated to a 32% and 19% reduction

in the central and temporal regions, respectively (Figure 4d).

Changes in intraocular pressure can be one mechanism that drives globe

expansion and retraction. Using an inflated ball as a model, morphological indicators of

a pressurized globe could include a taut ocular surface that resists deformation.

Conversely, using an under-inflated ball as a model, a previously-pressurized globe that

has lost pressure might exhibit wrinkles or folds of the surface and be more flaccid. The

surfaces of globes prior to St 16 were smooth and taut to the touch. In contrast, starting

at St 17, the exterior ocular surface was easily depressed with a pair of dull forceps and

wrinkles could be clearly seen in about half of the eyes examined (7/13 St 17 lizards)

(Figure 5). At St 18, wrinkling could still be detected (8/19 lizards), although it was not

as pronounced. By hatching, the majority of globes (9/10 lizards) were completely

smooth and once again the ocular surface was taut (Figure 5a). For developmental

stages between St 14 and hatching, wrinkled eyes tended to have reduced normalized

measurements along the lateromedial (z-axis$_N$) compared to smooth eyes (Figure 5c).

Ocular elongation & retraction in other foveated lizards

To test if ocular elongation and retraction occurs in other foveated lizards, ocular

morphogenesis was examined in the veiled chameleon lizard, *Chamaeleo calyptratus*.

This species was chosen because it possesses a single, prominent central fovea that is

completely devoid of all retina cell layers.[41] Chameleon embryos were collected at

several time points throughout development. To facilitate comparison with anoles,

chameleon embryos were staged following criteria described by Diaz et al., 2017 and

Diaz et al., 2019 and then matched to the anole using Sanger's morphological criteria.[38-

40] Only embryonic stages were used for this study; chameleon embryos collected just

prior to the expected hatching date are denoted as pre-hatch "pHch" (Table S2).

As observed for *A. sagrei*, ocular globe morphogenesis in *C. calyptratus* includes

pronounced asymmetric elongation along the lateromedial axis (Figure 6). In contrast

with *A. sagrei*, *C. calyptratus* embryos exhibit comparable expansion along both the

dorsoventral and nasotemporal axes (Figure 6). As with the anole, ocular elongation is

followed by a period of retraction that ends when the globe regains a spherical shape.

Compared with the anole, the elongated region in chameleon is less acute and occurs

over a broader area; in the anole, the elongated medial face appears more acute and

funnel-like (Compare St 13 chameleon in Figure 6B to St 13 anole in Figure 4b).

Despite this difference in morphology, the onset of ocular elongation and retraction

timing is nearly identical to anoles. For instance, ocular elongation begins around stage

6 and peaks by stage 14. This is followed by ocular retraction, which starts at stage 15

and plateaus just before hatching (Figure 6c). The chameleon fovea also develops

during the period of retraction (between stages 16-18) (Figure 6c), and its progression

also matches that of the anole (data not shown).

Discussion

The work we present here is a first step in establishing the brown anole as a new

model organism for fovea developmental studies. A primary motivation for choosing this

lizard is that the anole eye contains two fovea that represent the extremes of fovea

morphology: a central fovea with a pit devoid of all retina cell layers and a temporal

fovea with a shallow pit that retains the retinal layers. Most vertebrate species do not

have fovea, but in those that do, the fovea can differ greatly in pit shape, depth, and

diameter.[19,42-45] In humans, only the GCL and INL retinal layers are laterally displaced,

while the ONL is retained. This results in a broad and relatively shallow foveal pit with a

high density of photoreceptor cells at its center.[30,46-50] This is also true of most foveated

non-human primates. Although in some species, the GCL and/or INL are retained, and

as a result, only a rudimentary pit is present.[45,51-54] The fovea of birds is variable, and,

like the anole, some species are bifoveated, while others only possess a single fovea

located either in the central or temporal retinal regions.[44]

Anoles are the only squamate genus known to have a bifoveated retina. Among

the anole species studied, all have a prominent central fovea devoid of cell layers and a

shallower temporal fovea that retains these layers.[19,20,55] Slight variations in pit depth

have been observed across anoles and correlate with prey size.[19] For instance, anole

species that routinely eat smaller prey have considerably deeper temporal foveae. As in

birds, the location of the fovea in different squamate reptiles varies. In diurnal geckos

only a temporal fovea is present.[56-60] Although all layers are generally present in these

lizards, the degree of lateral displacement ranges from partial to full layer retention. In

Lygodactylus, which is a gekkonid, and *Gonatodes*, which is a sphaerodacylid, the GCL

is nearly absent, while the INL is only thinned and ONL packing is present at the foveal

pit center.[58] In contrast, *Phelsuma* geckos have only the shallowest of depressions

absent of any pronounced displacement of GCL and INL, resulting in a pit similar to the

temporal fovea of anoles.[58,59] In other lizards, including chameleons, the opposite is

generally true, with most having a large prominent central fovea devoid completely of

retinal cell layers.[61,62]

We observed four distinct phases of ocular morphogenesis in anoles and

chameleons. The first was a period of symmetrical growth that occurred during the first

week of embryonic development post-fertilization. At the end of this period, the retina

was thicker and exhibited a mounded appearance in the prospective foveal regions

(Figure 3g). This was followed by a second period, defined by asymmetrical growth,

where the regions that eventually gave rise to the fovea became strikingly elongated.

Coincidingly, this period also marks the gradually disappearance of retinal mounding

within the foveal regions (*Rasys et. al.,* in prep). By late development, when the foveae

take on their characteristic morphology, these regions appear to undergo retraction

coincident with retinal remodeling, i.e., pit formation and photoreceptor cell packing. The

fourth phase was characterized by a uniform expansion of the globe. In both the brown

anole and the veiled chameleon the regions that undergo elongation and retraction are

localized to areas of the retina where the foveae develop. Additionally, these foveal

regions, characterized by early retinal mounding, undergo retina differentiation and

lamination prior to the rest of the retina (*Rasys et. al.,* in prep). These observations

suggest a relationship between changes in retinal differentiation, ocular shape, and

foveal development.

We propose that ocular elongation followed by retraction are necessary steps in

the retinal remodeling needed to generate a fovea in vertebrates. Consistent with this

idea, evidence of asymmetrical globe development can be seen in the eyes of diurnal,

but not nocturnal, gecko species. Examples of this phenomenon can be seen in Figures

2 and 3 of the publication by Guerra-Fuentes and colleagues addressing the

embryology of the retinal pigmented epithelium in five species of sphaerodactyls,[32] and

in Figures 3 and 4 of the publication by Griffing, Gamble and Bauer addressing pigment

development in nocturnal gekkonids (*Gekko kuhli*, *Lepidodactylus lugubris*) and diurnal

gekkonids (*Phelsuma laticauda*, *Sphaerodactylus macrolepis*).[33] Among mammals,

asymmetric globe development has been observed only for foveated haplorrhine

primates but not for non-foveated primates or other mammalian species.[24-31] Together,

these observations suggest that fovea morphogenesis is similar among foveated

vertebrates. It will be interesting to learn if the eyes of foveated birds and fish undergo

similar morphogenetic changes during development.

In the brown anole, we noted that the magnitude of asymmetric ocular shape

changes during development appeared proportional to the extent of retinal remodeling

associated with formation of the morphologically dramatic central fovea compared to the

less distinct temporal fovea. We speculate that the process of elongation and retraction

of required for fovea formation, and that the relative extent to which an eye elongates

and retracts during fovea-genesis is directly proportional to the amount of retinal

remodeling required to make that particular fovea. If true, this may explain the variability

of fovea morphology present within each foveated species.

Several different mechanisms could mediate changes in the size and shape of

the ocular globe during development. We hypothesize that changes in intraocular

pressure (IOP) contribute to asymmetric ocular morphogenesis in the brown anole. This

idea arises from our observations that the embryonic anole eye appears to be

pressurized during peak periods of elongation and deflated during retraction. The idea

that IOP can drive ocular growth is not a new one. Previous studies have shown that

IOP plays a pivotal role in regulating normal ocular growth in chick[63] and increases in

IOP can lead to induced myopia (generalized axial elongation of the globe) in this

animal.[64,65] In foveated primates, Hendrickson and Springer proposed a model where

high IOP induces pit formation due to inherent increased elasticity present at the foveal

avascular zone, while "retinal stretching" induced by ocular growth, facilitates the

centripetal movement of photoreceptor cells towards the foveal center.[66-68] Although, it

is possible that lack of blood vessels would predispose this region of the primate retina

to be more susceptible to IOP and, therefore, form a pit, this cannot explain pit

formation in the lizard. Anoles have a retina that is entirely avascular.[43,55] This suggests

that if regional differences in retinal elasticity are present in the anole, they are unlikely

to be caused by avascular zones. Another challenge with this model, as applied to

anoles, is that it requires IOP to be high for pit formation to occur. In the brown anole, pit

formation occurs during the period that the eye is soft, indicating that IOP is low.

We propose, for anoles at least, that high IOP is involved in facilitating ocular

elongation, but that IOP is comparatively low during ocular retraction and pit formation.

As for Hendrickson and Springer's model, there must be additional mechanisms that

mediate regional differences in the elasticity of the foveal anlagen compared to other

regions of the ocular globe. We think it likely that regional and dynamic changes in the

elasticity of the tissues associated with the outer surface of the globe are required for

normal eye development in foveated lizards, and may be true for primates as well.

Acknowledgements

The authors thank Aaron Alcala, Sergio Minchey, Sukhada Samudra, Christina

Sabin, and Rebecca Ball of the Menke and Lauderdale research groups at the

University of Georgia, and Diana Baumann, Richard Kupronis, David Jewell, Alex

Muensch and Nikki Inlow of the Reptile and Aquatic Facility at the Stowers Institute for

Medical Research, for their help with animal husbandry, maintenance and care of the

anolis and chameleon colonies, respectively. The authors thank Drs. Jonathan

Eggenschwiler, Heike Kroeger and Robert Hufnagel, and Christina Sabin, Sukhada

Samudra, and Rida Osman for helpful discussions about this project and comments on

the manuscript.

References:

1.   Tsonis PA. *Animal models in eye research*. Academic Press; 2008.

2.   Wittbrodt J, Shima A, Schartl M. Medaka--a model organism from the far East.

Nat Rev Genet. Jan 2002;3(1):53-64. https://doi.org10.1038/nrg704.

3.   McKusick VA. Mendelian Inheritance in Man and its online version, OMIM. *Am J*

Hum Genet. Apr 2007;80(4):588-604. https://doi.org10.1086/514346.

4.   Miesfeld JB, Brown NL. Eye organogenesis: A hierarchical view of ocular

development. *Curr Top Dev Biol*. 2019;132:351-393.

https://doi.org10.1016/bs.ctdb.2018.12.008.

5.   Chow RL, Lang RA. Early eye development in vertebrates. *Annu Rev Cell Dev*

Biol. 2001;17:255-96.

6.   Fuhrmann S. Eye morphogenesis and patterning of the optic vesicle. *Curr Top*

Dev Biol. 2010;93:61-84. https://doi.org10.1016/b978-0-12-385044-7.00003-5.

7.   Saha MS, Servetnick M, Grainger RM. Vertebrate eye development. *Curr Opin*

Genet Dev. 1992;2(4):582-8.

8.   Uetz P, Stylianou A. The original descriptions of reptiles and their subspecies.

Zootaxa. Jan 24 2018;4375(2):257-264.

https://doi.org10.11646/zootaxa.4375.2.5.

9.   Losos JB, Schneider CJ. Anolis lizards. *Curr Biol*. Apr 28 2009;19(8):R316-8.

https://doi.org10.1016/j.cub.2009.02.017.

10.   Eckalbar WL, Lasku E, Infante CR, et al. Somitogenesis in the anole lizard and

alligator reveals evolutionary convergence and divergence in the amniote

segmentation clock. *Dev Biol*. Mar 1 2012;363(1):308-19.

https://doi.org10.1016/j.ydbio.2011.11.021.

11.   Park S, Infante CR, Rivera-Davila LC, Menke DB. Conserved regulation of

hoxc11 by pitx1 in Anolis lizards. *J Exp Zool B Mol Dev Evol*. May

2014;322(3):156-65. https://doi.org10.1002/jez.b.22554.

12.   Sanger TJ, Kircher BK. Model Clades Versus Model Species: Anolis Lizards as

an Integrative Model of Anatomical Evolution. *Methods Mol Biol*. 2017;1650:285-

297. https://doi.org10.1007/978-1-4939-7216-6_19.

13.   Sanger TJ, Revell LJ, Gibson-Brown JJ, Losos JB. Repeated modification of
early limb morphogenesis programmes underlies the convergence of relative
limb length in Anolis lizards. *Proc Biol Sci*. Feb 22 2012;279(1729):739-48.
https://doi.org10.1098/rspb.2011.0840.
14.   Sanger TJ, Seav SM, Tokita M, et al. The oestrogen pathway underlies the
evolution of exaggerated male cranial shapes in Anolis lizards. *Proc Biol Sci*. Jun
07 2014;281(1784):20140329. https://doi.org10.1098/rspb.2014.0329.
15.   Tollis M, Hutchins ED, Stapley J, et al. Comparative Genomics Reveals
Accelerated Evolution in Conserved Pathways during the Diversification of Anole
Lizards. *Genome Biol Evol*. Feb 1 2018;10(2):489-506.
https://doi.org10.1093/gbe/evy013.
16.   Infante CR, Mihala AG, Park S, et al. Shared Enhancer Activity in the Limbs and
Phallus and Functional Divergence of a Limb-Genital cis-Regulatory Element in
Snakes. *Dev Cell*. Oct 12 2015;35(1):107-19.
https://doi.org10.1016/j.devcel.2015.09.003.
17.   Infante CR, Rasys AM, Menke DB. Appendages and gene regulatory networks:
Lessons from the limbless. *Genesis*. Jan 2018;56(1).
https://doi.org10.1002/dvg.23078.
18.   Rasys AM, Park S, Ball RE, Alcala AJ, Lauderdale JD, Menke DB. CRISPR-
Cas9 Gene Editing in Lizards through Microinjection of Unfertilized Oocytes. *Cell*
Reports. Aug 27 2019;28(9):2288-+. https://doi.org10.1016/j.celrep.2019.07.089.
19.   Fite KV, Lister BC. Bifoveal vision in anolis lizards. *Brain Behav Evol*. 1981;19(3-
4):144-54. https://doi.org10.1159/000121639.
20.   Sannan NS, Shan X, Gregory-Evans K, Kusumi K, Gregory-Evans CY. Anolis
carolinensis as a model to understand the molecular and cellular basis of foveal
development. *Exp Eye Res*. Aug 2018;173:138-147.
https://doi.org10.1016/j.exer.2018.05.012.
21.   Chakraborty R, Read SA, Vincent SJ. Understanding myopia: Pathogenesis and
mechanisms. *Updates on Myopia*. Springer, Singapore; 2020:65-94.

22. Harper AR, Summers JA. The dynamic sclera: extracellular matrix remodeling in normal ocular growth and myopia development. *Exp Eye Res*. Apr 2015;133:100-11. https://doi.org10.1016/j.exer.2014.07.015.

23. Young TL. Molecular genetics of human myopia: an update. *Optom Vis Sci*. Jan 2009;86(1):E8-E22. https://doi.org10.1097/OPX.0b013e3181940655.

24. Badtke G. Entwicklungsmechanische Faktoren bei der Formgebung des embryonalen Augapfels. *Albrecht von Graefes Archiv für Ophthalmologie*. 1952/11/01 1952;152(6):671-688. https://doi.org10.1007/BF00685239.

25. Ehlers N, Matthiessen ME, Andersen H. The prenatal growth of the human eye. *Acta ophthalmologica*. 1968;46(3):329-349.

26. Pilleri G. [Comparative anatomical examination of the "protuberantia scleralis" of the eye]. *Acta Anat (Basel)*. 1960;41:131-7.

27. van L, Pilleri G. [Morphological research on the "scleral protuberance"]. *Albrecht Von Graefes Arch Ophthalmol*. 1961;163:1-9. https://doi.org10.1007/bf00684908.

28. Streeten BW. Development of the Human Retinal Pigment Epithelium and the Posterior Segment. *Archives of Ophthalmology*. Mar 01 1969;81(3):383-394.

29. Sondermann R. Die Bedeutung der Vererbung für die Entwicklung der Myopie. *Albrecht von Graefes Archiv für klinische und experimentelle Ophthalmologie*. 1950;151(1):200-208.

30. Hendrickson A. Development of Retinal Layers in Prenatal Human Retina. 2015;166:29-35.

31. Springer AD, Hendrickson aE. Development of the primate area of high acuity, 3: temporal relationships between pit formation, retinal elongation and cone packing. *Visual Neuroscience*. Mar 2005;22(2):171-185. https://doi.org10.1017/S095252380522206X.

32. Guerra-Fuentes RA, Bauer AM, Daza JD. The embryology of the retinal pigmented epithelium in dwarf geckos (Gekkota: Sphaerodactylinae): a unique developmental pattern. *BMC Developmental Biology*. 2014;14(1):29. https://doi.org10.1186/1471-213X-14-29.

33. Griffing AH, Gamble T, Bauer AM. Distinct patterns of pigment development underlie convergent hyperpigmentation between nocturnal and diurnal geckos

(Squamata: Gekkota). *BMC Evol Biol*. Mar 27 2020;20(1):40.

https://doi.org10.1186/s12862-020-01604-9.

34.   SANGER TJ, HIME PM, JOHNSON MA, DIANI J, LOSOS JB. Laboratory

protocols for husbandry and embryo collection of Anolis lizards. *Herpetological*

Review. 2008;39(1):58-63.

35.   Association AVM. AVMA Guidelines for the Euthanasia of Animals: 2020 Edition.

2020.

36.   Conroy CJ, Papenfuss T, Parker J, Hahn NE. Use of tricaine methanesulfonate

(MS222) for euthanasia of reptiles. *Journal of the American Association for*

Laboratory Animal Science : JAALAS. 2009;48(1):28-32.

37.   Diaz RE, Jr., Anderson CV, Baumann DP, et al. Captive Care, Raising, and

Breeding of the Veiled Chameleon (Chamaeleo calyptratus). *Cold Spring Harb*

Protoc. Aug 26 2015;2015(10):943-9. https://doi.org10.1101/pdb.prot087718.

38.   Diaz RE, Jr., Shylo NA, Roellig D, Bronner M, Trainor PA. Filling in the

phylogenetic gaps: Induction, migration, and differentiation of neural crest cells in

a squamate reptile, the veiled chameleon (Chamaeleo calyptratus). *Dev Dyn*.

Aug 2019;248(8):709-727. https://doi.org10.1002/dvdy.38.

39.   Sanger TJ, Losos JB, Gibson-Brown JJ. A developmental staging series for the

lizard genus Anolis: a new system for the integration of evolution, development,

and ecology. *J Morphol*. Feb 2008;269(2):129-37.

https://doi.org10.1002/jmor.10563.

40.   Diaz RE, Jr., Bertocchini F, Trainor PA. Lifting the Veil on Reptile Embryology:

The Veiled Chameleon (Chamaeleo calyptratus) as a Model System to Study

Reptilian Development. *Methods Mol Biol*. 2017;1650:269-284.

https://doi.org10.1007/978-1-4939-7216-6_18.

41.   Hulke JW. On the chameleon's retina; a further contribution to the minute

anatomy of the retina of reptiles. *Philosophical Transactions of the Royal Society*

of London. 1866;(156):223-229.

42.   Slonaker JR. *A comparative study of the area of acute vision in vertebrates*.

Boston: Ginn & company; 1897:1 p.l., p. [445]-502.

43.   Walls GL, Walls GL. The vertebrate Eye and Its Adaptive Radiation. 1942:1-785.

44.   Fite KV, Rosenfield-Wessels S. A comparative study of deep avian foveas. *Brain,*
2   *Behav Evol*. 1975;12:97-115.
3   45.   Ross CF. The Tarsier Fovea: Functionless Vestige or Nocturnal Adaptation? In:
4   Ross CF, Kay RF, eds. *Anthropoid Origins: New Visions*. Boston, MA: Springer
5   US; 2004:477-537.
6   46.   Hendrickson A. A morphological comparison of foveal development in man and
7   monkey. *Eye (Lond)*. 1992;6 (Pt 2):136-44. https://doi.org10.1038/eye.1992.29.
8   47.   Hendrickson a, Drucker D. The development of parafoveal and mid-peripheral
9   human retina. *Behav Brain Res*. Jul 31 1992;49(1):21-31.
https://doi.org10.1016/S0166-4328(05)80191-3.
48.   Hendrickson A, Possin D, Vajzovic L, Toth CA. Histologic development of the
human fovea from midgestation to maturity. *Am J Ophthalmol*. Nov
2012;154(5):767-778 e2. https://doi.org10.1016/j.ajo.2012.05.007.
49.   Hendrickson AE, Yuodelis C. The morphological development of the human
fovea. *Ophthalmology*. Jun 1984;91(6):603-12.
50.   Yuodelis C, Hendrickson A. A qualitative and quantitative analysis of the human
fovea during development. *Vision Res*. 1986;26(6):847-55.
51.   Webb SV, Kaas JH. The sizes and distribution of ganglion cells in the retina of
the owl monkey. Aotus trivirgatus. *Vision Research*. 1976;16(11):1247-1254.
https://doi.org10.1016/0042-6989(76)90049-3.
52.   Stone J, Stone J, Johnston E, Johnston E. The topography of primate retina: a
study of the human, bushbaby, and new- and old-world monkeys. [Case Report].
Journal of Comparative Neurology. Feb 20 1981;196(2):205-223.
https://doi.org10.1002/cne.901960204.
53.   DeBruyn EJ, Wise VL, Casagrande VA. The size and topographic arrangement
of retinal ganglion cells in the galago. *Vision Res*. 1980;20(4):315-27.
https://doi.org10.1016/0042-6989(80)90018-8.
54.   Collins CE, Hendrickson A, Kaas JH. Overview of the visual system of tarsius.
The Anatomical Record Part A: Discoveries in Molecular, Cellular, and
Evolutionary Biology. 2005;287A(1):1013-1025.
https://doi.org10.1002/ar.a.20263.

55. Makaretz M, Levine RL. A light microscopic study of the bifoveate retina in the lizard Anolis carolinensis general observation and convergence ratios. *Vision Res*. 1980;20:679-686.

56. Underwood G. Reptilian retinas. *Nature*. Feb 3 1951;167(4240):183-5.

57. Underwood G. On the classification and evolution of geckos. *Proceedings of the Zoological Society of London*. 1954/11/01 1954;124(3):469-492. https://doi.org10.1111/j.1469-7998.1954.tb07789.x.

58. Röll B. Gecko vision—retinal organization, foveae and implications for binocular vision. *Vision Research*. Jul 2001;41(16):2043-2056. https://doi.org10.1016/S0042-6989(01)00093-1.

59. Tansley K. The retina of a diurnal gecko. *Pflüger's Archiv für die gesamte Physiologie des Menschen und der Tiere*. 1961/05/01 1961;272(3):262-269. https://doi.org10.1007/BF00363014.

60. Tansley K. The gecko retina. *Vision Res*. 1964;4:33-37.

61. Hulke JW. On the Chameleon's Retina; A Further Contribution to the Minute Anatomy of the Retina of Reptiles. *Philosophical Transactions of the Royal Society of London*. 1866;156:223-229.

62. Detwiler SR, Laurens H. STUDIES ON THE RETINA THE STRUCTURE OF THE RETINA OF PHRYNOSOMA CORNUTUM. *J Comp Neurol*. 1920;32(3):347-356.

63. Schmid KL, Hills T, Abbott M, Humphries M, Pyne K, Wildsoet CF. Relationship between intraocular pressure and eye growth in chick. *Ophthalmic and Physiological Optics*. Jan 2003;23(1):25-33. https://doi.org10.1046/j.1475-1313.2003.00085.x.

64. Phillips JR, McBrien NA. Pressure-induced changes in axial eye length of chick and tree shrew: significance of myofibroblasts in the sclera. *Investigative Opthalmology & Visual Science*. Mar 2004;45(3):758-763. https://doi.org10.1167/iovs.03-0732.

65. Genest R, Chandrashekar N, Irving E. The effect of intraocular pressure on chick eye geometry and its application to myopia. *Acta of Bioengineering and Biomechanics*. 2012;14(2):3-8. https://doi.org10.5277/abb120201.

COMPLETE.

66.   Springer AD. New role for the primate fovea : A retinal excavation determines

photoreceptor deployment and shape. 1999;16(0):629-636.

67.   Springer AD, Hendrickson aE. Development of the primate area of high acuity. 1.

Use of finite element analysis models to identify mechanical variables affecting

pit formation. *Visual Neuroscience*. Jan 2004;21(1):53-62.

https://doi.orghttp://dx.doi.org/10.1017/S0952523804041057.

68.   Provis JM, Dubis AM, Maddess T, Carroll J. Adaptation of the central retina for

high acuity vision: cones, the fovea and the avascular zone. *Prog Retin Eye Res*.

Jul 2013;35:63-81. https://doi.org10.1016/j.preteyeres.2013.01.005.

Figure Legends

Figure 1. Adult anole eye. Top panel (a) shows adult male lizard with an enlarged view

of its left eye. Bottom panels show a lateral view (b) and dorsal view (c) of a right eye.

Directionality is designated by arrows and the letters T – temporal, N – nasal, D –

dorsal, and V – ventral. Markers indicate: white arrow heads with notches – dorsal

(green) and ventral (white) iris notches; asterisks (magenta)– individual sclera ossicle

sheets; narrow white arrow heads (b) – dorsal blood vessel, (c) temporal sclera ossicle

deformation; black arrow – optic nerve; crosshair – dorsal optical axis; and scale bars –

1 mm.

Figure 2. Anole eye organization and retinal layout. Left panel (a) shows a diagram

labeling structures present in a horizontally sectioned adult eye. Right panels show

magnified views of temporal (b) and center (c) foveae. Note: temporal fovea shown in

(b) is 150 µm deep to the plane of section in (a). Markers indicate GCL – ganglion cell

layer; INL – inner nuclear layer; ONL – outer nuclear layer; RPE – retina pigmented

epithelium; Ch – choroid; Scl – sclera; and scale bars – 1mm (a) and 50µm (b and c).

Figure 3. Developmental series of early, mid, and late embryonic stages of eye

morphogenesis. Top panel shows an array of early-stage embryos (PL1, PL3, and St 4)

during optic vesicle (a), len's pit (b), and lens separation (c). Bottom panel shows later

stages (St 5, 13, and Hch) along with histological sections through the eye's center

horizontal plane. Black arrow – temporal eye region; open arrows – center and closed

arrows – temporal retinal and fovea regions; and scale bars are 250 µm (a-c, g-i) and 1

mm (d-f).

Figure 4. Anole eyes undergo asymmetrical ocular elongation followed by retraction

during embryonic development. (a) Diagram illustrating orientation of y- (green), x-

(blue), and z-axis (magenta) ocular measurements. (b) Lateral (top) and dorsal (bottom)

views of whole right eyes from early (St 5), mid (St 13), and late (Hch) stages. Ocular

elongation and retraction phases are evident in the stage 13 embryo and hatchling,

respectively. All eyes are to scale with one another; scale bar is 1 mm. (c) Graph

displaying trends in ocular morphogenesis throughout development (St 4-Hch) and

adulthood (Adt F – adult female; Adt M – adult male). This graph was generated by

taking each lizard's ocular dimensions (y-axis and normalized x- and z-axis$_N$) and

dividing it by the mean (μ) hatchling ocular dimensions (y-, x-, and z-axis) to calculate a

ratio. (d) Direct comparison of ocular length along different axes reveals the degree of

ocular elongation (St 13-14) and subsequent retraction (St 18-Hch) in the anole eye.

Figure 5. The formation of surface wrinkles coincides with ocular retraction. (a) Dorsal

views of whole right eyes from stages 16 - Hatchling showing the progressive steps of

ocular retraction; scale bar – 1 mm. (b) An enlarged view of the St 17 eye from panel

(a); arrow heads mark folds/wrinkles present along the outer ocular surface. (c) ocular

dimensions along the z-axes relative to the hatchling eye; eyes where folds/wrinkles

were observed are indicated in magenta.

Figure 6. Chameleon eyes also undergo asymmetrical ocular elongation followed by

retraction during embryonic development. Diagram (a) demonstrates how y- (green), x-

(blue), and z-axis (magenta) ocular measurements were made. Image (b) shows lateral

(top) and dorsal (bottom) views of whole fixed right eyes from early (St 6), mid (St 13),

and late (St 18) embryonic development. Like the anole, ocular elongation (St 13) and

retraction (St 18) phases are evident. All eyes are to scale; scale bar is 1 mm. Graph

(c) summarizes trends in chameleon ocular morphogenesis throughout stages 4-pHch

(pHch – just prior to hatching). This graph was generated by taking each lizard's ocular

dimensions (y-axis and normalized x- and z-axis$_N$) and dividing it by the mean (μ) pHch

ocular dimensions (y-, x-, and z-axis) to calculate a ratio. Graph (d) compares the

degree of ocular elongation (St 14) and subsequent retraction (Hch and pHch) between

anoles (grey circles) and chameleons (color triangles: y-axis – green, x-axis$_N$ – blue,

and z-axis$_N$ – magenta). Calculations were made following same formula outlined above

in graph (c) for each respective species. Graph (e) indicates the relative percentage of

retraction occurring in chameleons between stage 14 and pHch.

Figure 1

Figure 2

a

Cornea

Limbus

Sclera ossicles

Iris

Ciliary body

Zonular fibers

Lens
Annular pad
Cortex
Nucleus

Temporal fovea region

Central fovea

Retina

Sclera

Text

Adult

b Temporal fovea

GCL

INL

ONL

Ch

O/IS

RPE

Scl

c Central fovea

GCL

INL

ONL

Ch

O/IS

RPE

Scl

Figure 3

Figure 4

c Trends of ocular morphogenesis

d Degree of ocular elongation and retraction

Figure 5

Figure 6

c Trends of ocular morphogenesis in chameleons

d Degree of ocular elongation and retraction between anoles and chameleons