Invivo and systematic analysis of random multigenic deletions associated

with human diseases during epithelial morphogenesis in Drosophila

Usha Nagarajan*[1#], Shanmugasundaram Pakkiriswami*[2#], Sandiya Srinivasan[3], Niveditha

Ramkumar[3], Rajeswari Rajaraman[3], Kumarasamy Thangaraj [3#]

[1]Central University of Haryana, Department of Biochemistry, Mahendragarh, Haryana, 123029
India
[2]Department of Biochemistry and Molecular Biology, Dalhousie University, Dalhousie Medicine
New Brunswick, Saint John, NB, E2L4L5,Canada,
[3] School of Chemical and Biotechnology (SCBT), SASTRA University, Thanjavur, 613401, India.
[4]Centre for Cellular and Molecular Biology (CCMB), Hyderabad, 500 007, India.
13
14 * Equal contribution
15 [#] Correspondence to: usha@cuh.ac.in sn374151@sdal.ca, thangs@ccmb.res.in

17 **Abstract**

Random loss of multigenic loci on chromosomes, a crucial drive for evolution,

occurs frequently in all living organisms. Analysis of such chromosomal

disruption and understanding the consequences of their impact on the growth

and development of multicellular organisms is challenging. In this report, we have

addressed this issue using invivo mosaic analysis of deficiency lines in Drosophila.

Genes on fly deficiency lines were compared with human orthologs for their

implications in disease development during cytoskeletal processes and epithelial

morphogenesis. The cytoskeletal phenotypes from the fly has been utilized to

predict the function of human orthologs. In addition, as these Drosophila

deficiency lines are equivalent to human microdeletions, based on the clonal

behaviour and phenotypes generated, a systematic analysis has been carried out

to establish the critical loci that correspond to Microdeletion Syndromes and

Mendelian Disorders in humans. Further we have drawn the synteny that exists

between these chromosomes and have identified critical region corresponding to

defects. A few potential candidates that might have an implication in epithelial

morphogenesis are also identified.

Introduction

Contiguous-gene deletion syndromes or multigenic deletion syndromes display
extremely variable clinical symptoms depending on the genes lost in the loci
(Rauch et al., 2001; Slavotinek, 2012). A great deal of evidence shows that
random loss of genes or spontaneous changes on chromosomes (like virus-
mediated mutagenesis in humans causing chromatid breakages, translocations,
duplications, deficiencies/deletions leads to developmental anomalies and growth
defects (Fortunato and Spector, 2003). "Less-is-more" hypothesis insists that gene
loss is an engine for evolutionary change and disease development (Olson, 1999;
Olson and Varki, 2003). Despite the fact that the adaptive role of gene loss has
formed the basis for evolution, impact of their adaptive biological functions on
individual integrity and stability remains elusive. Just as how loss and gain of
Homeotic genes and their clusters prove fundamental for the evolution of body
shape and size (Ruddle et al., 1994), the loss and gain of genes implicated in
epithelial morphogenesis also influence the integrity of body size and shape
(Schock and Perrimon, 2002). In a multicellular organism, epithelial
morphogenesis is an inevitable process in which epithelia, tissues that line the
organ and body surfaces, determine organ-to-body shape and size (Gumbiner
BM.1992; Schock and Perrimon, 2002). Of all the factors implicated in this
process, cytoskeletal components that balance rigidity and flexibility of the cell,
play a critical role in maintaining organ size and growth of the organism (Knust,
2005; Gibson and Perrimon, 2005; Shen and Dahmann, 2005; Donohoe et al.,
2018). Defects during this process lead to loss of tissue integrity; development of
genetic disorders and diseases (Olson and Varki, 2003; Knust, 2005 in Birchmeier
and Birchmeier, 2005).

Compared to loss of individual essential genes and frameshift mutations, gains
and losses of multiple (more than one) genes bring about greater dramatic

genetic repercussions. Random loss of contiguous (loss of genes in the same

order) genes would disrupt the normal development, affecting growth and

patterning of the tissue, which otherwise occur concurrently. Loss of such

contiguous genes that may occur due to any kind of genetic aberrations might

regulate both cell-autonomous and non-autonomous functions of the tissue

giving rise to heterogeneity in cell population (mosaicism). It is intriguing to

understand the influence of such sudden, random loss of genes causing

mosaicism and to compare their effect with other organisms.

Genomic studies suggest that genetic mosaicism is a very common phenomenon

during epithelial morphogenesis. Although such lesions to the genome are

random and generally asymptomatic in initial stages, affected individuals

subsequently develop varying levels of metabolic disorders and disabilities (Chial,

2008). Also, since such incidences occur sporadically, chances of undertaking

systematic analysis is very low or may not be feasible. In the past, rigorous

systematic analysis of genes (Reiter et al., 2001) and genetic screening (St

Johnson, 2002; Yamamoto et al., 2014) have been undertaken to understand

genes implicated in disease development and their functions. However, how

these individuals subjected to such sudden and random loss of genes manage

without their gene function and what impact does it have on the individual is not

clear. In multigenic deletions, the synergistic and antagonistic effect of genes in

maintaining homeostasis or disease development and the alternate pathways that

operate to compensate the loss is not known yet. Although it is intriguing to

understand the consequence of random loss of multiple genes that leads to

defects and genetic disorders, it is extremely challenging.

We hypothesized that those limitations can be overcome and unanswered

questions can be addressed by using the available *Drosophila* deficiency line

collections, one of the unique genetic tools well-utilized extensively for genetic

screening, mapping and characterization processes. Two types of molecularly

defined deficiency lines, whose details including ends of the genes lost as well as

cytological location on the chromosome are known, are available (Ryder et al.,

2004; Parks et al., 2004; Cook et al., 2012; Roote and Russell, 2012). In addition,

as chromosomes of these deletion-bearing fly stocks resemble those

chromosomes with microdeletion or chromosomal deletions of contiguous genes

(loss of genes in the same order) in other organisms, the collection has been

utilized for systematic and comparative analysis to establish the critical loci that

correspond to Mendelian Disorders and Microdeletion Syndromes in humans.

Based on this logic, in this work, we set out a pilot project by subjecting these

deficiency lines for mitotic clonal analyses in *Drosophila* wing imaginal discs to

identify and understand their role in cytoskeletal organization during epithelial

morphogenesis. Clonal analysis technique allows to analyse behaviour of two

different population of cells in a tissue (Lee and Luo, 2001; Wu and Luo, 2006;

Smith-Bolton et al., 2009; Rodriguez et al., 2012; Powell and Piddini, 2016). Here

we have utilized this property to understand the behaviour of randomly

eliminated genes, thereby identify the critical gene in that locus implicated in

epithelial morphogenesis. During this process of clonal analysis, we have also

followed the cytoskeletal defects produced by these deficiency lines. We also

undertook a systematic functional analysis of the genes in the fly deficiency lines,

compared them with human orthologs and analysed their implications in

cytoskeletal processes and epithelial morphogenesis. Further we have drawn the

synteny that exist between these chromosomes and identified critical region of

epithelial defects. Here, in this report, few potential candidates that might have

an implication in epithelial morphogenesis are discussed.

Results and Discussion

(i) Recombination of Deletion-bearing chromosomes with FRT chromosomes

and generation of mitotic clones

To address the behavioural property of the tissue subjected to multigenic

deletions we generated mitotic clones using the available deficiency lines in

Drosophila wing imaginal discs. We recombined X and II chromosome deficiency

lines with their respective FRT chromosomes. We have subjected ~67 FRT-

recombined X chromosome deficiency lines and 125 FRT-recombined second (2L

and 2R) chromosome deficiency lines for clonal analysis in III instar larval wing

imaginal discs. These deficiency lines produced clones of varied number and sizes

(Figure 1(A-O); Supplementary S1, Sheet2 – Sheet5). For instance, deficiency line

Df(1)Ex6237(BL-7711) with 21 genes deleted produced 103 clones, highest

average number of clones (Figure 1N) while deficiency line Df(1)ED6443(BL-9053)

with 45 genes deleted generated 0 to 1 (average =0.3), lowest number of clones.

Deficiency line Df(1)Exel6230(BL-7705) with 9 genes deleted developed the

biggest clones of 0.6 Sq. pixels, while deficiency line Df(1)Exel6221(BL-7699) with

14 genes deleted produced smallest clones of 0.04 Sq. pixels (Figure 1J),

Supplementary S1- Sheet2-3). Some of the deficiency lines did not generate

clones or generated few clones (<10) per wing disc. Interestingly few of the

deficiency lines showed clones specifically localized in wing discs, especially only

in pouch region (Figure1J) or only in notum of the wing disc (Figure 1F,

Supplementary 2(A-B).

In order to understand the clonal properties of the mitotic clones generated by

these deficiency lines, we have categorized the deficiency lines into very few (rare

or less than or upto 10 number of clones), few (upto an average of 10- 30

clones), moderate (upto 60 clones), and numerous (> 60 clones) clonal types and

based on the size of clones, deficiency lines were grouped into large, medium

and small types. On X chromosome, 37% of the deficiency lines generated

rare/very few clones and 27% generated few clones. Around 21% and 15% of

deficiency lines were categorized into moderate and numerous categories

respectively (Figure 2A, Supplementary S1- Sheet2). In addition, based on the

average size of the clones generated per wing disc, we have categorized the

deficiency lines from small through medium to large (Figure 2B, 2(E-F),

Supplementary Data S1 Sheet3).

To address the link between the number of genes lost in a deficiency line and

their corresponding clonal property, we have generated correlation graphs

between them. From the correlation graph, we find that depending on the

function and importance of genes deleted in a deficiency line, their clonal

properties and behaviour varied accordingly (Figure 2C, Supplementary S1-

Sheet2). For example, (i) on X chromosome there are four deficiency lines that

have very less number of genes lost on them but generated either rare clones

[Deficiency line (Df(1)Exel6244(BL-7717) which lost only 7 genes did not generate

clones at all while Df(1)Exel6221(BL-7699)which lost 14 genes generated very less

to 1 clone in a wing disc (Figure1J)] or very few clones [Df(1)Ex6241(BL-7715) lost

6 genes but generated on an average of only 4 clones (Supplementary Figure2C)

while Df(1)ED6579(BL-9518)(70.1) lost 10 genes but generated on an average of

10 clones per wingdisc (Supplementary Figure2D)]. (ii) Similarly, there are only

four deficiency lines in the moderate and numerous categories with more than 20

genes lost in them. Deficiency line Df(1)ED7170(BL-8898) lost 58 genes made on

an average 35 clones per wingdisc(Figure1K) and Df(1)ED6521(BL-9281) lost 25

genes generated on an average 46 clones per wingdisc (Figure(1L). Similarly, in

the numerous category, Df(1)ExelED6989(BL-9056)(76-1) that lost 32 genes made

on average of 65 clones per wingdisc (Supplementary Figure 2G). Of all these,

notable ones are Df(1)Exel6237(BL-7711) that lost 21 genes that made 103 clones,

the highest average number of clones per wingdisc (Figure1N) and

Df(1)Exel6223(BL-7770) that had lost only 3 genes but made 77 clones on an

average in a wingdisc (Supplementary Figure 2H). This deficiency line had the

least number of genes lost with the fourth highest number of clones

(Supplementary S1- Sheet2). There are nearly 8 deficiency lines on the X

chromosome and 26 deficiency lines on the II chromosome that did not

generated clones (Figure 2A and 2C, Figure 3A Supplementary S1 Sheet 2 and 4).

Similarly, we have performed clonal analysis on deficiency lines on II chromosome

and categorized them based on their number, size and distribution (Figure 3(A-D)

Supplementary Data S1 Sheet4-5). Nearly 27% of the deficiency lines were in

rare/ very few category while 23% of them were in few category. Around 31%

and 19% of deficiency lines were categorized into moderate and numerous

categories respectively. As in case of the X chromosome, the number of clones

generated by corresponding deficiency line depended on the gene type and

function. The deficiency line Df(2L)Exel6024, that is deficient in 19 genes, makes

the smallest clones that have an average size of 0.6 Sq. pixels. The largest clones,

with a mean size of 63, are made by the Deficiency line Df(2L)Exel9040, that lacks

only 2 genes. The line Df(2L)7042, that has lost 14 genes, makes on average,

clones sized 15 Sq. pixels while the line Df(2R)Exel6065, that has lost about 40

genes, makes but relatively small clones of average size 8 Sq. pixels (Figure 3(A-

D) Supplementary Data S1 Sheet4-5). These findings reinforce only the hypothesis

further that the clonal phenotype exhibited by the animals under study depends

on the function of the genes lost and their importance to the development

underway. We have utilized these clonal properties for systematic and

comparative analyses discussed in next sections to understand their implications

on human diseases.

2. Analysis of cytoskeletal changes associated with Drosophila deficiency lines

and functional analysis of their Human orthologs

To understand the role of genes in deficiency lines during epithelial

morphogenesis, we have analysed the cytoskeletal changes mediated by

generating mitotic clones in wing imaginal discs. Of the 66 deficiency lines on X

chromosomes subjected to clonal analysis, nearly 12 deficiency lines showed

cytoskeletal changes in the wing imaginal discs (Table 1). Although we generated

clones for both X and II chromosomes, we analysed the cytoskeletal defects for

entire X chromosomes deficiencies and part of the II chromosome deficiency lines

that showed visible defects either in larval or adult stages.

Based on the type and the degree of cytoskeletal changes, we found that clones

were either widely-distributed throughout the wingdisc or localized in wingdiscs.

Accordingly, we have classified them into six categories, namely, with (i)-no, (ii)-

mild, (iii)-severe morphological changes and with (iv)-outgrowths, (v)-extrusions

and (vi)-duplications (Table 1). Nearly 49 deficiency lines did not show any

cytoskeletal changes. Of the remaining deficiency lines, approximately 5

deficiency lines (Df(1)-Exel6226(BL-7703), Df(1)-Exel6233(BL-7707),

Df(1)Exel6290(BL-7753), Df(1)Exel6237(BL-7711), Df(1)ED6989(BL-9056) showed

very mild morphological changes. These deficiency lines generated medium-sized,

moderate to numerous clones in the wing discs. In the Df(1)Exel6226(BL-7703),

we could observe cell-autonomous cytoskeletal defects (Figure 4B(1-4)) while the

deficiency line Df(1)Exel6233(BL-7707), displayed non-cell autonomous

cytoskeletal defects involving cytoskeletal breakages and disruptions (Figure 4C(1-

4)). Similarly, for second chromosomes, we generated clones using conventional

method. Compared to the control wingdiscs (Figure 4D(1-4)), two deficiency lines

Df(2L)Exel8022(BL-7813) (Figure 4E(1-4)) and Df(2L)9040(BL-7815) (Figure 4F(1-4))

showed severe cytoskeletal defects. Two deficiency lines (Df(1)Exel6242(BL-7716)

Figure 5(A'-A", Figure 5C'))and Df(1)ED6584(BL-9348) (Figure 5(B'-B", Figure 5C")

showed cytoskeletal defects and extrusion of the clones associated with

outgrowths. Consistently, we find both deficiency lines show similar phenotypes

by both direct mosaic as well as MARCM techniques. Two deficiency lines showed

localized clones: Df(1)ED6584(BL-9348) produced clones near the periphery of

pouch region associated with the extrusion of rounded clonal cells (Figure 5B″

and 5C″); while the deficiency line Df(1)Exel6235(BL-7709) produced clones only

in the notum region displaying non-cell autonomous cytoskeletal rounding near

the clones, (Figure–5(D′-D″)). Similarly, three deficiency lines showed cytoskeletal

defects with outgrowths and duplication phenotypes. Of them, two from X

chromosome, (Df(1)ED6630(BL-8948) (Figure 6(B-D) and Df(1)ED6727(BL-8956)

(Figure 6(E-F)rarely produced clones, however, it showed severe morphological

changes, including outgrowth and duplication in the wingdiscs. One of the II

chromosome deficiency lines, Df(2L)9040(BL-7815), displayed outgrowth and

pouch duplication phenotype in the wingdisc (Figure7(A-D).

We have also undertaken a systematic functional analysis of the genes in the fly

deficiency lines, compared them with human orthologs and analysed their

implications in cytoskeletal processes and epithelial morphogenesis

(Supplementary Data S2). In the X chromosome of the fly, out of 1103 total

genes, there are 181 genes that are involved in such functions. Among the 181

genes, one-third includes transmembrane proteins, another one-third is involved

in cytoskeleton binding and organization, and others are involved in intracellular

vesicular transport, cell polarity establishment and maintenance of epithelial

morphogenesis. Similarly, in the II chromosome there are 431 genes out of 1245

which are implicated in cytoskeletal functions. We have compared these genes on

X and II chromosomes that have cytoskeletal implication with their respective

Human orthologs (Figure 8). In humans, 200 of the total orthologs are involved in

cytoskeletal functions. In some instances, the functions of the orthologs of

cytoskeletally implicated genes are known while in others it is known atleast in

the fly counterparts. About 122 genes and their orthologs both have cytoskeletal

functions. In numerous cases, the human orthologs of such genes have not been

functionally characterized as yet. For 18 genes on X chromosome, cytoskeletal

functions of fly orthologs are known but unknown in human orthologs (Table 2).

Nearly 11 genes have both fly and their corresponding human orthologs with

similar cytoskeletal functions. For example, Drosophila *kirre* of Df(1)ED6330 (Fig

1F) and its human ortholog *KIRREL3* is involved in cell adhesion and plasma

membrane fusion activities and the fly gene *SKPA* of Df(1)ED6443 and its human

ortholog *SKP1* is implicated in microtubule cytoskeletal organization. Similarly ,

gene linked to Notch regulation, Drosophila *sno* of Df(1)ED7170 (Fig 1L) and its

human ortholog *SBNO1* and gene connected to synaptic growth, fly *Efhc1.1* of

Df(1)ED7344 (Fig 1I) and its human ortholog *EFHC2* are implicated in cytoskeletal

morphogenesis. Few of the fly orthologs are uncharacterized while cytoskeletal

functions of their human orthologs are known. For example, fly orthologs

CG10804 and *CG16181* of human genes *TROVE2* and *MARCH5* respectively are

uncharacterized genes. These two genes are covered by the deficiency line

Df(1)Exel6233. In the same section above we have shown that Df(1)Exel6233

produced mild cytoskeletal defects (Fig 4). Therefore it is possible that the genes

CG10804 and *CG16181* may be implicated in cytoskeletal functions. Clonal

properties of 18 genes on respective deficiency lines are undertaken in this study

(Supplementary data S2(Sheet1-2); Table 2). The cytoskeletal phenotypes from the

fly can thus be utilized to predict the etymology of many rare, unexplored human

diseases.

3. A Systematic approach to compare genes on Drosophila deficiency lines

with Human Mendelian Diseases (MenDs) and Microdeletion Syndrome (MDs)

to draw synteny between them

We undertook a systematic approach for both X and II chromosomes to compare

the Drosophila genes on deficiency lines with their corresponding Human

Orthologs. We have classified them into two categories based on whether or not

they have available Human orthologs. There are 600 genes on X chromosome

and 1090 genes on Ⅱ chromosomes with Human orthologs. These groups with

Human orthologs were looked up for their association with disease enlisted in

Online Mendelian Inheritance in Man (OMIM) database. There are nearly 189

genes on X chromosome and 509 genes on Ⅱ chromosome with OMIM disease

association. Based on this criterion, we further categorized the diseases into

Mendelian disease and Microdeletion Syndrome. There are nearly 119 genes on

X chromosome and 227 on Ⅱ chromosome for Microdeletion syndrome (Figure

9(A-B), Supplementary data S3 (Sheet1-2)) and 70 genes on X chromosome and

202 genes on Ⅱ chromosome associated with Mendelian Disease (Figure 9(A-B),

Supplementary data S3(Sheet 7)).

Further we have classified and analysed the data in two ways. First, for any locus

on the human chromosome, we have identified all possible MDs or MenDs

pertaining to it. Therefore, any lesion on that particular locus can lead to any of

these disorders listed Supplementary data S3 (Sheet1-2), (Sheet 7)). Some of the

MDs are coded by more than one locus on either of these chromosomes while

some of them are either present on X or Ⅱ chromosome.   We have pooled and

compared the number of genes on both X and Ⅱ chromosome that represent the

corresponding MD syndrome to generate a comprehensive comparative chart

between them (Table 3, Figure9C).

Second, we found many human Microdeletion and Mendelian syndromes to

mirror the genetic deletions observed in the fly deficiency lines under study. We

could thus make comparison between them to identify etymologies of rare

variants of human genetic disorders. We have analysed the MDs by comparing

the order of genes as they are arranged on the human chromosomes (synteny).

In this study, we have found that there are number of genes in the order on

Drosophila chromosome to be represented in the same order on Human

chromosome leading to either Microdeletion syndromes–(Supplementary data

S3(Sheet 5-6)) or Mendelian disease (Supplementary data S3(Sheet 7)).

Interestingly there are also 16 Mendelian Disease genes on X chromosome of

Drosophila that correspond to genes on Human X chromosome (Table 4 or

Supplementary Data S3(Sheet 8)).

Until now, researchers and clinicians were finding it difficult to pin down the

respective genes that correspond to the disease manifestation. However, from

this comprehensive analysis, it is possible to identify the specific gene that lead

to MDs and MenDs. For example, we found that one of the MDs,

Holoproscencephaly, is caused by loss of several genes. Nearly there are 7 genes

on X and 23 genes on II chromosome that can lead to the manifestation of the

syndrome (Table 3, Supplementary Data S3 (Sheet1-2). Next highest are the MDs

like Autism (11 genes on X and 10 genes on II chromosome) and

Neurofibromatosis (10 on X chromosome and 13 on II chromosome).

Comparative loci analysis reveals that in the locus on human 2q37.3, either

Autism or Holoprosencephaly can occur. By comparing with the Drosophila

genes, it is possible to point out the genes and locus on the chromosome. While

loss of *per* (at 3B1-3B2) can lead to Autism, loss of *CG10353* (at 10F2-10F4) can

lead to Holoproscencephaly Supplementary Data S3 (Sheet1, Sheet3). Similarly,

lesions at loci 11p11.2 or 16p11.2 can lead to DiGeorge

Syndrome/Velocardiofacial syndrome and Holoproscencephaly. From this study

we find that loss of either *SPACA3* at 11p11.2 or *FAM57B* at 16p11.2 can lead to

DiGeorge Syndrome, while loss of either *PTPRI* at 11p11.2 or loss of *PHKG2* at

16p11.2 can lead to Holoproscencephaly (Supplementary data S3(Sheet1,

Sheet3)).

Similarly, the locus 5A12-5D1 encodes for several essential genes and therefore

any deficiency in this region may potentially give rise to MDs like Beckwith

Wiedemann Syndrome (BWS), Bracydactyly-Mental Retardation syndrome

(Bdmr1), (Branchiorenal syndrome, Bor1 and Campomelic Dysplasia). Deficiency

line Df(1)ED6802(BL-8949), pertaining to the locus 5C2-5C2, that encodes for

Carboxy peptidases A3 and B1 (CPA3) may lead to BWS as well as for Bdmr1.

This also suggests that Human orthologs of these genes could potentially be

involved in human X-linked MDs) (Supplementary data S3(Sheet1, Sheet3)).

Interestingly, we have found that there are MDs that are caused by loss of genes

only on X chromosome and not on II chromosome. For example, Rubinstein Taybi

Syndrome is caused by 10 loci, all located only on X chromosome and not on II

chromosome. Similarly, Potocki-Lupski Syndrome is caused by 5 loci on X

chromosome and not on II chromosome. Rett's syndrome and Tuberus sclerosis

are MDs caused by microdeletions in *MECP2* gene and *TSC1/2* respectively both

with single locus each on X chromosome. (Table3, Supplementary data S3(Sheet1,

Sheet3)). Corresponding deficiency lines analysed here are Df(1)ED6630-8948

(Figure 1E)　and Df(1)ED7170-9058(Figure 1K) that has the genes *CG2680* and

HDAC4 respectively representing the microdeletions causing Rett's syndrome and

Tuberus sclerosis.

From this analysis, in addition to the point that a particular MDS can be

represented by more than one locus, we also understand that, multiple MDs that

arise in a particular locus depend on the genes deleted and the order of

contiguous genes deleted. In addition, comprehensive and comparative analysis

between Human and Drosophila loci have led to identification of putative critical

region in human chromosome that could give rise to various MDs. This is

addressed in detail in the following section.

4. Identification of critical region or the gene implicated in epithelial defect

Although multiple genes are lost in the manifestation of MDs or Mendelian

diseases, there exists a critical region involving either an essential gene or a small

genomic region in that locus which is central in causing the disease. Since the

molecular ends of the deficiency lines used in the study were precisely defined

beforehand, we compared this information with the clonal phenotype or

behaviour generated to identify the critical region in that locus, necessary for the

maintenance of tissue integrity. We took two approaches while comparing the

overlapping deficiency lines, one based on the number and size of the clones

generated with respect to the number of genes eliminated (Supplementary S1

Sheet(2 and 4); and the other based on the cytoskeletal defects displayed by

them (Supplementary S3). For the former approach, using the deficiency lines on

X chromosome, we looked into the category with numerous clones (>60 clones

per wingdisc) generated by a smaller number of genes lost in them. Of the 10

deficiency lines in this category, deficiency line Df(1)Exel6237 (BDSC-7711)at 5C2-

5C6 with 21 genes lost including *Usp30* generated 103 clones, the highest

number of clones of all lines. Followed by that, Df(1)Exel6230(BDSC-7705) at 3A2-

3A4 with 9 genes lost generated biggest clones(with average size of 0.6 Sq.pixels)

and the second highest number of clones per wing disc (on an average 91

clones). Under this category, we also found an overlapping deficiency line

Df(1)Exel6231(BDSC-7706) for the same region at 3A2-3A3, that lost 6 genes but

produced 65 clones (6th highest number of clones) with average size of 0.4

Sq.pixels(one of the biggest clones). For the same locus, we also found two

bigger deficiency lines Df(1)ED11354 (BDSC-9345) at 2F6-3A4, with 15 genes lost,

but generating only 13 clones (with average size of 0.05 Sq.pixels); and

Df(1)ED411(BDSC-8031) at 3A3-3A8 with 18 genes lost, but produced only 3

clones on an average(with average of size 0.2 Sq.pixels). Of the four deficiency

lines spanning this locus, (Df(1)Exel6230(BDSC-7705) and Df(1)Exel6231(BDSC-

7706) show similar phenotype with large (0.6 and 0.4 Sq.pixels)and numerous

clones (91 and 65) respectively. By comparing these three deficiency lines

(Df(1)Exel6230(BDSC-7705), Df(1)Exel6231(BDSC-7706) with Df(1)ED411(BDSC-

8031), we found that 3A2-3A3, that contains the gene *Brother of ihog (Boi)* is the

critical region/gene responsible for producing high number of bigger clones

(Figure10A, 10E and 10H ; Supplementary Data S4 (Sheet1). It is understood that

the third biggest deficiency line Df(1)ED11354 (BDSC-9345) might include other

genes like *Raf,* another regulator of growth. *Boi* is suggested to be a negative

regulator of growth, that functions as a co-receptor for Patched(Ptc) in regulating

Hedgehog(Hh)-signaling mediated growth. Therefore, highest number of clones

generated is due to the elimination of *Boi,* the negative regulator of growth, in

that locus. Interestingly, loss of the gene *Cell adhesion associated, Oncogene*

regulated (CDON), human ortholog of Boi, localized at 11q24.7 is suggested to

cause MD Holoprosencephaly in humans.

As a second approach, on both X and II chromosomes, we looked into those

deficiency lines which generated cytoskeletal defects and wingdisc phenotypes

associated with them. On X chromosome, the deficiency line Df(1)ED6584(BDSC-

9348) (Figure11C) at 3A8-3B1 lost only 3 genes, but generated moderate number

of clone (average number of 36 and average size of 0.12 Sq.pixels). We compared

it with available overlapping deficiency lines Df(1)ED6679 (BDSC-9518)

(Figure11(A′-A″)) at 3A6-3A8 with 10 genes lost, that produced only 8 clones

(with average size of 0.08 Sq.pixels); and Df(1)ED411(BDSC-8031) (Figure11B) at

3A3-3A8 with 18 genes lost produced only 3 clones on an average(with average

of size 0.2 Sq.pixels). The clonal analysis of the deficiency line Df(1)ED6679

(BDSC-9518) by both conventional method Figure 11 A-A′) and MARCM

technique (Figure11(D′-D″)) showed several cytoskeletal defects in the hinge

region, with small rounded cells around the periphery of the hinge. However, the

clonal analysis of Df(1)ED6584(BDSC-9348) showed cell-autonomous extrusion of

the clones from the wing surface (Figure11(E′-E″)). By comparing the overlapping

region between them and the similar phenotype produced by Df(1)ED6679

(BDSC-9518), we could identify the critical region is coded by the gene

Shaggy(Sgg). As *Sgg* is a key component of β-catenin destruction complex so as

to function as a negative regulator of Wg-signaling pathway and Insuline-like

receptor signaling pathways, we corroborate that the phenotype and the critical

region involved in these deficiency lines is encoded by *Sgg* (Figure10B, 10F,

Figure11C, Figure11(D′-D″).

Similarly we have compared all other overlapping deficiency lines that produced

cytoskeletal defects and identified the possible critical regions responsible for the

phenotype (Figure10). The deficiency line Df(1)Exel6242(BDSC- 7716) localized at

10D1-10D7 showed outgrowth and protruding phenotype of the wing disc

(Figure 12B). When we looked for overlapping regions, we found the deficiency

line Df(1)ED7067 (BDSC-9154) localized adjacent to it but not completely

overlapping. Therefore we generated an FRT-recombination for a smaller region

using a deficiency line Df(1)Exel9050(BDSC-7759), losing 3genes at 10D5-10D6.

The cytoskeletal defect produced was similar to that produced by the deficiency

line Df(1)Exel6242(BDSC- 7716) (Figure 12(B-C)) and therefore we can suggest

that the critical region for the epithelial defect is because of the gene *α1,6-*

fucosyltransferase (FucT6) localized to 10D5-10D5. *FucT6* is an uncharacterized

gene suggested to be involved in N-glycan Fucosylation and N-linked

glycosylation (Roos et al., 2002; Paschinger et al., 2005; Frappaolo et al., 2018).

Similarly for overlapping regions, using deficiency lines Df(1)Exel6226(BDSC-7703)

at 1E3-1F3 that generated on average only 12 clones and Df(1)ED6521(BDSC-

9281) at 1E3-1F4 that generated 46 clones on an average, we have identified the

critical region encoding a previously uncharacterized gene as

N(alpha)acetyltransferase 30A (Naa30A) localized at 1F3-1F4. Along the same

lines, for deficiency lines Df(1)Exel6234(BDSC-7708) at 4F10-5A2 that generated

on an average 42 clones and Df(1)Exel6235 (BDSC-7709) at 5A2-5A6 that

generated 64 clones, we have narrowed it down on an uncharacterized gene

CG12730 localized at 5A2 to the critical gene/locus. There are few more

overlapping deficiency lines, but since the critical gene identified further included

more than two or three genes, and due to unavailability of stocks to generate

clones, we were unable to characterize them in this present study.

Using the same approach, for II chromosome, we compared two overlapping

deficiency lines that produced opposing phenotypes to each other. The

Deficiency line Df(2L)Exel7042(BDSC-7812)(Figure13B) at 30B10-30C1 that lost 14

genes produced 36 clones while an overlapping deficiency line

Df(2L)Exel6024(BDSC-7507) (Figure13C) at 30C1-30C9 that lost 19 genes

produced clones very rarely(No clones to 1 clone). Interestingly when we

compared the wingdiscs in both cases, we found them to be slightly bigger than

the control wings (Figure13(A2,B2,C2)). Therefore we took another overlapping

deficiency line Df(2L)9040(BDSC-7815) available around 30C1-30C1. On

generating clones, it produced overgrowth and duplication of

wingdiscs(Figure13(D'-D")). Since there were two genes, *Thioredoxin(Trx2)* and

*Glcuronyl Acetyl Transferase-S (GlcAT-S)*in that region, we generated

overexpression clones for both of them. We found that overexpression of *GlcAT-S*

clones and not *Trx2* (data not shown), produced cytoskeletal defects ((Figure13E).

We also used a P-element insertion at *GlcAT-S* that functions as a hypomorph,

recombined with FRT chromosome *(FRT40A-GlcAT-S^{KG})* for clonal analysis. We

found that GlcAT-S^{KG} reproduced the overgrowth and cytoskeletal

defects(Figure13G), indicating that *GlcAT-S* localized at 30C1-30C1, is the critical

region causing the epithelial defect. *GlcAT-S* is an uncharacterized gene,

suggested to function as a co-receptor for glypicans and glycoproteins in

regulating growth (Nagarajan et al., 2015).

Concluding Remarks

Surprisingly genetic loci of chromosomes, for instance Y chromosomes of

Humans, which are restricted from recombination and spurious exchange of

genetic material have been shown to undergo multigenic deletions/duplications that leads to polymorphism in the population. Though it has been intriguing to understand the consequence of such random loss of genetic material, it has remained challenging. To address this, we have utilized the fly deficiency lines. Collections of deletion-bearing Fly stocks are powerful tool utilized for mutagenesis and gene mapping. To make them even more useful, as a pilot attempt, here we have generated a novel deficiency kit comprising small-chromosomal deletion recombined onto FRT chromosomes. This collection currently comprises most of the X (~67) and partial II chromosome (~125) FRT-recombined deficiency lines. These stocks will be submitted and made available at the Bloomington Drosophila stock centre. In this study we have utilized these FRT recombined -deletion-bearing chromosomes for mosaic analysis in III instar wing disc to screen predominantly for cytoskeletal changes implicated in epithelial morphogenesis. In addition, researchers can also utilize this approach for any of the individual deficiency lines not included in this collection and a similar work can be extended either to other tissues for clonal analysis or to understand any other cellular processes like neurogenesis, growth processes, polarity-regulation etc. This approach helps to understand how genes behave if they are deleted in a group or when small region containing genes in order is lost. We found that overlapping deficiency lines spanning the chromosome with varied number of genes lost in them, produced clones of varied number and sizes. Clonal properties (size, number and distribution) and clonal behaviour (interaction with its neighbouring cells) of mitotic clones generated by a gene loss reflects its function. Therefore, we have further categorized them into groups based on their clonal properties to have better understanding of the gene function. Further, using these overlapping deficiency lines that generate cytoskeletal defects with phenotypes, we have identified few essential loci and genes critical for maintaining tissue integrity and growth. We also understand that those deficiency lines that do not generate clones suggest that these loci

may be a crucial region encoding essential genes. As these deletion-bearing

chromosomes are equivalent to human chromosomes with microdeletions, we

compared the such regions that caused microdeletion-syndromes and other

Mendelian disorders in humans with respective cytological locations on

Drosophila chromosomes. As contiguous-gene deletion syndromes display

extremely variable clinical symptoms depending on the genes lost in the loci, the

existing problem in diagnosing patients with MD/MenD or other Contiguous-

gene deletion syndromes or multigenic deletion syndromes is the difficulty in

identifying the critical loci for disease manifestation. This study is an example to

show how data from genomic information from multigenic syndrome patients or

MD or MenD patients can be pooled to identify the critical loci implicated in

disease manifestation. By correlating the genotype-phenotype data available for

multigenic deletion syndromes and genomic information it might be possible to

provide a better prognosis of the disease development.

Experimental Procedure

1. Clonal Analysis

(i) Drosophila Stocks – Drosophila stocks and crosses were maintained on a

standard yeast/cornmeal/corn syrup/malt extract/agar medium at 25°C, uncles

indicated otherwise. Deletion-bearing stocks/ Deficiency lines of Exelixis, Inc (Exel),

nearly ~208 fly stocks covering X chromosome and second chromosome were

obtained from Bloomington Drosophila Stock Centre. Other stocks used for this

study include (5130)-pin/cyo; UAS-mCD8GFP, (5131)-hs-FLP; FRTG13, UAS-

mCD8GFP, (5134)-hs-FLP, tubP-GAL80, FRT19A; UAS-mCD8GFP, (5074)-FRT40A;

T155GAL4, UAS-FLP, (5075)-FRTG13; T155GAL4, UAS-FLP, (5192)-FRT40A, tubP-

GAL80; (5132)-hs-FLP, tubP-GAL80, FRT19A; (5140)- FRTG13, tubP-GAL80; (5138)-

tubP-GAL4; (8862)- hs-FLP; (5136)- UAS-mCD8GFP (X);

(5130)- UAS-mCD8GFP (Ⅲ); (42725)- hs-FLP, UAS-mCD8GFP; FRT40A, tubP-GAL80;

tubP-GAL4. For generating clones by conventional method, FRT19A ; T155GAL4,

UAS-FLP, FRT40A; T155GAL4, UAS-FLP and FRTG13; T155GAL4, UAS-FLP were

used.

(ii) Recombination and Generation of clones

Sequence-Mapped deletion-bearing fly stocks (Deficiency lines) for X and II

chromosomes generated by DrosDel Project and Exelixis collection (Exel) (Ryder

et al., 2004; Parks et al., 2004; Cook et al., 2012; Roote and Russell, 2012)

obtained from Bloomington Drosophila Stock Centre (BDSC) were recombined

with their corresponding FRT chromosomes (FRT19A, FRT40A and FRTG13

chromosomes) and recombined stocks were selected on neomycin (G418)

containing media. We have successfully recombined ~67 X chromosome

deficiency lines and 125 second (2L and 2R) chromosome deficiency lines with

their respective FRT chromosomes. List of Deficiency lines with the number of

genes lost in the chromosome is provided (Supplementary data S1-Sheet-1).

Further FRT-Recombined deficiency lines were subjected to mitotic clonal analysis

using both conventional and Mosaic Analysis with Repressible Cell Marker

(MARCM) techniques (Lee and Luo, 1999, Lee and Luo, 2001; Blair, 2003; Wu and

Luo, 2006) (Supplementary Figure1) in the III instar wing imaginal discs (Figure

1(A-O)).

Analysing clones by conventional method was tedious, therefore we followed

clonal analysis by MARCM method. Total number of clones and size of each

clones per wing imaginal discs were counted and recorded using ImageJ

software. Nearly 10 wing discs (n=10) were imaged and replicates of three wing

discs were randomly chosen for averaging total number of clones and size of

each clone per wing imaginal discs.

Following protocol by Wu and Luo, 2006, MARCM Stock FRT19A (hs-FLP, tubP-

GAL80, FRT19A; tubP-GAL4; UAS-mCD8GFP), MARCM Stock FRT40A (hs-FLP, UAS-

mCD8GFP; FRT40A, tubP-GAL80; tubP-GAL4) and MARCM Stock FRTG13 (hs-FLP,

UAS-mCD8GFP; FRTG13, tubP-GAL80; tubP-GAL4) were assembled using above

stocks to generate MARCM clones. Individual FRT-recombined deletion-bearing

stocks were crossed with corresponding MARCM stocks. On the 3rd day AEL, first

instar larvae were heatshocked at 37^{O}C for 1hour and allowed to recover at 25^{O}C

after that. On 6th day AEL, from each mutant type generated, third instar larvae

were dissected and subjected to immunostaining (Usha and Shashidhara, 2010;

Szuperak et al., 2011).

(iii) Immunostaining and Clonal Analysis

Wing Imaginal discs were immunologically stained using Phalloidin followed by

Alexa594 and viewed under the confocal microscope. Their images were then

analysed for clone size, distribution and number using Image J software. The

clones of each of the genes mutated were analyzed in triplicates each for both

the fly lines. The average clone sizes and clone numbers were calculated in each

case, with the most approximate result in each case (For discs with <10 clones, all

clones were considered for mean clone number estimation; For discs with >50

clones, 50 of the clones were taken into consideration for mean estimation; as

well as for discs with intermediate number of clones, suitably, either 10, 20, 30 or

40 clones were considered for best estimate of the average). A triple blind test

was also performed to corroborate each of these results.

(2) Systematic Analysis

We began the systematic analysis by identifying the genes deleted in each of the

available deficiency lines from https://bdsc.indiana.edu/. We then built an

inventory with the deleted genes arranged according to their cytological location,

ensuring that there are no repetitions, against their respective deficiency lines.

Alongside, a list of their corresponding human orthologs was generated using

the gene summaries available at www.flybase.org → Orthologs → Human

orthologs .

Apart from this, we assessed each gene for cytoskeletal implications from the

gene ontology summaries updated on their respective flybase web pages and we

have categorized based on their function. (www.flybase.org → Gene Ontology →

Molecular Function / Biological Processes / Cellular Component). Similarly, we

also scrutinized the human orthologs for any cytoskeletal implications, under the

same categories, using the database available at

https://www.ncbi.nlm.nih.gov/gene/. For those genes that were involved in

functions that fell into more than one of these categories, we took their most

predominant function into account for grouping and analyzing.

We used the Molecular Genetics Laboratory group, MGenoma®'s webpage as a

reference for a list of microdeletion syndromes. From the table present, we

enlisted the genes that were deleted in each of the syndromes. We identified

among them, those human orthologs that we previously enlisted in the inventory

generated, and also their respective fly counterparts. For each of the

microdeletion syndromes, we used the database from Orphanet to find the

critical region, or the mutational hotspot. The name of the disease was given in

the search field and the molecular genetics section of the clinical genetics review

for each disease was referred to, in order to identify the mutational hotspot or

the critical region of the gene(s) deleted.

Figure 1-Distribution and size of MARCM clones. Representative Wing discs (A-O)

with Mitotic clones (GFP positive, counterstained with DAPI in blue) generated

using Deficiency lines on X chromosome recombined to FRT19A chromosomes.

Note the variation in size and distribution of the clones in wingdiscs.

Figure 2-Number and size of clones generated by deficiency lines on X

chromosome. Classification of clones based on number of clones (A) and size of

the clones (B). Note average number (D) and size of mitotic clones (F) generated

by deficiency lines spanning the X chromosome. Correlation between number of

genes lost in deficiency lines on X chromosomes with average number (C) and

size of mitotic clones (E) generated by them.

Figure 3-Number and size of clones generated by deficiency lines on Ⅱ

chromosome. Correlation between number of genes lost in deficiency lines on Ⅱ

chromosomes with average number (A) and size of mitotic clones (C) generated

by them. Note average number (B) and size of mitotic clones (D) generated by

deficiency lines spanning the X chromosome.

Figure 4 - Wing discs with cytoskeletal breakages and disruptions. Compared to

control wingdiscs with MARCM clones (A1-A6), GFP positively marked clones by X

chromosome deficiency line BL-7703 display cell autonomous (B1-B6) and

deficiency line BL-7707 display non-cell autonomous (C1-C6) cytoskeletal defects

as shown by Actin (Phalloidin, Red). Compared to control wingdiscs with

conventional clones (D1-D6), GFP negatively marked clones by Ⅱ chromosome

deficiency line BL-7813 display cell autonomous (E1-E6) and deficiency line BL-

7815 display non-cell autonomous (F1-F6) cytoskeletal defects as shown by Actin

(Phalloidin, Red). All panels are standard confocal XY sections.

Note panels B3, C3, E3 and F3 for the changes in Actin (Red) representing

cytoskeletal breakages and disruptions in wingpouch.

Figure 5 - Wing discs with cytoskeletal defects and extrusions. Two deficiency

lines BL-7716 (A′-A″, 5C′) and BL-9348 (B′-B″, 5C″) show similar and consistent

cytoskeletal defects in clones generated either by conventional method or

MARCM strategies. Cells and tissue adjacent to the clones are extruded out of

the surface. Localized cytoskeletal defects are produced either along the

periphery of the pouch region as shown by deficiency lines BL-9348 (5B″ and 5C″)

or only in the hinge region as shown by deficiency line BL-7709(D′-D″). Note the

rounding of the clones and associated with developmental defects.

Figure 6 - Wing discs with severe cytoskeletal defects associated with outgrowths

and duplications. Compared to the control (A) rare clone produced by two

deficiency lines BL-8948 (B) and BL-8956 (E-E″). Note the outgrowth of the

wingpouch in apical (E), medial (E′) and basal (E″) XY sections and XZ section of

the medial region (F).

Figure 7 - Wing discs with severe cytoskeletal defects associated with outgrowths

and duplications. Compared to control wingdiscs with conventional clones (A) or

MARCM clones(C) clones generated by deficiency line BL-7815(B and D)

consistently show a similar phenotype of outgrowth and duplication of wing

pouch.

Figure 8 - Classification and comparison of genes implicated in cytoskeletal

functions. In the X chromosome, 181 genes out of 1103 total genes and in II

chromosome, 431 genes out of 1245 genes are involved in mediating cytoskeletal

functions. Those functions are classified into five categories and percentage of

genes involved are compared and represented.

Figure 9 - Comparative analysis of Drosophila deficiency lines with Human

Mendelian Diseases and Microdeletion Syndrome.

A detailed Flowchart (A) and a sunburst representation(B) of systematic approach

to compare genes in the Drosophila Deficiency lines of X and II chromosome with

human orthologs . Analytical representation of comparison of genes on X and II

chromosomes coding for 52 human Microdeletion syndrome (c).

Figure 10 - Identification of critical region based on clonal behaviour. For the

locus 2F6-3A8, two big deficiency lines BL-9345 (10A, E) and BL-8031(10D-E)

produced a smaller number of clones while two smaller deficiency lines BL-7705

(B, E) and BL-7706(C, E) produced high number of big sized clones (B, E). By

comparing their clonal properties and cytoskeletal information, we identify that

the critical gene is *Boi* (E). Similarly, for the loci 3A3-3B1 and 30B10-30C9 (for

clones Fig 11, Fig12) the critical gene is *Shaggy (Sgg)* and *GlcAT-S(F)* respectively.

Schematic representation of the critical region identified in this study (H).

Figure 11 - Identification of critical region based on the cytoskeletal defects. On

the X chromosome, three deficiency lines BL-9518(A-A'), BL-8031(B) and BL-

9348(C) that produced moderate number of clones were compared for their

cytoskeletal defects. For the locus 3A3-3B1, severe phenotype produced by BL-

9348(C, D'-D") showing cell-autonomous extrusion of the clones (C, D'-D", E-E')

from the wing surface is encoded by the critical gene *Shaggy (Sgg)*.

Figure 12 - Identification of critical region using non-overlapping deficiency lines.

MARCM clones for the locus 10D1-10D7, a big deficiency line BL-7716 produced

a cytoskeletal defect (B). A smaller deficiency line, BL-7759 covering a gene ·-

1,6, fucosyltransferase (FUCT6) produced a similar cytoskeletal defect (C).

Figure 13 - Identification of critical region for locus with opposing phenotypes

For the locus 30B10-30C9, a big deficiency line (BL-7812) produced moderate

number of clones (B) while an overlapping deficiency lines produced rare or very

few clones (C). Note wingdiscs produced by both deficiency lines were slightly

larger than the control(A-C). A small, overlapping deficiency line (BL-7815)

containing two genes GlcAT-S and Trx-2 for the locus 30C1-30C1 prodeuced

overgrowth and duplication of the wingdisc (D'-D"). Overexpression clones of

GlcAT-S, produced cytoskeletal defects(F) and clones generated by FRT-

recombined GlcAT-S^{KG} hypomorph produced overgrowth and cytoskeletal

defects(G).

Supplementary Figure 1 - Mitotic clones. Control Wingdisc showing mitotic

clones generated (A) by conventional method where clones are negatively

marked, and counterstained with Phalloid (Actin, RED) and (B) by Mosaic Analysis

with Repressible Cell Marker (MARCM) techniques where clones are positively

marked (GFP) and counter stained with DAPI (Nucleus, Blue).

Supplementary Figure 2 - Variation in localization and number of clones.

Representative Wingdiscs with localized clones (A, B, E), few clones (C, D),

moderate clones (E, F) and high number of clones (G, H).

Supplementary Data Table S1 - Excel sheet with list of deficiency lines used for

recombination. X chromosome (Sheet1) and II chromosomes (Sheet 2)

(Supplementary Data Table S2) – Number of clones and average size of clones

Supplementary Data Figure S3)- Mitotic clones Range from no clones to big

clones

Authors Contributions

Conceptualization: UN, SP; and Methodology: UN, SP, SS, RR, NR; Investigation:

UN, SP, TK; Formal Analysis: UN, SP, SS, RR, NR; Writing- UN, SP, TK; Writing-

Review and Editing- UN, SP, NR, TK Funding Acquisition: UN; Supervision: UN, SP.

Acknowledgments

We gratefully acknowledge Dr Matt Gibson and Dr Marios Georgiou for their help

with fly experiments in their laboratories. We thank the Bloomington stock

centers and the fly community for the generosity with reagents. We thank Prof

Satish Kumar and Dr Vasanthi Dasari for helpful advice and critical comments on

the manuscript. We thank Science and Engineering Research Board (SERB),

Department of Science and Technology (DST), Government of India, for the

grants (YSS/2014/000896) and CRG/2018/003725 to UN.

Declaration of Interests:

765 The authors declare no competing interests.

Reference

Akiyama ,T., User, SD and Gibson, MC (2018). eLife ;7: e35258. DOI:

https://doi.org/10.7554/eLife.3525

Bejsovec, A. (2018). Wingless Signaling: A Genetic Journey from Morphogenesis

to Metastasis. Genetics 208(4): 1311-1336.

Blair SS (2003). Genetic mosaic techniques for studying *Drosophila* development.

Development 130: 5065-5072; doi: 10.1242/dev.00774

Camp, D., Haitian He, HB., Li, S., Althaus, IW., Holtz, AM., Allen, BL., Charron, F.,

van Meyel, DJ. (2014). Ihog and Boi elicit Hh signaling via Ptc but do not aid

Ptc in sequestering the Hh ligand. Development 141(20): 3879--3888.

Chial, H. (2008). Somatic mosaicism and chromosomal disorders. Nature

Education 1(1):69

Clevers, H. (2006). Wnt/beta-catenin signaling in development and

disease. Cell 127(3): 469-480.

Cook, RK., Christensen, SJ., Deal, JA., Coburn, RA., Deal, ME., Gresens, JM.,

Kaufman, TC., Cook, KR. (2012). The generation of chromosomal deletions to

provide extensive coverage and subdivision of the Drosophila melanogaster

genome. Genome Biol. 13(3): R21.

Donohoe CD, Csorda´s G, Correia A, Jindra M, Klein C, Habermann B, et al. (2018).

Atf3 links loss of epithelial polarity to defects in cell differentiation and

cytoarchitecture. PLoS Genet.14 (3): e1007241. https://doi.org/10.1371/journal.

pgen.1007241.

Frappaolo, A, Sechi,S, Kumagai, T, Karimpour-Ghahnavieh, A, Tiemeyer,M,

Giansanti, MG. (2018). Modeling Congental Disorders of N-Linked

Glycoprotein Glycosylation in Drosophila melanogaster. Front. Genet. 9():436.

Franco, B., Bogdanik, L., Bobinnec, Y., Debec, A., Bockaert, J., Parmentier, M.L.,

Grau, Y. (2004). Shaggy, the homolog of glycogen synthase kinase 3, controls

neuromuscular junction growth in Drosophila. J. Neurosci. 24(29): 6573--

6577.

Fortunato, EA and Spector DH (2003). Viral induction of site‑specific chromosome

damage. Rev Med Virol. 13(1) pg21-37. https://doi.org/10.1002/rmv.368.

Germani, F., Bergantinos C and Johnston LA(2018). Mosaic Analysis in *Drosophila*

Genetics 208(2) 473-490 ;https://doi.org/10.1534/genetics.117.300256.

Giansanti MG and Fuller MT (2012). What Drosophila spermatocytes tell us about

the mechanisms underlying cytokinesis. Cytoskeleton (Hoboken). 69(11):869-

81. doi: 10.1002/cm.21063.

Gibson, MC. And Perrimon N (2005). Extrusion and Death of DPP/BMP-

Compromised Epithelial Cells in the Developing Drosophila Wing. Science,

307(5716), 1785–1789. doi:10.1126/science.1104751

Gumbiner BM. (1992). Epithelial morphogenesis. Cell. 1;69(3):385-7.

Hsia, EYC., Zhang, Y., Tran, HS., Lim, A., Chou, YH., Lan, G., Beachy, PA., Zheng, X.

(2017). Hedgehog mediated degradation of Ihog adhesion proteins

modulates cell segregation in Drosophila wing imaginal discs. Nat.

Commun. 8(1): 1275.

Knust E (2005).Development of Epithelial Cell Polarity in Drosophila Volume 7

Chapter 2 p121. Edited by (Birchmeier W and Birchmeier C, Epithelial

Morphogenesis In Development And Disease(Cell adhesion and

communication) ISBN 0-203-30376-8

Lee, T and Luo L. (1999). Mosaic analysis with a repressible cell marker for studies

of gene function in neuronal morphogenesis. Neuron. 22(3):451-61.

Lee, T, Luo L. (2001). Mosaic analysis with a repressible cell marker (MARCM) for

Drosophila neural development. Trends Neurosci. 24(5):251-4.

Olson, MV (1999). When less is more: Gene loss as an engine of evolutionary
change. Am J Hum Genet 64: 18–23.
Olson, MV and Varki A (2003) Sequencing the chimpanzee genome: Insights
into human evolution and disease. Nat Rev Genet 4: 20–28.
Paciorkowski, AR, Keppler-Noreuil K, Robinson L, et al. (2013). Deletion 16p13.11
uncovers NDE1 mutations on the non-deleted homolog and extends the
spectrum of severe microcephaly to include fetal brain disruption. Am J Med
Genet A. 161A(7):1523–1530.
Parks, AL., Cook, KR., Belvin, M., Dompe, NA., Fawcett, R., Huppert, K., etal., (2004).
Systematic generation of high-resolution deletion coverage of the
Drosophila melanogaster genome. Nat Genet. 36(3):288-92.
Paschinger, K., Straudacher, E., Stemmer, U., Fabini, G., Wilson, I.B.H. (2005).
Fucosyltransferase substrate specificity and the order of fucosylation in
invertebrates. Glycobiology 15(5): 463--474.
Price, MA and Kalderon, D. (2002). Proteolysis of the Hedgehog signaling effector
Cubitus interruptus requires phosphorylation by Glycogen Synthase Kinase 3
and Casein Kinase 1. Cell 108(6): 823--835.
Powell, K and Piddini, E (2016). Chasing how cells out compete one another.
Journal of Cell Biology. 213(3): 291–292.
Rauch, A; Schellmoser, S; Kraus, C; Dörr, HG; Trautmann, U; Altherr, MR; Pfeiffer,
RA; Reis, A (2001). First known microdeletion within the Wolf-Hirschhorn
syndrome critical region refines genotype-phenotype correlation. American
Journal of Medical Genetics. **99**(4): 338-
42. doi:10.1002/ajmg.1203. PMID 11252005.
Reiter, LT., Potocki, L., Chien, S., Gribskov, M., Bier E. (2001). A systematic analysis
of human disease-associated gene sequences in Drosophila melanogaster.
Genome Res. 11(6):1114-25.
Roos, C., Kolmer, M., Mattila, P., Renkonen, R. (2002). Composition of Drosophila
melanogaster proteome involved in fucosylated glycan metabolism. J. Biol.
Chem. 277(5): 3168--3175.

Rodríguez, AV, Didiano D, Desplan C (2012). Power tools for gene expression and
clonal analysis in Drosophila Nature Methods 9: 47–55.
Roote, J and Russell, S. (2012). Toward a complete Drosophila deficiency
kit. Genome Biol. 13(3): 149.
Ruddle, FH, Bentleyl, KL, Murtha MT, Risch N (1994). Gene loss and gain in the
evolution of the vertebrates. Development Supp. 155-161.
Ryder, E., Blows, F., Ashburner, M., Bautista-Llacer, R., Coulson, D., Drummond, J.,
Webster, J., et al., (2004). The DrosDel collection: a set of P-element
insertions for generating custom chromosomal aberrations in Drosophila
melanogaster. Genetics 167(2): 797-813.
Ryder, E., Ashburner, M., Bautista-Llacer, R., Drummond, J., Webster, J., Johnson,
G., et al., (2007). The DrosDel deletion collection: a Drosophila genomewide
chromosomal deficiency resource. Genetics. 177(1):615-29.
Schock, F and Perrimon N. (2002). Molecular mechanisms of epithelial
morphogenesis. Annu Rev Cell Dev Biol. 18:463-93. DOI:
10.1146/annurev.cellbio.18.022602.131838.
Shen, J and Dahmann C (2005). Extrusion of Cells with Inappropriate Dpp
Signaling from Drosophila Wing Disc Epithelia. Science, 307(5716), 1789–
1790. doi:10.1126/science.1104784.
Smith-Bolton, RK., Worley, MI., Kanda, H and Hariharan, IK. (2009) Regenerative
growth in Drosophila imaginal discs is regulated by Wingless and Myc. Dev.
Cell 16: 797–809.
Slavotinek, A. (2012). Microdeletion Syndromes. In eLS, (Ed.).
doi:10.1002/9780470015902.a0005549.pub2
St Johnston, D., (2002). The art and design of genetic screens: Drosophila
melanogaster. Nat. Rev. Genet. 3: 176–188.

Sustar A., Bonvin M., Schubiger M and Schubiger G (2011) *Drosophila* twin spot
clones reveal cell division dynamics in regenerating imaginal discs.
Developmental Biology (356)2: 576-587.

Szuperak M, Salah S, Meyer E, Usha N, Ikmi A and Gibson M. (2011) Feedback
modulation of *Drosophila* BMP signaling by the novel extracellular protein,
Larval Translucida Development 138, 715-724.

Nagarajan U, Pakkiriswami S and Pillai AB. (2015) Sugar tags and tumorigenesis.
Frontiers in Cell and Developmental Biology 3(69), 1-5.

Usha N and Shashidhara LS. (2010) Interaction between Ataxin-2 Binding Protein
1 and Cubitus- interruptus during wing development in *Drosophila*.
Developmental Biology 341, 389–399.

Yamamoto S, Jaiswal M, Charng WL, Gambin T, Karaca E, Mirzaa G, et al., (2014).
A drosophila genetic resource of mutants to study mechanisms underlying
human genetic diseases. Cell. 25;159(1):200-214. doi:
10.1016/j.cell.2014.09.002.

Wu, JS and Luo L. (2006). A protocol for mosaic analysis with a repressible cell
marker (MARCM) in Drosophila. Nat Protoc. 1(6):2583-9.

Zhang W, Hong M, Bae GU, Kang JS, Krauss RS. (2011) Boc modifies the
holoprosencephaly spectrum of Cdo mutant mice. Dis Model
Mech. 4(3):368-80. doi: 10.1242/dmm.005744.

Figure 1-Distribution and size of MARCM clones. Representative Wing discs (A-O) with Mitotic clones (GFP positive, counterstained with DAPI in blue) generated using Deficiency lines on X chromosome recombined to FRT19A chromosomes. Note the variation in size and distribution of the clones in wingdiscs.

Figure 2-Number and size of clones generated by deficiency lines on X chromosome.
Classification of clones based on number of clones (A) and size of the clones (B). Note average
number (D) and size of mitotic clones (F) generated by deficiency lines spanning the X
chromosome. Correlation between number of genes lost in deficiency lines on X chromosomes
with average number (C) and size of mitotic clones (E) generated by them.

911
912

Figure 3-Number and size of clones generated by deficiency lines on II chromosome. Correlation between number of genes lost in deficiency lines on II chromosomes with average number (A) and size of mitotic clones (C) generated by them. Note average number (B) and size of mitotic clones (D) generated by deficiency lines spanning the X chromosome.

914
915

Figure 4 - Wing discs with cytoskeletal breakages and disruptions. Compared to control wingdiscs with MARCM clones (A1-A6), GFP positively marked clones by X chromosome deficiency line BL-7703 display cell autonomous (B1-B6) and deficiency line BL-7707 display non-cell autonomous (C1-C6) cytoskeletal defects as shown by Actin (Phalloidin, Red). Compared to control wingdiscs with conventional clones (D1-D6), GFP negatively marked clones by II chromosome deficiency line BL-7813 display cell autonomous (E1-E6) and deficiency line BL-7815 display non-cell autonomous (F1-F6) cytoskeletal defects as shown by Actin (Phalloidin, Red). All panels are standard confocal XY sections.

Note panels B3, C3, E3 and F3 for the changes in Actin (Red) representing cytoskeletal breakages and disruptions in wingpouch.

Figure 5 - Wing discs with cytoskeletal defects and extrusions. Two deficiency lines BL-7716 (A'-A", 5C') and BL-9348 (B'-B", 5C") show similar and consistent cytoskeletal defects in clones generated either by conventional method or MARCM strategies. Cells and tissue adjacent to the clones are extruded out of the surface. Localized cytoskeletal defects are produced either along the periphery of the pouch region as shown by deficiency lines BL-9348 (5B" and 5C") or only in the hinge region as shown by deficiency line BL-7709(D'-D"). Note the rounding of the clones and associated with developmental defects.

Figure 6 - Wing discs with severe cytoskeletal defects associated with outgrowths and duplications. Compared to the control (A) rare clone produced by two deficiency lines BL-8948 (B) and BL-8956 (E-E"). Note the outgrowth of the wingpouch in apical (E), medial (E') and basal (E") XY sections and XZ section of the medial region (F).

923
924

Figure 7 - Wing discs with severe cytoskeletal defects associated with outgrowths and duplications. Compared to control wingdiscs with conventional clones (A) or MARCM clones(C) clones generated by deficiency line BL-7815(B and D) consistently show a similar phenotype of outgrowth and duplication of wing pouch.

Figure 8 - Classification and comparison of genes implicated in cytoskeletal functions. In the X chromosome, 181 genes out of 1103 total genes and in II chromosome, 431 genes out of 1245 genes are involved in mediating cytoskeletal functions. Those functions are classified into five categories and percentage of genes involved are compared and represented.

Figure 9 - Comparative analysis of Drosophila deficiency lines with Human Mendelian Diseases and Microdeletion Syndrome. A detailed Flowchart (A) and a sunburst representation(B) of systematic approach to compare genes in the Drosophila Deficiency lines of X and II chromosome with human orthologs . Analytical representation of comparison of genes on X and II chromosomes coding for 52 human Microdeletion syndrome (c).

954
955

Figure 10 - Identification of critical region based on clonal behaviour. For the locus 2F6-3A8, two big deficiency lines BL-9345 (10A, E) and BL-8031(10D-E) produced a smaller number of clones while two smaller deficiency lines BL-7705 (B, E) and BL-7706(C, E) produced high number of big sized clones (B, E). By comparing their clonal properties and cytoskeletal information, we identify that the critical gene is *Boi* (E). Similarly, for the loci 3A3-3B1 and 30B10-30C9 (for clones Fig 11, Fig12) the critical gene is *Shaggy (Sgg)* and *GlcAT-S(F)* respectively. Schematic representation of the critical region identified in this study (H).

M=MARCM Clone Phalloidin WT Genes/Clones DAPI

C=Conventional Clone Phalloidin WT Genes/Clones DAPI

Figure 11 - Identification of critical region based on the cytoskeletal defects. On the X chromosome, three deficiency lines BL-9518(A-A'), BL-8031(B) and BL-9348(C) that produced moderate number of clones were compared for their cytoskeletal defects. For the locus 3A3-3B1, severe phenotype produced by BL-9348(C, D'-D") showing cell-autonomous extrusion of the clones (C, D'-D", E-E') from the wing surface is encoded by the critical gene *Shaggy (Sgg)*.

Figure 12 - Identification of critical region using non-overlapping deficiency lines. MARCM clones for the locus 10D1-10D7, a big deficiency line BL-7716 produced a cytoskeletal defect (B). A smaller deficiency line, BL-7759 covering a gene □-1,6, *fucosyltransferase (FUCT6)* produced a similar cytoskeletal defect (C).

Figure 13 - Identification of critical region for locus with opposing phenotypes
For the locus 30B10-30C9, a big deficiency line (BL-7812) produced moderate number of clones (B) while an overlapping deficiency lines produced rare or very few clones (C). Note wingdiscs produced by both deficiency lines were slightly larger than the control(A-C). A small, overlapping deficiency line (BL-7815) containing two genes GlcAT-S and Trx-2 for the locus 30C1-30C1 prodeuced overgrowth and duplication of the wingdisc (D'-D"). Overexpression clones of GlcAT-S, produced cytoskeletal defects(F) and clones generated by FRT-recombined GlcAT-S^{KG} hypomorph produced overgrowth and cytoskeletal defects(G).

Supplementary Figure 1: FRT-Recombined deficiency lines were subjected to mitotic clonal analysis using both conventional and Mosaic Analysis with Repressible Cell Marker (MARCM) techniques

Supplementary Figure 2 - Variation in localization and number of clones. Representative Wingdiscs with localized clones (A, B, E), few clones (C, D), moderate clones (E, F) and high number of clones (G, H).

Table 1: Category based on Morphological/cytoskeletal changes

	Category	Number of Deficiency lines (DLs)	List of DLs
Widely distributed clones in wingdisc with Morphological defects	Without any Morphological changes	49	All others
	Mild Morphological changes (medium-sized, moderate to numerous clones)	5	Df(1)-Exel6226(BL-7703)
			Df(1)-Exel6233(BL-7707)
			Df(1)Exel6290(BL-7753)
			Df(1)Exel6237(BL-7711)
			Df(1)ED6989(BL-9056)
	Severe morphological changes (cytoskeletal breakages and disruptions)	2	Df(1)Exel6226(BL-7703)
			Df(1)Exel6233(BL-7707)
		2	Df(2L)Exel8022(BL-7813)
			Df(2L)Exel9040(BL-7815)
	Morphological changes with outgrowths	2	Df(1)ED6630(BL-8948)
			Df(1)ED6727(BL-8956)
		1	Df(2L)Exel6029-BL-7512
	Morphological changes with extrusions and outgrowths	2	Df(1)Exel6242(BL-7716)
			Df(1)ED6584(BL-9348)
	Morphological changes with duplications	2	Df(1)ED6630(BL-8948)
			Df(1)ED6727(BL-8956)
		1	Df(2L)Exel9040(BL-7815)
Localized clones in wingdisc with Morphological defects	Only in Notum	2	Df(1)Exel6235(BL-7709)
	Only in Hinge		Df(1)ED6584(BL-9348)

Table 2: Comparison of fly genes implicated in cytoskeletal functions with corresponding human orthologs with unknown function

DEFICIENCY LINE	FLY GENE	HUMAN GENE	OMIM	CYTOSKELETAL IMPLICATION	
				IN FLY	IN HUMANS
Df(1)ED6443	SKPA	SKP1	601434	Microtubule Cytoskeletal Organization	_
Df(1)ED6330	kirre	KIRREL3; kirre like nephrin family adhesion molecule 3	607761	Cell Adhesion, Plasma Membrane Fusion	NIL
Df(1)Exel6227	CG11448	RILP1	614092	Dynein Light Chain And Small GTPase Binding	
Df(1)Exel8196	inc	KCTD5	611285	Cytoskeletal Binding	NIL
Df(1)ED6565-9299	pcx	PCNX1; pecanex homolog 1	617655	Notch Pathway	Transmembrane Protein
Df(1)Exel6233BL7707, Df(1)ED6712-BL9169	CG10804	TROVE2; TROVE domain family member 2	600063	Transporter	NIL
Df(1)Exel6233BL7707, Df(1)ED6712-BL9169	CG16781	MARCH5; membrane associated ring-CH-type finger 5	610637	Nil	NIL
Df(1)Exel6290	NAAT1	SLC6A5; solute carrier family 6 member 5	604159	Sodium Amino Acid Symporter	NIL
Df(1)ED7067	Drak	STK17B; serine/threonine kinase 17b	604727	Epithelial Morphogenesis, Stimulates Actomyosin Contractility	NIL
Df(1)ED7217	CG32628	AGBL2; ATP/GTP binding protein like 2	617345	Mitochondrial Organization	NIL
Df(1)ED7217	NnaD	AGBL2; ATP/GTP binding protein like 2	617345	Mitochondrial Organization	NIL
Df(1)ED6716	brn	B3GNT3	605863	Border Follicle Cell Migration, Maintenance	NIL

				Of Epithelial Cell Polarity, Cell Adhesion And Morphogenesis	
Df(1)ED7165	CG12717	SENP7	612846	Axonogenesis,Dendritic Spine Morphogenesis	NIL
Df(1)ED7165	Bap60	SMARCD1	601745	Myosin Binding	NIL
Df(1)ED7170-BL8898	Sno	SBNO1	614274	Notch Regulating	NIL
Df(1)ED7261	CG5334	MKRN1; makorin ring finger protein 1	607754	GTPase, Vesicle Fusion Regulation, Intracellular Protein Transport	NIL
Df(1)ED7261	CG32595	HARBI1; harbinger transposase derived 1	NIL	Centripetal Follicle Cell Migration	NIL
Df(1)ED7344	CG8509	PTPA; protein phosphatase 2 phosphatase activator	600756	Mitotic Spindle Organization	NIL
Df(1)ED7344-9220	Efhc1.1	EFHC2; EF-hand domain containing 2	300817	Synaptic Growth, Dendritic Morphogenesis	NIL

Table 3: Deficiency lines in Drosophila and Microdeletion Syndromes in Humans

Microdeletion_Syndromes	X	Ii	Prevalence in Humans	Mutation hotspot in Humans
17q21.31 microdeletion syndrome	2	4		KAT8 regulatory NSL complex subunit 1 - KANSL1
1p36 deletion syndrome	1	2		NA
1q21.1 deletion syndrome, 1.35 Mb	0	4		GJA5, GJA8 (CHD1L, BCL9, and PRKAB2)
3q29 microdeletion syndrome	2	3		NA
Adrenal hypoplasia, Congenital	0	1	25%	NROB1, GK, DMD, IL1RAPL1
Angelmann's syndrome	1	2		MBD5
Aniridia	1	3	17-22%	PAX6
Autism	11	10		ADNP
Basal Cell Nevus syndrome	0	1	1/6-21%	PTCH1, SUFU
Beckwith-Wiedemann Syndrome	2	17		H19, IGF2, CDKN1C, KCNQ1, KCNQ1OT1
Brachydactyly mental retardation syndrome	5	4		HDAC4
Branchiootorenal syndrome	1	6		EYA1, SIX5, SIX1
Bruton agammaglobulinemia	0	1	8%	BTK
Campomelic Dysplasia	1	3	rare	SOX9
Charcot-Marie-Tooth Syndrome	7	5		GJB1
Cleidocranial dysplasia	0	7	10%	RUNX2
Cornelia De Lange syndrome	0	4		HDAC8, NIPBL, RAD21, SMC1A, SMC3
Cri-du-chat syndrome	1	2		catenin delta 2 - CTNND2 semaphorin 5A - SEMA5A

Dandy Walker syndrome	1	6		ZIC1, ZIC4 (PROBABLY)
Diaphragmatic Hernia, Congenital	1	6	rare	*NR2F, CHD, RGMA, DISP1, GATA4 and SIAT8B*
Digeorge syndrome	6	4		TBX1
Down syndrome	0	4	rare	NA
Greig Cephalopolysyndactyly syndrome	2	1	5-10%	GLI3
Holoprosencephaly	7	23		SHH, ZIC2, SIX3, TGIF1, GLI2, PTCH1
Kabuki syndrome	0	1		KDM6A, KMT2D
Leri-Weill dyschondrosteosis	0	2	rare	SHOX
Mental retardation Autosomal Dominant 51	5	0		NA
Metachromatic leukodystrophy	0	1	<1%	ARSA
Microphthalmia, Syndromic 7	1	6		COX7B, HCCS, NDUFB11
Miller-Dieker Lissencephaly syndrome	3	4		PAFAH1B1
Muscular dystrophy, Becker Type	1	4		DMD
Nail-patella syndrome	4	6	5-10%	LMX1B
Neurofibromatosis	10	13		NF1
Neuropathy hereditary, with liability to pressure palsies	0	1		PMP22
Noonan Syndrome 1	2	5		MAP2K2, NF1 (PROBABLY)
Polycystic kidney disease, infantile severe, with Tuberus Sclerosis	0	15		PKD1, TSC2 (PROBABLY)
Potocki-Lupski syndrome	5	0		FLCN, RAI1
Potocki-Shaffer syndrome	4	1		ALX4, EXT2, PHF21A (PROBABLY)
Prader-Willi Syndrome	0	1		SIM1
Retinoblastoma Rb1	1	1	8-16%	RB1

Rett's syndrome	1	0	8%	MECP2
Rieger Syndrome Type I	1	3		PITX2, FOXC1
Rubinstein Taybi Syndrome	10	0		CREBBP, EP300
Saethre Chotzen syndrome	2	1		TWIST1
Split hand/foot malformation 1	4	6	rare	TP63
Towne's Brocks syndrome	2	8	5%	SALL1
Trichorhinophalangeal syndrome	0	1	3-4%	EXT1, RAD21, TRPS1
Tuberus Sclerosis	1	0	<2%	TSC1, TSC2
William-Beuren region duplication syndrome	3	5		ELN, NCF1
Wilm's Tumor 1, Mental retardation	1	0	14%	PAX6
Wolf-Hirschhorn Syndrome	0	3		CPLX1, CTBP1, FGFRL1, LETM1, NELFA, NSD2, PIGG

Table 4: Correlation of genes on X chromosomes of Drosophila and Human associated with Mendelian diseases

Drosophila Gene	Cytolocation on Fly genome	Human gene	OMIM	Cytolocation on Human genome	Mendelian Disorder associated
CG8184	13E8-13E12	HUWE1; HECT, UBA and WWE domain containing 1, E3 ubiquitin protein ligase	300697	Xp11.22	Mental retardation, X-linked syndromic, Turner type
		SPACA5; sperm acrosome associated 5	300593	Xp11.23	
regucalcin	11A1-11A1	RGN; regucalcin	300212	Xp11.3	
Efhc1.1	13E17-13E18	EFHC2; EF-hand domain containing 2	300817	Xp11.3	
		PHEX; phosphate regulating endopeptidase homolog X-linked	300550	Xp22.11	Hypophosphatemic rickets, X-linked dominant
		PRDX4; peroxiredoxin 4	300927	Xp22.11	
CG7024	4C14-4C14	PDHA1; pyruvate dehydrogenase E1 alpha 1 subunit	300502	Xp22.12	Pyruvate dehydrogenase E1-alpha deficiency
		PDHA1; pyruvate dehydrogenase E1 alpha 1 subunit	300502	Xp22.12	Pyruvate dehydrogenase E1-alpha deficiency
CG7766	8C8-8C9	PHKA2; phosphorylase kinase regulatory subunit alpha 2	300798	Xp22.13	Glycogen storage disease, type IXa1
cib		TMSB4X	300159	Xp22.2	
Moe	8B4-8B6	MSN; moesin	309845	Xq12	Immunodeficiency 50
		OPHN1; oligophrenin 1	300127	Xq12	Mental retardation, X-linked, with cerebellar hypoplasia and distinctive facial appearance

Klp3A	3A6-3A6	KIF4A; kinesin family member 4A	300521	Xp12	Mental retardation, X-linked 100
		PHKA1; phosphorylase kinase regulatory subunit alpha 1	311870	Xq13.1	Muscle glycogenosis
CG32683	9C1-9C4	ARR3; arrestin 3	301770	Xq13.1	Myopia 26, X-linked, female-limited
		DLG3; discs large MAGUK scaffold protein 3	300189	Xq13.1	Mental retardation, X-linked 90
nod	10C7-10C8	KIF4A; kinesin family member 4A	300521	Xq13.1	Mental retardation, X-linked 100
		NAPIL2	300026	Xq13.2	
		KLHL4; kelch like family member 4	300348	Xq21.31	
		NAPIL3	300117	xq21.32	
		RAB40AL; RAB40A like	300405	Xq22.1	
CG5941	5D2-5D2	MCTS1; MCTS1, re-initiation and release factor	300587	Xq24	
		TENM1; teneurin transmembrane protein 1	300588	Xq25	
CG12796	6D4-6D4	GPR119	300513	Xq26.1	
		SAGE1; sarcoma antigen 1	300359	Xq26.3	
Clic	12C5-12C6	CLIC2; chloride intracellular channel 2	300138	Xq28	Mental retardation, X-linked, syndromic 32
CG8565	13F9-13F10	SRPK3; SRSF protein kinase 3	301002	Xq28	
scu	16F6-16F6	HSD17B10; hydroxysteroid 17-beta dehydrogenase 10	300256	Xp11.22	HSD10 mitochondrial disease
CG2694	3B3-3B3	GCNA; germ cell nuclear acidic peptidase	300369	Xq13.1	
		ATP7A; ATPase copper transporting alpha	300011	Xq21.1	Menkes disease, Occipital horn syndrome, Spinal muscular atrophy, distal, X-linked 3

		OCRL; OCRL, inositol polyphosphate-5-phosphatase	300535	Xq26.1	Dent disease 2, Lowe syndrome
		F8A1; coagulation factor VIII associated 1	305423	Xq28	