1 **Invivo and systematic analysis of random multigenic deletions associated with**
2 **human diseases during epithelial morphogenesis using Drosophila**
3

4 **Usha Nagarajan*[1], Shanmugasundaram Pakkiriswami*[2] ,Sandiya Srinivasan[3^], Niveditha**
5 **Ramkumar[3^], Rajeswari Rajaraman[3^], Swaminathan Sethuraman[1], Kumarasamy Thangaraj *[3]**
6
[1]Central University of Haryana, Department of Biochemistry, Mahendragarh, Haryana, 123029
India
[2]Department of Biochemistry and Molecular Biology, Dalhousie University, Dalhousie Medicine
New Brunswick, Saint John, NB, E2L4L5,Canada,
[3] School of Chemical and Biotechnology (SCBT), SASTRA University, Thanjavur, 613401, India.
[4]Centre for Cellular and Molecular Biology (CCMB), Hyderabad, 500 007, India.
13
14 Correspondence to: usha@cuh.ac.in sn374151@sdal.ca, thangs@ccmb.res.in

15
Abstract

Random loss of multigenic loci on chromosomes, a crucial drive for evolution, occurs
frequently in all living organisms. Analysis of such chromosomal disruption and
understanding the consequences of their impact on the growth and development of
multicellular organisms is challenging. In this report, we have addressed this issue
using invivo mosaic analysis of deficiency lines in Drosophila. Genes on fly
deficiency lines were compared with human orthologs for their implications in
disease development during cytoskeletal processes and epithelial morphogenesis.
The cytoskeletal phenotypes from the fly has been utilized to predict the function of
human orthologs. In addition, as these Drosophila deficiency lines are equivalent to
human microdeletions, based on the clonal behaviour and phenotypes generated, a
systematic analysis has been carried out to establish the critical loci that correspond
to Microdeletion Syndromes and Mendelian Disorders in humans. Further we have
drawn the synteny that exists between these chromosomes and have identified
critical region corresponding to defects. A few potential candidates that might have
an implication in epithelial morphogenesis are also identified.

32
33
34
35
Introduction
Contiguous-gene deletion syndromes or multigenic deletion syndromes display
extremely variable clinical symptoms depending on the genes lost in the loci (Rauch
et al., 2001; Slavotinek, 2012). A great deal of evidence shows that random loss of
genes or spontaneous changes on chromosomes (like virus-mediated mutagenesis
in humans causing chromatid breakages, translocations, duplications,
deficiencies/deletions leads to developmental anomalies and growth defects
(Fortunato and Spector, 2003). "Less-is-more" hypothesis insists that gene loss is an
engine for evolutionary change and disease development (Olson, 1999; Olson and
Varki, 2003). Despite the fact that the adaptive role of gene loss has formed the
basis for evolution, impact of their adaptive biological functions on individual integrity
and stability remains elusive. Just as how loss and gain of Homeotic genes and their
clusters prove fundamental for the evolution of body shape and size (Ruddle et al.,

1994), the loss and gain of genes implicated in epithelial morphogenesis also
influence the integrity of body size and shape (Schock and Perrimon, 2002). In a
multicellular organism, epithelial morphogenesis is an inevitable process in which
epithelia, tissues that line the organ and body surfaces, determine organ-to-body
shape and size (Gumbiner BM.1992; Schock and Perrimon, 2002). Of all the
factors implicated in this process, cytoskeletal components that balance rigidity and
flexibility of the cell, play a critical role in maintaining organ size and growth of the
organism (Knust, 2005; Gibson and Perrimon, 2005; Shen and Dahmann, 2005;
Donohoe et al., 2018). Defects during this process lead to loss of tissue integrity;
development of genetic disorders and diseases (Olson and Varki, 2003; Knust, 2005
in Birchmeier and Birchmeier, 2005).
60
Compared to loss of individual essential genes and frameshift mutations, gains and
losses of multiple (more than one) genes bring about greater dramatic genetic
repercussions. Random loss of contiguous (loss of genes in the same order) genes
would disrupt the normal development, affecting growth and patterning of the tissue,
which otherwise occur concurrently. Loss of such contiguous genes that may occur
due to any kind of genetic aberrations might regulate both cell-autonomous and non-
autonomous functions of the tissue giving rise to heterogeneity in cell population
(mosaicism). It is intriguing to understand the influence of such sudden, random loss
of genes causing mosaicism and to compare their effect with other organisms.
70
Genomic studies suggest that genetic mosaicism is a very common phenomenon
during epithelial morphogenesis. Although such lesions to the genome are random
and generally asymptomatic in initial stages, affected individuals subsequently
develop varying levels of metabolic disorders and disabilities (Chial, 2008). Also,
since such incidences occur sporadically, chances of undertaking systematic
analysis is very low or may not be feasible. In the past, rigorous systematic analysis
of genes (Reiter et al., 2001) and genetic screening (St Johnson, 2002; Yamamoto
et al., 2014) have been undertaken to understand genes implicated in disease
development and their functions. However, how these individuals subjected to such
sudden and random loss of genes manage without their gene function and what
impact does it have on the individual is not clear. In multigenic deletions, the
synergistic and antagonistic effect of genes in maintaining homeostasis or disease
development and the alternate pathways that operate to compensate the loss is not
known yet. Although it is intriguing to understand the consequence of random loss of
multiple genes that leads to defects and genetic disorders, it is extremely
challenging.
87
We hypothesized that those limitations can be overcome and unanswered questions
can be addressed by using the available *Drosophila* deficiency line collections, one
of the unique genetic tools well-utilized extensively for genetic screening, mapping
and characterization processes. Two types of molecularly defined deficiency lines,
whose details including ends of the genes lost as well as cytological location on the

chromosome are known, are available (Ryder et al., 2004; Parks et al., 2004; Cook
et al., 2012; Roote and Russell, 2012). In addition, as chromosomes of these
deletion-bearing fly stocks resemble those chromosomes with microdeletion or
chromosomal deletions of contiguous genes (loss of genes in the same order) in
other organisms, the collection has been utilized for systematic and comparative
analysis to establish the critical loci that correspond to Mendelian Disorders and
Microdeletion Syndromes in humans.
100
Based on this logic, in this work, we set out a pilot project by subjecting these
deficiency lines for mitotic clonal analyses in *Drosophila* wing imaginal discs to
identify and understand their role in cytoskeletal organization during epithelial
morphogenesis. Clonal analysis technique allows to analyse behaviour of two
different population of cells in a tissue (Lee and Luo, 2001; Wu and Luo, 2006;
Smith-Bolton et al., 2009; Rodriguez et al., 2012; Powell and Piddini, 2016). Here we
have utilized this property to understand the behaviour of randomly eliminated
genes, thereby identify the critical gene in that locus implicated in epithelial
morphogenesis. During this process of clonal analysis, we have also followed the
cytoskeletal defects produced by these deficiency lines. We also undertook a
systematic functional analysis of the genes in the fly deficiency lines, compared them
with human orthologs and analysed their implications in cytoskeletal processes and
epithelial morphogenesis. Further we have drawn the synteny that exist between
these chromosomes and identified critical region of epithelial defects. Here, in this
report, few potential candidates that might have an implication in epithelial
morphogenesis are discussed.
117
Results and Discussion
(i) Recombination of Deletion-bearing chromosomes with FRT chromosomes
and generation of mitotic clones
To address the behavioural property of the tissue subjected to multigenic deletions
we generated mitotic clones using the available deficiency lines in Drosophila wing
imaginal discs. We recombined X and II chromosome deficiency lines with their
respective FRT chromosomes. We have subjected ~67 FRT-recombined X
chromosome deficiency lines and 125 FRT-recombined second (2L and 2R)
chromosome deficiency lines for clonal analysis in III instar larval wing imaginal
discs. These deficiency lines produced clones of varied number and sizes (Figure
1(A-O); Supplementary S1, Sheet2 – Sheet5). For instance, deficiency line
Df(1)Ex6237(BL-7711) with 21 genes deleted produced 103 clones, highest average
number of clones (Figure 1N) while deficiency line Df(1)ED6443(BL-9053) with 45
genes deleted generated 0 to 1 (average =0.3), lowest number of clones. Deficiency
line Df(1)Exel6230(BL-7705) with 9 genes deleted developed the biggest clones of
0.6 Sq. pixels, while deficiency line Df(1)Exel6221(BL-7699) with 14 genes deleted
produced smallest clones of 0.04 Sq. pixels (Figure 1J), Supplementary S1- Sheet2-
3). Some of the deficiency lines did not generate clones or generated few clones
(<10) per wing disc. Interestingly few of the deficiency lines showed clones

specifically localized in wing discs, especially only in pouch region (Figure1J) or only
in notum of the wing disc (Figure 1F, Supplementary 2(A-B).
139
In order to understand the clonal properties of the mitotic clones generated by these
deficiency lines, we have categorized the deficiency lines into very few (rare or less
than or upto 10 number of clones), few (upto an average of 10- 30 clones), moderate
(upto 60 clones), and numerous (> 60 clones) clonal types and based on the size of
clones, deficiency lines were grouped into large, medium and small types. On X
chromosome, 37% of the deficiency lines generated rare/very few clones and 27%
generated few clones. Around 21% and 15% of deficiency lines were categorized
into moderate and numerous categories respectively (Figure 2A, Supplementary S1-
Sheet2). In addition, based on the average size of the clones generated per wing
disc, we have categorized the deficiency lines from small through medium to large
(Figure 2B, 2(E-F), Supplementary Data S1 Sheet3).
151
To address the link between the number of genes lost in a deficiency line and their
corresponding clonal property, we have generated correlation graphs between them.
From the correlation graph, we find that depending on the function and importance of
genes deleted in a deficiency line, their clonal properties and behaviour varied
accordingly (Figure 2C, Supplementary S1- Sheet2). For example, (i) on X
chromosome there are four deficiency lines that have very less number of genes lost
on them but generated either rare clones [Deficiency line (Df(1)Exel6244(BL-7717)
which lost only 7 genes did not generate clones at all while Df(1)Exel6221(BL-
7699)which lost 14 genes generated very less to 1 clone in a wing disc (Figure1J)] or
very few clones [Df(1)Ex6241(BL-7715) lost 6 genes but generated on an average of
only 4 clones (Supplementary Figure2C) while Df(1)ED6579(BL-9518)(70.1) lost 10
genes but generated on an average of 10 clones per wingdisc (Supplementary
Figure2D)]. (ii) Similarly, there are only four deficiency lines in the moderate and
numerous categories with more than 20 genes lost in them. Deficiency line
Df(1)ED7170(BL-8898) lost 58 genes made on an average 35 clones per
wingdisc(Figure1K) and Df(1)ED6521(BL-9281) lost 25 genes generated on an
average 46 clones per wingdisc (Figure(1L). Similarly, in the numerous category,
Df(1)ExelED6989(BL-9056)(76-1) that lost 32 genes made on average of 65 clones
per wingdisc (Supplementary Figure 2G). Of all these, notable ones are
Df(1)Exel6237(BL-7711) that lost 21 genes that made 103 clones, the highest
average number of clones per wingdisc (Figure1N) and Df(1)Exel6223(BL-7770) that
had lost only 3 genes but made 77 clones on an average in a wingdisc
(Supplementary Figure 2H). This deficiency line had the least number of genes lost
with the fourth highest number of clones (Supplementary S1- Sheet2). There are
nearly 8 deficiency lines on the X chromosome and 26 deficiency lines on the II
chromosome that did not generated clones (Figure 2A and 2C, Figure 3A
Supplementary S1 Sheet 2 and 4).
179

Similarly, we have performed clonal analysis on deficiency lines on II chromosome
and categorized them based on their number, size and distribution (Figure 3(A-D)
Supplementary Data S1 Sheet4-5). Nearly 27% of the deficiency lines were in rare/
very few category while 23% of them were in few category. Around 31% and 19% of
deficiency lines were categorized into moderate and numerous categories
respectively. As in case of the X chromosome, the number of clones generated by
corresponding deficiency line depended on the gene type and function. The
deficiency line Df(2L)Exel6024, that is deficient in 19 genes, makes the smallest
clones that have an average size of 0.6 Sq. pixels. The largest clones, with a mean
size of 63, are made by the Deficiency line Df(2L)Exel9040, that lacks only 2 genes.
The line Df(2L)7042, that has lost 14 genes, makes on average, clones sized 15 Sq.
pixels while the line Df(2R)Exel6065, that has lost about 40 genes, makes but
relatively small clones of average size 8 Sq. pixels (Figure 3(A-D) Supplementary
Data S1 Sheet4-5). These findings reinforce only the hypothesis further that the
clonal phenotype exhibited by the animals under study depends on the function of
the genes lost and their importance to the development underway. We have utilized
these clonal properties for systematic and comparative analyses discussed in next
sections to understand their implications on human diseases.
198
2. Analysis of cytoskeletal changes associated with Drosophila deficiency
lines and functional analysis of their Human orthologs
To understand the role of genes in deficiency lines during epithelial morphogenesis,
we have analysed the cytoskeletal changes mediated by generating mitotic clones in
wing imaginal discs. Of the 66 deficiency lines on X chromosomes subjected to
clonal analysis, nearly 12 deficiency lines showed cytoskeletal changes in the wing
imaginal discs (Table 1). Although we generated clones for both X and II
chromosomes, we analysed the cytoskeletal defects for entire X chromosomes
deficiencies and part of the II chromosome deficiency lines that showed visible
defects either in larval or adult stages.
209
Based on the type and the degree of cytoskeletal changes, we found that clones
were either widely-distributed throughout the wingdisc or localized in wingdiscs.
Accordingly, we have classified them into six categories, namely, with (i)-no, (ii)-mild,
(iii)-severe morphological changes and with (iv)-outgrowths, (v)-extrusions and (vi)-
duplications (Table 1). Nearly 49 deficiency lines did not show any cytoskeletal
changes. Of the remaining deficiency lines, approximately 5 deficiency lines (Df(1)-
Exel6226(BL-7703), Df(1)-Exel6233(BL-7707), Df(1)Exel6290(BL-7753),
Df(1)Exel6237(BL-7711), Df(1)ED6989(BL-9056) showed very mild morphological
changes. These deficiency lines generated medium-sized, moderate to numerous
clones in the wing discs. In the Df(1)Exel6226(BL-7703), we could observe cell-
autonomous cytoskeletal defects (Figure 4B(1-4)) while the deficiency line
Df(1)Exel6233(BL-7707), displayed non-cell autonomous cytoskeletal defects
involving cytoskeletal breakages and disruptions (Figure 4C(1-4)). Similarly, for
second chromosomes, we generated clones using conventional method. Compared

to the control wingdiscs (Figure 4D(1-4)), two deficiency lines Df(2L)Exel8022(BL-
7813) (Figure 4E(1-4)) and Df(2L)9040(BL-7815) (Figure 4F(1-4)) showed severe
cytoskeletal defects. Two deficiency lines (Df(1)Exel6242(BL-7716) Figure 5(A'-A",
Figure 5C'))and Df(1)ED6584(BL-9348) (Figure 5(B'-B", Figure 5C") showed
cytoskeletal defects and extrusion of the clones associated with outgrowths.
Consistently, we find both deficiency lines show similar phenotypes by both direct
mosaic as well as MARCM techniques. Two deficiency lines showed localized
clones: Df(1)ED6584(BL-9348) produced clones near the periphery of pouch region
associated with the extrusion of rounded clonal cells (Figure 5B" and 5C"); while the
deficiency line Df(1)Exel6235(BL-7709) produced clones only in the notum region
displaying non-cell autonomous cytoskeletal rounding near the clones, (Figure 5(D'-
D")). Similarly, three deficiency lines showed cytoskeletal defects with outgrowths
and duplication phenotypes. Of them, two from X chromosome, (Df(1)ED6630(BL-
8948) (Figure 6(B-D) and Df(1)ED6727(BL-8956) (Figure 6(E-F)rarely produced
clones, however, it showed severe morphological changes, including outgrowth and
duplication in the wingdiscs. One of the II chromosome deficiency lines,
Df(2L)9040(BL-7815), displayed outgrowth and pouch duplication phenotype in the
wingdisc (Figure7(A-D).
242
We have also undertaken a systematic functional analysis of the genes in the fly
deficiency lines, compared them with human orthologs and analysed their
implications in cytoskeletal processes and epithelial morphogenesis (Supplementary
Data S2). In the X chromosome of the fly, out of 1103 total genes, there are 181
genes that are involved in such functions. Among the 181 genes, one-third includes
transmembrane proteins, another one-third is involved in cytoskeleton binding and
organization, and others are involved in intracellular vesicular transport, cell polarity
establishment and maintenance of epithelial morphogenesis. Similarly, in the II
chromosome there are 431 genes out of 1245 which are implicated in cytoskeletal
functions. We have compared these genes on X and II chromosomes that have
cytoskeletal implication with their respective Human orthologs (Figure 8). In humans,
200 of the total orthologs are involved in cytoskeletal functions. In some instances,
the functions of the orthologs of cytoskeletally implicated genes are known while in
others it is known atleast in the fly counterparts. About 122 genes and their orthologs
both have cytoskeletal functions. In numerous cases, the human orthologs of such
genes have not been functionally characterized as yet. For 18 genes on X
chromosome, cytoskeletal functions of fly orthologs are known but unknown in
human orthologs (Table 2).
261
Nearly 11 genes have both fly and their corresponding human orthologs with similar
cytoskeletal functions. For example, Drosophila *kirre* of Df(1)ED6330 (Fig 1F) and its
human ortholog *KIRREL3* is involved in cell adhesion and plasma membrane fusion
activities and the fly gene *SKPA* of Df(1)ED6443 and its human ortholog *SKP1* is
implicated in microtubule cytoskeletal organization. Similarly , gene linked to Notch
regulation, Drosophila *sno* of Df(1)ED7170 (Fig 1L) and its human ortholog *SBNO1*

and gene connected to synaptic growth, fly *Efhc1.1* of Df(1)ED7344 (Fig 1I) and its
human ortholog *EFHC2* are implicated in cytoskeletal morphogenesis. Few of the fly
orthologs are uncharacterized while cytoskeletal functions of their human orthologs
are known. For example, fly orthologs *CG10804* and *CG16181* of human genes
TROVE2 and *MARCH5* respectively are uncharacterized genes. These two genes
are covered by the deficiency line Df(1)Exel6233. In the same section above we
have shown that Df(1)Exel6233 produced mild cytoskeletal defects (Fig 4). Therefore
it is possible that the genes *CG10804* and *CG16181* may be implicated in
cytoskeletal functions. Clonal properties of 18 genes on respective deficiency lines
are undertaken in this study (Supplementary data S2(Sheet1-2); Table 2). The
cytoskeletal phenotypes from the fly can thus be utilized to predict the etymology of
many rare, unexplored human diseases.
280
3. A Systematic approach to compare genes on Drosophila deficiency lines
with Human Mendelian Diseases (MenDs) and Microdeletion Syndrome (MDs)
to draw synteny between them
We undertook a systematic approach for both X and II chromosomes to compare the
Drosophila genes on deficiency lines with their corresponding Human Orthologs. We
have classified them into two categories based on whether or not they have available
Human orthologs. There are 600 genes on X chromosome and 1090 genes on II
chromosomes with Human orthologs. These groups with Human orthologs were
looked up for their association with disease enlisted in Online Mendelian Inheritance
in Man (OMIM) database. There are nearly 189 genes on X chromosome and 509
genes on II chromosome with OMIM disease association. Based on this criterion, we
further categorized the diseases into Mendelian disease and Microdeletion
Syndrome. There are nearly 119 genes on X chromosome and 227 on II
chromosome for Microdeletion syndrome (Figure 9(A-B), Supplementary data S3
(Sheet1-2)) and 70 genes on X chromosome and 202 genes on II chromosome
associated with Mendelian Disease (Figure 9(A-B), Supplementary data S3(Sheet
7)).
298
Further we have classified and analysed the data in two ways. First, for any locus on
the human chromosome, we have identified all possible MDs or MenDs pertaining to
it. Therefore, any lesion on that particular locus can lead to any of these disorders
listed Supplementary data S3 (Sheet1-2), (Sheet 7)). Some of the MDs are coded by
more than one locus on either of these chromosomes while some of them are either
present on X or II chromosome.   We have pooled and compared the number of
genes on both X and II chromosome that represent the corresponding MD syndrome
to generate a comprehensive comparative chart between them (Table 3, Figure9C).
307
Second, we found many human Microdeletion and Mendelian syndromes to mirror
the genetic deletions observed in the fly deficiency lines under study. We could thus
make comparison between them to identify etymologies of rare variants of human
genetic disorders. We have analysed the MDs by comparing the order of genes as

they are arranged on the human chromosomes (synteny). In this study, we have
found that there are number of genes in the order on Drosophila chromosome to be
represented in the same order on Human chromosome leading to either
Microdeletion syndromes (Supplementary data S3(Sheet 5-6)) or Mendelian disease
(Supplementary data S3(Sheet 7)). Interestingly there are also 16 Mendelian
Disease genes on X chromosome of Drosophila that correspond to genes on Human
X chromosome (Table 4 or Supplementary Data S3(Sheet 8)).
319
Until now, researchers and clinicians were finding it difficult to pin down the
respective genes that correspond to the disease manifestation. However, from this
comprehensive analysis, it is possible to identify the specific gene that lead to MDs
and MenDs. For example, we found that one of the MDs, Holoproscencephaly, is
caused by loss of several genes. Nearly there are 7 genes on X and 23 genes on II
chromosome that can lead to the manifestation of the syndrome (Table 3,
Supplementary Data S3 (Sheet1-2). Next highest are the MDs like Autism (11 genes
on X and 10 genes on II chromosome) and Neurofibromatosis (10 on X chromosome
and 13 on II chromosome). Comparative loci analysis reveals that in the locus on
human 2q37.3, either Autism or Holoprosencephaly can occur. By comparing with
the Drosophila genes, it is possible to point out the genes and locus on the
chromosome. While loss of *per* (at 3B1-3B2) can lead to Autism, loss of *CG10353*
(at 10F2-10F4) can lead to Holoproscencephaly Supplementary Data S3 (Sheet1,
Sheet3). Similarly, lesions at loci 11p11.2 or 16p11.2 can lead to DiGeorge
Syndrome/Velocardiofacial syndrome and Holoproscencephaly. From this study we
find that loss of either *SPACA3* at 11p11.2 or *FAM57B* at 16p11.2 can lead to
DiGeorge Syndrome, while loss of either *PTPRI* at 11p11.2 or loss of *PHKG2* at
16p11.2 can lead to Holoprosencephaly (Supplementary data S3(Sheet1, Sheet3)).
338
Similarly, the locus 5A12-5D1 encodes for several essential genes and therefore any
deficiency in this region may potentially give rise to MDs like Beckwith Wiedemann
Syndrome (BWS), Bracydactyly-Mental Retardation syndrome (Bdmr1),
(Branchiorenal syndrome, Bor1 and Campomelic Dysplasia). Deficiency line
Df(1)ED6802(BL-8949), pertaining to the locus 5C2-5C2, that encodes for Carboxy
peptidases A3 and B1 (CPA3) may lead to BWS as well as for Bdmr1. This also
suggests that Human orthologs of these genes could potentially be involved in
human X-linked MDs) (Supplementary data S3(Sheet1, Sheet3)).
347
Interestingly, we have found that there are MDs that are caused by loss of genes
only on X chromosome and not on II chromosome. For example, Rubinstein Taybi
Syndrome is caused by 10 loci, all located only on X chromosome and not on II
chromosome. Similarly, Potocki-Lupski Syndrome is caused by 5 loci on X
chromosome and not on II chromosome. Rett's syndrome and Tuberus sclerosis are
MDs caused by microdeletions in *MECP2* gene and *TSC1/2* respectively both with
single locus each on X chromosome. (Table3, Supplementary data S3(Sheet1,
Sheet3)). Corresponding deficiency lines analysed here are Df(1)ED6630-8948

(Figure 1E)   and Df(1)ED7170-9058(Figure 1K) that has the genes *CG2680* and
HDAC4 respectively representing the microdeletions causing Rett's syndrome and
Tuberus sclerosis.
359
From this analysis, in addition to the point that a particular MDS can be represented
by more than one locus, we also understand that, multiple MDs that arise in a
particular locus depend on the genes deleted and the order of contiguous genes
deleted. In addition, comprehensive and comparative analysis between Human and
Drosophila loci have led to identification of putative critical region in human
chromosome that could give rise to various MDs. This is addressed in detail in the
following section.
367
4. Identification of critical region or the gene implicated in epithelial defect
Although multiple genes are lost in the manifestation of MDs or Mendelian diseases,
there exists a critical region involving either an essential gene or a small genomic
region in that locus which is central in causing the disease. Since the molecular ends
of the deficiency lines used in the study were precisely defined beforehand, we
compared this information with the clonal phenotype or behaviour generated to
identify the critical region in that locus, necessary for the maintenance of tissue
integrity. We took two approaches while comparing the overlapping deficiency lines,
one based on the number and size of the clones generated with respect to the
number of genes eliminated (Supplementary S1 Sheet(2 and 4); and the other based
on the cytoskeletal defects displayed by them (Supplementary S3). For the former
approach, using the deficiency lines on X chromosome, we looked into the category
with numerous clones (>60 clones per wingdisc) generated by a smaller number of
genes lost in them. Of the 10 deficiency lines in this category, deficiency line
Df(1)Exel6237 (BDSC-7711)at 5C2-5C6 with 21 genes lost including *Usp30*
generated 103 clones, the highest number of clones of all lines*.* Followed by that,
Df(1)Exel6230(BDSC-7705) at 3A2-3A4 with 9 genes lost generated biggest
clones(with average size of 0.6 Sq.pixels) and the second highest number of clones
per wing disc (on an average 91 clones). Under this category, we also found an
overlapping deficiency line Df(1)Exel6231(BDSC-7706) for the same region at 3A2-
3A3, that lost 6 genes but produced 65 clones (6[th] highest number of clones) with
average size of 0.4 Sq.pixels(one of the biggest clones). For the same locus, we also
found two bigger deficiency lines Df(1)ED11354 (BDSC-9345) at 2F6-3A4, with 15
genes lost, but generating only 13 clones (with average size of 0.05 Sq.pixels); and
Df(1)ED411(BDSC-8031) at 3A3-3A8 with 18 genes lost, but produced only 3 clones
on an average(with average of size 0.2 Sq.pixels). Of the four deficiency lines
spanning this locus, (Df(1)Exel6230(BDSC-7705) and Df(1)Exel6231(BDSC-7706)
show similar phenotype with large (0.6 and 0.4 Sq.pixels)and numerous clones (91
and 65) respectively. By comparing these three deficiency lines
(Df(1)Exel6230(BDSC-7705), Df(1)Exel6231(BDSC-7706) with Df(1)ED411(BDSC-
8031), we found that 3A2-3A3, that contains the gene *Brother of ihog (Boi)* is the
critical region/gene responsible for producing high number of bigger clones

(Figure10A, 10E and 10H ; Supplementary Data S4 (Sheet1). It is understood that
the third biggest deficiency line Df(1)ED11354 (BDSC-9345) might include other
genes like *Raf,* another regulator of growth. *Boi* is suggested to be a negative
regulator of growth, that functions as a co-receptor for Patched(Ptc) in regulating
Hedgehog(Hh)-signaling mediated growth. Therefore, highest number of clones
generated is due to the elimination of *Boi*, the negative regulator of growth, in that
locus. Interestingly, loss of the gene *Cell adhesion associated, Oncogene regulated*
(CDON), human ortholog of Boi, localized at 11q24.7 is suggested to cause MD
Holoprosencephaly in humans.
409
As a second approach, on both X and II chromosomes, we looked into those
deficiency lines which generated cytoskeletal defects and wingdisc phenotypes
associated with them. On X chromosome, the deficiency line Df(1)ED6584(BDSC-
9348) (Figure11C) at 3A8-3B1 lost only 3 genes, but generated moderate number of
clone (average number of 36 and average size of 0.12 Sq.pixels). We compared it
with available overlapping deficiency lines Df(1)ED6679 (BDSC-9518) (Figure11(A'-
A")) at 3A6-3A8 with 10 genes lost, that produced only 8 clones (with average size
of 0.08 Sq.pixels); and Df(1)ED411(BDSC-8031) (Figure11B) at 3A3-3A8 with 18
genes lost produced only 3 clones on an average(with average of size 0.2 Sq.pixels).
The clonal analysis of the deficiency line Df(1)ED6679 (BDSC-9518) by both
conventional method Figure 11 A-A') and MARCM technique (Figure11(D'-D"))
showed several cytoskeletal defects in the hinge region, with small rounded cells
around the periphery of the hinge. However, the clonal analysis of
Df(1)ED6584(BDSC-9348) showed cell-autonomous extrusion of the clones from the
wing surface (Figure11(E'-E")). By comparing the overlapping region between them
and the similar phenotype produced by Df(1)ED6679 (BDSC-9518), we could identify
the critical region is coded by the gene *Shaggy(Sgg).* As *Sgg* is a key component of
β-catenin destruction complex so as to function as a negative regulator of Wg-
signaling pathway and Insuline-like receptor signaling pathways, we corroborate that
the phenotype and the critical region involved in these deficiency lines is encoded by
Sgg (Figure10B, 10F, Figure11C, Figure11(D'-D").
431
Similarly we have compared all other overlapping deficiency lines that produced
cytoskeletal defects and identified the possible critical regions responsible for the
phenotype (Figure10). The deficiency line Df(1)Exel6242(BDSC- 7716) localized at
10D1-10D7 showed outgrowth and protruding phenotype of the wing disc (Figure
12B). When we looked for overlapping regions, we found the deficiency line
Df(1)ED7067 (BDSC-9154) localized adjacent to it but not completely overlapping.
Therefore we generated an FRT-recombination for a smaller region using a
deficiency line Df(1)Exel9050(BDSC-7759), losing 3genes at 10D5-10D6. The
cytoskeletal defect produced was similar to that produced by the deficiency line
Df(1)Exel6242(BDSC- 7716) (Figure 12(B-C)) and therefore we can suggest that the
critical region for the epithelial defect is because of the gene *α1,6-fucosyltransferase*

(FucT6) localized to 10D5-10D5. *FucT6* is an uncharacterized gene suggested to be
involved in N-glycan Fucosylation and N-linked glycosylation (Roos et al., 2002;
Paschinger et al., 2005; Frappaolo et al., 2018). Similarly for overlapping regions,
using deficiency lines Df(1)Exel6226(BDSC-7703) at 1E3-1F3 that generated on
average only 12 clones and Df(1)ED6521(BDSC-9281) at 1E3-1F4 that generated
46 clones on an average, we have identified the critical region encoding a previously
uncharacterized gene as *N(alpha)acetyltransferase 30A (Naa30A)* localized at 1F3-
1F4. Along the same lines, for deficiency lines Df(1)Exel6234(BDSC-7708) at 4F10-
5A2 that generated on an average 42 clones and Df(1)Exel6235 (BDSC-7709) at
5A2-5A6 that generated 64 clones, we have narrowed it down on an uncharacterized
gene *CG12730* localized at 5A2 to the critical gene/locus. There are few more
overlapping deficiency lines, but since the critical gene identified further included
more than two or three genes, and due to unavailability of stocks to generate clones,
we were unable to characterize them in this present study.
Using the same approach, for II chromosome, we compared two overlapping
deficiency lines that produced opposing phenotypes to each other. The Deficiency
line Df(2L)Exel7042(BDSC-7812)(Figure13B) at 30B10-30C1 that lost 14 genes
produced 36 clones while an overlapping deficiency line Df(2L)Exel6024(BDSC-
7507) (Figure13C) at 30C1-30C9 that lost 19 genes produced clones very rarely(No
clones to 1 clone). Interestingly when we compared the wingdiscs in both cases, we
found them to be slightly bigger than the control wings (Figure13(A2,B2,C2)).
Therefore we took another overlapping deficiency line Df(2L)9040(BDSC-7815)
available around 30C1-30C1. On generating clones, it produced overgrowth and
duplication of wingdiscs(Figure13(D'-D")). Since there were two genes,
Thioredoxin(Trx2) and *Glcuronyl Acetyl Transferase-S (GlcAT-S)*in that region, we
generated overexpression clones for both of them. We found that overexpression of
GlcAT-S clones and not *Trx2* (data not shown), produced cytoskeletal defects
((Figure13E). We also used a P-element insertion at *GlcAT-S* that functions as a
hypomorph, recombined with FRT chromosome *(FRT40A-GlcAT-S^{KG})* for clonal
analysis. We found that GlcAT-S^{KG} reproduced the overgrowth and cytoskeletal
defects(Figure13G), indicating that *GlcAT-S* localized at 30C1-30C1, is the critical
region causing the epithelial defect. *GlcAT-S*is an uncharacterized gene, suggested
to function as a co-receptor for glypicans and glycoproteins in regulating growth
(Nagarajan et al., 2015).
477
Concluding Remarks
Surprisingly genetic loci of chromosomes, for instance Y chromosomes of Humans,
which are restricted from recombination and spurious exchange of genetic material
have been shown to undergo multigenic deletions/duplications that leads to
polymorphism in the population. Though it has been intriguing to understand the
consequence of such random loss of genetic material, it has remained challenging.
To address this, we have utilized the fly deficiency lines. Collections of deletion-
bearing Fly stocks are powerful tool utilized for mutagenesis and gene mapping. To
make them even more useful, as a pilot attempt, here we have generated a novel

deficiency kit comprising small-chromosomal deletion recombined onto FRT
chromosomes. This collection currently comprises most of the X (~67) and partial II
chromosome (~125) FRT-recombined deficiency lines. These stocks will be
submitted and made available at the Bloomington Drosophila stock centre. In this
study we have utilized these FRT recombined -deletion-bearing chromosomes for
mosaic analysis in III instar wing disc to screen predominantly for cytoskeletal
changes implicated in epithelial morphogenesis. In addition, researchers can also
utilize this approach for any of the individual deficiency lines not included in this
collection and a similar work can be extended either to other tissues for clonal
analysis or to understand any other cellular processes like neurogenesis, growth
processes, polarity-regulation etc. This approach helps to understand how genes
behave if they are deleted in a group or when small region containing genes in order
is lost. We found that overlapping deficiency lines spanning the chromosome with
varied number of genes lost in them, produced clones of varied number and sizes.
Clonal properties (size, number and distribution) and clonal behaviour (interaction
with its neighbouring cells) of mitotic clones generated by a gene loss reflects its
function. Therefore, we have further categorized them into groups based on their
clonal properties to have better understanding of the gene function. Further, using
these overlapping deficiency lines that generate cytoskeletal defects with
phenotypes, we have identified few essential loci and genes critical for maintaining
tissue integrity and growth. We also understand that those deficiency lines that do
not generate clones suggest that these loci may be a crucial region encoding
essential genes. As these deletion-bearing chromosomes are equivalent to human
chromosomes with microdeletions, we compared the such regions that caused
microdeletion-syndromes and other Mendelian disorders in humans with respective
cytological locations on Drosophila chromosomes. As contiguous-gene deletion
syndromes display extremely variable clinical symptoms depending on the genes
lost in the loci, the existing problem in diagnosing patients with MD/MenD or other
Contiguous-gene deletion syndromes or multigenic deletion syndromes is the
difficulty in identifying the critical loci for disease manifestation. This study is an
example to show how data from genomic information from multigenic syndrome
patients or MD or MenD patients can be pooled to identify the critical loci implicated
in disease manifestation. By correlating the genotype-phenotype data available for
multigenic deletion syndromes and genomic information it might be possible to
provide a better prognosis of the disease development.
522
Experimental Procedure
1. Clonal Analysis
(i) Drosophila Stocks – Drosophila stocks and crosses were maintained on a
standard yeast/cornmeal/corn syrup/malt extract/agar medium at 25^{o}C, uncles
indicated otherwise. Deletion-bearing stocks/ Deficiency lines of Exelixis, Inc (Exel),
nearly ~208 fly stocks covering X chromosome and second chromosome were
obtained from Bloomington Drosophila Stock Centre. Other stocks used for this
study include (5130)-pin/cyo; UAS-mCD8GFP, (5131)-hs-FLP; FRTG13, UAS-

mCD8GFP, (5134)-hs-FLP, tubP-GAL80, FRT19A; UAS-mCD8GFP, (5074)-

FRT40A; T155GAL4, UAS-FLP, (5075)-FRTG13; T155GAL4, UAS-FLP, (5192)-

FRT40A, tubP-GAL80; (5132)-hs-FLP, tubP-GAL80, FRT19A; (5140)- FRTG13,

tubP-GAL80; (5138)- tubP-GAL4; (8862)- hs-FLP; (5136)- UAS-mCD8GFP (X);

(5130)- UAS-mCD8GFP (III); (42725)- hs-FLP, UAS-mCD8GFP; FRT40A, tubP-

GAL80; tubP-GAL4. For generating clones by conventional method, FRT19A ;

T155GAL4, UAS-FLP, FRT40A; T155GAL4, UAS-FLP and FRTG13; T155GAL4,

UAS-FLP were used.

(ii) Recombination and Generation of clones

Sequence-Mapped deletion-bearing fly stocks (Deficiency lines) for X and II

chromosomes generated by DrosDel Project and Exelixis collection (Exel) (Ryder et

al., 2004; Parks et al., 2004; Cook et al., 2012; Roote and Russell, 2012) obtained

from Bloomington Drosophila Stock Centre (BDSC) were recombined with their

corresponding FRT chromosomes (FRT19A, FRT40A and FRTG13 chromosomes)

and recombined stocks were selected on neomycin (G418) containing media. We

have successfully recombined ~67 X chromosome deficiency lines and 125 second

(2L and 2R) chromosome deficiency lines with their respective FRT chromosomes.

List of Deficiency lines with the number of genes lost in the chromosome is provided

(Supplementary data S1-Sheet-1). Further FRT-Recombined deficiency lines were

subjected to mitotic clonal analysis using both conventional and Mosaic Analysis with

Repressible Cell Marker (MARCM) techniques (Lee and Luo, 1999, Lee and Luo,

2001; Blair, 2003; Wu and Luo, 2006) (Supplementary Figure1) in the III instar wing

imaginal discs (Figure 1(A-O)).

Analysing clones by conventional method was tedious, therefore we followed clonal

analysis by MARCM method. Total number of clones and size of each clones per

wing imaginal discs were counted and recorded using ImageJ software. Nearly 10

wing discs (n=10) were imaged and replicates of three wing discs were randomly

chosen for averaging total number of clones and size of each clone per wing

imaginal discs.

Following protocol by Wu and Luo, 2006, MARCM Stock FRT19A (hs-FLP, tubP-

GAL80, FRT19A; tubP-GAL4; UAS-mCD8GFP), MARCM Stock FRT40A (hs-FLP,

UAS-mCD8GFP; FRT40A, tubP-GAL80; tubP-GAL4) and MARCM Stock FRTG13

(hs-FLP, UAS-mCD8GFP; FRTG13, tubP-GAL80; tubP-GAL4) were assembled

using above stocks to generate MARCM clones. Individual FRT-recombined

deletion-bearing stocks were crossed with corresponding MARCM stocks. On the 3[rd]

day AEL, first instar larvae were heatshocked at 37[O]C for 1hour and allowed to

recover at 25[O]C after that. On 6[th] day AEL, from each mutant type generated, third

instar larvae were dissected and subjected to immunostaining (Usha and

Shashidhara, 2010; Szuperak et al., 2011).

(iii) Immunostaining and Clonal Analysis

Wing Imaginal discs were immunologically stained using Phalloidin followed by Alexa594 and viewed under the confocal microscope. Their images were then analysed for clone size, distribution and number using Image J software. The clones of each of the genes mutated were analyzed in triplicates each for both the fly lines. The average clone sizes and clone numbers were calculated in each case, with the most approximate result in each case (For discs with <10 clones, all clones were considered for mean clone number estimation; For discs with >50 clones, 50 of the clones were taken into consideration for mean estimation; as well as for discs with intermediate number of clones, suitably, either 10, 20, 30 or 40 clones were considered for best estimate of the average). A triple blind test was also performed to corroborate each of these results.

(2) Systematic Analysis

We began the systematic analysis by identifying the genes deleted in each of the available deficiency lines from https://bdsc.indiana.edu/. We then built an inventory with the deleted genes arranged according to their cytological location, ensuring that there are no repetitions, against their respective deficiency lines. Alongside, a list of their corresponding human orthologs was generated using the gene summaries available at www.flybase.org → Orthologs → Human orthologs .
Apart from this, we assessed each gene for cytoskeletal implications from the gene ontology summaries updated on their respective flybase web pages and we have categorized based on their function. (www.flybase.org → Gene Ontology → Molecular Function / Biological Processes / Cellular Component). Similarly, we also scrutinized the human orthologs for any cytoskeletal implications, under the same categories, using the database available at https://www.ncbi.nlm.nih.gov/gene/. For those genes that were involved in functions that fell into more than one of these categories, we took their most predominant function into account for grouping and analyzing.

We used the Molecular Genetics Laboratory group, MGenoma®'s webpage as a reference for a list of microdeletion syndromes. From the table present, we enlisted the genes that were deleted in each of the syndromes. We identified among them, those human orthologs that we previously enlisted in the inventory generated, and also their respective fly counterparts. For each of the microdeletion syndromes, we used the database from Orphanet to find the critical region, or the mutational hotspot. The name of the disease was given in the search field and the molecular genetics section of the clinical genetics review for each disease was referred to, in order to identify the mutational hotspot or the critical region of the gene(s) deleted.

Figure 1-Distribution and size of MARCM clones. Representative Wing discs (A-O) with Mitotic clones (GFP positive, counterstained with DAPI in blue) generated using Deficiency lines on X chromosome recombined to FRT19A chromosomes. Note the variation in size and distribution of the clones in wingdiscs.

619
Figure 2-Number and size of clones generated by deficiency lines on X
chromosome. Classification of clones based on number of clones (A) and size of the
clones (B). Note average number (D) and size of mitotic clones (F) generated by
deficiency lines spanning the X chromosome. Correlation between number of genes
lost in deficiency lines on X chromosomes with average number (C) and size of
mitotic clones (E) generated by them.
626
627 Figure 3-Number and size of clones generated by deficiency lines on II chromosome.
628 Correlation between number of genes lost in deficiency lines on II chromosomes with
629 average number (A) and size of mitotic clones (C) generated by them. Note average
630 number (B) and size of mitotic clones (D) generated by deficiency lines spanning the
631 X chromosome.
632
Figure 4 - Wing discs with cytoskeletal breakages and disruptions. Compared to
control wingdiscs with MARCM clones (A1-A6), GFP positively marked clones by X
chromosome deficiency line BL-7703 display cell autonomous (B1-B6) and
deficiency line BL-7707 display non-cell autonomous (C1-C6) cytoskeletal defects as
shown by Actin (Phalloidin, Red). Compared to control wingdiscs with conventional
clones (D1-D6), GFP negatively marked clones by II chromosome deficiency line BL-
7813 display cell autonomous (E1-E6) and deficiency line BL-7815 display non-cell
autonomous (F1-F6) cytoskeletal defects as shown by Actin (Phalloidin, Red). All
panels are standard confocal XY sections.
Note panels B3, C3, E3 and F3 for the changes in Actin (Red) representing
cytoskeletal breakages and disruptions in wingpouch.
644
Figure 5 - Wing discs with cytoskeletal defects and extrusions. Two deficiency lines
BL-7716 (A'-A", 5C') and BL-9348 (B'-B", 5C") show similar and consistent
cytoskeletal defects in clones generated either by conventional method or MARCM
strategies. Cells and tissue adjacent to the clones are extruded out of the surface.
Localized cytoskeletal defects are produced either along the periphery of the pouch
region as shown by deficiency lines BL-9348 (5B" and 5C") or only in the hinge
region as shown by deficiency line BL-7709(D'-D"). Note the rounding of the clones
and associated with developmental defects.
653
654 Figure 6 - Wing discs with severe cytoskeletal defects associated with outgrowths
655 and duplications. Compared to the control (A) rare clone produced by two deficiency
656 lines BL-8948 (B) and BL-8956 (E-E"). Note the outgrowth of the wingpouch in apical
657 (E), medial (E') and basal (E") XY sections and XZ section of the medial region (F).
658
659 Figure 7 - Wing discs with severe cytoskeletal defects associated with outgrowths
660 and duplications. Compared to control wingdiscs with conventional clones (A) or
661 MARCM clones(C) clones generated by deficiency line BL-7815(B and D)
662 consistently show a similar phenotype of outgrowth and duplication of wing pouch.

Figure 8 - Classification and comparison of genes implicated in cytoskeletal functions. In the X chromosome, 181 genes out of 1103 total genes and in II chromosome, 431 genes out of 1245 genes are involved in mediating cytoskeletal functions. Those functions are classified into five categories and percentage of genes involved are compared and represented.

Figure 9 - Comparative analysis of Drosophila deficiency lines with Human Mendelian Diseases and Microdeletion Syndrome.
A detailed Flowchart (A) and a sunburst representation(B) of systematic approach to compare genes in the Drosophila Deficiency lines of X and II chromosome with human orthologs . Analytical representation of comparison of genes on X and II chromosomes coding for 52 human Microdeletion syndrome (c).

Figure 10 - Identification of critical region based on clonal behaviour. For the locus 2F6-3A8, two big deficiency lines BL-9345 (10A, E) and BL-8031(10D-E) produced a smaller number of clones while two smaller deficiency lines BL-7705 (B, E) and BL-7706(C, E) produced high number of big sized clones (B, E). By comparing their clonal properties and cytoskeletal information, we identify that the critical gene is *Boi* (E). Similarly, for the loci 3A3-3B1 and 30B10-30C9 (for clones Fig 11, Fig12) the critical gene is *Shaggy (Sgg)* and *GlcAT-S(F)* respectively. Schematic representation of the critical region identified in this study (H).

Figure 11 - Identification of critical region based on the cytoskeletal defects. On the X chromosome, three deficiency lines BL-9518(A-A'), BL-8031(B) and BL-9348(C) that produced moderate number of clones were compared for their cytoskeletal defects. For the locus 3A3-3B1, severe phenotype produced by BL-9348(C, D'-D") showing cell-autonomous extrusion of the clones (C, D'-D", E-E') from the wing surface is encoded by the critical gene *Shaggy (Sgg).*

Figure 12 - Identification of critical region using non-overlapping deficiency lines. MARCM clones for the locus 10D1-10D7, a big deficiency line BL-7716 produced a cytoskeletal defect (B). A smaller deficiency line, BL-7759 covering a gene ⬚-*1,6, fucosyltransferase (FUCT6)* produced a similar cytoskeletal defect (C).

Figure 13 - Identification of critical region for locus with opposing phenotypes
For the locus 30B10-30C9, a big deficiency line (BL-7812) produced moderate number of clones (B) while an overlapping deficiency lines produced rare or very few clones (C). Note wingdiscs produced by both deficiency lines were slightly larger than the control(A-C). A small, overlapping deficiency line (BL-7815) containing two genes GlcAT-S and Trx-2 for the locus 30C1-30C1 prodeuced overgrowth and duplication of the wingdisc (D'-D"). Overexpression clones of GlcAT-S, produced cytoskeletal defects(F) and clones generated by FRT-recombined GlcAT-S^{KG} hypomorph produced overgrowth and cytoskeletal defects(G).

707
708 Supplementary Figure 1 - Mitotic clones. Control Wingdisc showing mitotic clones
709 generated (A) by conventional method where clones are negatively marked, and
710 counterstained with Phalloid (Actin, RED) and (B) by Mosaic Analysis with
711 Repressible Cell Marker (MARCM) techniques where clones are positively marked
712 (GFP) and counter stained with DAPI (Nucleus, Blue).
713
714 Supplementary Figure 2 - Variation in localization and number of clones.
715 Representative Wingdiscs with localized clones (A, B, E), few clones (C, D),
716 moderate clones (E, F) and high number of clones (G, H).
717
718 Supplementary Data Table S1 - Excel sheet with list of deficiency lines used for
719 recombination. X chromosome (Sheet1) and II chromosomes (Sheet 2)
720 (Supplementary Data Table S2) – Number of clones and average size of clones
721 Supplementary Data Figure S3)- Mitotic clones Range from no clones to big clones
722

723 **Authors Contributions**
724 Conceptualization: UN, SP; and Methodology: UN, SP, SS, RR, NR; Investigation:
725 UN, SP, TK; Formal Analysis: UN, SP, SS, RR, NR; Writing- UN, SP, TK; Writing-
726 Review and Editing- UN, SP, NR, TK Funding Acquisition: UN; Supervision: UN, SP.
727

Acknowledgments
We gratefully acknowledge Dr Matt Gibson and Dr Marios Georgiou for their help
with fly experiments in their laboratories. We thank the Bloomington stock centers
and the fly community for the generosity with reagents. We thank Prof Satish Kumar
and Dr Vasanthi Dasari for helpful advice and critical comments on the manuscript.
We thank Science and Engineering Research Board (SERB), Department of Science
and Technology (DST), Government of India, for the grants (YSS/2014/000896) and
CRG/2018/003725 to UN.
736

737 **Declaration of Interests:**
738 The authors declare no competing interests.
739

Reference
Akiyama ,T., User, SD and Gibson, MC (2018). eLife ;7: e35258. DOI:
https://doi.org/10.7554/eLife.3525
Bejsovec, A. (2018). Wingless Signaling: A Genetic Journey from Morphogenesis to
Metastasis. Genetics 208(4): 1311-1336.
Blair SS (2003). Genetic mosaic techniques for studying *Drosophila* development.
Development 130: 5065-5072; doi: 10.1242/dev.00774
Camp, D., Haitian He, HB., Li, S., Althaus, IW., Holtz, AM., Allen, BL., Charron, F.,
van Meyel, DJ. (2014). Ihog and Boi elicit Hh signaling via Ptc but do not aid Ptc
in sequestering the Hh ligand. Development 141(20): 3879--3888.

Chial, H. (2008). Somatic mosaicism and chromosomal disorders. Nature Education 1(1):69

Clevers, H. (2006). Wnt/beta-catenin signaling in development and disease. Cell 127(3): 469-480.

Cook, RK., Christensen, SJ., Deal, JA., Coburn, RA., Deal, ME., Gresens, JM., Kaufman, TC., Cook, KR. (2012). The generation of chromosomal deletions to provide extensive coverage and subdivision of the Drosophila melanogaster genome. Genome Biol. 13(3): R21.

Donohoe CD, Csorda´s G, Correia A, Jindra M, Klein C, Habermann B, et al. (2018). Atf3 links loss of epithelial polarity to defects in cell differentiation and cytoarchitecture. PLoS Genet.14 (3): e1007241. https://doi.org/10.1371/journal. pgen.1007241.

Frappaolo, A, Sechi,S, Kumagai, T, Karimpour-Ghahnavieh, A, Tiemeyer,M, Giansanti, MG. (2018). Modeling Congential Disorders of N-Linked Glycoprotein Glycosylation in Drosophila melanogaster. Front. Genet. 9():436.

Franco, B., Bogdanik, L., Bobinnec, Y., Debec, A., Bockaert, J., Parmentier, M.L., Grau, Y. (2004). Shaggy, the homolog of glycogen synthase kinase 3, controls neuromuscular junction growth in Drosophila. J. Neurosci. 24(29): 6573--6577.

Fortunato, EA and Spector DH (2003). Viral induction of site specific chromosome damage. Rev Med Virol. 13(1) pg21-37. https://doi.org/10.1002/rmv.368.

Germani, F., Bergantinos C and Johnston LA(2018). Mosaic Analysis in *Drosophila* Genetics 208(2) 473-490 ;https://doi.org/10.1534/genetics.117.300256.

Giansanti MG and Fuller MT (2012). What Drosophila spermatocytes tell us about the mechanisms underlying cytokinesis. Cytoskeleton (Hoboken). 69(11):869-81. doi: 10.1002/cm.21063.

Gibson, MC. And Perrimon N (2005). Extrusion and Death of DPP/BMP-Compromised Epithelial Cells in the Developing Drosophila Wing. Science, 307(5716), 1785–1789. doi:10.1126/science.1104751

Gumbiner BM. (1992). Epithelial morphogenesis. Cell. 1;69(3):385-7.

Hsia, EYC., Zhang, Y., Tran, HS., Lim, A., Chou, YH., Lan, G., Beachy, PA., Zheng, X. (2017). Hedgehog mediated degradation of Ihog adhesion proteins modulates cell segregation in Drosophila wing imaginal discs. Nat. Commun. 8(1): 1275.

Knust E (2005).Development of Epithelial Cell Polarity in Drosophila Volume 7 Chapter 2 p121. Edited by (Birchmeier W and Birchmeier C, Epithelial Morphogenesis In Development And Disease(Cell adhesion and communication) ISBN 0-203-30376-8

Lee, T and Luo L. (1999). Mosaic analysis with a repressible cell marker for studies of gene function in neuronal morphogenesis. Neuron. 22(3):451-61.

Lee, T, Luo L. (2001). Mosaic analysis with a repressible cell marker (MARCM) for Drosophila neural development. Trends Neurosci. 24(5):251-4.

Olson, MV (1999). When less is more: Gene loss as an engine of evolutionary change. Am J Hum Genet 64: 18–23.

Olson, MV and Varki A (2003) Sequencing the chimpanzee genome: Insights into human evolution and disease. Nat Rev Genet 4: 20–28.

Paciorkowski, AR, Keppler-Noreuil K, Robinson L, et al. (2013). Deletion 16p13.11 uncovers NDE1 mutations on the non-deleted homolog and extends the

spectrum of severe microcephaly to include fetal brain disruption. Am J Med
Genet A. 161A(7):1523–1530.

Parks, AL., Cook, KR., Belvin, M., Dompe, NA., Fawcett, R., Huppert,
K., etal., (2004). Systematic generation of high-resolution deletion coverage of
the Drosophila melanogaster genome. Nat Genet. 36(3):288-92.

Paschinger, K., Straudacher, E., Stemmer, U., Fabini, G., Wilson, I.B.H. (2005).
Fucosyltransferase substrate specificity and the order of fucosylation in
invertebrates. Glycobiology 15(5): 463--474.

Price, MA and Kalderon, D. (2002). Proteolysis of the Hedgehog signaling effector
Cubitus interruptus requires phosphorylation by Glycogen Synthase Kinase 3
and Casein Kinase 1. Cell 108(6): 823--835.

Powell, K and Piddini, E (2016). Chasing how cells out compete one another.
Journal of Cell Biology. 213(3): 291–292.

Rauch, A; Schellmoser, S; Kraus, C; Dörr, HG; Trautmann, U; Altherr, MR; Pfeiffer,
RA; Reis, A (2001). First known microdeletion within the Wolf-Hirschhorn
syndrome critical region refines genotype-phenotype correlation. American
Journal of Medical Genetics. **99**(4): 338-
42. doi:10.1002/ajmg.1203. PMID 11252005.

Reiter, LT., Potocki, L., Chien, S., Gribskov, M., Bier E. (2001). A systematic analysis
of human disease-associated gene sequences in Drosophila melanogaster.
Genome Res. 11(6):1114-25.

Roos, C., Kolmer, M., Mattila, P., Renkonen, R. (2002). Composition of Drosophila
melanogaster proteome involved in fucosylated glycan metabolism. J. Biol.
Chem. 277(5): 3168--3175.

Rodríguez, AV, Didiano D, Desplan C (2012). Power tools for gene expression and
clonal analysis in Drosophila Nature Methods 9: 47–55.

Roote, J and Russell, S. (2012). Toward a complete Drosophila deficiency
kit. Genome Biol. 13(3): 149.

Ruddle, FH, Bentleyl, KL, Murtha MT, Risch N (1994). Gene loss and gain in the
evolution of the vertebrates. Development Supp. 155-161.

Ryder, E., Blows, F., Ashburner, M., Bautista-Llacer, R., Coulson, D., Drummond, J.,
Webster, J., et al., (2004). The DrosDel collection: a set of P-element insertions
for generating custom chromosomal aberrations in Drosophila
melanogaster. Genetics 167(2): 797-813.

Ryder, E., Ashburner, M., Bautista-Llacer, R., Drummond, J., Webster, J., Johnson,
G., et al., (2007). The DrosDel deletion collection: a Drosophila genomewide
chromosomal deficiency resource. Genetics. 177(1):615-29.

Schock, F and Perrimon N. (2002). Molecular mechanisms of epithelial
morphogenesis. Annu Rev Cell Dev Biol. 18:463-93. DOI:
10.1146/annurev.cellbio.18.022602.131838.

Shen, J and Dahmann C (2005). Extrusion of Cells with Inappropriate Dpp Signaling
from Drosophila Wing Disc Epithelia. Science, 307(5716), 1789–1790.
doi:10.1126/science.1104784.

Smith-Bolton, RK., Worley, MI., Kanda, H and Hariharan, IK. (2009) Regenerative
growth in Drosophila imaginal discs is regulated by Wingless and Myc. Dev.
Cell 16: 797–809.

Slavotinek, A. (2012). Microdeletion Syndromes. In eLS, (Ed.).
doi:10.1002/9780470015902.a0005549.pub2
St Johnston, D., (2002). The art and design of genetic screens: Drosophila
melanogaster. Nat. Rev. Genet. 3: 176–188.
Sustar A., Bonvin M., Schubiger M and Schubiger G (2011) *Drosophila* twin spot
clones reveal cell division dynamics in regenerating imaginal discs.
Developmental Biology (356)2: 576-587.
Szuperak M, Salah S, Meyer E, Usha N, Ikmi A and Gibson M. (2011) Feedback
modulation of *Drosophila* BMP signaling by the novel extracellular protein,
Larval Translucida Development 138, 715-724.
Nagarajan U, Pakkiriswami S and Pillai AB. (2015) Sugar tags and tumorigenesis.
Frontiers in Cell and Developmental Biology 3(69), 1-5.
Usha N and Shashidhara LS. (2010) Interaction between Ataxin-2 Binding Protein 1
and Cubitus- interruptus during wing development in *Drosophila*.
Developmental Biology 341, 389–399.
Yamamoto S, Jaiswal M, Charng WL, Gambin T, Karaca E, Mirzaa G, et al., (2014).
A drosophila genetic resource of mutants to study mechanisms underlying
human genetic diseases. Cell. 25;159(1):200-214. doi:
10.1016/j.cell.2014.09.002.
Wu, JS and Luo L. (2006). A protocol for mosaic analysis with a repressible cell
marker (MARCM) in Drosophila. Nat Protoc. 1(6):2583-9.

Zhang W, Hong M, Bae GU, Kang JS, Krauss RS. (2011) Boc modifies the
holoprosencephaly spectrum of Cdo mutant mice. Dis Model Mech. 4(3):368-
80. doi: 10.1242/dmm.005744.

866
867
868 Table 1 **Category based on Morphological/cytoskeletal changes**

	Category	Number of Deficiency lines (DLs)	List of DLs
	Without any Morphological changes	49	All others
	Mild Morphological changes (medium-sized, moderate to numerous clones)	5	Df(1)-Exel6226(BL-7703) Df(1)-Exel6233(BL-7707) Df(1)Exel6290(BL-7753) Df(1)Exel6237(BL-7711) Df(1)ED6989(BL-9056)
Widely distributed clones in wingdisc with Morphological defects	Severe morphological changes (cytoskeletal breakages and disruptions)	2	Df(1)Exel6226(BL-7703) Df(1)Exel6233(BL-7707)
		2	Df(2L)Exel8022(BL-7813) Df(2L)Exel9040(BL-

			7815)
	Morphological changes with outgrowths	2	Df(1)ED6630(BL-8948) Df(1)ED6727(BL-8956)
		1	Df(2L)Exel6029-BL-7512
	Morphological changes with extrusions and outgrowths	2	Df(1)Exel6242(BL-7716) Df(1)ED6584(BL-9348)
	Morphological changes with duplications	2	Df(1)ED6630(BL-8948) Df(1)ED6727(BL-8956)
		1	Df(2L)Exel9040(BL-7815)
Localized clones in wingdisc with Morphological defects	Only in Notum	2	Df(1)Exel6235(BL-7709) Df(1)ED6584(BL-9348)
	Only in Hinge		

869
870
871 **Table 3 Deficiency lines in Drosophila and Microdeletion Syndromes in**
872 **Humans**

Microdeletion Syndromes	X	li	Prevalence in Humans	Mutation hotspot in Humans
17q21.31 microdeletion syndrome	2	4		KAT8 regulatory NSL complex subunit 1 - KANSL1
1p36 deletion syndrome	1	2		NA
1q21.1 deletion syndrome, 1.35 Mb	0	4		GJA5, GJA8 (CHD1L, BCL9, and PRKAB2)
3q29 microdeletion syndrome	2	3		NA
Adrenal hypoplasia, Congenital	0	1	25%	NROB1, GK, DMD, IL1RAPL1
Angelmann's syndrome	1	2		MBD5
Aniridia	1	3	17-22%	PAX6
Autism	11	10		ADNP
Basal Cell Nevus syndrome	0	1	1/6-21%	PTCH1, SUFU
Beckwith-Wiedemann Syndrome	2	17		H19, IGF2, CDKN1C, KCNQ1, KCNQ1OT1
Brachydactyly mental retardation	5	4		HDAC4

syndrome				
Branchiootorenal syndrome	1	6		EYA1, SIX5, SIX1
Bruton agammaglobulinemia	0	1	8%	BTK
Campomelic Dysplasia	1	3	rare	SOX9
Charcot-Marie-Tooth Syndrome	7	5		GJB1
Cleidocranial dysplasia	0	7	10%	RUNX2
Cornelia De Lange syndrome	0	4		HDAC8, NIPBL, RAD21, SMC1A, SMC3
Cri-du-chat syndrome	1	2		catenin delta 2 - CTNND2 semaphorin 5A - SEMA5A
Dandy Walker syndrome	1	6		ZIC1, ZIC4 (PROBABLY)
Diaphragmatic Hernia, Congenital	1	6	rare	*NR2F, CHD, RGMA, DISP1, GATA4 and SIAT8B*
Digeorge syndrome	6	4		TBX1
Down syndrome	0	4	rare	NA
Greig Cephalopolysyndactyly syndrome	2	1	5-10%	GLI3
Holoprosencephaly	7	23		SHH, ZIC2, SIX3, TGIF1, GLI2, PTCH1
Kabuki syndrome	0	1		KDM6A, KMT2D
Leri-Weill dyschondrosteosis	0	2	rare	SHOX
Mental retardation Autosomal Dominant 51	5	0		NA
Metachromatic leukodystrophy	0	1	<1%	ARSA
Microphthalmia, Syndromic 7	1	6		COX7B, HCCS, NDUFB11
Miller-Dieker Lissencephaly syndrome	3	4		PAFAH1B1
Muscular dystrophy, Becker Type	1	4		DMD
Nail-patella syndrome	4	6	5-10%	LMX1B
Neurofibromatosis	10	13		NF1
Neuropathy hereditary, with liability to pressure palsies	0	1		PMP22

Noonan Syndrome 1	2	5			MAP2K2, NF1 (PROBABLY)
Polycystic kidney disease, infantile severe, with Tuberus Sclerosis	0	15			PKD1, TSC2 (PROBABLY)
Potocki-Lupski syndrome	5	0			FLCN, RAI1
Potocki-Shaffer syndrome	4	1			ALX4, EXT2, PHF21A (PROBABLY)
Prader-Willi Syndrome	0	1			SIM1
Retinoblastoma Rb1	1	1	8-16%		RB1
Rett's syndrome	1	0	8%		MECP2
Rieger Syndrome Type I	1	3			PITX2, FOXC1
Rubinstein Taybi Syndrome	10	0			CREBBP, EP300
Saethre Chotzen syndrome	2	1			TWIST1
Split hand/foot malformation 1	4	6	rare		TP63
Towne's Brocks syndrome	2	8	5%		SALL1
Trichorhinophalangeal syndrome	0	1	3-4%		EXT1, RAD21, TRPS1
Tuberus Sclerosis	1	0	<2%		TSC1, TSC2
William-Beuren region duplication syndrome	3	5			ELN, NCF1
Wilm's Tumor 1, Mental retardation	1	0	14%		PAX6
Wolf-Hirschhorn Syndrome	0	3			CPLX1, CTBP1, FGFRL1, LETM1, NELFA, NSD2, PIGG

873
874 **Table 4 Correlation of genes on X chromosomes of Drosophila and Human**
875 **associated with Mendelian diseases**

Drosophila Gene	Cytolocation on Fly genome	Human gene	OMIM	Cytolocation on Human genome	Mendelian Disorder associated

CG8184	13E8-13E12	HUWE1; HECT, UBA and WWE domain containing 1, E3 ubiquitin protein ligase	300697	Xp11.22	Mental retardation, X-linked syndromic, Turner type
		SPACA5; sperm acrosome associated 5	300593	Xp11.23	
regucalcin	11A1-11A1	RGN; regucalcin	300212	Xp11.3	
Efhc1.1	13E17-13E18	EFHC2; EF-hand domain containing 2	300817	Xp11.3	
		PHEX; phosphate regulating endopeptidase homolog X-linked	300550	Xp22.11	Hypophosphatemic rickets, X-linked dominant
		PRDX4; peroxiredoxin 4	300927	Xp22.11	
CG7024	4C14-4C14	PDHA1; pyruvate dehydrogenase E1 alpha 1 subunit	300502	Xp22.12	Pyruvate dehydrogenase E1-alpha deficiency
		PDHA1; pyruvate dehydrogenase E1 alpha 1 subunit	300502	Xp22.12	Pyruvate dehydrogenase E1-alpha deficiency
CG7766	8C8-8C9	PHKA2; phosphorylase kinase regulatory subunit alpha 2	300798	Xp22.13	Glycogen storage disease, type IXa1
cib		TMSB4X	300159	Xp22.2	
Moe	8B4-8B6	MSN; moesin	309845	Xq12	Immunodeficiency 50
		OPHN1; oligophrenin 1	300127	Xq12	Mental retardation, X-linked, with cerebellar hypoplasia and distinctive facial appearance

Klp3A	3A6-3A6	KIF4A; kinesin family member 4A	300521	Xp12	Mental retardation, X-linked 100
		PHKA1; phosphorylase kinase regulatory subunit alpha 1	311870	Xq13.1	Muscle glycogenosis
CG32683	9C1-9C4	ARR3; arrestin 3	301770	Xq13.1	Myopia 26, X-linked, female-limited
		DLG3; discs large MAGUK scaffold protein 3	300189	Xq13.1	Mental retardation, X-linked 90
nod	10C7-10C8	KIF4A; kinesin family member 4A	300521	Xq13.1	Mental retardation, X-linked 100
		NAPIL2	300026	Xq13.2	
		KLHL4; kelch like family member 4	300348	Xq21.31	
		NAPIL3	300117	xq21.32	
		RAB40AL; RAB40A like	300405	Xq22.1	
CG5941	5D2-5D2	MCTS1; MCTS1, re-initiation and release factor	300587	Xq24	
		TENM1; teneurin transmembrane protein 1	300588	Xq25	
CG12796	6D4-6D4	GPR119	300513	Xq26.1	
		SAGE1; sarcoma antigen 1	300359	Xq26.3	
Clic	12C5-12C6	CLIC2; chloride intracellular channel 2	300138	Xq28	Mental retardation, X-linked, syndromic 32
CG8565	13F9-13F10	SRPK3; SRSF protein kinase 3	301002	Xq28	
scu	16F6-16F6	HSD17B10; hydroxysteroid 17-beta dehydrogenase 10	300256	Xp11.22	HSD10 mitochondrial disease

CG2694	3B3-3B3	GCNA; germ cell nuclear acidic peptidase	300369	Xq13.1	
		ATP7A; ATPase copper transporting alpha	300011	Xq21.1	Menkes disease, Occipital horn syndrome, Spinal muscular atrophy, distal, X-linked 3
		OCRL; OCRL, inositol polyphosphate-5-phosphatase	300535	Xq26.1	Dent disease 2, Lowe syndrome
		F8A1; coagulation factor VIII associated 1	305423	Xq28	

876
877
878

FRT19A,Df(1)Exel6231-(BL-7706) FRT19A,Df(1)Exel6242-(BL-7716) FRT19A,Df(1)Exel6225-(BL-7702) FRT19A,Df(1)Exel6223-(BL-7700)

A B C D

FRT19A,Df(1)Exel6234-(BL-7708) FRT19A,Df(1)ED6630-(BL-8948) FRT19A,Df(1)ED6712-(BL-9169) FRT19A,Df(1)Exel6226-(BL-7703)

D E F G

FRT19A,Df(1)ED7344-(BL-9220) FRT19A,Df(1)ED7265-(BL-9414) FRT19A,Df(1)Exel6221-(BL-7699) FRT19A,Df(1)ED7170-(BL_8898)

H I J K

FRT19A,Df(1)E-(BL-9218) FRT19A,Df(1)ED7289-(BL-29732) FRT19A,Df(1)Ex6237-(BL-7711) FRT19A,Df(1)ED6396-(BL-9052)

L M N O

| | MARCM Clone | Phalloidin WT | Genes/Clones DAPI | | | |
| | 1 | 2 | 3 | 4 | 5 | 6 |

A Control

B BL-7703

C BL-7707

| | Conventional Clone | Phalloidin WT | Genes/Clones DAPI | | | |
| | 1 | 2 | 3 | 4 | 5 | 6 |

D Control

E BL-7813

F BL-7815

Conventional Clone · Phalloidin · WT · Genes/Clones · DAPI

A′ BL-7716
A″

B′ BL-9348
B″

MARCM Clone · Phalloidin · WT · Genes/Clones · DAPI

C′ BL-7716
C″ BL-9348

MARCM Clone · Phalloidin · WT · Genes/Clones · DAPI

D′ BL-7709
D″

Classification and Comparison of genes implicated in various cytoskeletal functions

A

Flow chart of Comparative analysis

X chromosome	II chromosome
60 Deficiency Lines	125 Deficiency Lines
1108 Genes	1769 Genes

With Human Orthologs		Without Human Orthologs	
600 genes	1090 genes	503 genes	679 genes

With OMIM + disease		Only OMIM No disease	
189 genes	509 genes	411 genes	581 genes

Mendelian Disease		Microdeletion Syndrome	
70 genes	282 genes	119 genes	227 genes

B

C

Comparison of genes coding for Microdeletion syndrome on X and II chromosomes

Legend:
2 3
5 6
8 9
11 12
14 15
17 18
20 21
23 24
26 27
29 30
32 33
35 36
38 39
41 42
44 45
47 48
50 51

A   B   C   D

	Deficiency Line	No. of genes deleted	No. of clones	Size of clones	Cytolocation	Critical gene
	Based on Clonal Behavior					
E	Df(1)Exel6230 (BDSC-7705)	9	91	0.6	3A2-3A4	*Boi*
	Df(1)Exel6231 (BDSC-7706)	6	65	0.4	3A2-3A3	
	Df(1)ED11354 (BDSC-9345)	15	13	0.05	2F6-3A4	
	Df(1)ED411 (BDSC-8031)	18	3	0.2	3A3-3A8	
	Based on Cytoskeletal defects					
F	Df(1)ED6584 (BDSC-9348)	3	36	0.12	3A8-3B1	*Sgg*
	Df(1)ED6679 (BDSC-9518)	10	8	0.08	3A6-3A8	
	Df(1)ED411 (BDSC-8031)	18	3	0.2	3A3-3A8	
G	Df(2L)Exel6024 (BDSC-7504)	19	36	0.6	30B10-30C1	
	Df(2L)Exel7042 (BDSC-7812)	14	1	0	30C1-30C9	
	Df(2L)9040 (BDSC-7815)	2	35	2.6	30C1-30C1	*GlcAT-S*

Cytological locations   2F6   3A2   3A3   3A4 3A6 3A8 3B1

H

Df(1)-9345   15
Df(1)-7705   9   boi
Df(1)-7706   6
Df(1)-8031   18
Df(1)-9348   3 Sgg
Df(1)-9518   10

M=MARCM Clone Phalloidin WT Genes/Clones DAPI

C=Conventional Clone Phalloidin WT Genes/Clones DAPI

M=MARCM Clone Phalloidin WT Genes/Clones DAPI

C=Conventional Clone Phalloidin WT Genes/Clones DAPI