A system to decipher the origins and relationships of all arthropod outgrowths

Authors: Heather S. Bruce[1]* and Nipam H. Patel[2,3]

Affiliations: 1. University of British Columbia, Vancouver, BC V6T1Z4 Canada. 2. Marine Biological Laboratory, 7 MBL St. Woods Hole, MA 02543 USA. 3. University of Chicago, Organismal Biology & Anatomy, 1027 E 57th Street, Chicago, IL 60637 USA.

*Correspondence to: heather.bruce@ubc.ca

Abstract: Arthropods (insects, "crustaceans", myriapods, and chelicerates) display a fascinating
15 diversity of ectodermal structures like plates, horns, helmets, knobs, carapaces, mimicry outgrowths, and wings. The origins and relationships of these structures has tantalized researchers for over a century: did these structures arise de novo in each lineage, or do they emerge from shared, ancestral primordia present in some or even all arthropods? One way to begin answering this is to assess the position and context of these structures on arthropod bodies:
20 do these structures emerge from proximal embryonic leg segments that were converted into the body wall, or do they represent dorsal, non-leg-derived structures? Here, the expression of *pannier*, *araucan*, and *drumstick* – genes previously shown to distinguish proximal leg segments in crustaceans and insects – are examined in a chelicerate representative, the tarantula *Acanthoscurria*. This gene expression comparison, together with over a century of gene
25 functional, morphological, embryological, and paleontological data, suggests that all arthropod leg segments correspond to each other in a one-to-one fashion, but that in many arthropod lineages, the base of the ancestral leg in the embryo flattens and expands to form the body wall of the adult. This would mean that many arthropod outgrowths which appear to stand on the lateral body wall, such as wings, tergal plates, and gin traps, are derived from the
30 ancestral/embryonic leg base, and likely arose from shared, ancestral primordia present on this leg base in all arthropods. The analysis detailed here means that a simple three- or four-gene in situ expression experiment with *pannier*, *araucan*, and *drumstick* can elucidate the homologies of any arthropod ectodermal structure of interest, including beetle horns, treehopper helmets, and more.

Key words

Arthropod, appendage, homology, *Parhyale, Tribolium, Acanthoscurria*

Introduction

40   The diversity of arthropod appendages has facilitated their success across every biome. There are four main groups of living arthropods: chelicerates (spiders, etc.), myriapods (millipedes, etc.), "crustaceans" (i.e. the paraphyletic clade of non-insect pancrustaceans, such as shrimps, etc.), and insects (beetles, etc.). Each group has different numbers of and names for their anatomical leg segments (Fig. 1)[1–5]. Understanding how the leg segments of the four

45   different arthropod groups correspond to each other has tantalized researchers for over a century[2,4,6–11], because this would answer generations of speculation about the origins and possible relationships of fascinating arthropod structures like wings, gills, helmets, and horns[6,12–23].

        In our 2020 paper, we sought to understand the leg segment relationships between two of

50   the four groups of arthropods, insects and crustaceans[24]. To do so, we compared a century of previous morphological and embryological work with the expression and function of several leg- and wing-patterning genes between insects and crustaceans, using original and published data. Authors have historically proposed a variety of leg segment deletions, duplications, and/or fusions to account for the different numbers of leg segments in the four arthropod groups[2,4,8,14,25–

55   32]. However, our work (Fig. 1e) suggested that insect and crustacean leg segments have a simple, one-to-one correspondence, and the apparent loss of two leg segments in insects is due to the embryonic expansion and flattening of two proximal leg segments to form the adult insect lateral body wall (pleura and lateral tergum; also called subcoxa and precoxa/epicoxa, respectively)[14,33–37]. We found that two genes – the Iroquois complex gene *araucan* (*ara*) and the GATA factor

60   *pannier* (*pnr*) – distinguish these proximal leg segments even if, as in insects, the leg segments

now function as body wall[24,38–45]. This molecular comparison supported previous morphological, embryological, and fossil studies which suggested that arthropods ancestrally have a total of 8 leg segments[2,4,8,34,46,47].

We therefore wondered whether this model held true for all arthropods. There is currently no gene function data for myriapods, so we focused on chelicerates. To determine whether chelicerates with 7 leg segments had converted an ancestral/embryonic proximal leg segment into the body wall, we investigated the expression of *pnr*, *ara,* and the arthropod joint marker *drumstick* in the tarantula spider *Acanthoscurria geniculata*. We then placed our results within the broader context of over a century of morphological, embryological, fossil, and molecular studies in all arthropods.

Results

Extinct arthropods at the base of the arthropod phylogenetic tree, as well as early-branching chelicerates[48,49] like sea spiders, camel spiders, and some mites, have 8 true (eudesmatic) leg segments[4,46], but later-branching chelicerates like spiders and harvestman have only 7 apparent leg segments[4]. This begs the question of whether and how the ancestral 8th leg segment was lost in these chelicerates: was this ancestral leg segment deleted, or was it converted into the body wall as insects have done?[8,24,28,33,34,36,47,50–54]

To elucidate the composition of proximal leg segments in chelicerates, the expression of *pnr*, *ara,* and *Dll* was compared between the crustacean *Parhyale*, the insect *Tribolium*, and the tarantula *Acanthoscurria geniculata* (Fig. 2)[24]. In *Drosophila*, *pannier* (*pnr*) patterns the dorsal midline, extra-embryonic membranes, and heart. Loss of *pnr* results in failure of embryonic dorsal closure, deletion of the dorsal midline ectoderm[43,55], and failure of the dorsal mesoderm to become the heart[56], whereas cells in the wing induced to express *pnr* form dorsal medial body

85   wall (notum)[39]. Iroquois complex genes, including *araucan-caupolican* (*ara-caup*), pattern the

left and right lateral notum in *Drosophila*, and loss transforms the lateral body wall (notum) into

wing hinge[38,40,41,57].

        From the *Acanthoscurria* transcriptome[58], OrthoFinder[59] identified two paralogs of *pnr* in

Acanthoscurria that were the reciprocal best blast hits to the single *pnr* ortholog in *Drosophila*,

90   *Tribolium*, and *Parhyale* (Fig. S1). However, only one of these was expressed in *Acanthoscurria*

embryos at the stages examined and was therefore presumed to be the conserved paralog of *pnr*.

Additionally, OrthoFinder[59] identified several Iroquois genes, but only one best *Acanthoscurria*

araucan gene, and one best *Acanthoscurria Distal-less* (*Dll*) which were the closest to their

respective orthologs in *Drosophila*, *Tribolium*, and *Parhyale*.

95   In order to facilitate the following comparisons, the names of each leg segment in insects,

crustaceans, and chelicerates have been replaced with numbers 1 – 8, counting from the terminal

claw (i.e. pretarsus, dactyl, etc.), as in Bruce and Patel 2020[24], but the reader may refer to Fig. 1

for quick look-up. For example, the trochanter of chelicerates, the basis of crustaceans, and the

coxa of insects are leg segment 6. For the insect representative used here, *Tribolium* (with has 6

100   moveable leg segments), the boundaries of the embryonic leg segments 7 and 8 that form the

adult lateral body wall (pleura and lateral tergum, respectively) are based on molecular and

morphological analyses in Kobayashi 2022[37], as well as on the extensive literature regarding the

"insect subcoxa theory", such as the location of the paracoxal suture in Matsuda 1970[36],

Kobayashi 2013[34], 2017[35], and 2018[60].

105   In *Parhyale*, *Tribolium,* and *Acanthoscurria, Dll* was found to be expressed in leg

segments 1 – 5 and *pnr* expressed in the most dorsal tissue (Fig. 2). Thus, in all arthropods for

which there is data, including the insects *Drosophila*[39,43,55,61,62], *Tribolium*[24], and *Bicyclus*[63]; the

crustaceans *Parhyale*[24], *Triops* (unpublished), and *Daphnia*[44]; the millipede *Glomeris*[64]; the

chelicerate *Parasteatoda* (called GATA1)[65]; and here the chelicerate *Acanthoscurria*, *pnr* marks

110   the "true" body wall (medial tergum) that is not derived from the leg base. In all arthropods

examined to date, *ara* is expressed in four possible domains: a dorsal armband on proximal leg

segment 8 that is adjacent to the *pnr* domain; a second armband at the joint between leg

segments 7 and 8; a circular patch of expression on the lateral side of leg segment 6; and in the

distal tip of leg segment 1. The crustaceans *Parhyale* and *Daphnia* express all four domains of

115   *ara*[24,44]. In *Tribolium*, *ara* is expressed in the first three of these domains but not in the distal tip

of the leg. In *Acanthoscurria*, *ara* is expressed in three of these domains: the dorsal armband

adjacent to the *pnr* domain; the second armband on proximal leg segment 7; and in the distal tip

of the leg. The lack of the lateral dot of *ara* in *Acanthoscurria* legs may be connected to the

evolution of exopods within the pancrustaceans. Thus, as predicted by the leg segment alignment

120   model in Bruce and Patel 2020 for insect and crustacean legs[24], the two armbands of *ara*

expression in *Acanthoscurria* bracket a region proximal to leg segment 7 (coxa) and adjacent to

pnr. This suggests that *Acanthoscurria,* like *Parhyale*, *Daphnia*, and *Tribolium*, did not delete

the ancestral, proximal 8th leg segment, but rather converted it into the body wall.

   To further test this hypothesis, the expression of *drumstick* was examined in

125   *Acanthoscurria*. *Drumstick* is in the *odd-skipped* family of genes, which are expressed in the

distal edge of each leg segment in all arthropods examined, including in *Parhyale*, *Tribolium*,

and the chelicerate spider *Parastatoda*[63,66–69]. In particular, *drumstick* is expressed in a sharp line

and only in the true (eudesmatic) leg joints, and not in the leg subdivisions that provide flexion

such as those of the tarsus. Thus, expression of *drumstick* can be used to count the number of

130   true leg segments present, and to molecularly distinguish true leg segments from superficial

subdivisions. In *Drosophila*, all members of the *odd-skipped* family of genes – *odd-skipped (odd), brother of odd with entrails limited (bowl), sister of odd and bowl (sob), and drumstick (drm)* – have been shown to have similar overlapping expression patterns in the leg joints and to variably induce cells to buckle and form a flexible joint[70]. OrthoFinder[59] identified a gene in

135 *Acanthoscurria* and in *Parhyale* which were each the reciprocal best BLAST hit to *drm* in *Drosophila* and *Tribolium*. This *Acanthoscurria drumstick* was the reciprocal best BLAST hit to *Cupiennius* (spider) *odd-related* 3 (*odd-r3*)[66]. As in other arthropods, *Parhyale drm* is expressed in 7 stripes, one on the distal side of each of the 8 leg segments, denoting the future location of leg joints (Fig. S4)[67,68]. Similarly, in *Acanthoscurria, drm* is expressed in the expected distal

140 regions of leg segments 1 – 7, but also in an additional ring proximal to leg segment 7 (Fig. 2G, H). This additional ring of *drm* notably occurs on the distal side of a leg-segment-like bulge, as expected if this bulge represents an embryonic 8th leg segment. This leg-segment-like bulge is visible in other studies and other spider species, including *Acanthoscurria* embryos in Pechmann 2009[71] Figure 3 and Pechmann 2020[72] Figure 6; as well as in *Cupiennius* in Pechmann 2010

145 Figure 16, and Wolff 2011 Figures 11 and 12. Given that *drm* specifically marks the distal side of leg segments[70], the ring of *drm* expression on the distal side of the leg-segment-like bulge suggests that it is indeed a leg segment. Together, the conserved expression of *pnr*, *ara,* and *drm* surrounding the embryonic leg-segment-like bulge suggest that *Acanthoscurria* has an additional, cryptic proximal 8th leg segment.

150 In all arthropods examined, Iroquois genes are expressed strongly around proximal leg segment 8, with accessory expression domains, like the lateral dot and the tip of the claw, being variable in different arthropod lineages. However, the conserved expression configuration is *pnr*

in a dorsal-medial stripe, which is apposed to a stripe of *iro*, with a second stripe of *iro* between

leg segments 7 and 8.

Discussion

The expression of *pnr*, *ara*, and *drm* reveals how the proximal leg segments of insects,

crustaceans, and chelicerates correspond to each other, which supports the hypothesis that

arachnids with 7 moveable leg segments possess an embryonic proximal 8th leg segment that

160   forms the lateral body wall in the adult. This is consistent with all available molecular,

embryological, and morphological data in chelicerates, as follows.

To align the distal leg segments in insects, crustaceans, and chelicerates, we turn to the

published molecular loss-of-function literature. Four leg patterning genes have been functionally

tested in representatives for insects, crustaceans, and chelicerates: *Distal-less* (*Dll*), *Sp6-9*,

165   *dachshund* (*dac*), and *homothorax* (*hth*). We note that the expression of these leg patterning

genes is quite variable, both between arthropod phyla as well as over the embryonic development

of each individual species[24,73]. For example, *Dll* is expressed in regions where it does not have a

leg phenotype (early proximal leg[74,75] and wing sensory hairs[76]), and conversely, *dac* is not

expressed in regions where it does have a phenotype (leg segment 3, but presumably *dac* is

170   expressed here earlier). Thus, the expression of these four genes is not as reliable as gene

function when comparing such phylogenetically distant groups, and can confound

interpretations. This is in contrast to the expression of *pnr*, *ara*, and *drm*, which – at the stage

when leg joints are forming – are surprisingly well-conserved in their expression domains

despite half a billion years of divergence.

175   We therefore focus on the loss-of-function data for *Dll*, *Sp6-9*, *dac*, and *hth* more than expression. Furthermore, given that RNAi produces a range of moderate to strong knockdowns, we focus on what appear to be strong phenotypes. For the following descriptions of leg segment deletion phenotypes, all interpretations are taken directly from the source material. For example, Setton and Sharma[77] state that *Sp6-9* RNAi in spider results in "truncation of all remaining

180   appendages up to the proximal-most leg segment, the coxa", which in the present nomenclature is leg segment 7. In several cases, the respective authors were also contacted to confirm the interpretations here (Prashant Sharma 2025; Natascha Turetzek 2022; Mathias Pechmann, July 2022; Graeme Mardon, 2020; Carlos Estella 2018). Furthermore, the respective authors' conclusions were visually confirmed from their original figures based on the shape of each leg

185   segment. For example, spider leg segment 7 (coxa) is short and has a medial bulge called the coxal endite. This endite is especially pronounced on the pedipalp, which has the same number of leg segments as walking legs[78] making the pedipalp endite a useful morphological marker for interpreting phenotypes. The interpretations from the source material are applied to the best of our ability in demarking leg segment boundaries in the figures reproduced here.

190   Based on the leg segment deletion phenotypes of *Dll*, *Sp6-9*, *hth*, and *dac,* the six distal leg segments of *Parhyale*, insects, and chelicerates (leg segments 1 – 6, counting from the distal claw) correspond to each other in a one-to-one fashion (Figs. 3, 4), as follows.

        Dll RNAi in chelicerates deletes distal leg segments 1 – 5[79–81] (Fig. 3D, and personal communication, Mathias Pechmann, July 2022), the same segments in which *Dll* is expressed

195   when the embryonic leg joints become visible. The *Dll* RNAi phenotype is confirmed morphologically by the presence of the coxal endite in the leg stump, and molecularly by the expression in the stump terminus of *dac1*, a marker of leg segment 6[71,77,82–84], the next most

proximal leg segment. Thus, *Dll* is required for the development of leg segments 1 – 5 in chelicerates, a crustacean, and insects (Fig. 3A – F)[24,79,85–90].

200   *Sp6-9* RNAi in chelicerates deletes distal leg segments 1 – 6[77,82] (Fig. 3J). This is confirmed morphologically by the presence of the coxal endite on the pedipalp stump, indicating that leg segment 7 is still present, and molecularly by the lack of *dac1* expression, a marker of leg segment 6, in the stump, despite *dac1* expression in the ventral nerve cord[77,82]. Thus, *Sp6-9* is required for the development of leg segments 1 – 6 in chelicerates, a crustacean, and insects (Fig.

205   3G – L)[24,77,82,91–93].

For *dac*, the available data is sparse and less clearcut, but suggests that *dac* patterns middle leg segments 3 – 5 (Fig. 4D-F). The strongest *dac* functional data for a chelicerate is in the harvestman *Phalangium opilio*[78]. Similar, but more moderate, effects are also seen in the spider, *Parasteatoda*[83], where only one of the two *dac* paralogs gives an observable phenotype

210   (Natascha Turetzek Zhang, personal communication). In the harvestman *Phalangium*, RNAi against the single copy of *dac* affects leg segments 3 – 5, such that segments 4 and 5 are deleted, and segment 3 is truncated (Fig. 4D). The crustacean, *Parhyale,* has two *dac* paralogs: *dac1* is more strongly expressed in a neural pattern and *dac2* is more strongly expressed in the medial leg[94]. Consistent with this expression difference, only *dac2* in *Parhyale* gives a discernible

215   phenotype; *dac1* may have a neural phenotype but this was not observable from cuticle preps. This is similar to the situation in spider, where only one of the two *dac* paralogs gives an observable phenotype, which suggests that one *dac* paralog in *Parasteatoda* and *Parhyale* may retain the medial leg patterning while the other controls other aspects of patterning. In *Parhyale*, CRISPR-Cas9 knockout of *dac2* completely deletes leg segments 3 – 5, such that leg segment 2

220   connects directly to 6[24], even though *dac* is not clearly expressed in segment 3 at the stages

examined[24]. In the insect *Drosophila*, loss of *dac* also affects leg segments 3 – 5 (as well as the

proximal part of 2), which are shorter and fused into a contiguous mass of tissue[95]. RNAi against

dac in the insect *Oncopeltus* deletes leg segment 3 and shortens leg segment 4[89], and RNAi

against *dac* in *Tribolium* deletes leg segments 3 and 4, as well as the proximal part of 2[96]. Thus,

225 *dac* appears to be required for the development of leg segments 3 – 5 in a chelicerate, a

crustacean, and insects (Fig. 4A – F), but insect *dac* function appears to have shifted slightly

distally[24,78,83,89,95,96,99]. Strong CRISPR-Cas9 phenotypes in chelicerates, myriapods, other

crustaceans, and non-*Drosophila* insects would clarify this ambiguity.

For *hth*, the only chelicerate for which there is functional data is the harvestman,

230 *Phalangium opilio*. In *Phalangium*, RNAi against *hth* shortens and fuses the proximal leg

segments, leaving the distal segments unaffected[97]. Given that the most severely affected

embryos died before hatching, their wrinkled cuticle obscured the joints, making it difficult to

discern with high certainty how many distal leg segments are unaffected. However, in the

normal-looking distal part of the leg, leg segment 1 and the joint its shares with leg segment 2

235 are present, so the distal half of leg segment 2 appears unaffected. Thus, *hth* is required for the

development of the proximal leg segments and leaves distal leg segments 1 and part of intact 2 in

arachnids as well as *Parhyale* and insects (Fig. 4H – J)[24,89,95,97–100].

The functional data for *Dll*, *Sp6-9*, *dac*, and *hth*, together with the conserved expression

patterns of *pnr*, *ara*, and *drm*, support a model wherein living arthropods ancestrally had 8 leg

240 segments, and that the reduced number of canonical leg segments in chelicerates, myriapods,

non-insect crustaceans, and insects appears to be due to the conversion of proximal embryonic

leg segments into lateral body wall in the adult.

The molecular data discussed here agrees with over a century of morphological, embryological, and paleontological observations suggesting that arthropods ancestrally had 8 leg segments. First, living arthropods have no more than 8 true (eudesmatic) leg segments posterior to the first antenna[2]; early-branching chelicerates[48,49,104] often have 8 leg segments[4]; many crustaceans have an additional proximal leg segment, known as the precoxa[2,3,10], corresponding to leg segment 8 here; and the fossil ancestors of all living arthropods had 8 leg segments[46]. Furthermore, in arthropods with fewer than 8 moveable leg segments, the embryonic leg base is observed to broaden and expand to form the adult lateral body wall[34,35,37,47,105,106], suggesting that the arthropod ancestor had more leg segments than the present number of moveable leg segments would indicate. Similarly, pleurites (hard plates floating in the lateral body wall to which muscles attach, often surrounded by soft cuticle) are found only in arthropods with fewer than 8 free leg segments (Fig. 6b), including insects[8,34,35,51,107], decapods (crustaceans)[1,25,108,109], millipedes[28,110], centipedes[1], and pauropods[105] (myriapods), and horseshoe crabs (chelicerate)[1]. Notably, pleurites appear to be derived from the embryonic leg base[1,34,105,107], thus accounting for the missing ancestral leg segments in these lineages. These morphological, embryological, and fossil observations have led previous authors to a similar conclusion within each arthropod lineage[4,8,10,46,51]. These are here combined with the added insight of modern molecular phylogenies[48,49,111,112] and gene expression and function data to infer that all arthropods ancestrally had 8 leg segments that correspond to each other in a one-to-one fashion (Fig. 6). For example, the insect coxa, crustacean basis, and spider trochanter correspond to each other, such that a simpler term for these would be "leg segment 6", while the insect trochanter, crustacean ischium, and spider femur could be referred to as leg segment 5.

265   The leg model presented here also explains another peculiarity: a vestigial leg joint within the adult insect lateral body wall. Arthropod leg joints can be defined molecularly by the expression of joint patterning genes like *odd-skipped* family genes[53,68,69,101,102] and *Serrate*[37,103]; and morphologically by the presence of a thin, flexible line of cuticle (called a suture) with muscle insertions[2,36]. The vestigial leg joint present in the lateral body wall of adult insects

270   satisfies both of these criteria: *odd-skipped* genes and *Serrate* are expressed in a ring here in embryos, and this region forms a morphological suture known as the paracoxal suture in the adults of many insect lineages onto which muscles insert[35–37], which represents the vestigial joint between leg segments 7 and 8 that forms the lateral body wall of the adult insect (Fig. 5b).

       The differences in the canonical leg segment numbers in the four groups of arthropods

275   are proposed to be due to the variable conversion of proximal leg segments into the body wall, likely to support more of the body weight on the legs during the transition from swimming to walking and from aquatic to terrestrial lifestyles. Chelicerates with 7 free leg segments, like spiders, have converted one leg segment into the body wall (leg segment 8), whereas chelicerates with 8 free leg segments, like sea spiders, camel spiders, and red velvet mites, appear to retain

280   the proposed ancestral number of 8 free and mobile leg segments[4]. Myriapods with 6 or 7 free leg segments[1,2,7,28,52,110,113,114] appear to have converted 2 or 1 leg segments into the body wall, respectively[105]. Insects, with 6 free leg segments, have converted two proximal leg segments into the body wall (leg segments 7 and 8). Crustaceans have variably converted the leg base into the lateral body wall. Many early-branching crustaceans[111,112], like branchiurids, mystacocarids and

285   myodocope ostracods, retain a free and mobile leg segment 8 (precoxa)[2,3,10,115–118]. In decapods like prawn and crab, the embryonic/ancestral precoxa forms the adult lateral body wall, such that the pleurobranch gills (and probably arthrobranch gills) that are carried on the adult lateral body

wall[23,119] are derived from the embryonic precoxa[10,108,109,119,120]. In amphipods like *Parhyale*, the embryonic precoxa forms much of the lateral body wall, but is obscured by the tergal and coxal plates[24]. However, the precoxa is revealed when wing genes like *vestigial* are knocked out, which deletes the tergal and coxal plates, leaving behind just the exposed, protruding precoxa[16]. Terrestrial isopods with 6 free leg segments have, like insects, converted two proximal leg segments into the body wall (leg segments 7 and 8), an evolutionary sequence which is easier to appreciate in this lineage because aquatic isopods still retain a free and mobile leg segment 7[121–124]. Mantis shrimp possess a free and mobile precoxa but only on the posterior three thoracic legs; the powerful raptorial legs do not have a free precoxa[3,125–127], which may have converted the precoxa into the body wall in order to provide a more robust base from which to exert force. Cladoceran crustaceans like the water flea *Daphnia* have partially converted the precoxa into the lateral body wall[10].

Insect embryologists since the late 1800s, working in several major clades of insects, have observed that the embryonic leg base flattens and expands to form the lateral body wall of the adult[8,34,36,37,47,54,128–131]. This has become known as the insect subcoxa theory, likely familiar to any entomology student. The analysis presented here suggests that this same morphogenetic sequence happens throughout arthropods, meaning that the insect subcoxa theory should more generally be called the arthropod subcoxa theory.

A system to decipher the origins and relationships of any arthropod outgrowth

In the model presented here, the function of the five leg patterning genes in conjunction with the expression of the proximal patterning genes *pnr* and *ara* and the joint marker *drm* can be used to infer the homologies of structures along the proximal-distal axis in the same way that

the well-known "Hox code" has been used to infer homologies along the anterior-posterior axis. Notably, given the highly conserved expression patterns of *pnr*, *ara*, and *drm* across arthropods, and the ease and low cost of in situ Hybridization Chain Reaction (HCR)[132,133] expression experiments, the homologies of any and every long-debated arthropod ectodermal structure can

315   now be elucidated with a single in situ HCR experiment using just three or four genes. For example, a simple four-gene HCR of *pnr, ara, drm*, and a wing gene like *vestigial* could answer definitively whether beetle horns and treehopper helmets are wing serial homologs that are carried on embryonic/ancestral leg segment 8. Similarly, a simple three-gene HCR of *pnr*, *ara*, and *drm* could answer definitely whether all "entomostraca"-like crustaceans truly lack

320   abdominal legs[134], or have simply truncated them as insects have[135]. Additionally, if insects have 6 free and 2 converted leg segments, such that *odd-skipped* genes are expressed in a stripe at the vestigial joint between segments 7 and 8 that form the lateral body wall, then other arthropods with 6 free and 2 converted leg segments should also express *odd-skipped* genes in a stripe at this vestigial joint, such as isopod crustaceans, and symphylan and pauropod

325   myriapods[2,52,105,113,114,121,122]. Furthermore, the expression of *pnr* and *ara* has thus far only been examined in arthropods that have partially or completely converted leg segment 8 into the lateral body wall. The expression of these genes should therefore be examined in arthropods where leg segment 8 is still free and mobile, such as sea spiders (Fig. 5), camel spiders (solifugids), red velvet (trombidiid) mites, and mantis shrimp (leg segment 8/precoxa is free[125–127], but the distal

330   leg segment number is reduced; however, leg segment 6/basipod is easily identifiable morphologically by the presence of the exopod[3,125,126]). If this model is accurate, then *pnr* should be expressed in the true body wall (perhaps only dorsally but not ventrally), *ara* should bracket the moveable 8th leg segment proximally and distally, and *odd-skipped* genes should only be

expressed where two leg segments meet, but not where the proximal 8th leg segment meets the

335   body wall. Similarly, a prediction of this model is that lateral outgrowths like wings, plates, gills,

carapaces, and horns should only form on leg-derived tissue (either free and moveable leg

segments, or on the leg-derived body wall), but not on the non-leg derived "true" body wall. A

related prediction then is that arthropods that retain 8 moveable leg segments, like sea spiders,

camel spiders (solifugids), red velvet (trombidiid) mites, and mantis shrimp, should not have

340   outgrowths of their body wall, which seems to be the case. While the mantis shrimp may seem to

contradict this prediction in that it has both a free precoxa and a carapace, it in fact proves the

point: the legs that possess a moveable precoxa (the three posterior thoracic legs) do not form the

carapace; the carapace is formed by the anterior legs that do not have a free precoxa[3,125,136].

345   **Determining the leg segment affinities of chelicerae**

An example of how this leg coordinate system can be put to work with a simple HCR

experiment is the chelicerae (anterior-most legs of chelicerates). Chelicerae are the namesake of

the chelicerates, and represent the appendages of the deuterocerebrum that correspond to the

antennae of insects, the first antennae (antennules) of crustaceans[137–140], and perhaps the great

350   appendages of fossil megacheiran arthropods[141]. In spiders, the chelicerae form fangs, but in sea

spiders, horseshoe crabs, scorpions, and mites, they form pincer claws. Chelicerae have been

proposed to have been 3-segmented in the chelicerate ancestor, but that the proximal segment

was lost in more derived spiders with 2-segmented chelicera[142,143]. The expression of *pnr, ara,*

drm, and *Dll* in *Acanthoscurria*, an early-branching spider, supports this hypothesis in that the

355   proximal-most leg segment 8 appears to form the body wall of the head, as follows (Fig. S3).

pnr is expressed dorsally around the head lobes, likely marking the non-leg-derived ancestral

body wall. *ara* is expressed in the same three domains as the other legs of *Acanthoscurria*: the dorsal armband that follows the *pnr* domain, the second armband, and the distal claw domain in the fang. *drm* is expressed in two bands, indicating the boundaries of three leg segments. But

360 which of the eight possible leg segments in particular? The proximal-most segment likely corresponds to leg segment 8, because it is bracketed by the dorsal *ara* armband that follows *pnr* and the ventral *ara* armband, which is also adjacent to the *drm* band, indicating the distal end of the segment. The distal-most segment of the chelicera likely corresponds to leg segment 1 (the claw) because of the distal, sock-like expression of *Dll* as well as *ara* in the fang. The identity of

365 the medial segment is more ambiguous. It expresses *Dll*, and therefore could correspond to leg segments 2, 3, 4, or 5. However, a comparison of *dac* expression in spiders narrows the possibilities because *dac* is co-expressed with *Dll* only in leg segment 5 (femur)[73,83,84,143].

Myriapod legs

370 Unfortunately, no gene functional data is available for myriapods, and in myriapod embryos, the non-specific background staining becomes too high after the early leg bud stages when the leg joints form, which precludes relating leg gene expression domains to specific morphological leg segments. However, the available gene expression data for early leg patterning is consistent with the model presented here.

375 In all arthropods including myriapods, *Dll* is expressed in distal sock-like domains of the mouthparts and legs[144–146], corresponding to leg segments 1-5. Consistent with the model presented here, the mandible of mandibulates (myriapods, crustaceans, and insects) appears to lack segments 1-5 (the telopod), and consist of only the two proximal-most leg segments, 7 and 8[68,77,145–147], with segment 8 latter forming the head lobes and conglomerated walls of the head,

380 and segment 7 forming the moveable part of the mandible. For example, *Dll* is expressed only in

the tooth-like structure of the mandible, and appears to have a sensory role rather than patterning

the actual leg-segments of the mandible[24,68,94,145–147]. Similarly, in all arthropods including

myriapods, *exd* and *hth* are expressed in proximal domains, including in the mandible[144,148],

whereas *dac* is expressed in a medial domain in non-truncated legs, but in the distal tip of the

385 mandible and maxilla[144]. *dac* expression in the mandible likely represents endite patterning[106,144],

similar to *Dll* expression here, while dac expression in the maxilla may correspond to endite

patterning or to leg segment 5 in these distally truncated leg types[94,149].

Furthermore, the expression domains of axial patterning genes, which also pattern legs,

are highly conserved across all arthropods, including myriapods[150,151]. For example, *engrailed*

390 and *hedgehog* are expressed in the posterior of each leg, while *cubitus interruptus* is expressed in

the anterior[152–154]. Similarly, *wingless* is expressed in a ventral stripe in each leg, while

decapentaplegic and *optomotor blind* are expressed dorsally in each leg[153,155,156].

No data is available for the expression of *pnr* or any of the Iroquois genes like *ara* in any

myriapods, which would facilitate the molecular identification of the proximal leg segments.

395 However, the morphological and embryological data suggests that in myriapods, like insects,

many crustaceans, and spiders, the embryonic leg base forms the adult body wall[2,105,110]. In

embryos of a myriapod representative, the symphylan *Hanseniella*, the proximal part of the

developing leg ("limb base") is observed to broaden and flatten to form the adult lateral body

wall[105] . Incorporation of proximal leg segments into the myriapod body wall would bring their

400 leg segment count to 8, in agreement with other arthropods.

A model and simple 3-gene test to decipher the origins and relationship of any

ectodermal structure in any arthropod opens up a powerful system for studying the origins of

novel structures, the evolvability of morphology and gene networks across vast phylogenetic

distances, and the convergent evolution of shared ancestral developmental fields.

410 **Materials and Methods**

<u>Embryo fixation</u>
Acanthoscurria embryos were fixed overnight in 5% formaldehyde/PBS and heptane. *Tribolium* embryos were fixed 30 min in 4% formaldehyde/PEM and heptane. *Parhyale* embryos fixed 40
415 min in 4% PFA in or artificial seawater. All embryos then dehydrated into 100% methanol. All embryos stored at -20°C.

<u>In situ HCR version 3.0</u>
In situs were performed as in Bruce et. al. 2021[133]. cDNA sequences were submitted to
420 Molecular Instruments[132], and the probe sets are available from the company, as below. cDNA sequences for Acanthoscurria *pnr, ara, Dll*, and *drm* Supplemental information. Probe sequences for the three *pannier* paralogs ordered as o-pools from IDT in Supplemental information.

Species	Gene	Accession number	Probe set numbers from Molecular Instruments
Acanthoscurria geniculata	araucan	-	PRE883
Acanthoscurria geniculata	*Distalless*	-	PRE884
Acanthoscurria geniculata	drumstick	-	PRG135
Tribolium castaneum	pannier	XM_008202266.2	PRD736
Tribolium castaneum	araucan	XM_008194186.2	PRD737
Tribolium castaneum	*Distalless*	NM_001039439.2	PRD738
Parhyale hawaiensis	pannier	MT103930.1	PRD223
Parhyale hawaiensis	araucan	MT103931.1	PRD221
Parhyale hawaiensis	*Distalless*	FJ617569.1	PRD002

 Imaging

Embryos imaged with Zeiss LSM880 confocal. Image processing done with Fiji-ImageJ. Fiji "Image Calculator > Subtract" method was used to remove high background from yolk autofluorescence. Figures processed using Adobe Photoshop 2020.

430 Gene orthology

OrthoFinder and gene phylogenies were used to determine orthology of Acanthoscurria *pannier*, *araucan*, and *drumstick*. *Acanthoscurria* transcriptome, kindly provided by Mathias Pechman, and *Parhyale* transcriptome from Sun 2022[157] were each first converted to proteomes using Transcdecoder.LongestORFs[158], then Transdecoder.Predict with the single_best_orf option

435 and blastp options using the Drosophila Reference Sequences proteome (NCBI), Tribolium Reference Sequences proteome (NCBI) to identify the single best ORF for each transcript. The resulting *Acanthoscurria* and *Parhyale* proteomes were then used with OrthoFinder against the above proteomes for *Drosophila* and *Tribolium* (which each contain the specific sequences of *pnr*, *ara*, and *drm* that have been previously verified by in situ to give the expected expression

440 pattern), to identify *pnr*, *ara*, and *drm* in *Acanthoscurria* and *Parhyale*.

The resulting Acanthoscurria proteome was also used to generate phylogenies for pnr and ara. For ara, the following sequences with accession numbers were used to generate sequence alignments, with the closely related Mohawk gene[159] as the outgroup:

NP_808263.2_Mohawk_Mus, XP_001811513.2_Mohawk_Tribolium,
445 NP_608502.1_CG11617_Mohawk_Drosophila, XP_008192408.1_araucan_Tribolium,
XP_008192405.1__araucan_Tribolium, NP_001163432.1_araucan_Drosophila,
KYB28757.1_araucan_Tribolium, KYB28756.1_araucan_Tribolium,
XP_008192400.1__araucan_Tribolium, AAT11862.1_iroquois_Tribolium,
AAB53640.1_mirror_Drosophila, XP_971676.1__caupolican_Tribolium,
450 KZS08873.1_caup_Daphnia_magna, NP_524046.2_caup_Drosophila, XP_971676.1__caup_Tribolium,
XP_016770700.1__caup_Apis, KFM58187.1_caup_Stegodyphus, NP_524045.2_araucan_Drosophila,
NP_001163432.1_araucan_Drosophila, XP_015182907.1__araucan_Polistes_dominula,
XP_018911022.1__ara_Bemisia_tabaci, JAG00978.1_araucan_Lygus_hesperus,
ODM99984.1_ara_Orchesella_cincta, XP_008192408.1__araucan_Tribolium,

455   XP_008192405.1_araucan_Tribolium, XP_008192400.1_araucan_Tribolium,
EKC34186.1_araucan_Crassostrea, NP_524047.2_mirror_Drosophila, EHJ65533.1_mirror_Danaus,
JAB79499.1_mirror_Ixodes, JAB72752.1_mirror_Ixodes, XP_015907356.1_caup_Parasteatoda,
EEC00349.1_IRX_Ixodes, XP_015782374.1_IRX-3_Tetranychus,
KFM58187.1_caupolican_Stegodyphus, XP_002400564.1_IRX_Ixodes_scapularis,
460   KZS08873.1_caup_Daphnia_magna, XP_021002308.1_IRX-6_Parasteatoda, XP_021002310.1_IRX-
6_Parasteatoda, XP_021002309.1_IRX-6_Parasteatoda, KZS08871.1_IRX-1_Daphnia_magna,
Parhyale.ara.phaw_50.282976aG352.1

For pnr, the following sequences with accession numbers were used to generate sequence

465   alignments, with *Caenorhabditis elegans* elt-1 as the outgroup:

Hyalella_grn_XP_018013183.1, Parhyale_grain_phaw_50.000203bG892.1,
Trinorchestia_grn_KAF2359614.1, Acanthoscurria_DN76703_c4_g1_i1,
Acanthoscurria_DN74902_c0_g1_i1, Acanthoscurria_DN81267_c7_g2_i1,
Stegodyphus_GATA4_KFM82098.1, Parasteatoda_GATA4_XP_015929395.1,
470   Acanthoscurria_DN76525_c1_g1_i1, Euperipatoides_GATA123_CUW78639.1,
Limulus_grn_XP_022242144.1, Limulus_GATA2_XP_022240403.1,
Limulus_GATA2_XP_013773452.2, Euperipatoides_GATA456_CUW78640.1,
Stegodyphus_GATA-4_KFM82100.1, Stegodyphus_pnr_XP_035228380.1, C_elegans_elt-
1_NP_001368482.1, Tigriopus_grn_TRY70078.1, Stegodyphus_grn_XP_035208929.1,
475   Parasteatoda_grn_XP_021003356.1, Trinorchestia_pnr_KAF2355501.1,
D.magna_grn_XP_032783972.1, Eurytemora_grn_XP_023338271.1,
Penaeus_grn_XP_027213737.1, Penaeus_grn_XP_027213731.1,
Parhyale_grain_comp168149_c1_seq6, Tribolium_grain_NP_001158260.1,
Drosophila_grain_NP_001262366.1, Centruroides_GATA-4_XP_023233756.1,
480   Eurytemora_grn_XP_023340853.1, Penaeus_pnr_XP_027229190.1,
Acanthoscurria_DN55779_c1_g1_i1, Hyalella_pnr_XP_018008986.1,
Parhyale_pannier_comp167878_c4_seq4, Ixodes_GATA-5_XP_029839700.1,
Limulus_pnr_XP_013775036.1, Limulus_GATA-4_XP_022242593.1,
Ixodes_pnr_XP_029842940.1, Parasteatoda_GATA-A_BBD75256.1,
485   Parasteatoda_LOC107456523_XP_015929888.1,
Parasteatoda_LOC107456523_XP_015929890.1, Ixodes_GATA-5_XP_029840357.1,
Acanthoscurria_DN77434_c10_g2_i1, Acanthoscurria_DN77434_c10_g2_i2,
Centruroides_pnr_XP_023233757.1, Centruroides_grn_XP_023227752.1,
Ixodes_grn_EEC05487.1, Parasteatoda_LOC107439679_XP_015907840.1,
490   Parasteatoda_GATA-B_BBD75257.1, Acanthoscurria_DN80076_c8_g1_i4, Limulus_GATA-
B_XP_022249933.1, Acanthoscurria_DN78099_c0_g1_i1, Acanthoscurria_DN77604_c0_g1_i1,
Centruroides_GATA-4_XP_023228374.1, Stegodyphus_GATA-5_KFM72207.1,
Stegodyphus_GATA-A_KFM57600.1, Parasteatoda_pnr_XP_015929399.1,
D.pulex_pnr_EFX78972.1, D.magna_GATA_KZS08761.1, Bombyx_pnr_QGA69977.1,
495   Tribolium_serpent_XP_008200495.1, Tigriopus_pnr_TRY71481.1, Glomeris_pannier,
Nasonia_pnr_XP_001607848.1, Tribolium_pannier_XP_008200486.1,
Ontho.sag_pnr_QDE10346.1, Drosophila_pannier_NP_001262620.1,
Apis_pnr_XP_001121210.2, Ontho.taur_pnr_XP_022901096.1

500   Sequence alignments were generated using clustal[160], mafft[161], and muscle[162]. Best model was selected using ProTest[163]. Trees were then generated in Geneious[164] with PHYML[165] and MrBayes[166] using the recommended model from ProtTest. Trees visualized with TreeView[167]

Data and materials availability: All data is available in the main text or the supplementary
505 materials.

Competing interests: Authors declare no competing interests.

510 **Acknowledgments:** HSB thanks Nipam H. Patel for reagents, space, and equipment for this work. HSB thanks Carsten Wolff, Rachel Thayer, Joanna Wolfe, Gregory Edgecomb, and Frank W. Smith for helpful feedback. Fixed *Acanthoscurria* embryos in methanol, *Acanthoscurria* transcriptome, and unlabeled images of Dll RNAi spider embryos were kindly provided by Mathias Pechmann (Institute for Zoology/Developmental Biology, Biocenter, University of
515 Cologne, Germany). HSB thanks Professor Yukimasa Kobayashi for analysis of *Drosophila* musculature to determine the location of the paracoxal suture boundary between leg segment 7 and 8 (Kobayashi's subcoxa 2 and 1, respectively). HSB thanks Aaron Vose for computational assistance.

520 **Author contributions**
HSB conceived, designed, and performed experiments, and wrote manuscript.

References

1. Snodgrass, R.E. (1952). A textbook of arthropod anatomy. (Ithaca, N.Y., Comstock Pub. Associates).

2. Boxshall, G.A. (2004). The evolution of arthropod limbs. Biol. Rev. *79*, 253–300. https://doi.org/10.1017/S1464793103006274.

530   3. Schram, F.R. (1986). Crustacea (Oxford University Press, USA).

4. Shultz, J.W. (1989). Morphology of locomotor appendages in Arachnida: evolutionary trends and phylogenetic implications. Zool J Linn Soc *97:1-56*.

5. Grimaldi, D., and Engel, M.S. (2005). Evolution of the Insects (Cambridge University Press).

535   6. Crampton, G. (1916). The phylogenetic origin and the nature of the wings of insects according to the paranotal theory. J NY Entomol. Soc. *24*, 1–39.

7. Brusca, R.C., and Brusca, G.J. (2003). Invertebrates (Sinauer Associates Incorporated).

8. Snodgrass, R.E. (1927). Morphology and mechanism of the insect thorax (City of Washington, Smithsonian Institution).

540   9. Sharov, A.G. (1966). Basic Arthropodan Stock: With Special Reference to Insects 1st ed. (Pergamon Press).

10. Hansen, H.J. (1925). Studies on Arthropoda II. (Copenhagen: Gyldendalske Boghandel).

11. Størmer, L. (1944). On the relationships and phylogeny of fossil and recent Arachnomorpha.

12. Damen, W.G.M., Saridaki, T., and Averof, M. (2002). Diverse adaptations of an ancestral
545   gill: a common evolutionary origin for wings, breathing organs, and spinnerets. Curr. Biol. *12*, 1711–1716.

13. Averof, M., and Cohen, S.M. (1997). Evolutionary origin of insect wings from ancestral gills. Nature *385*, 627–630. https://doi.org/10.1038/385627a0.

14. Kukalová-Peck, J. (1983). Origin of the insect wing and wing articulation from the
550   arthropodan leg. Can. J. Zool. *61*, 1618–1669. https://doi.org/10.1139/z83-217.

15. Niwa, N., Akimoto-Kato, A., Niimi, T., Tojo, K., Machida, R., and Hayashi, S. (2010). Evolutionary origin of the insect wing via integration of two developmental modules. Evol. Dev. *12*, 168–176. https://doi.org/10.1111/j.1525-142X.2010.00402.x.

16. Clark-Hachtel, C.M., and Tomoyasu, Y. (2020). Two sets of candidate crustacean wing
555   homologues and their implication for the origin of insect wings. Nat. Ecol. Evol. *4*, 1694–1702. https://doi.org/10.1038/s41559-020-1257-8.

17. Prud'homme, B., Minervino, C., Hocine, M., Cande, J.D., Aouane, A., Dufour, H.D., Kassner, V.A., and Gompel, N. (2011). Body plan innovation in treehoppers through the evolution of an extra wing-like appendage. Nature *473*, 83–86. https://doi.org/10.1038/nature09977.

18. Hu, Y., Linz, D.M., and Moczek, A.P. (2019). Beetle horns evolved from wing serial homologs. Science *366*, 1004–1007.

19. Linz, D.M., and Tomoyasu, Y. (2018). Dual evolutionary origin of insect wings supported by an investigation of the abdominal wing serial homologs in Tribolium. *115*, E658–E667. https://doi.org/10.1073/pnas.1711128115.

20. Zattara, E.E., Busey, H.A., Linz, D.M., Tomoyasu, Y., and Moczek, A.P. (2016). Neofunctionalization of embryonic head patterning genes facilitates the positioning of novel traits on the dorsal head of adult beetles. Proc. R. Soc. B *283*, 20160824. https://doi.org/10.1098/rspb.2016.0824.

21. Almudi, I., Vizueta, J., Wyatt, C.D.R., de Mendoza, A., Marlétaz, F., Firbas, P.N., Feuda, R., Masiero, G., Medina, P., Alcaina-Caro, A., et al. (2020). Genomic adaptations to aquatic and aerial life in mayflies and the origin of insect wings. Nat Commun *11*, 2631. https://doi.org/10.1038/s41467-020-16284-8.

22. Glassford, W.J., Johnson, W.C., Dall, N.R., Smith, S.J., Liu, Y., Boll, W., Noll, M., and Rebeiz, M. (2015). Co-option of an Ancestral Hox-Regulated Network Underlies a Recently Evolved Morphological Novelty. Dev. Cell *34*, 520–531. https://doi.org/10.1016/j.devcel.2015.08.005.

23. Boxshall, G.A., and Jaume, D. (2009). Exopodites, epipodites and gills in Crustaceans. Arthropod Syst. Phylog. *67*, 229–254.

24. Bruce, H.S., and Patel, N.H. (2020). Knockout of crustacean leg patterning genes suggests that insect wings and body walls evolved from ancient leg segments. Nat. Ecol. Evol. *4*, 1703–1712. https://doi.org/10.1038/s41559-020-01349-0.

25. Snodgrass, R.E. (1935). Principles of insect morphology. (McGraw-Hill Book Company, Inc).

26. Walossek, D., and Müller, K.J. (1990). Upper Cambrian stem-lineage crustaceans and their bearing upon the monophyletic origin of Crustacea and the position of Agnostus. Lethaia *Vol. 23, pp. 409–427*.

27. Wagele, J.W., and Bartolomaeus, T. (2014). Deep Metazoan Phylogeny: The Backbone of the Tree of Life. 216.

28. Backer, H., Fanenbruck, M., and Wägele, J.W. (2008). A forgotten homology supporting the monophyly of Tracheata: The subcoxa of insects and myriapods re-visited. Zoologischer Anzeiger - A Journal of Comparative Zoology *247*, 185–207. https://doi.org/10.1016/j.jcz.2007.11.002.

29. Størmer, L. (1939). Studies on Trilobite Morphology. Part I: The Thoracic Appendages and Their Phylogenetic Significance.

30. Deuve, T. (2001). The epipleural field in hexapods. Annales De La Societe Entomologique De France *37*, 195–231.

31. Haug, J.T., Maas, A., Haug, C., and Waloszek, D. (2013). EVOLUTION OF CRUSTACEAN APPENDAGES. In Functional Morphology and Diversity, Les Watling and M. Thiel, eds. (Oxford). https://doi.org/10.18476/pale.v12.a1.

32. Brusca, R.C. (2001). Origin of the Hexapoda. Journal of Crustacean Biology *21(4):1084-1086.* https://doi.org/10.1651/0278-0372(2001)021%25255B1084:OOTH%25255D2.0.CO;2.

33. Deuve, T. (2018). What is the epipleurite? A contribution to the subcoxal theory as applied to the insect abdomen. Ann. Soc. entomol. Fr. *54*, 1–26. https://doi.org/10.1080/00379271.2018.1431568.

34. Kobayashi, Y., Niikura, K., Oosawa, Y., and Takami, Y. (2013). Embryonic development of Carabus insulicola (Insecta, Coleoptera, Carabidae) with special reference to external morphology and tangible evidence for the subcoxal theory. Journal of Morphology *274*, 1323–1352.

35. Kobayashi, Y. (2017). Formation of Subcoxae-1 and 2 in Insect Embryos: The Subcoxal Theory Revisited. Proc Arthropod Embryol Soc Jpn *48*, 33–38.

36. Matsuda, R. (1970). Morphology and evolution of the insect thorax. Mem. Entomol. Soc. Can. *Volume 102*, 5–431. https://doi.org/10.4039/entm10276fv.

37. Kobayashi, Y., Bruce, H.S., Terada, K., Hirayama, A., Oosawa, Y., Niikura, K., and Takami, Y. (2022). Paranotal Lobes are Appendicular in Origin: Elucidation by Micro-CT Analysis of the Thoracic Muscular System in the Larvae of Carabus insulicola (Insecta, Coleoptera). Proceedings of the Arthropodan Embryological Society of Japan.

38. Ikmi, A., Netter, S., and Coen, D. (2008). Prepatterning the Drosophila notum: the three genes of the iroquois complex play intrinsically distinct roles. Dev. Biol. *317*, 634–648. https://doi.org/10.1016/j.ydbio.2007.12.034.

39. Calleja, M., Herranz, H., Estella, C., Casal, J., Lawrence, P., Simpson, P., and Morata, G. (2000). Generation of medial and lateral dorsal body domains by the pannier gene of Drosophila. Development *127*, 3971–3980.

40. Diez del Corral, R., Aroca, P., Gomez-Skarmeta, J.L., Cavodeassi, F., and Modolell, J. (1999). The Iroquois homeodomain proteins are required to specify body wall identity in Drosophila. Genes Dev. *13*, 1754–1761.

41. Cavodeassi, F., Diez del Corral, R., Campuzano, S., and Domínguez, M. (1999). Compartments and organising boundaries in the Drosophila eye: the role of the homeodomain Iroquois proteins. Development *126*, 4933–4942.

42. Carrasco-Rando, M., Tutor, A.S., Prieto-Sánchez, S., González-Pérez, E., Barrios, N., Letizia, A., Martín, P., Campuzano, S., and Ruiz-Gómez, M. (2011). Drosophila Araucan and Caupolican Integrate Intrinsic and Signalling Inputs for the Acquisition by Muscle Progenitors of the Lateral Transverse Fate. PLoS Genet *7*, e1002186-13. https://doi.org/10.1371/journal.pgen.1002186.

43. Herranz, H., and Morata, G. (2001). Embryonic functions of pannier. 10.

44. Bruce, H.S., and Patel, N.H. (2022). The Daphnia carapace and other novel structures evolved via the cryptic persistence of serial homologs. Curr. Biol. *32*. https://doi.org/10.1016/j.cub.2022.06.073.

45. Fisher, C.R., Kratovil, J.D., Angelini, D.R., and Jockusch, E.L. (2021). Out from under the wing: reconceptualizing the insect wing gene regulatory network as a versatile, general module for body-wall lobes in arthropods. Proc. R. Soc. B. *288*, 20211808. https://doi.org/10.1098/rspb.2021.1808.

46. Yang, J., Ortega-Hernández, J., Legg, D.A., Lan, T., Hou, J., and Zhang, X. (2018). Early Cambrian fuxianhuiids from China reveal origin of the gnathobasic protopodite in euarthropods. Nat Commun *9*, 1–9. https://doi.org/10.1038/s41467-017-02754-z.

47. Roonwal, M.L. (1937). Studies on the embryology of the African migratory locust, Locusta migratoria migratorioides Reiche and Frm.(Orthoptera, Acrididae). II. Organogeny. Philosophical Transactions of the Royal Society of \ldots *CCXXVII-B543*.

48. Ballesteros, J.A., Santibáñez-López, C.E., Baker, C.M., Benavides, L.R., Cunha, T.J., Gainett, G., Ontano, A.Z., Setton, E.V.W., Arango, C.P., Gavish-Regev, E., et al. (2022). Comprehensive Species Sampling and Sophisticated Algorithmic Approaches Refute the Monophyly of Arachnida. Molecular Biology and Evolution *39*, msac021. https://doi.org/10.1093/molbev/msac021.

49. Lozano-Fernandez, J., Tanner, A.R., Giacomelli, M., Carton, R., Vinther, J., Edgecombe, G.D., and Pisani, D. (2019). Increasing species sampling in chelicerate genomic-scale datasets provides support for monophyly of Acari and Arachnida. Nat Commun, 1–8. https://doi.org/10.1038/s41467-019-10244-7.

50. Weber, H. (1952). Morphologie, Histologie und Entwicklungsgeschichte Der Articulaten (G. Fischer).

51. Kukalova-Peck, J. (1978). Origin and evolution of insect wings and their relation to metamorphosis, as documented by the fossil record. J. Morphol.

52. Ewing, H.E. (1928). The legs and leg-bearing segments of some primitive arthropod groups, with notes on leg-segmentation in the Arachnida. Smithsonian Inst *80*, 1–41.

665   53. Coulcher, J.F., Edgecombe, G.D., and Telford, M.J. (2015). Molecular developmental evidence for a subcoxal origin of pleurites in insects and identity of the subcoxa in the gnathal appendages. Scientific reports *5*, 1–8.

54. Heymons, R. (1899). Beiträge zur Morphologie und Entwicklungsgeschichte der Rhynchoten.

670   55. Heitzler, P., Haenlin, M., Ramain, P., Calleja, M., and Simpson, P. (1996). A Genetic Analysis of Pannier, a Gene Necessary for Viability of Dorsal Tissues and Bristle Positioning in Drosophila. Genetics *143*, 1271–1286. https://doi.org/10.1093/genetics/143.3.1271.

56. Klinedinst, S.L., and Bodmer, R. (2003). Gata factor Pannier is required to establish
675   competence for heart progenitor formation. Development *130*, 3027–3038. https://doi.org/10.1242/dev.00517.

57. Villa-Cuesta, E., González-Pérez, E., and Modolell, J. (2007). Apposition of iroquois expressing and non-expressing cells leads to cell sorting and fold formation in the Drosophila imaginal wing disc. BMC Dev Biol *7*, 106–109. https://doi.org/10.1186/1471-
680   213X-7-106.

58. Pechmann, M. (2020). Embryonic development and secondary axis induction in the Brazilian white knee tarantula Acanthoscurria geniculata, C. L. Koch, 1841 (Araneae; Mygalomorphae; Theraphosidae). Dev Genes Evol *230*, 75–94. https://doi.org/10.1007/s00427-020-00653-w.

685   59. Emms, D.M., and Kelly, S. (2019). OrthoFinder: phylogenetic orthology inference for comparative genomics. Genome Biol *20*, 238. https://doi.org/10.1186/s13059-019-1832-y.

60. Kobayashi, Y. (2018). The paracoxal suture in insect embryos: Its state and importance for understanding the basalmost podomeres. 1–29. https://doi.org/10.13140/RG.2.2.24495.64165.

690   61. Winick, J., Abel, T., Leonard, M.W., M. Michelson, A., Chardon-Loriaux, I., Holmgren, R.A., Maniatis, T., and Engel, J.D. (1993). A GATA family transcription factor is expressed along the embryonic dorsoventral axis in *Drosophila melanogaster*. Development *119*, 1055–1065. https://doi.org/10.1242/dev.119.4.1055.

62. Gajewski, K., Zhang, Q., Choi, C.Y., Fossett, N., Dang, A., Kim, Y.H., Kim, Y., and Schulz,
695   R.A. (2001). Pannier is a Transcriptional Target and Partner of Tinman during Drosophila Cardiogenesis. Developmental Biology *233*, 425–436. https://doi.org/10.1006/dbio.2001.0220.

63. Matsuoka, Y., Murugesan, S.N., Prakash, A., and Monteiro, A. (2023). Lepidopteran prolegs are novel traits, not leg homologs. Science Advances *9*, eadd9389.
700   https://doi.org/10.1126/sciadv.add9389.

64. Dove, H., Tautz, D., and Roth, S. (2003). Neurogenesis in the millipede Glomeris marginata (Myriapoda: Diplopoda).

65. Leite, D.J., Schönauer, A., Blakeley, G., Harper, A., Garcia-Castro, H., Baudouin-Gonzalez, L., Wang, R., Sarkis, N., Nikola, A.G., Koka, V.S.P., et al. (2024). An atlas of spider development at single-cell resolution provides new insights into arthropod embryogenesis. EvoDevo *15*, 5. https://doi.org/10.1186/s13227-024-00224-4.

66. Prpic, N.-M., and Damen, W.G.M. (2009). Notch-mediated segmentation of the appendages is a molecular phylotypic trait of the arthropods. Developmental Biology *326*, 262–271. https://doi.org/10.1016/j.ydbio.2008.10.049.

67. O'Day, K.E. Notch signaling and segmentation in Parhyale hawaiensis. 195.

68. Thumecke, S., and Schroder, R. (2021). The odd-skipped related gene drumstick is required for leg development in the beetle Tribolium castaneum. Developmental Dynamics *251*, 1456–1471. https://doi.org/10.1002/dvdy.347.

69. Hao, I., Green, R.B., Dunaevsky, O., Lengyel, J.A., and Rauskolb, C. (2003). The odd-skipped family of zinc finger genes promotes Drosophila leg segmentation. Developmental Biology *263*, 282–295. https://doi.org/10.1016/j.ydbio.2003.07.011.

70. Mirth, C., and Akam, M. (2002). Joint development in the Drosophila leg: cell movements and cell populations. Developmental Biology *246*, 391–406. https://doi.org/10.1006/dbio.2002.0593.

71. Pechmann, M., and Prpic, N.-M. (2009). Appendage patterning in the South American bird spider Acanthoscurria geniculata (Araneae: Mygalomorphae). Dev Genes Evol *219*, 189–198. https://doi.org/10.1007/s00427-009-0279-7.

72. Pechmann, M. (2020). Embryonic development and secondary axis induction in the Brazilian white knee tarantula Acanthoscurria geniculata, C. L. Koch, 1841 (Araneae; Mygalomorphae; Theraphosidae). Dev Genes Evol *130*, 1735–20. https://doi.org/10.1007/s00427-020-00653-w.

73. Abzhanov, A., and Kaufman, T.C. (2000). Homologs of Drosophila appendage genes in the patterning of arthropod limbs. Dev. Biol. *227*, 673–689. https://doi.org/10.1006/dbio.2000.9904.

74. Requena, D., Álvarez, J.A., Gabilondo, H., Loker, R., Mann, R.S., and Estella, C. (2017). Origins and Specification of the Drosophila Wing. Current Biology *27*, 3826-3836.e5. https://doi.org/10.1016/j.cub.2017.11.023.

75. Estella, C. (2012). A dynamic network of morphogens and transcription factors patterns the fly leg. Curr. Top. Dev. Biol. *98*, 173–198. https://doi.org/10.1016/B978-0-12-386499-4.00007-0.

76. Gorfinkiel, N., Morata, G., and Guerrero, I. (1997). The homeobox gene Distal-less induces ventral appendage development in Drosophila. Genes & Development *11*, 2259–2271.

77. Setton, E.V.W., and Sharma, P.P. (2018). Cooption of an appendage-patterning gene cassette in the head segmentation of arachnids. *128*, 201720193–10. https://doi.org/10.1073/pnas.1720193115.

78. Sharma, P.P., Schwager, E.E., Giribet, G., Jockusch, E.L., and Extavour, C.G. (2013). Distal-less and dachshund pattern both plesiomorphic and apomorphic structures in chelicerates: RNA interference in the harvestman Phalangium opilio (Opiliones). Evol. Dev. *15*, 228–242. https://doi.org/10.1111/ede.12029.

79. Pechmann, M., Khadjeh, S., Turetzek, N., McGregor, A.P., Damen, W.G.M., and Prpic, N.-M. (2011). Novel Function of Distal-less as a Gap Gene during Spider Segmentation. PLoS Genet *7*, e1002342. https://doi.org/10.1371/journal.pgen.1002342.s011.

80. Schoppmeier, M., and Damen, W.G.M. (2001). Double-stranded RNA interference in the spider Cupiennius salei : the role of Distal-less is evolutionarily conserved in arthropod appendage formation. Dev Genes Evol *211*, 76–82. https://doi.org/10.1007/s004270000121.

81. Khila, A., and Grbić, M. (2007). Gene silencing in the spider mite Tetranychus urticae: dsRNA and siRNA parental silencing of the Distal-less gene. Dev Genes Evol *217*, 241–251. https://doi.org/10.1007/s00427-007-0132-9.

82. Konigsmann, T., Turetzek, N., Pechmann, M., and Prpic, N.-M. (2017). Expression and function of the zinc finger transcription factor Sp6–9 in the spider Parasteatoda tepidariorum. Dev Genes Evol *130*, 1–12. https://doi.org/10.1007/s00427-017-0595-2.

83. Turetzek, N., Pechmann, M., Schomburg, C., Schneider, J., and Prpic, N.-M. (2015). Neofunctionalization of a Duplicate dachshundGene Underlies the Evolution of a Novel Leg Segment in Arachnids. Mol. Biol. Evol. *33*, 109–121. https://doi.org/10.1093/molbev/msv200.

84. Prpic, N.-M., Janssen, R., Wigand, B., Klingler, M., and Damen, W.G.M. (2003). Gene expression in spider appendages reveals reversal of exd/hth spatial specificity, altered leg gap gene dynamics, and suggests divergent distal morphogen signaling. Developmental Biology *264*, 119–140. https://doi.org/10.1016/j.ydbio.2003.08.002.

85. Cohen, S.M., and Jurgens, G. (1989). Proximal-distal pattern formation in Drosophila: cell autonomous requirement for Distal-less gene activity in limb development. EMBO J. *8*, 2045–2055.

86. Cohen, B., Simcox, A.A., and Cohen, S.M. (1993). Allocation of the thoracic imaginal primordia in the Drosophila embryo. Development *117*, 597–608.

87. Campbell, G., and Tomlinson, A. (1998). The roles of the homeobox genes aristaless and Distal-less in patterning the legs and wings of Drosophila. Development *125*, 4483–4493.

88. Beermann, A., Jay, D.G., Beeman, R.W., Hülskamp, M., Tautz, D., and Jürgens, G. (2001). The Short antennae gene of Tribolium is required for limb development and encodes the orthologue of the Drosophila Distal-less protein. Development *128*, 287–297.

89. Angelini, D.R., and Kaufman, T.C. (2004). Functional analyses in the hemipteran Oncopeltus fasciatus reveal conserved and derived aspects of appendage patterning in insects. Developmental Biology *271*, 306–321. https://doi.org/10.1016/j.ydbio.2004.04.005.

90. Hiruta, C., Kakui, K., Tollefsen, K.E., and Iguchi, T. (2018). Targeted gene disruption by use of CRISPR/Cas9 ribonucleoprotein complexes in the water flea Daphnia pulex. Genes Cells *23*, 494–502. https://doi.org/10.1111/gtc.12589.

91. Beermann, A., Aranda, M., and Schröder, R. (2004). The Sp8 zinc-finger transcription factor is involved in allometric growth of the limbs in the beetle Tribolium castaneum. Development *131*, 733–742. https://doi.org/10.1242/dev.00974.

92. Schaeper, N.D., Prpic, N.-M., and Wimmer, E.A. (2009). A conserved function of the zinc finger transcription factor Sp8/9 in allometric appendage growth in the milkweed bug Oncopeltus fasciatus. Dev Genes Evol *219*, 427–435. https://doi.org/10.1007/s00427-009-0301-0.

93. Estella, C., and Mann, R.S. (2010). Non-Redundant Selector and Growth-Promoting Functions of Two Sister Genes, buttonhead and Sp1, in Drosophila Leg Development. PLoS Genet *6*, e1001001. https://doi.org/10.1371/journal.pgen.1001001.g007.

94. Bruce, H.S., and Patel, N.H. (2025). The molecular basis of arthropod appendage diversity. Arthropod Structure & Development.

95. Tavsanli, B.C., Ostrin, E.J., Burgess, H.K., Middlebrooks, B.W., Pham, T.A., and Mardon, G. (2004). Structure–function analysis of the Drosophila retinal determination protein Dachshund. Developmental Biology *272*, 231–247. https://doi.org/10.1016/j.ydbio.2004.05.005.

96. Yang, X., Zarinkamar, N., Bao, R., and Friedrich, M. (2009). Probing the Drosophila retinal determination gene network in Tribolium (I): The early retinal genes dachshund, eyes absent and sine oculis. Developmental Biology *333*, 202–214. https://doi.org/10.1016/j.ydbio.2009.02.040.

97. Sharma, P.P., Tarazona, O.A., Lopez, D.H., Schwager, E.E., Cohn, M.J., Wheeler, W.C., and Extavour, C.G. (2015). A conserved genetic mechanism specifies deutocerebral appendage identity in insects and arachnids. Proceedings of the Royal Society B: Biological Sciences *282*, 20150698–20150698. https://doi.org/10.1006/dbio.1999.9309.

98. Casares, F., and Mann, R.S. (2001). The ground state of the ventral appendage in Drosophila. Science *293*, 1477–1480. https://doi.org/10.1126/science.1062542.

99. Mardon, G., Solomon, N.M., and Rubin, G.M. (1994). dachshund encodes a nuclear protein required for normal eye and leg development in Drosophila. Development *120*, 3473–3486.

100. Wu, J., and Cohen, S.M. (1999). Proximodistal axis formation in the Drosophila leg: subdivision into proximal and distal domains by Homothorax and Distal-less. Development *126*, 109–117.

101. Coulter, D.E., Swaykus, E.A., Beran-Koehn, M.A., Goldberg, D., Wieschaus, E., and Schedl, P. (1990). Molecular analysis of odd-skipped, a zinc finger encoding segmentation gene with a novel pair-rule expression pattern. EMBO J *9*, 3795–3804.

102. Ward, E.J., and Coulter, D.E. (2000). odd-skipped is expressed in multiple tissues during Drosophila embryogenesis. Mechanisms of Development *96*, 233–236. https://doi.org/10.1016/S0925-4773(00)00389-0.

103. de Celis, J.F., Tyler, D.M., de Celis, J., and Bray, S.J. (1998). Notch signalling mediates segmentation of the Drosophila leg. Development *125*, 4617–4626. https://doi.org/10.1242/dev.125.23.4617.

104. Gainett, G., Klementz, B.C., Setton, E.V.W., Simian, C., Iuri, H.A., Edgecombe, G.D., Peretti, A.V., and Sharma, P.P. (2023). A plurality of morphological characters need not equate with phylogenetic accuracy: A rare genomic change refutes the placement of Solifugae and Pseudoscorpiones in Haplocnemata. Evolution & Development *n/a*, e12467. https://doi.org/10.1111/ede.12467.

105. Tiegs, O.W. (1940). The embryology and affinities of the symphyla based on a study of Hanseniella agilis. Q. J. Microsc. Sci. *82*, 1–208.

106. Coulcher, J.F., Edgecombe, G.D., and Telford, M.J. (2015). Molecular developmental evidence for a subcoxal origin of pleurites in insects and identity of the subcoxa in the gnathal appendages. Sci Rep *5*, 1–8. https://doi.org/10.1038/srep15757.

107. Komatsu, S., and Kobayashi, Y. (2012). Embryonic development of a whirligig beetle, Dineutus mellyi, with special reference to external morphology (insecta: Coleoptera, Gyrinidae). J. Morphol. *273*, 541–560. https://doi.org/10.1002/jmor.20005.

108. Fukuda, Y. (1994). Existence of Precoxa in the Thoracic Appendages of Decapod Crustaceans and its Functions. Proc. Japan. Soc. Sys. Zool. *52*, 47–64.

109. Borradaile, L.A. (1926). XXVII.— Notes upon Crustacean limbs. Journal of Natural History Series 11 *17*, 193–213. https://doi.org/10.1111/j.1096-3642.1922.tb00467.x.

110. Wesener, T. (2014). Sternites and spiracles - The unclear homology of ventral sclerites in the basal millipede order Glomeridesmida (Myriapoda, Diplopoda). Arthropod Structure, 9.

111. Lozano-Fernandez, J., Giacomelli, M., Fleming, J.F., Chen, A., Vinther, J., Thomsen, P.F., Glenner, H., Palero, F., Legg, D.A., Iliffe, T.M., et al. (2019). Pancrustacean Evolution Illuminated by Taxon-Rich Genomic-Scale Data Sets with an Expanded Remipede Sampling. Genome Biol. Evol. *11*, 2055–2070. https://doi.org/10.1093/gbe/evz097.

112. Bernot, J.P., Owen, C.L., Wolfe, J.M., Meland, K., Olesen, J., and Crandall, K.A. (2023). Major Revisions in Pancrustacean Phylogeny and Evidence of Sensitivity to Taxon Sampling. Molecular Biology and Evolution *40*, msad175. https://doi.org/10.1093/molbev/msad175.

113. Manton, S.M. (1966). … Mechanisms. Part 9, Functional Requirements and Body Design in Symphyla and Pauropoda and the Relationships Between Myriapoda and Pterygote ….

114. Minelli, A. (2011). The Myriapoda, Treatise on Zoology - Anatomy, Taxonomy, Biology. Volume 2 Diplopoda.

115. Boxshall, G. (1997). Comparative limb morphology in major crustacean groups: the coxa-basis joint in postmandibular limbs. Arthropod Relationships.

116. Tanzola, R.D., and Villegas-Ojeda, M.A. (2017). Argulus ventanensis sp. n.(Crustacea, Branchiura) parasite of Hypsiboas pulchellus tadpoles (Anura, Hylidae). Pan Am J A S *12*, 218–226.

117. Horne, D.J. (2005). Homology and homoeomorphy in ostracod limbs. Hydrobiologia *538*, 55–80. https://doi.org/10.1007/s10750-004-4937-5.

118. Schram, F.R., and Koenemann, S. (2020). Evolution and phylogeny of pancrustacea: a story of scientific method (Oxford University Press).

119. Hong, S.Y. (1988). Development of epipods and gills in some pagurids and brachyurans. Journal of Natural History Series 11 *22*, 1005–1040. https://doi.org/10.1071/MF9650369.

120. Calman, W.T. (1909). A Treatise on Zoology. Part. VII. Appendiculata. Third Fascicle. Crustacea. (Lankester. London).

121. Dreyer, H., and Wägele, J.W. (2002). The Scutocoxifera tax. nov. and the information content of nuclear ssu rDNA sequences for reconstruction of isopod phylogeny (Peracarida: Isopoda). Journal of Crustacean Biology *22(2):217-234*. https://doi.org/10.1651/0278-0372(2002)022%25255B0217:TSTNAT%25255D2.0.CO;2.

122. Gruner, H.E. (1954). Über das Coxalglied der Pereiopoden der Isopoden (Crustacea).

123. Brandt, A., and Poore, G.C.B. (2003). Higher classification of the flabelliferan and related Isopoda based on a reappraisal of relationships. Invert. Systematics *17*, 893. https://doi.org/10.1071/IS02032.

124. Wolff, C. (2009). The embryonic development of the malacostracan crustacean Porcellio scaber (Isopoda, Oniscidea). Dev Genes Evol *219*, 545–564. https://doi.org/10.1007/s00427-010-0316-6.

125. Watling, L., Hof, C.H.J., and Schram, F.R. (2000). The place of the Hoplocarida in the Malacostracan Pantheon. Journal of Crustacean Biology *20*, 1–11. https://doi.org/10.1163/1937240X-90000002.

126. Schram, F.R. (1968). Paleosquilla gen. nov.: A Stomatopod (Crustacea) from the Cretaceous of Colombia. J. Paleontol. *Vol. 42, No. 5 (Sep., 1968), pp. 1297–1301*.

127. Hyzny, M., Hoch, I., Schram, F.R., and Rybár, S. (2014). Crangopsis Salter, 1863 from the Lower Carboniferous (Mississippian) of the Ostrava Formation - the first record of Aeschronectida (Malacostraca: Hoplocarida) from continental Europe. Bull. Geosci., 707–717. https://doi.org/10.3140/bull.geosci.1458.

128. Crampton, G.C. (1909). A contribution to the comparative morphology of the thoracic sclerites of insects. In.

129. Uchifune, T., and Machida, R. (2005). Embryonic development of Galloisiana yuasai Asahina, with special reference to external morphology (insecta: Grylloblattodea). Journal of Morphology *266*, 182–207.

130. ANDERSON, D.T. (1963). The Embryology of Dacus tryoni 2. Development of lmaginal Discs in the Embryo. Development.

131. Mashimo, Y., and Machida, R. (2017). Embryological evidence substantiates the subcoxal theory on the origin of pleuron in insects. Sci Rep *7*, 1–9. https://doi.org/10.1038/s41598-017-12728-2.

132. Choi, H.M.T., Schwarzkopf, M., Fornace, M.E., Acharya, A., Artavanis, G., Stegmaier, J., Cunha, A., and Pierce, N.A. (2018). Third-generation in situhybridization chain reaction: multiplexed, quantitative, sensitive, versatile, robust. Development *145*, dev165753-122.

133. Bruce, H.S., Jerz, G., Kelly, S., McCarthy, J., Pomerantz, A., Senevirathne, G., Sherrard, A., A Sun, D., Wolff, C., and H Patel, N. (2021). Hybridization Chain Reaction (HCR) In Situ Protocol v1. protocols.io *Hybridization Chain Reaction (HCR) In Situ Protocol v1*, 10.17504/protocols.io.bunznvf6. https://dx.doi.org/10.17504/protocols.io.bunznvf6.

134. Fritsch, M., and Richter, S. How body patterning might have worked in the evolution of arthropods—A case study of the mystacocarid Derocheilocaris remanei (Crustacea, Oligostraca). Journal of Experimental Zoology Part B: Molecular and Developmental Evolution *n/a*. https://doi.org/10.1002/jez.b.23140.

135. Bruce, H.S., and Patel, N.H. (2023). Cryptic Persistence of Truncated Abdominal Legs in Insects Enabled Diverse Outgrowths with Novel Functions. Preprints.org. https://doi.org/10.20944/preprints202212.0268.v3.

136. Kunze, J.C. (1981). The Functional Morphology of Stomatopod Crustacea. Philosophical Transactions of the Royal Society of … *292(1059):255-328*.

137. Manuel, M., Jager, M., Murienne, J., Clabaut, C., and Guyader, H.L. (2006). Hox genes in sea spiders (Pycnogonida) and the homology of arthropod head segments. Dev Genes Evol *216*, 481–491. https://doi.org/10.1007/s00427-006-0095-2.

138. Gainett, G., Klementz, B.C., Blaszczyk, P.O., Bruce, H.S., Patel, N.H., and Sharma, P.P. (2023). Dual Functions of labial Resolve the Hox Logic of Chelicerate Head Segments. Mol. Biol. Evol. *40*, msad037. https://doi.org/10.1093/molbev/msad037.

139. Telford, M.J., and Thomas, R.H. (1998). Expression of homeobox genes shows chelicerate arthropods retain their deutocerebral segment. Proc. Natl. Acad. Sci. U.S.A. *95*, 10671–10675.

140. Abzhanov, A., Popadić, A., and Kaufman, T.C. (1999). Chelicerate Hox genes and the homology of arthropod segments. Evol. Dev. *1*, 77–89.

141. Ortega-Hern ndez, J., Janssen, R., and Budd, G.E. (2017). Origin and evolution of the panarthropod head - A palaeobiological and developmental perspective. Arthropod Structure and Development *46*, 354–379. https://doi.org/10.1016/j.asd.2016.10.011.

142. Shultz, J.W. (2007). A phylogenetic analysis of the arachnid orders based on morphological characters. Zool J Linn Soc *150*, 221–265.

143. Sharma, P.P., Schwager, E.E., Extavour, C.G., and Giribet, G. (2012). Evolution of the chelicera: a dachshund domain is retained in the deutocerebral appendage of Opiliones (Arthropoda, Chelicerata). Evol. Dev. *14*, 522–533. https://doi.org/10.1111/ede.12005.

144. Prpic, N.-M., and Tautz, D. (2003). The expression of the proximodistal axis patterning genes Distal-less and dachshund in the appendages of Glomeris marginata (Myriapoda: Diplopoda) suggests a special role of these genes in patterning the head appendages. Dev Biol *260*, 97–112. https://doi.org/10.1016/S0012-1606(03)00217-3.

145. Scholtz, G., Mittmann, B., and Gerberding, M. (1998). The pattern of Distal-less expression in the mouthparts of crustaceans, myriapods and insects: new evidence for a gnathobasic mandible and the common origin of Mandibulata. Int. J. Dev. Biol. *42*, 801–810.

146. Popadic, A., Rusch, D., Peterson, M., Rogers, B.T., and Kaufman, T.C. (1996). Origin of the arthropod mandible. Nature *380*, 395–395. https://doi.org/10.1038/380395a0.

147. Coulcher, J.F., and Telford, M.J. (2013). Comparative gene expression supports the origin of the incisor and molar process from a single endite in the mandible of the red flour beetle Tribolium castaneum. EvoDevo *4*, 1–12. https://doi.org/10.1186/2041-9139-4-1.

148. Brena, C. (2015). Myriapoda. In (null) (Springer Vienna), pp. 141–189. https://doi.org/10.1007/978-3-7091-1865-8_6.

149. Coulcher, J.F. (2011). Evolution of the arthropod mandible: a molecular developmental perspective. 1–251.

150. Janssen, R. (2011). Diplosegmentation in the pill millipede Glomeris marginata is the result of dorsal fusion: Millipede diplosegmentation. Evolution & Development *13*, 477–487. https://doi.org/10.1111/j.1525-142X.2011.00504.x.

951. Janssen, R., Budd, G.E., and Damen, W.G.M. (2011). Gene expression suggests conserved mechanisms patterning the heads of insects and myriapods. Developmental Biology *357*, 64–72. https://doi.org/10.1016/j.ydbio.2011.05.670.

152. Hughes, C.L., and Kaufman, T.C. (2002). Exploring Myriapod Segmentation: The Expression Patterns of even-skipped, engrailed, and wingless in a Centipede. Developmental Biology *247*, 47–61. https://doi.org/10.1006/dbio.2002.0683.

153. Janssen, R., Budd, G.E., Damen, W.G.M., and Prpic, N.-M. (2008). Evidence for Wg-independent tergite boundary formation in the millipede Glomeris marginata. Dev Genes Evol *218*, 361–370. https://doi.org/10.1007/s00427-008-0231-2.

154. Janssen, R., Prpic, N.-M., and Damen, W.G.M. (2004). Gene expression suggests decoupled dorsal and ventral segmentation in the millipede Glomeris marginata (Myriapoda: Diplopoda). Dev. Biol. *268*, 89–104. https://doi.org/10.1016/j.ydbio.2003.12.021.

155. Prpic, N.-M. (2004). Homologs of wingless and decapentaplegic display a complex and dynamic expression profile during appendage development in the millipede Glomeris marginata (Myriapoda: Diplopoda). Front. Zool. *1*, 6. https://doi.org/10.1186/1742-9994-1-6.

156. Prpic, N.-M., Janssen, R., Damen, W.G.M., and Tautz, D. (2005). Evolution of dorsal-ventral axis formation in arthropod appendages: H15 and optomotor-blind/bifid-type T-box genes in the millipede Glomeris marginata (Myriapoda: Diplopoda). Evol. Dev. *7*, 51–57. https://doi.org/10.1111/j.1525-142X.2005.05006.x.

157. Sun, D.A., Bredeson, J.V., Bruce, H.S., and Patel, N.H. (2022). Identification and classification of cis-regulatory elements in the amphipod crustacean Parhyale hawaiensis. Development *149*, dev200793. https://doi.org/10.1242/dev.200793.

158. TransDecoder Haas, BJ. https://github.com/TransDecoder/TransDecoder. GitHub. https://github.com/TransDecoder/TransDecoder/wiki/Home.

159. Burglin, T.R., and Affolter, M. (2016). Homeodomain proteins: an update. Chromosoma *125*, 497–521. https://doi.org/10.1007/s00412-015-0543-8.

160. Sievers, F. (2011). Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. Mol. Syst. Biol. *7*, 539. https://doi.org/10.1038/msb.2011.75.

161. Katoh, K., Misawa, K., Kuma, K., and Miyata, T. (2002). MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. Nucleic Acids Res *30*, 3059–3066. https://doi.org/10.1093/nar/gkf436.

162. Edgar, R.C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Research *32*, 1792–1797. https://doi.org/10.1093/nar/gkh340.

985   163. Darriba, D., Taboada, G.L., Doallo, R., and Posada, D. (2011). ProtTest 3: fast selection of best-fit models of protein evolution. Bioinformatics *27*, 1164–1165. https://doi.org/10.1093/bioinformatics/btr088.

164. Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A., Markowitz, S., Duran, C., et al. (2012). Geneious Basic: An integrated and
990   extendable desktop software platform for the organization and analysis of sequence data. Bioinformatics *28*, 1647–1649. https://doi.org/10.1093/bioinformatics/bts199.

165. Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., and Gascuel, O. (2010). New Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies: Assessing the Performance of PhyML 3.0. Systematic Biology *59*, 307–321.
995   https://doi.org/10.1093/sysbio/syq010.

166. Huelsenbeck, J.P., and Ronquist, F. (2001). MRBAYES: Bayesian inference of phylogenetic trees. Bioinformatics *17*, 754–755. https://doi.org/10.1093/bioinformatics/17.8.754.

167. Bianchini, G., and Sánchez-Baracaldo, P. (2024). TreeViewer: Flexible, modular software
1000   to visualise and manipulate phylogenetic trees. Ecology and Evolution *14*, e10873. https://doi.org/10.1002/ece3.10873.

168. Thomas, R.H., and Telford, M.J. (1999). Appendage development in embryos of the oribatid mite Archegozetes longisetosus (Acari, Oribatei, Trhypochthoniidae). Acta Zoologica, 80:193-200.

1005   169. Ito, A., Aoki, M.N., Yahata, K., and Wada, H. (2011). Complicated evolution of the caprellid (Crustacea: Malacostraca: Peracarida: Amphipoda) body plan, reacquisition or multiple losses of the thoracic limbs and pleons. Dev Genes Evol *221*, 133–140. https://doi.org/10.1007/s00427-011-0365-5.

170. Mittmann, B., and Scholtz, G. (2001). Distal-less expression in embryos of Limulus
1010   polyphemus (Chelicerata, Xiphosura) and Lepisma saccharina (Insecta, Zygentoma) suggests a role in the development of mechanoreceptors, chemoreceptors, and the CNS. Dev Genes Evol *211*, 232–243. https://doi.org/10.1007/s004270100150.

171. Williams, T.A. (1998). Distalless expression in crustaceans and the patterning of branched limbs. Dev Genes Evol *207*, 427–434.

1015   172. Panganiban, G., Sebring, A., Nagy, L., and Carroll, S. (1995). The development of crustacean limbs and the evolution of arthropods. Science *270*, 1363–1366.

173. Pabst, T., and Scholtz, G. (2009). The Development of Phyllopodous Limbs in Leptostraca and Branchiopoda. Journal of Crustacean Biology *29*, 1–12. https://doi.org/10.1651/08/3034.1.

1020 174. Copf, T. (2003). Posterior patterning genes and the identification of a unique body region in the brine shrimp Artemia franciscana. Development *130*, 5915–5927. https://doi.org/10.1242/dev.00835.

175. Kato, Y., Shiga, Y., Kobayashi, K., Tokishita, S., Yamagata, H., Iguchi, T., and Watanabe, H. (2011). Development of an RNA interference method in the cladoceran crustacean
1025 Daphnia magna. Dev Genes Evol *220*, 337–345. https://doi.org/10.1007/s00427-011-0353-9.

176. Shiga, Y., Yasumoto, R., Yamagata, H., and Hayashi, S. (2002). Evolving role of Antennapedia protein in arthropod limb patterning. Development *129*, 3555–3561.

177. Estella, C., Rieckhof, G., Calleja, M., and Morata, G. (2003). The role of buttonhead and
1030 Sp1 in the development of the ventral imaginal discs of Drosophila. Development *130*, 5929–5941. https://doi.org/10.1242/dev.00832.

178. Rogers, B.T., Peterson, M.D., and Kaufman, T.C. (2002). The development and evolution of insect mouthparts as revealed by the expression patterns of gnathocephalic genes. Evol. Dev. *4*, 96–110.

1035 179. Suzuki, Y., and Palopoli, M. (2001). Evolution of insect abdominal appendages: are prolegs homologous or convergent traits? Dev Genes Evol *211*, 486–492. https://doi.org/10.1007/s00427-001-0182-3.

180. Mito, T., Ronco, M., Uda, T., Nakamura, T., Ohuchi, H., and Noji, S. (2008). Divergent and conserved roles of extradenticle in body segmentation and appendage formation,
1040 respectively, in the cricket Gryllus bimaculatus. Developmental Biology *313*, 67–79. https://doi.org/10.1016/j.ydbio.2007.09.060.

181. Niwa, N., Saitoh, M., Ohuchi, H., Yoshioka, H., and Noji, S. (1997). Correlation between Distal-less expression patterns and structures of appendages in development of the two-spotted cricket, Gryllus bimaculatus. Zool. Sci. *14: 115-125*.

1045 182. Inoue, Y., Mito, T., Miyawaki, K., and Matsushima, K. (2002). Correlation of expression patterns of homothorax, dachshund, and Distal-less with the proximodistal segmentation of the cricket leg bud. Mechanisms of … *113 (2002) 141–148*.

183. Jockusch, E.L., Nulsen, C., Newfeld, S.J., and Nagy, L.M. (2000). Leg development in flies versus grasshoppers: differences in dpp expression do not lead to differences in the
1050 expression of downstream components of the leg patterning pathway. Development *127*, 1617–1626.

184. Schaeper, N.D., Wimmer, E.A., and Prpic, N.-M. (2013). Appendage patterning in the primitively wingless hexapods Thermobia domestica (Zygentoma: Lepismatidae) and Folsomia candida (Collembola: Isotomidae). Dev Genes Evol *223*, 341–350.
1055 https://doi.org/10.1007/s00427-013-0449-5.

185. Ohde, T., Masumoto, M., and Yaginuma, T. (2009). Embryonic RNAi analysis in the firebrat, Thermobia domestica: Distal-less is required to form caudal filament. Journal of insect … *78, 99–105*.

186. Galindo, M.I., Fernández-Garza, D., Phillips, R., and Couso, J.P. (2011). Control of Distal-less expression in the Drosophila appendages by functional 3′ enhancers. Developmental Biology *353*, 396–410. https://doi.org/10.1016/j.ydbio.2011.02.005.

187. Córdoba, S., Requena, D., Jory, A., Saiz, A., and Estella, C. (2016). The evolutionarily conserved transcription factor Sp1 controls appendage growth through Notch signaling. Development *143*, 3623–3631. https://doi.org/10.1242/dev.138735.

1060

1065

Arthropod legs

a Chelicerate (*Acanthoscurria*)
7 coxa · 4 patella · 3 tibia · 2 tarsus · 5 femur · 6 trochanter · 1 pretarsus

b Myriapod (*Oxidus*)
5 prefemur · 4 femur · 3 tibia · 2 tarsus · 7 coxa · 6 trochanter · 1 pretarsus

c Crustacean (*Parhyale*)
7 coxa · 2 propodus · 1 dactyl · 8 precoxa · 3 carpus · 6 basis · 5 ischium · 4 merus

d Insect (*Oncopeltus*)
3 tibia · 2 tarsus · 4 femur · 6 coxa · 5 trochanter · 1 pretarsus

Insect-crustacean leg segment homologies

e
Crustacean: Tergal plate · Body · Coxal plate · Gill
Insect: Wing · Body · Spiracle · Lobes

From Bruce & Patel 2020

Fig. 1. Arthropod legs. **a-d.** Chelicerates, myriapods, crustaceans, and insects have different numbers, shapes, and names for their leg segments. **e.** Leg segment correspondence between insect and crustacean leg segments. Colors indicate proposed homologies. Exites indicated with checker pattern. Panel modified from Bruce and Patel[24]. Phylogeny based on Lozano-Fernandez 2019[111] and Bernot 2023[112]. *Acanthoscurria* and *Oxidus* images from Wikipedia. Leg drawings adapted from Snodgrass 1952[1], except *Parhyale* leg, original image.

1070

A *Parhyale* — dorsal, tp, head, legs, *, 30um

B *Parhyale* — dorsal, tp, cp, gill, ventral, 8, 7, 6, 5, 4, 3, 2, 1, *
pnr, ara, Dll, DAPI

C *Tribolium* — dorsal, wing, head, L1

D *Tribolium* — dorsal, wing, ventral, 8, 7, 6, 5, 4, 2-3, 1

E *Acanthoscurria* — head, pp, L1, *

F *Acanthoscurria* — dorsal, 8, 7, 6, 5, 4, 3, 2, 1, ce, ventral, *, 100µm

G *Acanthoscurria* — 8, 7, 6, 5, 4, 3, 2, 1, ce, pp, L2

G' — head, ce, 8, 7, 6, 5, 4, 3, 2, 1, 150µm

H — dorsal, 8, 7, 6, 5, 4, 3, 2, 1, ce, ventral, 100µm
pnr, ara, Dll, odd, DAPI

1075 **Fig. 2.** Elucidating proximal leg segments in arthropods. Dissected right half of *Parhyale* (A), *Tribolium* (C), and *Acanthoscurria* (E) embryos. Dissected leg of *Parhyale* (B), *Tribolium* (D), and *Acanthoscurria* (F) embryos. *pnr* (red), *ara* (green), *Dll* (pink), DAPI (grey). In all arthropods for which there is data, leg segments 1 through 5 are identified by *Dll* expression at the embryonic stage where leg joints first become visible[13,24,71,73,76,78,80,81,87–89,92,143,144,146,168–186].

1080 *Dll* also expressed in *Parhyale* gill, coxal plate (cp) and tergal plate (tp). In all three arthropods, *pnr* marks the most dorsal domain. In all three arthropods, the two *ara* armband domains bracket a region proximal to leg segment 7. *ara* is also expressed in a smattering of ventral non-leg cells, and in *Parhyale* and *Acanthoscurria*, in the distal tip of the leg (*). *Tribolium* larvae have a fused tibia and tarsus, the tibiotarsus, here labelled 2-3[53]. In *Acanthoscurria*, leg segment 7 is easily

1085 identified by the coxal endite (ce) that bulges medially. Large dorsal cells in *Parhyale* and *Acanthoscurria* are yolk or extra-embryonic cells that exist prior to dorsal closure. (G – H) *Acanthoscurria drm* (yellow) is expressed in the distal region of each leg segment where the joint will later form. (G) Stage 12.5 *Acanthoscurria* embryos dissected away from yolk mass. Leg segment 7 is readily identified by the coxal endite. At this stage, proximal to leg segment 7,

1090 there is a leg-segment-like bulge (white curly brace), which expresses *drm* in its distal region (most visible in G'). By St 12.5, *pnr* expression in the walking legs and head is reduced and not visible through the other colors (compare with St 11.5 embryo in G). (H) dissected walking leg 1 from Stage 11.5 embryo where morphological bulges and subdivisions of the leg segments have not yet begun. *drm* encircles the distal region of each leg segment, including the hypothesized

1095 proximal 8th leg segment. *Dll* and *pnr* in F, H in same channel but have non-overlapping domains, *Dll* false colored. See Fig. S2 for separate *Dll* and *pnr Acanthoscurria* embryos. A, C from Bruce and Patel 2020[24]

1100

	Chelicerate (*P.tep*)	Crustacean (*P.haw*)	Insect (*T.cas, D.mel*)

WT

A — Pechmann 2011

B — Bruce 2020

C — Beerman 2001 — *Tribolium*

Dll

D — Pechmann 2011

E — Bruce 2020

F — Beerman 2001 — *Tribolium*

WT

G — Konigsman 2017

H — Bruce 2020

I — Estella 2010 — *Drosophila*

Sp6-9

J — Konigsman 2017

K — Bruce 2020

L — non-specific tissue — Estella 2010 — *Drosophila*

Fig. 3. *Dll* and *Sp6-9* function in arthropods. In all arthropods examined, including the chelicerate *Parasteatoda*, crustacean *Parhyale*, and insect *Tribolium*, *Dll* is expressed in and required for the development of leg segments 1 - 5, counting from the distal end of the leg[24,79,85–90]. In all arthropods examined, including *Parasteatoda*, *Parhyale,* and *Drosophila*, *Sp6-9* is expressed in and required for the development of leg segments 1 – 6[24,77,82,91–93]. In *Parasteatoda* embryos, leg segment boundaries are best approximations based on embryo morphology; the respective author's interpretations of their loss of function phenotypes, which were confirmed visually from their figures; as well as published gene expression boundaries for *Dll* and *Sp6-9* in *Parasteatoda* and other spiders and their respective author's interpretations[71,73,79,80,84]. In C, F, boundaries of leg segments 7 and 8 that form the lateral body wall of *Tribolium* are based on molecular and morphological analyses in Kobayashi 2022[37]; the remnant of the eudesmatic joint that ancestrally separated segments 7 and 8 corresponds to the paracoxal suture in Matsuda 1970[36], and Kobayashi 2013[34], 2017[35], and 2018[60]. In I, L, morphological boundary between leg segments 7 and 8 of *Drosophila* kindly provided by Professor Yukimasa Kobayashi's analyses of attachment points of muscles Iltpm4, 5, and 6, and Ipcm6 in the *Drosophila* thorax from Liu 2023, as well as the fact that no *Sp1* deletion phenotypes for body wall, pleura, or notum have been reported for *Drosophila*[93,177,187]. A, D from Pechmann 2011[79], note that strong Dll RNAi also deletes walking leg 1. B, E, H, K from Bruce and Patel 2020[24]. C, F from Beerman 2001[88]. G, J from Konigsman 2017[82]. I, L from Estella 2010[93]. L shows an animal with a clone of cells generated during embryogenesis with a genomic deletion of both Sp1 (i.e. Sp6-9) and btd (Sp5, the less active paralog[187]), and Estella 2010 conclude that the remaining leg "consisted of only a small patch of residual leg tissue. When leg tissue was observed, it was invariably associated with non-mutant tissue, suggesting that these clones were generated slightly later than those samples in which no leg tissue remained.". Abbreviations from source material: as, anterior spinneret; ch, chelicera; e, endite; L1-4, walking leg 1-4; pp, pedipalp; ps, posterior spinneret. See Fig. S2 for images without color.

1135

Fig. 4. *dac* and *hth* function in arthropods. A-C, WT legs. D-F, *dac* loss of function. In chelicerate *Phalangium*, crustacean *Parhyale*, and insect *Drosophila*, a strong *dac2* phenotype affects leg segments 3 – 5. Note that the *dac* phenotype in *Drosophila* extends partway into leg segment 2, such that the proximal part (tarsomeres 1 – 2, ta1, ta2) are fused with medial leg
1140 segments 3 – 5 into a contiguous mass. However, tarsal bristles are present on the distal end of the fused mass (blue shading) and the joint with tarsomere 4 is intact, suggesting that tarsomere 3 is partially present. The coxa (leg segment 6) and tarsomeres 4 and 5 (distal end of leg segment 2) remain intact. G-I, *hth* loss of function. In *Parhyale*, genomic loss of *hth* deletes the proximal leg segments, leaving only the distal 2 leg segments intact. In *Phalangium*, reduction of *hth*
1145 mRNA shortens and fuses the proximal leg segments, leaving the distal segments unaffected. Arrow in I, transverse row bristles, demarcating the distal end of leg segment 3. Scale bars 50μm except where indicated. A, D, G from Sharma 2013[78], 2015[97]. B, E, H from Bruce and Patel 2020[24]. C from Mardon 1994[99]. F, dac4/dac4 from Graeme Mardon, unpublished. Note this panel is zoomed in to 2x the magnification compared to the WT in C so that the reader may confirm
1150 the details of the phenotype. I from Casares 2001[98]. Abbreviations: T2-4, third thoracic legs. See Fig. S3 for this figure without color.

1155

Fig. 5. Proposed model for how all arthropod leg segments correspond to one another. A. Schematic of the leg segments and structures patterned by each gene. B. Morphology and homologies of arthropod leg segments based on leg gene function in insects, *Parhyale*, and chelicerates. Colors and patterns indicate proposed homologies. Exites (checker pattern); endites (stripe pattern). Drawings in B modified from Snodgrass 1952.

1160

Supplementary information

A system to decipher the origins and relationships of all arthropod outgrowths

Authors: Heather S. Bruce[1]* and Nipam H. Patel[2,3]

Affiliations: 1. University of British Columbia, Vancouver, BC V6T1Z4 Canada. 2. Marine Biological Laboratory, 7 MBL St. Woods Hole, MA 02543 USA. 3. University of Chicago, Organismal Biology & Anatomy, 1027 E 57th Street, Chicago, IL 60637 USA.

Acanthoscurria sequences and probes

Pool name	Sequence
Acanth.pnr.pnr.DN77604	GAGGAGGGCAGCAAACGGaaATCTAAATGGCCCCCCATGGTAAGA
Acanth.pnr.pnr.DN77604	ATGATGGTGGTGGTGGTGATGGTGAtaGAAGAGTCTTCCTTTACG
Acanth.pnr.pnr.DN77604	GAGGAGGGCAGCAAACGGaaGATTGTGATGTTGGTGGCTGGGAGA
Acanth.pnr.pnr.DN77604	AAGGGGTCAACGGAGATGACTGCTGtaGAAGAGTCTTCCTTTACG
Acanth.pnr.pnr.DN77604	GAGGAGGGCAGCAAACGGaaGAATAATTTGAAGAAGCTGGAGCTG
Acanth.pnr.pnr.DN77604	ACTCCGGTGGCGAAGGAAGCTATGCtaGAAGAGTCTTCCTTTACG
Acanth.pnr.pnr.DN77604	GAGGAGGGCAGCAAACGGaaATTCTTTGGCTTTCTTTTCCTTTTC
Acanth.pnr.pnr.DN77604	GCCTTTCTTTTTGATATCTGCTTGAtaGAAGAGTCTTCCTTTACG
Acanth.pnr.pnr.DN77604	GAGGAGGGCAGCAAACGGaaCGTGGTTATTTCTCCTCCAGAGAGT
Acanth.pnr.pnr.DN77604	ACCCACAGGCATTACAGACAGGTTCtaGAAGAGTCTTCCTTTACG
Acanth.pnr.pnr.DN77604	GAGGAGGGCAGCAAACGGaaCGCATGAGAGGTCTATTGACTCCAT
Acanth.pnr.pnr.DN77604	AGGGGACCTACCAGCCGCTTTTGCGtaGAAGAGTCTTCCTTTACG
Acanth.pnr.pnr.DN77604	GAGGAGGGCAGCAAACGGaaATTAACACATTCCCTCTCCTCGTGG
Acanth.pnr.pnr.DN77604	CCACAAGGGCGTAGATATGGCGCCGtaGAAGAGTCTTCCTTTACG
Acanth.pnr.pnr.DN77604	GAGGAGGGCAGCAAACGGaaTAGGATCTAAAAGTCCCCAAGGTGA
Acanth.pnr.pnr.DN77604	ATGGTGAATGTGAAAGACCGGACATtaGAAGAGTCTTCCTTTACG
Acanth.pnr.pnr.DN77604	GAGGAGGGCAGCAAACGGaaCCCGCACTCGATGTTGAAGCTAGAT
Acanth.pnr.pnr.DN77604	GTCCTACTCAAGTCATTCTGATCACtaGAAGAGTCTTCCTTTACG
Acanth.pnr.pnr.DN77604	GAGGAGGGCAGCAAACGGaaGGGCTGCGACACAGTTTGATGTCCC
Acanth.pnr.pnr.DN77604	TGCAGACTGATGATCTTGAGCACTTtaGAAGAGTCTTCCTTTACG
Acanth.pnr.pnr.DN77604	GAGGAGGGCAGCAAACGGaaGTGATGCGTACAGAGCGGCCGGCGA
Acanth.pnr.pnr.DN77604	ACGGTAACATGGTAAGAGCCCCGCCtaGAAGAGTCTTCCTTTACG
Acanth.pnr.pnr.DN77604	GAGGAGGGCAGCAAACGGaaGGTGCAGGGTAATTTCCAGCCATGC

Acanth.pnr.pnr.DN77604	TAAAGGTCACGACCCGAGTAAGAGGtaGAAGAGTCTTCCTTTACG
Acanth.pnr.pnr.DN77604	GAGGAGGGCAGCAAACGGaaGCCGTCAGATGGCGCAGTGGCGGTG
Acanth.pnr.pnr.DN77604	GGACCCACTCTCCGGGTCAGTTGTGtaGAAGAGTCTTCCTTTACG
Acanth.pnr.pnr.DN77604	GAGGAGGGCAGCAAACGGaaCCTGTAGACACGCCTTTGAGGCGCT
Acanth.pnr.pnr.DN77604	AAGGATGCACCTCCTCTCGCGGATCtaGAAGAGTCTTCCTTTACG
Acanth.pnr.pnr.DN77604	GAGGAGGGCAGCAAACGGaaGGTTGAGCAGAATTATGTGAGTGTT
Acanth.pnr.pnr.DN77604	TCATTTCCTTCAGCGGAAGGGCTCAtaGAAGAGTCTTCCTTTACG
Acanth.pnr.pnr.DN77604	GAGGAGGGCAGCAAACGGaaGGGAAGATCGGAATGCTCTTGAATT
Acanth.pnr.pnr.DN77604	GTGGTCTTCGAGGGACCAAGAATGTtaGAAGAGTCTTCCTTTACG
Acanth.pnr.pnr.DN77604	GAGGAGGGCAGCAAACGGaaGAATTTTCCACATCTTTATCCGATT
Acanth.pnr.pnr.DN77604	CGCTGATCGCCATTTACCCAGTTATtaGAAGAGTCTTCCTTTACG
Acanth.pnr.pnr.DN77604	GAGGAGGGCAGCAAACGGaaCTTTCCCATCTACCAGCAATGTTTT
Acanth.pnr.pnr.DN77604	ATCCATGCTCCATTATAAGTCGAATtaGAAGAGTCTTCCTTTACG
Acanth.pnr.pnr.DN77604	GAGGAGGGCAGCAAACGGaaGTTCCCTTTTATACTCTCTCCTCTT
Acanth.pnr.pnr.DN77604	GGTCAGTAGTCGTGCCTTGGAAAATtaGAAGAGTCTTCCTTTACG
Acanth.pnr.pnr.DN77604	GAGGAGGGCAGCAAACGGaaTACACCTGCTCTTCGCATTTTGTCG
Acanth.pnr.pnr.DN77604	TCCACATTTTGATCATGGCTCGTGAtaGAAGAGTCTTCCTTTACG
Acanth.pnr.B4.DN78099	CCTCAACCTACCTCCAACaaATGAGATTCTTTGCCATTTGAACAA
Acanth.pnr.B4.DN78099	TGGCCAAGTGTGCATGTGAGGTATAatTCTCACCATATTCGCTTC
Acanth.pnr.B4.DN78099	CCTCAACCTACCTCCAACaaTTGTAGTGATGGTATCTGATGAAGA
Acanth.pnr.B4.DN78099	TGGCGCCTGCTGCAGACTGTGTTGTatTCTCACCATATTCGCTTC
Acanth.pnr.B4.DN78099	CCTCAACCTACCTCCAACaaTGTTGATGTGACTGCTGATAAGGGA
Acanth.pnr.B4.DN78099	TGGGACTGGCACAGATGGTAGGTGGatTCTCACCATATTCGCTTC
Acanth.pnr.B4.DN78099	CCTCAACCTACCTCCAACaaGACGGTGGTGAAGGAGTTGAGCTAG
Acanth.pnr.B4.DN78099	TCAACAGGACTCCTCACTTGTCTTCatTCTCACCATATTCGCTTC
Acanth.pnr.B4.DN78099	CCTCAACCTACCTCCAACaaTGTCCTTCTCGGATGAGCCTTGTAC
Acanth.pnr.B4.DN78099	GGGCACTGCCATCAAAACTTAGCTTatTCTCACCATATTCGCTTC
Acanth.pnr.B4.DN78099	CCTCAACCTACCTCCAACaaGGGTGTTGCTGTTTTGGGCTTCCTT
Acanth.pnr.B4.DN78099	TGGGACTTCGGAAGACACAGCTGCTatTCTCACCATATTCGCTTC
Acanth.pnr.B4.DN78099	CCTCAACCTACCTCCAACaaGCAGGCGGACGGTTCATATTGTGCA
Acanth.pnr.B4.DN78099	CTGGTTTGAATTCCAGCTTTCCTCAatTCTCACCATATTCGCTTC
Acanth.pnr.B4.DN78099	CCTCAACCTACCTCCAACaaCAGGTTCTCCGTGATTGTTTCTCCT
Acanth.pnr.B4.DN78099	TATAATAGAGACCACATGCGTTGCAatTCTCACCATATTCGCTTC
Acanth.pnr.B4.DN78099	CCTCAACCTACCTCCAACaaCGAACAAGATAACCCCAGACGTCGT
Acanth.pnr.B4.DN78099	TAGTGTGGTGGTTGTAGTTCCACAGatTCTCACCATATTCGCTTC
Acanth.pnr.B4.DN78099	CCTCAACCTACCTCCAACaaAGTGGCCTGTTGACTCCGTTCATCC
Acanth.pnr.B4.DN78099	GCTGGCAGCCTCCTCTGAGGTTTCGatTCTCACCATATTCGCTTC
Acanth.pnr.B4.DN78099	CCTCAACCTACCTCCAACaaAGTGGCCGGTTCCATCTCTCCTCCA
Acanth.pnr.B4.DN78099	GGTATAATCCACATGCATTGCACAAatTCTCACCATATTCGCTTC
Acanth.pnr.B4.DN78099	CCTCAACCTACCTCCAACaaACATTCTCGGCCTTCGTTGAACTGA
Acanth.pnr.B4.DN78099	GGGGGTAGACACAGCTCCACAATTAatTCTCACCATATTCGCTTC

Acanth.pnr.B4.DN78099	CCTCAACCTACCTCCAACaaAGGTGTGCCATTGACGATTGCATTC
Acanth.pnr.B4.DN78099	TAATCGTGAGAATCTGCTCTTCGATatTCTCACCATATTCGCTTC
Acanth.pnr.B4.DN78099	CCTCAACCTACCTCCAACaaCGTTGTCCGTGATGTATGAGTGATA
Acanth.pnr.B4.DN78099	TGCCGTTCTCGAAGCTTGACCAATGatTCTCACCATATTCGCTTC
Acanth.pnr.B4.DN78099	CCTCAACCTACCTCCAACaaCGAGCCTCCGTCCCTAGTTTGTACC
Acanth.pnr.B4.DN78099	TGAGAATGCGTTGCTTCTTTGAAAAatTCTCACCATATTCGCTTC
Acanth.pnr.B4.DN78099	CCTCAACCTACCTCCAACaaGTTGTCGATAATGAGCTATGCGCAG
Acanth.pnr.B4.DN78099	ATGCCTGGAGTTTGGAATCCAAACCatTCTCACCATATTCGCTTC
Acanth.pnr.B4.DN78099	CCTCAACCTACCTCCAACaaCAGCAGTCCTATTTGGATCACCAGT
Acanth.pnr.B4.DN78099	GATGATACGTCTCGTGTACCTGAACatTCTCACCATATTCGCTTC
Acanth.pnr.B4.DN78099	CCTCAACCTACCTCCAACaaCATGGTTGCTCCTCCTTGATGCAGA
Acanth.pnr.B4.DN78099	ACCCCACGGGCTTGGGTTCGAACAAatTCTCACCATATTCGCTTC
Acanth.pnr.B4.DN78099	CCTCAACCTACCTCCAACaaACGGGACTAGCGTGAATGTATCCGT
Acanth.pnr.B4.DN78099	ACCATTGGCCTGGATGCTGGTAGGAatTCTCACCATATTCGCTTC
Acanth.pnr.B4.DN78099	CCTCAACCTACCTCCAACaaTATTCTGGAACATGTTGGGTACATA
Acanth.pnr.B4.DN78099	GATAAGATGGCATGTGGATAGAGGAatTCTCACCATATTCGCTTC
Acanth.pnr.B5.DN80076	CTCACTCCCAATCTCTATaaTCCACAGGCATTGCACACTGGTTCC
Acanth.pnr.B5.DN80076	TGTCTTCCTGTGAAGCTTGAAGTACaaCTACCCTACAAATCCAAT
Acanth.pnr.B5.DN80076	CTCACTCCCAATCTCTATaaTCTTTTGATTCTTGACTACAGGCCT
Acanth.pnr.B5.DN80076	GACCCATTCTTCGGGAAGCTGACAGaaCTACCCTACAAATCCAAT
Acanth.pnr.B5.DN80076	CTCACTCCCAATCTCTATaaATAGCACCACAATTGACACATTCGC
Acanth.pnr.B5.DN80076	CCGTCCCGCCTCCATAATGGCGTCGaaCTACCCTACAAATCCAAT
Acanth.pnr.B5.DN80076	CTCACTCCCAATCTCTATaaACTCGGGTCATAGCCCCATGAGGAA
Acanth.pnr.B5.DN80076	CAACTGAAGAGCAGTCATATTTGCTaaCTACCCTACAAATCCAAT
Acanth.pnr.B5.DN80076	CTCACTCCCAATCTCTATaaTAGGGGAAAGATCTTCACTCAGCTG
Acanth.pnr.B5.DN80076	ATGATGACGAAGGGGACATTTGTGTaaCTACCCTACAAATCCAAT
Acanth.pnr.B5.DN80076	CTCACTCCCAATCTCTATaaGTAGATACCGTTGGCACGTATAAAG
Acanth.pnr.B5.DN80076	GATGACAAGCCTAACAGTTTAGAGGaaCTACCCTACAAATCCAAT
Acanth.pnr.B5.DN80076	CTCACTCCCAATCTCTATaaTGCTGAATCAACACCAATATCCCTT
Acanth.pnr.B5.DN80076	CACAGAAGGTGTACTAACGGTATGTaaCTACCCTACAAATCCAAT
Acanth.pnr.B5.DN80076	CTCACTCCCAATCTCTATaaCAGAAGAATTTATACTTGCTAAAGT
Acanth.pnr.B5.DN80076	GCGGTGAAGATTCAATTGTTTGAGGaaCTACCCTACAAATCCAAT
Acanth.pnr.B5.DN80076	CTCACTCCCAATCTCTATaaGATCCGTGAACCTCAGGGAGGTCAA
Acanth.pnr.B5.DN80076	GAGGTGGGGCTTCGAATGTGCGGGAaaCTACCCTACAAATCCAAT
Acanth.pnr.B5.DN80076	CTCACTCCCAATCTCTATaaCTGTAAAGCGTTGTGAGAGTTTGTT
Acanth.pnr.B5.DN80076	ATGGTGAAGAATATCTCTGGTGTGAaaCTACCCTACAAATCCAAT
Acanth.pnr.B5.DN80076	CTCACTCCCAATCTCTATaaCAGCTCCCTCTCCTGCTACGTCTTC
Acanth.pnr.B5.DN80076	AAATATGAGGATGCATTTGGGCTGTaaCTACCCTACAAATCCAAT
Acanth.pnr.B5.DN80076	CTCACTCCCAATCTCTATaaGCACTGTGGAAATTCGACGAATGCA
Acanth.pnr.B5.DN80076	TCATGCTGCTGCTGTTGTTCATGGAaaCTACCCTACAAATCCAAT
Acanth.pnr.B5.DN80076	CTCACTCCCAATCTCTATaaTGGTTGGTTATCATCTGCCGAAAAG

Acanth.pnr.B5.DN80076	GTCCTCATGCTGGATATGAATGTGAaaCTACCCTACAAATCCAAT
Acanth.pnr.B5.DN80076	CTCACTCCCAATCTCTATaaTGATGGATATTGGGAGTTTCTCCCG
Acanth.pnr.B5.DN80076	TGCTCTTGATGATGATGATGAGGGGaaCTACCCTACAAATCCAAT
Acanth.pnr.B5.DN80076	CTCACTCCCAATCTCTATaaTATTACTAGAGTCTTTTGCATCTGC
Acanth.pnr.B5.DN80076	CCGGATTATAAGATGGGATACTACTaaCTACCCTACAAATCCAAT
Acanth.pnr.B5.DN80076	CTCACTCCCAATCTCTATaaTTCATCGAGGTTAGCTAAATGTGTT
Acanth.pnr.B5.DN80076	TAAATGGTCATGATGACGTCGCTCCaaCTACCCTACAAATCCAAT
Acanth.pnr.B5.DN80076	CTCACTCCCAATCTCTATaaTGGAGTAAAGAATCCTGTTCTGTAT
Acanth.pnr.B5.DN80076	CGCTGTTCACCCTGTCCGTGGAAGTaaCTACCCTACAAATCCAAT
Acanth.pnr.B5.DN80076	CTCACTCCCAATCTCTATaaAAATGTGCTCGTTCACGTGATCTCT
Acanth.pnr.B5.DN80076	GGTGCTCATTAGAATGGTGGTGATGaaCTACCCTACAAATCCAAT
Acanth.pnr.B5.DN80076	CTCACTCCCAATCTCTATaaTGGGACGCTATGTTCTGTTTCTACC
Acanth.pnr.B5.DN80076	TTCTGAGACGTGAGGAGAAAGATCAaaCTACCCTACAAATCCAAT
Acanth.pnr.B5.DN80076	CTCACTCCCAATCTCTATaaTCCAATTCTTGCACTCTCCTGTGAT
Acanth.pnr.B5.DN80076	TGTTCATTAACATCTTTGTCATCCTaaCTACCCTACAAATCCAAT

Acanthoscurria sequences

pannier

1180
>pannier_TRINITY_DN77604_c0_g1_i1
CAAAATGATGAAAATTCTCACGTTTCGCTTCATGATATTCATTCTTGGAGCAATCACGAT
AGTGCTGTAACAAGCCATTCTTGGGCAGATCATCCTTCTTCTCATAGTTCTAGTGTTCCA
AAAACATCATCATGGCCTGCGAAGCAACATCCAACTTCGTTGCATCAAGGTGTTGATCTT
CTGTGTCACGCCTGGGAAAAAACAAACGAGGAAAATCCAGAGCATTCACCTGGAGACTGG
1185
ATGAAAATGAAGTGACGACAAAATGCGAAGAGCAGGTGTACTTCACGAGCCATGATCAA
AATGTGGACGTAAGAGGAGAGAGTATAAAAGGGAACAAATTTTCCAAGGCACGACTACTG
ACCAACAAAACATTGCTGGTAGATGGGAAAGTAATTCGACTTATAATGGAGCATGGATCA
GAATCGGATAAAGATGTGGAAAATTCAGATAACTGGGTAAATGGCGATCAGCGTACAATT
CAAGAGCATTCCGATCTTCCCTCACATTCTTGGTCCCTCGAAGACCACGCCCATATCAAC
1190
GGTATAGTGTCACATTCTGCGCCTGAATCGTCTAATAACCCCTGGGAACACTCACATAAT
TCTGCTCAACCAATGAGCCCTTCCGCTGAAGGAAATGAAACACACCCTAGTGAGATAAAC
ATCAACCAATCGGCCTCTGAAACCCCCACCCACCAACAGCGCCTCAAAGGCGTGTCTACAG
GGTGATCCGCGAGAGGAGGTGCATCCTTCCGAGCAGCGGAATCACTCCAGCGTAGTTCAC
TTGCGATCGTCACAACAAGAACCCGACACCGCCACTGCGCCATCTGACGGCGGCACAACT
1195
GACCCGGAGAGTGGGTCCGACCACCCTAACAGTGTCCCAGTCATGGTCAGCACGTTTGGC
AGCATGGCTCCATACAGCATGGCTGGAAATTACCCTGCACCATCCTCTTACTCGGGTCGT
GACCTTTATTCTGGGAAACCATCTCCCGGCTACGGACATTCTGTGGACACCTCTTCTCCT
ACAACTTCGCCGGCCGCTCTGTACGCATCACCGGGCGGGGCTCTTACCATGTTACCGTAT
GTGACGGGTCAGTCAATGAGCCATCAATCCGTGGTCACCCAGGGCAGTCACGCATGGGGA
1200
CATCAAACTGTGTCGCAGCCCCCAAGTGCTCAAGATCATCAGTCTGCAAGTACATACGGA
ATTTCTGCTTTGTCGTCAGGATTGAACTTGGCTCAAAGTTCTCTTAATCTAGCTTCAACA
TCGAGTGCGGGATGTGATCAGAATGACTTGAGTAGGACAGGTGGATTTTCTACTTTCGGA
TCCACTGGACATGGTTATCTTCGTCCGGACATGACGTCACCTTGGGGACTTTTAGATCCT
ACAATGTCCGGTCTTTCACATTCACCATATTGTACAGATGGAATGGGACATCATTTAGGA
1205
CAAGATCGTGGGGACTATTTTGGCATCCACGAGGAGAGGGAATGTGTTAATTGCGGCGCC

```
      ATATCTACGCCCTTGTGGCGGCGAGACGGGACCGGCCACTACCTATGCAACGCCTGTGGA
      CTCTACAGCAAGATGAATGGAGTCAATAGACCTCTCATGCGACCGCAAAAGCGGCTGGTA
      GGTCCCCTAGTGACAGCGAATCGAAGAGCCGGTCAGATCTGCACAAACTGTGGAACAACC
      ACGACTACTCTCTGGAGGAGAAATAACCACGGTGAACCTGTCTGTAATGCCTGTGGGTTG
1210  TACTTCAAACTCCACGGGGTAAACAGGCCTCCCGCGATGAAGAAAGATGGCATACAGAAA
      AGGAAAAGAAAGCCAAAGAATAATCAAGCAGATATCAAAAAGAAAGGCCTGTCACCCATA
      TCAACAGGTGCGTCTAAGAGTTCCAATGAAGTCAGCAGCGGGAGCACAGCTCCAGCTTCT
      TCAAATTATTCGGGCATAGCTTCCTTCGCCACCGGAGTATCAGGAACGCCTTCAACACCA
      TCGTCTTCGTCTGCCGCAGTTACGTCTATTGTAGATTCTCCCAGCCACCAACATCACAAT
1215  CATCAGCAGTCATCTCCGTTGACCCCTTCTTCATCATCCTCTTCATCATCACCTTCTTCG
      TCGTTGTTGCAAAGACATATGCAGGGTCTTACCATGGGGGGCCATTTAGATCATCACCAT
      CACCACCACCACCATCAT

      >TRINITY_DN78099_c0_g1_i1
1220  ACTCTGAAGTGTATAGAAAAGGTAAATAAAATTCAAGCCTATTGTGAAGAGAATGAGAAT
      TATGGTTTTATATAACAGAGAGCAATGAATACAAAGGTAATATTCAGTGATTGTAACATA
      GTAAAGTTTTTATTGCAGCTTTATTAACACCTGCACAGAAATAACCCTGCTGTGGCTCAC
      TTTATATGAGGAAGTGCTGCATTTTTATTGGCCACAACACAGATATTCATGGCATATTTT
      TGGAATTTTTTTTTTCTGGATACATTTGAGTTCTAAATGAATACATATCTCACTTGCTGG
1225  ATGGACTTCGGTAACTGAGCATTGCTTTGCTGTGGTAAGTTACGAGACTCCGTAAAGCAC
      AGTTTGATGAACACAGGCTGAAAGAGAAGTTTTTCCAAACTTAGTACCTAACATTTTATT
      TGGGATTTCAAATGTCAAAATCTGTTATTTATGGCCAAGTGTGCATGTGAGGTATAAAT
      GAGATTCTTTGCCATTTGAACAAGGATTGGTTTGTCCTGAAATTGTTATGGACGATACAG
      GAGGTGGGTATTCACTCTGTATGGCGCCTGCTGCAGACTGTGTTGTCATTGTAGTGATGG
1230  TATCTGATGAAGATGATGGGACTGGCACAGATGGTAGGTGGCATGTTGATGTGACTGCTG
      ATAAGGGAGATTCAACAGGACTCCTCACTTGTCTTCCTGACGGTGGTGAAGGAGTTGAGC
      TAGGTGGGGCACTGCCATCAAAACTTAGCTTTTTGTCCTTCTCGGATGAGCCTTGTACAT
      CTGGGACTTCGGAAGACACAGCTGCTTTGGGTGTTGCTGTTTTGGGCTTCCTTTTCCTGG
      TTTGAATTCCAGCTTTCCTCATAGCAGGCGGACGGTTCATATTGTGCAGTTTATAATAGA
1235  GACCACATGCGTTGCATACAGGTTCTCCGTGATTGTTTCTCCTCCATAGTGTGGTGGTTG
      TAGTTCCACAGTTCGAACAAGATAACCCCAGACGTCGTGTTGCTGGCAGCCTCCTCTGAG
      GTTTCGACAGTGGCCTGTTGACTCCGTTCATCCTGTGGTATAATCCACATGCATTGCACA
      AGTAGTGGCCGGTTCCATCTCTCCTCCACAGGGGGGTAGACACAGCTCCACAATTAACAC
      ATTCTCGGCCTTCGTTGAACTGAGAATAATCGTGAGAATCTGCTCTTCGATTTAGGTGTG
1240  CCATTGACGATTGCATTCCATTGCCGTTCTCGAAGCTTGACCAATGTCCGTTGTCCGTGA
      TGTATGAGTGATAAGGTGAGAATGCGTTGCTTCTTTGAAAATACGAGCCTCCGTCCCTAG
      TTTGTACCCCCATGCCTGGAGTTTGGAATCCAAACCTGGTTGTCGATAATGAGCTATGCG
      CAGACGGATGATACGTCTCGTGTACCTGAACGGCAGCAGTCCTATTTGGATCACCAGTTT
      GACCCCACGGGCTTGGGTTCGAACAAGCCATGGTTGCTCCTCCTTGATGCAGAGTGACCA
1245  TTGGCCTGGATGCTGGTAGGAATACGGGACTAGCGTGAATGTATCCGTCTTGATAAGATG
      GCATGTGGATAGAGGACATATTCTGGAACATGTTGGGTACATACATATATAGAGAAGAAG
      CAGTTATCTTTATCGGCCGAAACCGTCTCTTGGGAAATAAAATACAATTTTTCGTGAAAA
      CGCTGACTTAAATGCCAATGCGCAGTGTGCCGCTGGATAAACTTGAAAACAGTAACATAT
      TTATCTGTTTTATAAGATGGAGAGAAAGAGAGGTGCAGGTTGCAGAAGAAACATCACAAG
1250  CAGCTTCGTTTACCACACTCTGTAACTATGTCATAAAAGTCCTTGAAGTTCAACTTCCTT
      TTAAGTTTGATCATGTGCTTATTCTACAAAGTTATTTAGAATTCCAGTTTCTGTTACAGC
      AGAACACTTCTTGAAAGACTCAAACACTATTTAGTTGTTTCTTTACTGAAGGAAGGGAAT
      ACTAATCTTTTTCTTCCTGATGTGAGATCGGCTTGAAAGTAGATAAACTTGTCCTATCTC
```

```
        TGTGGTCTAATTTTTTTATTTTAACTCTGTCACAGAATCACTTATAAAAGATCACAATAT
1255    TCTGTAGAATAATGGTATAAGATTATCTGTATGAGAAGAGGAAATATCTTCTTATGATAA
        AAAAAGAGTAACTCTCCTCTGGTAAAGAACCAGGAACTCTACATCAGTATGCAGATCTCA
        AGAATCTGAACCAGACTTTTTCTCATTCATGACTTTTTCTCAAGAATCTGAACCAATGAC
        TTTTTCTCATTCTAGACTTTTTCTCAAGAATCTGAACCAGACTTTTTCTCATTCATGACT
        TTTTCTCAAGAATCTGAAC
1260

        >TRINITY_DN80076_c8_g1_i1
        GGTGAGGCGAGAGGACCTCGACCCTCGACCCCATCGCAGGTCCAAACTGTGCTTATTTAC
        GCCTAAAACCACGTGGTCAGGGTGTAGCTGTATCTCGCATAAGTGGCCCCAAACAGTCCC
        AAGTAACGAACGATTTCGGAAGGAAGGAAGCAAGCCTCCCCGAGGAAACACGGGAACTAA
1265    AATGCGATGGGATATGGTCAAGTCTCAATGGTCAGTCAAATAGCCACTGATGATTTTCCT
        ACTGTTTTAAACATCGGGATAGCGGAGCATGATACTTCGGTACGTTCTTGGTTTATTCTT
        TGATTGAAGCAGAAGTTATCAAAGCAAGGTTATCGTTAGACGCAAGAAAGTTGCCTGAAA
        GGGATAATGTTATCATAGTGGTGCCATGGCGAGGTCTCAGGCAGAAGTGACAAGCAGACA
        GAATTTCTACCATCACAGGAGAGTGCAAGAATTGGAGAAGGATGACAAAGATGTTAATGA
1270    ACACGAGGTAGAAACAGAACATAGCGTCCCAGCTGATCTTTCTCCTCACGTCTCAGAAAT
        CAGAGATCACGTGAACGAGCACATTTTACATCACCACCATTCTAATGAGCACCGGGATAC
        AGAACAGGATTCTTTACTCCAGGACTTCCACGGACAGGGTGAACAGCGGGTAACACATTT
        AGCTAACCTCGATGAACAGGAGCGACGTCATCATGACCATTTAGATGCAGATGCAAAAGA
        CTCTAGTAATACAAGTAGTATCCCATCTTATAATCCGGAGTCGGGAGAAACTCCCAATAT
1275    CCATCATCCCCCTCATCATCATCATCAAGAGCATCTCTTTTCGGCAGATGATAACCAACC
        AAATCACATTCATATCCAGCATGAGGACCAACATCACCATCATCATCACTCCACCCATCA
        CTCTCATCCTCATCAGAATATTTTGATGCATTCGTCGAATTTCCACAGTGCGCTCCATGA
        ACAACAGCAGCAGCATGACGAAGATTCAGCAATACCTAATCACATGAATGATGATAATCA
        TCATCTCTCCCATCTTGAAGACGTAGCAGGAGAGGGAGCTGAAACAGCCCAAATGCATCC
1280    TCATATTTCTCACCAGCCTCATCATCTTCACGCCTCTCATCATCACCTTCAGGATCAAGA
        CGATCCAACAAACTCTCACAACGCTTTACAGCATCACACCAGAGATATTCTTCACCATGC
        GGAGAAGGAACTACAAGCCCACATGGATTCACCGGGAGAAGATCCTCGTATAAGCCTTGA
        CCTCCCTGAGGTTCACGGATCATTCCCGCACATTCGAAGCCCCACCTCTCCAAGAGATTA
        CATCCAGGATCGTCTTGCTACATATACTTTGCCCTCGGACTACAGAACTTTAGCAAGTAT
1285    AAATTCTTCTGATCCTCAAACAATTGAATCTTCACCGCTGAGAAATATGCTGTTCCCCTC
        TATGGGTTATTTGTCATCGTCACCCTTAACTTCTCAAAGGGATATTGGTGTTGATTCAGC
        ATTACATACCGTTAGTACACCTTCTGTGACATATCATCATTTGCCAGTAGCATCGGAATC
        ACCTGTTAGTCAAAGTGTTGCTAGTTCTTTATACGTGCCAACGGTATCTACACCCTCTAA
        ACTGTTAGGCTTGTCATCTTACGGAAGAATTCACGAGCATCCAAGCTCAGGAGCCGGGCA
1290    CTCTCTAATGTGGAGTCAGCTGAGTGAAGATCTTTCCCCTAAAACACAAATGTCCCCTTC
        GTCATCATCAGTATTATCCAGAAATGCTACAGTCGGCCATTCTTCTTCATACGTCAGCCA
        AGATATTTCCTCATGGGGCTATGACCCGAGTTCAGCAAATATGACTGCTCTTCAGTTGTC
        AATGGCATCTTCGCACTCTCCACCAGATGGCGACATGTTTGGCAGCCTAGAGAGCCGCGA
        ATGTGTCAATTGTGGTGCTATTTCGACGCCATTATGGAGGCGGGACGGAACTGGTCACTA
1295    CCTATGTAATGCATGTGGGCTTTACCATAGGATGAATGGCATGAATAGGCCTGTAGTCAA
        GAATCAAAAGAGGCTGTCAGCTTCCCGAAGAATGGGTCTGTACTGTTCTAACTGCCAAAC
        AACGAATACTTCTTTATGGAGAAGGAACAGTCAAGGGGAACCAGTGTGCAATGCCTGTGG
        ACTGTACTTCAAGCTTCACAGGAAGACATGAAATACAGTTTAACAGGAAGACCAATGGAG
        ACAACAAGTTTGTTAAATTCCTCGCAGTCATATTTCGTGTATACCAATACCAGTACAGGA
1300    CAAAGGACTCTTCCAGTCTTCACGTCAATGAAAGACTTGCCGAAATCTTCTTCCACTATG
        CATATTGTGAGTCCACCTTTACTTCATGGCGGTGGCAACACTAGTAATCTTCCTTTCTTG
```

```
CTTCAGACAGGAAGTAACACTGTGGTTGGATCACCGTCTTCCCCCAAGCATGAGCAAGGC
TCGGACTATCAGATTTCGTCAACTGGACACGCCACGTTACAGCAGGCGCATGTGCAAATT
GCAGGAACAGAAGGAACTCCCAATGACTTTGACCAACAACATGATCTGCTGCCGTGATTG
GTGAAAGATTCTGGACGGTCACCCGCTTTCTTGATGACATTTTCTACGCCTGTCAAAAAA
CGAGGGGAAAAATGCCATGTCAGTTGTCGTGTCAAAATGTGTGCGTCGTTTTAAAAATGT
ACATAAAATGAGTTAGAGGAGAATACGACGTGTAAGGGGATTAAGACTAAATCGTCTAAA
ATGTATGTATTTTTTTGTTTTTCGTTCTGTTTTTTTTTTTTTTTTTTTTTTATGT
```

1310 *araucan*

```
>ara.DN78359_c3_g1_i2
CGGGAGACGCCGTCGAACGCGAGCGGTCGTCTGCAACCGGAGTAACGCCAGACCCAGTTC
AGGCGTGGCGAAAGCGCAAGCGCTCGTCCGATCGAAGCTTAGTATCTTCCAGTCGGCGGC
ACAACGCTCAGCGGAGCAGGAACTGCTACCACACACCGCATCGAAGACCATTCCCGCGTC
CGTGCGATCTTCACTGAGCTACCCACCCCCTCCTTTTTTTTTCCATTGTCGCGCTTTAAA
AAAAAAAGGAAATCAGTTGGACAATGGACATCTAGTGGTCTTTCGTTTCTCCTCTAGGT
TAACAACAAACGGAATACTCAAATAAATCGTCACTTTCTGTGCGAAAGGTAAGCGATTTC
GTAGCAATTTCTCTCGAGTACATATTCGAAAATTACAAAACTGAAAGAGACGAATATCTT
AGAAGAAAAAAATACTTGTGACGAAGAGTTGGTGGATTTTTTTCATGCAAAAGAACAGCT
CATTAATGCATTTATTTGCCATCATGGAAAGTGCACTGTGAAAAATTATATCATCTTTCT
GAAAGTTCAAAAAGGGCTGAAACAATTTTTCCTGAAAAAAAAAAAAAACACCTTTAATAAT
TAAGCTTCTTTTTTTGCCAGTTAATGTGAACAGTATCTGCAAGATCTGTAATAAGTTGTT
CATGGTAGACAATATACCTGGAAAAATACAAGAAAATTACAAATTTGAAAAAAAAAACCC
TTCAATGTTGTAACATTACAATTAGAATTTTTCCTGTCATGCTGTAACTAAAACAGTGTT
TTGAATTTTGTCTCGGAGCATGTGACTTAACTGACACTTTTCGGACAAAACTTGTTACTA
ATTTAAAAAGTTTAAACTTCCTTAAAAGGTGAATTTGAAAATTTAAAAAGACCATGTGAT
CTTGAAGTAGCAGCCAATAGTTGCTGCTGGTTGTAAGAAATCGAAAGAAATAGACTGAGC
GACTTTCTGAGAGAGAGAGAGGAAAAAACAAAAAAAAGCACATCCCCTCCCCCTCTAAAT
GCGTCGACGATTTCCTCTAAAACTGTTGGACGCATTTTGCTTCATGGCGTGACATCAGCT
TGCTTAGACATCTTATGTTCTGCTATTTCTCACATTGCGTGACTAGAGTGAAAATGGTCT
TGAGTTAAAGGATTTTGTCGCTGAACGCCCACCACGAATCAATGGACTCTGGCATGAGAG
TTTCTACTTCTTGAAGATGTCATGTCCTCAATTTGGTTATAACGGGTCACCTTCTTCAGC
GCAGGAAAAATAAAATTGAATTGAATTGAACCGAGAGATAGAACAGACTCTGGGCATTCA
TTCGACACAGCTGCCAGAAATCCTGCACGAACATTTACGCCTCAGGAAGGTTTCGTGCCG
TATACAAATGCATCTTCATCTGTACCCTGTTCATCTCTTTTGATGGCCGGGCAGTCGGCG
ACAGGGCAGCAATTGCATGGTGGGTCCGGAAGTGGCACGGGCCAGCCGTGCTGTGAATCC
GGTCGCCCCGTGCTGACGGACCCCCTCACGGGGCAGACGGTGTGCTCGTGCCAGTACGAC
GCCCAATTGCTTAGCTACCAGCGGTTGGCTGCTTCGGGGTTGCCGCTCAATATGTACGGC
ACGGCGGCTGCAGCAGCGGCCGCGGCGGCGTACGGCGGAGAGCAAGGCTTCCTACCGCTC
GGAACTGAACAGTCAGCTTTCTACTCGCCATCGGCAAATGGCTTTGATCTAAAAGACAAT
CTAGAGGCTTGGCGGAGTTTGCCATACGCAGCTTCTATGTATTATCCGTACGACTCTGCA
GCACTAGCTGGTTATCCTTTTCCTAACGGGTACGGCGTCGACCTCAACGGTGCACGGAGG
AAAAACGCCACGCGAGAGACGACGAGCACTTTGAAGGCGTGGCTGAATGAGCACCGGAAG
AATCCTTACCCCACAAAGGGCGAGAAGATCATGTTGGCTATCATCACCAAGATGACCCTT
ACCCAAGTGTCCACGTGGTTCGCCAATGCGAGGAGGCGGCTCAAAAAAGAGAACAAAATG
ACCTGGTCTCCTCGTAATCGTTGCGACGACGACGATGCCGATCCCGACCCGGAGGACACG
GAGAGACCGAAAGATGTCGACAACAAAGGCCGCGACAAAGATGGCGTTGATGCGGATCGG
```

```
CAGAGGAGGGATGCCGGGGATAAGGAACCCGACACGAGGCAAGGAGGTTCCGATGATGGC
AAAGTTGATTTTGTCGACGTTGACGACGATTGCAAGTCATGTGCATCAACAAGTACTGAT
GGCCATATTATTAACTCTTCTGACAAAGAGAACTGCGTGTTAAGTGCTAGTGGTAAACCA
AGAGTTCCCAGTCCTGTATCAGATGCCTCCCTCGCCTCAGACTCTGAGTCCTCTCAGGCT
TCGGTACTAATGCCCGGTGTTGTAATTAGCAGTCAGCAGTCAACGGAACTTCCGGCCAAG
CCACGGATCTGGTCTATAGTGGACACAGCAACCTCAAACACTTCTGCGCCGAGCACGTCA
GTGAACTCTCCTCCCAGTGGTGGTGCTTCTAGACTAAGTCCCCTGCAGAGGGCAGCTTAT
CTTAGCCCATATCCTAAACCTACATCGTGGTATGCTGGAACTTTAGGTAACCTAAGTGCA
GTGACCAGTGGAGTAGGAGGAAGTTTCCCGTTGTCAGCAGCAGCCTTCACTTGTCCGCCA
ATAGTATCTTCGATGACAGGTTCTGTTGCGTCCTCTGGTTCATCGTCCCTTGTGGAATCC
CCTGCAGTGTCTTCGCTAAATAGACTAAGGACCGCAGCGGCGGCATTTCCGACAGCATCT
TCCGGAGAGCTGTCGTTACACAAGGCCGCTGCTGTCGCAGCTGCAGCAGCAGCAGCTGCG
GCGTCACCAACGGCTTCTCACACCCCTTCGGTATTCAACCTAAGCACTAAAGTTTCTGGA
GGTCATTCCAGCTCGCATGTATCTTCAGGATCAGCGGCGGCCATAGCTGCAGTGGCTGGA
AGTGGTAACACAGTGTCTTCCTCGAGCAGCGGAATATCGTCTTCGGCCAGCAGTCAAGGG
AGAGTGACTTCGTGAAAGCGAATTTTTCTCGGTGATTCTAACTGTACAGTATAAAAAAAA
AGATATGAAATAAAATTTCAGGAATGTAAATGTGTACCATATACTGTTGGGAATGGTTCA
TTTGTTTTCTGATGATGAGACCAATGTGATTCGCTGAAGGATCTCTTTTACATGCCGTGG
AAGATCTGCAAAGTTCCAAAGTGACCCATCGTTCAGTAAACCTCCTATGAATTTCGTACA
GAGCAAACTACTAAGCACAAAACAAACACTTTAATCAGACTTCACAATTCCCTCAGCCTG
TATCAGGAAAAACATTTTCTATAATTTCCTGCAGCATTTGTGCAACTACAATGAGTTGTA
AATATCTTTCCTGAGAATGTACAGCGTCTAATTTTGCAACTCTCAGGAACGAATTACAAT
TCGTCACTATGACTGCATTGAACAACTGAGAGAAAGAAAGAATTGCAGCGCCAATCAGAT
TCGAACACTACTGTACACATGAATGGACTAGATTAGATAAGGAAAAGAAAGACCTTATGT
AAAATTGTATTATAATATTGCAAAAGAGGAAATTATAAATTGTACTTCACAGTGTTTTGG
CCCACCATCTGAGAAAGCCAAAATACTTAAAAGAAATCGATTAAACAGGCCAAATTGTTA
TCGTTAAATTAATGCAGGATATAAGATATATCGTTATGATCGACATTATGCTTTTTCTAA
ACCAAAGTTGTACATACTTTCTAATAAATATATGTGTACAAACGTTCAGCTTCGAGAACG
TAAGCATGTTCTCCAAGAAGATTAAAAGAATAAATAAAAAGAAGTTATGACACGTAGAAC
CTGAATATCATCATGGGTCATTGATTCCTAAACTCCTCCCCCAGCGGGCGCTGGGAATTA
TTTATTATTTCTTGAACATTTAAGCAGGGTGTATAATATAGATCAGGCGGTATGATACCG
CCCCTGATTTCACTGAGGAAACCCCGTGTCACCCTCATTTCCTGTGCTGCATATCCATCA
AGTTATATGAACATGTGATTATGTCCAAATGTCAAAAAAAAGTAATGACAAACTGTTTAT
GTCACAATAAAAGAGCAGTGTTTCGAAATAAGGAAAAAAAAA
```

odd-skipped

```
>Acanthoscurria_odd-skipped_TRINITY_DN78272_c8_g1_i1 len=1979
AGAGAATTGTAAAAAGTGTCGTTCTCTAATTATTATTGAGTGATTTAAATGTGCTACAAT
GCCACTTTATTGTAAGAACAGAGGGAAGTTATTGTGTGGGCATTCAAATAAGTACACTGT
AGTAATAAAAGGTATAACTGTGAATTTTCAGGATCACCATATAAGATATACCCAGCAGAA
AAAAATGTATTCCATTACAAGTCGTTTTATACGTAATAAATTTTATGTTCCATTCGTTTT
CCTATGACATGTTGATTGTGGCATTTTGTATCATAAATACCTTATCTTTGGTATATACAT
ATATATATTGAAGCACCAGTTTTCATGACCAAGTACCATGATCTCCATTTTTTTTCACAT
ATATATAATTTCAAAATGTTAACTGTATAACCAATTAACTCTTGTTTGTCTTACATATAG
AATTTAATTCAAATATACAGTAAAGTCCACCAGGGATTTTGATTGTCCTTTTTCCTTTAT
ACATAAATATTCTAGAACTGGCATGTTCACAAACTACCTGCAGATGGGTGAACCAGATTT
AGCGGATGATTATACGAAGTTACATCGACCACAGCGTCGTCAGAAGACTGATGAGGTAGT
```

1395
```
CCATCACTTCGAAGGCTATGTGTGAGGCTGTGCCTCCGAAGATCGCAGTTCCGTCGGAAC
ACCTTCCCACAGGAATCGCAGTTGTATGGTTTAATGTCCGTGTGAGTGAGGAGATGGGTT
TTCAGGTTACTCCTTTGATTAAAGGTTCTGCCGCAGGTGGGACACTTATGAGGAGAGTCT
TCCATGTGGAGTATCCTGTGTACAGCTAAAGTCCGAGATTGGCAGAAGCCTTTGCCACAC
TCCGTGCATTTGAATGGTTTCTCTTTCGAATGAATGTACCTGTGATCTCTGAGATGGTCC
```
1400
```
TGTCTTCTGAATGCTTTGTGGCAGATGTCACATGTGTAAGGTCGTTCATCTGTGTGAGTC
CTCTCATGGATAAGTAAGTTATAGGATTTGGTGAACCTCCTTTGACAATATTTGCAGATG
AATTCTTTCTTTGGTCTCGATGACACCCTACCGCGGAAAAACCTCTTCTCGAACAACAAG
GAACCATTGGCACTCCCAAGAGGTGATTTCCTAAATATGCCAGCAATGGCTGGATTGTAG
AGGTGACCGAGAAAAAGAGGATGGGACGGGAAGATGGCTGGTTCTTGAAGCGGAGTAACT
```
1405
```
GTTTTTGGCACCTCGGAGGCTACCAGACTGTGAGGTGGGGATTTGACAGAGCTGGGATCT
TCAGCATCCTCCTCATCAGATCTCGTGGCTGACTCTGCCAGATGAGCAAAATCGAATTTC
CTTTTCTTCTTCTTTGGCGATGTGGGTGTTTTAGGCTGCGGAAGGACGACATCCTGTGGC
GGCACGGGTCTCGGAGGCAATCCGGAATAGCTGACGTCGCCACAGTCGTACGGCGACGGA
GGGAGATAGGCTGTGAAGGCCGATCCGCTGATGTTCAGGTGAAGAGGAATCGGGACGACG
```
1410
```
GGAGCCTGAGGAAGCATAGTCGTCGTTACCGCAGCAACAGCGGCCGTCGTCTCCCGTTGC
GGTGGTGGCCAGGTAGTGTTGACAACCGTCGGACGAAACAAGGGTGGGGAGGGGCTGCTG
TGACTACTCGTCGGCGTGAGAGGAGGCGTCGGCAGGAACCTTTGGTGGTGATGATGATGC
TGATCTCTACGTTCATCTTCTAGCAGGAGCGTGATGCTATTTGCAGCAGCAGCAGAGTCC
GGCATGGCGACTGCTGGGGTGCGGCGGGGTTACTCACACACTCTGGGGGCAAGCCAGCCC
```
1415
```
AGGCGGAATTCGGCAGTCTTCAGGAACAGCAGCAACAGGAGCCGGTTCCAAAGTGGTCCT
GTGTCGCACAGGGCAGTCAGCTACTGACTGGTAAAGCGAGCAGCAAGAGAGCAGCGGTCC
GGGACCGCGGAGGGGGAAGAATTGCTCTGCTAGAACCCGGATGTTGGCGTGGCTCAGCG
```

Distalless

```
>Dll_DN74055_c5_g1_i4
```
1420
```
AGCCTATCCTGGTAGTTTCAAGTACACGTTCGTCGACGTCCGACACGCGGGTGTGCAGAG
CAGGGCAAAGCAATCTTCCTTCTTTTTCTCTCAGTCGGGGTCCCGCTAAGCCGCCACCAC
GTTGAATTTTACGGAGTTCTCGGCCCCCTACCCGAGTATGGCGGGTACTGCGGATGGCCT
TGATCAGGATGTGACCAACAAGTCTGCATTCATGGAGATACAACAGCAGGGTCTGGCCGG
TATGCCTCATCATGCCCAAGGTTACCCTATCCGTTCGACCTACTCAAGCCAGCCGACGCA
```
1425
```
GCACGAGGCTGTGTTCGCCACGTCCCAACACCCCAGGCCGCTGGGGGCTTATCCGTTCCC
TATGAACAACTCACCTTTGAATACAAGTTTGCACAATTCCTACACACACCCAAGTCATCC
ATATCTCAGCACATATCCAGCAAATGTTCCTGGTTGTCCACCGTGTCCTTCACCTCCAAG
AGACGATAAGTCACAGTTGGAGGAAACACTCCGAGTGAATGGTAAAGGTAAAAAGATGAG
AAAACCGCGCACAATCTACTCCAGTCTACAGCTACAGCAACTGAACAGGAGGTTTCAGCG
```
1430
```
GACACAGTACCTTGCCTTACCGGAGCGGGCTGAACTCGCAGCGTCCCTGGGACTTACGCA
GACACAGGTGAAGATATGGTTCCAAAATAGGAGGTCCAAGTACAAGAAGATGCTCAAAGC
TCAACAGCAGCAACAGCAGCAAAACCCCCAAGCACCAGGACCGAACCAGACGCCCGGAGC
TACGCCGAATCTACAGCCGCCCAACCCAGCGTCAACGCCGCAGACCCCGCCTGAATCCCA
CGACGCCCACACACCTCCCTTAGCACCGCTCATGAATCCACAGCAGCAGCAGCCGCCGCC
```
1435
```
ACCGCAGCAGCAGCAACCTCCGCCGCAGGGACCCGGTGTCATGCCGACTTCGGCCGTCAG
TCCTCCTGCTATGAGTCCTCCCATCTCATCCTGGGACATGGCATCTTCCAAGGCAGCAGC
AATGAACACTTATATGCCTCAGTACTCTTGGTACCATCAGACGGACCCCTCAATGAGCCA
ACAAATTCTCACTTAATCCATTTTAAAACAAATAAGAATCATTTTTCCTGAGATGACGAA
AATGAAGTGTGAATGAATATATTCGTGTGTGTTCATACAAAGGCTTCCCCCCCCACTTCAG
```
1440
```
ATGGGTGGGTTGATGCATCACACGTTGGACATCGGCTGCCAACATCCCAGCCTAGTATCA
TCACAGTATTATGTGACGGTGCGGTGTCATGAAGTGCCGAAGAAGAAGAAGGGCAGCCTG
```

```
        TAAACAGCAACCCGTGCTCAGTCGGACGGCAGGATCGTGGACCGAAACATCCAAACGCCA
        GCCAGTCAGCCAGCCAGTCAGATAAAGAAACCAGTGTCACCCATTTCACAACCCTTCCTA
        ACAATCGTCGCAAACGACCTACTCCCCAGGCCCTTTTCTCCCCGAGTCCCATTCACAGAG
1445    GGACTTTACATCAACTAATATCAGCCATTTTAAAAATTCATCAGCTTCATTTGTGGAACT
        TCTTTGGCCAGTACTATATATACTTCTATGCCACAGTTCAACAGCCAAATGTAAACTTGA
        GAGGTAACAAGTCTTTCTTCGTTGTGTCCAAAGGGAACTGCATCCAACTCAAGAGAATGT
        CAGCAAACCGCACAGGATTGATGATCCCTGCTACACATCTAGTGAGATATTTCCAAAAAC
        ATCATGCCAGAATTATTAATTTATCTCGTACTAAGCACTGTCTTGGATATGTAATATCTA
1450    CTTGTTGTAAAGAAGATGGCAGTGTAGTTTTTGTACAAATGCTTCTTCGAGGTTTAGCTA
        TTTAAAATAAAAAAAAACAAAAACTCCATCTCTCAAGAAATAAGAAGTCTGCGACAGAAA
        TTGTAATGTATATTTAGTACAAATGATATCAATCATTCTTTTTCTTCATGTAAACAATAC
        AAAAATTGTTACTGAAACCGATTATATTATGCTAATTGTGAATGAATCTGTATCTATTTG
        TAAAATATTTTTATGGAAATTATTATAAGGGACAAGAAACTATATTGAACCTGATAAATA
1455    AGCGTACATATTATGCCTGCCTCAAAATTACTTCTTTTTCATGAATTAAGCCGCGAGATT
        TAATTTGTATAAAAGTCCACAAAATTGCGGCCGCTTGACATGCTCATGATAAATGCCACC
        ATTGCTGAGAAATGAATTCCAGTTACAATCGGAATATTCAATGATATTTTTACCCTCCCT
        TTCAATCGGAATTATTTCCATAACGCTTGGCCTCATGGATGACAGTTTCATCAATAAATG
        ATCAGGCACTGTTACAGATATCGCCAATCTTGCATCATACTAAAAAAAAAAAAAACTAATA
1460    CAAGTTCATATTCATTGCAACTCGTTCAACCCAGAAGAAAACAGCAGCACAGACGAATGA
        TTTAGTTTTAAATATGCCACAAATCAGGTAACAGATTCAAAATTACACGGAAGCAAGAGT
        TGTACCAAGTTTTCTTACCTGAAAAGATTACGAATGGTAAATAAAATTCTTCTGTGTAAA
        TAAGTGGTCATATGTTCGAGCTAAGTCATATCTATAAGACAAATTAAATAGCATAAATAC
        ACTGAATTATGAATCATCCCGCATTAGTACACAAGAGCAGTTTATATTTTAATTACCGTA
1465    AGAAATATACAGAAGAGTGGTGGCAGAACTGCGGAATAACTTTAACACAATTAATTAATA
        AAAATAAAAAAAAATCATACAAGATATTTTTCTTACAACGTTAGTTACTGGTAAAAGTAA
        GCATGGTAATTATGCTTAACACCAAATAATACAGTAGACAACTATTTCGCCTAATATCAA
```

1470   **Fig. S1.** Three orthologs of *pnr* were identified in *Acanthoscurria* with closest homology to *Drosophila*, *Tribolium*, and *Parhyale pnr* (Fig. S1-2). However, only one of these was expressed at the stages examined, Acanth_DN77604, and was presumed to be *pnr*. Consensus tree generated using Mafft, which gave similar topology to Clustal consensus tree.

1475

Fig. S2. In situ HCR of separate *pannier* and *Distalless* channels showing no overlap of expression in legs.

a

dorsal

lb

sto

ch

5 * 1

pnr
ara
Dll
odd-r3
DAPI

b

dorsal

lb

sto

ch

5 * 1

ara
DAPI

c

dorsal

lb

sto

ch

5 * 1

pnr
Dll
DAPI

Fig. S3. Leg segments in *Acanthoscurria* chelicerae. The expression of *pannier, araucan, odd-skipped-related-3,* and *Distalless* suggests the presence of 3 leg segments in the chelicerae, segments 1, 5, and 8.

1485   **Fig. S4.** Expression of *drm* (*odd-skipped 2*) in *Parhyale* St 22 embryo reproduced from Kira O'Day's thesis[67]. In the full-length thoracic legs, seven stripes are visible (six strong stripes and one proximal, fainter stripe that is most visible in the legs on the right), which separate 8 leg segments.