**Deconvolving clinically relevant cellular immune crosstalk from bulk**

**gene expression using CODEFACS and LIRICS**

Kun Wang[**,1], Sushant Patkar[**,1,2], Joo Sang Lee[1,3], E. Michael Gertz[1], Welles

Robinson[1,2], Fiorella Schischlik[1], David R. Crawford[1,4], Alejandro A. Schäffer[1], Eytan

Ruppin[#,1]

[1] Cancer Data Science Laboratory, National Cancer Institute, NIH, Bethesda, MD

[2] Department of Computer Science, University of Maryland, College Park, MD

[3] Department of Artificial Intelligence & Department of Precision Medicine, School of

Medicine, Sungkyunkwan University, Suwon, Republic of Korea

[4] Department of Cell Biology and Molecular Genetics, University of Maryland, College

Park, MD

[**] These authors contributed equally

[#] Corresponding author:

eytan.ruppin@nih.gov

Key words: deconvolution, immunotherapy, tumor microenvironment, ligand-receptor

interaction, cell-cell interaction, microsatellite instability

## Abstract

The tumor microenvironment (TME) is a complex mixture of cell-types that interact with each other to affect tumor growth and clinical outcomes. To accelerate the discovery of such interactions, we developed CODEFACS (COnfident DEconvolution For All Cell Subsets), a deconvolution tool inferring cell-type-specific gene expression in each sample from bulk expression measurements, and LIRICS (LIgand Receptor Interactions between Cell Subsets), a supporting pipeline that analyzes the deconvolved gene expression from CODEFACS to identify clinically relevant ligand-receptor interactions between cell-types. Using 15 benchmark test datasets, we first demonstrate that CODEFACS substantially improves the ability to reconstruct cell-type-specific transcriptomes from individual bulk samples, compared to the state-of-the-art method, CIBERSORTx. Second, analyzing the TCGA, we uncover cell-cell interactions that specifically occur in TME of mismatch-repair-deficient tumors and are associated with their high response rates to anti-PD1 treatment. These results point to specific T-cell co-stimulating interactions that enhance immunotherapy responses in tumors independently of their mutation burden levels. Finally, using machine learning, we identify a subset of cell-cell interactions that predict patient response to anti-PD1 therapy in melanoma better than recently published bulk transcriptomics-based signatures. CODEFACS offers a way to study bulk cancer and normal transcriptomes at a cell type-specific resolution, complementing single-cell transcriptomics.

**Introduction**

The importance of the tumor microenvironment (TME) in cancer has been recognized since the late 1800s[1]. The recent success of immune checkpoint blockade has further sparked interest in studying TME interactions that shape clinical outcomes following immunotherapy, aiming to find biomarkers of treatment response and new treatment opportunities[2]. One key step in studying these interactions is the characterization of the molecular profiles of different cell types in a patient's tumor sample. Fluorescence-activated cell sorting (FACS) and single-cell RNA sequencing have emerged as effective tools to address this challenge[3]. However, due to the cost of these procedures and scarcity of fresh tumor biopsies, the application of these approaches has remained limited. Given that bulk tumor gene expression from preserved biopsies is far more abundant, computational methods that can effectively extract cell-type-specific expression from such data, termed *deconvolution* algorithms, could be very helpful. If successful, such deconvolution methods can markedly advance our knowledge of the TME across many tumor types and different contexts, and beyond that, they may be readily applied to interrogate other large bulk expression datasets.

Several previous studies have developed a variety of expression deconvolution algorithms. DeMixT[4] was designed to estimate individual-specific expression for three cell components provided prior reference samples of two of these cell components. ISOpure[5] has aimed to derive sample-specific cancer cell expression with the assumption that the observed bulk gene expression profile is a mixture of predefined stromal and immune cell expression profiles that are shared across all the samples. Building on this work, Fox et al extended ISOpure to predict individual-specific non-tumor cell expression by subtracting cancer cell profiles from the bulk mixtures in a two-cell type model[6]. More recently, Newman et al[7] developed CIBERSORTx, the first approach that aims to predict the *sample-specific* gene expression of all cell types composing it by employing a set of novel deconvolution heuristics. As a proof of concept, Newman et al showed that CIBERSORTx can accurately reconstruct the cell-type-specific expression of genes in each input sample under certain modelling assumptions. This groundbreaking work has, however, some notable limitations: (1) The number of genes whose cell-type-specific

expression can be reconstructed in each sample is relatively small, especially for low-
abundance cell types, and (2) their approach does not provide confidence estimations of
the predictions made, while such estimations could be potentially useful in most
deconvolution applications in the absence of ground truth data.

Here, we introduce a new deconvolution algorithm and software, CODEFACS
(COnfident DEconvolution For All Cell Subsets), which markedly advances the ability to
successfully deconvolve bulk gene expression data. CODEFACS receives as input bulk
gene expression profiles of tumor samples and either pre-computed estimates of
abundance of expected tumor, immunological and stromal cell types in each sample, or
their prototypical molecular signatures, which serve as seeds for estimating the
abundance of each cell type in each sample. CODEFACS then predicts the cell type-
specific gene expression profiles of each sample. It is a heuristic approach aimed at
maximizing the number of genes in each cell type whose expression across the samples
can be confidently predicted via a heuristic method to estimate confidence. Using 15
benchmark datasets where the ground-truth is known, we show that CODEFACS robustly
improves over CIBERSORTx, both in terms of gene coverage and the individual gene
expression estimation accuracy. We additionally developed LIRICS (LIgand Receptor
Interactions between Cell Subsets), a pipeline that integrates the output of CODEFACS
with a database of prior immunological knowledge that we curated to infer the active cell-
cell interaction landscape in each sample. These data can then be analyzed in
conjunction with any sample-associated clinical annotations (e.g., response to treatment)
to infer the most important clinically relevant immune interactions between the cell types
in a given patient's cancer cohort.

Building on its enhanced coverage and accuracy, we next applied CODEFACS to
reconstruct the cell-type-specific transcriptomes of ~8000 tumor samples from 21 cancer
types in TCGA. Analyzing these fully deconvolved TCGA expression datasets using
LIRICS we find a shared repertoire of intercellular interactions enriched in the TME of
mismatch repair deficient tumors of different tissues of origin, which is associated with
improved overall patient survival and high response rates to anti-PD1 treatment,

independently of their mutation burden levels. Finally, using machine learning techniques,

we identify a subset of intercellular TME interactions that are predictive of response to

immune checkpoint blockade treatment in melanoma patients.

In summary, CODEFACS and LIRICS present a new way to analyze large bulk

RNA-seq datasets to study cellular crosstalk in the TME of each patient, and to learn

more about the association of different tumor-immune interactions with different clinical

measures. The potential scope of applications of both CODEFACS and LIRICS goes

beyond studying the TME, as these tools can be applied to study any disease of interest

given bulk gene expression data and relevant reference signatures of cell types involved.

**Results**

**Overview of CODEFACS and LIRICS**

CODEFACS is designed to characterize the tumor microenvironment by reconstructing

the cell-type-specific transcriptomes of each sample from bulk expression. It takes as

input the bulk RNA-seq expression values of a cohort of tumor samples and either the

estimations of the cell fractions of a pre-defined set of cell types in each sample or their

cell-type-specific molecular signature profiles, derived based on reference datasets or

from the literature[7].

CODEFACS then employs a heuristic approach that sequentially executes three

modules: *(module 1) high resolution deconvolution, (module 2) hierarchical deconvolution*

*and (module 3) imputation*. Each module is designed to predict the cell-type-specific

expression of genes in each sample; the second and third modules aim to overcome the

shortcomings of the previous modules. For a more detailed description of each module

and its modeling assumptions, see **Supplementary Note**. A key component of

CODEFACS is its *confidence ranking system*, which receives cell-type-specific

expression predictions from the different modules and labels them as high or low-

confidence estimations. Genes whose expression is determined with high confidence in

a given module are added to the output set, while low confidence predictions are

continued to be processed in subsequent modules (See **Fig. 1A** and **Supplementary**

**Note**). The final output of CODEFACS consists of two items: (a) a three-dimensional gene
expression matrix, where each entry represents the predicted gene expression a gene in
a given cell-type in a specific sample, and (b) a two-dimensional matrix of confidence
scores ranging from [0,1] representing which gene-cell-type pairs have most confident
predictions (~1) and which pairs have least confident predictions (~0; See **Fig. 1A,**
**Output**). These scores can be further investigated to assess the quality of predictions for
a given dataset.
140
Given fully deconvolved gene expression data from CODEFACS, one can use
LIRICS (LIgand Receptor Interactions between Cell Subsets) (**Fig. 1B**) to transform this
data into a biologically interpretable feature space of active ligand-receptor interactions
between cell types in each sample. Specifically, LIRICS takes the output of CODEFACS
and processes it in three steps: (step 1) The first step queries a database of all plausible
ligand-receptor interactions between any two cell types A and B, that we have
systematically assembled and curated from the literature. This database is publicly
available as part of LIRICS (see **Supplementary Note**, supplementary tables 1-3). (step
2) In the second step, given the deconvolved expression profiles of cell type A and cell
type B in a given bulk tumor sample, LIRICS denotes as 'active' or 'likely to occur' ('1')
the interactions where both the ligand and receptor are over-expressed in the relevant
cell-types in that sample, or otherwise 'inactive' ('0'). A ligand or receptor is considered to
be overexpressed in a given cell type if its expression exceeds the median expression in
that cell type (**Supplementary Note**). (step 3) Finally, a Fisher's enrichment analysis is
performed to test the association of the activity of specific ligand-receptor interactions
with any relevant phenotypes of interest (e.g., treatment response, mutational subtype,
etc.) (**Fig. 1B**). Furthermore, if required, one can collectively analyze the binary profiles
returned by LIRICS from multiple independent datasets to augment sample size and
increase statistical power. Finally, one can apply a false discovery rate (FDR) cut-off post
multiple hypothesis correction and visualize significantly enriched interactions in a
volcano plot as shown at the bottom of Fig 1B, or, alternatively, display their network
structure as shown in subsequent biological applications.
163

A

**CODEFACS**
*COnfident DEconvolution For All Cell Subsets*

**User Input**

Bulk tumor tissue

Samples

Bulk expression Matrix

+

Cell fractions

**Cell types**
- Tumor cell
- B cell
- T cell
- NK cell

| Module 1 | Module 2 | Module 3 |
|---|---|---|
| High Resolution Deconvolution | Hierachical Deconvolution | Imputation-Based Deconvolution |

**Confidence ranking system**

Expression & Confidence Matrix

1 Accurate
0 Less Accurate

**CODEFACS Output**

Individual specific cell type expression

Samples — Genes — Tumor cell
Samples — Genes — B cell
Samples — Genes — T cell
Samples — Genes — NK cell

+

Confidence Matrix

Cell types — Genes

1
0

B

**LIRICS**
*Ligand-Receptor Interaction between Cell Subsets*

Tumor cell

CD8 T cell

**Step 1**
Query plausible ligand-receptor interactions between two cell types

**Step 2**
Inferring activity of queried interactions in each sample using deconvolved gene expression

Samples

Interactions

- Likely to occur
- Unlikely to occur

**Step 3**
Cell type specific clinically relevant interactions inererence

Any clinically relevant phenotype

Interactions

Samples

- Likely to occur
- Unlikely to occur
- Case
- Control

-log(P-value)

Fold change

**Figure 1. Overview of CODEFACS and LIRICS. (A) CODEFACS** takes bulk gene expression profiles and prior knowledge of the cellular composition of each sample and executes a heuristic three step procedure to infer the deconvolved gene expression in each sample. In module 1, we perform a high-resolution deconvolution, which extends the CIBERSORTx algorithm. In module 2 (hierarchical deconvolution), bulk expression is modeled as a mixture of two components: a specific cell type of interest and all the remaining cell types. The expression for the cell type of interest is predicted by removing the estimated expression in the second component (using high-resolution deconvolution from module 1) from the bulk mixture. In module 3 – imputation-based deconvolution, we impute the cell-type-specific expression of a specific gene based on the predicted cell-type-specific expression of other high-confidence genes that are co-expressed with that gene in the bulk. Each module is designed to overcome the shortcomings of its predecessor based on their respective modeling assumptions. The confidence ranking system is responsible for classifying all the predictions at the end of each module into high-confidence or low-confidence predictions. Genes classified into the low-confidence class at the end of one module (e.g. module 1) are passed to the next module (e.g. module 2) for refinement. Finally, after all the three modules are executed, the prediction confidence levels are re-evaluated. The final output of CODEFACS consists of a 3-dimensional matrix with cell-type-specific gene expression predictions for each sample, along with estimated confidence scores of predictions for each gene in each cell type. For more details, see Supplementary note. **(B) LIRICS** takes the output of CODEFACS and processes it in three steps. In step 1, for each possible permutation of cell type pairs, LIRICS queries a literature-curated repository for enumerating all plausible ligand-receptor interactions between specified cell types. In step 2 this prior knowledge is integrated with the output from CODEFACS to infer which of the plausible cell-cell interactions are likely to occur or be "active" in each individual sample. The result is a binary matrix with rows representing each plausible cell-cell interaction and columns representing each patient's tumor sample. Finally, in step 3, given any clinically relevant phenotype (e.g. response to therapy, driver mutation status, etc..), one can perform a Fisher's enrichment analysis (shown at the bottom) to discover cell-grounded receptor-ligand interactions in the TME that are associated with the phenotype of interest. For more details, see Supplementary note.

**Evaluating the performance of CODEFACS**

To assess the accuracy of CODEFACS, we generated 15 benchmark datasets (see Methods) by merging publicly available single cell RNA-seq[8,9] and FACS sorted purified RNA-seq[10]. Thereafter, we applied CODEFACS to deconvolve these generated bulk datasets and define the accuracy of its predictions by computing the Kendall correlation between the predicted and ground truth expression in each cell type across individual samples (the Kendall correlation provides a less inflated measure of accuracy by accounting for ties in the data). In the main text, we show the results obtained on three benchmark bulk datasets: one derived form a FACS-sorted lung cancer data, one from a single cell melanoma RNA-seq data and from a single cell glioblastoma RNA-seq dataset, respectively. Each bulk sample from these datasets represents a real biopsy from a patient. We show that CODEFACS can predict the cell-type-specific expression of more genes than CIBERSORTx (with Kendal's correlation ≥ 0.3) (**Fig. 2A-C**), and its predictions are overall more accurate (**Fig. 2D-F**). The results for all the remaining 12 benchmark datasets, created via artificial mixing of single cell profiles and single cell RNA-Seq imputation (See Methods and **Supplementary Table 6** for more details), also show the superiority of CODEFACS (**supplementary Fig. S1 and S2**). Overall, we observe that the more abundant the cell type is, the better CODEFACS can predict its cell-type-specific gene expression (**supplementary Fig. S3**).

Next, we quantified how well the confidence scores it returns align with the Kendall scores that measure the true prediction accuracy with the ground truth for each (gene, cell-type) pair to validate the claim that our confidence scores can help the user filter out potentially noisy predictions in real bulk datasets. We quantified this using two metrics: Spearman correlation and a classification AUC (Area Under the ROC Curve). Across all the benchmark datasets analyzed, the Spearman correlation between confidence scores of genes in each cell-type and their corresponding Kendall scores (quantifying the true prediction accuracy) is strong and positive (**Fig. 2G** depicts the results for the FACS sorted lung cancer benchmark dataset, and **supplementary Fig. S4** depicts the results for the remaining benchmark datasets); To perform a classification-based quantification, we grouped the genes in each cell-type into two classes based on the correlation between

their predicted and actual expression, informative (prediction accuracy ≥ 0.1 and p-value
≤ 0.05) and uninformative (prediction accuracy < 0.1 or p-value > 0.05). We then tested
whether the confidence scores could be used to classify genes into these two classes for
each cell type. We find that the confidence score could effectively filter out uninformative
predictions (**Fig. 2H** depicts the results for the FACS sorted lung cancer benchmark
dataset, and **supplementary Fig. S5,** depicts for the remaining benchmark datasets).
206
Finally, to further evaluate CODEFACS on real bulk tumor data (where the ground
truth is unavailable), we applied it to deconvolve bulk expression data from 21 cancer
types (~8000 RNA-seq samples) in TCGA. To infer the cellular abundance of each cell
type in each sample which is required as input for CODEFACS, we made use of matched
bulk methylation data available for these samples and methylation-based reference
signature profiles of distinct cell types. These include 11 cell type signatures
(macrophages/dendritic cells--CD14+, B cells--CD19+, CD4+T cells, CD8+ T cells, T reg
cells, NK cells--CD56+, endothelial cells, fibroblasts, neutrophils, basophils, eosinophils
and tissue-specific tumor cells) obtained from MethylCIBERSORT [11]. Reassuringly, we
found strong Spearman correlations between the resulting predicted tumor cell fraction
and the tumor purity estimates derived from matched mutation and copy number data
(based on ABSOLUTE) for the same samples across 10 cancer types (Spearman
correlation: min=0.72, max=0.88, avg=0.8; **supplementary Fig. S6**). This testifies that
methylation-based cell fraction estimates indeed form a reliable basis for running
CODEFACS to deconvolve TCGA samples.
222
We then asked if CODEFACS can recover the expected cell-type-specific gene
expression signature of different cell types in a given cancer type. To this end, we
computed the Spearman correlation between (a) the mean deconvolved gene expression
of the top confidently deconvolved genes in a given cell type (confidence score ≥ 0.95)
and (b) the mean expression of these genes, which we derived from completely
independent single cell expression data of the same cancer type (Methods). We find that
(a) and (b) are substantially correlated (**Fig. 2I** depicts results for the TCGA-LUAD (lung
adenocarcinoma) dataset as an example and **supplementary Fig. S7** for the remaining

cancer types that have publicly available scRNA-seq data). The concordance level is

higher for cell types that are abundant (e.g., tumor cells and fibroblasts) and decreases

for less abundant cell types. Additionally, we observed that tumor cells have the largest

fraction of genes whose expression is predicted with high confidence, with the highest in

thyroid cancer (THCA, ~67.4% of all genes). Furthermore, 7 KEGG pathways are

significantly enriched (adjusted p-value < 0.01) with highly confident genes in tumor cells

(confidence score ≥0.95) across the 21 cancer types (**supplementary Fig. S8**). Those

pathways mostly involve RNA transport, spliceosome, DNA replication, and mismatch

repair.

**Figure 2. Evaluating the performance of CODEFACS. (A-C)** bar plots depicting the number of genes with a prediction accuracy (Kendall correlation) ≥ 0.3 with the ground truth for each cell type, estimated from bulk-generated samples of lung cancer (LUAD dataset; sample size = 26), melanoma (SKCM dataset 1; sample size = 28) and glioblastoma (GBM dataset 1; sample size = 24) benchmark datasets, as estimated by CODEFACS (yellow bars) and CIBERSORTx (blue bars). **(D-F)** boxplots depicting prediction accuracy distributions of all genes across different cell types in the lung cancer (LUAD with sample size 26), melanoma (SKCM with sample size 28) and glioblastoma (GBM with sample size 24) benchmark datasets, using CODEFACS (yellow) and CIBERSORTx (blue). A two-sided Wilcoxon signed rank test was performed to compare the prediction accuracies of CODEFACS and that of CIBERSORTx for each cell type in each dataset. *** denotes p-values < 2e-16. **(G)** Spearman correlations between prediction accuracies and confidence scores among cell types in the lung cancer benchmark dataset (LUAD dataset; sample size = 26). The y-axis indicates the spearman correlation coefficient value, while the x-axis indicates the cell type. **(H)** AUCs obtained in classifying informative and uninformative predictions among cell types in lung cancer benchmark dataset (LUAD dataset; sample size = 26). **(I)** bar plots depicting the Spearman correlations between mean deconvolved cell-type-specific expression in TCGA-LUAD and mean cell-type-specific expression derived from publicly available single cell datasets of LUAD. The y-axis indicates the Spearman correlation coefficient value, while the x-axis indicates the cell type. Details of all single cell RNA-Seq based pseudo bulk datasets can be found in supplementary table 6.

**Tumors with DNA mismatch repair deficiency have heightened T cell co-stimulation**
**that is independent of their tumor mutation burden levels**
In normal cells, DNA is constantly repaired in response to DNA damage or DNA
replication errors[12]. However, defects in specific DNA repair pathways in cancer cells may
result in the accumulation of many somatic mutations resulting in hypermutated tumors
(TMB ≥ 10-20 mutation/Mb)[13–15]. One of the sources of hypermutability is a mismatch
repair deficiency (MMRD), which leads to the accumulation of insertions and deletion
mutations in microsatellite regions of the genome due to uncorrected DNA replication
polymerase slippage events. This is known as microsatellite instability (MSI)[16,17]. Solid
tumors with mismatch repair deficiency were shown to be sensitive to immune checkpoint

blockade (ICB) therapy irrespective of tumor type, leading the FDA to approve MSI as the
first cancer type agnostic biomarker for patients receiving anti-PD1 treatment[18]. The
reason behind this general sensitivity to anti-PD1 treatment is not completely understood.
Prior work has led to the prevailing hypothesis that elevated tumor mutation burden in
mismatch repair deficient tumors leads to more neoantigens, and thus is more likely to
activate a host immune response against tumor cells[16,19–21]. However, not all tumor types
with elevated tumor mutation burden have similar response rates to anti-PD1[22,23], and
recent studies have revealed that T cells recognize and respond to only a few neoantigens
per tumor[24–27]. More generally, when looking at non-synonymous tumor mutation burden,
MSI and survival data across the TCGA collection (**Fig 3A,B,** borrowed from[28,29]. See
Methods), we see a significant association between hypermutability and survival benefit
of patients in solid tumor types with a frequent underlying mismatch repair deficiency (**Fig.
3C.** log-rank test p-value = 0.00084), in contrast to other tumor types (**Fig 3D**, log-rank
test p-value = 0.4). These survival differences can be partially explained by the mutual
exclusivity of microsatellite instability and chromosomal instability, which has been
previously linked with a worse prognosis[28,30,31]. Taken together, these findings motivated
us to further study cellular immune crosstalk in the tumor microenvironment of mismatch
repair deficient tumors to gain additional cell-type-specific insights into their sensitivity to
anti-PD1.

**Figure 3. (A)** The **landscape of tumor mutation burden and microsatellite instability across 18 different solid tumor types.** This panel plots the distribution of non-synonymous tumor

mutation burden on a logarithmic scale (Y-axis). All points above the horizontal line are typically
regarded as hyper-mutated tumors (> 10 mutations/Mb). All red points represent tumors with a
DNA mismatch repair deficiency detected via microsatellite instability (MSI). **(B) This panel**
**depicts the percentage of all tumor samples per cancer type with microsatellite instability**
**(Y-axis).** Tumor types marked with a * represent those where MSI is prevalent. **(C-D)**
**Comparison of overall survival of patients with tumors that are hypermutated vs not**
**hypermutated.** Left panel (**C**) In Gastro-Intestinal+Endometrial tumor types where MSI is
prevalent (marked with a * in panel B). Right panel (**D**) In other solid tumor types where tumors
rarely have an underlying mismatch repair deficiency. Statistical significance of differences in
survival was calculated using the log-rank test.
288
We hence aimed to identify cell-cell interactions that are differentially active
between microsatellite instable tumors (highlighted as red dots in Fig 3A) and
microsatellite stable tumors (highlighted as black dots in Fig 3A). To this end, we applied
CODEFACS to deconvolve the bulk gene expression of all solid tumors from TCGA and
integrated their predicted cell-type-specific gene expression levels with LIRICS. The top
50 interactions from this differential analysis (ordered by FDR adjusted p-value) are
shown in a network in **Fig. 4A**. These interactions are frequently active in mismatch repair
deficient tumors of distinct tumor types (**Supplementary Fig. S9**) and importantly, they
are more frequently active in hypermutated tumors with DNA mismatch repair deficiency
compared to other hypermutated tumors (**Fig. 4B, Fig. 3A**), testifying to their MSI
specificity. The top 50 MSI-specific interactions include the PDL1-PD1 checkpoint
interaction between tumor cells and CD8+ T-cells, but notably, T-cell activating/co-
stimulatory interactions such as the 41BBL-41BB interaction between Tumor cells and
CD8+ T cells, ULBP2-NKG2D between tumor cells and CD4+T cells, and chemotaxis
interactions involved in trafficking of lymphocytes in and around the tumor mass, such as
the CXCL9-CXCR3 chemokine interaction between macrophages and CD4+ T cells and
CCL3/4/5 – CCR5 interactions between various immune and stromal cell-types.
306
This shared heightened cellular crosstalk unique to the TME of mismatch repair
deficient tumors suggests that tumor infiltrating T cells can be activated by co-stimulatory
signals in the TME independent of overall tumor mutation burden, only to be kept in

balance by other immunoregulatory mechanisms such as the PD1-PDL1 checkpoint
interaction between CD8+ T cells and tumor cells. Our results indicate that when this
interaction is blocked by anti-PD1 treatment, the presence of other co-stimulatory
interactions can lead to the observed enhanced response of MMRD tumors to immune
checkpoint blockade therapy. This in turn raises the possibility that switching on specific
T cell co-stimulation signals in the TME may lead to better responses to anti-PD1
treatment independent of tumor mutation burden. Notably, recent pre-clinical studies have
shown that combination therapies aimed at enhancing such T-cell co-stimulating
interactions improve anti-tumor immune responses even in low TMB and highly immuno-
suppressive settings [32–37]. Currently, several clinical trials to assess the safety and
efficacy of these combinations are in progress [38–41].

**Figure 4. (A) Interaction network consisting of the top 50 interactions most highly activated**
**in TME of tumors with DNA mismatch repair deficiency.** Interactions highlighted in green
represent co-stimulatory interactions/having an activating effect on the target cell. Interactions
highlighted in red represent checkpoint interactions/having an inhibitory effect on the target cell.
Interactions highlighted in black represent pro-inflammatory/chemotaxis interactions involved in
inflammatory response and immune cell trafficking to tumor sites. Eos: Eosinophils, CAF: Cancer

associated fibroblasts. **(B) A volcano plot depicting on the x-axis the log2 fold change in the**
**frequency of occurrence of each cell-cell interaction in the TME of hypermutated tumors**
**with an underlying DNA mismatch repair deficiency vs other hypermutated tumors.** The y-
axis indicates the -log10 FDR adjusted p-value of the observed enrichment. Highlighted in red in
the scatter plot are the top 50 interactions that are most differentially active between all MSI vs
non-MSI tumors (shown in panel A).

**Machine learning guided discovery of cell-type-specific ligand-receptor**
**interactions that predict patient response to ICB therapy**
Given the small number of large publicly available transcriptomics datasets of patients
receiving immune checkpoint blockade therapy, we asked if we can effectively utilize this
large resource of deconvolved TCGA data we generated to transfer-learn cell-cell
interactions robustly predictive of response to immune checkpoint blockade therapy.
Specifically, since some mutations during tumor evolution can be immunogenic, we
hypothesized that one could potentially discover cell-type-specific ligand-receptor
interactions predictive of response to ICB therapy by a joint analysis of mutation and
deconvolved expression data from the TCGA. We focus on melanoma, currently the
tumor type best responding to ICB, where there are many independent publicly available
bulk expression datasets of patient's receiving anti-PD1 treatment and single-cell derived
cell-type-specific signatures, which serve as priors for the deconvolution of these bulk
datasets. Starting from the deconvolved TCGA-SKCM dataset as our training set (N=469),
we employed a genetic algorithm to find cell-type specific ligand-receptor interactions,
whose activation state best separates hypermutated melanoma tumors from non-
hypermutated tumors (**Fig 3A**), assuming that the former group captures more samples
with some immunogenic mutations than the latter (see Methods, **Fig 5A**). We term the
interactions identified in this process *melanoma mutation specific functional interactions*
(*MSFI),* and the network formed by these interactions is displayed in **Fig. 5B***.*

Having identified the MSFI interactions, we  applied CODEFACS to deconvolve
the bulk expression data of pre-treatment samples from the three largest publicly
available melanoma datasets where patients received anti-PD1 treatment (either
monotherapy or in combination with anti-CTLA4; Methods) [42–44]. We then employed

LIRICS to the respective deconvolved expression of each of these checkpoint datasets,

*without any additional training*, and simply quantified the number of MSFI interactions that

are active in each of these patients' tumor samples, which we denote as the tumor's *MSFI*

*score*. Remarkably, we find that the MSFI score of each sample can robustly stratify

patients into those that are likely to respond to ICB vs those that are unlikely to respond

(**Fig. 5C**, progression free survival log rank test p-value: 0.00057, **Fig. 5D**, overall survival

log rank test p-value: 0.0031). **Supplementary Fig. S10** depicts the survival differences

for the two treatment groups separately (anti-PD1 monotherapy and anti-CTLA4 + anti-

PD1 combination). Additionally, **Supplementary Fig. S11** and **S12** depict the survival

differences for each ICB dataset separately. As evident, our results improve over recent

bulk gene expression based predictors of melanoma ICB therapy response (IMPRES [45],

TIDE [46] and the melanocytic plasticity signature (MPS) scores [47]). We note that the

performance levels of the latter on bulk expression datasets, where their original RNAseq

reads have been uniformly aligned and normalized as described in the Methods, is lower

than that reported in the original publications, pointing to the potential sensitivity of

expression-based predictors to the processing used and the need to do that in a uniform,

generally accepted manner (see Discussion).

To further evaluate the predictive performance of the MSFI score, we tested its

ability to predict partial or complete responders vs stable or progressive disease patients

in these datasets. To this end we plotted the receiver operator area under the curve (AUC)

obtained using it for classifying the patients to partial or complete responders vs stable or

progressive disease, and compared its performance to that obtained with the three other

published predictors for the different treatment groups (**Fig. 5E**). On average, the MSFI

score achieves an AUC of 0.63 (for anti-PD1 monotherapy the AUCs obtained are 0.6,

0.77 and 0.52 for the three individual ICB datasets, and for the combination ICB treatment

its 0.77, 0.49 and 0.63). A similar performance could not be achieved if the placement of

the ligand and receptor between interacting cell-types in the MSFI network was swapped

(average AUC 0.58) or by randomly shuffling the interaction activity profiles (average AUC

~ 0.52), testifying that the selected cell-cell interactions are best predictive of response to

ICB. Comparing MSFI predictive performance in this response classification task to that

of the recent melanoma bulk expression-based predictors, TIDE achieves an average

AUC of 0.6 (for anti-PD1 monotherapy the AUCS are 0.46, 0.66 and 0.45 for the three

ICB datasets and 0.89, 0.55 and 0.61 for the combination), IMPRES achieves an average

AUC of 0.55 (for anti-PD1 monotherapy the AUCS are 0.59, 0.64 and 0.6 respectively for

the three ICB datasets and 0.61, 0.44 and 0.42 for the combination) and MPS achieves

an average AUC of 0.51 (for anti-PD1 monotherapy the AUCS are 0.61, 0.49 and 0.63

respectively for the three ICB datasets and 0.41, 0.59 and 0.38 for the combination).

Examining the MSFI network leads to interesting insights **(Fig. 5B)**. First, we find

an over-representation of cell-type-specific co-stimulatory/immune cell activating

interactions known from prior immunological literature (hypergeometric test p-value <

0.05)[48–58] (See Supplementary Table 1). Second, the MSFI network additionally includes

cytokine/chemokine interactions involved in pro-inflammatory response and the trafficking

of NK, T and B cells to the tumor site (responsible for better lymphocyte infiltration into

the tumor mass). Importantly, on fitting a multivariate Cox-proportional hazards model

with MSFI scores and TMB of each patient receiving anti-PD1 (wherever TMB data was

available), we see that a high MSFI score is significantly associated with improved

progression free survival (p-value: 0.013) and overall survival (p-value: 0.0258), whereas

high TMB is not (PFS p-value: 0.224, OS p-value: 0.477). These results further support

the findings from the TCGA analysis that heightened co-stimulatory and pro-inflammatory

signals in the TME can mobilize tumor infiltrating lymphocytes to generate an effective

host immune response upon immune checkpoint blockade independent of TMB.

**Figure 5. (A) Overview of the machine learning analysis employed to identify cell type specific interactions that are predictive of response to immune checkpoint blockade therapy. (B) A chord diagram of the resulting MSFI network.** Each individual interaction is represented by a link from the source cell type (ligand expressing cell type) to the target cell type (receptor expressing cell type) and the color of the link represents the color of the source cell type. For interactions that are activating/co-stimulatory, the sector in the corresponding target cell type is highlighted in green. For inhibitory/checkpoint interactions, the sector in the target cell type is in highlighted red. Interactions involved in chemotaxis are highlighted in black and those I

mediating a pro-inflammatory response are highlighted in blue, cell-adhesion interactions are

highlighted in grey. **(C) Kaplan-Meier plot depicting the progression free survival of the**

**combined set of melanoma patients receiving immune checkpoint blockade (N= 244).** On

the top, the patients are stratified into low-risk/high-risk groups based on the median value of

MSFI score from LIRICS. Second from top, patients stratified into low/high risk groups based on

median IMPRES score[45]. Third from top, patients stratified into low/high risk groups based on

median TIDE score[46], Bottom, patients stratified into low/high risk groups based on median MPS

score[47]. **(D) Kaplan-Meier plots depicting the overall survival of all melanoma patients**

**receiving immune checkpoint blockade (N= 244).** On the top, the patients are stratified into

low-risk/high-risk groups based on the median value of MSFI score from LIRICS. Second from

top, patients stratified into low/high risk groups based on median IMPRES score[45]. Third from top,

patients stratified into low/high risk groups based on median TIDE score[46], Bottom, patients

stratified into low/high risk groups based on median MPS score[47]. Survival differences among

patients that received anti-PD1 monotherapy vs anti-CTLA4 + anti-PD1 combination are shown

in supplementary figure 10 **(E) Area under the ROC curves in predicting Complete/Partial-**

**response (based on RECIST v1.1) to immune checkpoint blockade therapy for the different**

**scores.** X-axis marks patients grouped by dataset source and treatment regimen. PD1 mono

represents patients that received anti-PD1 monotherapy. PD1 + CTLA4 represents patients that

additionally received anti-CTLA4 besides anti-PD1.

## Discussion

This study presents a computational tool, CODEFACS and a pipeline, LIRICS, that

enables an (averaged) 'virtual single cell' characterization of the TME from bulk tumor

expression data. Applying these tools, we identify cell type specific ligand-receptors

interactions that are active and functionally important within individual tumor

microenvironments, in modifying patients' survival and response to ICB. Applying

CODEFACS to ~8000 tumors from TCGA, we estimate the cell-type-specific gene

expression profiles of each individual tumor sample thus enabling the analysis of the

TCGA at a cell-type-specific resolution. Integrating these data with LIRICS, we

systematically characterized the immune cellular crosstalk of the tumor

microenvironments of different tumor types. We identified a shared core of intercellular

TME interactions in DNA mismatch repair deficient tumors, which are associated with

improved patient survival and high sensitivity to immune checkpoint blockade therapy.

One potentially interesting implication of these findings is that immunomodulators
enhancing T-cell co-stimulation (e.g, via the 41BB receptor) might improve patients'
response to ICB irrespective of their tumor mutation burden. Finally, focusing on
melanoma, we show that one can bootstrap on the large deconvolved data resource from
TCGA using machine learning techniques to discover cell-cell interactions within the TME
that successfully predict patients' response to immune checkpoint blockade.

While we have provided a toolkit to discover clinically relevant cell-cell interactions
from bulk tumor expression, there are limitations that should be noted and potentially
further improved upon in the future. First, several recent publications have reported
discrepancies between different RNA-seq expression quantification methods based on
the reference transcriptome version used and choice of method (alignment based vs
alignment free)[59–63]. This can potentially affect reproducibility of findings. Hence,
whenever possible, we recommend that all bulk RNA-seq datasets are homogeneously
pre-processed using the same RNA-seq quantification method and reference
transcriptome before the application of our tools (See Methods), and any clinically
relevant findings are further followed up. Indeed, following this notion, in this study we
have pre-processed all bulk RNA-seq datasets in a uniform manner, using STAR (v2.7.6a)
+ RSEM (v1.3.3) and GENCODE v23 human genome annotation (See Methods for more
details).

Second, regarding limitations of CODEFACS itself (1) it requires prior information
about the cell type composition of the input tumors, or alternatively, knowledge of the
pertaining cell-types' gene expression or methylation signatures that can be used to infer
their abundances, and its accuracy depends on the accuracy of the latter. (2) its prediction
power is limited to subsets of the whole exome genes, and its performance deteriorates
for lowly-abundant cell types. However, the confidence scores provided partially alleviate
this limitation, allowing the user to rank genes in each cell-type by the quality of
predictions of the expression of each gene in a given cell-type. Third, regarding LIRICS,
it currently restricted to well defined ligand-receptor interactions between tumor, immune,
stromal and epithelial cell types and does not consider the spatial localization of cells in

the TME. The inclusion of the latter with the advent of forthcoming spatial transcriptomics
data is likely to lead to considerably more informative interaction inference approaches.

Although this work focuses on studying the tumor microenvironment, the tools
presented here can be applied to discover important cell-cell interactions in noncancerous
tissues under a variety of normal and disease states.  One interesting application that we
envision is the characterization of clinically relevant intercellular interactions occurring at
the maternal-fetal interface using corresponding bulk gene expression data and
pregnancy outcome information, whose elucidation may help treat and mitigate
preeclampsia and other pregnancy related complications. One can also use our tools to
study bulk gene expression data from pre-malignant tissue samples and compare them
against malignant samples to elucidate cell-cell interaction dynamics on the way
malignancy.  Finally, one can use our tools to deconvolve expression data from
autoimmune disorders to learn more about the underlying immune interactions. In sum,
the computational tools developed and presented in this study offer a cost-effective way
to study immune responses at a cell type specific resolution from the widely abundant
bulk gene expression in both health and disease, complementing first-line single cell
technologies in situations where the latter are less applicable.

**Data availability**
All deconvolved expression data generated in this study will be made publicly available
upon publication.

**Code availability**
The tools and all codes for reproducing the figures of this study will be made publicly
available upon publication.

**Methods**
**Data collection and pre-processing**
**Single cell RNA-seq datasets:** To benchmark the performance of CODEFACS, we first
set out to obtain publicly available single cell RNA-seq datasets where both tumor and

non-tumor cells were successfully isolated. This search led us to the identification of nine

such single cell RNA-seq datasets from the literature, each from a different cancer type.

Collection of additional single cell datasets was frozen after Dec 2019. Details of the

single cell datasets that we curated for this study in **Supplementary Table 4**. For each

dataset sequenced on the SmartSeq2 platform, the log normalized transcript counts for

each gene in each sequenced cell were made publicly available by the original authors.

For the application of deconvolution, these counts were transformed back to the

Transcripts Per Million (TPM) scale. For datasets sequenced on the 10x platform, UMI

counts for each gene were made publicly available and were scaled by the library size of

each cell and multiplied by a factor of 1 million to get expression values in TPM scale.

**Bulk RNA-seq datasets:** Gene expression and matching bulk tumor methylation data

from fresh frozen tumor biopsies in TCGA were downloaded from (http://xena.ucsc.edu/)

[64]. In addition, publicly available bulk expression data from formalin fixed paraffin

embedded tumor biopsies of melanoma patients receiving immune checkpoint blockade

treatment were downloaded from[42–44]. All bulk RNA-seq datasets were collected such that

they have a sufficiently large sample size to reliably perform complete deconvolution of

expression profiles (> 4 times the number of cell-types involved) [7]. Collection of datasets

was frozen after Dec 2019. Details on bulk RNA-seq datasets deconvolved in this study

can be found in **Supplementary Table 5**. To maintain consistency with the pipeline used

for preprocessing TCGA data, bulk gene expression levels in immune checkpoint

blockade datasets were re-quantified using STAR v2.7.6a and RSEM v1.3.3[65] with

GENCODE v23 human genome annotation[66]. Furthermore, to mitigate technical biases,

between-sample scaling factors were estimated using TMM method implemented in

edgeR[67] and TPM values in each sample were further rescaled by these scaling factors[68].

**Generation of simulated bulk RNA-seq datasets:** To evaluate the performance of

CODEFACS, we generated 14 different pseudo-bulk RNA-seq datasets from mixing

experiments with single cell data. Each sample in each benchmark dataset has matching

cell type specific gene expression profiles derived from averaging single cell RNA-seq

profiles of individual cells from the same sample and same cell type. These profiles serve

as the ground truth for the evaluation of deconvolution performance. To avoid any

circularity in our validations, for each of the single cell datasets involved, single cell data

from 4 randomly chosen patients were separated from the rest. These data were used to

derive reference gene expression signatures for each cell type. The mixing experiments

were then performed on single cell data of the remaining patients that were hidden from

the reference signature derivation process. In addition, we simulated technical replicates

for each pseudo-bulk sample, wherein we injected noise in the pseudo-bulk expression

of a few randomly chosen genes and then renormalized the expression data by the

sample library size. This procedure simulates mRNA composition noise that is commonly

observed in bulk RNA-seq datasets due to technical differences in sample preparation

[64-66]. The datasets and the mixing experiment used to generate the pseudo-bulk

samples are listed in **Supplementary Table 6**. In addition, we obtained a FACS sorted

lung cancer dataset which include purified RNA-seq for four cell types [10] and generated

a pseudo bulk correspondingly. In total, 15 benchmark datasets were generated.

**Deconvolving immune checkpoint blockade (ICB) and TCGA bulk RNA-seq**

**datasets using CODEFACS**

For the application of CODEFACS, molecular profiles of signature genes of each cell type

of interest are needed to estimate the relative cell fractions in the bulk. We used single

cell expression derived signatures as priors to deconvolve the melanoma ICB datasets.

To derive these signatures from single cell data, we first start out by obtaining the class

labels of each cell type of interest. These data are publicly available for each single cell

dataset we collected. Hence, we primarily use these labels in our study (unless further

refinement of labels into specific cell subtypes of interest is needed for a specific usage).

With a collection of single cell expression profiles and matching cell type labels as input,

we used CIBERSORT online tool to derive a cell-type-specific signature matrix.

Thereafter, we applied CODEFACS to ICB datasets with default parameters settings and

batch correction requirement specified. For TCGA deconvolution, we first estimated cell

fractions based on bulk methylation and then applied CODEFACS to corresponding bulk

gene expression for the 21 cancer types which have both types of data available. We

chose methylation signatures over expression-based signatures for TCGA analysis for

two reasons. First, single cell expression data with consistent cell types across 21 cancer

types are not available. Second, DNA methylation-based signatures are considered to be

more stable marks of cellular identity compared to dynamic RNA expression derived

signatures [69]. The methylation-based cell type signatures (**Supplementary Table 7**)

were obtained from MethylCIBERSORT [11]. We applied CODEFACS to TCGA datasets

with default parameters settings and without batch correction requirement specified. For

the details of CODEFACS algorithm and parameter settings, see Supplementary note 1.

**Machine learning for discovery of cell-cell interactions predictive of clinical**

**responses to immune checkpoint blockade in melanoma**

We used a genetic algorithm, which is a randomized heuristic search algorithm designed

to select optimal features for a prediction task given some user-defined fitness function

for training [70]. In this setting, the features are ligand-receptor interactions between cell-

types, the prediction task is predicting response to ICB treatment and the fitness function

is defined as the accuracy of predicting a user defined phenotype based on the total

number of interactions from those selected occurring in a given sample; accuracy is

quantified by the AUC. To aid in faster convergence of the genetic algorithm during

training, the size of the search space is reduced by first assessing the fold change of each

feature between the two classes specified in the training dataset and incorporated into

the fitness function (e.g., hypermutated vs non-hypermutated). After this step, only

features with a fold change >1 is passed to the genetic algorithm for further optimization.

The algorithm starts out by randomly generating sets of features. This is defined

as the seed population. These sets iteratively evolve via the phenomenon of natural

selection enforced by the user defined fitness function. Specifically, for each subsequent

iteration, features from the best performing sets, as determined by the user defined fitness

function, in the current iteration are mixed at random followed by random new feature

additions or dropouts (referred to as mutations) to build a new generation of feature sets

and the process repeats. Eventually, after a number of epochs, which we set to 100, the

fitness function converges to an optimum and the best set of features for the prediction

task is returned to the user. Since the fitness function landscape is often non-convex and

the training process is stochastic, we repeat the training process 500 times, each with a

randomly chosen seed population, and eventually choose frequently selected features

over all solutions to reach a solution we suspect is close to the global optimum solution.

For our plots, we set the threshold to frequency > 100 times. The probability of any feature

being selected more than 100 times by random chance based on this approach is

estimated to be < 0.01. Results are qualitatively similar for more stringent thresholds. The

genetic algorithm was implemented in R using the genalg package [71]

**Acknowledgements**

This research is supported in part by the Intramural Research Program of the National

Institutes of Health, National Cancer Institute, Center for Cancer Research. This work

utilized the computational resources of the NIH HPC Biowulf cluster. We would

additionally like to acknowledge Dr. Noam Auslander, Dr. Chi-Ping Day and other

members of the Cancer Data Science Laboratory for their helpful feedback on this work

**References:**

1.    Paget S. the Distribution of Secondary Growths in Cancer of the Breast. *Lancet*.

632            1889;133(3421):571-573. doi:10.1016/S0140-6736(00)49915-0

633    2.    Smyth MJ, Ngiow SF, Ribas A, Teng MWL. Combination cancer immunotherapies

634            tailored to the tumour microenvironment. *Nat Rev Clin Oncol*. 2016;13(3):143-158.

635            doi:10.1038/nrclinonc.2015.209

636    3.    Wagner A, Regev A, Yosef N. Revealing the vectors of cellular identity with single-cell

637            genomics. *Nat Biotechnol*. 2016;34(11):1145-1160. doi:10.1038/nbt.3711

638    4.    Wang Z, Cao S, Morris JS, et al. Transcriptome Deconvolution of Heterogeneous Tumor

639            Samples with Immune Infiltration. *iScience*. 2018;9:451-460.

640            doi:10.1016/j.isci.2018.10.028

641    5.    Quon G, Haider S, Deshwar AG, Cui A, Boutros PC, Morris Q. Computational purification

642            of individual tumor gene expression profiles leads to significant improvements in

643            prognostic prediction. *Genome Med*. 2013;5(3). doi:10.1186/gm433

644    6.    Fox NS, Haider S, Harris AL, Boutros PC. Landscape of transcriptomic interactions

645            between breast cancer and its microenvironment. *Nat Commun*. 2019;10(1):1-13.

646            doi:10.1038/s41467-019-10929-z

7.    Newman AM, Steen CB, Liu CL, et al. Determining cell type abundance and expression

from bulk tissues with digital cytometry. *Nat Biotechnol*. 2019;37(7):773-782.

doi:10.1038/s41587-019-0114-2

8.    Jerby-Arnon L, Shah P, Cuoco MS, et al. A Cancer Cell Program Promotes T Cell

Exclusion and Resistance to Checkpoint Blockade. *Cell*. 2018;175(4):984-997.e24.

doi:10.1016/j.cell.2018.09.006

9.    Ghoshdastider U, Naeini MM, Rohatgi N, et al. Data-driven inference of crosstalk in the

tumor microenvironment. *bioRxiv*. Published online January 1, 2019:835512.

doi:10.1101/835512

10.   Gentles AJ, Hui ABY, Feng W, et al. A human lung tumor microenvironment interactome

identifies clinically relevant cell-type cross-talk. *Genome Biol*. 2020;21(1):107.

doi:10.1186/s13059-020-02019-x

11.   Chakravarthy A, Furness A, Joshi K, et al. Pan-cancer deconvolution of tumour

composition using DNA methylation. *Nat Commun*. 2018;9(1):1-13. doi:10.1038/s41467-

018-05570-1

12.   Drake JW, Charlesworth B, Charlesworth D, Crow JF. Rates of spontaneous mutation.

*Genetics*. 1998;148(4):1667-1686. doi:10.1007/978-94-011-4385-1_17

13.   Loeb LA. Mutator Phenotype May Be Required for Multistage Carcinogenesis. *Cancer

Res*. 1991;51(12):3075-3079.

14.   Alexandrov LB, Nik-Zainal S, Wedge DC, et al. Signatures of mutational processes in

human cancer. *Nature*. 2013;500(7463):415-421. doi:10.1038/nature12477

15.   Muzny DM, Bainbridge MN, Chang K, et al. Comprehensive molecular characterization of

human colon and rectal cancer. *Nature*. 2012;487(7407):330-337.

doi:10.1038/nature11252

16.   Funkhouser WK, Lubin IM, Monzon FA, et al. Relevance, Pathogenesis, and testing

algorithm for mismatch repair-defective colorectal carcinomas: A report of the association

for molecular pathology. *J Mol Diagnostics*. 2012;14(2):91-103.

doi:10.1016/j.jmoldx.2011.11.001

17.   Cortes-Ciriano I, Lee S, Park WY, Kim TM, Park PJ. A molecular portrait of microsatellite

instability across multiple cancers. *Nat Commun*. 2017;8. doi:10.1038/ncomms15180

18.   Boyiadzis MM, Kirkwood JM, Marshall JL, Pritchard CC, Azad NS, Gulley JL. Significance

and implications of FDA approval of pembrolizumab for biomarker-defined disease. *J

Immunother Cancer*. 2018;6(1). doi:10.1186/s40425-018-0342-x

19.   Galon J, Costes A, Sanchez-Cabo F, et al. Type, density, and location of immune cells

within human colorectal tumors predict clinical outcome. *Science (80- )*.

2006;313(5795):1960-1964. doi:10.1126/science.1129139

20.    Llosa NJ, Cruise M, Tam A, et al. The vigorous immune microenvironment of

microsatellite instable colon cancer is balanced by multiple counter-inhibitory

checkpoints. *Cancer Discov*. 2015;5(1):43-51. doi:10.1158/2159-8290.CD-14-0863

21.    Le DT, Uram JN, Wang H, et al. PD-1 blockade in tumors with mismatch-repair

deficiency. *N Engl J Med*. 2015;372(26):2509-2520. doi:10.1056/NEJMoa1500596

22.    Yarchoan M, Hopkins A, Jaffee EM. Tumor Mutational Burden and Response Rate to PD-

1 Inhibition. *N Engl J Med*. Published online 2017. doi:10.1056/nejmc1713444

23.    Imakaev M. Limited evidence of tumour mutational burden as a biomarker of response to

immunotherapy. *bioRxiv*. Published online January 1, 2020:2020.09.03.260265.

doi:10.1101/2020.09.03.260265

24.    Robbins PF, Lu YC, El-Gamil M, et al. Mining exomic sequencing data to identify mutated

antigens recognized by adoptively transferred tumor-reactive T cells. *Nat Med*.

2013;19(6):747-752. doi:10.1038/nm.3161

25.    Bassani-Sternberg M, Bräunlein E, Klar R, et al. Direct identification of clinically relevant

neoepitopes presented on native human melanoma tissue by mass spectrometry. *Nat*

*Commun*. 2016;7. doi:10.1038/ncomms13404

26.    Roh W, Chen PL, Reuben A, et al. Integrated molecular analysis of tumor biopsies on

sequential CTLA-4 and PD-1 blockade reveals markers of response and resistance. *Sci*

*Transl Med*. 2017;9(379). doi:10.1126/scitranslmed.aah3560

27.    Kalaora S, Wolf Y, Feferman T, et al. Combined analysis of antigen presentation and T-

cell recognition reveals restricted immune responses in melanoma. *Cancer Discov*.

2018;8(11):1366-1375. doi:10.1158/2159-8290.CD-17-1418

28.    Taylor AM, Shih J, Ha G, et al. Genomic and Functional Approaches to Understanding

Cancer Aneuploidy. *Cancer Cell*. 2018;33(4):676-689.e3. doi:10.1016/j.ccell.2018.03.007

29.    Bonneville R, Krook MA, Kautto EA, et al. Landscape of Microsatellite Instability Across

39 Cancer Types. *JCO Precis Oncol*. 2017;(1):1-15. doi:10.1200/po.17.00073

30.    Davoli T, Uno H, Wooten EC, Elledge SJ. Tumor aneuploidy correlates with markers of

immune evasion and with reduced response to immunotherapy. *Science (80- )*. Published

online 2017. doi:10.1126/science.aaf8399

31.    Auslander N, Wolf YI, Koonin E V. Interplay between DNA damage repair and apoptosis

shapes cancer evolution through aneuploidy and microsatellite instability. *Nat Commun*.

2020;11(1). doi:10.1038/s41467-020-15094-2

32.    Chen S, Lee LF, Fisher TS, et al. Combination of 4-1BB agonist and PD-1 antagonist

716           promotes antitumor effector/memory CD8 T cells in a Poorly Immunogenic Tumor Model.

717           *Cancer Immunol Res*. Published online 2015. doi:10.1158/2326-6066.CIR-14-0118

718   33.   Ma HS, Poudel B, Torres ER, et al. A CD40 Agonist and PD-1 Antagonist Antibody

719           Reprogram the Microenvironment of Nonimmunogenic Tumors to Allow T-cell–Mediated

720           Anticancer Activity. *Cancer Immunol Res*. Published online 2019. doi:10.1158/2326-

721           6066.CIR-18-0061

722   34.   Barber A. Costimulation of Effector CD8+ T Cells: Which Receptor is Optimal for

723           Immunotherapy? *MOJ Immunol*. Published online 2014.

724           doi:10.15406/moji.2014.01.00011

725   35.   Melero I, Hirschhorn-Cymerman D, Morales-Kastresana A, Sanmamed MF, Wolchok JD.

726           Agonist antibodies to TNFR molecules that costimulate T and NK cells. *Clin Cancer Res*.

727           Published online 2013. doi:10.1158/1078-0432.CCR-12-2065

728   36.   Chester C, Sanmamed MF, Wang J, Melero I. Immunotherapy targeting 4-1BB:

729           Mechanistic rationale, clinical results, and future strategies. *Blood*. Published online 2018.

730           doi:10.1182/blood-2017-06-741041

731   37.   Philipson BI, O'Connor RS, May MJ, June CH, Albelda SM, Milone MC. 4-1BB

732           costimulation promotes CAR T cell survival through noncanonical NF-κB signaling. *Sci*

733           *Signal*. Published online 2020. doi:10.1126/scisignal.aay8248

734   38.   Siu LL, Steeghs N, Meniawy T, et al. Preliminary results of a phase I/IIa study of BMS-

735           986156 (glucocorticoid-induced tumor necrosis factor receptor–related gene [GITR]

736           agonist), alone and in combination with nivolumab in pts with advanced solid tumors. *J*

737           *Clin Oncol*. Published online 2017. doi:10.1200/jco.2017.35.15_suppl.104

738   39.   Tolcher AW, Sznol M, Hu-Lieskovan S, et al. Phase Ib study of utomilumab (PF-

739           05082566), a 4-1BB/CD137 agonist, in combination with pembrolizumab (MK-3475) in

740           patients with advanced solid tumors. *Clin Cancer Res*. Published online 2017.

741           doi:10.1158/1078-0432.CCR-17-1243

742   40.   Cohen EEW, Pishvaian MJ, Shepard DR, et al. A phase Ib study of utomilumab (PF-

743           05082566) in combination with mogamulizumab in patients with advanced solid tumors. *J*

744           *Immunother Cancer*. Published online 2019. doi:10.1186/s40425-019-0815-6

745   41.   Choi Y, Shi Y, Haymaker CL, Naing A, Ciliberto G, Hajjar J. T-cell agonists in cancer

746           immunotherapy. *J Immunother Cancer*. Published online 2020. doi:10.1136/jitc-2020-

747           000966

748   42.   Riaz N, Havel JJ, Makarov V, et al. Tumor and Microenvironment Evolution during

749           Immunotherapy with Nivolumab. *Cell*. 2017;171(4):934-949.e15.

750    doi:10.1016/j.cell.2017.09.028

751 43. Gide TN, Quek C, Menzies AM, et al. Distinct Immune Cell Populations Define Response

752    to Anti-PD-1 Monotherapy and Anti-PD-1/Anti-CTLA-4 Combined Therapy. *Cancer Cell*.

753    2019;35(2):238-255.e6. doi:10.1016/j.ccell.2019.01.003

754 44. Liu D, Schilling B, Liu D, et al. Integrative molecular and clinical modeling of clinical

755    outcomes to PD1 blockade in patients with metastatic melanoma. *Nat Med*.

756    2019;25(12):1916-1927. doi:10.1038/s41591-019-0654-5

757 45. Auslander N, Zhang G, Lee JS, et al. Robust prediction of response to immune

758    checkpoint blockade therapy in metastatic melanoma. *Nat Med*. 2018;24(10):1545-1549.

759    doi:10.1038/s41591-018-0157-9

760 46. Jiang P, Gu S, Pan D, et al. Signatures of T cell dysfunction and exclusion predict cancer

761    immunotherapy response. *Nat Med*. 2018;24(10):1550-1558. doi:10.1038/s41591-018-

762    0136-1

763 47. Pérez-Guijarro E, Yang HH, Araya RE, et al. Multimodel preclinical platform predicts

764    clinical response of melanoma to immunotherapy. *Nat Med*. 2020;26(5):781-791.

765    doi:10.1038/s41591-020-0818-3

766 48. Billadeau DD, Leibson PJ. ITAMs versus ITIMs: striking a balance during cell regulation.

767    *J Clin Invest*. 2002;109(2):161-168. doi:10.1172/jci14843

768 49. Staub E, Rosenthal A, Hinzmann B. Systematic identification of immunoreceptor tyrosine-

769    based inhibitory motifs in the human proteome. *Cell Signal*. 2004;16(4):435-456.

770    doi:10.1016/j.cellsig.2003.08.013

771 50. Murphy K, Weaver C. *Janeway's Immunobiology. Ninth Edition*.; 2018.

772    doi:10.1086/696793

773 51. Chen L, Flies DB. Molecular mechanisms of T cell co-stimulation and co-inhibition. *Nat*

774    *Rev Immunol*. 2013;13(4):227-242. doi:10.1038/nri3405

775 52. Campbell KS, Purdy AK. Structure/function of human killer cell immunoglobulin-like

776    receptors: Lessons from polymorphisms, evolution, crystal structures and mutations.

777    *Immunology*. 2011;132(3):315-325. doi:10.1111/j.1365-2567.2010.03398.x

778 53. Pende D, Falco M, Vitale M, et al. Killer Ig-like receptors (KIRs): Their role in NK cell

779    modulation and developments leading to their clinical exploitation. *Front Immunol*.

780    2019;10(MAY). doi:10.3389/fimmu.2019.01179

781 54. Ward-Kavanagh LK, Lin WW, Šedý JR, Ware CF. The TNF Receptor Superfamily in Co-

782    stimulating and Co-inhibitory Responses. *Immunity*. 2016;44(5):1005-1019.

783    doi:10.1016/j.immuni.2016.04.019

784    55.    Gonçalves CM, Henriques SN, Santos RF, Carmo AM. CD6, a rheostat-type signalosome

785            that tunes T cell activation. *Front Immunol*. 2018;9. doi:10.3389/fimmu.2018.02994

786    56.    Steri M, Orru V, Idda ML, et al. Overexpression of the cytokine BAFF and autoimmunity

787            risk. *N Engl J Med*. 2017;376(17):1615-1626. doi:10.1056/NEJMoa1610528

788    57.    Chen M, Lin X, Liu Y, et al. The function of BAFF on T helper cells in autoimmunity.

789            *Cytokine Growth Factor Rev*. 2014;25(3):301-305. doi:10.1016/j.cytogfr.2013.12.011

790    58.    Varin MM, Le Pottier L, Youinou P, Saulep D, Mackay F, Pers JO. B-cell tolerance

791            breakdown in Sjögren's Syndrome: Focus on BAFF. *Autoimmun Rev*. 2010;9(9):604-608.

792            doi:10.1016/j.autrev.2010.05.006

793    59.    Wu DC, Yao J, Ho KS, Lambowitz AM, Wilke CO. Limitations of alignment-free tools in

794            total RNA-seq quantification. *BMC Genomics*. 2018;19(1). doi:10.1186/s12864-018-4869-

795            5

796    60.    Sahraeian SME, Mohiyuddin M, Sebra R, et al. Gaining comprehensive biological insight

797            into the transcriptome by performing a broad-spectrum RNA-seq analysis. *Nat Commun*.

798            2017;8(1). doi:10.1038/s41467-017-00050-4

799    61.    Everaert C, Luypaert M, Maag JLV, et al. Benchmarking of RNA-sequencing analysis

800            workflows using whole-transcriptome RT-qPCR expression data. *Sci Rep*. 2017;7(1).

801            doi:10.1038/s41598-017-01617-3

802    62.    Teng M, Love MI, Davis CA, et al. A benchmark for RNA-seq quantification pipelines.

803            *Genome Biol*. 2016;17(1). doi:10.1186/s13059-016-0940-1

804    63.    Robert C, Watson M. Errors in RNA-Seq quantification affect genes of relevance to

805            human disease. *Genome Biol*. 2015;16(1). doi:10.1186/s13059-015-0734-x

806    64.    Goldman MJ, Craft B, Hastie M, et al. Visualizing and interpreting cancer genomics data

807            via the Xena platform. *Nat Biotechnol*. 2020;38(6):675-678. doi:10.1038/s41587-020-

808            0546-8

809    65.    Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq

810            quantification. *Nat Biotechnol*. 2016;34(5):525-527. doi:10.1038/nbt.3519

811    66.    Harrow J, Frankish A, Gonzalez JM, et al. GENCODE: The reference human genome

812            annotation for the ENCODE project. *Genome Res*. 2012;22(9):1760-1774.

813            doi:10.1101/gr.135350.111

814    67.    Robinson MD, McCarthy DJ, Smyth GK. edgeR: A Bioconductor package for differential

815            expression analysis of digital gene expression data. *Bioinformatics*. 2009;26(1):139-140.

816            doi:10.1093/bioinformatics/btp616

68.    Robinson MD, Oshlack A. A scaling normalization method for differential expression

analysis of RNA-seq data. *Genome Biol*. 2010;11(3):R25. doi:10.1186/gb-2010-11-3-r25

69.    Bird A. DNA methylation patterns and epigenetic memory. *Genes Dev*. 2002;16(1):6-21.

doi:10.1101/gad.947102

70.    Melanie M. An introduction to genetic algorithms By Melanie Mitchell. MIT Press,

Cambridge, MA. (1996). 205 pages. $30.00. *Comput Math with Appl*. 1996;32(6):133.

doi:10.1016/S0898-1221(96)90227-8

71.    Willighagen E, Ballings M, Ballings MM. Package 'genalg.' Published online 2015.