Deconvolving clinically relevant cellular immune crosstalk from bulk gene expression using CODEFACS and LIRICS

Kun Wang[**,1], Sushant Patkar[**,1,2], Joo Sang Lee[1,3], E. Michael Gertz[1], Welles Robinson[1,2], Fiorella Schischlik[1], David R. Crawford[1,4], Alejandro A. Schäffer[1], Eytan Ruppin[#,1]

[1] Cancer Data Science Laboratory, National Cancer Institute, NIH, Bethesda, MD

[2] Department of Computer Science, University of Maryland, College Park, MD

[3] Department of Artificial Intelligence & Department of Precision Medicine, School of Medicine, Sungkyunkwan University, Suwon, Republic of Korea

[4] Department of Cell Biology and Molecular Genetics, University of Maryland, College Park, MD

[**] co-first author

[#] Corresponding author:

eytan.ruppin@nih.gov

Key words: deconvolution, gene expression, immunotherapy, tumor microenvironment, ligand-receptor interaction, cell-cell interaction, microsatellite instability

Abstract

The tumor microenvironment (TME) is a complex mixture of cell types whose interactions affect tumor growth and clinical outcome. To discover such interactions, we developed CODEFACS (COnfident DEconvolution For All Cell Subsets), a tool deconvolving cell-type-specific gene expression in each sample from bulk expression, and LIRICS (LIgand Receptor Interactions between Cell Subsets), a statistical framework prioritizing clinically relevant ligand-receptor interactions between cell types from the deconvolved data. We first demonstrate the superiority of CODEFACS versus the state-of-the-art deconvolution method, CIBERSORTx. Second, analyzing the TCGA, we uncover cell-type-specific interactions of mismatch-repair-deficient tumors that are associated with their higher anti-PD1 response rates, including specific T-cell co-stimulating interactions that enhance immunotherapy response independently of the tumors mutation burden levels. Finally, we identify a subset of ligand-receptor interactions in the melanoma TME that predict patient response to anti-PD1 therapy better than recently published transcriptomics-based methods.

Introduction

The importance of the tumor microenvironment (TME) in cancer has been recognized since the late 1800s[1]. The recent success of immune checkpoint blockade has further sparked interest in studying the various cellular states of cell types and their interactions in the TME, aiming to find biomarkers of treatment response and new treatment opportunities[2]. One key step in elucidating the different cellular states of cell types *de novo* is the characterization of the molecular profiles of each cell type in *each patient's tumor sample*. Fluorescence-activated cell sorting (FACS) and single-cell RNA sequencing have emerged as effective tools to address this challenge[3]. The use of FACS is limited because it can only measure expression of a few protein markers simultaneously. The use of single-cell RNA sequencing has been limited due to its cost and the scarcity of fresh tumor biopsies. Since bulk tumor gene expression from preserved biopsies accompanied by clinical outcome metadata is abundant, computational methods that can effectively extract cell-type-specific expression from such data in each individual sample could be very helpful. If successful, such deconvolution methods could be used to identify clinically relevant cellular states of cell types and prioritize cell-cell interactions associated with patient survival and response to treatment. Furthermore, they may be readily applied to interrogate the troves of large bulk expression datasets that are in the public domain, to further advance what we can learn from them.

Recovery of cell-type-specific gene expression profiles from bulk gene expression is a computationally challenging problem because the standard formulation requires one to solve a large system of underdetermined linear equations. This problem is closely related to the problem of compressed sensing in signal processing[4]. Several recent studies have developed a variety of algorithms to address this challenge. DeMixT[5] was designed to estimate individual-specific expression for three cell components provided prior reference samples of two of these cell components. ISOpure[6] has aimed to derive sample-specific tumor cell expression only, with the assumption that the observed bulk gene expression profile is a mixture of predefined stromal and immune cell expression profiles that are shared across all the samples. Building on this work, Fox et al extended

ISOpure to predict individual-specific non-tumor cell expression by subtracting cancer cell
expression profiles from the bulk mixtures in a two-cell type model[7]. More recently,
Newman et al[8] developed CIBERSORTx, the first approach that aims to predict the
sample-specific gene expression of all cell types composing it by employing a set of novel
deconvolution heuristics. As a proof of concept, Newman et al[8] showed that
CIBERSORTx can accurately reconstruct the cell-type-specific expression of genes in
each input sample under certain modelling assumptions. This groundbreaking work has,
however, some notable limitations: (1) The number of genes whose cell-type-specific
expression can be reconstructed in each sample is quite small, especially for low-
abundance cell types, and (2) CIBERSORTx does not provide confidence estimates of
the predictions made. Confidence estimates on the expression predictions could be useful
in most deconvolution applications because of the absence of ground truth data.
79
Starting from the work of Newman et al[8], we introduce a new deconvolution
algorithm and software, CODEFACS (COnfident DEconvolution For All Cell Subsets) that
markedly advances our ability to successfully reconstruct cell-type-specific gene
expression of each bulk sample. CODEFACS receives as input bulk gene expression
profiles of tumor samples and either pre-computed estimates of abundance of expected
tumor, immunological and stromal cell types in each sample, or their prototypical
molecular signatures, which serve as seeds for estimating the abundance of each cell
type in each sample. CODEFACS then predicts the cell type-specific gene expression
profiles in each sample. It is a heuristic approach aimed at maximizing the number of
genes in each cell type whose expression across the samples can be estimated as
confidently predicted. Using 15 benchmark datasets where the ground-truth is known, we
show that CODEFACS robustly improves over CIBERSORTx, both in terms of gene
coverage and the individual gene expression estimation accuracy. We focus our
comparisons on CIBERSORTx because it is widely used (cited more than 350 times as
measured by Google Scholar since its publication in 2019) and to our knowledge, it is the
only widely used package that makes sample-specific estimates of gene expression for
each cell type.
97

We additionally introduce LIRICS (LIgand Receptor Interactions between Cell

Subsets), that integrates the output of CODEFACS with a database of prior

immunological knowledge that we curated to infer the cell-type-specific ligand-receptor

pairs that are likely to interact in a given sample. These data can then be analyzed in

conjunction with any sample-associated clinical annotations (e.g., response to treatment)

to systematically prioritize clinically relevant immune related interactions between any pair

of cell types in a given patient's cancer cohort. This opens up the possibility of identifying

phenotypic/clinical associations of individual interactions in large clinical tumor

expression cohorts.

Building on the enhanced coverage and accuracy of CODEFACS, we applied it to

reconstruct the cell-type-specific transcriptomes of ~8000 tumor samples from 21 cancer

types in the Cancer Genome Atlas (TCGA). Analyzing these fully deconvolved TCGA

expression datasets using LIRICS, we find a shared repertoire of cell-type-specific ligand-

receptor pairs that are likely to interact in the TME of mismatch repair deficient tumors of

different tissues of origin. These interactions are associated with improved overall patient

survival and high response rates to anti-PD1 treatment, independently of their mutation

burden levels. Finally, using machine learning techniques, we identify a subset of

intercellular TME interactions that are predictive of survival of melanoma patients

receiving immune checkpoint blockade therapy better than recently published

transcriptomics-based methods.

In summary, CODEFACS and LIRICS effectively build upon the statistical power

of large bulk RNA-seq datasets and prior immunological knowledge to characterize

sample specific cell-type-specific gene expression and learn more about the association

of different tumor-immune interactions with different clinical measures. Notably, the

potential scope of applications of both CODEFACS and LIRICS goes beyond studying

the TME, as these tools can be applied to study any disease of interest given bulk gene

expression data and relevant reference signatures of cell types involved.

Results

Overview of CODEFACS and LIRICS

CODEFACS is designed to characterize the tumor microenvironment by reconstructing

the cell-type-specific transcriptomes of each sample from bulk expression. It takes as

input the bulk RNA-seq expression values of a cohort of tumor samples and either the

estimations of the cell fractions of a pre-defined set of cell types in each sample or their

cell-type-specific molecular signature profiles, derived based on reference datasets or

from the literature[8].

CODEFACS then employs a heuristic approach that sequentially executes three

modules: *(module 1) high resolution deconvolution, (module 2) hierarchical deconvolution*

and (module 3) imputation. In module 1, we perform a high-resolution deconvolution,

which extends the CIBERSORTx algorithm. In module 2 (hierarchical deconvolution), bulk

expression is modeled as a mixture of two components: a specific cell type of interest and

all the remaining cell types. The expression for the cell type of interest is predicted by

removing the estimated expression in the second component (using high-resolution

deconvolution from module 1) from the bulk mixture. In module 3 – imputation-based

deconvolution, we impute the cell-type-specific expression of a specific gene based on

the predicted cell-type-specific expression of other high-confidence genes that are co-

expressed with that gene in the bulk. Each module is designed to overcome the

shortcomings of its predecessor based on their respective modeling assumptions. The

confidence ranking system is responsible for classifying all the predictions at the end of

each module into high-confidence or low-confidence predictions. Genes classified into

the low-confidence class at the end of one module (e.g. module 1) are passed to the next

module (e.g. module 2) for refinement. Finally, after all three modules are executed, the

prediction confidence levels are re-evaluated. The final output of CODEFACS consists of

a 3-dimensional matrix with cell-type-specific gene expression predictions for each

sample, along with a 2-dimensional matrix of estimated confidence scores of predictions

for each gene in each cell type. For more details, see Supplementary Note 1.

A key component of CODEFACS is its *confidence ranking system*, which receives
cell-type-specific expression predictions from the different modules and labels them as
high or low-confidence estimations. Genes whose expression is determined with high
confidence in a given module are added to the output set, while low confidence
predictions are continued to be processed in subsequent modules (See **Fig. 1A** and
Supplementary Note 1). The final output of CODEFACS consists of two items: **(a)** a
three-dimensional gene expression matrix, where each entry represents the predicted
gene expression a gene in a given cell-type in a specific sample, and **(b)** a two-
dimensional matrix of confidence scores ranging from [0,1] representing which gene-cell-
type pairs have the most confident predictions (**Fig. 1A, Output**).

Given fully deconvolved gene expression data from CODEFACS, one can use
LIRICS (LIgand Receptor Interactions between Cell Subsets) (**Fig. 1B**) to infer which cell-
type specific ligand-receptor pairs are likely to interact in each sample. Specifically,
LIRICS takes the output of CODEFACS and processes it in three steps: (*Step 1*) The first
step queries a database of all plausible ligand-receptor interactions between any two cell
types A and B, which we have systematically assembled and curated from the literature.
This database is publicly available as part of LIRICS (see **Supplementary Note 2**,
Supplementary Tables 1-3) and serves as a repository of all prior immunological
knowledge. (*Step 2*) In the second step, given the deconvolved expression profiles of cell
type A and cell type B in a given bulk tumor sample, LIRICS denotes as 'active' or 'likely
to occur' ('1') the interactions where both the ligand and receptor are over-expressed in
the relevant cell-types in that sample, or otherwise 'inactive' ('0'). A ligand or receptor is
considered to be overexpressed in a given cell type if its expression exceeds the median
expression in that cell type (**Supplementary Note 2**). (*Step 3*) In the third step, a Fisher's
enrichment analysis is performed to test the association of the activity of specific ligand-
receptor interactions with any relevant phenotypes of interes (e.g., treatment response,
mutational subtype, etc.) (**Fig. 1B**).

Figure 1. Overview of CODEFACS and LIRICS. (A) CODEFACS takes bulk gene expression

profiles and prior knowledge of the cellular composition of each sample and executes a heuristic

three-step procedure to infer the deconvolved gene expression in each sample, as described in

above and in more details in **Supplementary Note 1**. **(B) LIRICS** takes the output of CODEFACS

and processes it in three steps, as described in the main text and in detail in details, see

Supplementary Note 2.

Evaluating the performance of CODEFACS versus numerous different

benchmarks

To assess the accuracy of CODEFACS, we generated 15 benchmark datasets (see

Methods) by merging publicly available single cell RNA-seq[9,10] and FACS sorted purified

RNA-seq[11]. Thereafter, we applied CODEFACS to deconvolve these generated bulk

datasets and define the accuracy of its predictions by computing the Kendall correlation

between the predicted and ground truth expression in each cell type across individual
samples (the Kendall correlation provides a less inflated measure of accuracy by
accounting for ties in the data). In the main text, we show the results obtained on three of
these benchmark bulk datasets: one derived form a FACS-sorted lung cancer data, one
from a single cell melanoma RNA-seq data and from a single cell glioblastoma RNA-seq
dataset, respectively. Each bulk sample from these datasets represents a biopsy from a
patient. We show that CODEFACS predicts the cell-type-specific expression of many
more genes than CIBERSORTx (with Kendal's correlation ≥ 0.3) (**Fig. 2A-C**), and its
predictions are overall more accurate (**Fig. 2D-F**). The results for all the remaining 12
benchmark datasets, created via artificial mixing of single cell profiles and simulation of
batch effects between bulk and single cell expression data (See Methods and
Supplementary Table 4 for more details), further show the superiority of CODEFACS
(**Supplementary Fig. S1 and S2**). Overall, we observe that the more abundant the cell
type is, the better CODEFACS can predict its cell-type-specific gene expression
(**Supplementary Fig. S3**).

To test CODEFACS confidence scores can filter out potentially noisy predictions
we quantified how well do these scores align with the Kendall scores that measure the
prediction accuracy with the ground truth for each (gene, cell type) pair, as described
above. We quantified this using two metrics: Spearman correlation and a classification
Area Under the ROC Curve (AUC). As evident, this correlation is strong and positive (**Fig.**
2G depicts the results for the FACS sorted lung cancer benchmark dataset, and
Supplementary Fig. S4 depicts the results for the remaining benchmark datasets). To
perform a classification-based quantification, we grouped the genes in each cell type into
two classes based on the correlation between their predicted and actual expression, as
informative (prediction accuracy ≥ 0.1 and p-value ≤ 0.05) and *uninformative* (prediction
accuracy < 0.1 or p-value > 0.05). We then tested whether the confidence scores could
be used to classify genes into these two classes for each cell type. We found that the
confidence score could effectively filter out uninformative predictions (**Fig. 2H** depicts the
results for the FACS sorted lung cancer benchmark dataset, and **Supplementary Fig.**
S5, depicts for the remaining benchmark datasets).

Finally, to evaluate CODEFACS on real bulk tumor data, we applied it to

deconvolve bulk expression data from 21 cancer types (~8000 RNA-seq samples) in

TCGA. To infer the cellular abundance of each cell type in each sample which is required

as input for CODEFACS, we made use of matched bulk methylation data available for

these samples and methylation-based reference signature profiles of distinct cell types.

These include 11 cell type signatures (macrophages/dendritic cells--CD14+, B cells--

CD19+, CD4+T cells, CD8+ T cells, T regulatory cells, NK cells--CD56+, endothelial cells,

fibroblasts, neutrophils, basophils, eosinophils and tissue-specific tumor cells) obtained

from MethylCIBERSORT[12]. Reassuringly, we found high Spearman correlations between

the resulting predicted tumor cell fraction and the tumor purity estimates derived from

matched mutation and copy number data (based on ABSOLUTE) for the same samples

in each of 10 different cancer types (Spearman correlation: min=0.72, max=0.88, avg=0.8;

Supplementary Fig. S6).

We then asked if CODEFACS can recover the cell-type-specific gene expression

signature of different cell types in a given cancer type? To this end, we computed the

Spearman correlation between (a) the mean deconvolved gene expression of the top

confidently deconvolved genes in a given cell type (confidence score ≥ 0.95) and (b) the

mean expression of these genes, which we derived from completely independent single

cell expression data of the same cancer type (Methods). We find that (a) and (b) are

substantially correlated (**Fig. 2I** depicts results for the TCGA-LUAD (lung

adenocarcinoma) dataset as an example and **Supplementary Fig. S7** for the remaining

cancer types that have publicly available scRNA-seq data). The concordance level is

higher for cell types that are abundant (e.g., tumor cells and fibroblasts) and decreases

for less abundant cell types. Additionally, we observed that tumor cells have the largest

fraction of genes whose expression is predicted with high confidence, with the highest in

thyroid cancer (THCA, ~67.4% of all genes). Furthermore, seven KEGG pathways are

significantly enriched (adjusted p-value < 0.01) with highly confident genes in tumor cells

(confidence score ≥0.95) across the 21 cancer types (**Supplementary Fig. S8**). Those

pathways mostly involve RNA transport, spliceosome, DNA replication, and mismatch
repair.

266
267
268
269

Figure 2. Evaluating the performance of CODEFACS. (A-C) bar plots depicting the number of genes with a prediction accuracy (Kendall correlation) ≥ 0.3 with the ground truth for each cell type, estimated from bulk-generated samples of lung cancer (LUAD dataset; sample size = 26), melanoma (SKCM dataset 1; sample size = 28) and glioblastoma (GBM dataset 1; sample size = 24) benchmark datasets, as estimated by CODEFACS (yellow bars) and CIBERSORTx (blue bars). **(D-F)** boxplots depicting prediction accuracy distributions of all genes across different cell types in the lung cancer (LUAD with sample size 26), melanoma (SKCM with sample size 28) and glioblastoma (GBM with sample size 24) benchmark datasets, using CODEFACS (yellow) and CIBERSORTx (blue). A two-sided Wilcoxon signed rank test was performed to compare the prediction accuracies of CODEFACS and that of CIBERSORTx for each cell type in each dataset. *** denotes p-values < 2e-16. **(G)** Spearman correlations between prediction accuracies and confidence scores among cell types in the lung cancer benchmark dataset (LUAD dataset; sample size = 26). The y-axis indicates the spearman correlation coefficient value, while the x-axis indicates the cell type. **(H)** AUCs obtained in classifying informative and uninformative predictions among cell types in lung cancer benchmark dataset (LUAD dataset; sample size = 26). **(I)** bar plots depicting the Spearman correlations between mean deconvolved cell-type-specific expression in TCGA-LUAD and mean cell-type-specific expression derived from publicly available single cell datasets of LUAD. The y-axis indicates the Spearman correlation coefficient value, while the x-axis indicates the cell type. Details of all single cell RNA-Seq based pseudo bulk datasets can be found in **Supplementary Table 4.**

Tumors with DNA mismatch repair deficiency have heightened T cell co-stimulation
that is independent of their tumor mutation burden levels
In normal cells, DNA is constantly repaired in response to DNA damage or DNA
replication errors[13]. However, defects in specific DNA repair pathways in cancer cells may
result in the accumulation of many somatic mutations resulting in hypermutated tumors
(TMB ≥ 10-20 mutation/Mb)[14–16]. One cause of hypermutability is a mismatch repair
deficiency (MMRD), which leads to the accumulation of insertions and deletion mutations
in microsatellite regions of the genome due to uncorrected DNA replication polymerase
slippage events. This is known as microsatellite instability (MSI)[17,18]. Solid tumors with
mismatch repair deficiency were shown to be sensitive to immune checkpoint blockade

(ICB) therapy irrespective of tumor type, leading the FDA to approve MSI as the first
cancer type agnostic biomarker for patients receiving anti-PD1 treatment[19]. The reason
behind this general sensitivity to anti-PD1 treatment is not completely understood. Prior
work has led to the prevailing hypothesis that elevated tumor mutation burden in
mismatch repair deficient tumors leads to more neoantigens, and thus is more likely to
activate a host immune response against tumor cells[17,20–22]. However, not all tumor types
with elevated tumor mutation burden have similar response rates to anti-PD1[23,24], and
recent studies have revealed that T cells recognize and respond to only a few neoantigens
per tumor[25–28]. In the TCGA collection, there are three solid tumor types with a significant
association between hypermutability and survival benefit in patients. These three are the
MSI enriched solid tumor types (**Fig 3A,B,** borrowed from[29,30], MSI tumors highlighted as
red dots in Fig 3A, **Fig. 3C.** log-rank test p-value = 0.00084). The same cannot be said
about other solid tumor types (**Fig 3D**, log-rank test p-value = 0.4). Taken together, these
findings motivated us to further study cellular immune crosstalk in the tumor
microenvironment of mismatch repair deficient solid tumors to gain additional cell-type-
specific insights into their sensitivity to anti-PD1 treatment.

Figure 3. (A) The **landscape of tumor mutation burden and microsatellite instability across**

18 different solid tumor types. This panel plots the distribution of non-synonymous tumor

mutation burden on a logarithmic scale (Y-axis). All points above the horizontal line are typically

regarded as hyper-mutated tumors (> 10 mutations/Mb). All red points represent tumors with a

DNA mismatch repair deficiency detected via microsatellite instability (MSI). **(B) This panel**

depicts the percentage of all tumor samples per cancer type with microsatellite instability

(Y-axis). Tumor types marked with a * represent those where MSI is prevalent. **(C-D)**

Comparison of overall survival of patients with tumors that are hypermutated vs not

hypermutated. (**C**) Survival comparison in Gastro-Intestinal+Endometrial tumor types where MSI

is prevalent (marked with a * in panel B). (**D**) Survival comparison in all other solid tumor types

where tumors rarely have an underlying mismatch repair deficiency. Statistical significance of

differences in survival was calculated using the log-rank test.

To identify cell-cell interactions that are differentially active between microsatellite

instable tumors and microsatellite stable tumors (red vs black dots in Fig 3A), we applied

CODEFACS to deconvolve the bulk gene expression of all solid tumors from TCGA and

integrated their predicted cell-type-specific gene expression levels with LIRICS. The top

50 interactions from this differential analysis (ordered by FDR adjusted p-value) are

shown in a network in **Fig. 4A**. These interactions are frequently active in hypermutated

solid tumors with DNA mismatch repair deficiency compared to other hypermutated

tumors (**Fig. 4B, Supplementary Fig. S9**), testifying to their MSI specificity. They include

the well established PDL1-PD1 checkpoint interaction between tumor cells and CD8+ T

cells, but additionally, numerous T-cell activating/co-stimulatory interactions such as the

41BBL-41BB interaction between tumor cells and CD8+ T cells, ULBP2-NKG2D between

tumor cells and CD4+ T cells. They also encompass several chemotaxis related

interactions involved in trafficking of lymphocytes in and around the tumor mass, such as

the CXCL9-CXCR3 chemokine interaction between macrophages and CD4+ T cells and

CCL3/4/5 – CCR5 interactions between various immune and stromal cell-types.

Figure 4. (A) **Interaction network consisting of the top 50 interactions most highly activated**
in TME of tumors with DNA mismatch repair deficiency. Interactions highlighted in green
represent co-stimulatory interactions/having an activating effect on the target cell. Interactions
highlighted in red represent checkpoint interactions/having an inhibitory effect on the target cell.
Interactions highlighted in black represent pro-inflammatory/chemotaxis interactions involved in
inflammatory response and immune cell trafficking to tumor sites. Eos: Eosinophils, CAF: Cancer
associated fibroblasts. **(B) A volcano plot depicting on the x-axis the log2 fold change in the**
frequency of occurrence of each cell-cell interaction in the TME of hypermutated tumors
with an underlying DNA mismatch repair deficiency vs other hypermutated tumors. The y-
axis indicates the -log10 FDR adjusted p-value of the observed enrichment. Highlighted in red in
the scatter plot are the top 50 interactions that are most differentially active between all MSI vs
non-MSI tumors (shown in panel A).

Machine learning guided discovery of cell-type-specific ligand-receptor
interactions that predict patient response to ICB therapy

We set to discover cell-type-specific ligand-receptor interactions predictive of response
to ICB therapy by analyzing the TCGA deconvolved expression we have generated. We
focus on melanoma, currently the tumor type best responding to ICB, where there are
many independent publicly available bulk expression datasets of patient's receiving anti-

PD1 treatment. Starting from the deconvolved TCGA-SKCM dataset as our training set

(N=469), we employed a genetic algorithm to find cell-type specific ligand-receptor

interactions, whose activation state best separates hypermutated melanoma tumors from

non-hypermutated tumors (see Methods, **Fig 5A**). We term the interactions identified in

this process *melanoma hypermutation-specific functional interactions* (*MSFI*)*,* and the

network formed by these interactions is displayed in **Fig. 5B***.*

Having identified the MSFI interactions, we applied CODEFACS to deconvolve

the bulk expression data of pre-treatment samples from the three largest publicly

available melanoma datasets where patients received anti-PD1 treatment (either

monotherapy or in combination with anti-CTLA4; Methods)[43–45]. We then employed

LIRICS (step 1 and step 2) to the respective deconvolved expression of each of these

checkpoint datasets, *without any additional training*, and simply quantified the number of

MSFI interactions that are active in each of these patients' tumor samples, which we

denote as the tumor's *MSFI score*. Remarkably, we find that the MSFI score of each

sample can robustly stratify patients into those that are likely to respond to ICB vs those

that are unlikely to respond (**Fig. 5C**, progression free survival log rank test p-value:

0.00057, **Fig. 5D**, overall survival log rank test p-value: 0.0031). **Supplementary Fig.**

S10 depicts the survival differences for the two treatment groups separately (anti-PD1

monotherapy and anti-CTLA4 + anti-PD1 combination). Additionally, **Supplementary Fig.**

S11 and **S12** depict the survival differences for each ICB dataset separately. As evident,

our results improve over recent bulk gene expression based predictors of melanoma ICB

therapy response (IMPRES[46], TIDE[47] and the melanocytic plasticity signature (MPS)

scores[48]). We note that the performance levels of the latter on the more recent larger

bulk expression datasets, where their original RNAseq reads have been uniformly aligned

and normalized as described in the Methods, is lower than that reported in the original

publications, pointing to the growing awareness of the sensitivity of expression-based

predictors to batch effects, the normalization and alignment methods used, highlighting

the need to pre-process the raw sequencing data in a uniform manner in making

meaningful comparisons (see Discussion).

To further evaluate the predictive performance of the MSFI score, we tested its

ability to predict partial or complete responders vs stable or progressive disease patients

in these datasets. To this end, we plotted the receiver operator area under the curve (AUC)

obtained using the MSFI score for classifying the patients to partial or complete

responders vs stable or progressive disease, and compared its performance to that

obtained with the three other published predictors for the different treatment groups (**Fig.**

5E). On average, the MSFI score achieves an AUC of 0.63 (for anti-PD1 monotherapy

the AUCs obtained are 0.6, 0.77 and 0.52 for the three individual ICB datasets, and for

the anti-CTLA4 + anti-PD1 combination ICB treatment, the AUCs are 0.77, 0.49 and 0.63).

A similar performance could not be achieved if the placement of the ligand and receptor

between interacting cell-types in the MSFI network was swapped (average AUC 0.58) or

by randomly shuffling the interaction activity profiles (average AUC ~ 0.52), testifying that

the selected cell-cell interactions are best predictive of response to ICB. Comparing MSFI

predictive performance in this response classification task to that of the recent melanoma

bulk expression-based predictors, TIDE achieves an average AUC of 0.6 (for anti-PD1

monotherapy the AUCs are 0.46, 0.66 and 0.45 for the three ICB datasets and 0.89, 0.55

and 0.61 for the combination), IMPRES achieves an average AUC of 0.55 (for anti-PD1

monotherapy, the AUCs are 0.59, 0.64 and 0.6 respectively for the three ICB datasets

and 0.61, 0.44 and 0.42 for the combination) and MPS achieves an average AUC of 0.51

(for anti-PD1 monotherapy, the AUCs are 0.61, 0.49 and 0.63 respectively for the three

ICB datasets and 0.41, 0.59 and 0.38 for the combination).

Examining the MSFI network **(Fig. 5B)**, we find an over-representation of cell-

type-specific co-stimulatory/immune cell activating interactions known from prior

immunological literature (hypergeometric test p-value < 0.05)[49–59] (see **Supplementary**

Table 1), again highlighting the importance of co-stimulation in mediating successful anti-

tumor immune responses. Additionally, the MSFI network additionally includes

cytokine/chemokine interactions involved in pro-inflammatory response and the trafficking

of NK, T and B cells to the tumor site (responsible for better lymphocyte infiltration into

the tumor mass). Importantly, a multivariate Cox-proportional hazards model with MSFI

scores and TMB of each patient receiving anti-PD1 (wherever TMB data were available)

shows that the MSFI score remains significantly associated with improved progression free survival (p-value: 0.013) and overall survival (p-value: 0.0258) despite differences in mutation burden (TMB PFS p-value: 0.224, OS p-value: 0.477).

Taken together, these results suggest that while recognition of melanoma tumor neo-antigens serves as an initiating trigger, the activation of additional, specific, co-stimulatory and pro-inflammatory signals in the TME could further enhance an effective host immune response upon immune checkpoint blockade treatments.

Figure 5. (A) Overview of the machine learning analysis employed to identify cell type specific interactions that are predictive of response to immune checkpoint blockade therapy. (B) A chord diagram of the resulting MSFI network. Each individual interaction is represented by a link from the source cell type (ligand expressing cell type) to the target cell type (receptor expressing cell type) and the color of the link represents the color of the source cell type. For interactions that are activating/co-stimulatory, the sector in the corresponding target cell type is highlighted in green. For inhibitory/checkpoint interactions, the sector in the target cell type is highlighted in red. Interactions involved in chemotaxis are highlighted in black and those mediating a pro-inflammatory response are highlighted in blue, cell-adhesion interactions are highlighted in grey. **(C) Kaplan-Meier plot depicting the progression free survival of the combined set of melanoma patients receiving immune checkpoint blockade (N= 244).** On the top, the patients are stratified into low-risk/high-risk groups based on the median value of MSFI score from LIRICS. Second from top, patients stratified into low/high risk groups based on median IMPRES score[46]. Third from top, patients stratified into low/high risk groups based on median TIDE score[47], Bottom, patients stratified into low/high risk groups based on median MPS score[48]. **(D) Kaplan-Meier plots depicting the overall survival of all melanoma patients receiving immune checkpoint blockade (N= 244).** On the top, the patients are stratified into low-risk/high-risk groups based on the median value of MSFI score from LIRICS. Second from top, patients stratified into low/high risk groups based on median IMPRES score[46]. Third from top, patients stratified into low/high risk groups based on median TIDE score[47], Bottom, patients stratified into low/high risk groups based on median MPS score[48]. Survival differences among patients that received anti-PD1 monotherapy vs anti-CTLA4 + anti-PD1 combination are shown in **Supplementary Fig. S10 (E) Area under the ROC curves in predicting Complete/Partial-response (based on RECIST v1.1) to immune checkpoint blockade therapy for the different scores.** X-axis marks patients grouped by dataset source and treatment regimen. PD1 mono represents patients that received anti-PD1 monotherapy. PD1 + CTLA4 represents patients that additionally received anti-CTLA4 besides anti-PD1.

Discussion

This study presents a new computational tool, CODEFACS and a supporting immune interactions framework, LIRICS, that enable an (averaged) 'virtual single cell' characterization of the TME from bulk tumor expression data. Applying these tools across

the TCGA, we systematically identify, for the first time, cell-type-specific ligand-receptor

pairs that are likely to interact directly in the TME of specific patient populations and

prioritize those that are associated with patients' response to ICB. Specifically, we

identified a shared core of intercellular TME interactions in DNA mismatch repair deficient

tumors, which are associated with improved patient survival and high sensitivity to

immune checkpoint blockade therapy. Finally, focusing on melanoma, we show that one

can bootstrap on the large deconvolved data resource from TCGA using machine learning

techniques to discover cell-cell interactions within the TME that successfully predict

patients' response to immune checkpoint blockade.

The heightened cellular crosstalk unique to the TME of mismatch repair deficient

tumors suggests that T cells are being activated by co-stimulatory signals in addition to

immunogenic neo-antigens only to be kept in balance by other immunoregulatory

mechanisms such as the PD1-PDL1 checkpoint interaction between CD8+ T cells and

tumor cells. Our results suggest that when this interaction is blocked by anti-PD1

treatment, additional co-stimulatory interactions (such as the 41BBL-41BB between

tumor and T cells) lead to the observed enhanced response of MMRD tumors to immune

checkpoint blockade therapy. These results further support the possibility that switching

on specific T cell co-stimulation signals in the TME may lead to better responses to anti-

PD1 treatment independent of tumor mutation burden. Notably, recent pre-clinical studies

have shown that combination therapies aimed at enhancing such T-cell co-stimulating

interactions improve anti-tumor immune responses even in low TMB and highly immuno-

suppressive settings[33–38]. Currently, several clinical trials to assess the safety and efficacy

of these combinations are in progress[39–42].

While we have provided a toolkit to prioritize clinically relevant cell-cell interactions

from bulk tumor expression, there are limitations that should be noted and potentially

further improved upon in the future. First, several recent publications have reported

discrepancies between different RNA-seq expression quantification methods based on

the reference transcriptome version used and choice of method (alignment based vs

alignment free)[60–64]. This can potentially affect reproducibility of findings. Hence,

whenever possible, we recommend that all bulk RNA-seq datasets are homogeneously
pre-processed using the same RNA-seq quantification method and reference
transcriptome before the application of our tools. Indeed, following this notion, in this study
we have pre-processed all bulk RNA-seq datasets in a uniform manner, using STAR
(v2.7.6a) + RSEM (v1.3.3) and GENCODE v23 human genome annotation (Methods).
490

Second, CODEFACS itself has several limitations: (1) it requires prior information
about the cell type composition of the input tumors, or alternatively, knowledge of the
pertaining cell-types' gene expression or methylation signatures that can be used to infer
their abundances, and its accuracy depends on the accuracy of the latter. (2) It is
predictive typically for subsets of the whole exome and its performance deteriorates for
lowly-abundant cell types. However, the confidence scores provided help by allowing the
user to rank genes in each cell-type by the quality of predictions. Third, LIRICS is currently
restricted to well-defined ligand-receptor interactions between tumor, immune, stromal
and epithelial cell types and does not consider the spatial localization of cells in the TME.
The inclusion of the latter with the advent of forthcoming spatial transcriptomics data is
likely to lead to considerably more informative interaction inference approaches.
502

Although this work focuses on studying the tumor microenvironment, the tools
presented here can be applied to prioritize important cell-cell interactions in noncancerous
tissues under a variety of normal and disease states. One interesting application that we
envision is the characterization of clinically relevant intercellular interactions occurring at
the maternal-fetal interface using corresponding bulk gene expression data and
pregnancy outcome information, whose elucidation may help treat and mitigate
preeclampsia and other pregnancy related complications. One can also use our tools to
study bulk gene expression data from pre-malignant tissue samples and compare them
against malignant samples to elucidate cell-cell interaction dynamics on the way to
malignancy. Finally, one can deconvolve expression data from autoimmune disorders to
learn more about the underlying immune interactions. In sum, the computational tools
developed and presented here offer a cost-effective way to study immune responses at
a cell-type-specific resolution from the widely abundant bulk gene expression in both

health and disease, complementing first-line single cell technologies in situations where
the latter are less applicable.
518
519   **Data availability**
520   All deconvolved expression data generated in this study will be made publicly available
521   upon publication.
522
523   **Code availability**
524   The tools and all codes for reproducing the figures of this study will be made publicly
525   available upon publication.
526
Methods
Data collection and pre-processing
Single cell RNA-seq datasets: To benchmark the performance of CODEFACS, we first
set out to obtain publicly available single cell RNA-seq datasets where both tumor and
non-tumor cells were successfully isolated. This search led us to the identification of nine
such single cell RNA-seq datasets from the literature, each from a different cancer type.
Collection of additional single cell datasets was frozen after Dec 2019. Details of the
single cell datasets that we curated for this study in **Supplementary Table 5**. For each
dataset sequenced on the SmartSeq2 platform, the log normalized transcript counts for
each gene in each sequenced cell were made publicly available by the original authors.
For the application of deconvolution, these counts were transformed back to the
Transcripts Per Million (TPM) scale. For datasets sequenced on the 10x platform, UMI
counts for each gene were made publicly available and were scaled by the library size of
each cell and multiplied by a factor of 1 million to get expression values in TPM scale.
541
Bulk RNA-seq datasets: Gene expression and matching bulk tumor methylation data
from fresh frozen tumor biopsies in TCGA were downloaded from
(http://xena.ucsc.edu/)[65]. In addition, publicly available bulk expression data from formalin
fixed paraffin embedded tumor biopsies of melanoma patients receiving immune
checkpoint blockade treatment were downloaded from[43–45]. All bulk RNA-seq datasets

were collected such that they have a sufficiently large sample size to reliably perform

complete deconvolution of expression profiles (> 4 times the number of cell-types involved)

[8]. Collection of datasets was frozen after Dec 2019. Details on bulk RNA-seq datasets

deconvolved in this study can be found in **Supplementary Table 6**. To maintain

consistency with the pipeline used for preprocessing TCGA data, bulk gene expression

levels in immune checkpoint blockade datasets were re-quantified using STAR v2.7.6a

and RSEM v1.3.3[66] with GENCODE v23 human genome annotation[67]. Furthermore, to

mitigate technical biases, between-sample scaling factors were estimated using the TMM

method implemented in edgeR[68] and TPM values in each sample were further rescaled

by these scaling factors[69].

Generation of simulated bulk RNA-seq datasets: To evaluate the performance of

CODEFACS, we generated 14 different pseudo-bulk RNA-seq datasets from mixing

experiments with single-cell data. Each sample in each benchmark dataset has matching

cell type specific gene expression profiles derived from averaging single-cell RNA-seq

profiles of individual cells from the same sample and same cell type. These profiles serve

as the ground truth for the evaluation of deconvolution performance. To avoid any

circularity in our validations, for each of the single-cell datasets involved, single cell data

from 4 randomly chosen patients were separated from the rest. These data were used to

derive reference gene expression signatures for each cell type. The mixing experiments

were then performed on single-cell data of the remaining patients that were hidden from

the reference signature derivation process. In addition, we simulated technical replicates

for each pseudo-bulk sample, wherein we injected noise in the pseudo-bulk expression

of a few randomly chosen genes and then renormalized the expression data by the

sample library size. This procedure simulates mRNA composition noise that is commonly

observed in bulk RNA-seq datasets due to technical differences in sample preparation

[64-66]. The datasets and the mixing experiment used to generate the pseudo-bulk

samples are listed in **Supplementary Table 4**. In addition, we obtained a FACS sorted

lung cancer dataset which include purified RNA-seq for four cell types[11] and generated a

pseudo bulk correspondingly. In total, 15 benchmark datasets were generated.

Deconvolving immune checkpoint blockade (ICB) and TCGA bulk RNA-seq
datasets using CODEFACS

For the application of CODEFACS, molecular profiles of signature genes of each cell type of interest are needed to estimate the relative cell fractions in the bulk. We used single-cell-expression-derived signatures as priors to deconvolve the melanoma ICB datasets. To derive these signatures from single-cell data, we first start out by obtaining the class labels of each cell type of interest. These data are publicly available for each single-cell dataset we collected. Hence, we primarily use these labels in our study (unless further refinement of labels into specific cell subtypes of interest is needed for a specific usage). With a collection of single cell expression profiles and matching cell type labels as input, we used CIBERSORT online tool to derive a cell-type-specific signature matrix. Thereafter, we applied CODEFACS to ICB datasets with default parameters settings and batch correction requirement specified. For TCGA deconvolution, we first estimated cell fractions based on bulk methylation and then applied CODEFACS to corresponding bulk gene expression for the 21 cancer types which have both types of data available. We chose methylation signatures over expression-based signatures for TCGA analysis for two reasons. First, single cell expression data with consistent cell types across 21 cancer types are not available. Second, DNA methylation-based signatures are considered to be more stable marks of cellular identity compared to dynamic RNA expression derived signatures[70]. The methylation-based cell type signatures (**Supplementary Table 7**) were obtained from MethylCIBERSORT[12]. We applied CODEFACS to TCGA datasets with default parameters settings and without batch correction option specified. For the details of CODEFACS algorithm and parameter settings, see **Supplementary Note 1**.

Machine learning for discovery of cell-cell interactions predictive of clinical
responses to immune checkpoint blockade in melanoma

We used a genetic algorithm, which is a randomized heuristic search algorithm designed to select optimal features for a prediction task given some user-defined fitness function for training[71]. In this setting, the features are ligand-receptor interactions between cell-types, the prediction task is predicting response to ICB treatment and the fitness function is defined as the accuracy of predicting a user defined phenotype (used as pseudo-labels)

based on the total number of interactions from those selected occurring in a given sample;

accuracy is quantified by the AUC. To minimize the risk of overfitting to the TCGA dataset

and aid in faster convergence of the genetic algorithm during training, the size of the

feature space is reduced by filtering out complex and harder to measure interactions

involving receptors/ligands encoded by multiple genes and additionally assessing the fold

change of each interaction between the two classes specified in the training dataset and

incorporated into the fitness function (e.g., hypermutated vs non-hypermutated). After this

step, only interactions with a fold change >1 is passed to the genetic algorithm for further

optimization.

The algorithm starts out by randomly generating sets of features. This is defined

as the seed population. These sets iteratively evolve via the phenomenon of natural

selection enforced by the user defined fitness function. Specifically, for each subsequent

iteration, features from the best performing sets, as determined by the user defined fitness

function, in the current iteration are mixed at random followed by random new feature

additions or dropouts (referred to as mutations) to build a new generation of feature sets

and the process repeats. Eventually, after a number of epochs, which we set to 100, the

fitness function converges to an optimum and the best set of features for the prediction

task is returned to the user. Since the fitness function landscape is often non-convex and

the training process is stochastic, we repeat the training process 500 times, each with a

randomly chosen seed population, and eventually choose frequently selected features

over all solutions to reach a solution we suspect is close to the global optimum solution

and less likely to overfit. For our plots, we set the threshold to frequency > 100 times. The

probability of any feature being selected more than 100 times by random chance based

on this approach is estimated to be < 0.01. Results are qualitatively similar for more

stringent thresholds. The genetic algorithm was implemented in R using the genalg

package[72]

Acknowledgements

utilized the computational resources of the NIH HPC Biowulf cluster. We would additionally like to acknowledge Dr. Noam Auslander, Dr. Chi-Ping Day and other members of the Cancer Data Science Laboratory for their helpful feedback on this work.

Author contributions

K.W., S.P. A.A.S. and E.R. conceptualized and designed the project. A.A.S. and E.R. supervised the project. K.W. and S.P. implemented the software and performed the data analyses. K.W., S.P., A.A.S. and E.R. wrote the original draft with input from all authors. J.L., E.M.G., W.R., F.S., and D.R.C. contributed to data analyses, visualization, review and editing.

Competing interests

The authors declare no competing interests.

References:

1. Paget S. the Distribution of Secondary Growths in Cancer of the Breast. *Lancet*. 1889;133(3421):571-573. doi:10.1016/S0140-6736(00)49915-0

2. Smyth MJ, Ngiow SF, Ribas A, Teng MWL. Combination cancer immunotherapies tailored to the tumour microenvironment. *Nat Rev Clin Oncol*. 2016;13(3):143-158. doi:10.1038/nrclinonc.2015.209

3. Wagner A, Regev A, Yosef N. Revealing the vectors of cellular identity with single-cell genomics. *Nat Biotechnol*. 2016;34(11):1145-1160. doi:10.1038/nbt.3711

4. The fundamentals of compressed sensing. *Adv Comput Vis Pattern Recognit*. 2015;59:21-53. doi:10.1007/978-1-4471-6714-3_2

5. Ahn J, Yuan Y, Parmigiani G, et al. DeMix: Deconvolution for mixed cancer transcriptomes using raw measured data. *Bioinformatics*. 2013;29(15):1865-1871. doi:10.1093/bioinformatics/btt301

6. Quon G, Haider S, Deshwar AG, Cui A, Boutros PC, Morris Q. Computational purification of individual tumor gene expression profiles leads to significant improvements in prognostic prediction. *Genome Med*. 2013;5(3). doi:10.1186/gm433

7. Fox NS, Haider S, Harris AL, Boutros PC. Landscape of transcriptomic interactions between breast cancer and its microenvironment. *Nat Commun*. 2019;10(1):1-13. doi:10.1038/s41467-019-10929-z

8.   Newman AM, Steen CB, Liu CL, et al. Determining cell type abundance and expression
from bulk tissues with digital cytometry. *Nat Biotechnol*. 2019;37(7):773-782.
doi:10.1038/s41587-019-0114-2

9.   Jerby-Arnon L, Shah P, Cuoco MS, et al. A Cancer Cell Program Promotes T Cell
Exclusion and Resistance to Checkpoint Blockade. *Cell*. 2018;175(4):984-997.e24.
doi:10.1016/j.cell.2018.09.006

10.   Ghoshdastider U, Naeini MM, Rohatgi N, et al. Data-driven inference of crosstalk in the
tumor microenvironment. *bioRxiv*. Published online January 1, 2019:835512.
doi:10.1101/835512

11.   Gentles AJ, Hui ABY, Feng W, et al. A human lung tumor microenvironment interactome
identifies clinically relevant cell-type cross-talk. *Genome Biol*. 2020;21(1):107.
doi:10.1186/s13059-020-02019-x

12.   Chakravarthy A, Furness A, Joshi K, et al. Pan-cancer deconvolution of tumour
composition using DNA methylation. *Nat Commun*. 2018;9(1):1-13. doi:10.1038/s41467-
018-05570-1

13.   Erskine W, Kusmenoglu I, Muehlbauer FJ, Summerfield RJ. Breeding for increased
biomass and persistent crop residues in cool-season food legumes. *Genetics*.
2000;148(4):191-197. doi:10.1007/978-94-011-4385-1_17

14.   Loeb LA. Mutator Phenotype May Be Required for Multistage Carcinogenesis. *Cancer*
Res. 1991;51(12):3075-3079.

15.   Alexandrov LB, Nik-Zainal S, Wedge DC, et al. Signatures of mutational processes in
human cancer. *Nature*. 2013;500(7463):415-421. doi:10.1038/nature12477

16.   Muzny DM, Bainbridge MN, Chang K, et al. Comprehensive molecular characterization of
human colon and rectal cancer. *Nature*. 2012;487(7407):330-337.
doi:10.1038/nature11252

17.   Funkhouser WK, Lubin IM, Monzon FA, et al. Relevance, Pathogenesis, and testing
algorithm for mismatch repair-defective colorectal carcinomas: A report of the association
for molecular pathology. *J Mol Diagnostics*. 2012;14(2):91-103.
doi:10.1016/j.jmoldx.2011.11.001

18.   Cortes-Ciriano I, Lee S, Park WY, Kim TM, Park PJ. A molecular portrait of microsatellite
instability across multiple cancers. *Nat Commun*. 2017;8. doi:10.1038/ncomms15180

19.   Boyiadzis MM, Kirkwood JM, Marshall JL, Pritchard CC, Azad NS, Gulley JL. Significance
and implications of FDA approval of pembrolizumab for biomarker-defined disease. *J*
Immunother Cancer. 2018;6(1). doi:10.1186/s40425-018-0342-x

20. Galon J, Costes A, Sanchez-Cabo F, et al. Type, density, and location of immune cells
within human colorectal tumors predict clinical outcome. *Science (80-)*.
2006;313(5795):1960-1964. doi:10.1126/science.1129139
21. Llosa NJ, Cruise M, Tam A, et al. The vigorous immune microenvironment of
microsatellite instable colon cancer is balanced by multiple counter-inhibitory
checkpoints. *Cancer Discov*. 2015;5(1):43-51. doi:10.1158/2159-8290.CD-14-0863
22. Le DT, Uram JN, Wang H, et al. PD-1 Blockade in Tumors with Mismatch-Repair
Deficiency. *N Engl J Med*. 2015;372(26):2509-2520. doi:10.1056/nejmoa1500596
23. Yarchoan M, Hopkins A, Jaffee EM. Tumor Mutational Burden and Response Rate to PD-
1 Inhibition. *N Engl J Med*. 2017;377(25):2500-2501. doi:10.1056/nejmc1713444
24. Gurjao C, Tsukrov D, Imakaev M, Luquette LJ, Mirny LA. Limited evidence of tumour
mutational burden as a biomarker of response to immunotherapy. *bioRxiv*. Published
online January 1, 2020:2020.09.03.260265. doi:10.1101/2020.09.03.260265
25. Robbins PF, Lu YC, El-Gamil M, et al. Mining exomic sequencing data to identify mutated
antigens recognized by adoptively transferred tumor-reactive T cells. *Nat Med*.
2013;19(6):747-752. doi:10.1038/nm.3161
26. Bassani-Sternberg M, Bräunlein E, Klar R, et al. Direct identification of clinically relevant
neoepitopes presented on native human melanoma tissue by mass spectrometry. *Nat
Commun*. 2016;7. doi:10.1038/ncomms13404
27. Roh W, Chen PL, Reuben A, et al. Integrated molecular analysis of tumor biopsies on
sequential CTLA-4 and PD-1 blockade reveals markers of response and resistance. *Sci
Transl Med*. 2017;9(379). doi:10.1126/scitranslmed.aah3560
28. Kalaora S, Wolf Y, Feferman T, et al. Combined analysis of antigen presentation and T-
cell recognition reveals restricted immune responses in melanoma. *Cancer Discov*.
2018;8(11):1366-1375. doi:10.1158/2159-8290.CD-17-1418
29. Taylor AM, Shih J, Ha G, et al. Genomic and Functional Approaches to Understanding
Cancer Aneuploidy. *Cancer Cell*. 2018;33(4):676-689.e3. doi:10.1016/j.ccell.2018.03.007
30. Bonneville R, Krook MA, Kautto EA, et al. Landscape of Microsatellite Instability Across
39 Cancer Types. *JCO Precis Oncol*. 2017;(1):1-15. doi:10.1200/po.17.00073
31. Davoli T, Uno H, Wooten EC, Elledge SJ. Tumor aneuploidy correlates with markers of
immune evasion and with reduced response to immunotherapy. *Science (80-)*.
2017;355(6322). doi:10.1126/science.aaf8399
32. Auslander N, Wolf YI, Koonin E V. Interplay between DNA damage repair and apoptosis
shapes cancer evolution through aneuploidy and microsatellite instability. *Nat Commun*.

2020;11(1). doi:10.1038/s41467-020-15094-2

33.   Chen S, Lee LF, Fisher TS, et al. Combination of 4-1BB agonist and PD-1 antagonist

promotes antitumor effector/memory CD8 T cells in a Poorly Immunogenic Tumor Model.

Cancer Immunol Res. 2015;3(2):149-160. doi:10.1158/2326-6066.CIR-14-0118

34.   Ma HS, Poudel B, Torres ER, et al. A CD40 Agonist and PD-1 Antagonist Antibody

Reprogram the Microenvironment of Nonimmunogenic Tumors to Allow T-cell–Mediated

Anticancer Activity. *Cancer Immunol Res*. 2019;7(3):428-442. doi:10.1158/2326-

6066.CIR-18-0061

35.   Barber A. Costimulation of Effector CD8+ T Cells: Which Receptor is Optimal for

Immunotherapy? *MOJ Immunol*. 2014;1(2). doi:10.15406/moji.2014.01.00011

36.   Melero I, Hirschhorn-Cymerman D, Morales-Kastresana A, Sanmamed MF, Wolchok JD.

Agonist antibodies to TNFR molecules that costimulate T and NK cells. *Clin Cancer Res*.

2013;19(5):1044-1053. doi:10.1158/1078-0432.CCR-12-2065

37.   Chester C, Sanmamed MF, Wang J, Melero I. Immunotherapy targeting 4-1BB:

Mechanistic rationale, clinical results, and future strategies. *Blood*. 2018;131(1):49-57.

doi:10.1182/blood-2017-06-741041

38.   Philipson BI, O'Connor RS, May MJ, June CH, Albelda SM, Milone MC. 4-1BB

costimulation promotes CAR T cell survival through noncanonical NF-κB signaling. *Sci*

Signal. 2020;13(625). doi:10.1126/scisignal.aay8248

39.   Siu LL, Steeghs N, Meniawy T, et al. Preliminary results of a phase I/IIa study of BMS-

986156 (glucocorticoid-induced tumor necrosis factor receptor–related gene [GITR]

agonist), alone and in combination with nivolumab in pts with advanced solid tumors. *J*

Clin Oncol. 2017;35(15_suppl):104-104. doi:10.1200/jco.2017.35.15_suppl.104

40.   Tolcher AW, Sznol M, Hu-Lieskovan S, et al. Phase Ib study of utomilumab (PF-

05082566), a 4-1BB/CD137 agonist, in combination with pembrolizumab (MK-3475) in

patients with advanced solid tumors. *Clin Cancer Res*. 2017;23(18):5349-5357.

doi:10.1158/1078-0432.CCR-17-1243

41.   Cohen EEW, Pishvaian MJ, Shepard DR, et al. A phase Ib study of utomilumab (PF-

05082566) in combination with mogamulizumab in patients with advanced solid tumors. *J*

Immunother Cancer. 2019;7(1). doi:10.1186/s40425-019-0815-6

42.   Choi Y, Shi Y, Haymaker CL, Naing A, Ciliberto G, Hajjar J. T-cell agonists in cancer

immunotherapy. *J Immunother Cancer*. 2020;8(2). doi:10.1136/jitc-2020-000966

43.   Riaz N, Havel JJ, Makarov V, et al. Tumor and Microenvironment Evolution during

Immunotherapy with Nivolumab. *Cell*. 2017;171(4):934-949.e15.

doi:10.1016/j.cell.2017.09.028

44. Gide TN, Quek C, Menzies AM, et al. Distinct Immune Cell Populations Define Response

to Anti-PD-1 Monotherapy and Anti-PD-1/Anti-CTLA-4 Combined Therapy. *Cancer Cell*.

2019;35(2):238-255.e6. doi:10.1016/j.ccell.2019.01.003

45. Liu D, Schilling B, Liu D, et al. Integrative molecular and clinical modeling of clinical

outcomes to PD1 blockade in patients with metastatic melanoma. *Nat Med*.

2019;25(12):1916-1927. doi:10.1038/s41591-019-0654-5

46. Auslander N, Zhang G, Lee JS, et al. Robust prediction of response to immune

checkpoint blockade therapy in metastatic melanoma. *Nat Med*. 2018;24(10):1545-1549.

doi:10.1038/s41591-018-0157-9

47. Jiang P, Gu S, Pan D, et al. Signatures of T cell dysfunction and exclusion predict cancer

immunotherapy response. *Nat Med*. 2018;24(10):1550-1558. doi:10.1038/s41591-018-

0136-1

48. Pérez-Guijarro E, Yang HH, Araya RE, et al. Multimodel preclinical platform predicts

clinical response of melanoma to immunotherapy. *Nat Med*. 2020;26(5):781-791.

doi:10.1038/s41591-020-0818-3

49. Billadeau DD, Leibson PJ. ITAMs versus ITIMs: striking a balance during cell regulation.

J Clin Invest. 2002;109(2):161-168. doi:10.1172/jci14843

50. Staub E, Rosenthal A, Hinzmann B. Systematic identification of immunoreceptor tyrosine-

based inhibitory motifs in the human proteome. *Cell Signal*. 2004;16(4):435-456.

doi:10.1016/j.cellsig.2003.08.013

51. Murphy K, Weaver C. *Janeway's Immunobiology. Ninth Edition*.; 2018.

doi:10.1086/696793

52. Chen L, Flies DB. Molecular mechanisms of T cell co-stimulation and co-inhibition. *Nat*

Rev Immunol. 2013;13(4):227-242. doi:10.1038/nri3405

53. Campbell KS, Purdy AK. Structure/function of human killer cell immunoglobulin-like

receptors: Lessons from polymorphisms, evolution, crystal structures and mutations.

Immunology. 2011;132(3):315-325. doi:10.1111/j.1365-2567.2010.03398.x

54. Pende D, Falco M, Vitale M, et al. Killer Ig-like receptors (KIRs): Their role in NK cell

modulation and developments leading to their clinical exploitation. *Front Immunol*.

2019;10(MAY). doi:10.3389/fimmu.2019.01179

55. Ward-Kavanagh LK, Lin WW, Šedý JR, Ware CF. The TNF Receptor Superfamily in Co-

stimulating and Co-inhibitory Responses. *Immunity*. 2016;44(5):1005-1019.

doi:10.1016/j.immuni.2016.04.019

56. Gonçalves CM, Henriques SN, Santos RF, Carmo AM. CD6, a rheostat-type signalosome
that tunes T cell activation. *Front Immunol*. 2018;9. doi:10.3389/fimmu.2018.02994

57. Steri M, Orru V, Idda ML, et al. Overexpression of the cytokine BAFF and autoimmunity
risk. *N Engl J Med*. 2017;376(17):1615-1626. doi:10.1056/NEJMoa1610528

58. Chen M, Lin X, Liu Y, et al. The function of BAFF on T helper cells in autoimmunity.
Cytokine Growth Factor Rev. 2014;25(3):301-305. doi:10.1016/j.cytogfr.2013.12.011

59. Varin MM, Le Pottier L, Youinou P, Saulep D, Mackay F, Pers JO. B-cell tolerance
breakdown in Sjögren's Syndrome: Focus on BAFF. *Autoimmun Rev*. 2010;9(9):604-608.
doi:10.1016/j.autrev.2010.05.006

60. Wu DC, Yao J, Ho KS, Lambowitz AM, Wilke CO. Limitations of alignment-free tools in
total RNA-seq quantification. *BMC Genomics*. 2018;19(1). doi:10.1186/s12864-018-4869-
5

61. Sahraeian SME, Mohiyuddin M, Sebra R, et al. Gaining comprehensive biological insight
into the transcriptome by performing a broad-spectrum RNA-seq analysis. *Nat Commun*.
2017;8(1). doi:10.1038/s41467-017-00050-4

62. Everaert C, Luypaert M, Maag JLV, et al. Benchmarking of RNA-sequencing analysis
workflows using whole-transcriptome RT-qPCR expression data. *Sci Rep*. 2017;7(1).
doi:10.1038/s41598-017-01617-3

63. Teng M, Love MI, Davis CA, et al. A benchmark for RNA-seq quantification pipelines.
Genome Biol. 2016;17(1). doi:10.1186/s13059-016-0940-1

64. Robert C, Watson M. Errors in RNA-Seq quantification affect genes of relevance to
human disease. *Genome Biol*. 2015;16(1). doi:10.1186/s13059-015-0734-x

65. Goldman MJ, Craft B, Hastie M, et al. Visualizing and interpreting cancer genomics data
via the Xena platform. *Nat Biotechnol*. 2020;38(6):675-678. doi:10.1038/s41587-020-
0546-8

66. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq
quantification. *Nat Biotechnol*. 2016;34(5):525-527. doi:10.1038/nbt.3519

67. Harrow J, Frankish A, Gonzalez JM, et al. GENCODE: The reference human genome
annotation for the ENCODE project. *Genome Res*. 2012;22(9):1760-1774.
doi:10.1101/gr.135350.111

68. Robinson MD, McCarthy DJ, Smyth GK. edgeR: A Bioconductor package for differential
expression analysis of digital gene expression data. *Bioinformatics*. 2009;26(1):139-140.
doi:10.1093/bioinformatics/btp616

69. Robinson MD, Oshlack A. A scaling normalization method for differential expression

analysis of RNA-seq data. *Genome Biol*. 2010;11(3):R25. doi:10.1186/gb-2010-11-3-r25

70.   Bird A. DNA methylation patterns and epigenetic memory. *Genes Dev*. 2002;16(1):6-21.

doi:10.1101/gad.947102

71.   Melanie M. An introduction to genetic algorithms By Melanie Mitchell. MIT Press,

Cambridge, MA. (1996). 205 pages. $30.00. *Comput Math with Appl*. 1996;32(6):133.

doi:10.1016/S0898-1221(96)90227-8

72.   Willighagen E, Ballings M, Ballings MM. Package 'genalg.' Published online 2015.