Crosstalk between AML and stromal cells triggers acetate secretion through the metabolic rewiring of stromal cells

Nuria Vilaplana-Lopera[1], Ruba Almaghrabi[1*], Grigorios Papatzikas[1*], Vincent Cuminetti[2*], Mark Jeeves[1], Elena González[1], Alan Cunningham[3], Ayşegül Erdem[3], Frank Schnuetgen[4,5,6], Manoj Raghavan[1,7], Sandeep Potluri[1,7], Jean-Baptiste Cazier[1], Jan Jacob Schuringa[3], Michelle AC Reed[1], Lorena Arranz[2], Ulrich L Günther[1,8†] and Paloma Garcia[1,9†],

[1] Institute of Cancer and Genomic Sciences, University of Birmingham, Birmingham B15 2TT, United Kingdom
[2] Stem Cells, Ageing and Cancer Research Group, Department of Medical Biology, Faculty of Health Sciences, UiT – The Arctic University of Norway, 9019 Tromsø, Norway
[3] Department of Experimental Hematology, University Medical Center Groningen, University of Groningen, Hanzeplein 1, 9700 RB, Groningen, The Netherlands.
[4] Department of Medicine, Hematology/Oncology, University Hospital Frankfurt, Goethe-University, 60590 Frankfurt, Germany.
[5] Frankfurt Cancer Institute, Goethe-University Frankfurt, 60596 Frankfurt, Germany.
[6] German Cancer Consortium (DKTK), partner site Frankfurt/Mainz, and German Cancer Research Center (DKFZ), Heidelberg, Germany
[7] Centre for Clinical Haematology, University Hospitals Birmingham NHS Foundation Trust, Queen Elizabeth Hospital, Queen Elizabeth Medical Centre, Birmingham, United Kingdom.
[8] Institute of Chemistry and Metabolomics, University of Lübeck, 23562 Lübeck, Germany
[9] Corresponding author
* These authors contributed equally
† These two authors are co-senior authors.

Acute myeloid leukaemia (AML) cells interact and modulate components of their surrounding microenvironment into their own benefit. Stromal cells have been shown to support AML survival and progression through various mechanisms. Nonetheless, it is unclear whether AML cells could establish beneficial metabolic interactions with stromal cells. Here, we identify a novel metabolic crosstalk between AML and stromal cells where AML cells prompt stromal cells to secrete acetate for their own consumption to feed the tricarboxylic acid cycle (TCA). By performing transcriptome analysis, tracer-based metabolic NMR analysis and ROS measurements, we observe that stromal cells present a higher rate of glycolysis. We also find that acetate in stromal cells is derived from pyruvate via chemical conversion under the influence of reactive oxygen species (ROS) following ROS transfer from AML to stromal cells via gap junctions. Overall, we present a unique metabolic communication between AML and stromal cells that could potentially be exploited for adjuvant therapy.

Keywords: Leukaemic niche, acetate, AML, tracer-based NMR, metabolic crosstalk, ROS, acetate metabolism, gap junctions

Introduction

Acute myeloid leukaemia (AML) is a heterogeneous clonal disease characterised by a rapid proliferation of aberrant immature myeloid cells that accumulate in the bone marrow, and eventually in the blood and other organs, severely impairing normal haematopoiesis. AML cells show a highly adaptive metabolism that allows them to efficiently use a variety of nutrients to obtain energy and generate biomass (reviewed in [1]). This high metabolic plasticity confers AML cells a strong advantage against normal haematopoietic cells and has been related to AML aggressiveness. Although the metabolism in AML cells has been broadly investigated [1], fewer studies have focused on identifying metabolic alterations related to the interaction between AML and niche cells. For instance, AML cells are known to interact and modulate niche components for their own support by secreting soluble factors [2-6], via exosomes [7-9] or by establishing direct interactions, mediated by gap junctions [10] or tunnelling nanotubes [11]. These interactions with components of the niche provide AML cells with survival cues, chemoresistance and increased relapse in AML patients [2,4,7,10,12-14]. Furthermore, it has been reported that adipocytes in the niche secrete fatty acids, which are metabolised by AML cells through β-oxidation to obtain energy, protecting AML cells from apoptosis and ROS [15,16].

As a consequence of their highly proliferative demand, leukaemic stem cells (LSCs) [17-20] and, particularly, chemotherapy-resistant AML cells [21] present abnormally high levels of reactive oxygen species (ROS) [22,23]. How AML cells cope with high ROS levels has been intriguing and recent reports are shedding some light on whether the microenvironment plays a role in the redox metabolism of AML cells. For instance, it was reported that Nestin[+] bone marrow mesenchymal stem cells (BMSCs) support AML progression by increasing the bioenergetic capacity of AML cells and providing them with glutathione (GSH)-mediated antioxidant defence to balance the excess ROS [14]. Similarly, a recent study showed that co-culturing BMSCs with AML cells leads to a decrease in AML ROS levels due to an activation of the antioxidant enzyme GPx-3 in AML cells [24].

Our work provides new insight into the metabolic and redox crosstalk between AML and stromal cells, revealing a new metabolic interaction between AML and stromal cells. By combining transcriptomic and nuclear magnetic resonance (NMR) data, our results demonstrate that stromal cell metabolism is rewired in co-culture giving higher glycolysis and pyruvate decarboxylation, leading to acetate secretion. Our results also show that AML cells are able to transfer ROS to stromal cells by direct interaction through gap junctions and that these ROS can be used by stromal cells to generate and secrete acetate, which is utilised by AML cells as a biofuel by metabolising it via the

72   TCA cycle. Targeting ROS transfer via modulation of gap junctions to suppress the antioxidant

73   protection provided by stromal cells could serve as an adjuvant therapy to eradicate AML.

Results

Co-culturing AML and stromal cells in direct contact triggers acetate secretion by stromal cells

We first sought to determine whether interactions between AML and stromal cells in co-culture would result in differences in the consumption or production of extracellular metabolites. For this purpose, three AML cell lines (SKM-1, Kasumi-1 and HL-60) were cultured separately and in co-culture with MS-5, a stromal cell line capable of maintaining hematopoiesis [25], and the metabolic composition of the extracellular medium was analysed by ^{1}H-NMR and compared at different time points. The most striking difference found in co-culture compared to cells cultured separately was an increased secretion of acetate, which was common for the three cell lines used (Fig. 1a,b). Moreover, only stromal cells secreted acetate to a lower extent when cultured alone whereas AML cells did not secrete any acetate when cultured alone. Altogether, these findings suggest that acetate secretion is indeed a result of a direct interaction between AML and stromal cells. In addition, the observation that only stromal cells secrete acetate under these conditions suggests that stromal cells could be responsible for the increased acetate secretion found in co-culture.

Moreover, we examined the levels of other common extracellular metabolites, including glucose, lactate, glutamate and glutamine. As shown in Supplementary Fig. 1a higher consumption of glucose along with a higher secretion of lactate was observed in AML and stromal cells in co-culture compared to single cultures, suggesting a higher glycolytic flux in co-culture. However, the levels of glucose consumption and lactate production in co-culture were similar to the sum of the glucose consumption and lactate production levels of the AML and stromal cells in single cultures, suggesting that the overall increase in glycolysis in co-culture was just a result of culturing both cell types together. Additionally, we observed no variation in glutamate and glutamine levels suggesting that these metabolites are not involved in interactions that result from co-culture and are utilised depending on their availability.

The metabolic differences found in co-culture, could be due to altered proliferation under these conditions. To determine whether this was the case, a CFSE-based proliferation assay was performed in which AML cells and not stromal cells were stained and their growth was compared when cultured separately vs in co-culture. Over 48 hours, none of the AML cell lines tested presented differences in their proliferation rates (Supplementary Fig. 2a), thus confirming that the metabolic changes found in co-culture were caused by a mechanism independent from changes in proliferation.

Next, we aimed to determine whether cell-to-cell contact could play a role in the increased acetate secretion found in co-culture. For this we co-cultured AML and stromal cells separated by a permeable membrane, allowing cells to share the extracellular medium but impeding cell-to-cell

contact. Co-culturing cells using a permeable membrane blocked the increase in acetate secretion
observed under direct contact conditions (Fig. 1c). In fact, cells in co-culture presented lower levels
of acetate than MS-5 cells cultured alone revealing that direct cell-to-cell contact is required for
acetate secretion in co-culture.

To examine whether increased acetate secretion is specific for the interaction of AML cells with
stromal cells, we co-cultured AML cells with the cervical cancer cell line HeLa and compared the
levels of acetate of each cell type cultured alone. We found that there was no acetate secretion in
co-culture, also not by HeLa and AML cells cultured alone (Supplementary Fig. 2b), suggesting that
increased acetate secretion could be specific for the AML-stromal interaction.

Considering that our previous data seemed to indicate that MS-5 cells were responsible for acetate
secretion when co-cultured with AML cell lines, we decided to investigate how the levels of
extracellular acetate would vary after separating cells from co-culture. The three AML cell lines were
co-cultured for 24h with MS-5 cells prior to separation and were subsequently cultured in the same
spent media. Extracellular acetate levels in previously co-cultured MS-5 cells followed a similar trend
as before separation, suggesting that it is most likely that MS-5 cells are responsible for the
increased acetate secretion found in co-culture (Fig. 1d). Moreover, AML cells did not follow this
trend as they either maintained the levels of acetate seen prior to separating the cells (Kasumi-1 and
HL-60) or presented only a moderate increase (SKM-1) after being separated from co-culture.

We further investigated whether increased acetate secretion could take place in co-cultures, using
primary AML cells from patients, and whether acetate secretion in co-culture could be specific for
AML cells. To address this question, we isolated the CD34$^+$ population of four primary AML patient
samples, and three independent healthy donors, cultured them alone vs in co-culture with MS-5
cells, and analysed the composition of the extracellular medium at different time points. We found
that three out of four primary AML samples presented higher levels of acetate when co-cultured
with MS-5 cells compared to both MS-5 and AML cells cultured alone (Fig. 1e). Contrary, none of the
healthy donor samples showed an increased acetate secretion in co-culture suggesting that acetate
secretion is specific for AML cells in co-culture (Fig. 1f).

Moreover, we sought to determine whether in an experimental *in vivo* setting, increased acetate
production would be observed. For this, acetate levels were analysed in the bone marrow
extracellular fluid (BMEF) of mice transplanted either with MLL-AF9 leukaemic cells or with healthy
wild type hematopoietic cells. These experiments revealed higher amounts of acetate in the BMEF of
mice suffering from leukaemia compared to controls (Fig. 1g).

Fig 1. Acetate secretion by stromal cells increases in AML-stroma co-cultures of several

AML cell lines and primary AML cells in direct contact. a, Section of ^{1}H-NMR spectra,

corresponding to the methyl group of acetate, from extracellular medium samples of SKM-1 cells

cultured alone (blue), MS-5 cells cultured alone (red) and SKM-1 and MS-5 cells in co-culture (green)

after 24 hours. **b,** Extracellular acetate levels in AML cell lines (SKM-1, Kasumi-1 and HL-60) cultured

alone (blue), MS-5 cells cultured alone (red) and AML and MS-5 cells in co-culture in direct contact

(green) at 0, 20 and 24 hours of incubation. Each point represents the mean of n=3 independent

experiments and error bars represent standard deviations. **c,** Extracellular acetate levels in AML cell

lines cultured alone (blue), MS-5 cells cultured alone (red) and AML and MS-5 cells in co-culture

separated by a 0.4 µm permeable membrane (green) at 0, 20 and 24 hours of incubation. Each point

represents the mean of n=3 independent experiments and error bars represent standard deviations.

d, Extracellular acetate levels in AML cell lines and MS-5 cells in co-culture (black) for 24 hours and

after being separated and cultured alone in the same medium until 48 hours (blue for AML and red

for MS-5). Each point represents the mean of n=3 independent experiments and error bars

represent standard deviations.

For **b**, **c, d** and **g** unpaired Student's t-tests were applied for each condition (black brackets) and p-

values are represented by n.s. for not significant * for p-value<0.05, ** for p-value<0.01 and *** for

p-value<0.001.

e, Extracellular acetate levels in MS-5 cells cultured alone (black) and primary patient-derived AML

cells co-cultured with MS-5 cells (grey) at 48h. Each set of points represents an independent

experiment (n=4). **f,** Extracellular acetate levels in MS-5 cells cultured alone (black) and healthy

donor-derived peripheral blood mononuclear CD34[+] (PBMC) cells co-cultured with MS-5 cells (grey)

at 48h. Each set of points represents an independent experiment (n=3). **g,** Acetate levels in bone

marrow extracellular fluid of C57BL6/J mice 6 months after transplantation with bone marrow

nucleated cells isolated from control or MLL-AF9 transgenic mice. A Mann-Whitney test was applied

(black bracket) and p-value was represented as * for p-value<0.05 (p=0.0357).

AML cells consume and use acetate secreted by stromal cells to feed the TCA cycle

Following the finding that stromal cells are responsible for acetate secretion in co-culture, we decided to determine whether AML cells might be able to metabolise the secreted acetate. We first sought to define the concentration of secreted acetate in the extracellular medium in co-culture. For this, we compared a sample of extracellular medium from a co-culture of SKM-1 and MS-5 cells after 24 hours to a calibration curve (Supplementary Fig. 2c) which allowed us to determine that the concentration of acetate in co-culture, which was approximately of 3 mM. We then investigated whether SKM-1 cells can consume acetate both in physiological and co-culture concentrations. In both cases SKM-1 cells consumed acetate after 48 hours, and in physiological conditions also after 24 hours (Fig. 2a).

We used a tracer-based approach using [2-^{13}C]acetate to assess whether SKM-1 or MS-5 cells can utilise acetate (Fig. 2b,c and Supplementary Fig. 3a). Both cell types import ^{13}C labelled acetate as observed in NMR spectra (Fig. 2c). However, only SKM-1 cells showed ^{13}C label incorporation in several TCA cycle related metabolites, including aspartate, citrate, glutamate, 2-oxoglutarate, glutathione and proline, as well as in acetylcarnitine. Acetylcarnitine is known to be produced by cells when large amounts of acetyl-CoA are present in the mitochondria [26,27], which could be in line with this experiment in which a 4 mM concentration of [2-^{13}C]acetate was used. Overall, this data suggests that acetate in co-culture could be utilised by SKM-1 cells but not by MS-5 cells.

In order to determine whether AML cells can import and metabolise the secreted acetate in co-culture, we co-cultured AML and MS-5 cells for 24 hours, we then added [2-^{13}C]acetate to the spent extracellular medium and cultured cells for 8 hours before analysing the intracellular metabolites in each cell type (Fig. 2d). We found that only AML cells presented ^{13}C labelling in intracellular metabolites (Fig. 2e, Supplementary Fig. 3b). Moreover, the metabolisation pattern for all AML cells was similar to that observed for SKM-1 cells alone (Fig. 2c), where label was incorporated into TCA cycle-related metabolites. Overall, these results revealed that AML cells uptake and utilise the acetate secreted by stromal cells in co-culture as a substrate to feed into the TCA cycle.

Fig. 2. AML cells import the secreted acetate in co-culture to use it as a biofuel. a,

Extracellular acetate levels in SKM-1 cells cultured with 0.25 mM and 3 mM acetate medium after 0,

24 and 48 hours. Each point represents the mean of n=3 independent experiments and error bars

represent standard deviation. An unpaired Student's t-test was applied for each condition (black

brackets) and p-values are represented by * for p-value<0.05, ** for p-value<0.01 and *** for p-

value<0.001. **b,** Scheme of label distribution arising from [2-^{13}C]acetate in TCA cycle intermediates.

Black circles correspond to expected labelled positions. **c,** ^{13}C percentages of label incorporation in

polar metabolites from labelled acetate in SKM-1 and MS-5 cells after two hours of incubation with

[2-^{13}C]acetate (4 mM) derived from ^{1}H-^{13}C-HSQC NMR spectra. Bars represent the mean of the ^{13}C

percentage and error bars represent the standard deviations for n=3 independent experiments. **d,**

Experimental design for acetate labelling in co-culture. Briefly, AML cells are co-cultured with

stromal cells for 24 hours prior to addition of [2-^{13}C]acetate (extra 4 mM). After 8 hours of the

addition of acetate, cells are separated, and metabolites are extracted. **e,** ^{13}C percentages on polar

metabolites in SKM-1 and MS-5 cells in co-culture. Cells were co-cultured for 24 hours before the

addition of extra 4 mM sodium [2-^{13}C]acetate. Bars represent the mean of the ^{13}C percentage and

error bars represent the standard deviations of n=3 independent experiments. For C and E, ^{13}C

natural abundance is represented as a black bar at %^{13}C = 1.1.

Transcriptomic data highlights a metabolic rewiring of stromal cells in co-culture characterised by upregulation of glycolysis and downregulation of pyruvate dehydrogenase

After establishing that stromal cells are responsible for acetate secretion (Fig. 1d) and that AML cells consumed the secreted acetate in co-culture (Fig. 2d), we sought to elucidate the mechanism behind acetate secretion by MS-5 cells in co-culture. Thus, we set out to perform global gene expression profiling by RNA-seq comparing cells cultured alone and in co-culture (Supplementary Fig. 4a). With this approach we identified 587 differentially expressed genes (q-value<0.1) (Supplementary Fig. 4b); with 476 genes being upregulated and 111 genes being downregulated in co-culture.

Following clustering of differentially expressed genes, gene set enrichment analysis (GSEA) was performed (Fig. 3a) revealing a positive correlation with the expression of genes that are part of the glycolysis pathway (MSigDB: M5937) as well as the reactive oxygen species pathway (MsigDB: M5938) (Fig. 3b).

A closer examination of the genes involved in the glycolysis pathway revealed a major upregulation of several glycolysis-related genes in MS-5 cells in co-culture (Fig. 3c). Most of the genes involved in glucose transport (*Slc2a1* and *Slc2a4*) and glucose breakdown to pyruvate (*Pgm1*, *Hk2*, *Gpi1*, *Pfkl*, *Gapdh*, *Pgk1*, *Pgam1-2*, *Eno1-1b-2* and *Pkm*) were upregulated, including the gene encoding for the 6-phosphofructo-2-kinase/fructose-2,6-biphosphatase 3 (*Pfkfb3*), a well-known activator of glycolysis.

In contrast, individual examination of genes related to pyruvate metabolism revealed that the genes related to pyruvate conversion to acetyl-CoA were downregulated (Fig. 3c). Pyruvate transport into the mitochondria (*Mpc1-2*) and several pyruvate dehydrogenase (PDH) complex-related genes (*Pdha1*, *Pdhb* and *Pdhx*), involved in the conversion of pyruvate to acetyl-CoA, remained largely unaltered or were slightly downregulated by co-culture. Additionally, the pyruvate dehydrogenase kinases (*Pdk1*, *Pdk2* and *Pdk4*), which inhibit the activity of the PDH complex, were found to be upregulated by co-culture. Interestingly, the acetyl-CoA synthetases (ACSs), which can generate acetyl-CoA from acetate but have also been reported to perform the reverse reaction, encoded by *Acss2* and *Acss3*, were found to be slightly upregulated in co-culture. Pyruvate can also be converted to 2-oxaloacetate via pyruvate carboxylase (*Pcx*), to lactate by lactate dehydrogenase (*Ldha*), and to alanine via alanine transaminase (*Gpt*). Only *Ldha* was found to be moderately upregulated in co-culture. However, we did not observe a substantial increase in lactate production experimentally in co-culture. Altogether, transcriptomic data suggests that MS-5 cells in co-culture present a major upregulation of glycolysis and downregulation of the PDH complex.

To explore whether glycolysis is upregulated at a metabolic level in MS-5 cells in co-culture at the

metabolic level and whether acetate could derive from glucose, we performed [U-^{13}C]glucose tracing

on AML and MS-5 cells cultured alone and in co-culture and analysed the label incorporation in

glycolysis-related extracellular metabolites (Fig. 3d). For the three AML cell lines tested, extracellular

acetate presented significantly higher label incorporations from [U-^{13}C]glucose in co-culture

compared to cell types cultured alone, providing evidence that the secreted acetate in co-culture

derives from glucose. Lactate and alanine, which can be synthesised from pyruvate, did not show

significant increases in label incorporation from [U-^{13}C]glucose in co-culture compared to each cell

type cultured alone for all the AML cell lines, with the exception of alanine in HL-60. Overall, these

results are in agreement with the transcriptomic data (Fig. 3c), highlighting that glucose metabolism

is upregulated in co-culture but also confirming that acetate derives from glycolysis.

a **Top 25 hallmark gene sets**
(ranked by expression)

b Glycolysis

ROS

c Glycolysis

Pyruvate metabolism

d SKM-1

Kasumi-1

HL-60

Co-culture
MS-5 only
AML only

Fig. 3. Transcriptomic data reveals that stromal cell metabolism is shifted towards higher

glycolysis and ROS upon co-culture with AML cells. a, Top 25 GSEA hallmark gene sets ranked

by expression in MS-5 cells only vs co-culture with SKM-1 cells, analysed using the collection of

hallmark gene sets from Molecular Signature Database with a false discovery rate threshold at 5%. P-

values for each pathway are presented as –log(10)p-value. **b,** GSEA enrichment score plots of

glycolysis and ROS generated using Sleuth 0.30.0 R statistical package. **c,** Fold change values of

detected gene transcripts (TPMs) related to glycolysis and pyruvate metabolism. FC values are

represented as $\log_2$FC, red values indicate upregulation and blue values indicate downregulation in

MS-5 cells in co-culture. **d,** Label incorporation from [U-^{13}C]glucose into extracellular metabolites in

AML and MS-5 cells cultured alone or in co-culture after 24 hours. Bars represent the mean of n=3

independent experiments and error bars represent standard deviations.

AML cells rewire stromal cell metabolism transferring ROS via gap junctions to obtain acetate

Tracer-based data on MS-5 cells in co-culture revealed that acetate secreted in co-culture derives from glucose (Fig. 3d). However, transcriptomic data on MS-5 cells in co-culture suggested that acetate did not derive from pyruvate via acetyl-CoA (Fig. 3c). Moreover, acetate secretion was also observed in MS5 cells grown in thiamine-free media (Supplementary Fig. 5a), confirming that the acetate secretion was not dependent on keto acid dehydrogenases [28] and indicating that a different mechanism must be responsible. An alternative mechanism of acetate synthesis involving a non-enzymatic oxidative decarboxylation of pyruvate into acetate had previously been described [28-30]. This mechanism was reported to be mediated by ROS in mammalian cells and was linked to cells prone to overflow metabolism under the influence of high rates of glycolysis and excess pyruvate. Hence, we decided to investigate whether ROS might play a role in acetate secretion in our co-culture system.

We first modulated ROS levels in AML cells and MS-5 cells cultured alone and in co-culture and measured acetate production. Hydrogen peroxide was used to increase ROS levels and N-acetylcysteine (NAC) was used as a ROS scavenger. Extracellular acetate levels were measured by ^{1}H-NMR after 24 hours. Increasing ROS levels with peroxide resulted in a significant increase in acetate secretion, particularly in SKM-1 and Kasumi cells in co-culture (Fig. 4a, Supplementary Fig. 5b)). This experiment could not be carried out with HL-60 cells as peroxide treatment severely impaired the viability of HL-60 cells, as previously reported [31]. Additionally, when the ROS scavenger NAC was used, a significant decrease in the levels of acetate in both MS-5 cells cultured alone and in all the co-cultured cell lines was observed (Fig. 4a, Supplementary Fig. 5b). Interestingly, the production of acetate in cells in co-culture treated with NAC recovered the phenotype of MS-5 cells cultured alone, indicating that acetate synthesis in MS-5 cells in co-culture is mediated by ROS (Fig. 4a).

Next, we compared intracellular ROS levels in AML and stromal cells in co-culture and cultured alone by labelling cells with the H$_2$DCFDA fluorescent dye. Fluorescence measurements showed that ROS levels in the three AML cell lines used were significantly decreased in co-culture, whilst in the stromal cells ROS levels were significantly increased in two of the three co-cultures (Fig. 4b). These results suggested that AML cells might transfer ROS to stromal cells. We also performed the same experiment using three primary AML samples to corroborate the previous result. Fluorescence analysis showed decreased ROS levels in AML samples in co-culture and increased ROS levels in MS-5 cells in co-culture for the three primary AML samples analysed (Fig. 4c), suggesting ROS transfer from AML cells to stromal cells in co-culture.

To further test whether AML cells might transfer ROS through a contact-dependent mechanism, we compared the intracellular ROS levels in AML and stromal cells cultured alone and co-cultured without direct contact using a permeable membrane. Fluorescence measurements in both cell types revealed that ROS levels remained unaltered in contact-free co-cultures (Fig. 4d), indicating that ROS transfer could only occur via a contact-dependent mechanism.

As it had been previously reported that haematopoietic stem cells can transfer ROS to stromal cells via gap junctions to prevent senescence [32], we decided to examine gap junction genes in the transcriptome of MS-5 cultured alone and in co-culture. When individually examining the gap junction genes in MS-5 cells cultured alone vs co-culture, we found several gap junction genes upregulated in co-culture such as *Gja5, Gja8, Gjb5* and *Gjc2* (Fig. 4e). These results suggest that AML cells might establish gap junction interactions to transfer ROS to MS-5 cells when in co-culture. To test this hypothesis, we used the calcein-AM dye, which can only be transferred via gap junctions [10]. We labelled stromal cells with calcein-AM, cultured them with unlabelled AML cells and analysed the fluorescence of AML cells after three hours. We found that, for the three AML cell lines tested, the percentage of cells that had incorporated the calcein-AM dye from MS-5 cells was larger than 80% (Fig. 4f, Supplementary Fig. 5c), indicating that AML cells can establish gap junctions with stromal cells when co-cultured in direct contact.

Next, we decided to confirm that AML cells can transfer ROS via gap junctions by inhibiting the gap junctions using carbenoxolone (CBX), a well-known gap junction inhibitor [10,33,34]. We first confirmed that efficiency of inhibition by analysing the calcein-AM dye transfer in a control and treated co-culture of SKM-1 and MS-5 cells. The CBX treatment reduced the calcein-AM transfer more than 60% in co-culture (Fig. 4g). We then compared intracellular ROS levels in both AML and stromal cells treated with CBX and control. CBX treatment abrogated the decrease in ROS levels in AML cells (Fig. 4h) and the increase of ROS levels in stromal cells (Fig. 4i), indicating inhibition of ROS transfer in CBX treated co-cultures. Overall, this result provides strong evidence that ROS transfer occurs via gap junctions established between AML and stromal cells.

Fig. 4. Acetate secretion is linked to ROS transfer from AML to stromal cells via gap junctions. a, Extracellular acetate levels in SKM-1 (black) and MS-5 cells cultured alone (dark grey) and in co-culture (light grey) for 24 hours in a control medium, medium with 50 µM H_2O_2 or medium with 5 mM NAC. **b, c** and **d** Intracellular ROS levels measured by H_2DCFDA staining in **b** and **d** AML cells or **c** primary patient-derived AML cells and MS-5 cells cultured alone and in co-culture in **b** and **c** direct contact or **e,** fold change values of detected gene transcripts (TPMs) related to gap junctions. FC values are represented as log_2FC, red values indicate upregulation and blue values indicate downregulation in MS-5 cells in co-culture. **f,** Frequencies of Calcein-AM and CD33 positive AML cell lines after being in co-culture with Calcein-AM stained MS-5 cells for 3 hours. **g,** Frequencies of Calcein-AM and CD33 positive SKM-1 cells (treated or untreated with 200 µM carbenoxolone for 24 hours) after being in co-culture with Calcein-AM stained MS-5 cells for 3 hours. **h** and **i** Intracellular ROS levels measured by H_2DCFDA staining in AML cells (**h**) and MS-5 cells (**i**) cultured alone and in co-culture in direct contact treated or untreated with 200 µM carbenoxolone for 24 hours. For **a**, **b**, **c**, **d**, **f**, **g**, **h** and **i** bars represent the mean of n=3 independent experiments and error bars represent standard deviations. For **a**, **b**, **d**, **h** and **i** unpaired Student's t-test was applied for **a** each condition (H_2O_2/NAC) was compared to untreated; for **b**, **d**, **h** and **i** co-culture conditions were compared to cells cultured alone. P-values are represented by n.s. for not significant, * for p-value<0.05, ** for p-value<0.01 and *** for p-value<0.001.

Discussion

AML cells are known to interact and remodel niche cells through various mechanisms, including the secretion of soluble factors, cytokines or metabolites, resulting in a better support of AML cells at the expense of normal haematopoiesis. Yet, metabolic crosstalk between AML and stromal cells has not been reported before. Here, we have identified a novel metabolic communication between AML and stromal cells mediated by acetate. Our in vivo data using cells from a MLL-AF9 knock-in mouse model showed higher acetate levels in the bone marrow extracellular fluid in leukaemic compared to healthy control mice, indicative of a potential role for acetate in AML development. Our in vitro data suggests that AML cells can modulate stromal cells into secreting acetate in co-culture, by rewiring stromal cell metabolism, and can then utilise the secreted acetate to feed their own TCA cycle to generate energy. Co-cultures with primary cells suggest that this effect could be also occurring in at least a subset of primary AML cells.

Mechanistically, our data revealed that acetate secretion involves not only a higher glycolytic rate but also the non-enzymatic ROS-mediated conversion of pyruvate to acetate, as cells grown in thiamine-free media were still capable to produce acetate. Furthermore, our data indicates that AML cells can diminish their ROS levels by establishing gap junctions with stromal cells facilitating ROS transfer to stromal cells.

Studying the interactions between cancer cells and their microenvironment in terms of metabolism has become an exciting new field of cancer research. Our data indicates that AML cells can influence the metabolism of stromal cells causing increased acetate secretion, which was not observed in healthy counterparts. We presented several lines of evidence suggesting that stromal cells but not AML cells are responsible for acetate secretion: (i) stromal cells cultured alone secreted acetate, whilst AML cell lines and primary AML cells did not; (ii) after separating stromal cells from co-culture with AML cell lines, stromal cells continued to secrete acetate at a similar rate to that observed in co-culture; and, (iii) glycolysis was upregulated in stromal cells in co-culture, and glucose was found to be the precursor of the secreted acetate in co-culture.

Interestingly, acetate has been reported as an alternative fuel for cancer cells [35,36], especially under low oxygen conditions or lipid depletion [37-39], but it has not previously been described to participate in crosstalk between any type of cancer cells and their microenvironment. Nonetheless, other monocarboxylate metabolites, such as lactate and alanine, have been reported to participate in different types of metabolic interplay between stromal and cancer cells [40-42]. Our work reveals that aside from lactate and alanine, other monocarboxylate metabolites, such as acetate, can be utilised by leukaemic cells as a biofuel. Although acetate usage to feed the TCA cycle had already been

described in AML [43] and other types of cancer [38,44], this is, to our knowledge, the first report of AML

cells utilising acetate secreted by stromal cells in co-culture.

We have also proposed a mechanism for the altered stromal cellular metabolism, involving

increased glycolysis and the ROS-mediated chemical conversion of pyruvate to acetate.

Transcriptomics showed that glycolysis is upregulated in stromal cells in co-culture, and tracer-based

metabolism using [U-^{13}C]glucose, demonstrated that more glucose-derived label gets incorporated

in pyruvate and acetate in stromal cells in co-culture compared to either AML or stromal cells

cultured alone. Enrichment of hypoxia genes and elevated Pdk expression have been reported to be

related to higher glycolytic activity [45-47]. Similarly, other cancer cells [48-51] have also been shown to

modulate stromal metabolism by increasing glycolysis. Moreover, this mechanism was further

supported by modulating ROS levels in co-culture, as increasing or lowering ROS levels had a clear

influence on acetate secretion.

AML cells are known to exhibit high levels of ROS [52,53]. However, our data has shown that AML cells

in co-culture present lower levels of ROS than AML cells cultured alone, which is in agreement with

recent studies in primary AML and mesenchymal stromal cell co-cultures [14,24]. The current

mechanisms to describe this phenomenon involve mitochondrial transfer and activation of

glutathione-related antioxidant pathways [14,24], although previous data on haematopoietic stem cells

(HSCs) revealed that HSCs can directly transfer ROS via gap junctions to stromal cells [32]. Interestingly,

our data showed that the decrease in ROS levels was counteracted by inhibiting contact using a

permeable membrane. We also found several gap junction genes upregulated in stromal cell types in

co-culture. Moreover, we prevented ROS transfer from AML to stromal cells by using the gap

junction inhibitor CBX [10,33,34]. Thus, our results suggest that ROS transfer via gap junctions, at least

partially, mediates the mechanism behind AML cells presenting lower levels of ROS in co-culture.

Overall, this work reveals a unique and novel metabolic communication between AML and stromal

cells that involves acetate as the main crosstalk metabolite. We showed that AML cells are capable

of modulating the metabolism of stromal cells by transferring ROS via gap junctions resulting in an

increased secretion of acetate and its subsequent accumulation in the extracellular medium, which

correlates with the observation of higher acetate levels in the bone marrow extracellular fluid of

AML mice compared to control mice. Furthermore, we found that AML cells consume the secreted

acetate and use it as an energy source, by fluxing it into the TCA cycle. We believe our findings

provide a better understanding of how AML cells communicate with stromal cells and could serve as

a basis for the development of novel therapeutic strategies to target AML cells by modulating gap

junction formation as an adjuvant therapy.

Methods

Cell lines

The human AML cell lines (SKM-1, Kasumi-1 and HL-60), the mouse stromal cell line (MS-5) and the

human cervical cancer cell line (HeLa) were cultured in RPMI 1640 media supplemented with 15%

(v/v) FBS, 2 mmol/L L-glutamine and 100 U/ml Penicillin/Streptomycin (all from Thermo Fisher

Scientific). Co-cultures were plated in a 4:1 AML-stromal ratio, and 750.000 cells/ml density of AML

cells over confluent stromal cells. Prior to cell extraction for NMR, RNA collection or protein

extraction, a suspension of 10 million AML cells was collected, the stromal layer was washed with

PBS and was subject to mild trypsinisation with 1:5 dilution of 0.25% Trypsin 1 mM EDTA

(Thermofisher) to remove attached residual leukaemic cells before completely detaching stromal

cells with 0.25% Trypsin-EDTA.

Primary patient samples

AML and PBMC primary specimens' procedures were obtained in accordance with the Declaration of

Helsinki at the University Medical Center Groningen, approved by the UMCG Medical Ethical

Committee or at the University Hospital Birmingham NHS Foundation Trust, approved by the West

Midlands – Solihull Research Ethics Committee (10/H1206//58).

Additional information about the primary AML samples used in this study can be found in Annex 1.

Peripheral blood (PB) and bone marrow samples from AML patients and healthy donors were

obtained in heparin-coated vacutainers. Mononuclear cells were isolated using Ficoll-Paque (GE

Healthcare) and stored at -80°C.

For AML1-2 and PBMC1-2, samples were thawed and resuspended in newborn calf serum (NCS)

supplemented with DNase I (20 Units/mL), 4 mM $MgSO_4$ and heparin (5 Units/mL) and incubated on

37°C for 15 minutes (min). For AML1 and PBMC1, cells were sorted after thawing by fluorescence-

activated cell sorting (FACS) for the CD34+CD38- population using 10 µL of anti-CD34 APC (560940,

BD Biosciences), 10 µL of anti-CD38 FITC (555459, BD Biosciences) and 10 µL of DAPI (D1306, Thermo

Fisher Scientific) per 10 million cells. Cells were sorted using a Sony SH800S (Sony) sorter. For AML2

and PBMC2, cells were thawed and the CD34+ population was sorted by magnetic-activated cell

sorting (MACS) using 10 µL of anti-CD34 microbeads (130-046-702, Miltenyi) per million of expected

CD34 cells following manufacturer's protocol.

AML1-2, PBMC1-2 and MS-5 cells were cultured alone and in co-culture in a 4:1 AML/PBMC-stromal

ratio and 500,000 cells/mL density in α-MEM (Gibco) with 12.5% (v/v) FCS (Sigma-Aldrich), 1% (v/v)

Pen/Strep (Thermo Fisher Scientific), 12.5% (v/v) Horse serum (Invitrogen), 0.4% (v/v) β-
mercaptoethanol (Merck Sharp & Dohme BV) and 0.1% hydrocortisone (H0888, Sigma-Aldrich). For
AML1 and AML2 the medium was supplemented with 0.02 µg/mL of IL-3 (Sandoz), 0.02 µg/mL
NPlate (Amgen) and 0.02 µg/mL of G-CSF (Amgen). For PBMC1 and PBMC2 the medium was
supplemented with 100 ng/ml of human SCF (255-SC, Novus Biologicals), 100 ng/ml of NPlate, 100
ng/ml of FLT3 ligand (Amgen) and 20 ng/ml of IL-3. Samples of medium were collected at 0 and 48
hours.

AML3 - 7 and PBMC3 were thawed and kept in culture for 16-24 hours in Stem Span H3000 media
(STEMCELL Technologies) supplemented with 50 µg/ml ascorbic acid (Sigma-Aldrich), 50 ng/ml
human SCF (255-SC-010, R&D Systems), 10 ng/ml human IL-3 (203-IL-010, R&D Systems), 2 units/ml
human-erythropoietin (100-64, PeproTech), 40 ng/ml insulin-like growth factor 1 (IGF-1) (100-12,
PeproTech), 1 µM dexamethasone (D2915, Sigma-Aldrich) and 100 µg/ml primocin (ant-pm-2,
InvivogenCD34+ cKit+ cells were purified using magnetic microbeads (130-046-702 (CD34) and 130-
091-332 (CD117), Miltenyi Biotec). Cells were cultured in the supplemented Stem Span media for 24
hours prior to co-culture setting. Co-cultures were plated with a 4:1 leukaemic to stromal cells ratio
and a 300,000 cells/ml density in supplemented Stem Span medium. Samples of leukaemic/healthy
cells and MS-5 cells alone were also cultured in supplemented Stem Span media as controls. Samples
of medium were collected at 0 and 48 hours.

In vivo experiments

12 weeks old C57BL6/J female mice (The Jackson Laboratory) were lethally irradiated with 9Gy (2
doses of 4.5Gy separated by 3h) using an X-RAY source (Rad Source's RS 2000). Mice were
transplanted by intravenous injection 4h after with $2x10^6$ bone marrow (BM) nucleated cells isolated
as previously described [54] from leukemic male mice heterozygous for MLL-AF9 knock-in fusion
transgene or wild-type (WT) control male littermates [55]. Male transgenic and WT control MLL-AF9
were purchased from The Jackson Laboratory (Stock No: 009079) and were 6 months old when
euthanized to allow development of signs of leukemic transformation driven by the MLL-AF9 fusion
oncogene in transgenic animals. MLL-AF9 expression in BM cells derived from transgenic donors
results in development of AML with high blast counts in recipients. Recipients were euthanized 6
months after the transplant, bone marrow from femur and tibia was flushed with a syringe in 150 µL
of cold PBS and centrifuged at 15000g for 10 minutes at 4^oC. The supernatant made of bone marrow
extracellular fluid (BMEF) was kept for analysis of acetate level. Experiments were conducted with
the ethical approval of the Norwegian Food and Safety Authority. Animals were housed under

specific opportunistic and pathogen free environment at the Animal Facility of the University of Oslo,

Norway.

Proliferation analysis using CFSE

The CellTrace™ carboxyfluorescein succinimidyl ester (CFSE) Cell Proliferation Kit (C34570,

Invitrogen) was used to assess proliferation of AML cells following the manufacturer's protocol. AML

cells were stained and their fluorescence was assessed before dividing the bulk of cells into culturing

them alone or with MS-5 cells for 48 hours. Small aliquots of cells after 24 and 48 hours were

analysed by flow cytometry. Flow cytometry analysis was carried out in a CyAn ADP flow cytometer

(Beckman Coulter). Data analysis was performed using the FlowJo software package (BD).

Carbenoxolone treatment

Carbenoxolone (CBX) disodium salt (Sigma-Aldrich, C4790) was prepared fresh at 200 µM in cell

culture medium. Cells were resuspended in the CBX medium and cultured alone or in co-culture for

24 hours prior to ROS or Calcein-AM dye transfer experiments.

Cellular ROS measurements using DCFH-DA

Cellular ROS was measured by incubating cells with 100 µM 2',7'-Dichlorofluorescin diacetate (DCFH-

DA) (D6883, Sigma-Aldrich) in Hank's Balanced Salt Solution (HBSS) (Thermo Fisher Scientific) at 37°C

for 30 min protected from light. Cells were then harvested and stained with 5 µg/µL anti-human

CD33 eFluor 450 (eBioscience, P67.6) for 30 min at 4°C before flow cytometry analysis as previously

described.

Thiamine-free medium comparison

Thiamine-free medium (R9011-01, United States Biological) was prepared as per manufacturer

instructions and supplemented with 10% dialised FBS.

MS-5 cells were cultured in thiamine-free medium or control RPMI medium for 4 days prior to the

experiment. Cells were then seeded with fresh thiamine-free or control RPMI medium and incubated

for 24 hours. Samples of medium were collected at 0 and 24 hours and kept at -80ºC.

ROS-related treatments with H_2O_2 and NAC

SKM-1, Kasumi, HL-60 and MS-5 cells cultured alone and in co-culture were incubated for 24 hours in

50 µM H_2O_2 (Merck) complete cell culture medium, 5 mM N-acetylcysteine (NAC) (106425, Merck)

complete cell culture medium or control medium. Samples of medium were collected at 0 and 24
hours and kept at -80°C.

Calcein-AM dye transfer assay

Functional gap junction presence was evaluated using the fluorescent dye Calcein-AM green
(Invitrogen, C1430) adapting a previously established protocol [10]. MS-5 cells were stained with 500
nM Calcein-AM dye in complete cell culture medium for 1 hour at 37°C. Stained cells were washed
with serum-free medium for 30 min at 37°C before being co-cultured with AML cells for 3h. AML
cells were then harvested and stained with 5 μg/μL anti-human CD33 eFluor 450 (eBioscience,
P67.6) for 30 min at 4°C before flow cytometry analysis as previously described. Calcein-AM dye
transfer was quantified as the frequency of $CD33^+$ and $Calcein-AM^+$ cells.

Tracer-based NMR experiments

[U-^{13}C]Glucose (CC860P1, CortecNet) was added to RPMI 1640 medium without glucose (11879020,
Merck) to a final concentration of 2 g/L (as in the complete cell culture medium) and was
supplemented as usual with 15% (v/v) FBS, 2 mmol/L L-glutamine and 100 U/mL Pen/Strep. The
medium was prepared fresh and was filtered with a 0.2 μm syringe filter (Sartorius) before each
experiment.

4 mM sodium [2-^{13}C]acetate (279315-1G, Sigma-Aldrich) was added to complete cell culture medium
and the medium was filtered with a 0.2 μm syringe filter before each experiment. Cells were
incubated for the time indicated in each experiment before separation of cells and/or metabolite
extraction. **a**, **b**, **c**, **d**, **f**, **g**, **h** and **i** Unlabelled samples were prepared as a control for 2D-NMR
experiments and to measure acetate consumption by adding unlabelled sodium acetate trihydrate
(1.37012, Merck) to complete cell culture medium before metabolite extraction or collection of
medium samples.

Metabolite extraction

536 Suspension cells and detached adherent cells were washed with PBS before being rapidly
537 resuspended in 400 μl of HPLC grade methanol pre-chilled on dry ice. Samples were transferred to
538 glass vials and were subject to 10:8:10 methanol-water-chloroform extraction as described in
539 (Saborano 2019). Polar phase samples were evaporated in a SpeedVac concentrator and were
540 subsequently kept at -80°C.

Sample preparation for NMR

Medium samples or bone marrow extracellular fluid samples were mixed with a D_2O phosphate buffer containing TMSP ((3-trimethylsilyl)propionic-(2,2,3,3-d_4)-acid sodium salt) and NaN_3 to a final concentration of 0.1 M phosphate and 0.5 mM TMSP (3-trimethylsilyl)propionic-(2,2,3,3-d_4)-acid sodium salt) and 1.5 mM NaN_3 (all from Sigma) and transferred to 3 mm NMR tubes (Cortecnet).

Dried polar extracts for tracer-based NMR experiments were reconstituted in 50 μL of 0.1 M phosphate buffer in 100% D_2O with 3 mM NaN_3 and 0.5 mM TMSP. Samples were sonicated for 10 min and transferred to 1.7 mm NMR tubes using a micro pipet system. Samples were prepared freshly before the acquisition.

NMR acquisition and analysis

All NMR data were acquired on Bruker 600MHz spectrometers equipped with Avance-III consoles using cooled Bruker SampleJet autosamplers. For media samples, a 5mm triple resonance cryoprobe (TCI) z-axis pulsed field gradient (PFG) cryogenic probe was used, and for cell extracts, a TCI 1.7mm z-PFG cryogenic probe was used. Probes were equipped with a cooled SampleJet autosampler (Bruker) and automated tuning and matching.

For medium samples, spectra were acquired at 300K using 1D [1]H-NOESY (Nuclear Overhauser effect spectroscopy) pulse sequence with pre-saturation water suppression (noesygppr1d, standard pulse sequence from Bruker). The spectral width was 12 ppm, the number of data points was TD 32,768, the interscan delay was 4 sec and the NOE mixing time was 10 msec. The [1]H carrier was set on the water frequency and the [1]H 90º pulse was calibrated at a power of 0.256 W and had a typical length of ca 7-8 μs. 64 scans and 8 dummy scans were acquired and the total experimental time was 7.5 min. Spectra were processed using the MetaboLab [56] software within the MATLAB environment (MathWorks). A 0.3Hz exponential apodization function was applied to FIDs followed by zero-filling to 131,072 data points prior to Fourier transformation. The chemical shift was calibrated by referencing the TMSP signal to 0 ppm and spectra were manually phase corrected. The baseline was corrected by applying a spline baseline correction, the water and edge regions of the spectra were excluded before scaling the spectra using a probabilistic quotient normalization (PQN). Chenomx 7.0 software (Chenomx Inc.) and the human metabolome database (HMDB) were used to assign the metabolites present in the acquired spectra. Metabolite signal intensities were obtained directly from the spectra and were normalised to a control medium sample obtained at time 0 hours. For bone marrow extracellular fluid samples, metabolite signal areas were obtained directly from the spectra.

For ^{13}C-filtered ^1H-NMR experiments, [U-^{13}C]glucose labelled medium samples were analysed with

^{13}C-filtered ^1H-NMR spectroscopy as described in [57]. Spectra were acquired at 300 K using a double

gradient BIRD filter pulse sequence developed in-house [57]. A pulse program combining the $^1H[^{12}C]$

and the all-^1H experiments in scan-interleaved mode was used. The difference between the two FIDs

gave the $^1H[^{13}C]$ signal. The spectral width was 12 ppm, the number of data points was 16,384 in

each dimension and the relaxation delay was 5.3 sec. 256 scans with 64 dummy scans were acquired

and the experimental time was 15 min. ^{13}C-filtered ^1H-NMR spectra were processed in Topspin 4.0.5

(Bruker). Spectra were zero filled to 32,768 data points before Fourier transformation. Phase

correction was applied to the $^1H[-^{12}C]$ and the all-^1H spectra and the difference $^1H[^{13}C]$ spectrum was

obtained by aligning on chemical shift. Metabolites were selected in the $^1H[^{13}C]$ spectrum and

integrated in all the spectra ($^1H[^{12}C]$, $^1H[^{13}C]$ and all-^1H). Label incorporations (^{13}C percentages) were

calculated by dividing the signal areas obtained in the $^1H[^{13}C]$ spectrum by the ones obtained in the

all-^1H spectrum.

For ^1H-^{13}C-HSQC experiments, spectra were acquired using a modified version of Bruker's

hsqcgphprsp pulse program, with additional gradient pulses during the INEPT echo periods and using

soft 180º pulses for ^{13}C. For the ^1H dimension, the spectral width was 13.018 ppm with 2048

complex points. For the ^{13}C dimension, the spectral width was 160 ppm with 2048 complex points. 2

scans were acquired per spectrum and with an interscan delay of 1.5 sec. Non-uniform sampling

(NUS) with a 25% sampling schedule (generated using the Wagner's schedule generator (Gerhard

Wagner Lab, Harvard Medical School) with a tolerance of 0.01 and default values for the other

parameters) was used with 4096 increments yielding 8192 complex points after processing. The total

experimental time was 4 hours. Spectra were processed and phased with NMRPipe (National

Institute of Standards and Technology of the U.S.) (Appendix 8.1). MetaboLab was used to reference

the chemical shift using the signal for the methyl group of L-lactic acid, at 1.31/22.9 ppm in the ^1H

and ^{13}C dimensions, respectively. Identification of metabolites in 2D spectra was carried out using

the MetaboLab software which includes a chemical shift library for ca. 200 metabolites. Intensities

were obtained from signals in the spectra and were corrected for differences in cell numbers

contributing to the sample as follows: A ^1H-NMR spectrum was acquired for each sample and the

total metabolite area of this spectrum after removal of the water and TMSP reference signal was

calculated in MetaboLab. The intensities in the 2D spectra were then divided by the total metabolite

area of the corresponding ^1H-NMR spectrum. To obtain the % of ^{13}C in a metabolite, the normalized

intensity of a certain carbon in the labelled sample was divided by the normalized intensity of the

same carbon in the unlabelled sample and was multiplied by the natural abundance of ^{13}C (1.1%).

RNA extraction and sequencing

MS-5 cells were cultured alone or in co-culture with SKM-1 cells for 24 hours. Cells were separated

and washed with PBS prior to RNA extraction with TRIzol (Gibco) according to the manufacturer's

protocol. RNA was purified with RNeasy Plus Micro kit. Samples were sent to Theragenetex to be

sequenced with Novaseq 150bp PE with 40M reads.

Real-time PCR

Samples of RNA from SKM-1 and MS-5 co-cultures were collected and extracted using Trizol

(Invitrogen) following manufacturer's protocol. cDNA was synthesized using the M-MLV reverse

transcriptase (Promega) according to manufacturer's instructions. For gene expression analysis, qRT-

PCR of *Hk2* (NM_013820), *Pdhx* (NM_175094), *Pdk1* (NM_172665) and *Pdk2* (NM_133667) (all

KiCqStart™ primers KSPQ12012, Sigma Aldrich) were carried out using the SYBRGreen Master mix

(Thermo Fisher Scientific) and qRT-PCR of *B2m* (NM_009735.3, TaqMan® assays, Thermo Fisher

Scientific) was performed using TaqMan PCR Master Mix (Thermo Fisher Scientific). Reactions were

performed in a Stratagene Mx3000P and were run in triplicate. Relative gene expression was

calculated following the $2^{-\Delta\Delta Ct}$ method relative to the expression of *B2m*.

RNA sequencing

RNA samples of MS-5 cells cultured alone and in co-culture with SKM-1 cells extracted using Trizol

were purified using the Rneasy Plus Micro kit (Qiagen) according to manufacturer's protocol.

Transcriptome analysis was performed by Theragen Etex Co., LTD. (www.theragenetex.net). cDNA

libraries were prepared with the TruSeq Stranded mRNA Sample Prep Kit (Illumina) and RNA

sequencing was performed in a HiSeq2500 platform (Illumina). Quality control metrics were

obtained with FastQC 0.11.7 software (Babraham Bioinformatics). To quantify transcript abundances

using the Kallisto 0.43.0 software (Patcher Lab), read counts were aligned to the GRCm38 mouse

reference genome cDNA index (Ensembl rel.67). Gene-level differential expression analysis was

carried out with the R statistical package Sleuth 0.30.0 comparing the expression of cells cultured

alone vs co-culture. Differentially expressed genes were calculated using the Wald statistical test,

correcting for multiple testing with the Benjamin-Hochberg method. The false discovery rate (FDR)

threshold was set at 1% (q-values < 0.01). Ensembl gene transcripts were annotated to Entrez IDs

and official gene symbols with the R statistical package BioMart 2.40.3. To normalise for sequencing

depth and gene length, transcripts per million (TPM) expression values were calculated. Gene Set

Enrichment Analysis (GSEA) was performed with the R statistical package fgsea 1.10.0. The collection

of hallmark gene sets from the Molecular Signature Database was used for the GSEA, setting the FDR

threshold at 5%. Data was deposited in GEO (GSE163478).

References

1.   Kreitz, J*., et al.* Metabolic Plasticity of Acute Myeloid Leukemia. *Cells* **8**(2019).

2.   Schelker, R.C*., et al.* TGF-β1 and CXCL12 modulate proliferation and chemotherapy
sensitivity of acute myeloid leukemia cells co-cultured with multipotent mesenchymal
stromal cells. *Hematology* **23**, 337-345 (2018).

3.   Passaro, D*., et al.* Increased Vascular Permeability in the Bone Marrow Microenvironment
Contributes to Disease Progression and Drug Response in Acute Myeloid Leukemia. *Cancer*
Cell **32**, 324-341 e326 (2017).

4.   Zeng, Z*., et al.* Inhibition of CXCR4 with the novel RCP168 peptide overcomes stroma-
mediated chemoresistance in chronic and acute leukemias. *Mol Cancer Ther* **5**, 3113-3121
(2006).

5.   Carey, A*., et al.* Identification of Interleukin-1 by Functional Screening as a Key Mediator of
Cellular Expansion and Disease Progression in Acute Myeloid Leukemia. *Cell reports* **18**,
3204-3218 (2017).

6.   Zhang, T.Y*., et al.* IL-6 blockade reverses bone marrow failure induced by human acute
myeloid leukemia. **12**, eaax5104 (2020).

7.   Wang, B*., et al.* Exosomes derived from acute myeloid leukemia cells promote
chemoresistance by enhancing glycolysis-mediated vascular remodeling. *J Cell Physiol* **234**,
10602-10614 (2019).

8.   Kumar, B*., et al.* Acute myeloid leukemia transforms the bone marrow niche into a leukemia-
permissive microenvironment through exosome secretion. *Leukemia* **32**, 575-587 (2018).

9.   Hornick, N.I*., et al.* AML suppresses hematopoiesis by releasing exosomes that contain
microRNAs targeting c-MYB. *Science signaling* **9**, ra88 (2016).

10.   Kouzi, F*., et al.* Disruption of gap junctions attenuates acute myeloid leukemia
chemoresistance induced by bone marrow mesenchymal stromal cells. *Oncogene* **39**, 1198-
1212 (2020).

11.   Omsland, M., Bruserud, Ø., Gjertsen, B.T. & Andresen, V. Tunneling nanotube (TNT)
formation is downregulated by cytarabine and NF-κB inhibition in acute myeloid leukemia
(AML). *Oncotarget* **8**, 7946-7963 (2017).

12.   Ye, H*., et al.* Leukemic Stem Cells Evade Chemotherapy by Metabolic Adaptation to an
Adipose Tissue Niche. *Cell Stem Cell* **19**, 23-37 (2016).

13.   Moschoi, R*., et al.* Protective mitochondrial transfer from bone marrow stromal cells to
acute myeloid leukemic cells during chemotherapy. *Blood* **128**, 253-264 (2016).

14.   Forte, D*., et al.* Bone Marrow Mesenchymal Stem Cells Support Acute Myeloid Leukemia
Bioenergetics and Enhance Antioxidant Defense and Escape from Chemotherapy. *Cell Metab*
32, 829-843 e829 (2020).

15.   Shafat, M.S*., et al.* Leukemic blasts program bone marrow adipocytes to generate a
protumoral microenvironment. *Blood* **129**, 1320-1332 (2017).

16.   Tabe, Y*., et al.* Bone Marrow Adipocytes Facilitate Fatty Acid Oxidation Activating AMPK and
a Transcriptional Network Supporting Survival of Acute Monocytic Leukemia Cells. *Cancer*
Res **77**, 1453-1464 (2017).

17.   Baccelli, I*., et al.* Mubritinib Targets the Electron Transport Chain Complex I and Reveals the
Landscape of OXPHOS Dependency in Acute Myeloid Leukemia. *Cancer Cell* **36**, 84-99.e88
(2019).

18.   Pollyea, D.A*., et al.* Venetoclax with azacitidine disrupts energy metabolism and targets
leukemia stem cells in patients with acute myeloid leukemia. *Nat Med* **24**, 1859-1866 (2018).

19.   Molina, J.R*., et al.* An inhibitor of oxidative phosphorylation exploits cancer vulnerability.
Nature Medicine **24**, 1036-1046 (2018).

20.   Lagadinou, E.D*., et al.* BCL-2 inhibition targets oxidative phosphorylation and selectively
eradicates quiescent human leukemia stem cells. *Cell Stem Cell* **12**, 329-341 (2013).

21.   Farge, T.*, et al.* Chemotherapy-Resistant Human Acute Myeloid Leukemia Cells Are Not
692   Enriched for Leukemic Stem Cells but Require Oxidative Metabolism. *Cancer Discov* **7**, 716-
693   735 (2017).
694   22.   Li, L.*, et al.* Altered Hematopoietic Cell Gene Expression Precedes Development of Therapy-
695   Related Myelodysplasia/Acute Myeloid Leukemia and Identifies Patients at Risk. *Cancer Cell*
696   **20**, 591-605 (2011).
697   23.   Hole, P.S.*, et al.* Overproduction of NOX-derived ROS in AML promotes proliferation and is
698   associated with defective oxidative stress signaling. *Blood* **122**, 3322-3330 (2013).
699   24.   Vignon, C.*, et al.* Involvement of GPx-3 in the Reciprocal Control of Redox Metabolism in the
700   Leukemic Niche. *Int J Mol Sci* **21**(2020).
701   25.   Itoh, K.*, et al.* Reproducible establishment of hemopoietic supportive stromal cell lines from
702   murine bone marrow. *Exp Hematol* **17**, 145-153 (1989).
703   26.   Childress, C.C., Sacktor, B. & Traynor, D.R. Function of carnitine in the fatty acid oxidase-
704   deficient insect flight muscle. *J Biol Chem* **242**, 754-760 (1967).
705   27.   Stephens, F.B., Constantin-Teodosiu, D. & Greenhaff, P.L. New insights concerning the role of
706   carnitine in the regulation of fuel metabolism in skeletal muscle. *J Physiol* **581**, 431-444
707   (2007).
708   28.   Liu, X.*, et al.* Acetate Production from Glucose and Coupling to Mitochondrial Metabolism in
709   Mammals. *Cell* **175**, 502-513 e513 (2018).
710   29.   Vysochan, A., Sengupta, A., Weljie, A.M., Alwine, J.C. & Yu, Y. ACSS2-mediated acetyl-CoA
711   synthesis from acetate is necessary for human cytomegalovirus infection. *Proc Natl Acad Sci*
712   *U S A* **114**, E1528-E1535 (2017).
713   30.   Tiziani, S.*, et al.* Metabolomic profiling of drug responses in acute myeloid leukaemia cell
714   lines. *PLoS One* **4**, e4251 (2009).
715   31.   Nogueira-Pedro, A.*, et al.* Hydrogen peroxide (H2O2) induces leukemic but not normal
716   hematopoietic cell death in a dose-dependent manner. *Cancer Cell International* **13**, 123
717   (2013).
718   32.   Taniguchi Ishikawa, E.*, et al.* Connexin-43 prevents hematopoietic stem cell senescence
719   through transfer of reactive oxygen species to bone marrow stromal cells. *Proc Natl Acad Sci*
720   *U S A* **109**, 9071-9076 (2012).
721   33.   Davidson, J.S., Baumgarten, I.M. & Harley, E.H. Reversible inhibition of intercellular
722   junctional communication by glycyrrhetinic acid. *Biochem Biophys Res Commun* **134**, 29-36
723   (1986).
724   34.   Davidson, J.S. & Baumgarten, I.M. Glycyrrhetinic acid derivatives: a novel class of inhibitors
725   of gap-junctional intercellular communication. Structure-activity relationships. *J Pharmacol*
726   *Exp Ther* **246**, 1104-1107 (1988).
727   35.   Lyssiotis, C.A. & Cantley, L.C. Acetate fuels the cancer engine. *Cell* **159**, 1492-1494 (2014).
728   36.   Comerford, S.A.*, et al.* Acetate dependence of tumors. *Cell* **159**, 1591-1602 (2014).
729   37.   Gao, X.*, et al.* Acetate functions as an epigenetic metabolite to promote lipid synthesis under
730   hypoxia. *Nat Commun* **7**, 11960 (2016).
731   38.   Schug, Z.T.*, et al.* Acetyl-CoA synthetase 2 promotes acetate utilization and maintains cancer
732   cell growth under metabolic stress. *Cancer Cell* **27**, 57-71 (2015).
733   39.   Yoshii, Y.*, et al.* Cytosolic acetyl-CoA synthetase affected tumor cell survival under hypoxia:
734   the possible function in tumor acetyl-CoA/acetate metabolism. *Cancer Sci* **100**, 821-827
735   (2009).
736   40.   Sousa, C.M.*, et al.* Pancreatic stellate cells support tumour metabolism through autophagic
737   alanine secretion. *Nature* **536**, 479-483 (2016).
738   41.   Sonveaux, P.*, et al.* Targeting lactate-fueled respiration selectively kills hypoxic tumor cells in
739   mice. *J Clin Invest* **118**, 3930-3942 (2008).

42. Whitaker-Menezes, D., *et al.* Evidence for a stromal-epithelial "lactate shuttle" in human
tumors: MCT4 is a marker of oxidative stress in cancer-associated fibroblasts. *Cell Cycle* **10**,
1772-1783 (2011).
43. Saborano, R., *et al.* A framework for tracer-based metabolism in mammalian cells by NMR.
Sci Rep **9**, 2520 (2019).
44. Mashimo, T., *et al.* Acetate is a bioenergetic substrate for human glioblastoma and brain
metastases. *Cell* **159**, 1603-1614 (2014).
45. Testa, U., Labbaye, C., Castelli, G. & Pelosi, E. Oxidative stress and hypoxia in normal and
leukemic stem cells. *Experimental Hematology* **44**, 540-560 (2016).
46. Kocabas, F., *et al.* Hypoxic metabolism in human hematopoietic stem cells. *Cell & bioscience*
5, 39 (2015).
47. Takubo, K., *et al.* Regulation of glycolysis by Pdk functions as a metabolic checkpoint for cell
cycle quiescence in hematopoietic stem cells. *Cell Stem Cell* **12**, 49-61 (2013).
48. Shan, T., *et al.* Cancer-associated fibroblasts enhance pancreatic cancer cell invasion by
remodeling the metabolic conversion mechanism. *Oncol Rep* **37**, 1971-1979 (2017).
49. Migneco, G., *et al.* Glycolytic cancer associated fibroblasts promote breast cancer tumor
growth, without a measurable increase in angiogenesis: evidence for stromal-epithelial
metabolic coupling. *Cell Cycle* **9**, 2412-2422 (2010).
50. Pavlides, S., *et al.* The reverse Warburg effect: aerobic glycolysis in cancer associated
fibroblasts and the tumor stroma. *Cell Cycle* **8**, 3984-4001 (2009).
51. Cruz-Bermudez, A., *et al.* Cancer-associated fibroblasts modify lung cancer metabolism
involving ROS and TGF-beta signaling. *Free Radic Biol Med* **130**, 163-173 (2019).
52. Hole, P.S., *et al.* Overproduction of NOX-derived ROS in AML promotes proliferation and is
associated with defective oxidative stress signaling. *Blood* **122**, 3322-3330 (2013).
53. Robinson, A.J., *et al.* Reactive Oxygen Species Drive Proliferation in Acute Myeloid Leukemia
via the Glycolytic Regulator PFKFB3. *Cancer Res* **80**, 937-949 (2020).
54. Arranz, L., *et al.* Neuropathy of haematopoietic stem cell niche is essential for
myeloproliferative neoplasms. *Nature* **512**, 78-81 (2014).
55. Corral, J., *et al.* An Mll-AF9 fusion gene made by homologous recombination causes acute
leukemia in chimeric mice: a method to create fusion oncogenes. *Cell* **85**, 853-861 (1996).
56. Ludwig, C. & Gunther, U.L. MetaboLab--advanced NMR data processing and analysis for
metabolomics. *BMC Bioinformatics* **12**, 366 (2011).
57. Reed, M.A.C., Roberts, J., Gierth, P., Kupce, E. & Gunther, U.L. Quantitative Isotopomer Rates
in Real-Time Metabolism of Cells Determined by NMR Methods. *Chembiochem* **20**, 2207-
2211 (2019).
775

Acknowledgements

N. Vilaplana-Lopera, G. Papatzikas, A. Cunningham, and A. Erdem were supported by the EU grant HaemMetabolome H2020-MSCA-ITN-2015-675790. U. Günther, P. Garcia, J. J. Schuringa, J.-B. Cazier, and F. Schnütgen acknowledge support from the European Commission (HaemMetabolome [EC-675790]). This work was further supported by the Deutsche Forschungsgemeinschaft (DFG, German Research foundation) SFB815, TP A10 (F.S.). We also acknowledge the Wellcome Trust for supporting access to NMR instruments at the Henry Wellcome Building for Biomolecular NMR in Birmingham (grant number 208400/Z/17/Z). The mouse work was supported by a joint meeting grant of the Northern Norway Regional Health Authority and UiT (2014/5668) to L. Arranz. We thank A. Villatoro and L.M. Gonzalez for technical assistance.

Author Contributions

N.V-L. conceived and performed the experiments, acquired, analysed and interpreted the data, and wrote the manuscript. V.C., R.A., E.G., A.C., A.E., F.S., M.R., L.A. performed the experiments and edited the manuscript. G.P. analysed the RNA-seq data. J.B helped with the experimental design. S.P. and M.R. provided with patient samples. V.C. and L.A. provided mouse bone marrow extracellular fluid samples. J.J.S. provided patient samples, helped with the experimental design, interpretation and critical discussion of the data and edited the manuscript. U.G and P.G. conceived and performed the experiments, acquired and interpreted the data, wrote the manuscript, and managed the project.

EXTENDED DATA: **Crosstalk between AML and stromal cells triggers**

acetate secretion through the metabolic rewiring of stromal cells

Nuria Vilaplana-Lopera[1], Ruba Almaghrabi[1], Grigorios Papatzikas[1], Vincent Cuminetti[2], Mark Jeeves[1],

Elena González[1], Alan Cunningham[3], Ayşegül Erdem[3], Frank Schnuetgen[4,5,6], Manoj Raghavan[1,7],

Sandeep Potluri[1,7], Jean-Baptiste Cazier[1], Jan Jacob Schuringa[3], Michelle AC Reed[1], Lorena Arranz[2],

Ulrich L Günther[1,8*] and Paloma Garcia[1,9*],

[1] Institute of Cancer and Genomic Sciences, University of Birmingham, Birmingham B15 2TT, United Kingdom

[2] Stem Cell Ageing and Cancer Research Group, Department of Medical Biology, Faculty of Health Sciences, UiT – The Arctic
University of Norway, 9019 Tromsø, Norway

[3] Department of Experimental Hematology, University Medical Center Groningen, University of Groningen, Hanzeplein 1,
9700 RB, Groningen, The Netherlands.

[4] Department of Medicine, Hematology/Oncology, University Hospital Frankfurt, Goethe-University, 60590 Frankfurt,
Germany.

[5] Frankfurt Cancer Institute, Goethe-University Frankfurt, 60596 Frankfurt, Germany.

[6] German Cancer Consortium (DKTK), partner site Frankfurt/Mainz, and German Cancer Research Center (DKFZ),
Heidelberg, Germany

[7] Centre for Clinical Haematology, University Hospitals Birmingham NHS Foundation Trust, Queen Elizabeth Hospital,
Queen Elizabeth Medical Centre, Birmingham, United Kingdom.

[8] Institute of Chemistry and Metabolomics, University of Lübeck, 23562 Lübeck, Germany

[9] Corresponding author

[*] These two authors are co-senior authors.

Extended Data Figure 1. Other extracellular metabolite levels in co-cultures with SKM-1 and MS-5 cells

Sections of [1]H-NMR spectra from extracellular medium samples of SKM-1 cells cultured alone (blue), MS-5 cells cultured alone (red) and SKM-1 and MS-5 cells in co-culture (green) after 24 hours of culture, corresponding to glucose, lactate, glutamate and glutamine.

(**a**) CFSE cell proliferation analysis in AML cell lines alone or in co-culture with MS-5 cells. The population of living cells was gated, and 1500 cells for each condition were randomly selected and plotted. The geometric mean for each population and time point was compared between cells alone or in co-culture by performing an unpaired unpaired Student's t-test and p-values were represented as n.s. for not significant. Each histogram is representative of n=3 independent experiments. (**b**) Extracellular acetate levels in SKM-1 cells cultured alone (blue), HeLa cells cultured alone (red) and AML and HeLa cells in co-culture (green) at 0, 20 and 24 hours. Each point represents the mean of n=3 independent experiments and error bars represent standard deviation. An unpaired Student's t-test was applied for each condition (black brackets) and p-values are represented by n.s. for not significant. (**c**) Linear regression of acetate concentrations and detected intensities in ^{1}H-NMR

847   spectra (Intensity=$1.38 \cdot 10^{6}$ ± $1.6 \cdot 10^{4}$ x Acetate concentration + $2.27 \cdot 10^{5}$ ± $3.3 \cdot 10^{4}$; R^{2}=0.9987). The

848   intensities detected in samples of co-culture were interpolated to obtain the estimate acetate

849   concentration in co-culture (3.09 mM). Each point represents the mean of n=3 independent

850   experiments and error bars represent standard deviation.

S3

Extended Data Figure 3. Label incorporation from [2-^{13}C]acetate in SKM-1 and MS-5 cells cultured alone

(**a**) Example of metabolites assigned in a ^{1}H-^{13}C-HSQC spectrum from a polar extract from SKM-1 cells cultured with [2-^{13}C]acetate for 2 hours. ^{13}C percentages were calculated in 13 carbons from 11 different metabolites. The exact chemical shifts for each carbon are: acetate C2, 1.91 ppm (^{1}H) – 26.1 ppm (^{13}C); aspartate C2, 3.89 ppm (^{1}H) – 55.0 ppm (^{13}C); aspartate C3, 2.82 ppm (^{1}H) – 39.4 ppm (^{13}C); acetylcarnitine C9, 2.15 ppm (^{1}H) – 23.4 ppm (^{13}C); citrate C2, 2.52 ppm (^{1}H) – 48.2 ppm (^{13}C);

fumarate C2, 6.51 ppm (^{1}H) − 138.2 ppm (^{13}C); glutamate C4, 2.34 ppm (^{1}H) − 36.1 ppm (^{13}C);

glutamine C4, 2.44 ppm (^{1}H) −33.7 ppm (^{13}C); oxoglutarate C4, 2.43 ppm (^{1}H) − 33.4 ppm (^{13}C);

glutathione C4, 2.55 ppm (^{1}H) − 34.2 ppm (^{13}C); malate C2, 4.28 ppm (^{1}H) − 73.2 ppm (^{13}C); malate

C3, 2.68 ppm (^{1}H) − 45.4 ppm (^{13}C); and proline C4, 2 ppm (^{1}H) − 26.6 ppm (^{13}C). (**b**) ^{13}C percentages

on polar metabolites in Kasumi-1 and HL-60 cells in co-culture with MS-5 cells. Cells were co-

cultured for 24 hours before the addition of extra 4 mM sodium [2-^{13}C]acetate. Bars represent the

mean of the ^{13}C percentage and error bars represent the standard deviation of n=3 independent

experiments. ^{13}C natural abundance is represented as a black bar at %^{13}C = 1.1.

S4 a

b

c

Extended Data Figure 4. **PCA component analysis, heat map of differentially expressed**

genes and qPCR for MS-5 cells cultured alone and in co-culture with SKM-1 cells

(**a**) PCA plot showing the clustering of individual samples of MS-5 cells cultured alone (MO) and MS-5

cells in co-culture (MC). The x and y axis values represent the variation between the sample groups.

Generated using Sleuth 0.30.0 R statistical package. (**b**) Heat map of differentially expressed genes in

MS-5 only (red) vs MS-5 co-culture (blue) calculated using the Wald statistical test, correcting for

multiple testing comparison employing the Benjamini-Hochberg method using a false discovery rate

threshold of 1% (q-value<0.01). Generated using Sleuth 0.30.0 R statistical package. **(c)** Quantitative

PCR mRNA expression values for RNA sequencing validation of MS-5 cells cultured alone and in co-

culture with SKM-1 cells. The gene set chosen was *Hk2*, *Pdhx*, *Pdk1* and *Pdk2*. mRNA quantification

was normalized to *B2m* house-keeping gene. Bars represent the mean and error bars represent the

SEM of three independent experiments. An unpaired Student's t-test was applied for each gene and

p-values are represented by * for p-value<0.05, ** for p-value<0.01 and *** for p-value<0.001.

S5

(**a**) Extracellular acetate levels in MS-5 cells cultured in control medium (black) or in thiamine-depleted medium (grey) after 24 hours. (**b**) Extracellular acetate levels in Kasumi (navy blue) and HL-60 (red) cultured alone and in co-culture (light blue and light red) for 24 hours in a control medium, medium with 50 μM H_2O_2 or medium with 5 mM NAC. Acetate levels for HL-60 cells cultured alone or under co-culture with 50 μM H_2O_2 are not shown as treatment 50 μM H_2O_2 severely impaired their viability. N.A. = not analysed. For **a** and **b** each point represents the mean of n=3 independent experiments and error bars represent standard deviation. An unpaired Student's t-test was applied for each condition (black brackets) and p-values are represented by n.s. for not significant. * for p-value<0.05 and ** for p-value<0.01. (**c**) Representative flow cytometry diagrams showing CD33 and calcein-AM green fluorescence in SKM-1 cells cultured alone or SKM-1 cells in co-culture with Calcein-AM-stained MS-5 cells after 3 hours. Frequencies of gated cell populations are indicated.

899   **Extended Data Table 1.** **Primary AML samples' additional information**

Samples obtained from Martini Hospital, UMCG

(used in figure 1e)

Patient	Tissue of origin	Type of AML (FAB)	Karyotype	Other mutations	Risk
AML1	PB	M5	Normal	FLT3-ITD; NPM1; DNMTA3; IDH2	Adv
AML2	PB	M1	Unknown	-	Adv

Samples obtained from University Hospital Birmingham NHS Foundation Trust, UoB

(used in figure 1e and 4c)

Patient	Tissue of origin	Type of disease	Karyotype	Other mutations
AML3	PB	Secondary AML from MDS	46,XY	TET2; SRSF2; RUNX1; CEBPA; ASXL1
AML4	PB	Secondary AML from MDS	46,XY	NPM1 neg FLT3 neg
AML5	PB	AML M1 (FAB)	46 XX	FLT3-ITD, 2xWT1, CEBPA
AML6	PB	AML M2 (FAB)	46 XX	-
AML7	PB	AML M4 (FAB)	46 XY	-