Title Page

Optimum growth temperature declines with body size within fish species

Running title: Optimum growth temperature declines with size

Max Lindmark[a,1], Jan Ohlberger[b], Anna Gårdmark[c]

[a] Swedish University of Agricultural Sciences, Department of Aquatic Resources, Institute of

Coastal Research, Skolgatan 6, Öregrund 742 42, Sweden

[b] School of Aquatic and Fishery Sciences (SAFS), University of Washington, Box 355020, Seattle,

WA 98195-5020, USA

[c] Swedish University of Agricultural Sciences, Department of Aquatic Resources, Skolgatan 6,

SE-742 42 Öregrund, Sweden

[1] Author to whom correspondence should be addressed. Current address:

Max Lindmark, Swedish University of Agricultural Sciences, Department of Aquatic Resources,

Institute of Marine Research, Turistgatan 5, Lysekil 453 30, Sweden, Tel.: +46(0)104784137,

email: max.lindmark@slu.se

Keywords: body growth, metabolic rate, consumption rate, temperature-size rule, metabolic

theory of ecology, climate change, warming, scaling, allometry

Abstract

According to the temperature-size rule, warming of aquatic ecosystems is generally predicted to

increase individual growth rates but reduce asymptotic body sizes of ectotherms. However, we

lack a comprehensive understanding of how growth and key processes affecting it, such as

consumption and metabolism, depend on both temperature and body mass within species. This

limits our ability to inform growth models, link experimental data to observed growth patterns,

and advance mechanistic food web models. To examine the combined effects of body size and

temperature on individual growth, as well as the link between maximum consumption, metabolism

and body growth, we conducted a systematic review and compiled experimental data on fishes

from 52 studies that combined body mass and temperature treatments. By fitting hierarchical

models accounting for variation between species, we estimated how maximum consumption and

metabolic rate scale jointly with temperature and body mass within species. We found that whole-

organism maximum consumption increases more slowly with body mass than metabolism, and is

unimodal over the full temperature range, which leads to the prediction that optimum growth

temperatures decline with body size. Using an independent dataset, we confirmed this negative

relationship between optimum growth temperature and body size. Small individuals of a given

population may therefore exhibit increased growth with initial warming, whereas larger

conspecifics could be the first to experience negative impacts of warming on growth. These

findings help advance mechanistic models of individual growth and food web dynamics and

improve our understanding of how climate warming affects the growth and size structure of aquatic

ectotherms.

Introduction

Individual body growth is a fundamental process powered by metabolism, and thus depends on body size and temperature (Brown et al., 2004). It affects individual fitness and life history traits, such as maturation size, population growth rates (Savage et al., 2004), and ultimately energy transfer across trophic levels (Andersen et al., 2009; Barneche & Allen, 2018). Therefore, understanding how growth scales with body size and temperature is important for predicting the impacts of global warming on the structure and functioning of ecosystems.

Global warming is predicted to lead to declining body sizes of organisms (Daufresne et al., 2009; Gardner et al., 2011). The temperature size-rule ('TSR') states that warmer rearing temperatures lead to faster developmental times (and larger initial size-at-age or size-at-life-stage), but smaller adult body sizes in ectotherms (Atkinson, 1994; Ohlberger, 2013). This relationship is found in numerous experimental studies (Atkinson, 1994), is reflected in latitudinal gradients (Horne et al., 2015), and is stronger in aquatic than terrestrial organisms (Forster et al., 2012; Horne et al., 2015). Support for the TSR exists in fishes, in particular in young fish, where reconstructed individual growth histories often reveal positive correlations between growth rates and temperature in natural systems (Baudron et al., 2014; Huss et al., 2019; Neuheimer et al., 2011; Thresher et al., 2007). However, whether the positive effect of warming on growth is indeed limited to small individuals within a species, as predicted by the temperature size-rule, is less clear. Negative correlations between maximum size, asymptotic size or size-at-age of old fish and temperature have been found in commercially exploited fish species (Baudron et al., 2014; Ikpewe et al., 2020; van Rijn et al., 2017). However, other studies, including large scale experiments, controlled experiments and latitudinal studies or observational data on unexploited species, have found no or less clear negative relationships between maximum size, growth of old fish or mean

size and temperature (Audzijonyte et al., 2020; Barneche et al., 2019; Denderen et al., 2020; Huss

et al., 2019; van Dorst et al., 2019) and differences between species may be related to life history

traits and depend on local environmental conditions (Denderen et al., 2020; Wang et al., 2020).

While the support for TSR is mixed, and the underlying mechanisms are not well understood

(Audzijonyte et al., 2019; Neubauer & Andersen, 2019; Ohlberger, 2013), theoretical growth

models, such as Pütter growth models (Pütter, 1920), including the von Bertalanffy growth model

(VBGM) (von Bertalanffy, 1957), commonly predict declines in asymptotic body mass with

temperature and declines in optimum growth temperature with body mass, in line with the TSR

(Morita et al., 2010; Pauly, 2021; Pauly & Cheung, 2018a; Perrin, 1995). Yet, the physiological

basis of these models has been questioned, as the commonly applied scaling parameters (mass

exponents) tend to differ from empirical estimates (Lefevre et al., 2018; Marshall & White, 2019).

Hence, despite attempting to describe growth from first principles, Pütter growth models can also

be viewed as phenomenological. In more mechanistic growth models, the difference between

energy gain and expenditure is partitioned between somatic growth and gonads (Essington et al.,

2001; Jobling, 1997; Kitchell et al., 1977; Ursin, 1967). Energy gain is normally the amount of

energy extracted from consumed food, and expenditure is defined as maintenance, activity and

feeding metabolism. These components of the energetics of growth are found in dynamic energy

budget models (Kitchell et al., 1977; Kooijman, 1993), including those used in physiologically

structured population models (PSPMs) (de Roos & Persson, 2001) and size-spectrum models

(Blanchard et al., 2017; Hartvig et al., 2011; Maury & Poggiale, 2013). Therefore, it is important

to understand how consumption and metabolism rates scale with body mass and temperature in

order to understand if and how body growth of large fish within populations is limited by

temperature, and to evaluate the physiological basis of growth models.

Moreover, the effect of body mass and temperature on growth dynamics should be evaluated

over ontogeny at the intraspecific level (within species), which better represents the underlying

process than interspecific data (among species) (Marshall & White, 2019). For instance, we are

not aware of an interspecific relationship between optimum growth temperature and body mass,

but within species it may have a large effect on growth dynamics. Despite this, intraspecific body

mass and temperature scaling is often inferred from interspecific data, and we know surprisingly

little about average relationship between consumption and metabolic exponents within species

(Marshall & White, 2019). Importantly, how physiological rates depend on mass and temperature

within species can differ from the same relationships across species (Glazier, 2005; Jerde et al.,

2019; Rall et al., 2012). Across species, rates are often assumed and found to scale as power

functions of mass with exponents of 3/4 for whole organism rates, exponentially with temperature,

and with independent mass and temperature effects (e.g., in the Arrhenius fractal supply model

(AFS) applied in the metabolic theory of ecology, MTE (Brown et al., 2004; Downs et al., 2008;

Gillooly et al., 2001)). In contrast, within species, deviations from a general 3/4 mass exponent are

common (Barneche et al., 2019; Bokma, 2004; Clarke & Johnston, 1999; Jerde et al., 2019), rates

are typically unimodally related to temperature, activation energies can vary a lot (Dell et al., 2011;

Englund et al., 2011; Pawar et al., 2016; Rall et al., 2012; Uiterwaal & DeLong, 2020) and the

effects of mass and temperature can be interactive (García García et al., 2011; Glazier, 2005;

Lindmark et al., 2018; Ohlberger et al., 2012; Xie & Sun, 1990) (but see Jerde *et al.* (2019)).

Extensions of the MTE include fitting multiple regression models where coefficients for mass and

temperature are estimated jointly (Downs et al., 2008), as well as fitting non-linear models that

can capture the de-activation of biological rates at higher temperatures (Dell et al., 2011; Englund

et al., 2011; Padfield et al., 2017; Schoolfield et al., 1981). To advance our understanding of the

intraspecific properties of mass- and temperature dependence of biological rates, intraspecific data

with variation in both mass and temperature are needed.

In this study, we analyzed how maximum consumption and metabolic rate of fish scale

intraspecifically with mass and temperature, and how optimum growth temperature scales with

size. We performed a systematic literature review by searching the Web of Science Core Collection

to compile datasets on individual-level maximum consumption, metabolic and growth rate of fish

from experiments in which the effect of fish body mass is replicated across multiple temperatures

within species (total n=3672, with data from 20, 34 and 13 species for each rate, respectively). We

then fit hierarchical Bayesian models to estimate general intraspecific scaling parameters while

accounting for variation between species. The estimated mass dependence and temperature

sensitivity of consumption and metabolism were used to quantify average changes in net energy

gain (and hence, growth, assumed proportional to net energy gain) over temperature and body

mass. Lastly, we compared our predicted changes in optimum growth temperature over body mass

with an independent experimental dataset on optimum growth temperatures data across individuals

of different sizes within species.

Materials and methods

Data acquisition

We searched the literature for experimental studies evaluating the temperature response of

individual maximum consumption rate (feeding rate at unlimited food supply, *ad libitum*), resting,

routine and standard oxygen consumption rate as a proxy for metabolic rate (Nelson, 2016) and

growth rates across individuals of different sizes within species. We used three different searches

on the Web of Science Core Collection (see *SI Appendix*, for details). In order to estimate how

these rates depend on body size and temperature within species, we selected studies that

experimentally varied both body size and temperature (at least two temperature treatments and at

least two body masses). The average number of unique temperature treatments (temperature

rounded to nearest ℃) by species is 7.2 for growth and 4.3 for consumption (below peak

temperatures) and metabolism data. The criteria for both mass and temperature variation in the

experiments reduce the number of potential data sets, as most experimental studies use either size

or temperature treatments, not both. However, these criteria allow us to fit multiple regression

models and estimate the effects of mass and temperature jointly, and to evaluate the probability of

interactive mass- and temperature effects within species. Following common practice we excluded

larval studies, which represents a life stage exhibiting different constraints and scaling

relationships than non-larval life stages (Glazier, 2005).

Studies were included if (i) a unique experimental temperature was recorded for each trial

(±1°C), (ii) fish were provided food at *ad libitum* (consumption and growth data) or if they were

unfed (resting, standard or routine metabolic rate), and (iii) fish exhibited normal behavior during

the experiments. We used only one study per species and rate to ensure that all data within a given

species are comparable as measurements of these rates can vary between studies due to e.g.

measurement bias, differences in experimental protocols, or because different populations were

studied (Armstrong & Hawkins, 2008; Jerde et al., 2019). In cases where we found more than one

study for a given rate and species, we selected the most suitable study based on our pre-defined

criteria (for details, see *SI Appendix*). We ensured that the experiments were conducted at

ecologically relevant temperatures (*SI Appendix,* Figs. S1, S3). A more detailed description of the

search protocol, data selection, acquisition, quality control, collation of additional information and

standardizing of rates to common units can be found in *SI Appendix*.

We compiled four datasets: maximum consumption rate, metabolic rate, growth rate and the

optimum growth temperature for each combination of body mass group and species. We compiled

a total of 746 measurements of maximum consumption rate (of which 666 are below peak), 2699

measurements of metabolic rate and 227 measurements of growth rate (159 below peak, 45

optimum temperatures for species-size group combinations) from published articles for each rate,

from 20, 34 and 13 species, respectively, from different taxonomic groups, habitats and lifestyles

(Table S1-S2). We requested original data from all corresponding authors of each article. In cases

where we did not hear from the corresponding author, we extracted data from tables or figures

using Web Plot Digitizer (Rohatgi, 2012).

Model fitting

Model description

To each dataset, we fit hierarchical models with different combinations of species-varying

coefficients, meaning they are estimated with shrinkage. This reduces the influence of outliers

which could occur in species with small samples sizes (Gelman & Hill, 2007; Harrison et al.,

2018). The general form of the model is:

$$y_{ij} \sim N(\mu_{ij}, \sigma) \tag{1}$$

$$\mu_{ij} = \beta_{0j} + \sum_{p=1}^{n}(\beta_p \times x_{ip}) \tag{2}$$

$$\beta_{0j} \sim N(\mu_{\beta_0}, \sigma_{\beta_0}) \tag{3}$$

where y_{ij} is the ith observation for species j for rate y, β_{0j} is a species-varying intercept, x_{ip} is a

predictor and β_p is its coefficient, with $p = 1, \ldots, n$, where n is the number of predictors

considered in the model (mass, temperature, and their interaction). Predictors are mean centered

to improve interpretability (Schielzeth, 2010). Species-level intercepts follow a normal distribution

with hyperparameters μ_{β_0} (global intercept) and σ_{β_0} (between-species standard deviation). For

most models we also allow the coefficient β_p to vary between species, such that β_p becomes β_{pj}

and x_{ip} x_{ijp}, where $\beta_{pj} \sim N(\mu_{\beta_p}, \sigma_{\beta_p})$. For each dataset, we evaluate multiple combinations of

species-varying coefficients (from varying intercept to n varying coefficients). We used a mix of

flat, weakly informative, and non-informative priors. For the temperature and mass coefficients

we used the predictions from the MTE as the means of the normal prior distributions (Brown et

al., 2004), but with large standard deviations (see *SI Appendix*, Table S3). Below we describe how

the model in Eqns. 1-3 is applied to each data set.

Mass- and temperature dependence of maximum consumption and metabolic rate below peak

temperatures

Peak temperature (optimum in the case of growth) refers to the temperature at which the rate was

maximized, by size group. For data below peak temperatures, we assumed that maximum

consumption and metabolic rate scale allometrically (as a power function of the form $I = i_0 M^{b_0}$)

with mass, and exponentially with temperature. Hence, after log-log (natural log) transformation

of mass and the rate, and temperature in Arrhenius temperature ($1/kT$ in unit eV^{-1}, where k is

Boltzmann's constant [8.62×10^{-5} eV K^{-1}]), the relationship between the rate and its predictors

becomes linear. This is similar to the MTE, except that we estimate all coefficients instead of

correcting rates, and allow intercepts and slopes to vary across species.

When applied to Eqns. 1-3, y_{ij} is the ith observation for species j of the natural log of the rate

(maximum consumption or metabolic rate), and the predictors are m_{ij} (natural log of body mass),

$t_{A,ij}$ (Arrhenius temperature, $1/kT$ in unit eV^{-1}), both of which were mean-centered, and their

interaction. Body mass is in g, consumption rate in g day^{-1} and metabolic rate in mg O_2 h^{-1}. We

use resting or routine metabolism (mean oxygen uptake of a resting unfed fish only showing some

spontaneous activity) and standard metabolism (resting unfed and no activity, usually inferred

from extrapolation or from low quantiles of routine metabolism, e.g. lowest 10% of measurements)

to represent metabolic rate (Beamish, 1964; Ohlberger et al., 2007). Routine and resting

metabolism constitute 58% of the data used and standard metabolism constitutes 42%. We

accounted for potential differences between these types of metabolic rate measurements by adding

two dummy coded variables, $type_r$ and $type_s$, the former taking the value 0 for standard and 1

for a routine or resting metabolic rate measurement, and vice versa for the latter variable. Thus,

for metabolism, we replace the overall intercept β_{0j} in Eqns. 2-3 with β_{0rj} and β_{0sj}. β_{0sj} is forced

to 0 for a species that has a routine or resting metabolic rate and β_{0rj} is forced to 0 for a species

with standard metabolic rate data. We assume these coefficients vary by species following normal

distributions with global means $\mu_{\beta_{0r}}$ and $\mu_{\beta_{0s}}$, and standard deviations $\sigma_{\beta_{0r}}$ and $\sigma_{\beta_{0s}}$, i.e.

$\beta_{0rj} \sim N(\mu_{\beta_{0r}}, \sigma_{\beta_{0r}})$ and $\beta_{0sj} \sim N(\mu_{\beta_{0s}}, \sigma_{\beta_{0s}})$.

Mass- and temperature dependence of consumption including beyond peak temperatures

Over a large temperature range, many biological rates are unimodal. We identified such tendencies

in 10 out of 20 species in the consumption data set. To characterize the decline in consumption

rate beyond peak temperature, we fit a mixed-effects version of the Sharpe Schoolfield equation

(Schoolfield et al., 1981) as expressed in (Padfield et al., 2020), to equations 1-2 with y_{ij} as

rescaled consumption rates (C). Specifically, we model μ_{ij} in Eq. 1 with the Sharpe-Schoolfield

equation:

230
$$\mu_{ij} = \frac{C_{0_j}(T_C)e^{E_j\left(\frac{1}{kT_C}-\frac{1}{kT}\right)}}{1+e^{E_h\left(\frac{1}{kT_h}-\frac{1}{kT}\right)}} \tag{4}$$

231
$$E_j \sim N(\mu_E, \sigma_E) \tag{5}$$

232
$$C_{0_j} \sim N(\mu_{C_0}, \sigma_{C_0}) \tag{6}$$

where $C_{0_j}(T_C)$ is the rate at a reference temperature T_C in Kelvin [K] (here set to 263.15, note it is

on centered scale), E_j [eV] is the activation energy, E_h [eV] characterizes the decline in the rate

past the peak temperature and T_h [K] is the temperature at which the rate is reduced to half (of the

rate in the absence of deactivation) due to high temperatures. We assume E_j and C_{0_j} vary across

species according to a normal distribution with means μ_E and μ_{C_0}, and standard deviations σ_E and

σ_{C_0} (Eq. 5-6). Prior to rescaling maximum consumption (in unit g day^{-1}) by dividing $C_{i,j}$ with the

mean within species $\bar{C}_j$, we isolate the effect of mass by dividing consumption with $m^{0.63}$, which

is the estimated allometric relationship from the log-linear model. Temperature, T, is centered by

subtracting the temperature at peak consumption. This was estimated separately for each species

using the Sharpe-Schoolfield equation but without group-varying coefficients in a frequentist

framework (see '*Parameter estimation*'). The rescaling is done to control for differences in the

optimum temperature between species, which if not accounted for obscures the average

relationship between consumption and temperature among species.

Mass-dependence of optimum growth temperature

To evaluate how the optimum temperature ($t_{opt,ij}$, in degrees Celsius) for individual growth

depends on body mass, we fit Eqns. 1-3 with y_{ij} as the mean-centered optimum growth

temperature within species ($t_{opt,ij} = T_{opt,ij} - \bar{T}_{opt,j}$), to account for species being adapted to

different thermal regimes. m_{ij}, the predictor variable for this model, is the natural log of the ratio

between mass and mass at maturation acquired from FishBase (Froese & Pauly, 2019), within

species: $m_{ij} = \ln(M_{ij}/M_{mat,j}) - \overline{\ln(M_{ij}/M_{mat,j})}$. This rescaling is done because we are

interested in examining relationships within species over "ontogenetic size", and because we do

not want to confound that effect with any relationship that might occur across species that have

different asymptotic sizes. We consider both the intercept and the effect of mass to potentially vary

between species.

Parameter estimation

We fit the models in a Bayesian framework, using R version 4.0.2 (R Core Team, 2020) and JAGS

(Plummer, 2003) through the R-package '*rjags*' (Plummer, 2019). We used 3 Markov chains with

5000 iterations for adaptation, followed by 15000 iterations burn-in and 15000 iterations sampling

where every 5^{th} iteration saved. Model convergence was assessed by visually inspecting trace plots

and potential scale reduction factors ($\hat{R}$) (*SI Appendix*). $\hat{R}$ compares chain variance with the pooled

variance, and values <1.1 suggest all three chains converged to a common distribution (Gelman et

al., 2003). We relied heavily on the R packages within '*tidyverse*' (Wickham et al., 2019) for data

processing, as well as '*ggmcmc*' (Fernández-i-Marín, 2016), '*mcmcviz*' (Youngflesh, 2018) and

'*bayesplot*' (Gabry et al., 2019) for visualization. Frequentist single-species Sharpe-Schoolfield

models were fitted using the packages '*rTPC*' (Padfield & O'Sullivan, 2020) and '*nls.multstart*'

(Padfield & Matheson, 2020)

Model comparison

We compared the parsimony of models with different hierarchical structures, and with or without

mass-temperature interactions, using the Watanabe-Akaike information criterion (WAIC) (Vehtari

et al., 2017; Watanabe, 2013), which is based on the posterior predictive distribution. We report

WAIC for each model descried above (Table S4-S5), and examine models with ΔWAIC values <

2, where ΔWAIC is each models difference to the lowest WAIC across models, in line with other

studies (Olmos et al., 2019).

Net energy gain

The effect of temperature and mass dependence of maximum consumption and metabolism

(proportional to biomass gain and losses, respectively) (Essington et al., 2001; Kitchell et al., 1977;

Ursin, 1967) on growth is illustrated by visualizing the net energy gain. The model for the net

energy gain (growth) can be viewed as an empirical temperature-dependent Pütter-type model.

Pütter-type models are the simplest growth models based on a dynamic energy budget, and make

strong assumptions about mass-scaling of key life-history and physiological processes (e.g.,

maturation and assimilation). However, Pütter-type models are among the most commonly applied

growth models in ecology and fisheries, they tend to fit data reasonably well (Marshall & White,

2019), and are suitable for illustrating the consequences of non-linear consumption rates due to

their simplicity (in contrast to more complex and parameter-rich dynamic energy budget models,

e.g. (Cuenco et al., 1985; Kitchell et al., 1977)). A Pütter model is the result of two antagonistic

allometric processes, biomass gains and biomass losses:

$$\frac{dM}{dt} = H(T)M^a - K(T)M^b \tag{7}$$

where M is body mass and T is temperature, H and K the allometric constants and a and b the

exponents of the processes underlying gains and losses, respectively. We convert metabolism from

oxygen consumption [mg O_2 h^{-1} day^{-1}] to g day^{-1} by assuming 1 kcal = 295 mg O_2 (based on

an oxycaloric coefficient of 14.2 J mg^{-1} O_2) (Hepher, 1988), 1 kcal = 4184 J and an energy content

of 5.6 kJ g^{-1} (wet weight) (Rijnsdorp & Ibelings, 1989). Hence, to convert our estimated intercept

for routine/resting metabolic rates from unit $\log$(mg O_2 h^{-1} day^{-1}) to g day^{-1}, we multiply

$\exp(\mu_{\beta_0})$ with 0.0608063. Consumption rate is already in unit g day^{-1}. Consumption and

metabolic rates are calculated for two sizes (5 and 1000 g, which roughly correspond to the 25[th]

percentile of both datasets and the maximum mass in the consumption data, respectively), using

the global allometric relationships found in the log-log models fit to sub-peak temperatures. These

allometric functions are further scaled with the temperature correction factors r_c for consumption

and r_m for metabolism. r_c is based on the Sharpe-Schoolfield model and r_m is given by the

temperature dependence of metabolic rate from the log-linear model. Because r_c and r_m are fitted

to data on different scales, we divide these functions by their maximum. Lastly, we rescale the

product between the allometric functions and r_c and r_m such that the rate at 19°C (mean temperature

in both data sets) equals the temperature-independent rate.

Results

We identified that within species of fish, metabolic rates increase faster with body mass than

maximum consumption rates, and neither of these rates conform to the commonly predicted 3/4

scaling with body mass (Fig. 1). We also quantified the unimodal relationship of consumption rate

over the full temperature range (Fig. 2). Combined, these scaling relationships lead to the

prediction, based on Pütter-type growth models, that optimum growth temperature declines with

body size (Fig. 3). The prediction of declining optimum growth temperatures with size was

confirmed by our analysis of independent experimental growth rate data. We find that within

species the optimum growth temperature declines with body size by 0.31°C per unit increase in the

natural log of relative body mass (Fig. 4). Below we present the underlying results in more detail.

We found that the average intraspecific mass exponent for consumption rate is smaller (0.63

[0.55, 0.71]) than that for metabolic rate (0.79 [0.74, 0.84]), based on the non-overlapping

Bayesian 95% credible intervals (Fig. 1). It is also probable that the scaling exponents differ from

3/4 (that is predicted by the MTE), because >99% of the posterior distribution of the mass exponent

of maximum consumption is below 3/4, and 95% of the posterior distribution of the mass exponent

of metabolic rate is above 3/4. Activation energies of maximum consumption rate and metabolism

are both similar (0.69 [0.54, 0.85] and 0.62 [0.57, 0.67] respectively; Fig. 1) and largely fall within

the prediction from the MTE (0.6-0.7 eV) (Brown et al., 2004). The global intraspecific intercept

for maximum consumption rate is estimated to be -0.34 [-0.85, 0.17], and for routine and resting

metabolic rate it is estimated to be 1.85 [1.68, 2.04], and for standard metabolic rate it is 1.29 [0.97,

1.61] (*SI Appendix,* Fig. S6). Models where all coefficients varied by species were favored in terms

of WAIC (M5 and M1, for consumption and metabolism, respectively) (*SI Appendix*, Table S4).

We found statistical support for a species-varying mass and temperature interaction for metabolic

rate; 98% of the posterior distribution of the global interaction coefficient μ_{β_3} is above 0 (*SI*

Appendix, Fig. S4). The estimated coefficient is 0.0018 [0.015, 0.037] on the Arrhenius

temperature scale, which corresponds to a decline in the mass scaling exponent of metabolic rate

by 0.0026 °C^{-1}. The selected model for maximum consumption rate did not include an interaction

term between mass and temperature (M5).

We estimated the parameters of the Sharpe-Schoolfield equation (Eq. 4) for temperature-

dependence of consumption including data beyond peak temperature as: activation energy, $E_j =$

0.73 [0.54, 0.94], rate at reference temperature, C_{0_j} = 0.79 [0.58, 0.99], temperature at which the

rate is reduced to half (of the rate in the absence of deactivation) due to high temperatures, T_h =

0.75 [-0.86, 2.37], and the rate of the decline past the peak, E_h = 1.89 [1.68, 2.1]. This shows that

the relationship between consumption rate and temperature is unimodal and asymmetric, where

the decline in consumption rate at high temperatures is steeper than the increase at low

temperatures (Fig. 2).

The above results provide empirical support for the two criteria outlined in Morita et al., (2010)

that result in declining optimum temperatures with size, i.e. (i) smaller whole organism mass

exponent for consumption than metabolism (Fig. 1) and (ii) that growth reaches an optimum over

temperature. In our case, the second criterion is met because consumption reaches a peak over

temperature (Fig. 2) (in contrast to Morita *et al*. (2010), who assumed consumption to be linearly

related to temperature, based on data from Atkinson (1994)). We illustrate the consequence of

these findings in Fig. 3, which shows that the optimum temperature for net energy gain is reached

at a lower temperature for a larger fish because of the difference in mass exponents of consumption

and metabolism and because consumption is unimodally related to temperature. Assuming growth

is proportional to net energy gain, this predicts that optimum growth temperature declines with

body size.

Using independent data from growth trials across a range of body sizes and temperatures, we

also find statistical support for a decline in optimum growth temperature with body mass within

species, because 92% of the posterior density of the global slope estimate (μ_{β_1}) is below 0. The

models with and without species-varying slopes were indistinguishable in terms of WAIC (*SI*

Appendix, Table S5), and we present the results for the species-varying intercept and slope model,

due to slightly better model diagnostics (*SI Appendix*, Fig. S19-S22). The global relationship is

given by the model: $T_{opt} = -0.074 - 0.31 \times m$, where m is the natural log of the rescaled body

mass, calculated as the species-specific ratio of mass to maturation mass.

Discussion

In this study, we systematically analyzed the intraspecific scaling of consumption, metabolism and

growth with body mass and temperature. We found strong evidence for declining optimum growth

temperatures as individuals grow in size, based on two independent approaches. First, we find

differences in the intraspecific mass-scaling of consumption and metabolism, and a unimodal

temperature dependence of consumption, which lead to predicted declines in optimum temperature

for net energy gain (and hence body growth) with size. Second, we confirm this prediction using

intraspecific growth rate. Our analysis thus demonstrates the importance of understanding

intraspecific scaling relationships when predicting responses of fish populations to climate

warming.

That warming increases growth and development rates but reduces maximum or adult size is

well known from experimental studies and is referred to as the temperature-size rule (TSR). Yet,

the mechanisms underlying the TSR remain poorly understood. Pütter-type growth models,

including the von Bertalanffy growth equation (VBGE), predict that the asymptotic size declines

with warming if the ratio of the coefficients for energy gains and losses (H/K in Eq. 7) (Pauly &

Cheung, 2018a) declines with temperature. However, the assumptions underlying the VBGE were

recently questioned because of the lack of empirical basis for the scaling exponents and the effects

of those on the predicted effects of temperature on asymptotic size (Lefevre et al., 2018; Marshall

& White, 2019). Specifically, the allometric exponent of energy gains (a) is assumed to be smaller

than that of energetic costs (b) (Eq. 7). This is based on the assumption that anabolism scales with

the same power as surfaces to volumes ($a = 2/3$) and catabolism, or maintenance metabolism, is

proportional to body mass ($b = 1$) (Pauly & Cheung, 2018b; von Bertalanffy, 1957). In contrast,

maintenance costs are commonly thought to instead be proportional to standard metabolic rate,

which in turn often is proportional to intake rates at the interspecific level (Brown et al., 2004;

Marshall & White, 2019). This leads to $a \approx b$, resulting in unrealistic growth trajectories and

temperature dependencies of growth dynamics in Pütter models (Lefevre et al., 2018; Marshall &

White, 2019). However, similar to how the existence of large fishes in tropical waters does not

invalidate the hypothesis that old individuals of large-bodied fish may reach smaller sizes with

warming, interspecific scaling parameters cannot reject or support these model predictions on

growth *within* species. We show that the average intraspecific whole-organism mass scaling

exponent of metabolism is larger than that of maximum consumption, i.e., the inequality $a < b$

holds at the intraspecific level. By contrast, Pawar *et al.* (2012) estimated larger mass exponents

for consumption than metabolic rate (0.84 and 1.04 in 2D and 3D foraging) from interspecific data,

which reveals the importance of parameterizing processes occurring over ontogeny with

intraspecific rather than interspecific data. When accounting for the smaller intraspecific mass

exponent of consumption, and the unimodal thermal response of consumption, the thermal

response of net energy gain is characterized by the optimum temperature being a function of body

size (Morita et al., 2010). Therefore, empirically derived intraspecific parameterizations of simple

growth models result in predictions in line with the TSR, in this case via declines in optimum

growth temperatures over ontogeny rather than declines in asymptotic sizes.

Declines in optimum growth temperatures over ontogeny as a mechanism for TSR-like growth

dynamics do not rely on the assumption that the ratio of the coefficients for energy gains and losses

declines with temperature. In fact, we find that when using data from sub-peak temperatures only,

the predicted average intraspecific activation energy of metabolism and consumption do not differ

substantially, which implies there is no clear loss or gain of energetic efficiency with warming

within species below temperature optima. This is in contrast to other studies, e.g. Lemoine &

Burkepile (2012) and Rall *et al.* (2010). However, it is in line with the finding that growth rates

increase with temperature (e.g. Angilletta & Dunham, 2003), which is difficult to reconcile from

a bioenergetics perspective if warming always reduced net energy gain. Our analysis instead

suggests that the mismatch between gains and losses occurs when accounting for unimodal

consumption rates over temperature. The match, or mismatch, between the temperature

dependence of feeding vs. metabolic rates is a central question in ecology that extends from

experiments to meta-analyses to food web models (Fussmann et al., 2014; Lemoine & Burkepile,

2012; Lindmark et al., 2019; Rall et al., 2010; Vasseur & McCann, 2005). Our study highlights

the importance of accounting for non-linear thermal responses for two main reasons. First, the

thermal response of net energy gain reaches a peak at temperatures below the peak for

consumption. Secondly, as initial warming commonly leads to increased growth rates, the effect

of warming on growth rates itself depends on temperature, and growth should therefore not be

assumed to be monotonically related to temperature.

Life-stage dependent optimum growth temperatures have previously been suggested as a

component of the TSR (Ohlberger, 2013). Although previous studies have found declines in

optimum growth temperatures with body size in some species of fishes and other aquatic

ectotherms (Björnsson et al., 2007; Handeland et al., 2008; Panov & McQueen, 1998; Steinarsson

& Imsland, 2003; Wyban et al., 1995), others have not (Brett et al., 1969; Elliott & Hurley, 1995).

Using systematically collated growth data from experiments with variation in both size and

temperature treatments (13 species), we find that for an average fish, the optimum growth

temperature declines as it grows in size. This finding emerges despite the small range of body sizes

used in the experiments (only 10% of observations are larger than 50% of maturation size) (*SI*

Appendix, Fig. S2). Individuals of such small relative size likely invest little energy in

reproduction, which suggests that physiological constraints at warmer temperatures contribute to

reduced growth performance of large compared to small fish, in addition to increasing investment

into reproduction (Barneche et al., 2018).

Translating results from experimental data to natural systems is challenging because maximal

feeding rates, unlimited food supply, lack of predation, and constant temperatures do not reflect

natural conditions, yet they affect growth rates (Brett et al., 1969; Huey & Kingsolver, 2019;

Lorenzen, 1996). In addition, total metabolic costs in the wild also include additional costs for

foraging and predator avoidance. It is, however, typically found that standard metabolic rate is

proportional to routine metabolic rate (exhibit the same mass-scaling relationships (Kitchell et al.,

1977; Messmer et al., 2017)). This assumption is also commonly applied to the relationship

between maximum consumption rates and consumption rates in the wild (i.e., they are related via

a constant), which has some empirical support (Neuenfeldt et al., 2020). However, the assumption

of proportionality has not been tested thoroughly, despite it being identified as an important area

for research over 40 years ago (Kitchell et al., 1977). If, for example, large individuals in wild

populations feed more efficiently than small ones (e.g., due to being less sensitive to predation),

the prediction about declining optimum growth temperature with size based on lab experiments

may be altered.

Moreover, intraspecific growth rates may not appear to be unimodally related to temperature

when measured over a temperature gradient across populations within a species (Denderen et al.,

2020), because each population can be adapted to local climate conditions and thus display

different temperature optima. However, each population likely has a thermal optimum for growth,

which differs between individuals of different size. Hence, each population might have a unimodal

relationship with temperature that differs from other populations of the same species. This

highlights the importance of understanding the time scale of environmental change in relation to

that of immediate physiological responses, acclimation, adaptation and community reorganization

for the specific prediction about climate change impacts. In natural systems, climate warming may

also result in stronger food limitation (Huey & Kingsolver, 2019; Ohlberger et al., 2011). Hence,

as optimum growth temperatures decline not only with size but also food availability (Brett, 1971;

Brett et al., 1969), and realized consumption rates often are a fraction of the maximum

consumption rate (20-70%) (Kitchell *et al.* 1977; Neuenfeldt *et al.* 2020), species may be

negatively impacted by warming even when controlled experiments show they can maintain

growth capacity at these temperatures. Supporting this point is the observation that warming

already has negative or lack of positive effects on body growth in populations living at the edge of

their physiological tolerance in terms of growth (Huss et al., 2019; Neuheimer et al., 2011).

Whether the largest fish of a population will be the first to experience negative effects of

warming, as suggested by the finding that optimum growth temperature declines with body size,

depends on the environmental temperatures they typically experience compared to smaller

conspecifics. For instance, large fish may inhabit colder temperatures compared to small fish due

to ontogenetic habitat shifts (Lloret-Lloret et al., 2020; Werner & Hall, 1988); see also Heincke's

law (Audzijonyte & Pecl, 2018; Heincke, 1913). Yet, there is already empirical evidence of the

largest individuals in natural populations being the first to suffer from negative impacts of

warming, either from increased mortality (Peralta-Maraver & Rezende, 2021; Pörtner & Knust,

2007), or not being able to increase growth rates as smaller conspecifics tend to do (Huss et al.,

2019; van Dorst et al., 2019). Hence, assuming that warming affects all individuals of a population

equally is a simplification that can bias predictions of the biological impacts of climate change.

The interspecific scaling of fundamental ecological processes with body mass and temperature

has been used to predict the effects of warming on body size, size structure, and population and

community dynamics (Cheung et al., 2013; Gilbert et al., 2014; Morita et al., 2010; Vasseur &

McCann, 2005). We argue that a contributing factor to the discrepancy between mechanistic

growth models, general scaling theory, and empirical data has been the lack of data synthesis at

the intraspecific level. The approach presented here can help overcome limitations of small data

sets by borrowing information across species in a single modelling framework, while accounting

for the intraspecific scaling of rates. Accounting for the faster increase in whole-organism

metabolism than consumption with body size, the unimodal thermal response of consumption, and

resulting size-dependence of optimum growth temperatures is essential for understanding what

causes observed growth responses to global warming. Acknowledging these mechanisms is also

important for improving predictions on the consequences of warming effects on fish growth for

food web functioning, fisheries yields and global food production in warmer climates.

Acknowledgements

We thank Hiroki Yamanaka, Dennis Tomalá Solano, Vanessa Messmer, Björn Björnsson, Albert

Imsland, Tomas Árnasson, Yiping Luo, Takeshi Tomiyama and Myron Peck for generously

providing data; Magnus Huss and Ken Haste Andersen, for providing useful comments on earlier

versions of the manuscript; Daniel Padfield and Wilco Verberk for helpful discussions; Matthew

Low and Malin Aronsson for an introduction to Bayesian inference. This study was supported by

grants from the Swedish Research Council FORMAS (no. 217-2013-1315) and the Swedish

Research Council (no. 2015-03752) (both to AG).

Author contributions

507 ML conceived the study; ML, JO, AG designed research; ML performed research with input from

508 JO and AG; ML, JO, AG wrote the paper and contributed to revisions of the manuscript.

Data accessibility statement

511 All data and R code (lists of studies in literature search, data preparation, analyses, and figures)

512 have been deposited in GitHub (https://github.com/maxlindmark/scaling) and Zenodo

513 (https://doi.org/10.5281/zenodo.5806330).

References

Andersen, K. H., Beyer, J. E., & Lundberg, P. (2009). Trophic and individual efficiencies of
size-structured communities. *Proceedings of the Royal Society B: Biological Sciences*,
276(1654), 109–114. https://doi.org/10.1098/rspb.2008.0951
Angilletta, M. J., & Dunham, A. E. (2003). The temperature-size rule in ectotherms: Simple
evolutionary explanations may not be general. *The American Naturalist*, *162*, 332–342.
Armstrong, J. D., & Hawkins, L. A. (2008). Standard metabolic rate of pike, *Esox lucius*:
Variation among studies and implications for energy flow modelling. *Hydrobiologia*,
601(1), 83–90. https://doi.org/10.1007/s10750-007-9268-x
Atkinson, D. (1994). Temperature and organism size—A biological law for ectotherms? In
Advances in Ecological Research (Vol. 25, pp. 1–58). Elsevier.
https://doi.org/10.1016/S0065-2504(08)60212-3
Audzijonyte, A., Barneche, D. R., Baudron, A. R., Belmaker, J., Clark, T. D., Marshall, C. T.,
Morrongiello, J. R., & van Rijn, I. (2019). Is oxygen limitation in warming waters a valid
mechanism to explain decreased body sizes in aquatic ectotherms? *Global Ecology and*
Biogeography, *28*(2), 64–77. https://doi.org/10.1111/geb.12847
Audzijonyte, A., & Pecl, G. T. (2018). Deep impact of fisheries. *Nature Ecology & Evolution*,
2(9), 1348–1349. https://doi.org/10.1038/s41559-018-0653-9
Audzijonyte, A., Richards, S. A., Stuart-Smith, R. D., Pecl, G., Edgar, G. J., Barrett, N. S.,
Payne, N., & Blanchard, J. L. (2020). Fish body sizes change with temperature but not all

species shrink with warming. *Nature Ecology & Evolution*, *4*, 809–814. https://doi.org/10.1038/s41559-020-1171-0

Barneche, D. R., & Allen, A. P. (2018). The energetics of fish growth and how it constrains food-web trophic structure. *Ecology Letters*, *21*(6), 836–844. https://doi.org/10.1111/ele.12947

Barneche, D. R., Jahn, M., & Seebacher, F. (2019). Warming increases the cost of growth in a model vertebrate. *Functional Ecology*, *33*(7), 1256–1266. https://doi.org/10.1111/1365-2435.13348

Barneche, D. R., Robertson, D. R., White, C. R., & Marshall, D. J. (2018). Fish reproductive-energy output increases disproportionately with body size. *Science*, *360*(6389), 642–645. https://doi.org/10.1126/science.aao6868

Baudron, A. R., Needle, C. L., Rijnsdorp, A. D., & Marshall, C. T. (2014). Warming temperatures and smaller body sizes: Synchronous changes in growth of North Sea fishes. *Global Change Biology*, *20*(4), 1023–1031. https://doi.org/10.1111/gcb.12514

Beamish, F. W. H. (1964). Respiration of fishes with special emphasis on standard oxygen consumption II. Influence of weight and temperature on respiration of several species'. *Canadian Journal of Zoology/Revue Canadienne de Zoologie*, *42*(2), 177–188.

Björnsson, B., Steinarsson, A., & Árnason, T. (2007). Growth model for Atlantic cod (*Gadus morhua*): Effects of temperature and body weight on growth rate. *Aquaculture*, *271*(1–4), 216–226. https://doi.org/10.1016/j.aquaculture.2007.06.026

Blanchard, J. L., Heneghan, R. F., Everett, J. D., Trebilco, R., & Richardson, A. J. (2017). From bacteria to whales: Using functional size spectra to model marine ecosystems. *Trends in Ecology & Evolution*, *32*(3), 174–186. https://doi.org/10.1016/j.tree.2016.12.003

Bokma, F. (2004). Evidence against universal metabolic allometry. *Functional Ecology*, *18*(2), 184–187.

Brett, J. R. (1971). Energetic responses of salmon to temperature. A study of some thermal relations in the physiology and freshwater ecology of sockeye salmon (*Oncorhynchus nerka*). *Integrative and Comparative Biology*, *11*(1), 99–113. https://doi.org/10.1093/icb/11.1.99

Brett, J. R., Shelbourn, J. E., & Shoop, C. T. (1969). Growth rate and body composition of fingerling sockeye salmon, *Oncorhynchus nerka*, in relation to temperature and ration size. *Journal of the Fisheries Research Board of Canada*, *26*(9), 2363–2394. https://doi.org/10.1139/f69-230

Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., & West, G. B. (2004). Toward a metabolic theory of ecology. *Ecology*, *85*(7), 1771–1789. https://doi.org/10.1890/03-9000

Cheung, W. W. L., Sarmiento, J. L., Dunne, J., Frölicher, T. L., Lam, V. W. Y., Deng Palomares, M. L., Watson, R., & Pauly, D. (2013). Shrinking of fishes exacerbates impacts of global ocean changes on marine ecosystems. *Nature Climate Change*, *3*(3), 254–258. https://doi.org/10.1038/nclimate1691

Clarke, A., & Johnston, N. M. (1999). Scaling of metabolic rate with body mass and temperature in teleost fish. *Journal of Animal Ecology*, *68*, 893–905.

Cuenco, M. L., Stickney, R. R., & Grant, W. E. (1985). Fish bioenergetics and growth in aquaculture ponds: I. Individual fish model development. *Ecological Modelling*, *27*(3), 169–190. https://doi.org/10.1016/0304-3800(85)90001-8

Daufresne, M., Lengfellner, K., & Sommer, U. (2009). Global warming benefits the small in
aquatic ecosystems. *Proceedings of the National Academy of Sciences, USA*, *106*(31),
12788–12793. https://doi.org/10.1073/pnas.0902080106

de Roos, A. M., & Persson, L. (2001). Physiologically structured models – from versatile
technique to ecological theory. *Oikos*, *94*(1), 51–71. https://doi.org/10.1034/j.1600-
0706.2001.11313.x

Dell, A. I., Pawar, S., & Savage, V. M. (2011). Systematic variation in the temperature
dependence of physiological and ecological traits. *Proceedings of the National Academy
of Sciences*, *108*(26), 10591–10596. https://doi.org/10.1073/pnas.1015178108

Denderen, D. van, Gislason, H., Heuvel, J. van den, & Andersen, K. H. (2020). Global analysis
of fish growth rates shows weaker responses to temperature than metabolic predictions.
Global Ecology and Biogeography, *29*(12), 2203–2213.
https://doi.org/10.1111/geb.13189

Downs, C. J., Hayes, J. P., & Tracy, C. R. (2008). Scaling metabolic rate with body mass and
inverse body temperature: A test of the Arrhenius fractal supply model. *Functional
Ecology*, *22*(2), 239–244. https://doi.org/10.1111/j.1365-2435.2007.01371.x

Elliott, J. M., & Hurley, M. A. (1995). The functional relationship between body size and growth
rate in fish. *Functional Ecology*, *9*(4), 625. https://doi.org/10.2307/2390153

Englund, G., Öhlund, G., Hein, C. L., & Diehl, S. (2011). Temperature dependence of the
functional response. *Ecology Letters*, *14*(9), 914–921. https://doi.org/10.1111/j.1461-
0248.2011.01661.x

Essington, T. E., Kitchell, J. F., & Walters, C. J. (2001). The von Bertalanffy growth function,
bioenergetics, and the consumption rates of fish. *Canadian Journal of Fisheries and
Aquatic Sciences*, *58*(11), 2129–2138. https://doi.org/10.1139/cjfas-58-11-2129

Fernández-i-Marín, X. (2016). ggmcmc: Analysis of MCMC Samples and Bayesian Inference.
Journal of Statistical Software, *70*(1), 1–20. https://doi.org/10.18637/jss.v070.i09

Forster, J., Hirst, A. G., & Atkinson, D. (2012). Warming-induced reductions in body size are
greater in aquatic than terrestrial species. *Proceedings of the National Academy of
Sciences*, *109*(47), 19310–19314. https://doi.org/10.1073/pnas.1210460109

Froese, R., & Pauly, D. (2019). *Editors. FishBase. World Wide Web electronic publication.
Www.fishbase.org, (12/2019)*.

Fussmann, K. E., Schwarzmüller, F., Brose, U., Jousset, A., & Rall, B. C. (2014). Ecological
stability in response to warming. *Nature Climate Change*, *4*(3), 206–210.
https://doi.org/10.1038/nclimate2134

Gabry, J., Simpson, D., Vehtari, A., Betancourt, M., & Gelman, A. (2019). Visualization in
Bayesian workflow. *Journal of the Royal Statistical Society: Series A (Statistics in
Society)*, *182*(2), 389–402. https://doi.org/10.1111/rssa.12378

García García, B., Cerezo Valverde, J., Aguado-Giménez, F., García García, J., & Hernández,
M. D. (2011). Effect of the interaction between body weight and temperature on growth
and maximum daily food intake in sharpsnout sea bream (*Diplodus puntazzo*).
Aquaculture International, *19*(1), 131–141. https://doi.org/10.1007/s10499-010-9347-2

Gardner, J. L., Peters, A., Kearney, M. R., Joseph, L., & Heinsohn, R. (2011). Declining body
size: A third universal response to warming? *Trends in Ecology & Evolution*, *26*(6), 285–
291. https://doi.org/10.1016/j.tree.2011.03.005

Gelman, A., Carlin, J., Stern, H., & Rubin, D. (2003). *Bayesian Data Analysis. 2nd edition*.
Chapman and Hall/CRC.

Gelman, A., & Hill, J. (2007). *Data Analysis Using Regression and Multilevel/Hierarchical Models*. Cambridge University Press.

Gilbert, B., Tunney, T. D., McCann, K. S., DeLong, J. P., Vasseur, D. A., Savage, V. M., Shurin, J. B., Dell, A. I., Barton, B. T., Harley, C. D., Kharouba, H. M., Kratina, P., Blanchard, J. L., Clements, C., Winder, M., Greig, H. S., & O'Connor, M. I. (2014). A bioenergetic framework for the temperature dependence of trophic interactions. *Ecology Letters*, *17*(8), 902–914. https://doi.org/10.1111/ele.12307

Gillooly, J. F., Brown, J. H., West, G. B., Savage, V. M., & Charnov, E. L. (2001). Effects of size and temperature on metabolic rate. *Science*, *293.5538*, 2248-2251.

Glazier, D. S. (2005). Beyond the "3/4-power law": Variation in the intra- and interspecific scaling of metabolic rate in animals. *Biological Reviews of the Cambridge Philosophical Society*, *80*(4), 611–662. https://doi.org/10.1017/S1464793105006834

Handeland, S. O., Imsland, A. K., & Stefansson, S. O. (2008). The effect of temperature and fish size on growth, feed intake, food conversion efficiency and stomach evacuation rate of Atlantic salmon post-smolts. *Aquaculture*, *283*(1), 36–42. https://doi.org/10.1016/j.aquaculture.2008.06.042

Harrison, X. A., Donaldson, L., Correa-Cano, M. E., Evans, J., Fisher, D. N., Goodwin, C. E. D., Robinson, B. S., Hodgson, D. J., & Inger, R. (2018). A brief introduction to mixed effects modelling and multi-model inference in ecology. *PeerJ*, *6*, e4794. https://doi.org/10.7717/peerj.4794

Hartvig, M., Andersen, K. H., & Beyer, J. E. (2011). Food web framework for size-structured populations. *Journal of Theoretical Biology*, *272*(1), 113–122. https://doi.org/10.1016/j.jtbi.2010.12.006

Heincke, F. (1913). Rapports et procès-verbaux des reunions. *Conseil International Pour l'Exploration de La Mer*, *16*, 1–70.

Hepher, B. (1988). *Nutrition of Pond Fishes*. Cambridge University Press.

Horne, C. R., Hirst, Andrew. G., & Atkinson, D. (2015). Temperature-size responses match latitudinal-size clines in arthropods, revealing critical differences between aquatic and terrestrial species. *Ecology Letters*, *18*(4), 327–335. https://doi.org/10.1111/ele.12413

Huey, R. B., & Kingsolver, J. G. (2019). Climate warming, resource availability, and the metabolic meltdown of ectotherms. *The American Naturalist*, *194*(6), E140–E150. https://doi.org/10.1086/705679

Huss, M., Lindmark, M., Jacobson, P., van Dorst, R. M., & Gårdmark, A. (2019). Experimental evidence of gradual size-dependent shifts in body size and growth of fish in response to warming. *Global Change Biology*, *25*(7), 2285–2295. https://doi.org/10.1111/gcb.14637

Ikpewe, I. E., Baudron, A. R., Ponchon, A., & Fernandes, P. G. (2020). Bigger juveniles and smaller adults: Changes in fish size correlate with warming seas. *Journal of Applied Ecology*, *Early View*. https://doi.org/10.1111/1365-2664.13807

Jerde, C. L., Kraskura, K., Eliason, E. J., Csik, S. R., Stier, A. C., & Taper, M. L. (2019). Strong Evidence for an Intraspecific Metabolic Scaling Coefficient Near 0.89 in Fish. *Frontiers in Physiology*, *10*, 1166. https://doi.org/10.3389/fphys.2019.01166

Jobling, M. (1997). Temperature and growth: Modulation of growth rate via temperature change. In C. M. Wood & D. G. McDonald (Eds.), *Global Warming: Implications for Freshwater and Marine Fish* (Vol. 61, pp. 225–254). Cambridge University Press.

Kitchell, J. F., Stewart, D. J., & Weininger, D. (1977). Applications of a bioenergetics model to yellow perch (*Perca flavescens*) and walleye (*Stizostedion vitreum vitreum*). *Journal of the Fisheries Board of Canada*, *34*(10), 1922–1935.

Kooijman, S. A. L. M. (1993). *Dynamic energy budgets in biological systems*. Cambridge University Press.

Lefevre, S., McKenzie, D. J., & Nilsson, G. E. (2018). In modelling effects of global warming, invalid assumptions lead to unrealistic projections. *Global Change Biology*, *24*(2), 553–556. https://doi.org/10.1111/gcb.13978

Lemoine, N. P., & Burkepile, D. E. (2012). Temperature-induced mismatches between consumption and metabolism reduce consumer fitness. *Ecology*, *93*(11), 2483–2489. https://doi.org/10.1890/12-0375.1

Lindmark, M., Huss, M., Ohlberger, J., & Gårdmark, A. (2018). Temperature-dependent body size effects determine population responses to climate warming. *Ecology Letters*, *21*(2), 181–189. https://doi.org/10.1111/ele.12880

Lindmark, M., Ohlberger, J., Huss, M., & Gårdmark, A. (2019). Size-based ecological interactions drive food web responses to climate warming. *Ecology Letters*, *22*(5), 778–786. https://doi.org/10.1111/ele.13235

Lloret-Lloret, E., Navarro, J., Giménez, J., López, N., Albo-Puigserver, M., Pennino, M. G., & Coll, M. (2020). The seasonal distribution of a highly commercial fish is related to ontogenetic changes in its feeding strategy. *Frontiers in Marine Science*, *7*. https://doi.org/10.3389/fmars.2020.566686

Lorenzen, K. (1996). The relationship between body weight and natural mortality in juvenile and adult fish: A comparison of natural ecosystems and aquaculture. *Journal of Fish Biology*, *49*(4), 627–642. https://doi.org/10.1111/j.1095-8649.1996.tb00060.x

Marshall, D. J., & White, C. R. (2019). Have we outgrown the existing models of growth? *Trends in Ecology & Evolution*, *34*(2), 102–111. https://doi.org/10.1016/j.tree.2018.10.005

Maury, O., & Poggiale, J.-C. (2013). From individuals to populations to communities: A dynamic energy budget model of marine ecosystem size-spectrum including life history diversity. *Journal of Theoretical Biology*, *324*, 52–71. https://doi.org/10.1016/j.jtbi.2013.01.018

Messmer, V., Pratchett, M. S., Hoey, A. S., Tobin, A. J., Coker, D. J., Cooke, S. J., & Clark, T. D. (2017). Global warming may disproportionately affect larger adults in a predatory coral reef fish. *Global Change Biology*, *23*(6), 2230–2240.

Morita, K., Fukuwaka, M., Tanimata, N., & Yamamura, O. (2010). Size-dependent thermal preferences in a pelagic fish. *Oikos*, *119*(8), 1265–1272. https://doi.org/10.1111/j.1600-0706.2009.18125.x

Nelson, J. A. (2016). Oxygen consumption rate v. rate of energy utilization of fishes: A comparison and brief history of the two measurements. *Journal of Fish Biology*, *88*(1), 10–25. https://doi.org/10.1111/jfb.12824

Neubauer, P., & Andersen, K. H. (2019). Thermal performance of fish is explained by an interplay between physiology, behaviour and ecology. *Conservation Physiology*, *7*(1). https://doi.org/10.1093/conphys/coz025

Neuenfeldt, S., Bartolino, V., Orio, A., Andersen, K. H., Andersen, N. G., Niiranen, S., Bergström, U., Ustups, D., Kulatska, N., & Casini, M. (2020). Feeding and growth of

Atlantic cod (*Gadus morhua* L.) in the eastern Baltic Sea under environmental change. *ICES Journal of Marine Science*, *77*(2), 624–632. https://doi.org/10.1093/icesjms/fsz224

Neuheimer, A. B., Thresher, R. E., Lyle, J. M., & Semmens, J. M. (2011). Tolerance limit for fish growth exceeded by warming waters. *Nature Climate Change*, *1*(2), 110–113. https://doi.org/10.1038/nclimate1084

Ohlberger, J. (2013). Climate warming and ectotherm body size – from individual physiology to community ecology. *Functional Ecology*, *27*(4), 991–1001. https://doi.org/10.1111/1365-2435.12098

Ohlberger, J., Edeline, E., Vollestad, L. A., Stenseth, N. C., & Claessen, D. (2011). Temperature-driven regime shifts in the dynamics of size-structured populations. *The American Naturalist*, *177*(2), 211–223. https://doi.org/10.1086/657925

Ohlberger, J., Mehner, Thomas., Staaks, Georg., & Hölker, Franz. (2012). Intraspecific temperature dependence of the scaling of metabolic rate with body mass in fishes and its ecological implications. *Oikos*, *121*(2), 245–251. https://doi.org/10.1111/j.1600-0706.2011.19882.x

Ohlberger, J., Staaks, G., & Hölker, F. (2007). Effects of temperature, swimming speed and body mass on standard and active metabolic rate in vendace (*Coregonus albula*). *Journal of Comparative Physiology, B*, *177*(8), 905–916.

Olmos, M., Payne, M. R., Nevoux, M., Prévost, E., Chaput, G., Pontavice, H. D., Guitton, J., Sheehan, T., Mills, K., & Rivot, E. (2019). Spatial synchrony in the response of a long range migratory species (*Salmo salar*) to climate change in the North Atlantic Ocean. *Global Change Biology*, *26*(3), 1319–1337. https://doi.org/10.1111/gcb.14913

Padfield, D., Castledine, M., & Buckling, A. (2020). Temperature-dependent changes to host–parasite interactions alter the thermal performance of a bacterial host. *The ISME Journal*, *14*(2), 389–398. https://doi.org/10.1038/s41396-019-0526-5

Padfield, D., Lowe, C., Buckling, A., Ffrench-Constant, R., Jennings, S., Shelley, F., Ólafsson, J. S., & Yvon-Durocher, G. (2017). Metabolic compensation constrains the temperature dependence of gross primary production. *Ecology Letters*, *20*(10), 1250–1260. https://doi.org/10.1111/ele.12820

Padfield, D., & Matheson, G. (2020). *nls.multstart: Robust Non-Linear regression using AIC Scores* (R package version 1.2.0.) [Computer software]. https://CRAN.R-project.org/package=nls.multstart

Padfield, D., & O'Sullivan, H. (2020). *RTPC: Functions for fitting thermal performance curves* (R package version 1.0.0.) [Computer software]. https://github.com/padpadpadpad/rTPC

Panov, V. E., & McQueen, D. J. (1998). *Effects of temperature on individual growth rate and body size of a freshwater amphipod*. *76*, 1107–1116.

Pauly, D. (2021). The gill-oxygen limitation theory (GOLT) and its critics. *Science Advances*, *7*(2), eabc6050. https://doi.org/10.1126/sciadv.abc6050

Pauly, D., & Cheung, W. W. L. (2018a). Sound physiological knowledge and principles in modeling shrinking of fishes under climate change. *Global Change Biology*, *24*(1), e15–e26. https://doi.org/10.1111/gcb.13831

Pauly, D., & Cheung, W. W. L. (2018b). On confusing cause and effect in the oxygen limitation of fish. *Global Change Biology*, *24*(11), e743–e744. https://doi.org/10.1111/gcb.14383

Pawar, S., Dell, A. I., Savage, V. M., & Knies, J. L. (2016). Real versus Artificial Variation in the Thermal Sensitivity of Biological Traits. *The American Naturalist*, *187*(2), E41–E52. https://doi.org/10.1086/684590

Pawar, S., Dell, A. I., & Van M. Savage. (2012). Dimensionality of consumer search space drives trophic interaction strengths. *Nature*, *486*(7404), 485–489. https://doi.org/10.1038/nature11131

Peralta-Maraver, I., & Rezende, E. L. (2021). Heat tolerance in ectotherms scales predictably with body size. *Nature Climate Change*, *11*(1), 58–63. https://doi.org/10.1038/s41558-020-00938-y

Perrin, N. (1995). About Berrigan and Charnov's life-history puzzle. *Oikos*, 137–139.

Plummer, M. (2003). JAGS: A program for analysis of Bayesian graphical models using Gibbs sampling. *Working Papers*, 8.

Plummer, M. (2019). *Rjags* (R package version 4-10) [Computer software]. https://CRAN.R-project.org/package=rjags

Pörtner, H. O., & Knust, R. (2007). Climate change affects marine fishes through the oxygen limitation of thermal tolerance. *Science*, *315*(5808), 95–97.

Pütter, A. (1920). Studien über physiologische Ähnlichkeit VI. Wachstumsähnlichkeiten. *Pflüger's Archiv für die gesamte Physiologie des Menschen und der Tiere*, *180*(1), 298–340. https://doi.org/10.1007/BF01755094

R Core Team. (2020). *R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing*. https://www.R-project.org/

Rall, B. C., Brose, U., Hartvig, M., Kalinkat, G., Schwarzmuller, F., Vucic-Pestic, O., & Petchey, O. L. (2012). Universal temperature and body-mass scaling of feeding rates. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, *367*(1605), 2923–2934. https://doi.org/10.1098/rstb.2012.0242

Rall, B. C., Vucic-Pestic, O., Ehnes, R. B., Emmerson, M., & Brose, U. (2010). Temperature, predator–prey interaction strength and population stability. *Global Change Biology*, *16*(8), 2145–2157. https://doi.org/10.1111/j.1365-2486.2009.02124.x

Rijnsdorp, A. D., & Ibelings, B. (1989). Sexual dimorphism in the energetics of reproduction and growth of North Sea plaice, Pleuronectes platessa L. *Journal of Fish Biology*, *35*(3), 401–415. https://doi.org/10.1111/j.1095-8649.1989.tb02992.x

Rohatgi, A. (2012). *WebPlotDigitalizer: HTML5 based online tool to extract numerical data from plot images. Version 4.1. [WWW document] URL https://automeris.io/WebPlotDigitizer (accessed on January 2019)*.

Savage, V. M., Gillooly, J. F., Brown, J. H., West, G. B., & Charnov, E. L. (2004). Effects of body size and temperature on population growth. *The American Naturalist*, *163*(3), 429–441.

Schielzeth, H. (2010). Simple means to improve the interpretability of regression coefficients: Interpretation of regression coefficients. *Methods in Ecology and Evolution*, *1*(2), 103–113. https://doi.org/10.1111/j.2041-210X.2010.00012.x

Schoolfield, R. M., Sharpe, P. J. H., & Magnuson, C. E. (1981). Non-linear regression of biological temperature-dependent rate models based on absolute reaction-rate theory. *Journal of Theoretical Biology*, *88*(4), 719–731. https://doi.org/10.1016/0022-5193(81)90246-0

Steinarsson, A., & Imsland, A. K. (2003). Size dependent variation in optimum growth temperature of red abalone (*Haliotis rufescens*). *Aquaculture*, *224*(1–4), 353–362. https://doi.org/10.1016/S0044-8486(03)00241-2

Thresher, R. E., Koslow, J. A., Morison, A. K., & Smith, D. C. (2007). Depth-mediated reversal
of the effects of climate change on long-term growth rates of exploited marine fish.
Proceedings of the National Academy of Sciences, USA, *104*(18), 7461–7465.
Uiterwaal, S. F., & DeLong, J. P. (2020). Functional responses are maximized at intermediate
temperatures. *Ecology*, *101*(4), e02975. https://doi.org/10.1002/ecy.2975
Ursin, E. (1967). A Mathematical Model of Some Aspects of Fish Growth, Respiration, and
Mortality. *Journal of the Fisheries Research Board of Canada*, *24*(11), 2355–2453.
https://doi.org/10.1139/f67-190
van Dorst, R. M., Gårdmark, A., Svanbäck, R., Beier, U., Weyhenmeyer, G. A., & Huss, M.
(2019). Warmer and browner waters decrease fish biomass production. *Global Change*
Biology, *25*(4), 1395–1408. https://doi.org/10.1111/gcb.14551
van Rijn, I., Buba, Y., DeLong, J., Kiflawi, M., & Belmaker, J. (2017). Large but uneven
reduction in fish size across species in relation to changing sea temperatures. *Global*
Change Biology, *23*(9), 3667–3674. https://doi.org/10.1111/gcb.13688
Vasseur, D. A., & McCann, K. S. (2005). A mechanistic approach for modelling temperature-
dependent consumer-resource dynamics. *The American Naturalist*, *166*(2), 184–198.
Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model evaluation using leave-
one-out cross-validation and WAIC. *Statistics and Computing*, *27*(5), 1413–1432.
https://doi.org/10.1007/s11222-016-9696-4
von Bertalanffy, L. (1957). Laws in metabolism and growth. *The Quarterly Review of Biology*,
32(3), 217–231.
Wang, H.-Y., Shen, S.-F., Chen, Y.-S., Kiang, Y.-K., & Heino, M. (2020). Life histories
determine divergent population trends for fishes under climate warming. *Nature*
Communications, *11*(1), 4088. https://doi.org/10.1038/s41467-020-17937-4
Watanabe, S. (2013). A Widely Applicable Bayesian Information Criterion. *Journal of Machine*
Learning Research, *14*(Mar), 867–897.
Werner, E. E., & Hall, D. J. (1988). Ontogenetic habitat shifts in bluegill: The foraging rate-
predation risk trade-off. *Ecology*, *69*(5), 1352–1366. https://doi.org/10.2307/1941633
Wickham, H., Averick, M., Bryan, J., Chang, W., D'Agostino McGowan, L., François, R.,
Grolemund, G., & Alex, H. (2019). *Welcome to the tidyverse*. 1686.
https://doi.org/10.21105/joss.01686
Wyban, J., Walsh, W. A., & Godin, D. M. (1995). Temperature effects on growth, feeding rate
and feed conversion of the Pacific white shrimp (*Penaeus vannamei*). *Aquaculture*,
138(1), 267–279. https://doi.org/10.1016/0044-8486(95)00032-1
Xie, Xiaojun., & Sun, Ruyung. (1990). The Bioenergetics of the Southern Catfish (*Silurus*
meridionalis Chen). I. Resting Metabolic Rate as a Function of Body Weight and
Temperature. *Physiological Zoology*, *63*(6), 1181–1195.
Youngflesh, C. (2018). MCMCvis: Tools to Visualize, Manipulate, and Summarize MCMC
Output. *Journal of Open Source Software*, *3*(24), 640.
https://doi.org/10.21105/joss.00640

 Figures

Figure 1. *Natural log of maximum consumption rate (A) and metabolic rate (B) against body mass on a logarithmic x-axis. Lines and equation are global predictions on the centered data, x-axes are non-centered, (routine metabolic rate in panel B) at the average temperature in each data set (both 19°C, but note the model is fitted using mean-centered Arrhenius temperature), hence the temperature terms are omitted. Red dashed lines indicate a slope of 3/4, corresponding to the prediction from the metabolic theory of ecology. Gray bands correspond to 80% and 95% credible*

intervals. Species are grouped by color (legend not shown, n=20 for consumption and n=34 for

metabolism, respectively). C) Global and species-level effects of mass and temperature on

maximum consumption rate and metabolic rate. Horizontal lines show the posterior medians of

the global mass exponents and activation energies of maximum consumption and metabolism (μ_{β_1}

and $-\mu_{\beta_2}$ (negative since we present activation energies rather than the coefficient) in Eq. 3 for

the mass and temperature coefficients, respectively). The shaded horizontal rectangles correspond

to the posterior median ± 1.96 standard deviations. Points and triangles show the posterior

medians for each species-level coefficient (for maximum consumption rate and metabolic rate,

respectively), and the vertical bars show their 80% and 95% credible interval.

Figure 2. *Mass-corrected maximum consumption rate increases until a maximum is reached, after which it declines steeper than the initial rate of increase. Maximum consumption rates are relative to the average maximum consumption rates within species and temperature is the difference between the experimental temperature and the temperature where maximum consumption peaks (also by species). The black line shows the posterior median of predictions from the Sharpe-Schoolfield model. Gray bands correspond to 80% and 95% credible intervals. Colors indicate species.*

Figure 3. *Illustration of predicted whole-organism maximum consumption rate (green), metabolic*

rate (orange) and the difference between them ('net gain'; purple) for two body sizes (top=5g,

center=1000g) over temperature (A-B) (see 'Materials and Methods'), and the distribution of

optimum growth temperatures (C), i.e., the temperature at the peak of the net gain curve. Each

line (50 in total) is a draw from the posterior distribution of the respective mass-scaling exponent,

to illustrate uncertainty in the criteria leading to declining optimum growth temperatures with

size. The thick solid lines correspond to the posterior median estimate (as reported in the

'Results').

Figure 4. *Experimental data demonstrating optimum growth temperature declines with intraspecific body mass. The plot shows the optimum temperature within species (rescaled by subtracting the mean optimum temperature from each observation, by species) as a function of the natural log of rescaled body mass (ratio of mass to maturation mass within species). The solid line shows the global prediction (using μ_{β_0} and μ_{β_1}), and shaded areas correspond to 80% and 95% credible intervals. Colors indicate species and the area of the circle corresponds to body mass in unit g.*