A structural basis for inhibition of the complement initiator protease C1r by Lyme disease spirochetes

Running Title: The molecular interface of BBK32/C1r

Ryan J. Garrigues[1], Alexandra D. Powell Pierce[2], Michal Hammel[3],

Jon T. Skare[2, *], and Brandon L. Garcia[1,*]

[1]Department of Microbiology and Immunology, Brody School of Medicine, East Carolina University, Greenville, North Carolina, United States of America

[2]Department of Microbial Pathogenesis and Immunology, College of Medicine, Texas A&M University, Bryan/College Station, Texas, United States of America

[3]Molecular Biophysics and Integrated Bioimaging, Lawrence Berkeley National Laboratory, Berkeley, CA, USA

Author Contributions

*These authors contributed equally to this work. Correspondence to Brandon L. Garcia (garciabr18@ecu.edu) and Jon T. Skare (jskare@tamu.edu).

J.T.S and B.L.G. designed research; R.J.G., A.D.P., and M.H. performed research; R.J.G., A.D.P., M.H., J.T.S., and B.L.G. analyzed data; R.J.G., A.D.P., M.H., J.T.S, and B.L.G wrote the paper.

Abstract

Complement evasion is a hallmark of extracellular microbial pathogens such as *Borreliella burgdorferi*, the causative agent of Lyme disease. Lyme disease spirochetes express nearly a dozen outer surface lipoproteins that bind complement components and interfere with their native activities. Among these, BBK32 is unique in its selective inhibition of the classical pathway. BBK32 blocks activation of this pathway by selectively binding and inhibiting the C1r serine protease of first component of complement, C1. To understand the structural basis for BBK32-mediated C1r inhibition, we performed co-crystallography and size exclusion chromatography-coupled small angle x-ray scattering experiments, which revealed a molecular model of BBK32-C in complex with activated human C1r. Structure-guided site-directed mutagenesis was combined with surface plasmon resonance binding experiments and assays of complement function to validate the predicted molecular interface. The studies reported here, for the first time, provide a structural basis for classical pathway-specific inhibition by a human pathogen.

Introduction

The complement system consists of dozens of interacting soluble and surface-associated proteins that together form a primary arm of the innate immune system (*1–4*). Activation of complement on a microbial surface leads to opsonization, recruitment of professional phagocytes, and in many cases, direct cellular lysis. Complement activation involves a cascading series of proteolytic reactions catalyzed by specialized serine proteases that cleave circulating inert complement components into activated fragments. The cascade can be initiated in multiple ways, leading to a canonical grouping of complement activation into three pathways known as the classical, lectin, and alternative pathways. The classical pathway is initiated by a multiprotein

complex known as the first component of complement, C1 (*5–8*). C1 is composed of C1q, two

molecules of C1r, and two molecules of C1s (C1qr$_2$s$_2$). C1q serves as the pattern recognition

molecule of the classical pathway, binding to immune complexes and non-antibody ligands

resulting in zymogen activation of the attached C1r and C1s proteases. The lectin pathway is

similarly initiated by pattern recognition, whereby mannan-binding lectin (MBL), ficolins,

collectin-10 or collectin-11 bind foreign carbohydrate structures and cause activation of the MBL-

associated proteases (MASPs) (*9, 10*). In contrast, the alternative pathway does not rely on pattern

recognition but is instead continuously activated at a low level by a process known as 'tick-over'

(*11*). All three initiating pathways converge at the cleavage of the central molecule of the cascade,

complement component C3. C3 activation leads to formation of surface-bound enzyme complexes

known as convertases that drive complement amplification and terminal pathway activation,

ultimately resulting in the formation of the terminal pathway complete complex, C5b-9, also

known as the membrane attack complex (MAC) (*1, 2*).

Complement activation is a finely tuned process controlled by endogenous regulators of

complement activity (RCAs) (*12*). RCAs are critical in preventing inappropriate complement

activation on the surface of healthy host cells. Indeed, aberrant activation or overactivation of

complement causes or exacerbates a wide range of human autoimmune, inflammatory, and

neurodegenerative diseases (*13, 14*). Microbes lack RCAs and thus many human pathogens have

evolved sophisticated complement evasion systems to prevent complement-mediated attack (*15*).

A prototypical example of a bacterial complement evasion system is found in *Borreliella*

spirochetes (*16, 17*). *Borreliella* gen. nov. encompasses the etiological agents of Lyme disease

including the major genospecies *Borreliella burgdorferi, B. afzelii,* and *B. garinii* (*18*). The

Borreliella complement evasion repertoire is overlapping and is categorized by two general

mechanisms (*16*, *17*). The first mechanism involves functional recruitment of host RCAs to the

spirochete, surface such as factor H (FH) (*19–28*). Among this group are the complement regulator

acquiring surface proteins (CRASPs) which include three distinct classes of FH-binding proteins

called CspA (CRASP-1), CspZ (CRASP-2), and three OspEF-related family paralogs known as

ErpP (CRASP-3), ErpC (CRASP-4), and ErpA (CRASP-5) (*19–28*). The second general

mechanism of complement evasion by outer surface *Borreliella* proteins involves a direct mode of

inhibition. Known inhibitors from this group include *B. bavariensis* BGA66 and BGA71 (*29*), a

FH-binding independent activity for CspA (*30*), and a protein of unknown molecular identity with

CD59-like activity that each interact with components of C5b-9 and prevent MAC formation (*31*).

In addition, two *Borreliella* outer surface lipoproteins are known to directly interfere with

upstream initiation steps of complement: OspC which binds to C4b and prevents classical/lectin

pathway proconvertase formation (*32*), and BBK32 which selectively blocks classical pathway

activation (*33*).

B. burgdorferi BBK32 is an outer surface-localized lipoprotein upregulated during the

vertebrate infection phase of the *B. burgdorferi* lifecycle (*34*). We have previously shown that a

bbk32 mutant is significantly attenuated in murine models of infection due to defects in

dissemination and the inability of the borrelial cells to maintain a normal bacterial load (*35*, *36*).

BBK32 contains non-overlapping N-terminal binding sites for host glycosaminoglycans (GAGs)

and fibronectin and the contribution of BBK32 to Lyme disease pathogenesis is likely related to

its multifunctionality (*37*, *38*). Interactions by BBK32 with GAGs and fibronectin at the host

endothelium exhibit dragging and tethering, respectively, thereby promoting adherence during to

the shear force of blood flow, resulting in the hematogenous dissemination of *B. burgdorferi* via

extravasation (*39*, *40*). More recently, we identified a potent complement inhibitory activity for

the C-terminal domain of BBK32 (BBK32-C, hereafter) (*33*). BBK32-C binds directly to the C1

complex and blocks the earliest proteolytic events of the classical pathway of complement (*33*).

The classical pathway-selective inhibitory activity of BBK32 is currently unique among known

microbial complement evasion proteins.

We have previously shown that BBK32-C binds the human C1 complex by specifically

recognizing the C1r subcomponent with high-affinity (*33*). Recently we obtained a high-resolution

crystal structure of BBK32-C and mapped its binding site on human C1r to the serine protease

(SP) domain (*41*). Collectively, these studies showed that BBK32-C blocks C1r autoactivation,

as well as C1r-mediated cleavage of its partner protease in the C1 complex, C1s (*33*). However,

the structural basis and mechanism of action for BBK32-mediated C1r inhibition remains

unknown. To gain further insight into the nature of the BBK32/C1r protein-protein interaction, we

carried out a series of co-crystallography and size exclusion chromatography-coupled small angle

x-ray scattering (SEC-SAXS) experiments followed by a structure-guided mutagenesis strategy

involving biophysical, biochemical, and microbiological approaches. These investigations reveal

the structural basis for C1r inhibition by Lyme disease spirochetes, provide insight into BBK32's

C1r selectivity, and reveal a convergent inhibitory mechanism between a spirochete and a

hematophagous eukaryotic organism.

110
111 **Results**
112
The Structure of BBK32-C in Complex with Human C1r.

We have previously shown that BBK32-C interacts with high affinity with an autolytic C-

terminal region of human C1r corresponding to residues 300-705 (UNIPROT numbering) (*41*).

Co-crystals of BBK32-C in complex with this fragment of C1r produced crystal samples that

generally did not yield diffraction data beyond 10 Å limiting resolution. Ultimately, a single

sample was produced that diffracted to 4.1 Å. Unfortunately, this dataset was characterized by

significant data pathologies including severe anisotropy and low completeness. Despite this, a

reasonable final solution (PDB: 7MZT) could be obtained for a single copy of the BBK32-C/C1r

complex for which readily interpretable electron density maps were observed (**Fig. 1 A, S1 A-C**

and **Table S1**). A large protein-protein interface between BBK32-C and the SP domain of C1r,

totaling 1,340 Å^2 buried surface area (b.s.a.), was present in this structure (**Fig. 1 A, D, S1 A**).

However, underlying data quality issues prevented a complete modeling of the asymmetric unit in

this crystal (see *Materials and Methods* for a detailed description). This led to elevated R_{free}/R_{work}

values (36.9%/37.2%) outside of the normal range for high quality crystallographic solutions (i.e.

< 30% R_{work}). Despite extensive effort to overcome these issues, including co-crystallization trials

with a variety of SP-containing C1r constructs, ultimately an improved crystallographic model

could not be obtained. We thus turned to an orthogonal approach involving the study of the

BBK32-C in complex with recombinant C1r-CCP2-SP in solution by SEC-SAXS.

SEC-SAXS curves were obtained for recombinant BBK32-C (**Fig. S2 A, D**), a recombinant

autoactivated C1r domain truncation construct (hereafter, C1r-CCP2-SP) (**Fig. S2 B, E**), and an

equimolar mixture of BBK32-C and C1r-CCP2-SP (**Figs. S2C, S3**). Radius of gyration (R_g) values

of 21.3, 24.0, and 27.8 Å were calculated from the Guinier region of the scattering curves for

BBK32-C, C1r-CCP2-SP, and BBK32-C/C1r-CCP2-SP samples, respectively (**Table 1**, **Figs. S2**

D, E, S3 insets). SAXS and multi-angle light scattering (MALS) determined molecular weights

(i.e., MW_{SAXS} and MW_{MALS}) are consistent with BBK32-C and C1r-CCP2-SP behaving as

monomeric species in solution (**Table 1**). For the BBK32-C/C1r-CCP2-SP sample, MW_{SAXS} =

55.6 kDa (*42*) and MW_{MALS} = ~55-65 kDa were determined, which are close to the expected

molecular weight of a 1:1 complex ($MW_{sequence}$ = 54.6 kDa). Analysis of each scattering curve

using a pair-wise distribution function, P(r), is also consistent with complex formation as

evidenced by the shape of the overlaid P(r) profiles and increased maximum particle size (D_{max})

of the BBK32-C/C1r-CCP2-SP samples relative to BBK32-C or C1r-CCP2-SP alone (**Fig. 1B,**

Table 1).

Next, we compared the solution state of each unbound protein to available high-resolution

crystal structures of BBK32-C (PDB: 6N1L (*41*)) and the activated form of C1r-CCP2-SP (PDB:

1MD8 (*43*)). First, MODELLER (*44*) as implemented in UCSF Chimera (*45*), was used to model

missing loops into each structure, and theoretical scattering profiles of each model were then

generated and compared against the raw scattering profiles using Fast X-ray Scattering (FoXS)

(*46*, *47*). Good agreement between theoretical and experimental SAXS curves were obtained for

BBK32-C (χ^2 = 2.08) and C1r-CCP2-SP (χ^2 = 1.06) atomistic models (**Fig. S2 D, E, Table 1**).

Collectively, analysis of the experimental SAXS curves shows that solution states of unbound

BBK32-C and activated C1r-CCP2-SP are in agreement with previously published crystal

structures and that BBK32-C/C1r-CCP2-SP forms a 1:1 stoichiometric complex in solution.

To better define the molecular interface formed between BBK32 and human C1r, we employed

a SAXS-restrained macromolecular computational docking approach using FoXSDock (*48*). This

method – which relies on the availability of high-resolution structures of the individual complex

components – involves initial docking using a geometric shape-matching algorithm, coarse SAXS

filtering of the resulting docking models, scoring of the interface energies, and scoring the fit of

the model's theoretical SAXS profiles to the raw SAXS profile of the BBK32-C/C1r-CCP2-SP

complex (*47*). BBK32-C and C1r-CCP2-SP were docked as a ligand-receptor pair and each model

was compared against the BBK32-C/C1r-CCP2-SP SAXS curve. The top four models (as judged

by Z scores (*48*)), each exhibited nearly identical protein-protein interfaces involving BBK32-C

contact residues originating from helix 1, 5 and the short loop connecting helix 1 and 2. Consistent

with our previous mapping study (*33*), all C1r interactions in these four models involve only

residues within the C1r SP domain. We then reconstructed a molecular envelope using DAMMIF

(*49*, *50*), which produced a distinctive chevron shape (**Figs. 1D,** gray envelope). Superimposition

of the top scoring FoxsDock model onto the shape envelope results in a close match, further

supporting the validity of the interface (**Fig. 1D**). Minimal molecular dynamics simulations using

BILBOMD (*51*) were performed on the best scoring model (χ^2 = 1.28) to eliminate potential

clashes between side chains at the interface, further improving the model's SAXS score (χ^2 =

1.24) (**Fig. 1D, Table 1**). A structural alignment of the co-crystal structure to the top three scored

SAXS-derived structures reveals a highly similar protein-protein interface, with the primary

differences being related to the orientation of the BBK32-C helical bundle that does not participate

in C1r interaction (**Fig. 1 C, E**). Furthermore, a model of 7MZT that lacks the C1r-CCP1 domain

fits closely to the experimental SAXS curve obtained for BBK32-C/C1r-CCP2-SP, as judged by

FoXS (χ^2 = 1.20) (**Fig. S3**).

Analysis and Validation of the BBK32-C/C1r Structure

Protein-protein interaction analysis was carried out using the SAXS-derived structure of

BBK32-C in complex with C1r-CCP2-SP using the programs PISA and CCP4 CONTACT (*52*,

53). The model predicts an interaction involving 28 contact residues on the BBK32-C side and 34

contact residues from human C1r, resulting in of 1,270 $Å^2$ b.s.a. (**Fig. 1E, 2A**). To validate the

predicted interface, we carried out structure-guided site-directed mutagenesis. A total of eleven

BBK32-C residues comprising > 50% of the total surface area buried by BBK32-C were changed

to alanine in single and/or double mutant forms (**Fig. 2A**, red). These mutants were then evaluated

for direct relative binding to C1r-CCP2-SP by surface plasmon resonance (SPR).

To evaluate BBK32 mutants on the same surface, we initially captured HIS-C1r-CCP2-SP

uniformly on an Ni-NTA sensorchip and binding of BBK32-C was monitored using a single-cycle

injection strategy (*54*) (**Fig. 2B**). Binding of BBK32-C to the HIS-C1r-CCP-2SP biosensor is

characterized by a long complex half-life (**Fig. 2B**, blue lines). However, this feature combined

with a small upward baseline drift of the reference-subtracted sensorgram, which was related to

the affinity-capture-based surface, resulted in poorly determined equilibrium dissociation K_D

values when fitted to a single-cycle kinetic model of binding. To overcome this limitation, and to

better match the expected physiological orientation of the interaction under study, we evaluated

C1r-CCP2-SP analyte injections over immobilized BBK32 sensorchips generated by amine-

coupling chemistry (**Fig. 2C**). Baseline drift was minimal using this covalently-coupled biosensor.

BBK32-C biosensors were created at two immobilization densities for better comparison to

individual site-directed mutant surface densities. The resulting low and high density sensorgrams

fit well to a 1:1 Langmuir kinetic binding model as judged by residual plots (**Fig. 2C** blue lines,

Figs. S4, **S5,** and **Table S2**). Consistent with our previous SPR binding measurements of BBK32-

C for C1, C1r, an autolytic fragment of C1r, and HIS-tagged recombinant domain truncations of

C1r (*33*, *41*), BBK32-C bound to activated C1r-CCP2-SP with high affinity, exhibiting an

equilibrium dissociation constant (K_D) of 0.45 nM (high density surface) and 0.78 nM for the low

density surface (**Fig. 2C, Table S2**).

Each BBK32 site-directed mutant was then evaluated by generating individual biosensors,

carrying out a C1r-CCP2-SP titration series, calculating an associated K_D, and comparing to the

BBK32-C biosensor that most closely matched wild-type surface density (**Fig. 2C, S4, S5**). Three

different double alanine mutants involving mutagenesis of small contiguous contact surfaces

resulted in a 653-fold (K327A-T331A), 127-fold (T242A-Y245A), and 615-fold (Y323A-E324A)

reduction in affinity for C1r-CCP2-SP, whereas a fourth such mutant retained an affinity similar

to wild-type (L236A-T239A) (**Figs. 2C, S4, S5, Table S2**). In the case of T242A-Y245A and

Y323A-E324A, single mutants retained wild-type-like affinity, indicating that the observed

affinity loss in the double mutants was due to a cumulative disruption of a larger contact surface

area (**Fig. 2C**). In contrast, a single K327A mutant resulted in a 153-fold loss in affinity, whereas

the paired T331A single mutant had near wild-type affinity (**Fig. 2C**). Additional mutagenesis was

directed at two arginine residues (R248 and R337). While a single R337A mutant exhibited wild-

type like affinity, mutation of R248 resulted in a 13-fold loss in affinity for C1r-CCP2-SP (**Fig.**

2B, C, S4, S5). Although they do not form a contiguous binding surface, a pairing of the two

strongest loss-of-affinity single mutants (i.e., R248A-K327A), produced a double mutant where

no detectable binding signal was observed (**Fig. 2B, C, S5**). In all cases, loss of affinity was driven

by increased dissociation rate constants (k_d) (**Table S2**). Similar qualitative results were obtained

for the single and double mutants of R248A and K327A as analytes over the HIS-C1r-CCP2-SP

biosensor (**Fig. 2B**), indicating that surface orientation and/or surface density did not greatly

influence the measurement of relative binding strength for these mutants. Taken together, these

data show that many BBK32 residues predicted to contact C1r-CCP2-SP in the SAXS structure

(**Fig. 2A**) exhibit a loss of affinity when mutated to alanine and that R248 and K327 are critical

hotspot residues for mediating C1r-CCP2-SP interaction.

BBK32 Occludes Subsites Within the Active Site of C1r

Having taken steps to validate the structural models, and in so doing, identifying key

interactions that mediate BBK32-C/C1r complex formation, we then analyzed the structure for

potential clues into the mechanism of C1r inhibitory action by BBK32. C1r is a modular serine

protease with a chymotrypsin-like SP domain (*55*). We noted that the BBK32 binding site on

human C1r is in close proximity to the C1r catalytic site (**Fig. 3A**). Although limited contacts are made in the model with active site residues (28 Å^2 b.s.a.), much more extensive interactions are made with the specificity pocket subsites S1 (176 Å^2 b.s.a.) and S1' (167 Å^2 b.s.a.) (**Fig. 3B**). Residues involved in S1 (red) and S1' (blue) subsite interactions were mapped onto the BBK32-C surface (**Fig. 3B**). These include three residues (T242, Y245, and R248) that were shown to have loss of affinity effects when mutated to alanine (**Fig. 2**).

The steric occlusion of the S1 and S1' site by BBK32 in the model predicts that it would also be capable of directly blocking the hydrolysis of small peptide C1r substrates, as well as the larger natural C1r substrates, C1r and C1s proenzymes. Previously, we have shown that BBK32 blocks cleavage of proenzyme C1s by isolated C1r enzyme using an SDS-PAGE-based readout (*33*). We adapted this assay to one that utilizes a colorimetric substrate specific for C1s enzyme (i.e. Z-L-Lys thiobenzyl), and found that BBK32-C, but not R248A-K327A, could prevent activation of proenzyme C1s by C1r enzyme. Similar results were obtained when C1r-CCP2-SP cleavage of a colorimetric peptide substrate that requires only P2-P1-P1' access (i.e. Z-Gly-Arg-thiobenzyl) was monitored (**Fig. 3C**). Furthermore, BBK32-C, but not R248A-K327A, prevented cleavage of this substrate by spontaneously autoactivated full-length proenzyme C1r (**Fig. 3C**). To further evaluate the interaction of BBK32-C with the proenzyme form of C1r, we produced two forms of zymogen-stabilized C1r-CCP2-SP mutants. Both the scissile loop (i.e. R463Q) and catalytic serine-based (S654A) zymogen forms of C1r-CCP2-SP bound similarly to BBK32-C, and like autoactivated C1r-CCP2-SP, exhibited slow dissociation rates (**Fig. 3D**). Together, these data show that BBK32 directly blocks both small and large C1r substrates from proteolytic cleavage and that BBK32 recognizes both proenzyme and activated enzyme forms of C1r.

R248 and K327 are Critical for BBK32-C Complement Inhibitory Activity

Next we tested our panel of structure-based recombinant mutants for their ability to block

serum-based human complement activation in an ELISA assay that detects the classical pathway

activation product C4b. BBK32-C exhibited a half-maximal inhibitory concentration (IC_{50}) of 15

nM in this assay format (**Fig. 4A, Table S2**), consistent with our previously reported value of 5.6

nM (*33*). With the exception of E324A, which is discussed below, near wild-type IC_{50} values were

measured for all BBK32-C mutants that retained high affinity for C1r-CCP2-SP, although we

noted a modest reduction in inhibitory activity in the Y245A (~5-fold), N251A (~6-fold) and

Y323A (~2-fold) single mutants (**Fig. S6 and Table S2**). Importantly, in all cases where mutants

exhibited a strong loss of affinity for C1r-CCP2-SP, they were also greatly impaired in their ability

to block classical pathway activation. For example, the contiguous surface double mutants Y323A-

E324A and K327A-T331A failed to block complement at any concentration, while a ~240-fold

weaker IC_{50} value for T242A-Y245A was measured (**Fig. S6 and Table S2**). Likewise, the R248A

mutant was attenuated ~28-fold, and neither the K327A nor the R248A-K327A mutant blocked at

any of the concentrations tested (**Figs. 4A, and Table S2**).

Based on the SAXS-derived BBK32-C/C1r-CCP2-SP structure we were able to identify

several BBK32 residues that are important for its activity. In particular, substitution of R248 and

K327 to alanine in single mutants, or as a double mutant, results in a significant reduction of both

C1r-binding and inhibitory activity. To study these residues in a more physiological setting, we

produced site-directed BBK32 'knock-in' strains in the *B. burgdorferi* B314 genetic background

(**Fig. 4B**). *B. burgdorferi* B314 lacks linear plasmids that encode many of the outer surface proteins

associated with infection, including BBK32 (*56*, *57*). Previously we have shown that when wild-

type BBK32 is expressed heterologously on the surface of *B. burgdorferi* B314, it protects these

normally serum-sensitive spirochetes from human complement-mediated bacteriolysis (*33*, *41*).

Introduction of the R248A mutant into *B. burgdorferi* B314 resulted in a minimal reduction in

inhibitory activity relative to wild-type BBK32, while spirochetes expressing K327A were

significantly more sensitive to human serum. Spirochetes transformed with the *bbk32* R248A-

K327A allele lost all protective activity and were indistinguishable from the vector-only control

(**Fig. 4C**). These data confirmed that the R248 and K327 residues are critical for eliciting

complement inhibitory activity in the context of full-length BBK32 on the outer surface of *B.*

burgdorferi. Western immunoblots showed that full length BBK32 was produced equivalently in

all mutants tested, indicating that the enhanced sensitivity of the R248-K327A double mutant was

not the result of proteolytic instability (**Fig. 4B**).

BBK32-C Targets Non-Conserved C1r Loops Outside of the Active Site

We have previously shown that BBK32 is highly selective for C1r-binding relative to C1s

(*33*). To gain insight into this phenomenon, and with no comparable C1r inhibitors having been

reported, we instead compared the BBK32-C/C1r-CCP2-SP SAXS structure to that of a previously

published crystal structure of a protein inhibitor of both the classical and lectin complement

pathways, known as gigastasin (*58*). Gigastasin differs in selectivity from BBK32 as it blocks C1s,

MASP-1 and MASP-2, but is a relatively weak inhibitor of C1r (*58*). Interestingly, the

gigastasin/C1s protein-protein interface shares common features with the BBK32-C/C1r-CCP2-

SP SAXS structure presented here. Namely, both inhibitors form extensive interactions with the

S1 and S1' subsites of their respective target enzymes. Furthermore, BBK32 and gigastasin each

interact with three or more residues in the A and D loops, as well as loop 2 (**Fig. 5**). However,

BBK32 differs from gigastasin in that it makes extensive interactions with the B loop of C1r. This

loop contains an insertion sequence making it longer in C1r compared to most other complement

serine proteases, including C1s (**Fig. 5**). Interestingly, the B loop also is thought to block the

binding of the antistasin domain of gigastasin making it a determinant of inhibitor specificity (*43*,

58). Of the 16 residues that comprise the C1r B loop, 10 form contacts with BBK32-C and none

are identical at the homologous position in C1s (**Fig. 5**). Considering the entire interface, 25 of the

34 C1r contact residues predicted by the BBK32-C/C1r-CCP2-SP SAXS structure are non-

identical in C1s (**Fig. 5**). Most of these arise from surfaces involving the S1' site and four C1r

loops (i.e. A, B, D, E loops), suggesting these interactions may underlie the selectivity of BBK32

for C1r. Given the vastly different three-dimensional structures of BBK32 and gigastasin (**Fig. 5**),

it appears that a bloodborne prokaryote (*B. burgdorferi*) and a blood-feeding eukaryote

(*Haementeria ghilianii*) have evolved complement inhibitors with at least partially convergent

mechanisms of action, while also evolving differential target selectivity.

Discussion

In the absence of regulation, complement activation at the microbial surface leads to

deposition of C3b or C4b on the microbe surface, recognition by innate immune cells due to the

presence of immobilized C3b and C4b, recruitment of leukocytes by the anaphylatoxins C3a and

C5a, stimulation of the adaptive immune response by each of these fragments, and direct

bactericidal action by C5b-9 (i.e., MAC) (*1*, *2*). While intracellular pathogens may be protected by

complement within an intracellular niche, extracellular pathogens like *Borreliella* spirochetes face

a continuous threat from the complement systems of their hosts.

Borreliella first encounter complement during the bloodmeal of an infected *Ixodes* sp. tick.

Following transmission from their arthropod vector into the vertebrate host, spirochetes must

continue to protect themselves from complement during hematogenous dissemination, a challenge

that persists once they colonize distant tissues. To evade complement attack, Lyme disease

spirochetes express nearly a dozen different outer surface lipoproteins that directly interact with

complement proteins and modulate their native activities (*16*, *17*). While we have previously

shown that *B. burgdorferi bbk32* is required for optimal infectivity in naïve mice, it is likely that

C1r-specific inhibition by BBK32 also protects *B. burgdorferi* from targeted antibody-dependent

complement-mediated killing, a hallmark of the classical pathway. This contention is supported

by the persistent infection observed despite the presence of high titered antibody responses against

B. burgdorferi that can passively protect against borrelial infection and kill *B. burgdorferi in vitro*

in a complement-dependent manner (*59–61*).

In this study we aimed to determine the structural basis for the inhibition of C1r by *B.*

burgdorferi BBK32. A clear limitation of our study is the low resolution inherent to SAXS

experimental data. Importantly, the model presented here assumes limited conformational

rearrangement of BBK32-C and C1r-CCP2-SP relative to their unbound forms as previously

determined by crystallography. While analysis of the SAXS data for BBK32-C and C1r-CCP2-SP

suggests that each unbound protein adopts a similar structure in solution as they do in their

respective crystal structures (**Fig. S2**), we recognize that our approach would be unable to detect

larger scale conformational changes induced by BBK32-C/C1r-CCP2-SP complex formation. We

note that a highly similar interface is observed in the co-crystal structure presented here (**Fig. 1**),

although we acknowledge that the high R values and associated issues in the crystallographic data

quality are an additional limitation. Nonetheless, the validity of the SEC-SAXS-derived structure

is supported by its ability to successfully guide the identification of many BBK32 residues

important for driving both affinity and inhibitory activity (**Figs. 2-4**). Furthermore, the structure

provides a basis for an inhibitory mechanism that is consistent with the inhibitory activities and

selectivity of BBK32 (**Figs. 3-5**). Taken together, our data suggests that BBK32 inhibits human

C1r by simultaneously targeting its active site, as well as functionally critical, less conserved, non-

active site residues on loops of the C1r SP domain.

Like Lyme disease spirochetes, other bloodborne pathogens and blood feeding eukaryotes have

evolved complement evasion systems. For example, potent direct inhibitors of complement are

produced in the salivary glands of several species of ticks, including *Ixodes* sp. (*62–66*). Similarly,

blood-feeding insects like sandflies (*67*) and mosquitos (*68*), along with other human extracellular

pathogens such as *Staphylococcus aureus* (*15*, *69*, *70*), all produce a wide array of mechanistically

distinct protein inhibitors of complement. However, among known complement evasion

molecules, relatively few target the classical pathway with specificity (*70*), and to our knowledge

BBK32 remains the only reported selective inhibitor of C1r. Despite this, our study identifies a

surprising convergence of function between *B. burgdorferi* BBK32 and a complement protease

inhibitor of the classical and lectin pathways previously identified in the giant Amazonian leech

(i.e. gigastasin). However, gigastasin is related to a class of inhibitors known as antistasins (*71*),

that preferentially bind to the active form of their enzyme targets. This is in contrast to BBK32,

which recognizes both proenzyme and enzyme forms of C1r (**Figs. 2, 3**). While each inhibitor

functions, at least in part, by steric occlusion of the subsites upon enzyme-inhibitor complex

formation (*58*), it appears that differences in non-active site loops may drive selectivity of these

inhibitors (**Fig. 5**).

Borreliella spirochetes vary widely in their susceptibility to serum-based complement-

mediated bacteriolysis (*72*). While the categorizations are general and not without exceptions, *B.*

burgdorferi is typically considered serum-resistant whereas *B. garinii* is susceptible to killing by

human serum. The overall susceptibility of a spirochete to a particular host complement system is

related to the collective activities of the larger *Borreliella* complement evasion system (i.e.,

CRASPs, OspC, BBK32, etc.). However, insight into this host association phenomenon may be gained from analysis of a single *Borreliella* complement inhibitor, in this case BBK32. While the C1r-CCP2-SP binding affinity of nearly all mutants in our study correlated with the relative inhibitory activity, a curious exception was BBK32-E324A. This mutant was reduced only two-fold in affinity for C1r-CCP2-SP (**Fig. 2, S4**), but by several orders of magnitude in inhibitory activity (**Fig. S6, Table S2**). BBK32-E324 is located next to K327 on the face of alpha helix 5 of BBK32 and, like K327, is positioned across from the B loop of C1r in the model. In our previous study we coincidentally probed the function of E324 while investigating the reduced complement inhibitory activity of a BBK32 homologue from *B. garinii*, termed BGD19 (*41*). In that study, a triple mutant chimeric recombinant protein termed BXK32-C was created to convert BBK32 to BGD19 at three non-conserved residue positions (i.e., BBK32-E308K-Q319K-E324Q) (*41*). This mutant shifted from BBK32-like to BGD19-like inhibitory activity, leading us to conclude that at least one of these three surface-exposed residues was important for the complement inhibitory activity of BBK32. Unlike R248 and K327, which drive human C1r affinity and are universally conserved across publicly available *B. burgdorferi* genome sequences, position 324 is less conserved (**Fig. S7**). While the underlying reasons for the disconnect between the binding and inhibitory activities of these proteins remain unclear, we speculate that variation in position 324 in naturally occurring BBK32 sequences may allow for optimal inhibition of C1r in other vertebrates as the sequence of the C1r B loop is one of the least conserved features within vertebrate C1r. Further studies, including high-resolution structural information involving synthetic or naturally occurring BBK32 sequences such as *B. garinii* BGD19, along with mechanistic studies involving non-human C1r or extensive C1r mutagenesis, will be needed to further test this possibility.

For the first time, this study provides a structural basis for classical pathway complement

evasion by a human pathogen. Among naturally occurring complement inhibitors, *B. burgdorferi*

BBK32 is unique in its ability to selectively target and inhibit the initiator protease of the classical

pathway, C1r, and, as such, the structure-function studies reported here provide new insight into

an evolutionarily optimized mechanism of action for C1r inhibition. This knowledge improves our

understanding of complement evasion by bacterial pathogens, increases our understanding of

naturally occurring complement inhibitors, and provides a potential basis for the development of

novel complement directed therapeutics that are classical pathway-specific.

Materials and Methods

Strains, Plasmids constructed, and Oligonucleotides.

All bacterial strains, plasmid constructs, and oligonucleotides used are listed in

supplemental **Table S3**. *B. burgdorferi* strain B314, with all recombinant plasmid constructs, was

grown in complete BSK-II media containing 300 µg/ml kanamycin as previously described (*41*).

E. coli cells were grown in LB media with kanamycin at 50 µg/ml.

Plasmid constructs for usage in *B. burgdorferi* strain B314 were constructed in a step-wise

manner. First, pCD100 was used as template to obtain the *bbk32*-R248A allele through use of the

oligonucleotides R248A mutant F and R248A mutant R that face in the opposite orientation and

contain the sequence to change codon 248. To facilitate this, GXL polymerase (Takara Bio) was

used with the following parameters: a 15 second denaturing step at 98°C, an annealing temperature

dependent on the lowest melting temperature of primers used for a specific reaction, and a 68°C

extension step for 1min/kb. The reaction was then digested with *Dpn*I (NEB), to digest methylated

parental DNA, and transformed into NEB5-alpha competent *E. coli*. The resulting plasmid was

designated pAP5. The *bbk32*-K327A encoding construct was obtained by synthesizing a 241 bp

fragment starting at nucleotide 861 and encompassing the remainder of *bbk32* and a 40 base pair

region of downstream sequence (IDT gBlocks). This fragment of *bbk32* contains native sequence

with the exception of the K327A codon. The K327A containing fragment and either pCD100 (for

pAP6 construction) or pAP5 (for pAP7 construction) were used as template with oligonucleotide

primers described in **Table S3** for PCR amplification. The resulting PCR products were assembled

using the NEBuilder HiFi DNA Assembly Cloning Kit as described in (*41*). The construct

encoding the *bbk32*-K327A allele was named pAP6. The final construct that encoded the *bbk32*-

R248A/K327A double mutant was designated pAP7. All plasmids were sequenced to ensure that

the desired mutations were obtained and that no additional mutations were introduced into *bbk32*.

Transformations of strain B314 with the aforementioned plasmid constructs were done as

described in (*73*, *74*).

For the production of all recombinant BBK32 proteins, DNA fragments for corresponding

site directed mutations in BBK32-C (residues 206-348, *B. burgdorferi* B31), for expression in *E.*

coli, were *E. coli* codon optimized and synthesized commercially by IDT Technologies gBlock

Gene Fragment service. The R337A mutant contained the C-terminal 6 residues in the full length

BBK32 sequence making it residues 206-354 (*33*). IDT gBlocks were designed with 5' BamHI,

3' NotI sites and stop codon, which then were subsequently cloned into pT7HMT as previously

described (*33*, *75*). Wild-type BBK32-C and C1r-CCP2-SP were expressed and purified form

previously generated constructs (*33*, *41*, *75*).

Proteins

Recombinant BBK32-C proteins were expressed and purified as previously described (*33*,

41), with the following modifications. After elution in Ni-NTA elution buffer (20 mM Tris (pH

8.0), 500 mM NaCl, 500 mM imidazole (pH 8.0)), proteins were exchanged into Ni-NTA-binding

buffer (20 mM Tris (pH 8.0), 500 mM NaCl, 10 mM Imidazole (pH 8.0)) using a Hi-Prep Desalting

26/10 column (GE Healthcare). To remove the HIS-myc affinity tag, proteins were incubated with

HIS tagged tobacco etch virus (TEV) protease and 5mM β-mercaptoethanol at room temperature

overnight. The digested proteins were then passed over a 5mL HisTrap-FF column and the flow-

through was collected for further purification by gel filtration chromatography using a HiLoad

Superdex 75 26/600 pg column (GE Healthcare). The monodisperse peak corresponding to

BBK32-C was pooled following analysis of chromatography fractions by SDS-PAGE and

exchanged into HBS (10mM HEPES (pH 7.3), 140mM NaCl). C1r-CCP2-SP, C1r-CCP2-SP

zymogen mutants, and His-C1r-CCP2-SP were purified according to methods as described

previously, with a final polishing step involving gel filtration chromatography on a HiLoad

Superdex 75 column (GE Healthcare) (*41*, *55*, *76*). His-tagged proteins were not subjected to HIS-

TEV cleavage. An autoproteolytic fragment of human C1r corresponding to residues 300-705

(UNIPROT: P00736) used in the co-crystallography experiment was obtained as previously

described (*41*). Full length complement proteins C1r proenzyme, C1r enzyme, and C1s proenzyme

were purchased from Complement Technology. In order to test for global changes in protein

folding that may be induced by site-directed mutagenesis, recombinant proteins were evaluated

for major changes in size by size exclusion chromatography (**Fig. S8**). Proteins were separated on

a HiLoad Superdex 200 Increase 10/300 column (GE Healthcare) and plotted against BioRad

Standards to determine size. Monodisperse peaks for all BBK32-C related samples exhibited a

ranging between 13.2 and 16.4 kDa.

Statistical analysis

Statistical analysis of assays was performed on GraphPad version 8. Calculations of IC_{50}

for classical pathway ELISA was determined with a normalized variable four-parameter fit.

Statistical significance for colorimetric enzyme assays were analyzed using a paired t-test.

Significance for *B. burgdorferi* serum sensitivity assays were analyzed using a two-way ANOVA

with a Šidák correction for multiple comparisons.

Size exclusion chromatography-coupled Small Angle X-ray Scattering Data Collection

Volumes of 60 µl samples containing either 8.0 mg mL^{-1} BBK32-C, 2.0 mg mL^{-1} C1r-

CCP2-SP, or 2.0 mg mL^{-1} BBK32-C/C1r-CCP2-SP 1:1 Complex were prepared in 10 mM HEPES

pH 7.3, 140 mM NaCl buffer. SEC-SAXS-MALS were collected at the ALS beamline 12.3.1

LBNL, Berkeley, California [77]. X-ray wavelength was set at λ=1.127 Å and the sample-to-

detector distance was 2100 mm resulting in scattering vectors, q, ranging from 0.01 Å^{-1} to 0.4 Å^{-1}

. The scattering vector is defined as $q = 4\pi\sin\theta/\lambda$, where 2θ is the scattering angle. All experiments

were performed at 20°C and data was processed as described [78]. Briefly, a SAXS flow cell was

directly coupled with an online Agilent 1260 Infinity HPLC system using a Shodex KW802.5

column. The column was equilibrated with running buffer (10 mM HEPES (pH 7.3), 140 mM

NaCl) with a flow rate of 0.5 mL min^{-1}. 55 µL of each sample was run through the SEC and three

second X-ray exposures were collected continuously during a 30-minute elution. The SAXS

frames recorded prior to the protein elution peak were used to subtract all other frames. The

subtracted frames were investigated by radius of gyration (Rg) derived by the Guinier

approximation $I(q) = I(0) \exp(-q^2Rg2/3)$ with the limits qRg<1.5. The elution peak was mapped

by comparing integral of ratios to background and Rg relative to the recorded frame using the

program SCÅTTER. Uniform Rg values across an elution peak represent a homogenous assembly

(**Fig. S2**). Final merged SAXS profiles, derived by integrating multiple frames across the elution

peak, where used in PRIMUS [79] for further analysis including Guinier plot which determined

aggregation free state (**Figs. S2D-E, S3**). The program GNOM (*80*) was used to compute the pair

distribution function (P(r)) (**Fig. 1B**). The distance r where P(r) approaches zero intensity identifies

the maximal dimension of the macromolecule (D_{max}). P(r) functions were normalized to peak

maxima. Eluent was measured in line with a series of UV at 280 and 260 nm, multi-angle light

scattering (MALS), quasi-elastic light scattering (QELS), and refractometer detector. MALS

experiments were performed using an 18-angle DAWN HELEOS II light scattering detector

connected in tandem to an Optilab refractive index concentration detector (Wyatt Technology).

System normalization and calibration was performed with bovine serum albumin using a 45 μL

sample at 10 mg mL-1 in the same SEC running buffer and a dn/dc value of 0.19. The light

scattering experiments were used to perform analytical scale chromatographic separations for

MW_{MALS} determination of the principle peaks in the SEC analysis (**Fig. S2A-C**). UV, MALS, and

differential refractive index data was analyzed using Wyatt Astra 7 software to monitor the

homogeneity of the sample across the elution peak complementary to the above-mentioned SEC-

SAXS signal validation (**Fig. S2**).

SAXS Solution Structure Modeling

High resolution crystal structures for BBK32-C (PDB:6N1L) and C1r-CCP2-SP

(PDB:1MD8) were used for structural modeling (*41*, *43*). Missing residues, including the GSTGS

sequence cloning artifact, were added using COOT (*81*). In order to test multiple configurations

of residues added in this manner, structures were imported into UCSF-Chimera and the

MODELLER (*82*) tool was used to make five models. Each model was tested against SAXS data

in FoXS and the top scoring model was accepted. The BBK32-C/C1r-CCP2-SP complex was

determined by using FoXSDock (*47*) to perform rigid docking the highest overall score of the

almost 5000 models were accepted. Minimal molecular dynamics (MD) simulations   using

BILBOMD (*51*) was performed on the best scoring model to eliminate potential clashes between

the residues side chains at the interface. The experimental SAXS profiles were then compared to

theoretical scattering curves generated from molecular models using the FOXS (*46*, *48*). Shape

envelope reconstruction for the BBK32-C/C1r-CCP2-SP sample was carried out using DAMMIF

as implemented in the ATSAS software suite (*49*, *83*). The final model was produced from the

average of ten models filtered for low occupancy using default values for DAMAVER/DAMFILT

(*50*).

Crystallization, Structure Determination, Refinement and Analysis

Purified BBK32-C and a purified autolytic fragment of human C1r (Complement

Technologies) were mixed at 5% molar excess BBK32-C and concentrated to 3.5 mg ml^{-1} in 10

mM HEPES (pH 7.3) 140 mM NaCl. Small plate-like crystals were obtained by vapor diffusion in

hanging drops at 20 °C by mixing 1 µl of protein solution with 1 µl of precipitant solution that had

been diluted into 3 µl of double deionized water. The precipitant solution contained 0.2 M sodium

formate and 27.5% (v/v) PEG 3,350. Crystals appeared in 2-5 days and were harvested and

cryoprotected in the precipitant solution supplemented with 5% (v/v) glycerol. Monochromatic X-

ray diffraction data were collected at 1.00 Å using beamline 22-ID of the Advanced Photon Source

(Argonne National Laboratory) using an Eiger 16M detector. While several samples were

harvested from this condition (and subsequent replications of this condition), only two samples

diffracted X-rays beyond ~10 Å and ultimately a single dataset was collected on one of these

crystals. Diffraction data were integrated, scaled, and reduced using the HKL2000 software suite

(*84*) and a high-resolution cut-off of 4.1 Å was chosen on the basis of I/σI and CC$_{1/2}$ (**Table 1**).

Consistent with the weak diffraction patterns observed, and difficulty in obtaining diffraction data

in certain crystal orientations, PHENIX XTRIAGE (*85*) analysis suggested several data

pathologies. This included severe anisotropy and low overall completeness. Additionally, strong

evidence of translational non-crystallographic symmetry was observed by Patterson map analysis.

Initial space group analysis suggested primitive tetragonal symmetry, however the final solution

was obtained in space group P 21 21 2 (**Table S1**). Molecular replacement was carried out with

PHASER in the Phenix software suite (*85*). Although translational non-crystallographic symmetry

was detected, the option to account for this automatically during MR was deselected after initial

failed solutions. An initial search was performed using a model of the monomeric activated form

of recombinant C1r-CCP1-CCP2-SP (PDB: 2QY0, (*86*)) where four C-terminal loop residues were

removed from Chain A. This search produced a solution in P 21 21 2 (LLG:248; TFZ: 18.9). This

solution was fixed, and a monomeric copy of BBK32-C was used for a subsequent search

producing a new solution (LLG: 424; TFZ: 17.8). Initial refinement of this solution using a single

round of individual B-factors was performed (R_{work} = 42.5% / R_{free} = 44.4%) producing readily

interpretable *2 F*o-*F*c density maps for both BBK32-C and C1r. Analysis of the asymmetric unit

suggested two copies of the complex were probable (one copy: 74.3% solvent content / Matthews

coeff.: 4.79 vs. two copies: 48.7% / 2.39). Additional exhaustive searches of various combinations

of the complex, complex subcomponents, or individual domains yielded highly scoring solutions,

however, presented unrealistic packing that could not be resolved by occupancy refinement.

Higher and lower symmetry space groups were explored with no improvement. While we

acknowledge other possibilities exist, our judgement is that two copies of the complex existed in

the crystal, but the second copy cannot be confidently placed due to the underlying poor data

quality and associated data pathologies of the crystal sample. Extensive efforts failed at producing

better quality crystal samples, leading us to turn to an alternative SAXS-based approach. However,

interpretable density of the single copy of the complex (**Fig. 1A, S1**) prompted us to carry forward

with refinement in the absence of the second suspected asymmetric unit copy. Analysis of the

contacts between BBK32-C and C1r in the obtained solution revealed three interfaces. Two of

these interfaces were sized consistent with crystallographic contacts (346 $Å^2$, and 344 $Å^2$) and one

large interface formed by a symmetry-mate of 1,108 $Å^2$, which was rearranged into the asymmetric

unit. Using this model, successive rounds of manual building using COOT (*81*) and iterative

individual B-factor refinement using PHENIX.REFINE resulted in the final model 36.7%/37.2%

(R_{work}/ R_{free}). A description of crystal cell constants, diffraction data quality, and properties of the

final model can be found in **Table S1**. Representations of protein structures and electron density

maps were generated in PyMol (www.pymol.org/).

Surface Plasmon Resonance

SPR experiments were performed on a Biacore T200 instrument at 25º C at a flowrate of

30 µL min^{-1} with a running buffer of HBS-T (10 mM HEPES (pH 7.3), 140 mM NaCl, 0.005%

Tween 20). Proteins were immobilized either on a NiD 200M Ni^{2+} chelator-modified sensorchips

or CMD 200 biosensor chips (Xantec Bioanalytics) using standard amine coupling as previously

described (*33*). For experiments involving immobilization of HIS-tagged species of C1r-CCP2-

SP, NiD 200M surfaces were conditioned for 5 min with 0.5 M Na-EDTA (pH 8.5), followed by

equilibration in HBS-T for 2 min. A solution of 5 mM nickel chloride was then injected for 2 min.

Next, HIS-tagged ligands were captured on individual flow cells at densities between 300-400 RU.

Single cycle injection curves were performed at five concentrations of BBK32-C or BBK32-C

site-directed mutants (1, 3.2, 10, 32, 500 nM). For experiments using amine coupling,

immobilization densities for each protein used in the study are presented in **Table S2**. Single cycle

injection curves were performed with a five concentration five-fold injection series of (0-500 nM)

C1r-CCP2-SP with 2 min association and a final dissociation time of 30 min. BBK32-C was

immobilized under two immobilization densities on adjacent flow cells on the same chip: BBK32-

$C_{High\ Density}$ (2011.4 RU) and BBK32-$C_{Low\ Density}$ (604.7 RU). Subsequent regeneration was

performed by three 30 s injections of a regeneration solution (0.1 M glycine (pH 2.2), 2.5 M NaCl).

Kinetic analysis was performed on each injection series using a 1:1 (Langmuir) binding model was

applied using Biacore T200 Evaluation Software (GE Healthcare). Equilibrium dissociation

constants (K_D) were determined from resulting fits and are shown as mean K_D. Quality of fits were

assessed by inspection of residual plots (**Fig. S4 .S5**).

Classical pathway inhibitory assay

Classical pathway inhibition was determined using an ELISA-based assay as previously

described (*33, 41, 87*). Briefly, high-binding 96 well plates (Grenier Bio-One) were coated with

human IgM (MP Biomedical) and activation of classical pathway was monitored using 2% normal

human serum (NHS; Innovative Research) and C4b detection was determined by monoclonal

antibody (Santa Cruz Biotechnology). Dose-dependent inhibition was performed using a 12-point,

two-fold dilution series of BBK32-C and mutants ranging from 1.1 to 2200 nM. Each assay was

performed at least in duplicate and normalized to in-column positive (no inhibitor) and negative

(no serum) controls.

Serum complement sensitivity assay

B. burgdorferi strain B314 were grown at 32°C in 1% CO_2 to early- to mid-log phase and

diluted in complete BSK-II media to a concentration of 1×10^6 cell/ml. *B. burgdorferi* cells were

incubated in normal human serum (NHS; Complement Technology) at a final concentration of

15%. All strains were tested in duplicate independently four times. Samples were incubated at

37°C with gentle agitation for 1.5 hours with viability assessed by dark field microscopy.

Sensitivity was assessed by lack of motility, overt membrane damage, and/or cell lysis. The data

is represented as the average of all fields counted for biological replicates (40 minimum) in each

of the four independent assays.

Immunoblots

Western immunoblots were conducted to detect BBK32 from the parent and mutant

constructs produced in *B. burgdorferi* strain B314. Protein lysates were resolved by SDS-12.5%

PAGE and transferred to PVDF membranes. Membranes were incubated with either a monoclonal

antibody to BBK32 (diluted at 1:20,000) or *B. burgdorferi* strain B31 FlaB (Affinity BioReagents;

diluted at 1:10,000), washed, and then incubated with a 1:10,000 dilution of Goat anti-mouse IgG-

HRP (Thermo Fisher). The blots were processed to visualize the antigens as previously described

(*41*)

C1s proenzyme activation assay

Inhibition of C1r enzyme mediated activation of C1s proenzyme was assessed by a

colorimetric peptide cleavage assay. 2.2 µM BBK32-C or R248A-K327A was incubated with 25

nM C1r enzyme, 100 µM DTNB (5,5'-Dithiobis-(2-Nitrobenzoic Acid), Ellman's Reagent), and

C1s specific active site analog 100 µM Z-L-Lys-thiobenzyl (MP Biomedical). Lastly 3.13 nM

proenzyme C1s was added in HBS-T buffer (10 mM HEPES (pH7.3), 140 mM NaCl, .005%

Tween 20). Absorbance was monitored at 405 nm at 37º C on a Versamax plate reader for 1 hr in

triplicate. Data were in column normalized to no inhibitor and no C1s proenzyme controls.

C1r peptide cleavage enzyme assays

C1r-CCP2-SP or C1r proenzyme by BBK32-C or R248A-K327A, a colorimetric peptide

cleavage assay was employed (*55*). Single dose inhibition of 25 nM C1r-CCP2-SP or 25 nM C1r

proenzyme with 2.2 µM BBK32-C or R248A-K327A was performed in the presence of 300 µM

Z-Gly-Arg-thiobenzyl (MP Biomedical). Esterolytic cleavage was assayed by 100 μM DTNB

(5,5'-Dithiobis-(2-Nitrobenzoic Acid), Ellman's Reagent) in HBS-T buffer (10 mM HEPES

(pH7.3), 140 mM NaCl, .005% Tween 20). Absorbance was monitored at 405 nm at 25º C on a

Versamax plate reader for 1 hr in triplicate. Data were in column normalized to no inhibitor and

no C1r-CCP2-SP controls, with 100% cleavage defined as the amount of Z-Gly-Arg-thiobenzyl

cleaved by C1r-CCP2-SP in 1 hr at 25º C. Experimentation was performed in triplicate.

Data Availability

SAXS data and models are deposited in the Small Angle Scattering Biological Data Bank

(www.sasdb.org) with accession codes SASDKX7 (BBK32-C/C1r-CCP2-SP), SASDKV7 (C1r-

CCP2-SP), and SASDKW7 (BBK32-C). The co-crystal structure of BBK32-C/C1r$_{300-705}$ was

deposited into the Protein Data Bank, Research Collaboratory for Structure Bioinformatics,

Rutgers University (www.rcsb.org/) under the PDB code 7MZT.

Acknowledgments

SAXS data was collected at SIBYLS which is supported by the DOE-BER IDAT DE-

AC02-05CH11231 and NIGMS ALS-ENABLE (P30 GM124169 and S10OD018483). X-ray

diffraction data were collected at Southeast Regional Collaborative Access Team 22-ID beamline

at the Advanced Photon Source, Argonne National Laboratory. Supporting institutions may be

found at www.ser-cat.org/members.html. Use of the Advanced Photon Source was supported by

the US Department of Energy, Office of Science, Office of Basic Energy Sciences, under Contract

W-31-109-Eng-38. We thank Dr. Brian Geisbrecht (Kansas State University) for helpful

discussion related to this work. This study was supported by Public Health Service Grants

AI133367 and AI146930 (to JTS and BLG).

 Figures and Tables

Figure 1. Structures of BBK32-C in complex with human C1r fragments.

A) The co-crystal structure of BBK32-C in complex with an autolytic fragment of C1r (residues

300-705) at 4.1Å (PDB: 7MZT) is shown with 2*Fo-Fc* density contoured at 1.2 σ for the BBK32-

C polypeptide (Chain I; pink mesh) and C1r (Chains A, B; blue mesh). **B)** Normalized pair

distribution curves (P(r)) of C1r-CCP2-SP/BBK32-C complex (red), C1r-CCP2-SP (green), and

BBK32-C (blue) derived from SEC-SAXS experiments. **C)** An overlay of the top three BBK32-

C/C1r-CCP2-SP SAXS models obtained from FoXS Dock as judged by χ^2 where C1r-CCP2-SP

are colored in shades of green and BBK32-C in shades of blue. PDB: 7MZT was aligned to the

C1r SP domain of the SAXS structures (grey). The protein-protein interface in each structure is

similar with the largest differences being attributed to the orientation of the helical bundle that

does not participate in the interface. **D)** *Ab initio* SAXS envelope (gray) aligned with SAXS

solution structure from FoXSDock, refined with BILBOMD. **E)** Contact residues as judged by the

EMBL PDBePISA server for the BBK32-C/C1r-CCP2-SP SAXS structure (left, cyan) and the

BBK32/C1r co-crystal structure (right, red).

Figure 2. SPR binding assays with structure-guided site-directed BBK32 mutants. A) Surface

representation of BBK32-C with non-contact surface areas (gray) and BBK32-C/C1r-CCP2-SP

contact area (cyan). Residues mutated for this study (red) are indicated in **Table S2**. Site-directed

mutants used in **Fig. 2B-C** are denoted with arrows. **B)** SPR experiments were performed by

capturing His-C1r-CCP2-SP on a Ni-NTA sensor chip. BBK32-C and selected site-directed

mutants were injected over the His-C1r-CCP2-SP surface in a concentration series (1.0, 3.2, 10.0,

31.6, 100 nM). Injection phases are denoted with arrows. **C)** SPR experiments were performed in

the reverse orientation compared to panel **B** by immobilizing each BBK32 protein to sensor chips

using amine coupling chemistry. A fivefold concentration series of C1r-CCP2-SP (0.8, 4.0, 20.0,

100, 500 nM) was sequentially injected over each immobilized BBK32-C mutant. Arrows indicate

injection phases of each C1r-CCP2-SP concentration. Representative sensorgrams are shown with

ligands of similar immobilization density grouped with either high immobilization density

BBK32-C (i.e., BBK32-C$_{\text{High Density}}$) or low immobilization density BBK32-C (i.e., BBK32-C$_{\text{Low}}$

$_{\text{Density}}$). BBK32-C curves are reproduced in each respective panel for comparison. SPR experiments

were performed in duplicate. Dissociation constants (K_{D}) were calculated by kinetic fits with a 1:1

Langmuir model (**Fig. S4**, **S5, Table S2**). Additional site-directed mutants analyzed by SPR and

are shown in **Fig. S5**.

Figure 3. BBK32-C occludes the S1 and S1' subsites on C1r. A-B) C1r-CCP2-SP shown with the catalytic site (yellow), substrate specificity subsites S1 (red) and S1' (blue) as determined by Budayova and colleagues (*43*) using nomenclature provided by Schechter and Berger (*88*). BBK32-C shown with residues that contact the C1r S1 site (red) and S1' site (blue). **C)** The ability of BBK32-C or BBK32-R248A-K327A to directly inhibit cleavage of a small peptide or a natural substrate was evaluated by colorimetric substrate-based enzyme assays. BBK32-C significantly blocked C1r enzyme cleavage of C1s proenzyme, autocatalysis of C1r proenzyme and C1r-CCP2-

SP enzyme. All experiments were performed in triplicate and normalized to positive (no inhibitor)

and negative (no enzyme) controls. Statistical significance was evaluated using paired t-tests with

p-values noted above each experiment. **D)** To evaluate BBK32-C binding to zymogen-stabilized

forms of C1r-CCP2-SP, SPR experiments were performed by capturing His-C1r-CCP2-SP, His-

C1r-CCP2-SP-R463Q (R463Q), or His-C1r-CCP2-SP-S654A (S654A) on a Ni-NTA sensor chip.

BBK32-C was sequentially injected over each surface in a concentration series (1.0, 3.2, 10.0,

31.6, 100 nM). BBK32-C sensorgrams are reproduced from **Fig. 2B** for comparison. Arrows

indicate injection phases of each C1r-CCP2-SP concentration.

Figure 4. BBK32 R248 and K327 are critical for complement inhibitory activity of BBK32.

A) ELISA-based complement activation assays were performed to determine the ability of

BBK32-C mutants to inhibit the classical pathway of complement *in vitro*. A two-fold

concentration series of each BBK32-C mutant was used and performed in duplicate. Inhibitory

data for additional site-directed mutants are presented in **Figure S6**. IC$_{50}$ values and 95%

confidence intervals associated with a normalized four-parameter dose-inhibition fitting procedure

are shown in **Table S2**. **B)** Western blots of lysates from *B. burgdorferi* strains B314 containing

distinct *bbk32* alleles were probed with monoclonal antibodies to BBK32 (α-BBK32) or FlaB (a-

FlaB), the latter as a loading control. Vector refers to the plasmid only backbone sample

(B314/pBBE22*luc*), BBK32 is the wildtype *bbk32* construct (B314/pCD100), R248A is the *bbk32*-

R248A isolate (B314/pAP5), K327A is the *bbk32*-K327A isolate (B314/pAP6), and R248A-

K327A is the *bbk32*-R248A-K327A double mutant (B314/pAP7). Numbers on the left refer to the

molecular mass of markers (in kDa). **C)** BBK32 mutant proteins were tested for their ability to

confer resistance to normal human serum (NHS). The serum sensitive strain B314 was transformed

separately with vector only (black), wildtype *bbk32* (blue), *bbk32* R248A (green), *bbk32* K327A

(purple), and *bbk32* R248A-K327A (red) and incubated with NHS. Sensitivity was scored as a

ratio of the affected cells relative to the total cells viewed via darkfield microscopy. Heat

inactivated NHS (hiNHS) was used as a control (right). P-values between *bbk32* samples are

indicated above. Statistical significance was assessed using a two-way ANOVA.

Figure 5. Comparison of BBK32 to gigastasin reveals convergent mechanisms of action. A)

Ribbon diagrams of BBK32-C (PDB:6N1L), C1r-CCP2-SP (PDB:1MD8), Gigastasin/C1s-CCP2-

742   SP (PDB: 5UBM) with colored interactions within 4 Å, as judged by the program CCP4i

743   CONTACTS. The BBK32-C/C1r-CCP2-SP interactions are shown highlighted in red and

744   C1s/Gigastasin loop interactions in blue. **B)** Sequence alignment of serine protease domains of C1r

745   and C1s loops are designated with bars and are according to C1r structure.

747 **Table 1. MALS and SAXS parameters**

Sample	Maximal dimension D_{max} (Å)	R_g (Å)	MW_{seq} (kDa)	MW_{SAXS} (kDa)	MW_{MALS} (kDa)	SAXS model fit χ^2
BBK32-C	~75	21.3 ± 0.9	16.9	19.9 ± 1.58	~19	2.08
C1r-CCP2-SP	~80	24.0 ± 0.9	37.7	31.7 ± 1.75	~40-60	1.06
BBK32-C/ C1r-CCP2-SP Complex	~85	27.8 ± 2.2	54.6	55.6 ± 4.13	~55-65	1.24

References

1. N. S. Merle, S. E. Church, V. Fremeaux-Bacchi, L. T. Roumenina, Complement System Part I - Molecular Mechanisms of Activation and Regulation. *Front. Immunol.* **6**, 262 (2015).

2. N. S. Merle, R. Noe, L. Halbwachs-Mecarelli, V. Fremeaux-Bacchi, L. T. Roumenina, Complement System Part II: Role in Immunity. *Front. Immunol.* **6**, 257 (2015).

3. M. J. Walport, Complement. First of two parts. *N. Engl. J. Med.* **344**, 1058–1066 (2001).

4. M. J. Walport, Complement. Second of two parts. *N. Engl. J. Med.* **344**, 1140–1144 (2001).

5. R. J. Ziccardi, Activation of the early components of the classical complement pathway under physiologic conditions. *J. Immunol. (Baltimore, Md. 1950)*. **126**, 1769–1773 (1981).

6. U. Deniz, C. H. Stuart, de Kreuk Bart-Jan, I. K. Roman, N. de Jong Rob, J. B. Frank, S. Janine, J. K. Abraham, H. S. Thomas, W. H. I. Parren Paul, G. Piet, Structures of C1-IgG1 provide insights into how danger pattern recognition activates complement. *Science (80-.)*. **359**, 794–797 (2018).

7. T. H. Sharp, A. L. Boyle, C. A. Diebolder, A. Kros, A. J. Koster, P. Gros, Insights into IgM-mediated complement activation based on in situ structures of IgM-C1-C4b. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 11900–11905 (2019).

8. C. A. Diebolder, F. J. Beurskens, R. N. de Jong, R. I. Koning, K. Strumane, M. A. Lindorfer, M. Voorhorst, D. Ugurlar, S. Rosati, A. J. R. Heck, J. G. J. van de Winkel, I. A. Wilson, A. J. Koster, R. P. Taylor, E. O. Saphire, D. R. Burton, J. Schuurman, P. Gros, P. W. H. I. Parren, Complement is activated by IgG hexamers assembled at the cell surface. *Science*. **343**, 1260–1263 (2014).

9. T. R. Kjaer, S. Thiel, G. R. Andersen, Toward a structure-based comprehension of the lectin pathway of complement. *Mol. Immunol.* **56** (2013), pp. 413–422.

10. J. Dobó, A. Kocsis, P. Gál, Be on Target: Strategies of Targeting Alternative and Lectin Pathway Components in Complement-Mediated Diseases. *Front. Immunol.* **9**, 1851 (2018).

11. M. K. Pangburn, H. J. Müller-Eberhard, The alternative pathway of complement. *Springer Semin. Immunopathol.* **7**, 163–192 (1984).

12. C. Q. Schmidt, J. D. Lambris, D. Ricklin, Protection of host cells by complement regulators. *Immunol. Rev.* **274**, 152–171 (2016).

13. D. Ricklin, J. D. Lambris, Complement in immune and inflammatory disorders: pathophysiological mechanisms. *J. Immunol. (Baltimore, Md. 1950)*. **190**, 3831–3838 (2013).

14. D. Ricklin, D. C. Mastellos, E. S. Reis, J. D. Lambris, The renaissance of complement therapeutics. *Nat. Rev. Nephrol.* **14**, 26–47 (2018).

15. J. D. Lambris, D. Ricklin, B. V Geisbrecht, Complement evasion by human pathogens.
Nat. Rev. Microbiol. **6**, 132–142 (2008).

16. P. Kraiczy, Hide and Seek: How Lyme Disease Spirochetes Overcome Complement
Attack. *Front. Immunol.* **7**, 385 (2016).

17. J. T. Skare, B. L. Garcia, Complement Evasion by Lyme Disease Spirochetes. *Trends*
Microbiol. (2020), , doi:10.1016/j.tim.2020.05.004.

18. M. Adeolu, R. S. Gupta, A phylogenomic and molecular marker based proposal for the
division of the genus Borrelia into two genera: the emended genus Borrelia containing
only the members of the relapsing fever Borrelia, and the genus Borreliella gen. nov.
containing the members o. *Antonie Van Leeuwenhoek.* **105**, 1049–1072 (2014).

19. P. Kraiczy, J. Hellwage, C. Skerka, H. Becker, M. Kirschfink, M. M. Simon, V. Brade, P.
F. Zipfel, R. Wallich, Complement resistance of Borrelia burgdorferi correlates with the
expression of BbCRASP-1, a novel linear plasmid-encoded surface protein that interacts
with human factor H and FHL-1 and is unrelated to Erp proteins. *J. Biol. Chem.* **279**,
2421–2429 (2004).

20. K. Hartmann, C. Corvey, C. Skerka, M. Kirschfink, M. Karas, V. Brade, J. C. Miller, B.
Stevenson, R. Wallich, P. F. Zipfel, P. Kraiczy, Functional characterization of BbCRASP-
2, a distinct outer membrane protein of Borrelia burgdorferi that binds host complement
regulators factor H and FHL-1. *Mol. Microbiol.* **61**, 1220–1236 (2006).

21. B. Stevenson, K. Tilly, P. A. Rosa, A family of genes located on four separate 32-kilobase
circular plasmids in Borrelia burgdorferi B31. *J. Bacteriol.* **178**, 3508–3516 (1996).

22. B. Stevenson, N. El-Hage, M. A. Hines, J. C. Miller, K. Babb, Differential binding of host
complement inhibitor factor H by Borrelia burgdorferi Erp surface proteins: A possible
mechanism underlying the expansive host range of Lyme disease spirochetes. *Infect.*
Immun. (2002), doi:10.1128/IAI.70.2.491-497.2002.

23. J. V McDowell, J. Wolfgang, E. Tran, M. S. Metts, D. Hamilton, R. T. Marconi,
Comprehensive analysis of the factor h binding capabilities of borrelia species associated
with lyme disease: delineation of two distinct classes of factor h binding proteins. *Infect.*
Immun. **71**, 3597–3602 (2003).

24. J. V McDowell, K. M. Hovis, H. Zhang, E. Tran, J. Lankford, R. T. Marconi, Evidence
that the BBA68 protein (BbCRASP-1) of the Lyme disease spirochetes does not
contribute to factor H-mediated immune evasion in humans and other animals. *Infect.*
Immun. **74**, 3030–3034 (2006).

25. T. Bykowski, M. E. Woodman, A. E. Cooley, C. A. Brissette, R. Wallich, V. Brade, P.
Kraiczy, B. Stevenson, Borrelia burgdorferi complement regulator-acquiring surface
proteins (BbCRASPs): Expression patterns during the mammal-tick infection cycle. *Int. J.*
Med. Microbiol. IJMM. **298 Suppl**, 249–256 (2008).

26. M. R. Kenedy, S. R. Vuppala, C. Siegel, P. Kraiczy, D. R. Akins, CspA-mediated binding
of human factor H inhibits complement deposition and confers serum resistance in
Borrelia burgdorferi. *Infect. Immun.* **77**, 2773–2782 (2009).

27. E. A. Rogers, S. V Abdunnur, J. V McDowell, R. T. Marconi, Comparative analysis of the
properties and ligand binding characteristics of CspZ, a factor H binding protein, derived
from Borrelia burgdorferi isolates of human origin. *Infect. Immun.* **77**, 4396–4405 (2009).

28. M. R. Kenedy, D. R. Akins, The OspE-related proteins inhibit complement deposition and
enhance serum resistance of Borrelia burgdorferi, the lyme disease spirochete. *Infect.*
Immun. **79**, 1451–1457 (2011).

29. C. Hammerschmidt, Y. Klevenhaus, A. Koenigs, T. Hallström, V. Fingerle, C. Skerka, K.
M. Pos, P. F. Zipfel, R. Wallich, P. Kraiczy, BGA66 and BGA71 facilitate complement
resistance of Borrelia bavariensis by inhibiting assembly of the membrane attack complex.
Mol. Microbiol. **99**, 407–424 (2016).

30. T. Hallström, C. Siegel, M. Mörgelin, P. Kraiczy, C. Skerka, P. F. Zipfel, CspA from
Borrelia burgdorferi inhibits the terminal complement pathway. *MBio*. **4**, 13 (2013).

31. M. Pausa, V. Pellis, M. Cinco, P. G. Giulianini, G. Presani, S. Perticarari, R. Murgia, F.
Tedesco, Serum-resistant strains of Borrelia burgdorferi evade complement-mediated
killing by expressing a CD59-like complement inhibitory molecule. *J. Immunol.*
(Baltimore, Md. 1950). **170**, 3214–3222 (2003).

32. J. A. Caine, Y.-P. Lin, J. R. Kessler, H. Sato, J. M. Leong, J. Coburn, Borrelia burgdorferi
outer surface protein C (OspC) binds complement component C4b and confers
bloodstream survival. *Cell. Microbiol.* **19** (2017), doi:10.1111/cmi.12786.

33. B. L. Garcia, H. Zhi, B. Wager, M. Höök, J. T. Skare, Borrelia burgdorferi BBK32
Inhibits the Classical Pathway by Blocking Activation of the C1 Complement Complex.
PLoS Pathog. **12**, e1005404 (2016).

34. M. He, B. K. Boardman, D. Yan, X. F. Yang, Regulation of expression of the fibronectin-
binding protein BBK32 in Borrelia burgdorferi. *J. Bacteriol.* **189**, 8377–8380 (2007).

35. J. Seshu, M. D. Esteve-Gassent, M. Labandeira-Rey, J. H. Kim, J. P. Trzeciakowski, M.
Höök, J. T. Skare, Inactivation of the fibronectin-binding adhesin gene bbk32 significantly
attenuates the infectivity potential of Borrelia burgdorferi. *Mol. Microbiol.* **59**, 1591–1601
(2006).

36. J. A. Hyde, E. H. Weening, M. Chang, J. P. Trzeciakowski, M. Höök, J. D. Cirillo, J. T.
Skare, Bioluminescent imaging of Borrelia burgdorferi in vivo demonstrates that the
fibronectin-binding protein BBK32 is required for optimal infectivity. *Mol. Microbiol.* **82**,
99–113 (2011).

37. W. S. Probert, J. H. Kim, M. Höök, B. J. Johnson, Mapping the ligand-binding region of
Borrelia burgdorferi fibronectin-binding protein BBK32. *Infect. Immun.* **69**, 4129–4133
(2001).

38. J. R. Fischer, K. T. LeBlanc, J. M. Leong, Fibronectin binding protein BBK32 of the
Lyme disease spirochete promotes bacterial attachment to glycosaminoglycans. *Infect.*
Immun. **74**, 435–441 (2006).

39. T. J. Moriarty, M. Shi, Y.-P. Lin, R. Ebady, H. Zhou, T. Odisho, P.-O. Hardy, A. Salman-
Dilgimen, J. Wu, E. H. Weening, J. T. Skare, P. Kubes, J. Leong, G. Chaconas, Vascular

binding of a pathogen under shear force through mechanistically distinct sequential
interactions with host macromolecules. *Mol. Microbiol.* **86**, 1116–1131 (2012).

40. R. Ebady, A. F. Niddam, A. E. Boczula, Y. R. Kim, N. Gupta, T. T. Tang, T. Odisho, H.
Zhi, C. A. Simmons, J. T. Skare, T. J. Moriarty, Biomechanics of Borrelia burgdorferi
Vascular Interactions. *Cell Rep.* **16**, 2593–2604 (2016).

41. J. Xie, H. Zhi, R. J. Garrigues, A. Keightley, B. L. Garcia, J. T. Skare, Structural
determination of the complement inhibitory domain of Borrelia burgdorferi BBK32
provides insight into classical pathway complement evasion by Lyme disease spirochetes.
PLoS Pathog. **15**, e1007659 (2019).

42. R. P. Rambo, J. A. Tainer, Accurate assessment of mass, models and resolution by small-
angle scattering. *Nature* (2013), doi:10.1038/nature12070.

43. M. Budayova-Spano, W. Grabarse, N. M. Thielens, H. Hillen, M. Lacroix, M. Schmidt, J.
C. Fontecilla-Camps, G. J. Arlaud, C. Gaboriaud, Monomeric structures of the zymogen
and active catalytic domain of complement protease c1r: further insights into the c1
activation mechanism. *Struct. (London, Engl. 1993)*. **10**, 1509–1519 (2002).

44. B. Webb, A. Sali, *Curr. Protoc. Bioinforma.*, in press,
doi:10.1002/0471250953.bi0506s47.

45. E. F. Pettersen, T. D. Goddard, C. C. Huang, G. S. Couch, D. M. Greenblatt, E. C. Meng,
T. E. Ferrin, UCSF Chimera - A visualization system for exploratory research and
analysis. *J. Comput. Chem.* **25**, 1605–1612 (2004).

46. D. Schneidman-Duhovny, M. Hammel, J. A. Tainer, A. Sali, Accurate SAXS profile
computation and its assessment by contrast variation experiments. *Biophys. J.* **105**, 962–
974 (2013).

47. D. Schneidman-Duhovny, M. Hammel, J. A. Tainer, A. Sali, *FoXS, FoXSDock and*
MultiFoXS: Single-state and multi-state structural modeling of proteins and their
complexes based on SAXS profiles (2016;
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4987932/), vol. 44.

48. D. Schneidman-Duhovny, M. Hammel, A. Sali, Macromolecular Docking Restrained by a
Small Angle X-Ray Scattering Profile (2010), doi:10.1016/j.jsb.2010.09.023.

49. D. Franke, D. I. Svergun, DAMMIF, a program for rapid ab-initio shape determination in
small-angle scattering. *J. Appl. Crystallogr.* (2009), doi:10.1107/S0021889809000338.

50. V. V. Volkov, D. I. Svergun, in *Journal of Applied Crystallography* (2003).

51. M. Pelikan, G. L. Hura, M. Hammel, Structure and flexibility within proteins as identified
through small angle X-ray scattering. *Gen. Physiol. Biophys.* (2009),
doi:10.4149/gpb_2009_02_174.

52. M. D. Winn, C. C. Ballard, K. D. Cowtan, E. J. Dodson, P. Emsley, P. R. Evans, R. M.
Keegan, E. B. Krissinel, A. G. W. Leslie, A. McCoy, S. J. McNicholas, G. N. Murshudov,
N. S. Pannu, E. A. Potterton, H. R. Powell, R. J. Read, A. Vagin, K. S. Wilson, Overview
of the CCP4 suite and current developments. *Acta Crystallogr. Sect. D Biol. Crystallogr.*

(2011), , doi:10.1107/S0907444910045749.

53.   E. Krissinel, K. Henrick, Inference of Macromolecular Assemblies from Crystalline State.
J. Mol. Biol. **372**, 774–797 (2007).

54.   R. Karlsson, P. S. Katsamba, H. Nordin, E. Pol, D. G. Myszka, Analyzing a kinetic
titration series using affinity biosensors. *Anal. Biochem.* **349**, 136–147 (2006).

55.   J. Kardos, P. Gal, L. Szilagyi, N. M. Thielens, K. Szilagyi, Z. Lorincz, P. Kulcsar, L. Graf,
G. J. Arlaud, P. Zavodszky, The Role of the Individual Domains in the Structure and
Function of the Catalytic Region of a Modular Serine Protease, C1r. *J. Immunol.* **167**,
5202–5208 (2001).

56.   A. Sadziene, B. Wilske, M. S. Ferdows, A. G. Barbour, The cryptic ospC gene of Borrelia
burgdorferi B31 is located on a circular plasmid. *Infect. Immun.* **61**, 2192–2195 (1993).

57.   A. Sadziene, D. D. Thomas, A. G. Barbour, Borrelia burgdorferi mutant lacking Osp:
biological and immunological characterization. *Infect. Immun.* **63**, 1573–1580 (1995).

58.   S. S. Pang, L. C. Wijeyewickrema, L. Hor, S. Tan, E. Lameignere, E. M. Conway, A. M.
Blom, F. C. Mohlin, X. Liu, R. J. Payne, J. C. Whisstock, R. N. Pike, The Structural Basis
for Complement Inhibition by Gigastasin, a Protease Inhibitor from the Giant Amazon
Leech. *J. Immunol. (Baltimore, Md. 1950)*. **199**, 3883–3891 (2017).

59.   J. L. Schmitz, S. D. Lovrich, S. M. Callister, R. F. Schell, Depletion of complement and
effects on passive transfer of resistance to infection with Borrelia burgdorferi. *Infect.*
Immun. **59**, 3815–3818 (1991).

60.   S. W. Barthold, L. K. Bockenstedt, Passive immunizing activity of sera from mice
infected with Borrelia burgdorferi. *Infect. Immun.* **61**, 4696–4702 (1993).

61.   E. Fikrig, L. K. Bockenstedt, S. W. Barthold, M. Chen, H. Tao, P. A. Salaam, S. R.
Telford, R. A. Flavell, Sera from patients with chronic lyme disease protect mice from
lyme borreliosis. *J. Infect. Dis.* **169**, 568–574 (1994).

62.   M. P. Reichhardt, S. Johnson, T. Tang, T. Morgan, N. Tebeka, N. Popitsch, J. C. Deme,
M. M. Jore, S. M. Lea, An inhibitor of complement C5 provides structural insights into
activation. *Proc. Natl. Acad. Sci. U. S. A.* (2020), doi:10.1073/pnas.1909973116.

63.   K. Tyson, C. Elkins, H. Patterson, E. Fikrig, A. de Silva, Biochemical and functional
characterization of Salp20, an Ixodes scapularis tick salivary protein that inhibits the
complement pathway. *Insect Mol. Biol.* **16**, 469–479 (2007).

64.   T. J. Schuijt, J. W. R. Hovius, N. D. Van Burgel, N. Ramamoorthi, E. Fikrig, A. P. Van
Dam, The tick salivary protein Salp15 inhibits the killing of serum-sensitive Borrelia
burgdorferi sensu lato isolates. *Infect. Immun.* (2008), doi:10.1128/IAI.00232-08.

65.   N. C. S. Silva, V. F. Vale, P. F. Franco, N. F. Gontijo, J. G. Valenzuela, M. H. Pereira, M.
R. V. Sant'Anna, D. S. Rodrigues, W. S. Lima, B. Fux, R. N. Araujo, Saliva of
Rhipicephalus (Boophilus) microplus (Acari: Ixodidae) inhibits classical and alternative
complement pathways. *Parasites and Vectors* (2016), doi:10.1186/s13071-016-1726-8.

66. T. J. Schuijt, J. Coumou, S. Narasimhan, J. Dai, K. Deponte, D. Wouters, M. Brouwer, A.
Oei, J. J. T. H. Roelofs, A. P. van Dam, T. van der Poll, C. V. Veer, J. W. Hovius, E.
Fikrig, A tick mannose-binding lectin inhibitor interferes with the vertebrate complement
cascade to enhance transmission of the lyme disease agent. *Cell Host Microbe*. **10**, 136–
146 (2011).

67. V. P. Ferreira, V. Fazito Vale, M. K. Pangburn, M. Abdeladhim, A. Ferreira Mendes-
Sousa, I. V. Coutinho-Abreu, M. Rasouli, E. A. Brandt, C. Meneses, K. F. Lima, R.
Nascimento Araújo, M. Horácio Pereira, M. Kotsyfakis, F. Oliveira, S. Kamhawi, J. M. C.
Ribeiro, N. F. Gontijo, N. Collin, J. G. Valenzuela, SALO, a novel classical pathway
complement inhibitor from saliva of the sand fly Lutzomyia longipalpis. *Sci. Rep.* (2016),
doi:10.1038/srep19300.

68. A. F. Mendes-Sousa, D. C. Queiroz, V. F. Vale, J. M. C. Ribeiro, J. G. Valenzuela, N. F.
Gontijo, J. F. Andersen, An Inhibitor of the Alternative Pathway of Complement in Saliva
of New World Anopheline Mosquitoes. *J. Immunol.* (2016),
doi:10.4049/jimmunol.1600020.

69. B. L. Garcia, K. X. Ramyar, D. Ricklin, J. D. Lambris, B. V Geisbrecht, Advances in
understanding the structure, function, and mechanism of the SCIN and Efb families of
Staphylococcal immune evasion proteins. *Adv. Exp. Med. Biol.* **946**, 113 (2012).

70. B. L. Garcia, S. A. Zwarthoff, S. H. M. Rooijakkers, B. V Geisbrecht, Novel Evasion
Mechanisms of the Classical Complement Pathway. *J. Immunol.* **197**, 2051–2060 (2016).

71. C. Dunwiddie, N. A. Thornberry, H. G. Bull, M. Sardana, P. A. Friedman, J. W. Jacobs,
E. Simpson, Antistasin, a leech-derived inhibitor of factor Xa. Kinetic analysis of enzyme
inhibition and identification of the reactive site. *J. Biol. Chem.* **264**, 16694–16699 (1989).

72. P. Kraiczy, Travelling between Two Worlds: Complement as a Gatekeeper for an
Expanded Host Range of Lyme Disease Spirochetes. *Vet. Sci.* **3** (2016),
doi:10.3390/vetsci3020012.

73. D. S. Samuels, D. Drecktrah, L. S. Hall, Genetic Transformation and Complementation.
Methods Mol. Biol. **1690**, 183–200 (2018).

74. J. A. Hyde, E. H. Weening, J. T. Skare, Genetic transformation of Borrelia burgdorferi.
Curr. Protoc. Microbiol. **Chapter 12** (2011), doi:10.1002/9780471729259.mc12c04s20.

75. B. V Geisbrecht, S. Bouyain, M. Pop, An optimized system for expression and
purification of secreted bacterial proteins. *Protein Expr. Purif.* **46**, 23–32 (2006).

76. B. R. Rushing, D. L. Rohlik, S. Roy, D. A. Skaff, B. L. Garcia, Targeting the Initiator
Protease of the Classical Pathway of Complement Using Fragment-Based Drug
Discovery. *Molecules* (2020), doi:10.3390/molecules25174016.

77. K. N. Dyer, M. Hammel, R. P. Rambo, S. E. Tsutakawa, I. Rodic, S. Classen, J. A. Tainer,
G. L. Hura, High-throughput SAXS for the characterization of biomolecules in solution: A
practical approach. *Methods Mol. Biol.* (2014), doi:10.1007/978-1-62703-691-7_18.

78. G. L. Hura, A. L. Menon, M. Hammel, R. P. Rambo, F. L. Poole, S. E. Tsutakawa, F. E.
Jenney, S. Classen, K. A. Frankel, R. C. Hopkins, S. J. Yang, J. W. Scott, B. D. Dillard,

M. W. W. Adams, J. A. Tainer, Robust, high-throughput solution structural analyses by
small angle X-ray scattering (SAXS). *Nat. Methods* (2009), doi:10.1038/nmeth.1353.

79.   P. V. Konarev, V. V. Volkov, A. V. Sokolova, M. H. J. Koch, D. I. Svergun, PRIMUS: A
Windows PC-based system for small-angle scattering data analysis. *J. Appl. Crystallogr.*
(2003), doi:10.1107/S0021889803012779.

80.   D. I. Svergun, Determination of the regularization parameter in indirect-transform
methods using perceptual criteria. *J. Appl. Crystallogr.* (1992),
doi:10.1107/S0021889892001663.

81.   P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of Coot. *Acta*
Crystallogr. D. Biol. Crystallogr. **66**, 486–501 (2010).

82.   G. Q. Dong, H. Fan, D. Schneidman-Duhovny, B. Webb, A. Sali, Structural
bioinformatics Optimized atomic statistical potentials: assessment of protein interfaces
and loops. **29**, 3158–3166 (2013).

83.   K. Manalastas-Cantos, P. V. Konarev, N. R. Hajizadeh, A. G. Kikhney, M. V. Petoukhov,
D. S. Molodenskiy, A. Panjkovich, H. D. T. Mertens, A. Gruzinov, C. Borges, C. M.
Jeffries, D. I. Svergun, D. Franke, ATSAS 3.0 : expanded functionality and new tools for
small-angle scattering data analysis . *J. Appl. Crystallogr.* (2021),
doi:10.1107/s1600576720013412.

84.   Z. Otwinowski, W. Minor, (Academic Press, 1997;
http://www.sciencedirect.com/science/article/pii/S007668799776066X), vol. 276 of
Methods in Enzymology, pp. 307–326.

85.   P. D. Adams, P. V Afonine, G. Bunkóczi, V. B. Chen, I. W. Davis, N. Echols, J. J. Headd,
L.-W. Hung, G. J. Kapral, R. W. Grosse-Kunstleve, A. J. McCoy, N. W. Moriarty, R.
Oeffner, R. J. Read, D. C. Richardson, J. S. Richardson, T. C. Terwilliger, P. H. Zwart,
PHENIX: a comprehensive Python-based system for macromolecular structure solution.
Acta Crystallogr. D. Biol. Crystallogr. **66**, 213–221 (2010).

86.   J. Kardos, V. Harmat, A. Palló, O. Barabás, K. Szilágyi, L. Gráf, G. Náray-Szabó, Y.
Goto, P. Závodszky, P. Gál, Revisiting the mechanism of the autoactivation of the
complement protease C1r in the C1 complex: Structure of the active catalytic region of
C1r. *Mol. Immunol.* **45**, 1752–1760 (2008).

87.   A. Roos, L. H. Bouwman, J. Munoz, T. Zuiverloon, M. C. Faber-Krol, F. C. den Houten,
N. Klar-Mohamad, C. E. Hack, M. G. Tilanus, M. R. Daha, Functional characterization of
the lectin pathway of complement in human serum. *Mol. Immunol.* **39**, 655–668 (2003).

88.   I. Schechter, A. Berger, On the size of the active site in proteases. I. Papain. *Biochem.*
Biophys. Res. Commun. (1967), doi:10.1016/S0006-291X(67)80055-X.

1018