A prioritized and validated resource of mitochondrial proteins in *Plasmodium* identifies leads to unique biology

Selma L. van Esveld[1,2], Lisette Meerstein-Kessel[1,3,‡], Cas Boshoven[4,‡], Jochem F. Baaij[1], Konstantin Barylyuk[5], Jordy P. M. Coolen[4], Ronald A.J. Duim[1], Bas E. Dutilh[1,6], Daniel R. Garza[1,7], Marijn Letterie[4], Nicholas I. Proellochs[4], Michelle N. de Ridder[4], Prashanna Balaji Venkatasubramanian[1], Laura E. de Vries[4], Ross F. Waller[5], Taco W.A. Kooij[4], Martijn A. Huynen[1,2]

[1]*Center for Molecular and Biomolecular Informatics, Radboud Institute for Molecular Life Sciences, Radboudumc, Nijmegen, the Netherlands*
[2]*Radboud Center for Mitochondrial Medicine, Radboudumc, Nijmegen, the Netherlands*
[3]*Radboud Institute for Health Sciences, Radboudumc, Nijmegen, the Netherlands.*
[4]*Department of Medical Microbiology, Radboudumc Center for Infectious Diseases, Radboud Institute for Molecular Life Sciences, Radboudumc, Nijmegen, the Netherlands*
[5]*Department of Biochemistry, University of Cambridge, Cambridge, UK*
[6]*Theoretical Biology and Bioinformatics, Science for Life, Utrecht University, Utrecht, the Netherlands*
[7]*Laboratory of Molecular Bacteriology (Rega Institute), Department of Microbiology, Immunology and Transplantation, KU Leuven, Leuven, Belgium*

‡*These authors contributed equally to this work and should be considered co-second authors.*

Abstract

Plasmodium species have a single mitochondrion that is essential for their survival and has been successfully targeted by anti-malarial drugs. Most proteins are imported into this organelle and our picture of the *Plasmodium* mitochondrial proteome remains incomplete. Many data sources contain information about mitochondrial localization, including proteome and gene expression profiles, orthology to mitochondrial proteins from other species, co-evolutionary relationships, and amino acid sequences, each with different coverage and reliability. To obtain a comprehensive, prioritized list of *Plasmodium falciparum* mitochondrial proteins, we rigorously analyzed and integrated eight datasets using Bayesian statistics into a predictive score per protein for mitochondrial localization. At a corrected false discovery rate of 25%, we identified 295 proteins with a sensitivity of 65% and a specificity of 98%. They include proteins that have not been identified as mitochondrial in other eukaryotes but have characterized homologs in bacteria that are involved in metabolism or translation. Mitochondrial localization of seven *Plasmodium berghei* orthologs was confirmed by epitope labeling and co-localization with a mitochondrial marker protein. One of these belongs to a newly identified apicomplexan mitochondrial protein family that in *P. falciparum* has four members. With the experimentally validated mitochondrial proteins and the complete ranked *P. falciparum* proteome, which we have named PlasmoMitoCarta, we present a resource to study unique proteins of *Plasmodium* mitochondria.

Introduction

Although all mitochondria evolved from a single endosymbiotic event (1), they display a large variety among the eukaryotes. For example, while in most species the mitochondrial organelle still retains its genome, the number of proteins encoded varies from 100 in *Andalucia godoyi* (2) to three in myzozoan species like *Plasmodium falciparum* that only encode cytochrome b, cytochrome c oxidase subunits 1 and 3, and two fragmented rRNAs (3). Moreover, mitochondria-like organelles of a variety of anaerobic species have lost the

organellar genome altogether (4). Also, mitochondrial proteomes vary broadly in size, ranging from e.g. ~1500 different proteins in mammalian mitochondria (5) to a few members of only a single biochemical pathway found in some mitosomes (6). Within this variety, the *P. falciparum* mitochondrion, based on the size of its genome, the size of the nuclear genome, and the complexity of its oxidative phosphorylation, is predicted to have a relatively small proteome. Although several mitochondrial pathways have already been identified in *P. falciparum* (7), and databases like PlasmoDB contain valuable information about mitochondrial localization of individual proteins (8), a complete resource that weighs and combines all the relevant information about possible mitochondrial localization is lacking.

Plasmodium mitochondria are particularly interesting because their function is variable between different life-cycle stages. In the sexual blood stage, functional oxidative phosphorylation is essential for colonization and development inside the mosquito hosts (9, 10), while in asexual blood stages the only essential function of the respiratory chain is to recycle ubiquinone for pyrimidine biosynthesis (11). This is further highlighted by the observation that in asexual blood-stage parasites no cristae are detectable in the mitochondrial membrane, while in gametocyte mitochondria cristae-like structures that typically accumulate respiratory chain complexes are observed (10). A recent complexome profiling study demonstrated that the level of complex V components in asexual blood-stage parasites was only 3% of the level in gametocytes (12). Another intriguing aspect of mitochondria in *Plasmodium* and closely related species is that they are the only mitochondria that have no detectable homolog of the FeS cluster assembly protein frataxin (13), a protein that occurs even in mitosomes (14).

Knowledge on the mitochondrial proteome could provide viable targets for the development of new drugs against this deadly parasite, as apicomplexan mitochondrial proteomes contain unique proteins that are not present in the animal hosts (15). Recently for example, an unusual prohibitin-like protein, PHBL, was identified in *Plasmodium berghei* that might provide a transmission-blocking drug target as it is unique to Myzozoa and *phbl⁻* parasites fail to transmit (16). Furthermore, respiratory protein complex III has proven to be a good drug target (17) and inhibitors to the mitochondrial enzymes dihydroorotate dehydrogenase (DHODH) and NADH type II oxidoreductase (PfNDH2) have been identified (18).

For years, it has been difficult to perform proteomics experiments to identify the *Plasmodium* mitochondrial proteome due to the lack of organelle isolation methods that reliably separate mitochondria from the apicoplast (19). The definition of mitochondrial functions relied heavily upon homology studies (7, 20) and microscopic examination of individual proteins (Supplemental Table 1). With the use of two biotin tagging approaches, 422 mitochondrial matrix proteins were identified in *Toxoplasma gondii*, a species related to *Plasmodium* that also has an apicoplast and a mitochondrion (15). Besides these mitochondria-targeted approaches, hyperLOPIT, a whole-cell biochemical fractionation technique, defined the subcellular localization of the *T. gondii* proteome. This provided among others a set of 220 soluble and 168 membrane-bound *T. gondii* mitochondrial proteins (21). Alongside these proteomic data, there is a wealth of other data sources providing information on mitochondrial localization in *P. falciparum*. The challenge remains to use and combine these heterogeneous datasets to reliably predict the mitochondrial proteome.

Here, we integrated eight datasets containing proteomic, gene expression, orthology to mitochondrial proteins from other species, phylogenetic distribution, and

amino acid sequence data in a Bayesian manner as has been applied before such as for the
definition of the mitochondrial proteome of humans (5, 22-25). We used a naïve Bayesian
classifier to combine the data. This method exploits the relative strengths of the various
datasets while maintaining transparency about the contribution of each dataset to the
final Bayesian posterior probability score for being mitochondrial. For each protein, the
contribution of each dataset to the prediction was determined using gold standards of
Plasmodium proteins that are either known to be mitochondrial or non-mitochondrial.
After data integration, we obtained a list of 295 P. falciparum mitochondrial proteins. We
experimentally validated these predictions with seven proteins of varying probability
scores, and collectively this predicted proteome indicates that the P. falciparum
mitochondria are unique relative to mitochondria of model organisms.
110
111 **Materials and Methods**
112
Animal experiments. All animal experiments were performed in accordance with the
Dutch Experiments on Animals Act (Wod), the Directive 2010/63/EU from the European
Union and the European ETS 123 convention and approved by the Radboud University
Animal Welfare Body (IvD) and Animal Experiment Committee (RUDEC; 2015-0142), and
the Central Authority for Scientific Procedures on Animals (CCD; AVD103002016424). In
this study, we used outbred male and female NMRI mice (Envigo).
119
Nuclear encoded reference proteome. All datasets used in this study were mapped to
the *P. falciparum* 3D7 reference proteome version 3.1 from the Sanger Institute,
downloaded December 2017. This version, including isoforms, contains 5431 proteins
encoded by 5357 genes. As we are interested in the proteins encoded by nuclear genes,
all apicoplast and mitochondrial encoded proteins were excluded, leaving a reference of
5324 genes. When a dataset contains informational data points for two or more protein
isoforms of one gene, the value chosen for that gene was the one that comes closest to
the expected values for mitochondrial proteins in that specific dataset.
128
Assembly of positive and negative lists for benchmarking. Bayesian data integration
depends on gold standards of proteins known to be mitochondrial or non-mitochondrial.
We constructed two gold standards to assess the predictive value of the individual
datasets. Furthermore, we built two alternative sets for the construction of the CLIME (co-
evolution) and WICCA (co-expression) datasets to prevent that the standards chosen to
train those two predictors were also used to evaluate them. These four sets are unique,
have no overlapping genes, and are available in Supplemental Table 1. To construct these
lists, we commenced with a systematic review of all literature available via PubMed (as per
30-08-2018) using the following broad search string "(mitochondri* OR apicoplast OR
plastid) AND (plasmodium OR malaria)". Given the intimate functional and physical
relation between mitochondrion and apicoplast, we reasoned that the positive gold
standard list should only include genes encoding proteins that had been unambiguously
and singularly assigned to the mitochondrion through fluorescence microscopy and co-
localization with confirmed mitochondrial markers or via immune-electron microscopy (54
genes). Any proteins for which multiple studies showed different results were excluded.
To facilitate discrimination from the apicoplast, we also made a gold standard apicoplast
list (72 genes). Finally, all proteins that were unambiguously non-mitochondrial as
demonstrated in the same papers (170 genes) were combined with the apicoplast-positive

list to form the basis for the negative gold standard. This negative set was then complemented with an additional 346 non-overlapping genes encoding non-apicoplast and non-mitochondrial proteins identified through extensive literature review and published online by the Ralph lab (26). The 588 genes encoding non-mitochondrial proteins were randomly assigned to the negative gold standard (294 genes) that was used for the Bayesian data integration and the alternative negative gold standard (294 genes) that was used to train the CLIME and WICCA approaches. For the latter purpose, an alternative positive standard was compiled of 146 genes, non-overlapping with the first gold standard, but associated with one or more mitochondrial GO-terms (GO:0000275, GO:0005739-43, GO:0005746, GO:0005750, GO:0005753, GO:0005758, GO:0005759, GO:0006122, GO:0006839, GO:0006850, GO:0031966, GO:0033108, GO:0042775, GO:0044429, GO:0044455).

Datasets to predict mitochondrial and non-mitochondrial proteins. To predict mitochondrial proteins and non-mitochondrial proteins, we collected and constructed eight datasets (Table 1) that contain features typical of either mitochondrial or apicoplast proteins. Some proteins, like lipoate protein ligase 2 (PF3D7_0923600) (27) and serine hydroxymethyltransferase isoforms (PF3D7_1235600) (28), are dual localized to the mitochondrion and apicoplast, highlighting the need for the second category of datasets. Each dataset is described in detail below, including how the dataset was processed to obtain per *P. falciparum* protein a single score relevant to whether the protein is mitochondrial or not.

Table 1: Input datasets for Bayesian integration.

Dataset	Abbreviations used in figures	Size of dataset	Fraction identified (%)	Fraction gold identified (%)	P - value
Dataset based on microarray evidence					
Co-expression, built using in-house developed WICCA tool (rank ≤ 2400) (https://wicca.cmbi.umcn.nl/)	Co-expression	5310	45.6	83.0	2.0E-08
Datasets based on ortholog evidence					
hyperLOPIT (TAGM.MCMC.joint.mitochondrion_max ≥ 4.78E-05) (21)	hyperLOPIT	2056	16	94.9	3.3E-28
Mitochondrial ortholog	Mito ortholog	5280	8.7	75.5	2.6E-32

| Evolutionary inference, built using CLIME (NN score ≤ -1.5) (29) | CLIME | 4687 | 9.0 | 38.5 | 6.9E-09 |
| Absence of a *Cryptosporidium* ortholog (score ≤ 4) | Crypto ortholog | 5280 | 67.5 | 77.4 | 7.9E-02 |

Datasets based on amino acid sequence evidence

Mitochondrial targeting signal, built using PFMpred tool (SVM ≥ -0.67) (30)	Mitochondrial TS	5324	59.5	83.3	1.4E-04
Absence of an Apicoplast Targeting Signal, based on PlasmoAP (31)	Apicoplast TS	5324	90.5	96.3	1.0E-01
Isoelectric point (Patrickios pI ≥ 4.2)	pI	5324	55.1	83.3	1.1E-05

172 The size of datasets indicates the number of *Plasmodium* genes that could potentially
173 have been measured in this dataset. Fractions indicate the actually identified genes of this
174 set that fall in bins with a positive mitochondrial score (Fig 1A). P values (Fisher's exact
175 test) indicate overrepresentation of identified gold standard genes compared to random
176 expectation.
177
Dataset based on microarray evidence
Co-expression. Co-expressed genes tend to code for proteins that functionally interact.
For mammals, co-expression analyses have been successfully used in the discovery of
mitochondrial proteins (5, 32). To determine if the expression pattern of a *Plasmodium*
gene correlates with mitochondrial protein coding genes, we developed a co-expression
tool: WeIghted Co-expression Calculation tool for plAsmodium genes (WICCA, available at
https://wicca.cmbi.umcn.nl/). In short, the tool uses the co-expression of a group of input
genes with each other to weigh 83 microarray experiments for their predictive value for
that group. Datasets that show high co-expression of the input genes are more likely to be
relevant for the system that the input genes are part of, and receive a higher weight. It
then uses these weights to combine the expression datasets and calculate one co-
expression score per gene with the input system, in this case, one score per gene for the
level of co-expression with mitochondrial protein coding genes. We used the co-
expression ranking obtained with the alternative positive standard of 146 genes as input
for WICCA to weigh the co-expression datasets. The methods of this tool were based on
the WeGET method (http://weget.cmbi.umcn.nl/, (33)), methodological details on WICCA
can be found in the supplement.
195
Datasets based on ortholog evidence
hyperLOPIT. Hyperplexed Localization of Organelle Proteins by Isotope Tagging
(hyperLOPIT) is a proteomics technique that analyzes protein distributions upon
biochemical fractionation. It enables the identification of the subcellular localization of

thousands of proteins (34, 35). Barylyuk *et al.* used this technique on the apicomplexan
T. gondii and, among others, classified proteins identified in all three hyperLOPIT
experiments to be part of the mitochondrial soluble matrix and mitochondrial membrane.
This classification was based on t-augmented Gaussian mixture models (TAGM) in
combination with maximum a posteriori prediction (TAGM-MAP) and Markov-chain
Monte-Carlo (TAGM-MCMC) methods (21). Marker proteins used for the classification were
arbitrarily set to 1 in the class that they belong to and to 0 for all other classes. Using
BLAST (36), or if that produced no homologs HHpred (37), in combination with selecting
for best bidirectional hits, we determined a list of orthologs between *T. gondii* and
P. falciparum. The 3,832 *T. gondii* proteins that got a TAGM-MCMC probability for
mitochondrial soluble matrix and mitochondrial membrane were mapped to the
corresponding *P. falciparum* ortholog when available. This resulted in a list of 2,057
P. falciparum identifiers with two TAGM-MCMC probabilities, one for mitochondrial matrix
localization and one for mitochondrial membrane localization. As the input datasets for
the Bayesian integration should be independent, both TAGM-MCMC probabilities cannot
be included as separate datasets. Therefore the maximum TAGM-MCMC probability (so
either the soluble or the membrane value) for each protein was used to create one input
hyperLOPIT dataset.

Mitochondrial ortholog. We also used orthologs from more distantly related
species than *T. gondii*, as also at larger phylogenetic distances subcellular localizations
tend to be conserved (38). Using BLAST (36) (all versus all blast for proteomes with e-value
of 100) in combination with OrthoMCL (39) (percent match cut of 50, e-value exponent cut
off -5) to detect orthologs, or if that produced no result, HHpred (37) (default settings, e-
value cut off 0.01, three iterations, only best-bi-directional hits included), one-to-one
orthologs between *P. falciparum* and either *Homo sapiens*, *Arabidopsis thaliana* or
Saccharomyces cerevisiae were determined. When the orthologous protein of at least one
species was annotated to be mitochondrial in a published mitochondrial compendium (*H.
sapiens* MitoCarta2.0 (5), *A. thaliana* (40), *S. cerevisiae* (41)), then that *P. falciparum* protein
was included in the mitochondrial ortholog dataset. *P. falciparum* proteins that had
orthologs in plastids that were more similar than their orthologs in mitochondria were
excluded to prevent e.g. the apicoplast *P. falciparum* pyruvate dehydrogenase (PDH) that
is closely related to the *A. thaliana* plastid PDH to be included in the mitochondrial
orthologs.

Evolutionary inference. Clustered by inferred models of evolution (CLIME) uses
homology across model species (138 eukaryotic species and 1 prokaryotic outgroup) to
identify evolutionary conserved clusters. We downloaded the pre-computed CLIME
analyses of *P. falciparum* genes from http://gene-clime.org/ (29) and mapped the IDs to
our reference proteome. We used the CLIME matrix listing the presence/absence of
orthologs of *P. falciparum* in other species, with the alternative positive and negative sets
as training data, to train a perceptron with two hidden layers to obtain a single score per
protein (details in supplement). The resulting output scores on our test set formed the
CLIME input for the Bayesian integration.

Absence of a *Cryptosporidium* ortholog. Like *Plasmodium*, the genus
Cryptosporidium belongs to the phylum of Apicomplexa. *Cryptosporidium* species lack an
apicoplast and contain only a remnant mitochondrial-like organelle (42). Therefore it can
be expected that apicoplast proteins and many mitochondrial proteins will not have an
ortholog in *Cryptosporidium*. MetaPhOrs is an online tool that contains orthology and
paralogy predictions obtained from multiple phylogenetic trees (43) and was used to

assess orthology for *P. falciparum* genes in three *Cryptosporidium* species (*hominis*,
parvum, and *muris*). We calculated a combined score by multiplying two MetaPhOrs
metrics, one for the confidence level (see below) and one for the number of hits between
the four species, and used this combined score as input for the Bayesian integration. In
detail, the MetaPhOrs results include a consistency score (ranging from 0 with no overlap
to 1 if all trees contain the same protein relationship/orthology information) and an
evidence level for the number of independent databases (the theoretical maximum is 13,
but for our species combination it was four). The multiplication of consistency and
evidence level results in an arbitrary score (range 0-4) that indicates the confidence in
calling two proteins orthologs. *P. falciparum* proteins with a high score are more likely to
have an ortholog in *Cryptosporidium*. Notice that there is some overlap with the
Evolutionary inference as that includes one *Cryptosporidium* species. Nevertheless,
Cryptosporidium species show quite some variation in the complexity of their mitosomes
and the overlap in the predictions is limited (see below).
262
Datasets based on amino acid sequence evidence
Mitochondrial targeting signal. The canonical mitochondrial import system requires an
amphipathic α-helical N-terminal targeting sequence. As *Plasmodium* has a distinct amino
acid usage pattern (44), a *P. falciparum* specific tool, PFMpred (30), was chosen to predict
mitochondrial localization based on the amino acid sequence. This is a support vector
machine (SVM) based tool and was used in split-amino-acid-composition mode (Matthews
correlation coefficient 0.73) that allows separate calculations for the N-terminus, C-
terminus, and remainder of the protein. The tool reports one SVM-score per protein and
indicates whether the protein is predicted to be mitochondrial or not. The SVM-scores per
gene are used as the mitochondrial targeting signal dataset.
Absence of an apicoplast targeting signal. As for mitochondrial proteins, the
majority of the apicoplast proteins are encoded by the nucleus and targeted to the
apicoplast by an N-terminal signal and a transit peptide. PlasmoAP (31) apicoplast
localization predictions based on amino acid sequences were downloaded from
PlasmoDB (8) in January 2018. This binary dataset was used in the prediction to reduce
apicoplast protein contamination among the top scoring proteins.
Isoelectric point. Nuclear encoded mitochondrial proteins need to cross the
negatively charged mitochondrial membrane and need to function properly inside the
mitochondrial environment that has a slightly higher pH compared to the cytosol (45). A
density plot (Supplemental Fig 1) by Patrikios isoelectric points (46) of the human
mitochondrial proteome (MitoCarta2.0 (5)) showed that mitochondrial proteins on
average have a higher pI than the remainder of the proteome. Isoelectric points for the
P. falciparum proteome were calculated using the Patrickios algorithm (47) and directly
used as pI scores.
287
Bayesian integration to predict mitochondrial proteins. Using the eight datasets
described above and the gold standard evaluation sets, a mitochondrial score for each
P. falciparum reference protein was calculated. This score is the log odds of the likelihood
that a protein is localized in the mitochondrion relative to it being localized somewhere
else in the cell. For categorical data (mitochondrial ortholog, apicoplast targeting signal),
the dataset is separated into bins that represent the two categories. Continuous data (the
other six datasets) were binned in a systematic way using a custom python script
(https://github.com/JordyCoolen/Binning) (settings: --bins 0 --score 1). In short, this script

optimizes for each dataset the distribution of the bins such that the sum of log ratio scores
of all bins combined (see below) is close to zero, as this will result in the best separation of
positive and negative bins. The number of bins was varied to a maximum of five bins to
achieve the most optimal separation without expanding the number such that there are
too few proteins per bin to obtain reliable estimates of the log odds of that bin.
The fractions of gold standard proteins per bin are determined. The log ratio of
these fractions determines the score for all other proteins in that respective bin. The
mitochondrial score is based on the sum of log ratios of the individual datasets and is
calculated as follows:
305

$$
mitochondrial\ score \\
= log_2 \left(\frac{P_{mitochondrial}}{P_{non-mitochondrial}} \right) + \sum_{i=1}^{n} log_2 \left(\frac{P(data_i | mitochondrial)}{P(data_i | non - mitochondrial)} \right)
$$

$$
with\ \frac{P(data_i | mitochondrial)}{P(data_i | non - mitochondrial)} = \frac{mitochondrial\_pos_i}{mitochondrial\_neg_i}
$$

307
where mitochondrial_pos and mitochondrial_neg are the fractions of the positive and
negative gold standard genes in sample i respectively. If there were no gold standard
genes found in a certain bin, that positive or negative gold standard fraction was set to
0.5/'total number of negative set genes in the complete dataset' to prevent division by
zero and allow calculation of the log ratio. The O_{prior}, $log_2(P_{mitochondrial}/P_{non-mitochondrial})$, is based
on the estimation that 536 proteins out of the 5357 protein coding genes (~10%) encode
a mitochondrial protein. It does not affect the ranking of potential mitochondrial proteins.
To assess the performance of the integration, a false discovery rate (FDR) was
calculated. As this FDR depends on gold standard genes and the ratio gold
positives/negatives is not similar to the ratio of mitochondrial protein coding genes versus
non-mitochondrial coding genes in the genome, the FDR was corrected (cFDR) using the
following formula:
320

$$
cFDR = \frac{1 - specificity}{1 - specificity + sensitivity \cdot O_{prior}}
$$

321
322
Cross-validation. To assess the ability of the integrated predictor to discriminate known
mitochondrial proteins from non-mitochondrial proteins, ten-fold cross-validation was
performed. The gold standards (both negative and positive) were subsampled ten times,
thereby creating ten sets of nine-tenth of the gold standard genes. Each gold standard
gene was left out once in one of the ten sets. Data integration was performed with each of
these ten sets and the ranks of the one-tenth left out gold standard genes were retrieved.
The ten-fold cross-validated ROC curve was constructed based on those ranks.
330
Intraspecies homologs. The human mitochondrial proteome has expanded since the
divergence from yeast mainly due to gene duplications that create mitochondrial paralogs
(38). It is therefore interesting to see if the predicted *P. falciparum* mitochondrial proteins
have intraspecies homologs, and whether we can identify mitochondrial gene families.
With HHpred (37), using default settings and a 1E-5 E-value cut-off, the homologs within

the genome of *P. falciparum* were determined and included as a column in Supplemental
Table 2.
338
Validation of candidate proteins. To validate the predictions, we selected 14 genes with
unusual or novel mitochondrial functions within the top 295 of genes that fall within the
25% cFDR cut-off. Of these, 13 were selected based on maximum transcript levels during
asexual blood-stage development of *P. berghei* strain ANKA cl15cy1 that exceeded 100
FPKM (48). In addition, we selected one particularly interesting gene, PF3D7_1004900 that
had a lower transcript level. We used experimental genetics (49) to generate *P. berghei*
lines expressing 3xHA-tagged copies of the selected proteins and colocalized them with
an established mitochondrial marker (16) to determine their subcellular localization by
immunofluorescence microscopy. Details on plasmid construction (successful for 13 out of
14 targets, excluding PF3D7_1142800) and the generation of the lines can be found as
supplemental information. To assess colocalization, the freshly harvested transgenic
parasites were allowed to settle on a poly-L-lysine coated coverslip for 10 minutes and
then fixed with 4% EM-grade paraformaldehyde (Fisher Scientific) and 0.0075% EM-grade
glutaraldehyde (Sigma-Aldrich) in microtubule stabilizing buffer (MTSB, 10 mM MES, 150
mM NaCl, 5 mM EGTA, 5 mM glucose, 5 mM $MgCl_2$ pH 6.9) for 20 minutes (50). Next,
parasites were permeabilized with 0.1% Triton X-100 in PBS for 10 minutes. Samples in
which we imaged mOrange and mCherry were hereafter stained with DAPI (1:300) for 15
minutes and mounted on a microscope slide with VectaShield (Vector laboratories). All
other samples were blocked in 3% FCS-PBS for 1 hour. Samples were incubated overnight
at 4°C with rat anti-HA (1:500, ROAHAHA Roche) and chicken anti-GFP (1:1000, Thermo
Fisher) antibodies and for 1 hour with 1:500 goat anti-rat Alexa fluor™ 594 (Invitrogen) and
goat anti-chicken Alexa fluor™ 488 (Thermo Fisher) conjugated antibodies at room
temperature. Nuclei were stained with DAPI (1:300) for 15 minutes and coverslips were
mounted on a microscope slide with VectaShield (Vector laboratories). Samples were
imaged with an SP8 confocal microscope (Leica, 63x oil lens) or LSM900 confocal
microscope (Zeiss, 63x oil lens). Images were processed minimally and similarly with Fiji
(51).
366
367 **Tools for data analysis.** Plots, statistics, and calculations were performed with the R
368 statistical package (52) and additional packages gplots (53), ggplot2 (54), ROCR (55), scales
369 (56) and reshape (57). The separation of the datasets into bins was performed using a
370 python script (see above).
371
Results
We integrated eight features of proteomic, gene expression, orthology and amino acid
composition data, to predict mitochondrial localization in *P. falciparum*. All the individual
features had a predictive value for mitochondrial localization (Fig 1A and 1B), including, as
expected, a negative score for the "*Cryptosporidium* ortholog" and "apicoplast targeting
signal" datasets (Supplemental Fig 2A). Notice that for the co-expression feature it is the
bin with the low-rank values that contains most mitochondrial proteins because this bin
contains the proteins with the highest co-expression values. The best performing input
datasets are the ones that translate mitochondrial localization between species: the
hyperLOPIT dataset and the mitochondrial ortholog dataset, followed by the
mitochondrial targeting signal and co-expression datasets (Fig 1B). Correlation analyses
showed that the features are largely independent off each other, allowing us to use naïve

Bayesian data integration. The highest correlations were observed between features that
examine the phylogenetic distribution of the proteins: CLIME and *Cryptosporidium*
ortholog datasets (Supplemental Fig 2B). By integrating all data, we achieved both a
better sensitivity and specificity in predicting mitochondrial proteins with an AUC of 0.958
than we achieved for individual data sets (Fig 1B). The exception to this is the hyperLOPIT
dataset that on its own performs best with an AUC of 0.960 (Fig 1B). However, this dataset
contains information on only 2,056 (39%) of the nuclear encoded *Plasmodium* proteins,
highlighting the value of additional datasets and their integration to predict a complete
mitochondrial proteome. The distribution of the log-odds ratios (Fig 1C) showed that the
mitochondrial score separated the positive and negative gold standards well from each
other and that there are additional proteins with a high score that were not part of either,
which are potential new mitochondrial proteins. We use a cFDR of 25% (Fig 2A) as a
threshold for highly probable new mitochondrial proteins and identify 295 proteins (Fig
2B) with a sensitivity of 65% and a specificity of 98%. The ranked list of the complete
nuclear encoded proteome is available in Supplemental Table 2.
Overall, the hyperLOPIT dataset predicts 388 *T. gondii* mitochondrial proteins with
a TAGM-MCMC location probability of 99% or higher. Of those, 290 have a *P. falciparum*
ortholog and of these orthologs, we find 217 in our predicted mitochondrial proteome.
We predicted 52 additional mitochondrial proteins that do not have a *T. gondii* ortholog
and 26 proteins that are present in the hyperLOPIT dataset but fell outside their 99% cut
off for mitochondrial localization.
405

406
407 **Fig 1. Predictive values of the individual datasets and the integrated ranked list. (**A)
408 Barplot of each dataset with the fraction of the gold standards on the y-axis and the bins
409 on the x-axis. In green, the mitochondrial score per bin is indicated to show the predictive
410 value of each bin. (B) ROC-curves comparing the performance to identify mitochondrial

proteins of the prediction (10-fold cross-validated) to the performance of the individual datasets. For comparison, the area under the curve values are indicated in the legend. (C) Boxplot that visualizes the distribution of the gold-standard positives, negatives, and remainder of the proteome (x-axis) over the calculated mitochondrial score of the prediction (y-axis).

Fig 2. Integration produces a ranked proteome with 295 likely mitochondrial proteins at a 25% cFDR-cut. (A) Density plot of the mitochondrial score with a colored bar at the bottom indicating the scores of individual gold standards. The arrow indicates the 25% cFDR cut-off. (B) The top 295 proteins falling in the 25% cFDR cut-off ranked on their mitochondrial score with color indicating if the protein is part of a gold standard. Black arrows indicated the ranking of the seven candidate proteins that we experimentally

confirmed to be mitochondrial (Fig 3A and 3B), dark grey arrows indicate the ranking of
the three proteins with suggested mitochondrial localization (Fig 3C), and light grey
arrows the ranking of the three proteins with inconclusive results (Supplemental Fig 4B).
(C) Categorical representation of the 295 proteins identified at the 25% cFDR cut-off. First,
genes were separated into categories based on the gold standard, the remainder on
mitochondrial and apicoplast GO term, and subsequently the remainder on gene function
annotations.
431
Validation of candidates. We aimed to validate the mitochondrial candidate list by
tagging orthologs of the selected unusual or unique predicted mitochondrial proteins in
the efficient transfection model *P. berghei* (see the section 'new mitochondrial proteins' for
the *P. falciparum* orthologs of these proteins). Mitochondrial localization was assessed by
fluorescent microscopy using an experimental genetic approach developed in our lab
(16). This method allows the endogenous tagging of a target protein with a combined
fluorescent and epitope tag, while simultaneously introducing a strong mitochondrial
targeted GFP marker by fusion of the promoter and N terminus of HSP70-3
(PBANKA_0914400) to GFP. Initially, we tagged 6 proteins with a mOrange-3xHA tag. Live
imaging revealed very poor and undefined signals (data not shown). Next, we fixed blood
samples and stained them with anti-HA antibodies. Using this approach, only
PBANKA_0310100 and PBANKA_1203200 were convincingly shown to localize to the
mitochondrion (Fig 3A), while other proteins demonstrated mostly undetectable or
undefined signals (Supplemental Fig 4A). Since many of the selected proteins are
relatively small compared to the mOrange-tag and are expected to be imported into the
mitochondrion, we anticipated that interference of the tag with import or proper folding
could lead to the observed weak and undefined signals. To assess this, we selected two
similarly small and abundantly expressed targets, PBANKA_0310100 (127 amino acids) and
PBANKA_1024800 (144 amino acids), to test different tags. For PBANKA_0310100, which
was already successfully localized to the mitochondrion, these included tags consisting of
only the fluorescent protein mOrange or only the 3xHA epitope tag, either with or without
a linker sequence. For PBANKA_1024800, we also included the combined mCherry-cMyc
tag that has previously been used successfully to tag mitochondrial proteins (16). We
found that the fluorescent protein tag indeed interfered with the localization of
PBANKA_1024800, while the 3xHA tag, with or without a linker, supported a mitochondrial
localization (Supplemental Fig 5B). Surprisingly, using the mOrange-tag without the
subsequent HA-tag also led to an undefined rather punctuate mislocalization of
PBANKA_0310100 (Supplemental Fig 5B). Based on these observations, we decided to tag
the remaining 11 proteins with a linker-3xHA-tag (Fig 3B, 3C and Supplemental Fig 4B).
Five of these were also convincingly localized to the mitochondria (Fig 3B), while an
additional three proteins showed very weak staining patterns suggestive of a possible
mitochondrial localization (Fig 3C). The signals of the remaining three tagged proteins
were unable to be distinguished from the background and therefore no subcellular
localization could be assigned (Supplemental Fig 4B).
466

Fig 3. Validation of mitochondrial candidate proteins. Immunofluorescent analysis of tagged candidate proteins of interest (POI) in *P. berghei* parasites (PBANKA_ID indicated vertically). POI were stained with anti-HA antibody (red, first column), mitochondria were stained with anti-GFP antibody binding to the organelle marker (Mito, green, second column). The DNA was stained with DAPI (blue in merge). (A) Representative images of candidate proteins tagged with a mOrange-3xHA tag, imaged on a SP8 confocal microscope (Leica). (B-C) Representative images of candidate proteins tagged with a linker-3xHA tag, imaged on a LSM900 confocal microscope (Zeiss). (C) Images with the brightest signal for three lowly expressed candidate proteins. Pixel intensities of the POI and mitochondrial signal were normalized and plotted over the line indicated in the merge panel. Scale bar, 2 µm.

New mitochondrial proteins. A categorical representation of the identified proteins (Fig 2C) shows that 138 of the 295 proteins have a previous annotation of being mitochondrial as they are either part of the gold standard, have a mitochondrial GO term, or are described as putative mitochondrial protein. A smaller number of proteins, 16, were part of the negative set or have an apicoplast GO-term & no mitochondrial GO term. The remaining 141 proteins, of which 76 proteins are annotated as conserved hypothetical proteins, contain new mitochondrial candidates.

For 14 of the 76 conserved proteins with an unknown function, we found a
mitochondrial ortholog using the sensitive homology detection tool HHpred (37) that can
improve their annotation (Supplemental Table 2). For example, the human and yeast
orthologs of PF3D7_0306000 (Bayesian rank 41) are both cytochrome b-c1 complex
subunit 8, making it very likely that the *Plasmodium* protein is also cytochrome b-c1
complex subunit 8, as confirmed by recent complexome profiling (12). Other similar
examples include PF3D7_1357600 (rank 283) that is likely a mitochondrial member of the
39S ribosomal protein family L53, and PF3D7_1309900 (rank 78) that is likely succinate
dehydrogenase assembly factor 2.
Besides the possibility to improve annotations using well-characterized
mitochondrial orthologs, the predicted 295 proteins contain biologically interesting
proteins that can add new mitochondrial functions to *P. falciparum* and related species.
Therewith, the prediction provides targets for experimental research to uncover new
biology. These proteins can be divided into five categories: (i) proteins with known
mitochondrial orthologs and an unknown mitochondrial role in *P. falciparum*, (ii) proteins
without a mitochondrial ortholog but with homology to proteins with a known function
and conservation of critical residues that suggest conservation of that function, (iii)
proteins with homology to proteins with a known function from which residues known to
be essential for catalytic activity or for binding substrates residues have been lost, (iv) new
proteins of a known mitochondrial protein family and (v) proteins of a new mitochondrial
protein family. Below, we describe several proteins per category, including their domain
structure, the conservation of active site residues, and their phylogenetic distribution.
509
Proteins with known mitochondrial orthologs and an unknown mitochondrial role in
P. falciparum
PF3D7_0413500 (rank 109) contains a PGM domain and appears to be orthologous to the
human serine/threonine-protein phosphatase, PGAM5, of which the catalytic residue
H105 (58) is conserved. In human cells, PGAM5 is localized to the mitochondrial outer
membrane (59) where it plays a role in apoptosis, mitofission, and mitophagy (reviewed in
(60)) among others via interactions with BCL2-like protein 1 (61). We experimentally
confirmed PF3D7_0413500's mitochondrial location (PBANKA_0715500 in Fig 3B), making
it an interesting protein to study in the context of *Plasmodium* mitochondrial dynamics, a
process about which little is known. Originally, we flagged two additional proteins,
PF3D7_0213200 (rank 164) and PF3D7_0611300 (rank 166), as interesting due to potential
homology to BCL2, based on sequence similarity and a similar alpha-helical structure
predicted with HHpred (37). PF3D7_0213200 was assigned an apicoplast GO-term, but in
our analysis is a potential mitochondrial protein. Analysis with TMHMM (62) indicates that
it probably is a transmembrane protein with an in-out topology. The protein
PF3D7_0611300 already was assigned a mitochondrial GO-term based on a screening in
T. gondii (63), but was, to our knowledge, not confirmed as mitochondrial localized with
with experiments characterizing this protein in *P. falciparum*. Although we rejected BCL2
homology for both proteins after a detailed examination of the residues conserved in the
BCL2 family, we confirmed the mitochondrial localization of PF3D7_0213200
(PBANKA_0310100 in Fig 3A) and PF3D7_0611300 (PBANKA_0109600 in Fig 3B). Indeed,
complexome profiling revealed that the latter is part of the ATP synthase complex (12).
532
533 *Proteins with a non-mitochondrial homolog and conservation of critical residues*

PF3D7_0913400 (rank 30) is homologous to the bacterial elongation factor P. Although eukaryotes contain initiation factor 5A as a cytoplasmic elongation factor P homolog, this protein family was until now not observed in mitochondria. PF3D7_0913400 contains both the N-terminal KOW-like domain and the central P/YeiP domain of elongation factor P, but not the C-terminal OB domain. Elongation factor P plays a role in translation elongation and specifically in the translation of stretches of 2 or more prolines (64). A basic residue at position 34 of elongation factor P (numbering from the *Escherichia coli* PDB structure 3A5Z_F (65)) that after post-translational modification interacts with the CCA-end of the tRNA, is conserved in PF3D7_0913400. Notably, mitochondrial encoded *P. falciparum* cytochrome C oxidase subunit 1 contains a proline pair at position 135-136, making it a potential target for elongation factor P.

PF3D7_0812200 (rank 144) is homologous to the bacterial DegP protease family. It contains both a serine endoprotease domain and a PDZ domain. The DegP from *E. coli* functions in acid resistance in the periplasm. It is able to refold after acid stress and subsequently can cleave proteins misfolded due to acid stress (66). The active site residue (S210 in PDB structure 1KY9 (67)) is conserved in PF3D7_0812200, but residues involved in allosteric loop interactions, including R187 are not. Orthologs of PF3D7_0812200 appear, among the eukaryotes, limited to the Apicomplexa and Chromerida.

PF3D7_0503900 (rank 242) has a C-terminal dioxygenase domain (E= 1.3E-16). The closest homolog in human is the phytanoyl-CoA dioxygenase that resides in the peroxisome, an organelle absent from *Plasmodium*, where it is involved in α-oxidation of branched-chain fatty acids. Important residues in the active site that are involved in iron binding, H175, H177, H264 (68), are conserved in PF3D7_0503900, suggesting that it is a functional enzyme. Nevertheless, levels of sequence conservation are low (14% identity with the human phytanoyl-CoA dioxygenase) and we did not detect conservation of substrate binding sites with the phytanoyl-CoA dioxygenase. The co-localization analysis suggested a mitochondrial localization for PF3D7_0503900 (PBANKA_1103500 in Fig 3C).

PF3D7_1121500 (rank 250) contains a papain-like NLpC/P60 superfamily domain (Peptidase_C92 in PFAM). This protein family contains, among others, cysteine peptidases and amidases. Of the four residues essential for activity in BcPPNE from *Bacillus cereus* (69), the most similar experimentally characterized homolog of PF3D7_1121500, three are conserved (H49, E64, and Y164).

Proteins with homology to proteins with a known function from which residues known to be essential for catalytic activity or for binding substrates residues have been lost

PF3D7_1004900 (rank 206) is homologous to the protein component of the signal recognition particle (SRP), a ribonucleoprotein that recognizes and targets specific proteins to the ER in eukaryotes. PF3D7_1004900 contains the C-terminal M domain (E= 5.8E-25), but not the other, SRP54 and SRP54_N domains of this protein. In SRP the M domain binds both the SRP RNA and the signal sequence of the target protein. Inspection of individual amino acids did not reveal conservation of RNA binding amino acids, e.g. the arginines in helix M4 of the M domain (70), are not conserved, but did reveal some conservation of hydrophobic amino acids lining the groove in which the signal peptide is located. These hydrophobic amino acids have been implicated in interactions with the signal peptide, specifically V323, I326, L329 in the M1 helix and L418 in the M5 helix (positions relative to *Sulfolobus solfataricus* structure 3KL4 (70, 71)), suggesting a possible conservation of interaction of PF3D7_1004900 with a hydrophobic α-helix. We

experimentally confirmed the mitochondrial location of PF3D7_1004900
(PBANKA_1203200 in Fig 3A).
PF3D7_1246700 (rank 217) is a member of the pyridoxamine 5'-phosphate oxidase
(PNPOx) like protein family that, besides pyridoxamine 5'-phosphate oxidase, also
contains the general stress protein 26 family. The most similar experimentally
characterized protein is general stress protein 26 from *Xanthomonas citri* encoded by gene
Xac2369. The latter binds FAD and FMN but not pyridoxal 5'-phosphate, indicating that it
does not function as a PNPOx (72). Relative to the unpublished structure of a protein from
this family from *Nostoc punctiforme* (PDB: 2I02), that does contain an FMN, we did not
observe conservation of the amino acids like W111, that line the FMN binding pocket,
casting doubt of a function of PF3D7_1246700 that includes the binding of FMN.
592
New members of mitochondrial protein families
PF3D7_1417900 (rank 107), PF3D7_1142800 (rank 167) and PF3D7_0927100 (rank 244)
potentially contain a CHCH domain. They all have two pairs of cysteines that are very well
conserved among their detectable homologs, and the nine amino acids between the
cysteines are predicted to form an α-helix (37). Nevertheless, as nothing else but the
cysteines appears to be conserved, the E-value with the PFAM CHCH domain entry, or with
any known mitochondrial protein, is not significant (E>0.01). Jackhmmer analysis (73)
detects homologs among the Apicomplexa, including a homolog in *Cryptosporidium*
muris, but not outside this taxon. Known proteins with a CHCH domain are localized to the
mitochondrial intermembrane space where they can participate in disulfide bond
formation, suggesting that PF3D7_1417900, PF3D7_1142800, and PF3D7_0927100 might
also be in the intermembrane space. Complexome profiling revealed that the first two are
part of the ATP synthase complex (12). We tested PF3D7_1417900 and PF3D7_0927100
experimentally and confirmed them to be mitochondrial (PBANKA_1024800 and
PBANKA_0827900 in Fig 3B respectively). Thus, based on our data and the complexome
profiling data all three proteins are confirmed to be mitochondrial, increasing their
likelihood that they indeed contain a CHCH domain.
Plasmodium contains a cytochrome C heme lyase (PF3D7_1224600, rank 126) and a
cytochrome C1 heme lyase (PF3D7_1203600, rank 251), which are both essential during
blood-stage development and have non-overlapping functions (74). We uncovered a third
homolog in this family at rank 183, PF3D7_1121200 (E=0.0096, HHpred (37)). Although the
homology covers domains II, III, and IV of heme lyases, of which domain II has been
implicated in binding Heme (75), residues that have been implicated in heme binding, like
H154 (coordinates for human protein (75)) and residues of domain I are not conserved in
PF3D7_1121200. The protein also occurs in ciliates and dinoflagellates but has no
orthologs outside of those taxa and therewith appears to be an alveolate-specific
duplication.
620
A new Apicomplexan mitochondrial protein family
We assessed to what extent *Plasmodium* mitochondrial proteins were part of families with
multiple mitochondrial members that resulted from duplications (Supplemental Table 2).
Such "intra compartmental protein duplications" have also increased the size of the
human mitochondrial proteome (38). Among the mitochondrial homologs, we uncovered
a new family in the Apicomplexa that in *Plasmodium* species has four representatives:
PF3D7_0821900 (rank 203), PF3D7_1336100 (rank 254), PF3D7_1134400 (rank 278), and
PF3D7_1228000 (rank 503). The family is characterized by a well-conserved WPP motif at

its N-terminus (Supplemental Fig 7), and we therefore name it the WPP family. Although PF3D7_1228000 does not rank sufficiently high to fall within the 25% cFDR cut-off, all proteins of the family are likely mitochondrial given their overall high ranking and the tendency of members of protein families to be localized in the same cellular location. For the protein family itself, we did not detect homologs outside the Apicomplexa and *Vitrella*. In several species including *Plasmodium coatneyi* (PCOAH_00054040), *Babesia bovis* (BBOV_III010100), and *Theileria orientalis* (MACL_00002472), the ortholog of PF3D7_1228000 is predicted to be fused with an rRNA pseudourydilate synthase hinting at a potential function of this family in rRNA maturation. Nevertheless, there are no experimental data supporting those gene fusions. We unequivocally confirmed the mitochondrial location of PF3D7_0821900 (PBANKA_0708900 in Fig 3B) and the co-localization analysis also suggested a mitochondrial localization for PF3D7_1336100 and PF3D7_1134400 (PBANKA_1350000 and PBANKA_0914000 in Fig 3C respectively). PF3D7_1228000 was excluded from the analysis due to low transcript levels.

Discussion

In this work, we ranked all nuclear encoded proteins of *P. falciparum* for their likelihood of being mitochondrially targeted and defined the top 295 proteins (cFDR 25%) as likely mitochondrial proteins (Supplemental Table 2). Despite the large variation between the datasets in the coverage of the nuclear encoded proteome and in the ability to identify mitochondrial proteins (Table 1, Fig 1), the probabilistic approach we used was able to integrate the data in an unbiased way by assessing the distribution of gold standard genes within each dataset during the scoring process.

Quality of the predictions. Several observations indicate the high quality of the predictions. First of all, the 10-fold cross-validation with an area under the curve of 0.958 shows the high predictive value and robustness of the prediction. Second, half of the predicted proteins have a previous annotation of being mitochondrial, even though these were not all used in weighing the datasets. Third, of thirteen proteins that we set out to validate experimentally, the seven for which we obtained a high enough expression to convincingly locate them in the cell, were all mitochondrial, while for three others the experimental data was consistent with a mitochondrial localization but equivocal due to low expression levels. These seven plus three were well distributed over the order of predicted new mitochondrial proteins (Fig 2B). Fourth, even though our top 295 contains seven proteins of the negative gold standard that was not manually supervised for all entries, manual examination of those seven showed that the majority are likely mitochondrial after all. Six of them were detected in the hyperLOPIT study (21) and of those five were annotated as mitochondrial (legend Supplemental Table 2). Finally, several proteins that we predicted to be mitochondrial have, since we made those predictions, been shown to be mitochondrial using other methods. Specifically, complexome profiling of mitochondria-enriched fractions has elucidated the composition of mitochondrial complexes II, III, IV, and V (12). We identify in our top 295 three members of complex II (of which two with unknown function), nine members of complex III (of which three with unknown function), ten members of complex IV (of which six with unknown function), and eight of complex V (all with unknown function). In general, most members of the protein complexes (described in (12)) were retrieved by our mitochondrial score (Supplementary Tables 2 and 5). Of the 24 electron transport chain members that were newly discovered by complexome profiling (12), 19 were in our top 295 (Supplemental Tables 2 and 5). The

677 results outlined above reinforce the robustness of the predictions and that the ranked list
678 of the complete nuclear encoded *P. falciparum* proteome, which we name
679 PlasmoMitoCarta, is a useful resource for *Plasmodium* researchers.
680
New mitochondrial biology. The results of the integration provide ample material for
exploring new mitochondrial biology as 101 out of the 295 proteins do not have a
detectable mitochondrial ortholog in yeast, human, or plant. Some, however, do have
homologs in bacteria that include conservation of critical residues, for example, the
elongation factor P whose function, if confirmed, would be unique to mitochondria. Also
outside the cutoff are interesting candidates for mitochondrial proteins, like
PF3D7_0213000 (rank 328) that has an ortholog in *T. gondii* whose mitochondrial
localization was confirmed microscopically (21) and PF3D7_1026200 (rank 332) that
contains an RNA binding THUMP domain and a UPF0020 RNA methyltransferase domain.
The most similar homolog of PF3D7_1026200 with experimentally determined function is
the bifunctional methyltransferase RlmKL from *E. coli*. PF3D7_1026200 is homologous to
the N-terminal methyltransferase domain of RlmKL (E=5E-38) that methylates G2445 in
the *E. coli* large subunit rRNA (76), and this nucleotide is conserved in the *P. falciparum*
mitochondrial rRNA (77).
As part of this study, we provide two other valuable resources. First, we included
the ortholog information gathered for this study that can improve the annotation of
P. falciparum proteins and provide researchers with additional hints for the function of a
protein (Supplemental Table 2). In addition, the WICCA tool that we developed for this
study is available online. It to allows any researcher to assess co-expression of *P. falciparum*
genes with a set of genes of four *Plasmodium* species across data from 83 microarray
expression experiments, by weighing the datasets for their relevance to the set of genes.
This can aid in the discovery of novel members of known pathways and molecular
systems.
704
Size of the mitochondrial proteome. We predict 295 mitochondrial proteins at the strict
cut off, set at cFDR 25%. When using the posterior probabilities (2^mitochondrial score) of
all *P. falciparum* genes to calculate the estimated number (E) of mitochondrial proteins by
summing them all up (E=sum($P_{posterior}$/($P_{posterior}$+1), see ref (78) for a detailed explanation),
we predicted the total size of the mitochondrial proteome to be 466 proteins. It therefore
appears that our prior odds of ~10%, which affects the total size calculation,
overestimated the number of mitochondrial proteins. When we lower the prior odds (see
ref (78)) to 8.7% (430 proteins) the number is consistent with the estimated size of the
proteome based on all data. In contrast, the presence of five likely mitochondrial proteins
in our gold negative sets leads to an underestimate of the size of the mitochondrial
proteome. We decided not to manually weed out such inconsistencies, as this cannot be
done systematically for all datasets, and applying this only to the apparent inconsistencies
would result in circular arguments.
The predicted total size of 466 proteins is relatively small compared to the
mitochondrial proteomes in human, plant, and yeast that consist of at least 900 proteins
(5, 40, 41). However, small mitochondrial proteomes have been reported before, for
example, 573 proteins in the distantly related ciliate *Tetrahymena thermophila* (79). In
agreement with our findings, a recent study on the mitochondrial proteome of *T. gondii*
that was not included in our integration reports a proteome of 421 proteins (15).
724

725 In conclusion, we combined the information of eight datasets with the curated gold
726 standards to rank all nuclear encoded *P. falciparum* proteins on their likelihood of being
727 mitochondrial in PlasmoMitoCarta. The value of this ranked proteome is shown by *in vivo*
728 validation of top scoring proteins in the closely related species *P. berghei* and will provide
729 an important resource for future investigations.
730

Funding
SLvE and CB were supported by PhD fellowships from the Radboud Institute for Molecular
Life Sciences, Radboudumc (Radboudumc JO ronde 2014 and #19-015a). NIP was
supported by a Marie-Slodowska Curie grant (790085), and TWAK and LEdV by the
Netherlands Organisation for Scientific Research (NWO-VIDI 864.13.009). BED was
supported by the Netherlands Organization for Scientific Research (NWO) Vidi grant
864.14.004 and European Research Council (ERC) Consolidator grant 865694: DiversiPHI.
KB was supported by a Leverhulme Trust and Isaac Newton Trust Fellowship (ECF-2015-
562), and RFW by the Wellcome Trust (214298/Z/18/Z).

Competing interests
The authors declare no competing interests.

REFERENCES

1.   Gray MW, Burger G, Lang BF. Mitochondrial evolution. Science (New York, NY).
1999;283(5407):1476-81.
2.   Burger G, Gray MW, Forget L, Lang BF. Strikingly bacteria-like and gene-rich
mitochondrial genomes throughout jakobid protists. Genome biology and evolution.
2013;5(2):418-38.
3.   Wilson RJ, Williamson DH. Extrachromosomal DNA in the Apicomplexa. Microbiology
and molecular biology reviews : MMBR. 1997;61(1):1-16.
4.   Makiuchi T, Nozaki T. Highly divergent mitochondrion-related organelles in
anaerobic parasitic protozoa. Biochimie. 2014;100:3-17.
5.   Calvo SE, Clauser KR, Mootha VK. MitoCarta2.0: an updated inventory of mammalian
mitochondrial proteins. Nucleic acids research. 2016;44(D1):D1251-7.
6.   Jedelský PL, Doležal P, Rada P, Pyrih J, Smíd O, Hrdý I, et al. The minimal proteome in
the reduced mitochondrion of the parasitic protist Giardia intestinalis. PloS one.
2011;6(2):e17285.
7.   van Dooren GG, Stimmler LM, McFadden GI. Metabolic maps and functions of the
Plasmodium mitochondrion. FEMS microbiology reviews. 2006;30(4):596-630.
8.   Aurrecoechea C, Brestelli J, Brunk BP, Dommer J, Fischer S, Gajria B, et al. PlasmoDB:
a functional genomic database for malaria parasites. Nucleic acids research.
2009;37(Database issue):D539-43.
9.   Goodman CD, Siregar JE, Mollard V, Vega-Rodríguez J, Syafruddin D, Matsuoka H, et
al. Parasites resistant to the antimalarial atovaquone fail to transmit by mosquitoes. Science
(New York, NY). 2016;352(6283):349-53.
10.   Sturm A, Mollard V, Cozijnsen A, Goodman CD, McFadden GI. Mitochondrial ATP
synthase is dispensable in blood-stage Plasmodium berghei rodent malaria but essential in
the mosquito phase. Proceedings of the National Academy of Sciences of the United States
of America. 2015;112(33):10216-23.

11.   Painter HJ, Morrisey JM, Mather MW, Vaidya AB. Specific role of mitochondrial
electron transport in blood-stage Plasmodium falciparum. Nature. 2007;446(7131):88-91.
12.   Evers F, Cabrera-Orefice A, Elurbe DM, Lindert MK-t, Boltryk SD, Voss TS, et al.
Composition and stage dynamics of mitochondrial complexes in Plasmodium falciparum.
bioRxiv. 2020:2020.10.05.326496.
13.   Pastore C, Adinolfi S, Huynen MA, Rybin V, Martin S, Mayer M, et al. YfhJ, a
molecular adaptor in iron-sulfur cluster formation or a frataxin-like protein? Structure
(London, England : 1993). 2006;14(5):857-67.
14.   Goldberg AV, Molik S, Tsaousis AD, Neumann K, Kuhnke G, Delbac F, et al.
Localization and functionality of microsporidian iron-sulphur cluster assembly proteins.
Nature. 2008;452(7187):624-8.
15.   Seidi A, Muellner-Wong LS, Rajendran E, Tjhin ET, Dagley LF, Aw VY, et al. Elucidating
the mitochondrial proteome of Toxoplasma gondii reveals the presence of a divergent
cytochrome c oxidase. eLife. 2018;7.
16.   Matz JM, Goosmann C, Matuschewski K, Kooij TWA. An Unusual Prohibitin Regulates
Malaria Parasite Mitochondrial Membrane Potential. Cell reports. 2018;23(3):756-67.
17.   Fry M, Pudney M. Site of action of the antimalarial hydroxynaphthoquinone, 2-
[trans-4-(4'-chlorophenyl) cyclohexyl]-3-hydroxy-1,4-naphthoquinone (566C80). Biochemical
pharmacology. 1992;43(7):1545-53.
18.   Stocks PA, Barton V, Antoine T, Biagini GA, Ward SA, O'Neill PM. Novel inhibitors of
the Plasmodium falciparum electron transport chain. Parasitology. 2014;141(1):50-65.
19.   Kobayashi T, Sato S, Takamiya S, Komaki-Yasuda K, Yano K, Hirata A, et al.
Mitochondria and apicoplast of Plasmodium falciparum: behaviour on subcellular
fractionation and the implication. Mitochondrion. 2007;7(1-2):125-32.
20.   Seeber F, Limenitakis J, Soldati-Favre D. Apicomplexan mitochondrial metabolism: a
story of gains, losses and retentions. Trends in parasitology. 2008;24(10):468-78.
21.   Barylyuk K, Koreny L, Ke H, Butterworth S, Crook OM, Lassadi I, et al. A
Comprehensive Subcellular Atlas of the Toxoplasma Proteome via hyperLOPIT Provides
Spatial Context for Protein Functions. Cell Host Microbe. 2020;28(5):752-66.e9.
22.   Jansen R, Yu H, Greenbaum D, Kluger Y, Krogan NJ, Chung S, et al. A Bayesian
networks approach for predicting protein-protein interactions from genomic data. Science
(New York, NY). 2003;302(5644):449-53.
23.   van der Lee R, Feng Q, Langereis MA, Ter Horst R, Szklarczyk R, Netea MG, et al.
Integrative Genomics-Based Discovery of Novel Regulators of the Innate Antiviral Response.
PLoS computational biology. 2015;11(10):e1004553.
24.   Meerstein-Kessel L, van der Lee R, Stone W, Lanke K, Baker DA, Alano P, et al.
Probabilistic data integration identifies reliable gametocyte-specific proteins and transcripts
in malaria parasites. Scientific reports. 2018;8(1):410.
25.   van Esveld SL, Cansız-Arda Ş, Hensen F, van der Lee R, Huynen MA, Spelbrink JN. A
Combined Mass Spectrometry and Data Integration Approach to Predict the Mitochondrial
Poly(A) RNA Interacting Proteome. Frontiers in cell and developmental biology. 2019;7:283.
26.   Woodcroft BJ, McMillan PJ, Dekiwadia C, Tilley L, Ralph SA. Determination of protein
subcellular localization in apicomplexan parasites. Trends in parasitology. 2012;28(12):546-
54.
27.   Günther S, Wallace L, Patzewitz EM, McMillan PJ, Storm J, Wrenger C, et al.
Apicoplast lipoic acid protein ligase B is not essential for Plasmodium falciparum. PLoS
pathogens. 2007;3(12):e189.

28.   Read M, Müller IB, Mitchell SL, Sims PF, Hyde JE. Dynamic subcellular localization of
isoforms of the folate pathway enzyme serine hydroxymethyltransferase (SHMT) through
the erythrocytic cycle of Plasmodium falciparum. Malaria journal. 2010;9:351.
29.   Li Y, Calvo SE, Gutman R, Liu JS, Mootha VK. Expansion of biological pathways based
on evolutionary inference. Cell. 2014;158(1):213-25.
30.   Verma R, Varshney GC, Raghava GP. Prediction of mitochondrial proteins of malaria
parasite using split amino acid composition and PSSM profile. Amino acids. 2010;39(1):101-
10.
31.   Foth BJ, Ralph SA, Tonkin CJ, Struck NS, Fraunholz M, Roos DS, et al. Dissecting
apicoplast targeting in the malaria parasite Plasmodium falciparum. Science (New York, NY).
2003;299(5607):705-8.
32.   Baughman JM, Nilsson R, Gohil VM, Arlow DH, Gauhar Z, Mootha VK. A
computational screen for regulators of oxidative phosphorylation implicates SLIRP in
mitochondrial RNA homeostasis. PLoS genetics. 2009;5(8):e1000590.
33.   Szklarczyk R, Megchelenbrink W, Cizek P, Ledent M, Velemans G, Szklarczyk D, et al.
WeGET: predicting new genes for molecular systems by weighted co-expression. Nucleic
acids research. 2016;44(D1):D567-73.
34.   Mulvey CM, Breckels LM, Geladaki A, Britovšek NK, Nightingale DJH, Christoforou A,
et al. Using hyperLOPIT to perform high-resolution mapping of the spatial proteome. Nature
protocols. 2017;12(6):1110-35.
35.   Christoforou A, Mulvey CM, Breckels LM, Geladaki A, Hurrell T, Hayward PC, et al. A
draft map of the mouse pluripotent stem cell spatial proteome. Nature communications.
2016;7:8992.
36.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search
tool. Journal of molecular biology. 1990;215(3):403-10.
37.   Zimmermann L, Stephens A, Nam SZ, Rau D, Kübler J, Lozajic M, et al. A Completely
Reimplemented MPI Bioinformatics Toolkit with a New HHpred Server at its Core. Journal of
molecular biology. 2018;430(15):2237-43.
38.   Szklarczyk R, Huynen MA. Expansion of the human mitochondrial proteome by intra-
and inter-compartmental protein duplication. Genome biology. 2009;10(11):R135.
39.   Li L, Stoeckert CJ, Jr., Roos DS. OrthoMCL: identification of ortholog groups for
eukaryotic genomes. Genome research. 2003;13(9):2178-89.
40.   Rao RS, Salvato F, Thal B, Eubel H, Thelen JJ, Møller IM. The proteome of higher plant
mitochondria. Mitochondrion. 2017;33:22-37.
41.   Morgenstern M, Stiller SB, Lübbert P, Peikert CD, Dannenmaier S, Drepper F, et al.
Definition of a High-Confidence Mitochondrial Proteome at Quantitative Scale. Cell reports.
2017;19(13):2836-52.
42.   Putignani L, Tait A, Smith HV, Horner D, Tovar J, Tetley L, et al. Characterization of a
mitochondrion-like organelle in Cryptosporidium parvum. Parasitology. 2004;129(Pt 1):1-18.
43.   Pryszcz LP, Huerta-Cepas J, Gabaldón T. MetaPhOrs: orthology and paralogy
predictions from multiple phylogenetic evidence using a consistency-based confidence
score. Nucleic acids research. 2011;39(5):e32.
44.   Bender A, van Dooren GG, Ralph SA, McFadden GI, Schneider G. Properties and
prediction of mitochondrial transit peptides from Plasmodium falciparum. Molecular and
biochemical parasitology. 2003;132(2):59-66.

45.   Alberts BJ, A.; Lewis, J.; Raff, M.; Roberts, K.; Walter, P. Molecular Biology of the Cell
New York: Garland Science; 2002 [4th edition:[The Mitochondrion]. Available from:
https://www.ncbi.nlm.nih.gov/books/NBK26894/.
46.   Kozlowski LP. Proteome-pI: proteome isoelectric point database. Nucleic acids
research. 2017;45(D1):D1112-d6.
47.   Patrickios CS. Polypeptide Amino Acid Composition and Isoelectric Point: 1. A Closed-
Form Approximation. Journal of Colloid and Interface Science. 1995;175(1):256-60.
48.   Otto TD, Böhme U, Jackson AP, Hunt M, Franke-Fayard B, Hoeijmakers WA, et al. A
comprehensive evaluation of rodent malaria parasite genomes and gene expression. BMC
biology. 2014;12:86.
49.   Matz JM, Kooij TW. Towards genome-wide experimental genetics in the in vivo
malaria model parasite Plasmodium berghei. Pathogens and global health. 2015;109(2):46-
60.
50.   Deligianni E, Morgan RN, Bertuccini L, Kooij TW, Laforge A, Nahar C, et al. Critical role
for a stage-specific actin in male exflagellation of the malaria parasite. Cellular microbiology.
2011;13(11):1714-30.
51.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an
open-source platform for biological-image analysis. Nature methods. 2012;9(7):676-82.
52.   R Core Team. R: A Language and Environment for Statistical Computing. Vienna,
Austria: R Foundation for Statistical Computing; 2015.
53.   Warnes GR, Bolker B, Bonebakker L, Gentleman R, Liaw WHA, Lumley T, et al. gplots:
Various R Programming Tools for Plotting Data. 2016.
54.   Wickham H. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York;
2016.
55.   Sing T, Sander O, Beerenwinkel N, Lengauer T. ROCR: visualizing classifier
performance in R. Bioinformatics (Oxford, England). 2005;21(20):3940-1.
56.   Wickham H. scales: Scale Functions for Visualization. 2017.
57.   Wickham H. Reshaping data with the reshape package. Journal of Statistical
Software. 2007;21(12).
58.   Takeda K, Komuro Y, Hayakawa T, Oguchi H, Ishida Y, Murakami S, et al.
Mitochondrial phosphoglycerate mutase 5 uses alternate catalytic activity as a protein
serine/threonine phosphatase to activate ASK1. Proceedings of the National Academy of
Sciences of the United States of America. 2009;106(30):12301-5.
59.   Lo SC, Hannink M. PGAM5 tethers a ternary complex containing Keap1 and Nrf2 to
mitochondria. Experimental cell research. 2008;314(8):1789-803.
60.   Ma K, Chen G, Li W, Kepp O, Zhu Y, Chen Q. Mitophagy, Mitochondrial Homeostasis,
and Cell Fate. Frontiers in cell and developmental biology. 2020;8:467.
61.   Lo SC, Hannink M. PGAM5, a Bcl-XL-interacting protein, is a novel substrate for the
redox-regulated Keap1-dependent ubiquitin ligase complex. The Journal of biological
chemistry. 2006;281(49):37893-903.
62.   Krogh A, Larsson B, von Heijne G, Sonnhammer EL. Predicting transmembrane
protein topology with a hidden Markov model: application to complete genomes. Journal of
molecular biology. 2001;305(3):567-80.
63.   Sidik SM, Huet D, Ganesan SM, Huynh MH, Wang T, Nasamu AS, et al. A Genome-
wide CRISPR Screen in Toxoplasma Identifies Essential Apicomplexan Genes. Cell.
2016;166(6):1423-35.e12.

64.   Ude S, Lassak J, Starosta AL, Kraxenberger T, Wilson DN, Jung K. Translation
elongation factor EF-P alleviates ribosome stalling at polyproline stretches. Science (New
York, NY). 2013;339(6115):82-5.

65.   Yanagisawa T, Sumida T, Ishii R, Takemoto C, Yokoyama S. A paralog of lysyl-tRNA
synthetase aminoacylates a conserved lysine residue in translation elongation factor P.
Nature structural & molecular biology. 2010;17(9):1136-43.

66.   Fu X, Wang Y, Shao H, Ma J, Song X, Zhang M, et al. DegP functions as a critical
protease for bacterial acid resistance. The FEBS journal. 2018;285(18):3525-38.

67.   Krojer T, Garrido-Franco M, Huber R, Ehrmann M, Clausen T. Crystal structure of
DegP (HtrA) reveals a new protease-chaperone machine. Nature. 2002;416(6879):455-9.

68.   McDonough MA, Kavanagh KL, Butler D, Searls T, Oppermann U, Schofield CJ.
Structure of human phytanoyl-CoA 2-hydroxylase identifies molecular mechanisms of
Refsum disease. The Journal of biological chemistry. 2005;280(49):41101-10.

69.   Xu Q, Rawlings ND, Chiu HJ, Jaroszewski L, Klock HE, Knuth MW, et al. Structural
analysis of papain-like NlpC/P60 superfamily enzymes with a circularly permuted topology
reveals potential lipid binding sites. PloS one. 2011;6(7):e22013.

70.   Mary C, Scherrer A, Huck L, Lakkaraju AK, Thomas Y, Johnson AE, et al. Residues in
SRP9/14 essential for elongation arrest activity of the signal recognition particle define a
positively charged functional domain on one side of the protein. RNA (New York, NY).
2010;16(5):969-79.

71.   Janda CY, Li J, Oubridge C, Hernández H, Robinson CV, Nagai K. Recognition of a
signal peptide by the signal recognition particle. Nature. 2010;465(7297):507-10.

72.   Hilario E, Li Y, Niks D, Fan L. The structure of a Xanthomonas general stress protein
involved in citrus canker reveals its flavin-binding property. Acta crystallographica Section D,
Biological crystallography. 2012;68(Pt 7):846-53.

73.   Potter SC, Luciani A, Eddy SR, Park Y, Lopez R, Finn RD. HMMER web server: 2018
update. Nucleic acids research. 2018;46(W1):W200-w4.

74.   Posayapisit N, Songsungthong W, Koonyosying P, Falade MO, Uthaipibull C,
Yuthavong Y, et al. Cytochrome c and c1 heme lyases are essential in Plasmodium berghei.
Molecular and biochemical parasitology. 2016;210(1-2):32-6.

75.   Babbitt SE, San Francisco B, Bretsnyder EC, Kranz RG. Conserved residues of the
human mitochondrial holocytochrome c synthase mediate interactions with heme.
Biochemistry. 2014;53(32):5261-71.

76.   Wang KT, Desmolaize B, Nan J, Zhang XW, Li LF, Douthwaite S, et al. Structure of the
bifunctional methyltransferase YcbY (RlmKL) that adds the m7G2069 and m2G2445
modifications in Escherichia coli 23S rRNA. Nucleic acids research. 2012;40(11):5138-48.

77.   Feagin JE, Harrell MI, Lee JC, Coe KJ, Sands BH, Cannone JJ, et al. The fragmented
mitochondrial ribosomal RNAs of Plasmodium falciparum. PloS one. 2012;7(6):e38320.

78.   van Dam TJP, Kennedy J, van der Lee R, de Vrieze E, Wunderlich KA, Rix S, et al.
CiliaCarta: An integrated and validated compendium of ciliary genes. PloS one.
2019;14(5):e0216705.

79.   Smith DG, Gawryluk RM, Spencer DF, Pearlman RE, Siu KW, Gray MW. Exploring the
mitochondrial proteome of the ciliate protozoon Tetrahymena thermophila: direct analysis
by tandem mass spectrometry. Journal of molecular biology. 2007;374(3):837-63.