Tumor Microenvironment Alters Chemoresistance of Hepatocellular Carcinoma Through CYP3A4 Metabolic Activity

Alican Özkan[1,11]*, **Danielle L. Stolley[2]**, **Erik N. K. Cressman[3]**, **Matthew McMillin[4,5]**, **Sharon DeMorrow[4,5,6]**, **Thomas E. Yankeelov[2,7,8,9,10]**, **Marissa Nichole Rylander[1,2,7]**

[1] Department of Mechanical Engineering, The University of Texas, Austin, TX, 78712, United States

[2] Department of Biomedical Engineering, The University of Texas, Austin, TX, 78712, United States

[3] Department of Interventional Radiology, The University of Texas MD Anderson Cancer Center, Houston, TX, 77030. United States

[4] Department of Internal Medicine, Dell Medical School, The University of Texas at Austin, Austin, TX, United States

[5] Central Texas Veterans Health Care System, Temple, TX, United States

[6] Division of Pharmacology and Toxicology, College of Pharmacy, The University of Texas at Austin, Austin, TX 78712, United States

[7] Oden Institute for Computational Engineering and Sciences, The University of Texas, Austin, TX, 78712, United States

[8] Departments of Diagnostic Medicine, The University of Texas, Austin, TX, 78712, United States

[9] Department of Oncology, The University of Texas, Austin, TX, 78712, United States

[10] Livestrong Cancer Institutes, Dell Medical School, The University of Texas, Austin, TX, 78712, United States

[11] Current address: Wyss Institute for Biologically Inspired Engineering at Harvard University, Boston, MA, 02115, United States.

* Correspondence: Alican Özkan, alican.ozkan@wyss.harvard.edu

Keywords: Hepatocellular Carcinoma, Cirrhosis, Desmoplasia, Chemoresistance, Hypoxia, 3D cell culture, CYP Metabolism, Tissue Engineering, Drug Metabolization

Abstract

Variations in tumor biology from patient to patient combined with the low overall survival rate of hepatocellular carcinoma (HCC) present significant clinical challenges. During the progression of chronic liver diseases from inflammation to the development of HCC, microenvironmental properties, including tissue stiffness and oxygen concentration, change over time. This can potentially impact drug metabolism and subsequent therapy response to commonly utilized therapeutics, such as doxorubicin, multi-kinase inhibitors (e.g., sorafenib), and other drugs, including immunotherapies. In this study, we utilized four common HCC cell lines embedded in 3D collagen type-I gels of varying stiffnesses to mimic normal and cirrhotic livers with environmental oxygen regulation to quantify the impact of these microenvironmental factors on HCC chemoresistance. In general, we found that HCC cells with higher baseline levels of cytochrome p450-3A4 (CYP3A4) enzyme expression, HepG2 and C3Asub28, exhibited a cirrhosis-dependent increase in doxorubicin chemoresistance. Under the same conditions, HCC cell lines with lower CYP3A4 expression, HuH-7 and Hep3B2, showed a decrease in doxorubicin chemoresistance in response to an increase in microenvironmental stiffness. This differential therapeutic response was correlated with the regulation of CYP3A4 expression levels under the influence of stiffness and oxygen variation. In all tested HCC cell lines, the addition of sorafenib lowered the required doxorubicin dose to induce significant levels of cell death, demonstrating its

potential to help reduce systemic doxorubicin toxicity when used in combination. These results suggest
that patient-specific tumor microenvironmental factors, including tissue stiffness, hypoxia, and
CYP3A4 activity levels, may need to be considered for more effective use of chemotherapeutics in
HCC patients.

1   Introduction

Cancer is the second-highest cause of mortality in the United States, lagging just slightly behind
cardiovascular disease in 2019 [1]. Among all cancer types, hepatocellular carcinoma (HCC) has the
second-lowest 5-year survival rate (17.7%) and has shown the highest increase in mortality among all
cancers over the past seven years [2],[3]. Complicating treatment, HCC is commonly diagnosed at an
intermediate or an advanced stage and often occurs secondary to underlying chronic liver disease and
cirrhosis [4]. The prognosis is poor as treatment options are limited by compromised liver function due
to underlying disease. Despite screening efforts for at-risk patients, most are not surgical candidates
for partial resection, and the availability of full liver transplantation is very low relative to the need[5].

These issues mean that systemic and localized drug-based therapies play a significant role in current
standard therapy for HCC. Despite these therapeutic interventions, the survival rate for HCC remains
low, partially attributed to the variable efficacy of current treatment methods based on underlying
factors [6–8]. As such, stratifying patients for the most effective treatment is critical because of three
factors; the degree of tumor burden, the degree of liver dysfunction, and highly variable treatment
efficacy between patients [9]. For example, the tyrosine kinase inhibitor sorafenib has shown modest
success in selected patients as a systemic treatment [10], but effectiveness is tempered by poor tolerance
of the drug in many instances[9]. Localized delivery of drugs through transarterial chemoembolization
(TACE) combines delivery of drugs such as doxorubicin with embolization to promote localized
ischemia and hypoxia. TACE blocks the arterial blood supply of a tumor through particulate or viscous
liquid agents such as degradable starch microspheres, drug-eluting beads, or ethiodized oil. This is a
well-established technique that allows a high local dose while simultaneously increasing the residence
of chemotherapeutic drugs in the target area, cutting off the supply of nutrients, and also limiting
exposure and toxicity for the rest of the body [11]. This has emerged as the standard of care for
intermediate-stage HCC. However, tumor cells in the hypoxic environment may undergo phenotypic
adaptations that aid survival. Such changes may account for the high rate of persistent, viable tumor
cells observed after TACE in previous studies [12]. Therefore, while a substantial survival benefit can be
realized, there is still much room for improvement and understanding of the changes that occur in the
tumor cells during embolization[6–8].

It is well established that many of the difficulties in treating HCC may stem from the numerous tumor
microenvironment (TME) changes that occur in underlying chronic liver disease and the rapid
progression of HCC. The modulation of the TME has been shown to impact drug metabolism
significantly and is thought to be a major contributor to the known differential response of patients to
chemotherapy [13],[14]. Furthermore, induction of hypoxia in the TME due to stiffening of the extracellular
matrix (ECM) and embolization during treatment can alter the chemoresistance of the tumor cells,
further impacting the treatment efficiency in intermediate and advanced stage HCC [15–17]. The
modulation of response and the individual impact of these TME features have yet to be fully
characterized in a three-dimensional (3D) HCC-TME model.

The increase in microenvironmental stiffness resulting from fibrosis, usually culminating in cirrhosis,
is a hallmark of most chronic liver diseases and is observed in 80-90% of HCC patients [4]. The most
notable hallmark of liver cirrhosis that impacts cellular and tissue function is increased collagen

deposition from activated hepatic stellate cells, increasing the stiffness and compression modulus of
liver tissue [18]. Similarly, the progression of HCC is marked by further localized stiffening. This
desmoplastic reaction is attributed to further differentiation of liver stellate cells into myofibroblasts
[18], resulting in additional deposition of collagen[19]. This stiffening of the ECM, in conjunction with high
tumor cell density, further reduces local oxygen and nutrient diffusion. Limited oxygen availability has
been shown to alter the outcomes of the chemotherapeutic treatments by affecting the drug transporters'
p-glycoprotein (MDR1, multidrug resistance 1), drug targets (topoisomerase II), or by initiating drug-
induced apoptosis [20]. Subsequent alterations in the cancer cells' response to chemotherapy can occur
through modulation of chemoresistance markers and hepatocyte metabolic enzymes such as
cytochrome P450 (CYP450), primarily the CYP3A4 subgroup [21]. Thus CYP3A4 expression can
potentially serve as an indicator for predicting chemotherapeutic response [22]. This highlights a potential
mechanism of differential tumor chemoresistance through the modulation of CYP3A4 under different
microenvironmental conditions.

Numerous *in vitro* models have been used to study the impact of TME modulation on HCC
treatment response. Two-dimensional (2D) cell monolayers have documented an increase in HepG2
cell survival following exposure to doxorubicin during hypoxia compared to normoxic conditions [23].
However, traditional 2D cell culture models do not allow for adequate representation of the
physiological diffusion and associated transport barriers found in the 3D extracellular
microenvironment, limiting clinical translation. Tumor spheroid models have been utilized as a more
representative system for assessing HCC drug response in direct cell-cell contacts and the subsequent
decrease of HCC chemoresistance [24]. However, the lack of ECM in these models severely curtails the
study of the interactions with and subsequent tuning of the ECM components of the TME [25].

Previous efforts that address the importance of ECM microenvironmental properties on the
chemotherapeutic response, employ tunable hydrogels models, composed of cells cultured in ECMs of
collagen, fibrin, alginate, and Matrigel[TM]. These models have been used to investigate the ECM's role
in *in vitro* studies of chemotherapy response, drug transport, cell invasion, and differentiation [25,26].
Culturing different breast cancer cell lines in an alginate hydrogel showed doxorubicin
chemoresistance was altered in particular phenotypes when ECM stiffness was increased [27]. With
respect to metabolic activity, one study investigated Ifosfamide metabolism by C3A HCC cells with
different levels of CYP3A4 expression when cultured in polylactic acid (PLA) [28]. However, this study
reported treatment efficacy only with glioblastoma cells and did not explore or discuss the treatment
response of liver cells. Another study demonstrated that culturing Caco-2 colorectal cancer cells on the
top of a 3D collagen-Matrigel[TM] blended hydrogel with dynamic flow conditions increased CYP3A4
expression drastically compared to 2D monolayers without flow [29]. Similarly, another study showed
that the CYP3A4 activity of HepaRG HCC cells increased when cells were cultured in hyaluronan-
gelatin or wood-derived nanofibrillar cellulose ECMs relative to 2D monolayer culture [30]. Research
has shown culturing U251 and U87 glioblastoma cells in 3D PLA scaffold under hypoxia exhibited
higher resistance to doxorubicin and greater production of basic fibroblast growth factor (bFGF) and
vascular endothelial growth factor (VEGF)[31]. Furthermore, U87, U251, and SNB19 glioblastoma cells
have been shown to be more resistant to temozolomide when cultured in scaffold-free spheroids under
hypoxic conditions compared to comparable spheroids under normoxic conditions [32]. Other studies
have used native ECM (collagen and fibrin) and non-native polymers (agar, acrylamide, and polylactic
acid) to recapitulate 3D breast and hepatic tumor microenvironments. However, these studies did not
investigate the regulation of cellular metabolism, including CYP3A4 activity, under different
microenvironmental conditions [26,27,33,34]. Recent work extending these efforts has demonstrated
significant potential in utilizing collagen tunability to replicate native microenvironment properties

(pH, stiffness, fiber properties, and porosity) [35,36], demonstrating promise in replicating physiological
TME to investigate the modulation of CYP3A4 activity and chemoresistance.

In this study, we utilized a collagen type I hydrogel to investigate the influence of ECM stiffness,
oxygen concentration, cell type, and the availability of a 3D microenvironment (2D vs. 3D) on drug
metabolism and response to two common chemotherapeutic agents, doxorubicin and sorafenib, both
individually and in combination. We used four established HCC cell lines (HepG2, C3Asub28, HuH-
7, and Hep3B2), which present with different basal metabolic profiles of CYP3A4 expression to
quantify the impact of modulations of the TME. Our results demonstrate the importance of the
contribution of a 3D ECM in drug design and dosing based on the significant differences seen in drug
response and metabolism when tumor cells are cultured in 3D collagen type I compared to traditional
2D culture. Further, we show that variations in the liver TME, including liver stiffness and hypoxia,
results in altered drug metabolism and subsequent drug efficacy. Tissue stiffness, varied by using
collagen concentrations comparable to healthy and cirrhotic liver stiffnesses [37], caused an alteration in
chemoresistance and drug metabolism. Environmental oxygen regulation to simulate normoxic and
hypoxic conditions produces a similarly significant, general effect on both chemoresistance and drug
metabolism. However, the TME regulation was not consistent for every cell type investigated, shown
in the heterogeneous regulation of cell chemoresistance and drug metabolism in the HCC cell lines.
This highlights a potential clinically translational impact of HCC genetic polymorphisms and different
etiologies on treatment outcome. Specifically, we identify that the basal cellular CYP3A4 metabolism
can be differentially regulated by TME hypoxia and tissue stiffness, thus impacting the efficacy of
commonly used HCC therapeutics. This relationship provides a potential explanation of the poor
outcomes of drugs in HCC clinical trials and may eventually lead to improve outcomes for HCC
patients.

2   Methods

2.1   Cell Culture

Human hepatocellular carcinoma cell lines HepG2 (HB-8065™, ATCC®, Manassas, VA), HuH-7,
Hep3B2.1.7, and HepG2 derived C3A with enhanced expression of CYP3A4 mRNA and CYP3A4-
mediated activity (C3Asub28) [38] were used in this study. HuH-7 cells express a mutated form of tumor-
suppressive protein p53, leading to an increased half-life and accumulation of the protein in cell nuclei.
This has been shown to correlate with increased chemoresistance [39]. Hep3B2.1.7 was used as an
example of an HCC tumor with hepatitis B DNA in the genome and subsequent mutations, including
partially deleted and suppressed expression of p53. All cells were cultured with DMEM supplemented
with 10% heat-inactivated fetal bovine serum (FBS, F4135, Sigma Aldrich, MO) and 1%
Penicillin/Streptomycin (P/S, Invitrogen, CA). Normoxic conditions were similarly generated by
culturing cells in standard cell culture conditions in a normoxic incubator (21% O_2, 5% CO_2, 37˚C
Thermo Fisher Scientific, Rochester, NY, USA). Hypoxic conditions were simulated by placing cells
in a sealed incubator (1% O_2, 5% CO_2). All cells were grown to approximately 70% confluence and
used within the first eight passages.

2.2   Preparation and Tuning of Collagen

Type I collagen isolated from rat tail tendons (donated by the University of Texas at Austin - Institute
for Cellular and Molecular Biology) was used to recapitulate the tissue microenvironment as it is the
primary ECM component of human tissue, including the liver [26]. As we have previously published, a
stock solution of type I collagen was prepared by dissolving excised rat tail tendons in an HCl solution
at a pH of 2.0 for 12 hours at 23˚C [26]. The solution was then centrifuged at 4˚C for 45 minutes at 30000

g, and the supernatant was collected, lyophilized, and stored at -20°C. The lyophilized collagen was
mixed with diluted 0.1% glacial acetic acid, maintained at 4°C, and vortexed every 24 hours for three
days to create a collagen stock solution. Finally, collagen was centrifuged at 4°C for 10 minutes at 2700
rpm to remove air bubbles. Collagen concentrations of 4 and 7 mg/ml were used to replicate normal
and cirrhotic liver stiffness, respectively, which we have previously demonstrated to match native liver
compression moduli after cells have reached their native morphology [26,37]. Collagen solutions were
adjusted to pH 7.4 with 1X DMEM, 10X DMEM (Sigma Aldrich, St. Louis, MO), and 1N NaOH
(Fisher Scientific, Pittsburgh, PA.). Following this, the collagen mixture was mixed with the intended
HCC cell lines uniformly at a concentration of $1x10^6$ cells/ml. Each suspension was dispensed as 50
μL aliquots in 96 well plates and allowed to polymerize. For 2D monolayer samples, identical numbers
of cells in 100 μL of media were dispensed into the wells of a 96 well plate. Cell media was changed
every two days.

2.3   Confined Compression Test

Cirrhotic stiffening in the TME has been shown to alter the chemoresistance of many cancer cell types
[40]. As a result, the mechanical properties of the TME, such as compression modulus, also increase. The
compression modulus of 3D collagen hydrogels was measured with quasi-steady uniaxial unconfined
compression (Instron, Norwood, MA) [26] to ensure there is no significant difference between collagen
gels with different HCC cell lines three days after seeding. In the analysis, the hydrogels are assumed
to be linear under the deformation conditions, and the slope of the stress-strain curve represents the
compression modulus. Hydrogels were prepared as described in the previous section, and 500 μL of
hydrogels were placed inside 24 well plates. Polymerized collagen samples were punched (9.53 mm
diameter) to remove the concave meniscus at the sample edge. Samples were compressed using a 20
mm diameter load cell of a flat steel surface. Load cells (10 N Static Load Cell, 2519-10N) were
lowered approximately 2.5 mm away from the flat surface and displaced 2 mm at a rate of 0.0085 mm/s
to achieve 0.1% strain/s over the range of 0-20% strain. Stress was calculated from the force response
divided by the initial area of collagen sample (71.26 mm^2). All measurements were performed at room
temperature (23°C) and the total duration of each experiment was less than 4 minutes. The data was
analyzed using Matlab® (MathWorks, Natick, MA).

2.4   Dosing With Chemotherapeutics

Doxorubicin and sorafenib have been commonly used in ongoing clinical trials for HCC treatment
either alone or in combination [41]. Prepared samples were exposed to doxorubicin (D1515, Sigma-
Aldrich, St. Louis, MO) with or without sorafenib (HY-10201, MedChemExpress LLC, Monmouth
Junction, NJ,) for 24 and 48 h. A broad range (1 nM – 200μM) of doxorubicin concentrations were
used to determine the response across the different HCC cell lines. Three different doses of sorafenib
were used to replicate previously tested effective concentrations: none (0 μM), standard (11 μM), and
high (22 μM) [42]. Samples were washed with warm (37°C) 1x phosphate buffered saline (PBS) three
times to remove the excess drug after the treatment. Cell culture media was used as a negative control.

Measuring Viability: Cell viability 72 hours after the completion of drug treatment was measured with
Cell Titer Blue (G8081, Promega, Fitchburg, WI) cell viability assay to quantify the response of HCC
cells to chemotherapeutic treatment under varying TME conditions (such as stiffness and hypoxia).
Briefly, cell media was mixed at a ratio of 5:1 with assay solution and incubated at 37°C for 1 hr. A
Cytation 3 plate reader was used to read fluorescence (Ex: 530 nm/Em: 620 nm) of the assay. Findings
were normalized to control (cells treated with drug-free media) to obtain percent cell viability. The
data gathered from cell viability assays after doxorubicin treatment was fitted using the cftool function
of Matlab® (MathWorks, Natick, MA) and half-maximal inhibitory concentration (IC50) value was

calculated and reported as chemoresistance indicator. As sorafenib has been used as a prodrug used for
angiogenesis treatment, mainly by inhibiting vascular endothelial growth factor (VEGFR), platelet-
derived growth factor receptor (PDGFR), and rapidly Accelerated Fibrosarcoma (RAF), but it also has
a minor direct cytotoxic side-effect in some instances [15]. For that reason, treatment efficacy of
standalone sorafenib and combined treatments were reported as fold viability change compared to
untreated control for the two clinically relevant doses used, standard (11 μM), and high (22 μM).
Additionally, for combined treatment, the minimum required doxorubicin dose to initiate toxicity was
reported. For this analysis, statistical comparison ($p < 0.05$) of viability under the tested doses and
untreated samples were compared, and the minimum dose was reported.

2.5 CYP3A4 Activity Measurement

Overexpression of CYP3A4 has been shown to decrease the efficacy of chemotherapeutics and is one
of the main challenges in the treatment of HCC tumors [22]. To relate the chemotherapeutic response to
the metabolic activity of HCC cells under the influence of hypoxia and different stiffnesses, the
expression of CYP3A4 was quantified using P450-Glo™ Assay and Screening Systems (V9001,
Promega, Madison, WI) in response to different oxygen concentrations, normoxic and hypoxic, and
different spatial conditions, 2D, 3D normal, and 3D cirrhotic. HCC cells were cultured in 2D or 3D, as
previously described for three days before CYP3A4 measurements to allow cells to reach native
morphology [37]. Prepared samples were washed twice with PBS and incubated with 50 μL of CYP3A4
substrate Luciferin-IPA (3 μM, dissolved in 1X DMEM). After 1 hour of incubation at 37˚C, 50 μL of
Luciferin Detection Reagent was added and pipetted up and down several times to ensure cell lysis.
After 20 minutes of incubation at room temperature, cell supernatants were transferred to a 96-well
opaque white luminometer plate (white polystyrene; Costar, Corning Incorporated) and luminescence
was measured using a Cytation 3 plate reader (BioTek Instruments, Inc., VT). Reagents without cells
were included as background controls. Metabolic activities were calculated by subtracting the
background luminescence and normalizing to the seeded cell number.

2.6 Statistical Analysis

248 Two-tailed student's t-test assuming unequal variance was performed in Matlab® (MathWorks, Natick,
249 MA) to compare samples, and a p-value less than 0.05 was considered significant for variation. Data
250 are reported as mean ± standard deviation unless otherwise indicated. All experiments were replicated
251 a minimum of four times.

3 Results

253 HCC cells were cultured for three days to reach native morphology in a monolayer in a tissue culture
254 plate (2D) or in rat tail-derived collagen type I hydrogels (3D). Cells were treated with doxorubicin
255 with or without sorafenib for 24 or 48 hours. Viability was assessed 72 hours after the end of drug
256 treatment, as described in Figure 1a.

Figure 1: HCC cells cultured in a monolayer or in collagen hydrogels of varying stiffness. **a)** Experimental procedure outline. Cells were allowed to adhere and reach their native morphology before the treatment for 24 or 48 hours. **b)** Compression modulus of the HCC hydrogels with 4 and 7 mg/ml collagen concentrations, which replicate normal and cirrhotic tissues, respectively. The compression modulus did not vary when different HCC cells were cultured in collagen hydrogels. Compression modulus values were significantly different when different collagen concentrations were used to replicate normal and cirrhotic conditions. Dashed lines represent patient compression modulus of healthy (red) and cirrhotic (green) tissues. [43,44]. Selected collagen concentrations replicated native tissues successfully at different preload strains. * $p < 0.05$, ** $p < 0.01$.

3.1 Native 3D Microenvironment Compression Modulus

Hepatocytes uniformly embedded inside rat tail-derived collagen type I hydrogels were allowed to reach their native morphology for three days. Afterwards, stiffness of the collagen hydrogels was quantified using a uniaxial compression test with 5% and 10% preload strains as presented in Figure 1b to determine any potential impact of different HCC cell lines on collagen stiffness. Our results demonstrate that increasing collagen concentration significantly elevated the compression modulus of the collagen gels ($p < 0.05$). However, we found no significant difference between compression modulus between collagen hydrogels with the different HCC cell lines over the timeframe considered. Collagen hydrogels at 4 mg/ml concentration produce a compression modulus comparable to tissue in a normal hepatic microenvironment, which has been reported to be at 0.64 ± 0.08 kPa and 1.08 ± 0.16 kPa for 5% and 10% preload strains, respectively [43]. At 4 mg/ml collagen concentration, the average

compression modulus was found to be 0.66 ± 0.07 kPa and 0.11 ± 0.01 kPa for 5% and 10% preload
strains, respectively. Likewise, at 7 mg/ml collagen concentration, collagen hydrogels achieved a
compression modulus comparable to a human hepatic tumor microenvironment, which has been
reported to be 3 kPa under 5% preload strain [44]. The significant difference ($p<0.05$) of compression
moduli between 7 mg/ml and 4 mg/ml collagen hydrogels showed we could replicate the
microenvironment stiffness using these collagen properties. In our findings, using a 7 mg/ml collagen
concentration, the average compression modulus was found to be 2.70 ± 0.22 kPa for 5% preload strain.
The reported collagen compression moduli for this study are also consistent with our previously
reported collagen compression modulus values[26]. In the same study, we also showed that collagen
concentration does not alter the diffusivity of solutes, thereby demonstrating that differential response
to chemotherapy is not likely due to physical diffusion differences of the collagen concentrations [26].

3.2   **Matrix Stiffness and Hypoxia Modulate Doxorubicin Chemoresistance in Hepatic 2D and**
3D Cultures

HCC cell viability in response to doxorubicin treatment under different microenvironmental conditions
was measured. The resulting half-maximal inhibitory concentrations (IC50) of doxorubicin was
quantified for HepG2, C3Asub28, HuH-7, and Hep3B2 cells in 2D monolayers and 3D collagen
hydrogels at 4 mg/mL (3D-normal) or 7 mg/mL (3D-cirrhotic) to demonstrate the impact of the tumor
structural microenvironment on the hepatic response to this drug. Furthermore, the impact of oxygen
concentration was quantified through the environmental regulation of normoxic (21% O_2) and hypoxic
(1% O_2) conditions for 2D and 3D cultures. The IC50 values of doxorubicin after 24-hour treatments
are summarized in Figure 2. IC50 values for 48-hour treatment durations can be found in Fig S-I as
minimal changes to IC50 values were observed with the increased treatment duration. Overall, we
demonstrated that the IC50 values of doxorubicin against C3Asub28 cells were the highest compared
to other HCC cells, in agreeance with the sub-strain's increased drug metabolism through CYP3A4.

Figure 2: Half-maximal inhibitory concentrations (IC50) values of doxorubicin against HCC cells in different microenvironments. HCC cells have stiffness and oxygen-dependent resistance/sensitivity to 24-hour doxorubicin treatment. Cirrhosis and hypoxia altered IC50 findings differently among the HCC cell lines. 3D-normal: 4 mg/ml collagen gel, 3D-Cirrhosis: 7 mg/ml collagen gel. Statistical significance was compared to normoxic conditions. *p<0.05.

3.3 Comparison of Doxorubicin IC50 for 2D vs. 3D Culture under Normoxic Conditions

Studies have demonstrated that 3D matrix stiffness plays an important role in modulation of the cellular response to chemotherapeutics in some cancer cell lines, providing a significantly different response compared to standard 2D culture methods [27,45]. However, few groups demonstrate the variation of IC50 values of doxorubicin between standard 2D-monolayer and 3D culture under varying microenvironmental conditions. In the context of the 3D microenvironment, we were able to determine the influence of normal liver stiffness (4 mg/mL collagen) under normoxic conditions (21% O_2) compared to 2D-normoxic monolayers as calculated by the fold change in chemoresistance due to the presence of the 3D microenvironment. In general, we demonstrate that IC50 values of doxorubicin against HepG2 increased in 3D compared to 2D but did not change for the C3Asub28 cell line. In contrast, 3D culture decreased IC50 values of doxorubicin for HuH-7 and Hep3B2 cell lines compared to 2D monolayers.

We observed that IC50 of HepG2 cells after a 24-hour doxorubicin treatment was 3.09-fold higher in
the 3D normal-normoxic environment (IC50 = 1.13 μM) compared to the 2D-normoxic (IC50 = 3.48
μM). The IC50 of doxorubicin against C3Asub28 cells was consistently the highest, but it was the only
cell line that did not lead to a significant change in IC50 of doxorubicin in response to the 3D normal-
normoxic compared to 2D-normoxic. The efficacy of doxorubicin on HuH-7 cells cultured in 3D
normal-normoxic microenvironment was higher than the HepG2 and C3Asub28 phenotypes. Overall,
we observed that the IC50 of doxorubicin against HuH-7 cells was lower ($p<0.05$) in 3D normal-
normoxic compared to 2D-normoxic by 62.61-fold. Conversely, IC50 values of doxorubicin against
Hep3B2 cells were the lowest overall and significantly decreased 4.02-fold 3D normal-normoxic
compared to 2D-normoxic.

3.4   The Influence of Microenvironmental Stiffness on Doxorubicin Chemoresistance under
331   **Normoxic Conditions**

To isolate the impact that microenvironmental stiffness plays in the regulation of chemoresistance of
different HCC cell lines, we analyzed the impact that the shift from normal (4 mg/mL) to cirrhotic (7
mg/mL) collagen concentration has on doxorubicin IC50 values under normoxic conditions for a 24-
hour treatment duration. Overall, we found that the increase in microenvironmental stiffness, modeled
by the higher collagen concentration, increased the IC50 values of doxorubicin against the HCC cell
lines, HepG2 and C3Asub28, that had a higher basal chemoresistance to doxorubicin. Variations in
fold changes for 24-hour treatment were reported in Figure S-II. 48-hour treatment duration resulted
in marginal differences in IC50 values compared to 24-hour treatment duration (Figure S-III). The
IC50 of doxorubicin against HepG2 (3.24-fold) and C3Asub28 (2.22-fold) cells cultured in 3D
cirrhotic-normoxic conditions increased compared to cells cultured in 3D normal-normoxic conditions.
However, for both HuH-7 and Hep3B2.17, we did not see any significant difference in IC50 values of
doxorubicin against these cells ($p>0.05$), between 3D normal-normoxic and 3D cirrhotic-normoxic
conditions.

3.5   Influence of Hypoxia on Doxorubicin Chemoresistance

Oxygen concentration in HCC liver tumors can change due to reduced blood flow, increased
cell density, and environmental stiffening [17,23]. This has been shown to alter both the proliferation rate
and chemoresistance of tumor cells. We isolated and quantified the influence of hypoxia on
chemoresistance, as measured by IC50 values, when cells were cultured in 3D with normal liver
stiffness (4 mg/ml) and in 2D monolayers both under hypoxic conditions. Overall, the introduction of
hypoxic conditions to 2D monolayers or 3D collagen hydrogels with normal stiffness (4 mg/ml)
showed a change in response between the HCC cell lines for 24-hour treatment durations and saw
minimal changes for 48-hour treatment durations as shown in Figure S-III. Hypoxia consistently
increased IC50 values of doxorubicin against HepG2 cells but decreased IC50 values of doxorubicin
against C3Asub28 IC50 compared to normoxic conditions for all stiffnesses. The response of HuH-7
and Hep3B2 cells to doxorubicin under hypoxia was variable depending on if they were cultured in 2D
or 3D.

We next isolated the impact that hypoxia plays in the regulation of chemoresistance of different HCC
cell lines in 3D microenvironments. We first analyzed the effect that the shift from 4 mg/mL to 7
mg/mL collagen concentration had on doxorubicin IC50 values under hypoxic (1% O_2) conditions and
then quantified the impact that the shift from normoxia to hypoxia had within each collagen
concentration. Within both normal (4 mg/mL) and cirrhotic (7 mg/mL) liver stiffness, the introduction
of hypoxia resulted in a general increase in doxorubicin IC50 values, except for C3Asub28 which
shows the opposite trend. However, we did not observe a statistical significance of doxorubicin IC50

values when Hep3B2 cells cultures in 3D normal-hypoxic or 3D cirrhotic-hypoxic conditions relative
to their normoxic counterparts. All of the observed trends in IC50 fold changes suggested that HCC
cells, depending on the underlying phenotype (denoted by the different HCC cell lines utilized), can
display a differential change in IC50 values of doxorubicin dependent upon microenvironmental
stiffness and oxygen availability.

3.6   Matrix Stiffness and Hypoxia Modulate Cell Viability in Response to Sorafenib Treatment in Hepatic 3D Cultures

Sorafenib is not primarily utilized to induce direct cell death, its mechanism is to inhibit kinases
responsible for promoting angiogenesis and cell growth, mainly VEGFR, PDGFR, and RAF[15]. As
such, even high doses are not sufficient to terminate 50% of the HCC population rendering the
calculation of IC50 values impossible. We investigated the direct cytotoxic impact of sorafenib to
establish baseline values for more clinically relevant combined therapeutic administration of sorafenib
and doxorubicin. Clinically relevant standard (11 μM) and high (22 μM) doses of sorafenib reported
in previous studies were tested on four different HCC cell lines [42]. We observed many of the same
trends previously observed for doxorubicin chemoresistance (Figure 3). We note that C3Asub28
demonstrated the lowest levels of cell death. We also demonstrate that in C3asub28 cells that an
increase in microenvironmental stiffness correlated to a general increase in sorafenib chemoresistance.
However, the effects of hypoxia were more varied, and sorafenib chemoresistance differed depending
on the cell line, sorafenib dose, and microenvironmental stiffness. In our comparison of sorafenib
treated cells in 3D to the untreated 3D controls, we established that cell viability is not significantly
impacted at standard, 11 μM, Sorafenib doses for many of the cell lines regardless of matrix stiffness
or oxygen concentration at 24-hour treatment durations. However, higher doses of sorafenib, 22 μM,
shows a much higher direct cytotoxic effect across all HCC cell lines and microenvironmental
conditions tested. The influence of hypoxia on sorafenib chemoresistance has a markedly different
dynamic than what we have previously seen with doxorubicin.

3.7   Sorafenib Improves Minimum Required Dose of Doxorubicin for Acute Toxicity

Combined administration of doxorubicin and sorafenib in HCC patients has been commonly used due
to their potential synergistic effect [41,46]. Doxorubicin is used to inhibit tumor proliferation[47], while
sorafenib is used to inhibit angiogenesis and tumor cell proliferation in the TME [15]. Cells cultured in
3D normal and cirrhotic hydrogels were investigated for the impact of doxorubicin-sorafenib
combination therapy to determine how the introduction of sorafenib influenced the resulting cell
viability. Cell viability in normal and cirrhotic tissue with a 24-hour treatment duration of both drugs
and the influence of hypoxia are presented in Figure 3 and discussed in this section. Treatment efficacy
is reported as fold viability change compared to untreated controls reported in Figure S-IV. In addition,
we explore how the addition of sorafenib can lower the minimum required doxorubicin dose to produce
a significant change in cell viability from untreated controls. For this analysis, statistical comparison
(p<0.05) of viability under tested doses and untreated samples were compared, and minimum dose to
induce cytotoxicity was reported. We found minimal differences between 24- and 48-hour treatments
with these drugs and present 48-hour treatment results in Figure S-V.

HepG2 cells treated with combination therapy demonstrated a general decrease in the minimum
required concentration of doxorubicin to initiate statistically significant cell death compared to the non-
drug treated control. Under 3D normal-normoxic conditions, the required minimum doxorubicin dose
decreased from 1 μM to 0.5 μM and 0 μM for standard and high doses of sorafenib, respectively.
Similarly, 3D cirrhotic-normoxic conditions with sorafenib reduced the minimum doxorubicin dose
from 10 μM to 1 μM for both standard and high sorafenib doses. Under hypoxic conditions, we found

that the trend held. HepG2 cells in the cirrhotic-hypoxic conditions in combination therapy showed a
decreased minimum doxorubicin concentration from 10 μM to 1 μM compared to 3D cirrhotic-
normoxic conditions for both standard and high doses of sorafenib. Under normal-hypoxic conditions,
the required minimum dose of doxorubicin decreased from 5 μM to 0.5 μM and 0 μM compared to 3D
normal-normoxic conditions for standard and high doses of sorafenib, respectively. Although the
HepG2 phenotype showed a higher chemoresistance to doxorubicin in hypoxic and cirrhotic culture
conditions relative to normoxic and normal, the combination with sorafenib significantly diminished
the required dose of doxorubicin to initiate toxicity.

The C3Asub28 cell line demonstrated the highest levels of chemoresistance out of all of the selected
HCC cell lines, however like all other cell lines, the combination of sorafenib with doxorubicin
considerably lowered the required doxorubicin dose required to induce a significant decrease in cell
viability compared to samples treated only with doxorubicin. Without sorafenib, 10 μM and 50 μM
doxorubicin concentrations were required to initiate tumor cell toxicity for normal-normoxic and
cirrhotic-normoxic microenvironments, respectively. Under hypoxia, the required dose of doxorubicin
decreased from 50 μM to 5 μM in cirrhotic-hypoxic conditions only. For the normal-hypoxic condition,
the minimum required concentration to initiate tumor cell toxicity did not change (5 μM) for the high
and standard doses of sorafenib compared to the control. On the other hand, for the cirrhotic-hypoxic
condition, doxorubicin effectiveness to initiate tumor cell toxicity improved by 2.82 ($p=0.05$) and
19.88-fold ($p=0.04$) for the standard and high doses of sorafenib, respectively compared to samples
treated only with doxorubicin. Given that the C3Asub28 cell line required the highest dose of
doxorubicin, the combination with sorafenib significantly lowered the required dose of doxorubicin to
initiate tumor cell toxicity.

HuH-7 cells under the normal-normoxic condition showed a decrease in the minimum required dose
of doxorubicin from 0.1 μM to 0.001 μM and 0 μM for a significant decrease in cell viability compared
to normal-hypoxic conditions for standard and high doses of sorafenib respectively. Under the
cirrhotic-normoxic condition, standard sorafenib dose did not improve the minimum required
doxorubicin concentration. However, the high dose of sorafenib (22 μM) improved the minimum
required doxorubicin concentration to initiate toxicity from 1 μM to 0.1 μM. For the normal-hypoxic
condition, the required minimum dose of doxorubicin to initiate toxicity decreased from 0.5 μM to 0
μM for both standard and high doses of sorafenib compared to normal-normoxic conditions. A similar
trend was observed or the cirrhotic-hypoxic condition, the minimum required dose of doxorubicin
decreased from 0.1 μM to 0 μM for both standard and high doses of sorafenib compared to cirrhotic-
normoxic conditions.

For Hep3B2 cells, the standard dose of sorafenib (11 μM) under normal-normoxic conditions caused
a decrease in the minimum required dose of doxorubicin for a significant decrease ($p=0.05$) in cell
viability by 1.33-fold compared to no sorafenib. However, only the high dose of sorafenib (22 μM)
treatment of Hep3B2 cells improved the required doxorubicin dose in the cirrhotic-normoxic condition
from 0.1 μM to 0 μM. In both normal-hypoxic and cirrhotic-hypoxic conditions, the combined
treatment showed an improved minimum doxorubicin concentration from 0.1 μM to 0 μM for both
standard and high doses of sorafenib to initiate toxicity in cells.

Figure 3: Combined doxorubicin and sorafenib treatment efficacy of HCC cell lines under the influence of hypoxia and cirrhosis. HCC cell types show sensitivity to combined sorafenib and doxorubicin treatment for 24 hours. Cells cultured in 3D normal and cirrhotic hydrogels in hypoxic and normoxic conditions. Cell viability was analyzed and plotted as a percentage of the untreated control. A combination of doxorubicin with sorafenib improves treatment efficacy. The influence of cirrhosis decreased HCC cell death.

3.8   Cirrhosis and Hypoxia Regulated HCC Metabolic Activity

The CYP3A4 enzyme is one of the major mechanisms of drug metabolism for cancer therapeutics,
including sorafenib and doxorubicin, in the liver. These enzymes can metabolize drugs before they
have the chance to cause their intended direct cytotoxic effects on the cells.[14]. CYP3A4 metabolic
activity of the HCC cell lines was measured in 2D monolayers and in 3D collagen I hydrogels in
response to different stiffness and hypoxic conditions. Regulation of CYP3A4 by cirrhosis and hypoxia
is presented in Figure 4. In agreement with the previously published literature, the C3Asub28 cell line
expressed much higher CYP3A4 expression than other HCC cell lines[38]. In our study, C3Asub28
CYP3A4 expression is 7.17 ± 2.73 fold higher than the HepG2 cell line, which is within the range of
previously published work (6.1 ± 0.2 fold) [38]. The introduction of hypoxia significantly downregulated
CYP3A4 expression compared to normoxia in all microenvironments (p=0.03). CYP3A4 expression
of C3Asub28 cells did not change with culture in 3D normal-normoxic and 3D normal-hypoxic
conditions than similar 2D conditions (p=0.78).

On the other hand, 3D cirrhotic-normoxic and cirrhotic-hypoxic culture upregulated CYP3A4
expression by 1.50 (p=0.007) and 1.65 (p=0.01) fold compared to 3D normal-normoxic and normal-
hypoxic conditions, respectively. HepG2 cell lines expressed significantly lower (p=0.02) CYP3A4
expression compared to the C3Asub28 cell line. However, unlike the C3Asub28 cell line, hypoxia
upregulated CYP3A4 expression in HepG2 cell lines compared to normoxia for both 2D and 3D
microenvironments (p=0.04). 3D normal-normoxic and normal-hypoxic culture upregulated CYP3A4
expression compared to 2D normoxic and 2D hypoxic culture by 1.90 (p=0.03) and 1.35 (p=0.003)
fold, respectively. Similarly, 3D cirrhotic-normoxic and cirrhotic-hypoxic culture upregulated
CYP3A4 expression compared to 3D normal-normoxic and normal-hypoxic stiffness by 2.13
(p=0.004) and 2.00 (p=0.0002) fold, respectively. HuH-7 cells expressed relatively lower values of
CYP3A4, and none was detected in 3D culture conditions. Furthermore, the Hep3B2 cell line did not
express any CYP3A4 expression in all culture conditions. These results show that HCC cells express
different basal levels of CYP3A4 and that this expression is can be directly regulated by TME oxygen
concentration and stiffness.

Figure 4: Regulation of CYP3A4 is responsible for drug metabolization by HCC cell lines in response to hypoxic and cirrhotic conditions. Presence of cirrhosis and hypoxia altered CYP3A4 activity differently between HCC cell lines. Data represented as mean ± standard deviation. * denotes significance for cirrhotic and normal 3D samples compared to the 2D monolayer. * $p < 0.05$, ** $p < 0.01$.

4   Discussion

In this study, we establish that chemoresistance can be regulated by hypoxic and cirrhotic conditions in the TME through direct modulation of CYP3A4 expression. This regulation can differentially alter the efficacy of chemotherapeutic drugs in HCC cell lines, which potentially has clinical translation to patient-specific HCC treatments. We examined the direct impact of TME stiffness and oxygen concentration, variables commonly associated with HCC tumors on cellular response to chemotherapeutics. This impact was measured by determining the difference in cell viability in response to chemotherapeutic treatment and regulation of the expression of the primary drug-metabolizing enzyme CYP3A4. Using a collagen-based hydrogel system, we observed that 3D culture alone significantly modulates resistance to doxorubicin and sorafenib in HepG2, C3Asub28, and HuH-7 cells but not for Hep3B2 cells. This stiffness-dependent resistance was not observed in similar Hep3B2 cultures, which warrants further investigation into potential phenotypic and genotypic differences in this specific cell line that might elucidate this response. This differential regulation of chemoresistance in different cell lines of the same cancer type is not unique to HCC. This phenomenon has also been observed in other cancers: The chemoresistance indicator, ethoxyresorufin, was upregulated in 3D culture conditions compared to 2D culture in the C3A cell line. This is in agreement with HepG2 and C3Asub28 doxorubicin treatment findings [48]. Equivalently, the IC50 of C3A cells after treatment with paracetamol, trovafloxacin, and fialuridine were found to be higher in 2D culture conditions compared to 3D culture conditions. This result corresponds with our HuH-7 and Hep3B2 doxorubicin chemoresistance findings [24]. Overall, these studies suggest that drug efficacy in 2D vs. 3D conditions depends on cell phenotype and drug type. This coincides with our findings in which HepG2 and C3Asub28 cell lines had greater chemoresistance in 3D culture compared to 2D but showed the converse trend for HuH-7 and Hep3B2 cells.

Cirrhosis and desmoplastic stiffening in the TME has been shown to be potential factors regulating drug chemoresistance [49]. In our study, HepG2 chemoresistance to sorafenib and doxorubicin increased

in response to cirrhotic stiffness relative to their culture in a matrix of normal stiffness. The C3Asub28
cell line demonstrated a higher chemoresistance to doxorubicin in response to cirrhotic conditions than
other tested HCC cells. Similar to the HepG2 cell line, a rise in stiffness also increased the
chemoresistance to doxorubicin of the C3Asub28 cell line. However, sorafenib alone did not alter the
cell viability of the C3Asub28 cell line for the considered doses potentially attributed to the cell line's
high baseline CYP3A4 metabolic activity. The resistance of HuH-7 and Hep3B2 cells to doxorubicin
was not altered in response to cirrhosis, whereas CYP3A4 expression of these cell lines did not change
in response to cirrhosis. However, the same cell lines were shown to have higher chemoresistance to
sorafenib in response to cirrhosis. Although CYP3A4 carries out the majority of metabolic activity in
hepatocytes, other minor cytochromes, such as CYP1A2, 2A6, 1A2, and 2C9, exist and may also alter
the drug chemoresistance to an as yet unknown extent. The additional CYP expression present in HCC
cells may be responsible for the differential effect of cirrhosis on chemoresistance differences between
sorafenib and doxorubicin [50]. Furthermore, studies in literature showed that a rise in stiffness does not
always increase the chemoresistance of cancer cells [49]. The increase in TME stiffness may improve
IC50 of MDA-MB-231 triple-negative breast cancer cells to doxorubicin [27]. However, the same study
showed that MCF-7 HER2+ breast cancer cells did not show a stiffness-dependent resistance to
doxorubicin. This study hypothesized the increase of stiffness altered chemoresistance differently
because MCF-7 remained in an epithelial phenotype, but MDA-MB-231 had a mesenchymal
phenotype. Similarly, stiffness induces chemoresistance of BxPC-3 and Suit2-007 pancreatic cancer
cells to paclitaxel, but not to gemcitabine [51]. These studies hypothesized the differential effect of
chemoresistance to different drugs between cell lines could be related to phenotypical differentiation
from epithelial to mesenchymal phenotype. In addition, HepG2, HuH-7, and Hep3B2 cell lines have
different phenotypic profiles and differentiation levels, potentially explaining differences in their
chemoresistance and metabolic activity in response to cirrhosis [52].

Hypoxia is known to be one of the regulating factors of chemoresistance [53]. In our study, HepG2
chemoresistance to doxorubicin and sorafenib increased in response to hypoxia compared to the
normoxic condition as measured both by increased IC50 values and CYP3A4 expression. However,
for C3Asub28 and HuH-7, chemoresistance decreased in response to hypoxia compared to normoxia.
Previous studies have also demonstrated the differential effect of hypoxia on drug efficacy, depending
on the cell line and phenotype. The presence of hypoxia has been shown to upregulate hypoxia-induced
factor (HIF1-α), but this alters the CYP isoforms differently in various medulloblastoma cell lines [54].
The molecular pathway still could not be explained in this study, but it has been hypothesized that
nuclear receptors, namely PPARα, PPARγ, or ER-α, as well as the constitutive androstane and
pregnane X receptors, have found to be altered differently under hypoxia [54]. In addition to this,
chemoresistance does not always increase in response to hypoxia [53]. The same study also showed the
regulation of chemoresistance under hypoxia is not universal between ovarian, renal, breast, lung, and
lymphoma cancer cell lines and varies for different drugs. This supports our data showing the
differential effects of hypoxia on doxorubicin IC50 values of the tested HCC cell lines [53]. Also, hypoxia
increases HepG2 chemoresistance to doxorubicin, in confirmation with our study, but not to rapamycin
[23]. In parallel to this, a significant decrease in apoptotic cells induced by cisplatin was reported under
hypoxic conditions for HepG2 and MHCC97L cell lines, which is in agreement with what we observed
with increased chemoresistance of HepG2 cells under hypoxic conditions [55]. It has also been showed
that hypoxia downregulates drug-metabolizing enzymes and subsequently the chemoresistance of the
HepaRG hepatoma cell line, which agrees with our findings of C3Asub28 CYP3A4 modulation under
hypoxia [56].

Consequently, the differential role of hypoxia on molecular chemoresistance expressions and drug
efficacy has been reported. It has been shown that HIF1-α is upregulated due to a lack of oxygen in

TME. This may or may not induce drug transporters such as MDR1 and targets of delivered drugs
(topoisomerase II) in each cell line [20]. Additionally, possible nuclear receptors have been proposed to
regulate CYP3A4 in response to hypoxia through the expression of HIF1-α and p53 expression[57].
Alternation of molecular drug transport mechanisms could be the reason why we observed variable
chemoresistance between different HCC cell lines under hypoxia.

Our study showed CYP3A4 expression is regulated by microenvironment stiffness and hypoxia for
HepG2, C3Asub28, and HuH-7 cell lines providing a potential mechanism connecting the TME to the
chemotherapeutic response. The regulation of CYP3A4 resulted in a significant impact on the efficacy
of doxorubicin and sorafenib, whose trends in the regulation mirror the observed changes in cell
viability in response to the drugs. Doxorubicin IC50 was higher, and sorafenib terminated less HCC
population for HepG2 and C3Asub28 cells in cirrhotic, 7 mg/ml, microenvironments in general
compared to healthy, 4 mg/mL, stiffness reflecting the CYP3A4 expression in those
microenvironments. However, we did not see any significant IC50 change in response to doxorubicin
for the HuH-7 cell line when cultured in normal and cirrhotic 3D microenvironments. In addition,
CYP3A4 expression of HuH-7 did not change when cultured in normal and cirrhotic 3D
microenvironments, which is parallel with IC50 findings.

Moreover, in general, we demonstrate that hypoxia increases doxorubicin IC50 against HepG2 cells
but C3Asub28 cells. The change of doxorubicin IC50 was parallel to the regulation of CYP3A4
expression. Hypoxia upregulated CYP3A4 expression of HepG2 cells but downregulated CYP3A4
expression of C3Asub28 cells. This work further confirms the upregulated metabolic expression of
CYP3A4 decreases doxorubicin efficacy in confirmation with previous studies across multiple cancer
types, including liver[58], colorectal[29], breast[59], and prostate[60]. Inhibition of CYP3A4 activity of human
primary hepatocytes has shown to suppress human pregnane X receptor (hPXR) agonist-induced
chemoresistance [61]. Overexpression of CYP3A4 in tumor tissues and chemoresistance to therapeutics
has been shown in clinical practices [62]. Similarly, there is evidence that therapeutic efficacy of drugs
have been diminished through CYP3A4 enzyme expressed by hepatocellular carcinoma cells[63]. These
studies agree with our findings on the regulation of chemoresistance based on CYP3A4 activity.
However, our work presents the significant finding that the regulation of CYP3A4 expression can be
directly tied to the tumor microenvironment. We demonstrate that CYP3A4 activity can be regulated
by oxygen concentration and TME stiffness, subsequently altering the metabolism of the
chemotherapeutic drugs in HCC cell lines. Possible nuclear receptor pathways regulating CYP3A4 in
response to hypoxia through HIF1-α, p53, PPARα, VDR, FXR, and LXR have been proposed and
hypothesized that hypoxia could affect CYP3A4 at different degrees [57]. Furthermore, TME stiffness
has been hypothesized to alter CYP3A4 differentially through yes-associated protein (YAP) pathway
[64]. This likely has a direct clinical translation to *in vivo* HCC regulation enforced by the clinical
observations of highly variable patient to patient HCC CYP3A4 expression [65]. The majority of the
current HCC treatments result in poor treatment outcomes, as such, the consideration of tumor
microenvironment properties (such as stiffness variation due to the changing fibrosis scores of patients,
presence of different levels of hypoxia as a result of this desmoplastic stiffening in the TME), and CYP
expression levels could potentially bring benefits to outcomes of HCC treatment, and provide a basis
for personalized HCC treatment [62,66].

Overall, this work demonstrates TME stiffness and oxygen concentration modulates CYP3A4
expression of HCC cells and, consequently, their chemoresistance to doxorubicin and sorafenib
treatment. We determined the existence of a stiffness-dependent resistance to doxorubicin and
sorafenib, depending on the differential genetics of the HCC, such as phenotypical changes from
epithelial to mesenchymal [27]. HepG2 and C3Asub28 cells showed a higher chemoresistance to

doxorubicin and sorafenib under cirrhotic conditions. Conversely, we did not observe a change in
chemoresistance for HuH-7 and Hep3B2 cell lines to doxorubicin in response to cirrhotic conditions,
which may be due to underlying genotypic differences including differentiation levels, which alter
metabolism pathways through glucose, glutamine, and glutamate [52]. Hypoxia demonstrated a much
different impact on the HCC cells, upregulating the chemoresistance of HepG2 and HuH-7, but
downregulating chemoresistance of the C3Asub28 cell line. We did not observe a significant variation
of chemoresistance in the Hep3B2 cell line in 3D culture. Drug metabolism, measured by CYP3A4
expression, mirrored effective chemoresistance measured by IC50 values in cell vitality assays. We
saw an increase in CYP3A4 expression in the 3D culture of C3Asub28 and HepG2 cell lines compared
to 2D, but this expression decreased for the HuH-7 cell line for both hypoxic and normoxic conditions.
At a minimum, the presence of a 3D culture system significantly modulates the response of HCC to
chemotherapy over the standard 2D methods. Previously it has been shown 3D culture alters the
integrin ligands (such as AKT and RAF), which are the targeting for doxorubicin and sorafenib,
differently among different cell lines compared to 2D culture [49].

Further, the stiffness of the microenvironment, seen in HCC patients with cirrhosis, modulates drug
resistance and should be taken into consideration when determining treatment options and doses. While
CYP3A4 maintains the majority of the drug metabolism in the liver, other enzymes such as CYP2A6,
1A2, and 2C9 has shown to have a minor contribution to drug metabolism [62], and provide further
insight in IC50 of Hep3B2 variation we observed between 2D and 3D microenvironments but not
captured a difference in CYP3A4 expression. Further expansion of this work is needed to investigate
the varied response between existing HCC lines or patient-specific primary tumor cells that might
provide insight into the known differential effectiveness of standard HCC treatments *in vivo* [66].
Potentially, other drugs used to treat HCC in clinical practice such as lenvatinib could be tested in this
system to observed its potential effect on HCC treatment [67].

5   Acknowledgment

The authors would like to thank Dr. Wei Li (Department of Mechanical Engineering, the University of
Texas at Austin) and Dr. Christopher Sullivan (Department of Molecular Sciences) for generously
gifting C3Asub28 cells and HuH-7, respectively. This work was completed with support from the
Veterans Health Administration and with resources and the use of facilities at the Central Texas
Veterans Health Care System, Temple, Texas. The contents do not represent the views of the U.S.
Department of Veterans Affairs or the United States Government. T.E.Y. is a CPRIT scholar in cancer
research.

6   Funding

The authors acknowledge the support of the Cancer Prevention Research Institute of Texas (CPRIT)
for funding part of this work through grants RR160005, National Cancer Institutes for funding through
R21EB019646, R01CA186193, R01CA201127-01A1, U24CA226110 and U01CA174706, National
Institute of Diabetes and Digestive and Kidney Diseases funding through awards R01DK082435 and
R01DK112803 and Department of Veterans Affairs Biomedical Laboratory Research and
Development Service funding through award BX003486.

7   Author contributions

Conceptualization: A.O. and M.N.R. conceived of the idea for the study. Supervision: E.N.K.C, T.E.Y.
and M.N.R. supervised the project. E.N.K.C., S.D., M.M., T.E.Y. and M.N.R provided feedback and
assistance with manuscript preparation. Investigation: A.O., D.L.S., and M.N.R were responsible for

performing the studies and analyzing the experimental data. Writing: A.O. wrote the initial draft of the
paper. All authors discussed the results and revised the manuscript.

8   References

1.   Grandhi MS, Kim AK, Ronnekleiv-Kelly SM, Kamel IR, Ghasebeh MA, Pawlik TM.
Hepatocellular carcinoma: From diagnosis to treatment. *Surg Oncol*. 2016;25(2):74-85.
doi:10.1016/j.suronc.2016.03.002

2.   Petrick JL, Kelly SP, Altekruse SF, McGlynn KA, Rosenberg PS. Future of hepatocellular
carcinoma incidence in the United States forecast through 2030. *J Clin Oncol*.
2016;34(15):1787-1794. doi:10.1200/JCO.2015.64.7412

3.   Siegel RL, Miller KD, Jemal A. Cancer Statistics, 2018. *CA CANCER J CLIN*. 2018;68(1):7-
30. doi:10.3322/caac.21387

4.   Davis GL, Dempster J, Meler JD, et al. *Hepatocellular Carcinoma: Management of an
Increasingly Common Problem*. Vol 21.; 2008. doi:10.1080/08998280.2008.11928410

5.   Marrero JA, Kulik LM, Sirlin CB, et al. Diagnosis, Staging, and Management of Hepatocellular
Carcinoma: 2018 Practice Guidance by the American Association for the Study of Liver
Diseases. *Hepatology*. 2018;68(2):723-750. doi:10.1002/hep.29913

6.   Abou-Alfa GK, Niedzwieski D, Knox JJ, et al. Phase III randomized study of sorafenib plus
doxorubicin versus sorafenib in patients with advanced hepatocellular carcinoma (HCC):
CALGB 80802. *J Clin Oncol*. 2016;34:192-192. doi:10.1200/jco.2016.34.4_suppl.192

7.   Grazie LM. et al. Chemotherapy for hepatocellular carcinoma: The present and the future, *World
J Hepatol*. 2017;9(21):9-21.

8.   Lohitesh K, Chowdhury R, Mukherjee S. Resistance a major hindrance to chemotherapy in
hepatocellular carcinoma: An insight. *Cancer Cell Int*. 2018;18(1):44. doi:10.1186/s12935-018-
0538-7

9.   LU L-C, CHEN P-J, YEH Y-C, et al. Prescription Patterns of Sorafenib and Outcomes of
Patients with Advanced Hepatocellular Carcinoma: A National Population Study. *Anticancer
Res*. 2017;37(5):2593-2599. doi:10.21873/anticanres.11604

10.   Llovet JM, Ricci S, Mazzaferro V, et al. Sorafenib in advanced hepatocellular carcinoma. *N
Engl J Med*. 2008;359(4):378-390. doi:10.1056/NEJMoa0708857

11.   Schicho A, Hellerbrand C, Krüger K, et al. Impact of Different Embolic Agents for Transarterial
Chemoembolization (TACE) Procedures on Systemic Vascular Endothelial Growth Factor
(VEGF) Levels. *J Clin Transl Hepatol*. 2016;4(4):288-292. doi:10.14218/jcth.2016.00058

12.   Marin HL, Furth EE, Olthoff K, Shaked A, Soulen MC. Histopathologic outcome of neoadjuvant
image-guided therapy of hepatocellular carcinoma. *J Gastrointest Liver Dis*. 2009;18(2):169-
176. Accessed June 11, 2019.

13.   De Montellano PRO. Cytochrome P450-activated prodrugs. *Future Med Chem*. 2013;5(2):213-

228. doi:10.4155/fmc.12.197

14.   Rodriguez-Antona C, Ingelman-Sundberg M. Cytochrome P450 pharmacogenetics and cancer.
Oncogene. 2006;25(11):1679-1691. doi:10.1038/sj.onc.1209377

15.   Keating GM, Santoro A. *Sorafenib: A Review of Its Use in Advanced Hepatocellular*
Carcinoma. Vol 69.; 2009. doi:10.2165/00003495-200969020-00006

16.   Xu W, Zhang X, Yu J, et al. Targeting the vasculature in hepatocellular carcinoma treatment:
Starving versus normalizing blood supply. *Clin Transl Gastroenterol*. 2017;8(6):e98.
doi:10.1038/ctg.2017.28

17.   Lencioni R, Petruzzi P, Crocetti L. Chemoembolization of hepatocellular carcinoma. *Semin*
Intervent Radiol. 2013;30(1):3-11. doi:10.1055/s-0033-1333648

18.   Koyama Y, Brenner DA. Liver inflammation and fibrosis. *J Clin Invest*. 2017;127(1):55-64.
doi:10.1172/JCI88881

19.   Affo S, Yu L-X, Schwabe RF. The Role of Cancer-Associated Fibroblasts and Fibrosisin Liver
Cancer. *Annu Rev Pathol Mech Dis*. 2016;12(1):153-186. doi:10.1146/annurev-pathol-052016-
100322

20.   Doktorova H, Hrabeta J, Khalil MA, Eckschlager T. Hypoxia-induced chemoresistance in
cancer cells: The role of not only HIF-1. *Biomed Pap*. 2015;159(2):166-177.
doi:10.5507/bp.2015.025

21.   Beckwitt CH, Clark AM, Wheeler S, et al. Liver 'organ on a chip.' *Exp Cell Res*.
2018;363(1):15-25. doi:10.1016/j.yexcr.2017.12.023

22.   Rodríguez-Antona C, Leskelä S, Zajac M, et al. Expression of CYP3A4 as a predictor of
response to chemotherapy in peripheral T-cell lymphomas. *Blood*. 2007;110(9):3345-3351.
doi:10.1182/blood-2007-02-075036

23.   Bowyer C, Lewis AL, Lloyd AW, Phillips GJ, MacFarlane WM. Hypoxia as a target for drug
combination therapy of liver cancer. *Anticancer Drugs*. 2017;28(7):771-780.
doi:10.1097/CAD.0000000000000516

24.   Webb SD, Colley HE, Sharma P, Murdoch C, Gaskell H, Williams DP. Characterization of a
functional C3A liver spheroid model. *Toxicol Res*. 2016;5(4):1053-1065.
doi:10.1039/c6tx00101g

25.   Khawar IA, Park JK, Jung ES, Lee MA, Chang S, Kuh HJ. Three Dimensional Mixed-Cell
Spheroids Mimic Stroma-Mediated Chemoresistance and Invasive Migration in hepatocellular
carcinoma. *Neoplasia*. 2018;20(8):800-812. doi:10.1016/j.neo.2018.05.008

26.   Antoine EE, Vlachos PP, Rylander MN. Tunable collagen I hydrogels for engineered
physiological tissue micro-environments. *PLoS One*. 2015;10(3).
doi:10.1371/journal.pone.0122500

27.   Brock A, Joyce MH, Suggs LJ, et al. Phenotypic Basis for Matrix Stiffness-Dependent

Chemoresistance of Breast Cancer Cells to Doxorubicin. *Front Oncol*. 2018;8:337. doi:10.3389/fonc.2018.00337

28. Ma L, Barker J, Zhou C, et al. Towards personalized medicine with a three-dimensional micro-scale perfusion-based two-chamber tissue model system. *Biomaterials*. 2012;33(17):4353-4361. doi:10.1016/j.biomaterials.2012.02.054

29. Kim HJ, Ingber DE. Gut-on-a-Chip microenvironment induces human intestinal cells to undergo villus differentiation. *Integr Biol*. 2013;5(9):1130-1140. doi:10.1039/c3ib40126j

30. Malinen MM, Kanninen LK, Corlu A, et al. Differentiation of liver progenitor cell line to functional organotypic cultures in 3D nanofibrillar cellulose and hyaluronan-gelatin hydrogels. *Biomaterials*. 2014;35(19):5110-5121. doi:10.1016/j.biomaterials.2014.03.020

31. Kim JW, Ho WJ, Wu BM. *The Role of the 3D Environment in Hypoxia-Induced Drug and Apoptosis Resistance*. Vol 31.; 2011.

32. Musah-Eroje A, Watson S. A novel 3D in vitro model of glioblastoma reveals resistance to temozolomide which was potentiated by hypoxia. *J Neurooncol*. 2019;142(2):231-240. doi:10.1007/s11060-019-03107-0

33. Adriani G, Pavesi A, Kamm RD. Studying TCR T cell anti-tumor activity in a microfluidic intrahepatic tumor model. In: *Methods in Cell Biology*. Vol 146. ; 2018:199-214. doi:10.1016/bs.mcb.2018.05.009

34. Ahn J, Ahn J, Yoon S, Nam YS, Son M, Oh J. Human three-dimensional in vitro model of hepatic zonation to predict zonal hepatotoxicity. *J Biol Eng*. 2019;5:1-15. doi:10.1186/s13036-019-0148-5

35. Özkan A, Stolley D, Cressman ENK, et al. The Influence of Chronic Liver Diseases on Hepatic Vasculature: A Liver-on-a-chip Review. *Micromachines*. 2020;11(5):487. doi:10.3390/mi11050487

36. Stolley DL, Crouch AC, Özkan A, et al. Combining chemistry and engineering for hepatocellular carcinoma: Nano-scale and smaller therapies. *Pharmaceutics*. 2020;12(12):1-18. doi:10.3390/pharmaceutics12121243

37. Ozkan A, Ghousifam N, Hoopes PJ, Yankeelov TE, Rylander MN. In vitro vascularized liver and tumor tissue microenvironments on a chip for dynamic determination of nanoparticle transport and toxicity. *Biotechnol Bioeng*. 2019;116(5):1201-1219. doi:10.1002/bit.26919

38. Küblbeck J, Reinisalo M, Mustonen R, Honkakoski P. Up-regulation of CYP expression in hepatoma cells stably transfected by chimeric nuclear receptors. *Eur J Pharm Sci*. 2010;40(4):263-272. doi:10.1016/j.ejps.2010.03.022

39. Hientz K, Mohr A, Bhakta-Guha D, Efferth T. *The Role of P53 in Cancer Drug Resistance and Targeted Chemotherapy*. Vol 8.; 2017. doi:10.18632/oncotarget.13475

40. Godugu C, Patel AR, Desai U, Andey T, Sams A, Singh M. AlgiMatrix^{TM} Based 3D Cell Culture System as an In-Vitro Tumor Model for Anticancer Studies. *PLoS One*. 2013;8(1).

doi:10.1371/journal.pone.0053708

41.   Abou-Alfa GK, Shi Q, Knox JJ, et al. Assessment of Treatment with Sorafenib Plus Doxorubicin
vs Sorafenib Alone in Patients with Advanced Hepatocellular Carcinoma: Phase 3 CALGB
80802 Randomized Clinical Trial. *JAMA Oncol*. 2019;5(11):1582-1588.
doi:10.1001/jamaoncol.2019.2792

42.   Coriat R, Nicco C, Chéreau C, et al. Sorafenib-induced hepatocellular carcinoma cell death
depends on reactive oxygen species production in vitro and in vivo. *Mol Cancer Ther*.
2012;11(10):2284-2293. doi:10.1158/1535-7163.MCT-12-0093

43.   Hey-Chi Hsu, Wen-Chun Yeh, Pei-Ming Yang, et al. Young's modulus measurements of human
liver and correlation with pathological findings. In: *2001 IEEE Ultrasonics Symposium.*
Proceedings. An International Symposium (Cat. No.01CH37263). Vol 2. IEEE; 2002:1233-
1236. doi:10.1109/ultsym.2001.991942

44.   Yeh WC, Li PC, Jeng YM, et al. Elastic modulus measurements of human liver and correlation
with pathology. *Ultrasound Med Biol*. 2002;28(4):467-474. doi:10.1016/S0301-
5629(02)00489-1

45.   Ozcelikkale A, Shin K, Noe-Kim V, et al. Differential response to doxorubicin in breast cancer
subtypes simulated by a microfluidic tumor model. *J Control Release*. 2017;266:129-139.
doi:10.1016/j.jconrel.2017.09.024

46.   Merle P, Blanc JF, Phelip JM, et al. Doxorubicin-loaded nanoparticles for patients with
advanced hepatocellular carcinoma after sorafenib treatment failure (RELIVE): a phase 3
randomised controlled trial. *Lancet Gastroenterol Hepatol*. 2019;4(6):454-465.
doi:10.1016/S2468-1253(19)30040-8

47.   Cox J, Weinman S. Mechanisms of doxorubicin resistance in hepatocellular carcinoma. *Hepatic*
Oncol. 2015;3(1):57-59. doi:10.2217/hep.15.41

48.   Amitay-Shaprut S, Cohen S, Elkayam T, Dvir-Ginzberg M, Harel T. Enhancing the Drug
Metabolism Activities of C3A— A Human Hepatocyte Cell Line—By Tissue Engineering
Within Alginate Scaffolds. *Tissue Eng*. 2006;12(5):1357-1368. doi:10.1089/ten.2006.12.1357

49.   Shin JW, Mooney DJ. Extracellular matrix stiffness causes systematic variations in proliferation
and chemosensitivity in myeloid leukemias. *Proc Natl Acad Sci U S A*. 2016;113(43):12126-
12131. doi:10.1073/pnas.1611338113

50.   Ashida R, Okamura Y, Ohshima K, et al. CYP3A4 Gene Is a Novel Biomarker for Predicting a
Poor Prognosis in Hepatocellular Carcinoma. *Cancer Genomics Proteomics*. 2017;14(6):445-
453. doi:10.21873/cgp.20054

51.   Rice AJ, Cortes E, Lachowski D, et al. Matrix stiffness induces epithelial-mesenchymal
transition and promotes chemoresistance in pancreatic cancer cells. *Oncogenesis*.
2017;6(7):352. doi:10.1038/oncsis.2017.54

52.   Nwosu ZC, Battello N, Rothley M, et al. Liver cancer cell lines distinctly mimic the metabolic
gene expression pattern of the corresponding human tumours. *J Exp Clin Cancer Res*.

2018;37(1). doi:10.1186/s13046-018-0872-6

53.   Strese S, Fryknäs M, Larsson R, Gullbo J. *Effects of Hypoxia on Human Cancer Cell Line*
Chemosensitivity. Vol 13.; 2013. doi:10.1186/1471-2407-13-331

54.   Valencia-Cervantes J, Huerta-Yepez S, Aquino-Jarquin G, et al. Hypoxia increases
chemoresistance in human medulloblastoma DAOY cells via hypoxia-inducible factor 1α-
mediated downregulation of the CYP2B6, CYP3A4 and CYP3A5 enzymes and inhibition of
cell proliferation. *Oncol Rep*. 2019;41(1):178-190. doi:10.3892/or.2018.6790

55.   Chi KL, Zhen FY, Ho DW, et al. An Akt/hypoxia-inducible factor-1α/platelet-derived growth
factor-BB autocrine loop mediates hypoxia-induced chemoresistance in liver cancer cells and
tumorigenic hepatic progenitor cells. *Clin Cancer Res*. 2009;15(10):3462-3471.
doi:10.1158/1078-0432.CCR-08-2127

56.   Boudjema K, Ishida S, Hori T, et al. Drug-metabolising enzymes are down-regulated by hypoxia
in differentiated human hepatoma HepaRG cells: HIF-1α involvement in CYP3A4 repression.
Eur J Cancer. 2009;45(16):2882-2892. doi:10.1016/j.ejca.2009.07.010

57.   Yuan X, Lu H, Zhao A, Ding Y, Min Q, Wang R. Transcriptional regulation of CYP3A4 by
nuclear receptors in human hepatocytes under hypoxia. *Drug Metab Rev*. 2020;0(0):1-10.
doi:10.1080/03602532.2020.1733004

58.   Zhou F, Lu L, Peng X, et al. Sorafenib Metabolism Is Significantly Altered in the Liver Tumor
Tissue of Hepatocellular Carcinoma Patient. *PLoS One*. 2014;9(5):e96664.
doi:10.1371/journal.pone.0096664

59.   Breslin S, O'Driscoll L. *The Relevance of Using 3D Cell Cultures, in Addition to 2D Monolayer*
Cultures, When Evaluating Breast Cancer Drug Sensitivity and Resistance. Vol 7.; 2016.
doi:10.18632/oncotarget.9935

60.   Chen TC, Sakaki T, Yamamoto K, Kittaka A. The roles of cytochrome P450 enzymes in prostate
cancer development and treatment. *Anticancer Res*. 2012;32:291-298.

61.   Abbott KL, Chaudhury CS, Chandran A, et al. Belinostat, at its clinically relevant
concentrations, inhibits rifampicin-induced CYP3A4 and MDR1 gene expressions. *Mol*
Pharmacol. 2019;95(3):324-334. doi:10.1124/mol.118.114587

62.   Molina-Ortiz D, Camacho-Carranza R, González-Zamora JF, et al. Differential expression of
Cytochrome P450 enzymes in normal and tumor tissues from childhood rhabdomyosarcoma.
PLoS One. 2014;9(4). doi:10.1371/journal.pone.0093261

63.   Hu DG, Mackenzie PI, Lu L, Meech R, McKinnon RA. Induction of human UDP-
Glucuronosyltransferase 2B7 gene expression by cytotoxic anticancer drugs in liver cancer
HepG2 Cells. *Drug Metab Dispos*. 2015;43(5):660-668. doi:10.1124/dmd.114.062380

64.   Speer JE, Wang Y, Fallon JK, Smith PC, Allbritton NL. Evaluation of human primary intestinal
monolayers for drug metabolizing capabilities. *J Biol Eng*. 2019;13(1). doi:10.1186/s13036-
019-0212-1

65.   Yan T, Lu L, Xie C, et al. Severely impaired and dysregulated cytochrome P450 expression and
836   activities in hepatocellular carcinoma: Implications for personalized treatment in patients. *Mol*
837   *Cancer Ther*. 2015;14(12):2874-2886. doi:10.1158/1535-7163.MCT-15-0274

838   66.   Rodríguez-Perálvarez M, Luong TV, Andreana L, Meyer T, Dhillon AP, Burroughs AK. A
839   systematic review of microvascular invasion in hepatocellular carcinoma: Diagnostic and
840   prognostic variability. *Ann Surg Oncol*. 2013;20(1):325-339. doi:10.1245/s10434-012-2513-1

841   67.   Chang Lee R, Tebbutt N. Systemic treatment of advanced hepatocellular cancer: new hope on
842   the   horizon.   *Expert*   *Rev*   *Anticancer*   *Ther*.   2019;19(4):343-353.
843   doi:10.1080/14737140.2019.1585245