Dynamics of sex-biased gene expression over development

in the stick insect *Timema californicum*

6   Jelisaveta Djordjevic[1], Zoé Dumas[1], Marc Robinson-Rechavi[1,2],

7   Tanja Schwander[1,a], Darren James Parker[1,2,a]

[1] Department of Ecology and Evolution, University of Lausanne, Lausanne,

Switzerland

[2] Swiss Institute of Bioinformatics, Lausanne, Switzerland

[a] equal contribution

Correspondence should be addressed to jelisaveta.djordjevic@unil.ch or

DarrenJames.Parker@unil.ch

Abstract

Sexually dimorphic phenotypes are thought to arise primarily from sex-biased gene expression during development. Major changes in developmental strategies, such as the shift from hemimetabolous to holometabolous development, are therefore expected to have profound consequences for the dynamics of sex-biased gene expression. However, no studies have previously examined sex-biased gene expression over development in hemimetabolous insects, precluding comparisons between developmental strategies. Here we characterized sex-biased gene expression at three developmental stages in a hemimetabolous stick insect (*Timema californicum*): hatchlings, juveniles, and adults. As expected, the proportion of sex-biased genes gradually increased over development, mirroring the gradual increase of phenotypic sexual dimorphism. Sex-biased genes identified at early developmental stages were generally consistently male- or female- biased at later stages, suggesting their importance in sexual differentiation. Additionally, we compared the dynamics of sex-biased gene expression over development in *T. californicum* to those of the holometabolous fly *Drosophila melanogaster* by reanalyzing publicly available RNA-seq data from third instar larval, pupal and adult stages. In *D. melanogaster,* 84% of genes were sex-biased at the adult stage (compared to only 20% in *T. californicum*), and sex-biased gene expression increased abruptly at the adult stage when morphological sexual dimorphism is manifested. Our findings support the prediction that the dynamics of sex-biased gene expression over development differ extensively between holometabolous and hemimetabolous insect species.

Introduction

Males and females often have divergent evolutionary interests, resulting in sex-specific selection pressures and ultimately the evolution of sexually dimorphic phenotypes (Lande 1980; Khila *et al.* 2012). Studies investigating how sexually dimorphic phenotypes are generated have shown the importance of differential gene expression between the sexes, suggesting that sex-specific selection is the major driving force behind the evolution of sex-biased gene expression (Chauhan *et al.* 2016; Mank 2017). The relationship between sexual dimorphism and sex-biased gene expression has been largely studied at the adult stage when sexual dimorphism is completely manifested and reproductive interests between males and females are most different (Pointer *et al.* 2013; Chauhan *et al.* 2016). However, sex-biased gene expression has also been found at early developmental stages of many species, well before any phenotypic sexual dimorphism becomes apparent (Lowe *et al.* 2015; Paris *et al.* 2015). This suggests that expression patterns in early developmental stages are also under sex-specific selection pressures (Ingleby *et al.* 2015).

Because we lack detailed studies of sex-biased gene expression over development, we do not know how consistent or dynamic the expression of the sex-biased genes is, nor how these dynamics relate to changes in phenotypic sexual dimorphism. For example, although we expect to see an overall increase in sex-biased gene expression over development (Ingleby *et al.* 2015; Mank 2017), it is not clear whether this results from the gradual increase of sex-bias of a set of genes or if sex-biased genes at early stages are largely different to those at later stages? Furthermore, sex-biased gene expression can be considered to reflect a broad measure of sexual dimorphism, including physiological and behavioral traits, and may therefore be more representative than dimorphism quantified using external

morphology. However, whether the extent of sexual dimorphism and of sex-biased

gene expression are generally correlated remains poorly known.

In insects there are two major developmental strategies, and developmental

patterns of sex-biased gene expression could differ between them. Holometabolous

insects have morphologically and ecologically distinct larval, pupal, and adult stages,

and phenotypic sexual dimorphism is commonly prominent only at the adult stage.

Contrastingly, hemimetabolous insects go through gradual morphological changes

and sexual differentiation, and the nymphal stages morphologically resemble adults

(Chen *et al.* 2010). These distinct dynamics for sexual morphological differentiation

are expected to be mirrored by sex-biased gene expression, with abrupt versus

gradual increases in the proportion of sex-biased genes over development. However,

studies of sex-biased gene expression during development have thus far only been

conducted in holometabolous insects (Magnusson *et al.* 2011; Rago *et al.* 2020), with

the main focus on *Drosophila* (Perry *et al.* 2014; Ingleby *et al.* 2015), precluding

comparisons between species with different developmental strategies.

Here, we studied the dynamics of sex-biased gene expression across

development in hemimetabolous insects, *Timema californicum.* These are sexually

dimorphic walking-stick insects, found in California (Vickery and Sandoval 2001).

Males are smaller than females, with different coloration, and have conspicuous cerci

used for holding on to the female during copulation. We selected three developmental

stages (Fig. 1a): the hatchling stage where sexes are phenotypically identical, the third

juvenile stage with minor phenotypic differences, and the adult stage, with pronounced

sexual dimorphism. We investigated whether sex-biased genes are recruited in a

stage-specific manner, and if the magnitude of sex-bias increases with development.

We further explored sequence evolution rates of sex-biased genes in pre-adult stages,

to determine if they show elevated rates as commonly observed for sex-biased genes

at adult stages in many species (Ellegren and Parsch 2007; Grath and Parsch 2012;

Perry *et al.* 2014).

We then compared the dynamics of sex-biased gene expression over

development in the hemimetabolous stick insect to those of the holometabolous fly

Drosophila melanogaster by reanalyzing publicly available RNA-seq data from 3rd.

instar larvae, pupal and adult stages, using the same pipeline as for *T. californicum.*

In *D. melanogaster*, both larval and pupal stages have little morphological sexual

dimorphism, while the adult stage has extensive dimorphism (Fig. 1b). We

hypothesized that sex biased gene expression follows the establishment of

morphological sexual dimorphism, with a gradual increase in sex-biased gene

expression over development in *T. californicum*, and an abrupt increase at the adult

stage in *D. melanogaster*.

Fig. 1. Life cycles of (a) the stick insect *T. californicum* and (b) the fly *D. melanogaster.* *represents a stage present only in females, as *Timema* males have one moult less than females. Hatchling and egg photographs of *Timema* were kindly provided by Bart Zijlstra (http://www.bartzijlstra.com), juvenile and adult stages by Jelisaveta Djordjevic. Life cycle of *D. melanogaster* modified from Weigmann *et al.* (2003).

Results

Sex-biased gene expression increases over development in *T. californicum*

Sex-biased gene expression gradually increased over the three developmental stages, with 0.2% (26) of the expressed genes sex-biased at the hatchling stage, 4.7% (568) at the juvenile, and 20.3% (2485) at the adult stage (Fig. 2). The hatchling stage had more female than male biased genes ($x^2_{(1)}$ =12.46, p<0.05), while the juvenile and adult stages had more male biased genes ($x^2_{\text{juvenile (1)}}$ =457.92, $x^2_{\text{adult (1)}}$ = 77.554, p<0.05) (Supplemental Table 4). The majority (85% and 91%) of sex biased genes at the two preadult stages had strong sex bias ($\geq$2 log$_2$FC, P$_{adj}$ <0.05), in contrast to the adult stage where sex bias was slight (57% of the sex-biased genes had <2 log$_2$FC, P$_{adj}$ <0.05) (Fig. 2). The intensity of sex-bias varied greatly over development with 1137 genes showing a significant sex by development interaction (P$_{adj}$<0.05).

To characterize the dynamics of sex-biased expression over development, we then classified the 2671 genes which are sex-biased at one or more developmental stages into 13 categories, dependent on their expression patterns over development (Supplemental Fig. 1). This classification showed that sex-biased genes are added gradually over development, with genes sex-biased at earlier stages generally

remaining sex-biased in the same direction at later stages (Supplemental Fig. 2). Only

a single gene shifted from male-biased to female-biased expression (Supplemental

Fig. 1). Out of 26 sex-biased genes at the hatchling stage, 18 (69%) were also

significantly sex-biased in at least one of the later stages, with 9 (35%) remaining sex-

biased throughout development (Fig. 3), while out of 568 sex-biased genes at the

juvenile stage, 390 (69%) stayed sex-biased at the adult stage (Fig. 3). This

classification is conservative, as it depends on the sex-biased gene expression

detection threshold, meaning sex-biased gene expression may be missed in some

stages inflating the number of differences we see. This is supported by the fact that

only around 65% of genes that are sex-biased in two stages show a significant sex by

development stage interaction (Supplemental table 8, also see Supplemental Fig. 1).

Sex-biased genes at the hatchling stage are particularly interesting because they

reveal sexual differentiation prior to visible morphological differences. We therefore

looked for functional annotations of the 26 genes sex-biased at the hatchling stage in

the *T. californicum* reference genome (Jaron *et al.* 2020). However, only 16 out of the

26 sex-biased genes had functional annotations and only one (*vitellogenin receptor*)

had a clear link to sexual differentiation (Fig. 3b). While key genes known to play a

role in insect sex-determination and differentiation pathways (*doublesex, transformer-*

2 and *sex-lethal*) were expressed at the hatchling stage, *doublesex* had very low

expression, preventing analysis of sex-bias, and *transformer-2* and *sex-lethal* did not

feature sex-biased expression.

Sex-biased genes are enriched for development-related processes

Genes sex-biased at the hatchling stage were enriched for GO-terms related to developmental processes (e.g., "regulation of cell development", "regulation of cell proliferation", "cuticle development"), and GO-terms related to female specific functions such as oogenesis ("oocyte construction", "oocyte development") (Supplemental Fig. 3a, and Supplemental Table 10). At the juvenile stage, sex-biased genes were enriched for GO-terms related to metabolic processes, in particular to catabolism (e.g., "lipid catabolic processes", "cellular catabolic processes", "regulation of catabolic processes") (Supplemental Fig. 3b, and Supplemental table 11). At the adult stage, sex-biased genes were enriched for GO-terms related to diverse metabolic and physiological processes with no clear association to sexual differentiation (Supplemental Fig. 3c, and Supplemental table 12); however, several terms were related to chemosensory and olfactory behavior, which may play a role in mate detection. Furthermore, sex-biased genes at the adult stage were enriched for pigmentation (e.g., "developmental pigmentation", "eye pigmentation"), which is a sexually dimorphic trait in *Timema californicum*.

Fig. 2 | Gene expression (log$_2$CPM) in *T. californicum* males and females at the hatchling (left), juvenile (middle) and adult stage (right). The number of differentially

expressed genes (SBG) is shown at the bottom right corner of each plot, as well as the total number of expressed genes at each stage (T). Genes are classified based on their sex-bias into seven categories: "slight FB"- female bias (<2 log$_2$FC), "strong FB"- female bias (≥2 log$_2$FC), "female limited"- with no expression in males, "slight MB"- male bias (<2 log$_2$FC), "strong MB"- male bias (≥2 log$_2$FC), "male limited"- no expression in females, "Not DE"- not differentially expressed genes.

Fig. 3 | Sex-biased genes shared across development. (**a**) Venn-diagram of sex-biased genes in three developmental stages: hatchling (yellow), juvenile (purple), and adult (green). The number of genes shared between all three stages was greater than expected by chance (observed overlap= 9, expected overlap= 0.21, Exact test of multi set interactions: p= 4.66x10^{-13}). (**b**) A heatmap showing the 26 sex-biased genes at the hatchling stage with their annotations and their expression in three developmental stages. Genes in red are female-biased, blue are male-biased. Saturation of the colors increases with the log$_2$FC. Genes without a label have no annotation, annotation in purple is for genes that were sex-biased in all three stages

Sex-biased genes have more stage-specific expression

Expression levels of sex-biased genes tend to be specific to only one developmental

stage while unbiased genes have a more constant expression level across

development (Fig. 4). Male-biased genes have the most stage specific expression in

both males and females. Female-biased genes are only slightly more stage specific

than un-biased genes, but the trend is consistent and the difference is significant

notably in both male and female adults.

Fig. 4 | Tau index of gene expression at three developmental stages in males (a) and

females (b). Tau range from zero (similarly expressed over development) to one (gene

expressed in only one stage). Three gene categories are depicted with different colors;

female-biased in red, male-biased in blue, and un-biased in grey. Boxplots represent

the median, lower and upper quartiles, and whiskers the minimum and maximum

values (in the limit of 1.5x interquartile range). Adjusted p values of Wilcoxon rank sum

tests are summarized above the box plots (*** = $p < 0.001$, * = $p < 0.05$, ˙ p=0.06).

Sex-biased genes have faster sequence evolution rates

Male-biased genes have faster rates of sequence evolution (dN/dS) compared to both un-biased and female-biased genes, at juvenile and adult stages (Fig. 5). At the hatchling stage, female-biased genes evolve significantly faster than un-biased genes, but not at the other two developmental stages. Elevated rates of sequence evolution are likely the result of reduced purifying selection, but stronger positive selection may also contribute to fast sequence evolution in some genes. To test this possibility, we repeated our analysis after excluding 19 (11 F-biased and 8 M-biased) genes for which a previous study (Jaron *et al.* 2020) detected signatures of positive selection. Excluding these 19 genes did not change our results (Supplemental Table 7a). More generally, we did not find an enrichment of positively selected genes among those with sex-biased expression (Fisher's exact test; p-value=0.39, odds ratio: 1.1) (Supplemental Table 7b).

We performed partial correlation analyses to test the effect of sex bias strength on sequence evolutionary rate, controlling for the average expression level of the gene (Supplemental Table 6). For both male-biased and female-biased genes, stronger sex-bias is associated with faster sequence evolution at the adult stage (Fig. 6). Partial correlations are not significant at juvenile or hatchling stages.

Fig. 5 | Divergence rates (the ratio of nonsynonymous to synonymous substitutions, d_N/d_S), at three developmental stages; hatchling, juvenile and adult stage. Boxplots represent the median, lower and upper quartiles, and whiskers the minimum and maximum values (in the limit of 1.5x interquartile range). Adjusted p values of Wilcoxon rank sum tests are summarized above the box plots (*** = $p < 0.001$, * = $p < 0.05$).

Fig. 6 | Spearman's rank correlations between the divergence rate of genes, d_N/d_S, and their strength of the sex bias $|\log_2\text{FC}|$ at three developmental stages. Note that all genes are displayed on each plot but only male- respectively female-biased ones are used for correlation tests.

Sex-biased gene expression increases abruptly in adult *D. melanogaster*

In order to compare the dynamics of sex-biased gene expression over development between hemimetabolous and holometabolous insects, we reanalyzed publicly available data from *D. melanogaster* (Ingleby *et al.* 2016)*,* using the same pipeline as for *T. californicum.* This experiment was chosen for comparison as it has a similar design, i.e. whole-body RNA-seq data from several developmental stages. In addition, although *D.* melanogaster from this experiment were lab-reared, they had a similar between sample variance as *T. californicum* (biological coefficient of variation at the adult stage: *D. melanogaster* = 0.274, *T. californicum* = 0.348).

In contrast to the gradual increase in the amount of sex-biased expression over

development observed in *T. californicum,* sex-biased gene expression increased

abruptly at the adult stage in *D. melanogaster* (Fig. 8), from 22% to 84% of sex-biased

genes (Fig. 7). Overall, at each developmental stage, there were more male than

female-biased genes (Supplemental table 5). The majority (85% and 63%) of sex-

biased genes at the two preadult stages had strong sex bias ($\geq$2 log$_2$FC, P_{adj}<0.05) in

contrast to the adult stage where weaker sex-bias was slightly more common (50.5%

of the sex-biased genes had <2 log$_2$FC, P_{adj} <0.05) (Supplemental table 5). Out of

2113 sex-biased genes at the larval stage, 1262 (60%) were significantly sex-biased

in at least one of the later stages, with 1226 (58%) remaining sex-biased throughout

development (Fig. 7b), while out of 2643 sex-biased genes at the pupal stage, 2423

(92%) stayed sex-biased at the adult stage (Fig. 7b).

Fig. 7 | (a) Gene expression (log$_2$CPM) in males and females at the larval (left), pupal

(middle) and adult stage (right). The number of differentially expressed genes (SBG)

is shown at bottom right corner of each plot, as well as the total number of expressed

genes at each stage (T). Genes are classified based on their sex-bias into seven categories: "slight FB"- female bias (<2 log$_2$FC), "strong FB"- female bias (≥ 2 log$_2$FC), "female limited"- with no expression in males, "slight MB"- male bias (<2 log$_2$FC), "strong MB"- male bias (≥ 2 log$_2$FC), "male limited"- no expression in females, "Not DE"- not differentially expressed genes. **(b)** Venn-diagram of sex-biased genes in three developmental stages: larval (yellow), pupal (purple), and adult (green). The number of genes shared between all three stages was greater than expected by chance (observed overlap= 1226, expected overlap= 334.7), Exact test of multi set interactions: p-value≈ 0).

Fig. 8 | Percentage of sex-biased genes (SBG) in *T. californicum* (left panel) and *D. melanogaster* (right panel) at three developmental stages. Note that the immature developmental stages in the two species are not homologous but serve to illustrate the gradual *versus* abrupt increase in sex-biased gene expression in adults.

Discussion

In the stick insect *T. californicum,* gene expression differs between males and females before morphological sexual dimorphism. Over development, most of the sex-biased genes are added gradually and cumulatively, meaning that once a gene gains sex-biased expression in *T. californicum* it generally remains sex-biased at later developmental stages. In addition, genes were consistent in their sex-bias direction; only a single gene shifted from male- to female-biased expression. Surprisingly, such a gradual and consistent addition of sex-bias does not appear to be a general pattern over animal development. For example, in humans and mosquitos, some stages have bursts of sex-biased gene expression, suggesting that specific developmental stages contribute discretely to sexual differentiation, and that sexual phenotypes are not gradually established (Magnusson *et al.* 2011; Shi *et al.* 2016).

Although only 26 genes were sex-biased in *T. californicum* at the hatchling stage, they all had a strong sex-bias and nine of them stayed significantly sex-biased at both the juvenile and adult stages. As such these 26 genes likely represent key genes involved in the sexual differentiation of *Timema*. Most of the sex-biased genes at the hatchling stage were female-biased. This might suggest that the process of building a female phenotype begins earlier in development. Later developmental stages had more male-biased than female- biased genes. A similar shift from female- to male-bias in gene expression during development was found in *Nasonia* jewel wasps, where little male-biased expression was detected until the pupal stage, during the activation of spermatogenesis. Similar to *Timema*, the shift was interpreted as a different developmental timing of the two sexes (Rago *et al.* 2020). In *Timema*, male-biased genes were also expressed in a relatively stage-specific manner, in contrast to

a more constant expression of female-biased genes. This further supports the idea

that *Timema* individuals first develop along the female trajectory, i.e. that the female

phenotype is the "default" in *Timema*, and may indicate greater pleiotropy of female-

biased genes. Additional support for this interpretation comes from the lower

sequence divergence rates of female-biased genes than male-biased genes,

implicating stronger selective constraints on female- than male- biased genes. In

addition, we observed that genes with greater sex-bias have faster sequence

divergence rates. This is largely due to relaxed purifying selection, with positive

selection contributing little to the accelerated evolutionary rate of sex-biased genes.

These findings are consistent with the idea that sex-biased genes evolve from genes

with few evolutionary constraints and that are relatively "dispensable" (Mank and

Ellegren 2009; Catalan *et al.* 2018).

Sex-biased genes in *Timema* hatchlings were enriched for processes related

to oogenesis. Moreover, one of the most consistently and strongly female-biased

genes across development was annotated as *vitellogenin receptor*. In insects,

vitellogenin receptor transfers yolk protein precursors into the oocytes and is

necessary for oocyte and early embryo development (Raikhel and Dhadialla 1992;

Cho and Raikhel 2001). The role of these processes in *Timema* hatchlings remain to

be investigated but *vitellogenin receptor* expression may indicate a very early onset of

oocyte development in *Timema*, consistent with studies in other stick insects (Taddei

et al. 1992). Sex-biased genes in juvenile and adult stages were not enriched for

processes obviously related to sexual traits. However, some processes such as

pigmentation and chemosensory/olfactory behavior could have a sex-related role.

Pigmentation is a sexually dimorphic trait in *Timema* (Sandoval 2008), while

chemosensory and olfactory behavior are important for mate recognition and mating

(Nosil *et al.* 2007; Schwander *et al.* 2013). We also checked the expression of two key

genes involved in insect sex-determination pathways: *transformer* and *doublesex*. In

holometabolous insects, *transformer* acts as a splicing regulator of *doublesex*, a sex

differentiation master switch gene (Geuverink and Beukeboom 2014). The *Doublesex-*

Transformer role in sex-differentiation is conserved among Diptera, Coleoptera and

Hymenoptera (Verhulst *et al.* 2010; Wexler *et al.* 2019), and likely play the same role

in sex differentiation in hemimetabolous insects (Zhuo *et al.* 2018; Wexler *et al.* 2019).

In *T. californicum,* we found that *doublesex* was expressed in all three developmental

stages, but its expression was too low to examine sex-bias. *Transformer* is not

annotated in the available *T. californicum* genome; a *Transformer-2 homolog* is

annotated but did not feature sex-bias at any of the three developmental stages. As

such, whether or not these classic sex-determining genes influence sexual

differentiation in *Timema* is unclear, and requires future studies.

We hypothesized that the dynamics of sex biased gene expression would be

strongly affected by hemimetabolous *vs* holometabolous development in insects, with

a gradual increase of sex-biased gene expression over development in the former and

an abrupt increase at the adult stage in the latter. Hemimetabolous insects such as *T.*

californicum have multiple nymphal stages that progressively resemble the adult

stage, together with a gradual increase in sexual dimorphism. On the other hand, a

holometabolous insect like *D. melanogaster* has more monomorphic larval stages and

pupae, with no resemblance to the sexually dimorphic adult stage. The change in

sexually dimorphic gene expression across development indeed mirrored these

different developmental patterns, with the proportion of sex-biased genes increasing

gradually in *T. californicum* as sexual dimorphism became more pronounced, whereas

sexually dimorphic gene expression increased abruptly at the adult stage in *D.*

melanogaster.

Although the change in sexually dimorphic gene expression across

development fits with our expectations, we were surprised to find a much higher

proportion of sex-biased genes in *D. melanogaster* than in *T. californicum* overall. This

difference is clearest at the adult stage when most of the expressed genes in *D.*

melanogaster were sex-biased (84%), compared to only 20% in *T. californicum*.

Similar to *Drosophila*, other holometabolous insects also featured sex-bias for more

than half of the genes expressed at the adult stage (Baker *et al.* 2011; Rago *et al.*

2020), while all studied hemimetabolous insects had a much smaller proportion of sex-

biased genes (<10%) (Pal and Vicoso 2015). Why there is such a difference in the

prevalence of sex-biased genes between species with different developmental

strategies is still an open question. One potential explanation is that the proportion of

sex-biased genes at the adult stage may be less constrained under holo- than

hemimetabolous development. For example, the complete reorganization of cells and

tissues during metamorphosis in holometabolous insects may allow for more drastic

phenotypic changes between the sexes, while incomplete metamorphosis constrains

the evolution of distinct sexual phenotypes. Data on additional species are required to

confirm that this difference is a general phenomenon and not simply due

idiosyncrasies of the few species yet investigated. Holometabolous insects studied are

often established lab-models, with high fecundity, and may thus be characterized by

comparatively large gonads, which would result in large proportions of sex-biased

genes in adults. By contrast, all hemimetabolous insects studied for sex-biased

expression are non-model species. However, the choice of model species is unlikely

to be the sole explanation for the different amount of sex bias in holo- and

hemimetabolous species. Indeed, in cricket, *Gryllus bimaculatus*, and *Timema* stick insects, even the most sexually differentiated tissue (gonads) feature sex-bias of fewer than 30% of genes (Parker *et al.* 2019; Whittle *et al.* 2020), suggesting that extensive differences are not completely driven by large gonads.

Overall, our results describe the dynamics of sex-biased gene expression over development in a hemimetabolous insect, and suggest profound differences in these dynamics between holometabolous and hemimetabolous species. Generating sexually dimorphic phenotypes is a developmental process, and we show that dynamics of sex-biased gene expression mirror this development with sex-biased genes being added gradually over development, and with the majority of genes sex-biased at early stages remaining sex-biased in later stages.

Material and methods

Sample collection and preservation

Timema californicum eggs hatch in early winter. Upon hatching, insects moult several times before reaching maturity in late spring (Sandoval 1994). Developmental stages (especially adults and hatchlings) do not co-occur temporally, hence samples from different stages cannot be collected simultaneously. Adults (4 males, 4 females) and juveniles (4 males, 3 females) were collected in California (Saratoga County) in 2015 and 2016, respectively. They were fed with artificial medium for two days to prevent contamination with plant cells from the gut, and subsequently frozen at -80°C. Hatchlings were obtained from the eggs laid by field-collected adults in 2015. Hatchlings were flash frozen and stored at -80 °C. Sexes cannot be distinguished morphologically at the hatchling stage, thus single individuals were extracted to allow

389 for later identification of sex via genotyping (see below) and 5 male and 4 female

390 hatchlings were used.

RNA extraction and sequencing

Individuals (whole-bodies) from all developmental stages were mechanically

homogenized with beads (Sigmund Linder) in liquid nitrogen. We then added 900 ul

Trizol (Life Technologies), followed by 180 ul chloroform and 350 ul ethanol. The

aqueous layer was then transferred to RNeasy MinElute Columns (Qiagen). Upon

RNA extraction, samples were treated with DNase Turbo Kit (Life Tech) following the

manufacturer's protocol. Total RNA from the hatchlings was extracted with

MagMaxTM Express Robot (AB Applied Biosystems) using a MagMAXTM-96 Total

RNA Isolation Kit from Ambion (Life Technologies) following the manufacturer's

protocol, but without DNAse to be able to determine the sex of hatchlings via

genotyping. The quality and quantity of the extracted RNA was then measured using

a NanoDrop and Bioanalyzer. Library preparations (one for each individual) were done

using the Illumina TruSeq Stranded Total RNA kit, upon which samples were

sequenced together in six lanes. Paired-end sequencing with a read length of 100 bp

was done on a HiSeq2000 platform at the GTF (Genomic Technologies Facility,

Centre of Integrative Genomics, Lausanne, Switzerland).

Hatchling sex identification

To identify the sex of *T. californicum* hatchlings, we developed microsatellite markers

on the *T. californicum* X chromosome scaffolds reported by Parker et al. (in prep).

Timema have an XX/X0 sex determination system (Schwander and Crespi 2009)

meaning X-linked regions will be present in two copies in females and one copy in

males. Consequently, females can feature heterozygous genotypes for polymorphic

X-linked markers, while males are invariably hemizygous. Candidate microsatellite

markers were designed with msatcommander (v. 1.08, default options) (Faircloth

2008). A selection of four candidate markers were tested for polymorphism and sex-

linkage using 15 adult males and 15 adult females in two Multiplex PCR Kit reactions

(Qiagen; see Supplemental Table 1 for primer sequences and Supplemental Table 2

for PCR conditions). Microsatellite alleles were then determined with an ABI3100

machine (Applied Biosystem) and Genemapper v.4.1 (Applied Biosystems 2009).

Across microsatellite markers, all 15 tested females were heterozygous at minimum

two markers, whereas males were invariably hemizygous at all markers, confirming

the predictions for *Timema* X-linked markers (Supplemental table 2). We thus

genotyped the nine hatchlings at the four microsatellite markers to identify their sex,

and were able to identify four females and five males that were used for the

transcriptome study (Supplemental table 2).

Raw data quality control, mapping and read counting

The quality of the reads was checked using FastQC v.0.11.2 (Andrews 2014). We

used Cutadapt v. 2.3 with Python v. 3.5.2 (Martin 2011) to remove adapter sequences.

Low quality bases at both ends of the reads were trimmed when below a Phred score

of 10. Bases with an average Phred score below 20 in a 4bp sliding window were

trimmed. Finally, reads with a length below 80 bp were removed Trimmomatic v. 0.36

(Bolger *et al.* 2014). Trimmed reads were then mapped to the reference *T. californicum*

genome from Jaron *et al.* (2020) using STAR v. 2.6.0c (Dobin *et al.* 2013). HTSeq

v.0.9.1 (Anders *et al.* 2014) was used to count the number of reads uniquely mapped

to each gene, with the following options (htseq-count -f bam -r name -s reverse -t gene

-i ID -m union --nonunique none).

Drosophila melanogaster data

D. melanogaster RNA-seq whole body data from Ingleby *et al.* (2016) was downloaded from SRA (accession number SRP068235). To mirror the data available for *Timema* and to facilitate comparisons, four replicates per sex and three developmental stages from different hemiclonal lines were used. A complete list of samples is provided in the supplementary materials (Supplemental Table 3). Although reads were deposited as paired-end, we found that most of them were unpaired, thus we used only the forward reads for analysis. Reads were trimmed, mapped to the reference genome (FlyBase, r6.23) (Gramates *et al.* 2017) and counted using the methods described above.

Differential gene expression analysis

Differential expression between the sexes was performed using edgeR v.3.16.5 (Robinson *et al.* 2009; McCarthy *et al.* 2012) in RStudio v.1.2.5019 (Team 2019). This analysis fits a generalized linear model following a negative binomial distribution to the expression data and calculates a p-value associated with the hypothesis that gene counts differ between experimental groups. EdgeR implements a normalization method (trimmed mean of M values, TMM) to account for composition bias. We analyzed data separately for each developmental stage. We filtered out genes with low counts; we required a gene to be expressed in a majority of male or female libraries dependent on the number of replicates per sex (i.e. a minimum three libraries when replicate number is >= 4, two libraries when replicate number is = 3), with expression level > 0.5 CPM (counts per million). To test for differential gene expression between the sexes, we used a generalized linear model with quasi-likelihood F- test, with

contrasts for each stage between females and males (Chen *et al.* 2016). To correct

for multiple testing, we applied the Benjamini-Hochberg method (Benjamini and

Hochberg 1995) with an alpha of 5%. Genes with significantly higher expression in

males were considered as male-biased, genes with significantly higher expression in

females as female-biased, and the genes without significant differential expression

between sexes as un-biased. To identify genes showing a significant sex by

developmental stage interaction we used a similar GLM modelling approach but with

all stages included. We used SuperExactTest v.0.99.4 (Wang *et al.* 2015) to test if the

overlap of sex-biased genes between the three developmental stages was greater

than what is expected by chance. To visualize the overlap of sex-biased genes

between developmental stages we used VennDiagram v.1.6.20 (Chen and Boutros

2011). We visualized the log$_2$FC sex-biased gene expression across developmental

stages, using pheatmap v.1.0.12 (Kolde 2018)

Stage specific gene expression

Median gene expression (CPM) values were calculated for each developmental stage

and sex. These values were used to calculate Tau, an index of specificity of gene

expression over development (Yanai *et al.* 2004; Liu and Robinson-Rechavi 2018)

Values of Tau range from zero (broadly expressed over development) to one (gene

expressed in only one stage). To visualize the results we made boxplots using ggplot2

v. 3.3.2 (Wickham 2016).

Divergence rates

To calculate sequence divergence rates, we used the 7098 one-to-one orthologs

between *T. californicum* and *T. poppensis* from Jaron *et al.* (2020). Coding sequences

were codon-aligned using prank v.170427 (Loytynoja 2014). For each of the orthologs,

the number of nonsynonymous substitutions per sites (d_N), the number of synonymous

substitutions per site (d_S) and their ratio (d_N/d_S) was calculated using PAML v4.9 (Yang

2007). To test for differences in sequence divergence rates (d_N/d_S) between different

gene categories (female-biased, male-biased and un-biased), we used Wilcoxon tests

(ggpubr v.0.2.5, in R) with p-values adjusted for multiple comparisons using the

Benjamini-Hochberg method (Haynes 2013). Note that some of the *T. californicum*

sex-biased genes had no identified ortholog in *T. poppensis*, thus we do not have

sequence evolution rates for all sex-biased genes. Sequence divergence rates were

obtained for 14 out of 26 SBG at the hatchling stage, 323/568 at the juvenile stage,

and 1286/2485 at the adult stage. To account for the influence of positive selection we

repeated our analyses after removing genes with evidence for positive selection in the

T. californicum lineage (identified in Jaron *et al.* (2020)).

Furthermore, we used a partial correlation between the strength of the sex bias

($\log_2$FC) and the sequence divergence rate (d_N/d_S) on sex-biased genes only, while

controlling for the average expression level (CPM) of the gene in all libraries (ppcor

v.01 (Kim 2015), in R) in order to determine if genes with higher $\log_2$FC have faster

sequence divergence rates.

Functional analysis of sex-biased genes

We performed Gene Ontology (GO) enrichment analysis using TopGO v.2.26.0 on

sex-biased genes obtained with EdgeR (see above) at each developmental stage. We

used a functional annotation derived from the blasting sequences to *Drosophila*

melanogaster database. Only GO terms with minimum ten annotated genes were

used. Enrichment of terms was determined using a weighted Kolmogorov-Smirnov-

like statistical test, equivalent to the gene set enrichment analysis (GSEA method).

We applied the "elim" algorithm, which considers the Gene Ontology hierarchy, i.e. it

first assesses the most specific GO terms, and then more general ones (Alexa *et al.*

2006). Our analysis focused on gene sets in the Biological Processes (BP), GO

category. We considered terms as significant when $p < 0.05$. Enriched GO terms were

then semantically clustered using ReviGO (Supek *et al.* 2011) to aid interpretation.

Data availability

519 *Timema californicum* raw reads have been deposited in SRA under PRJNA678950

520 bioproject, with accession codes SRR13084978-SRR13085001 (Supplemental table

521 9). *Timema californicum* mapped read counts are provided in Supplemental table 13

522 and fold changes in Supplemental table 14. *Drosophila melanogaster* mapped read

523 counts are provided in Supplemental table 15.

Code availability

526 Scripts with code for the analysis in this study are available at

527 https://github.com/JelisavetaDjordjevic/Sex_biased_gene_expression

Acknowledgment

530 This study was supported by Swiss National Science Foundation Grant

531 31003A_182495.

Bibliography

Alexa A, Rahnenführer J, Lengauer T. 2006. Improved scoring of functional groups from gene expression data by decorrelating GO graph structure. Bioinformatics 22:1600-1607.

Anders S, Pyl PT, Huber W. 2014. HTSeq—a Python framework to work with high-throughput sequencing data. Bioinformatics 31:166-169.

FastQC. v.0.11.2. [Internet]. 2014. Available from: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Baker DA, Nolan T, Fischer B, Pinder A, Crisanti Aa, Russell S. 2011. A comprehensive gene expression atlas of sex- and tissue- specificity in the malaria vector, *Anopheles gambiae*. BMC genomics 12:12.

Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. Journal of the Royal statistical society: series B (Methodological) 57:289-300.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30:2114-2120.

Catalan A, Macias-Munoz A, Briscoe AD. 2018. Evolution of sex-biased gene expression and dosage compensation in the eye and brain of *Heliconius* butterflies. Mol Biol Evol 35:2120-2134.

Chauhan P, Wellenreuther M, Hansson B. 2016. Transcriptome profiling in the damselfly *Ischnura elegans* identifies genes with sex-biased expression. BMC genomics 17:985.

Chen H, Boutros PC. 2011. VennDiagram: a package for the generation of highly-customizable Venn and Euler diagrams in R. BMC Bioinformatics 12:35.

Chen S, Yang P, Jiang F, Wei Y, Ma Z, Kang L. 2010. De novo analysis of transcriptome dynamics in the migratory locust during the development of phase traits. PLOS ONE 5:e15633.

Chen Y, Lun AT, Smyth GK. 2016. From reads to genes to pathways: differential expression analysis of RNA-Seq experiments using Rsubread and the edgeR quasi-likelihood pipeline. F1000Res 5:1438.

Cho K-H, Raikhel AS. 2001. Organization and developmental expression of the mosquito vitellogenin receptor gene. Insect Molecular Biology 10:465-474.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29:15-21.

Ellegren H, Parsch J. 2007. The evolution of sex-biased genes and sex-biased gene expression. *Nat Rev Genet* 8:689-698.

Faircloth BC. 2008. Msatcommander: detection of microsatellite repeat arrays and automated, locus-specific primer design. Mol Ecol Resour 8:92-94.

Geuverink E, Beukeboom LW. 2014. Phylogenetic distribution and evolutionary dynamics of the sex determination genes *doublesex* and *transformer* in Insects. Sexual Development 8:38-49.

Gramates LS, Marygold SJ, Santos GD, Urbano JM, Antonazzo G, Matthews BB, Rey AJ, Tabone CJ, Crosby MA, Emmert DB, et al. 2017. FlyBase at 25: looking to the future. Nucleic Acids Res 45:D663-D671.

Grath S, Parsch J. 2012. Rate of amino acid substitution is influenced by the degree and conservation of male-biased transcription over 50 myr of *Drosophila* evolution. Genome Biol Evol 4:346-359.

Haynes W. 2013. "Benjamini–Hochberg method". In: Dubitzky W, Wolkenhauer O, Cho K-H, Yokota H, editors. *Encyclopedia of Systems Biology*. New York, NY: Springer New York. p. 78-78.

Ingleby FC, Ilona F, Morrow EH. 2015. Sex-biased gene expression and sexual conflict throughout development. Cold Spring Harbor Laboratory Press 7:a017632.

Ingleby FC, Webster CL, Pennell TM, Flis I, Morrow EH. 2016. Sex-biased gene expression in *Drosophila melanogaster* is constrained by ontogeny and genetic architecture. bioRxiv:034728.

Jaron KS, Parker DJ, Anselmetti Y, Van PT, Bast J, Dumas Z, Figuet E, François CM, Hayward K, Rossier V, et al. 2020. Convergent consequences of parthenogenesis on stick insect genomes. bioRxiv 2020.11.20.391540

Khila A, Abouheif E, Rowe L. 2012. Function, developmental genetics, and fitness consequences of a sexually antagonistic trait. Science 336:585-589.

Kim S. 2015. ppcor: an R package for a fast calculation to semi-partial correlation coefficients. Communications for statistical applications and methods 22:665-674.

Kolde R. 2018. pheatmap: Pretty Heatmaps, R package version 1.0.12.

Lande R. 1980. Sexual dimorphism, sexual selection, and adaptation in polygenic characters. Evolution 34:292-305.

Liu J, Robinson-Rechavi M. 2018. Developmental constraints on genome evolution in four bilaterian model species. Genome Biol Evol 10:2266-2277.

Lowe R, Gemma C, Rakyan VK, Holland ML. 2015. Sexually dimorphic gene expression emerges with embryonic genome activation and is dynamic throughout development. BMC genomics 16:295.

Loytynoja A. 2014. Phylogeny-aware alignment with PRANK. Methods Mol Biol 1079:155-170.

Magnusson K, Mendes AM, Windbichler N, Papathanos PA, Nolan T, Dottorini T, Rizzi E, Christophides GK, Crisanti A. 2011. Transcription regulation of sex-biased genes during ontogeny in the malaria vector *Anopheles gambiae*. PLOS ONE 6:e21572.

Mank JE. 2017. The transcriptional architecture of phenotypic dimorphism. Nat Ecol Evol 1:6.

Mank JE, Ellegren H. 2009. Are sex-biased genes more dispensable? Biol Lett 5:409-412.

Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet.journal 17:10-12.

McCarthy DJ, Chen Y, Smyth GK. 2012. Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation. Nucleic Acids Res 40:4288-4297.

Nosil P, Crespi BJ, Gries R, Gries G. 2007. Natural selection and divergence in mate preference during speciation. Genetica 129:309.

Pal A, Vicoso B. 2015. The X Chromosome of Hemipteran Insects: Conservation, Dosage Compensation and Sex-Biased Expression. Genome Biol Evol 7:3259-3268.

Paris M, Villalta JE, Eisen MB, Lott SE. 2015. Sex bias and maternal contribution to gene expression divergence in *Drosophila* blastoderm Embryos. PLoS Genet 11:e1005592.

Parker DJ, Bast J, Jalvingh K, Dumas Z, Robinson-Rechavi M, Schwander T. 2019. Sex-biased gene expression is repeatedly masculinized in asexual females. Nat Commun 10:4638.

Perry JC, Harrison PW, Mank JE. 2014. The ontogeny and evolution of sex-biased gene expression in *Drosophila melanogaster*. Mol Biol Evol 31:1206-1219.

Pointer MA, Harrison PW, Wright AE, Mank JE. 2013. Masculinization of gene
expression is associated with exaggeration of male sexual dimorphism. PLoS Genet
9:e1003697.
Rago A, Werren JH, Colbourne JK. 2020. Sex biased expression and co-expression
networks in development, using the hymenopteran *Nasonia vitripennis*. PLoS Genet
16:e1008518.
Raikhel A, Dhadialla T. 1992. Accumulation of Yolk Proteins in Insect Oocytes. Annual
Review of Entomology 37:217-251.
Robinson MD, McCarthy DJ, Smyth GK. 2009. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics 26:139-
140.
Sandoval CP. 2008. Differential visual predation on morphs of *Timema cristinae*
(Phasmatodeae:Timemidae) and its consequences for host range. Biological Journal
of the Linnean Society 52:341-356.
Sandoval CP. 1994. Geographic, ecological and behavioral factors affecting spatial
variation in color morph frequency in the walking-stick, *Timema cristinae*.
Schwander T, Crespi BJ. 2009. Multiple direct transitions from sexual reproduction to
apomictic parthenogenesis in *Timema* stick insects. Evolution 63:84-103.
Schwander T, Crespi JB, Gries R, Gries G. 2013. Neutral and selection-driven decay
of sexual traits in asexual stick insects. Proceedings of the Royal Society B: Biological
Sciences 280:20130823.
Shi L, Zhang Z, Su B. 2016. Sex biased gene expression profiling of human brains at
major developmental stages. Sci Rep 6:21181.
Supek F, Bosnjak M, Skunca N, Smuc T. 2011. REVIGO summarizes and visualizes
long lists of gene ontology terms. PLOS ONE 6:e21800.
Taddei C, Chicca M, Maurizii MG, Scali V. 1992. The germarium of panoistic ovarioles
of *Bacillus rossius* (Insecta Phasmatodea): Larval differentiation. Invertebrate
Reproduction & Development 21:47-56.
Team R. 2019. RStudio: Integrated Development Environment for R.
Verhulst EC, van de Zande L, Beukeboom LW. 2010. Insect sex determination: it all
evolves around *transformer*. Curr Opin Genet Dev 20:376-383.
Vickery VR, Sandoval CP. 2001. Descriptions of three new species of *Timema*
(Phasmatoptera: Timematodea: Timematidae) and notes on three other species.
Journal of Orthoptera Research 10:53-61.
Wang M, Zhao Y, Zhang B. 2015. Efficient test and visualization of multi-set
intersections. Sci Rep 5:16923.
Weigmann K, Klapper R, Strasser T, Rickert C, Technau G, Jäckle H, Janning W,
Klämbt C. 2003. FlyMove- a new way to look at development of *Drosophila*. Trends in
Genetics 19:310-311.
Wexler J, Delaney EK, Belles X, Schal C, Wada-Katsumata A, Amicucci MJ, Kopp A.
2019. Hemimetabolous insects elucidate the origin of sexual development via
alternative splicing. eLife 8:e47490.
Whittle CA, Kulkarni A, Extavour CG. 2020. Sex-biased genes expressed in the cricket
brain evolve rapidly. bioRxiv 2020.07.07.192039.
Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis: Springer-Verlag New
York.
Yanai I, Benjamin H, Shmoish M, Chalifa-Caspi V, Shklar M, Ophir R, Bar-Even A,
Horn-Saban S, Safran M, Domany E, et al. 2004. Genome-wide midrange
transcription profiles reveal expression level relationships in human tissue
specification. Bioinformatics 21:650-659.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. Mol Biol Evol
24:1586-1591.
Zhuo JC, Hu QL, Zhang HH, Zhang MQ, Jo SB, Zhang CX. 2018. Identification and
functional analysis of the doublesex gene in the sexual development of a
hemimetabolous insect, the brown planthopper. Insect Biochem Mol Biol 102:31-42.
685