Dynamics of sex-biased gene expression over development

in the stick insect *Timema californicum*

Jelisaveta Djordjevic[1], Zoé Dumas[1], Marc Robinson-Rechavi[1,2],

Tanja Schwander[1,a], Darren James Parker[1,2,a]

[1] Department of Ecology and Evolution, University of Lausanne, Lausanne, Switzerland

[2] Swiss Institute of Bioinformatics, Lausanne, Switzerland

[a] equal contribution

Correspondence should be addressed to jelisaveta.djordjevic@unil.ch or

DarrenJames.Parker@unil.ch

Abstract

Sexually dimorphic phenotypes are thought to arise primarily from sex-biased gene

expression during development. Major changes in developmental strategies, such as the shift

from hemimetabolous to holometabolous development, are therefore expected to have

profound consequences for the dynamics of sex-biased gene expression. However, no studies

have previously examined sex-biased gene expression over development in hemimetabolous

insects, precluding comparisons between developmental strategies. Here we characterized

sex-biased gene expression at three developmental stages in a hemimetabolous stick insect

(*Timema californicum*): hatchlings, juveniles, and adults. As expected, the proportion of sex-

biased genes gradually increased over development. Sex-biased genes identified at early

developmental stages were generally consistently male- or female- biased at later stages,

suggesting their importance in sexual differentiation. We then compared the dynamics of sex-

biased gene expression over development in *T. californicum* to those of the holometabolous

fly *Drosophila melanogaster* by reanalyzing publicly available RNA-seq data from third instar

larval, pupal and adult stages*.* In *D. melanogaster,* sex-biased gene expression increases

abruptly at the adult stage when morphological sexual dimorphism is manifested. This

supports the prediction that sex-biased gene expression mirrors phenotypic sexual

dimorphism. Our study details for the first time the dynamics of sex-biased gene expression

over development in a hemimetabolous insect and suggests that these dynamics differ

extensively between holometabolous and hemimetabolous species.

Introduction

Males and females often have divergent evolutionary interests, resulting in sex-specific selection pressures and ultimately the evolution of sexually dimorphic phenotypes (Khila et al., 2012; Lande, 1980). Studies investigating how sexually dimorphic phenotypes are generated have shown the importance of differential gene expression between the sexes, suggesting that sex-specific selection is the major driving force behind the evolution of sex-biased gene expression (Chauhan et al., 2016; Mank, 2017). The relationship between sexual dimorphism and sex-biased gene expression has been largely studied at the adult stage when sexual dimorphism is completely manifested and reproductive interests between males and females are most different (Chauhan et al., 2016; Pointer et al., 2013). However, sex-biased gene expression has also been found at early developmental stages of many species, well before any phenotypic sexual dimorphism becomes apparent (Lowe et al., 2015; Paris et al., 2015). This suggests that expression patterns in early developmental stages are also under sex-specific selection pressures (Ingleby et al., 2015).

Because we lack detailed studies of sex-biased gene expression over development, we do not know how consistent or dynamic the expression of the sex-biased genes is, nor how these dynamics relate to changes in phenotypic sexual dimorphism. For example, although we expect to see an overall increase in sex-biased gene expression over development (Ingleby et al., 2015; Mank, 2017), it is not clear if this results from the gradual increase of sex-bias of a set of genes or if sex-biased genes at early stages are largely different to those at later stages. Furthermore, sex-biased gene expression can be considered to reflect a broad measure of sexual dimorphism, including physiological and behavioral traits, and may therefore be more representative than dimorphism quantified using external morphology.

However, whether the extent of sexual dimorphism and sex-biased gene expression are

generally correlated remains poorly known.

In insects, patterns of sex-biased gene expression over development could differ

depending on developmental strategy. Holometabolous insects have morphologically and

ecologically distinct larval, pupal, and adult stages, and phenotypic sexual dimorphism is

commonly prominent only at the adult stage. Contrastingly, hemimetabolous insects go

through gradual morphological changes and sexual differentiation, and the nymphal stages

morphologically resemble adults (Chen et al., 2010). These distinct dynamics for the

morphological differentiation of the sexes are expected to be mirrored by sex-biased gene

expression, with abrupt versus gradual increases in the proportion of sex-biased genes over

development. However, studies of sex-biased gene expression during development have thus

far only been conducted in holometabolous insects, with the main focus on *Drosophila*

(Ingleby et al., 2015; Magnusson et al., 2011; Perry et al., 2014; Rago et al., 2020), precluding

comparisons between species with different developmental strategies.

Here, we studied the dynamics of sex-biased gene expression across development in

a hemimetabolous insect, *Timema californicum.* These are sexually dimorphic walking-stick

insects, found in California (Vickery and Sandoval, 2001). Males are smaller than females, with

different coloration, and have conspicuous cerci used for holding on to the female during

copulation. We selected three developmental stages (Figure 1a): the hatchling stage where

sexes are phenotypically identical, the third juvenile stage with minor phenotypic differences,

and the adult stage, with pronounced sexual dimorphism. We investigated whether sex-

biased genes are recruited in a stage-specific manner, and if the magnitude of sex-bias

increases with development. We further explored sequence evolution rates of sex-biased

genes in pre-adult stages, to determine if they show elevated rates as commonly observed

for sex-biased genes at adult stages in many species (Ellegren and Parsch, 2007; Grath and

Parsch, 2012; Perry et al., 2014).

We then compared the dynamics of sex-biased gene expression over development in

the hemimetabolous stick insect to those of the holometabolous fly *Drosophila melanogaster*

by reanalyzing publicly available RNA-seq data from 3$^{rd.}$ instar larvae, pupal and adult stages,

using the same pipeline as for *T. californicum.* In *D. melanogaster*, both larval and pupal stages

have little sexual dimorphism, while the adult stage has extensive sexual dimorphism (Figure

1b). We hypothesized that sex biased gene expression follows the establishment of

morphological sexual dimorphism, with a gradual increase in sex-biased gene expression over

development in *T. californicum*, and an abrupt increase at the adult stage in *D. melanogaster*.

Figure 1. Life cycles of (a) the stick insect *T. californicum* and (b) the fly *D. melanogaster.*

*represents a stage present only in females, as *Timema* males have one moult less than

females. Hatchling and egg photographs of *Timema* were kindly provided by Bart Zijlstra

(http://www.bartzijlstra.com), juvenile and adult stages by Jelisaveta Djordjevic. Life cycle of

D. melanogaster modified from (Weigmann et al., 2003)

Material and methods

Sample collection and preservation

Timema californicum eggs hatch in early winter. Upon hatching, insects moult several times

before reaching maturity in late spring (Sandoval, 1994). Developmental stages (especially

adults and hatchlings) do not co-occur temporally, hence samples from different stages

cannot be collected simultaneously. Adults (4 males, 4 females) and juveniles (4 males, 3

females) were collected in California (Saratoga County) in 2015 and 2016, respectively. They

were fed with artificial medium for two days to prevent contamination with plant cells from

the gut, and subsequently frozen at -80°C. Hatchlings were obtained from the eggs laid by

field-collected adults in 2015. Hatchlings were flash frozen and stored at -80 °C. Sexes cannot

be distinguished morphologically at the hatchling stage, thus single individuals were extracted

to allow for later identification of sex via genotyping (see below) and 5 male and 4 female

hatchlings were used.

RNA extraction and sequencing

Individuals (whole-bodies) from all developmental stages were mechanically homogenized

with beads (Sigmund Linder) in liquid nitrogen. We then added 900 ul Trizol (Life

Technologies), followed by 180 ul chloroform and 350 ul ethanol. The aqueous layer was then

transferred to RNeasy MinElute Columns (Qiagen). Upon RNA extraction, samples were

treated with DNase Turbo Kit (Life Tech) following the manufacturer's protocol. Total RNA

from the hatchlings was extracted with MagMax™ Express Robot (AB Applied Biosystems)

using a MagMAX™-96 Total RNA Isolation Kit from Ambion (Life Technologies) following the

manufacturer's protocol, but without DNAse to be able to determine the sex of hatchlings via

genotyping. The quality and quantity of the extracted RNA was then measured using a

NanoDrop and Bioanalyzer. Library preparations (one for each individual) were done using

the Illumina TruSeq Stranded Total RNA kit, upon which samples were sequenced together in

six lanes. Paired-end sequencing with a read length of 100 bp was done on a HiSeq2000

platform at the GTF (Genomic Technologies Facility, Centre of Integrative Genomics,

Lausanne, Switzerland).

Hatchling sex identification

To identify the sex of *T. californicum* hatchlings, we developed microsatellite markers on the

T. californicum X chromosome scaffolds reported by Parker et al. (in prep). *Timema* have an

XX/X0 sex determination system (Schwander and Crespi, 2009) meaning X-linked regions will

be present in two copies in females and one copy in males. Consequently, females can feature

heterozygous genotypes for polymorphic X-linked markers, while males are invariably

hemizygous. Candidate microsatellite markers were designed with msatcommander (v. 1.08,

default options) (Faircloth, 2008). A selection of four candidate markers were tested for

polymorphism and sex-linkage using 15 adult males and 15 adult females in two Multiplex

PCR Kit reactions (Qiagen; see Supplemental Table 1. for primer sequences and Supplemental

table 2. for PCR conditions). Microsatellite alleles were then determined with an ABI3100

machine (Applied Biosystem) and Genemapper v.4.1 (Applied Biosystems 2009). Across

microsatellite markers, all 15 tested females were heterozygous at minimum two markers,

whereas males were invariably hemizygous at all markers, confirming the predictions for

Timema X-linked markers (Supplemental table 2). We thus genotyped the nine hatchlings at

the four microsatellite markers to identify their sex, and were able to identify four females

and five males that were used for the transcriptome study (Supplemental table 2).

Raw data quality control, mapping and read counting

The quality of the reads was checked using FastQC v.0.11.2 (Andrews, 2014). We used

Cutadapt v. 2.3 with Python v. 3.5.2 (Martin, 2011) to remove adapter sequences. Low quality

bases at both ends of the reads were trimmed when below a Phred score of 10. Bases with

an average Phred score below 20 in a 4bp sliding window were trimmed. Finally, reads with a

length below 80 bp were removed Trimmomatic v. 0.36 (Bolger et al., 2014). Trimmed reads

were then mapped to the reference *T. californicum* genome from (Jaron et al., 2020) using

STAR v. 2.6.0c (Dobin et al., 2013). HTSeq v.0.9.1 (Anders et al., 2014) was used to count the

number of reads uniquely mapped to each gene, with the following options (htseq-count -f

bam -r name -s reverse -t gene -i ID -m union --nonunique none).

***Drosophila melanogaster* data**

D. melanogaster RNA-seq whole body data from (Ingleby et al., 2016) was downloaded from

SRA (accession number SRP068235). To mirror the data available for *Timema* and to facilitate

comparisons, four replicates per sex and three developmental stages from different

hemiclonal lines were used. A complete list of samples is provided in the supplementary

materials (Supplemental Table 3). Although reads were deposited as paired-end, we found

most of them were unpaired, thus we used only the forward reads for analysis. Reads were

trimmed, mapped to the reference genome r6.23 (Gramates et al., 2017) and counted using

the methods described above.

Differential gene expression analysis

Differential expression between the sexes was performed using edgeR v.3.16.5 (McCarthy et al., 2012; Robinson et al., 2009) in RStudio v.1.2.5019 (Team, 2019). This analysis fits a generalized linear model following a negative binomial distribution to the expression data and calculates a p-value associated with the hypothesis that gene counts differ between experimental groups. EdgeR implements a normalization method (trimmed mean of M values, TMM) to account for composition bias. We analyzed data separately for each developmental stage. We filtered out genes with low counts; we required a gene to be expressed in n-1 libraries of males or females (where n is number of replicates per stage), with the expression level > 0.5 CPM (counts per million). To test for differential gene expression between the sexes, we used a generalized linear model with quasi-likelihood F-test, with contrasts for each stage between females and males (Chen et al., 2016). To correct for multiple testing, we applied the Benjamini- Hochberg method with an alpha of 5%. Genes with significantly higher expression in males were considered as male- biased, genes with significantly higher expression in females as female- biased, and the genes without significant differential expression between sexes as un-biased. To identify genes showing a significant sex by developmental stage interaction we used a similar GLM modelling approach but with all stages included. We used SuperExactTest v.0.99.4 (Wang et al., 2015) to test if the overlap of sex-biased genes between the three developmental stages was greater than what is expected by chance. To visualize the overlap of sex-biased genes between developmental stages we used VennDiagram v.1.6.20 (Chen and Boutros, 2011). We visualized the $\log_2$FC sex-biased gene expression across developmental stages, using the pheatmap: Pretty Heatmaps v.1.0.12 (Kolde, 2018)

Stage specific gene expression

Median gene expression (CPM) values were calculated for each developmental stage and sex.

These values were used to calculate Tau, an index of specificity of gene expression over

development (Liu and Robinson-Rechavi, 2018; Yanai et al., 2004) Values of Tau range from

zero (broadly expressed over development) to one (gene expressed in only one stage). To

visualize the results we made boxplots using ggplot2 v. 3.3.2 (Wickham, 2016)

Divergence rates

To calculate sequence divergence rates, we used the 7098 one-to-one orthologs between *T.*

californicum and *T. poppensis* from (Jaron et al., 2020). Coding sequences were codon-aligned

using prank v.170427 (Loytynoja, 2014). For each of the orthologs, the number of

nonsynonymous substitutions per sites (d_N), the number of synonymous substitutions per site

(d_S) and their ratio (d_N/d_S) was calculated using PAML v4.9 (Yang, 2007). To test for differences

in sequence divergence rates (d_N/d_S) between different gene categories (female-biased,

male-biased and un-biased), we used Wilcoxon tests (ggpubr v.0.2.5, in R) with p-values

adjusted for multiple comparisons using the Benjamini-Hochberg method (Haynes, 2013).

Note that some of the *T. californicum* sex-biased genes had no identified ortholog in *T.*

poppensis, thus we do not have sequence evolution rates for all sex-biased genes. Sequence

divergence rates were obtained for 14 out of 26 SBG at the hatchling stage, 323/568 at the

juvenile stage, and 1286/2485 at the adult stage. To account for the influence of positive

selection we repeated our analyses after removing genes with evidence for positive selection

in the *T. californicum* lineage (identified in (Jaron et al., 2020)).

Furthermore, we used a partial correlation between the strength of the sex bias (log_2FC) and

the sequence divergence rate (d_N/d_S) on sex-biased genes only, while controlling for the

average expression level (CPM) of the gene in all libraries (ppcor v.01 (Kim, 2015), in R) in

order to determine if genes with higher log_2FC have faster sequence divergence rates.

Functional analysis of sex-biased genes

We performed Gene Ontology (GO) enrichment analysis using TopGO v.2.26.0 on sex-biased

genes obtained with EdgeR (see above) at each developmental stage. We used a functional

annotation derived from the blasting sequences to *Drosophila melanogaster* database. Only

GO terms with minimum ten annotated genes were used. Enrichment of terms was

determined using a weighted Kolmogorov-Smirnov-like statistical test, equivalent to the gene

set enrichment analysis (GSEA method). We applied the "elim" algorithm, which considers

the Gene Ontology hierarchy, i.e. it first assesses the most specific GO terms, and then more

general ones (Alexa et al., 2006). Our analysis focused on gene sets in the Biological Processes

(BP), GO category. We considered terms as significant when p< 0.05. Enriched GO terms were

then semantically clustered using ReviGO (Supek et al., 2011) to aid interpretation.

Results

Sex-biased gene expression increases over development in *T. californicum*

Sex-biased gene expression gradually increased over the three developmental stages, with

0.2% (26) of the expressed genes sex-biased at the hatchling stage, 4.7% (568) at the juvenile,

and 20.3% (2485) at the adult stage (Figure 2). The hatchling stage had more female than

male biased genes ($x^2_{(1)}$ =12.46, p<0.05), while the juvenile and adult stages had more male

biased genes ($x^2_{\text{juvenile (1)}}$ =457.92, $x^2_{\text{adult (1)}}$ = 77.554, p<0.05) (Supplemental Table 4). The

majority (85% and 91%) of sex biased genes at the two preadult stages had strong sex bias

($\geq$2 log$_2$FC, P_{adj}<0.05) in contrast to the adult stage where sex bias was slight (57% of the sex-

biased genes had <2 log$_2$FC, P_{adj}<0.05) (Figure 2). The intensity of sex-bias varied greatly over

development with 1137 genes showing a significant sex by development interaction

(P_{adj}<0.05).

To characterize the dynamics of sex-biased expression over development, we then classified

all 2671 genes sex-biased at one or more developmental stages into 13 categories dependent

on their expression patterns over development (Supplemental Figure 1). This classification

revealed that sex-biased genes are added gradually over development, with genes sex-biased

at earlier stages generally remaining sex-biased in the same direction at later stages

(Supplemental Figure 2). Only a single gene shifted from male-biased to female-biased

expression. Out of 26 sex-biased genes at the hatchling stage, 34% remained sex-biased

throughout development, while out of 567 sex-biased genes at the juvenile stage, 67% stayed

sex-biased at the adult stage. This classification is conservative, as it depends on the sex-

biased gene expression detection threshold.

Sex-biased genes at the hatchling stage are particularly interesting because they reveal sexual

differentiation prior to visible morphological differences. We therefore looked for functional

annotations of the 26 sex-biased genes at the hatchling stage in the *T. californicum* reference

genome from (Jaron et al., 2020). However, only 16 out of the 26 sex-biased genes had

functional annotations and only one (*vitellogenin receptor*) had a clear link to sexual

differentiation (Figure 3b). While key genes known to play a role in insect sex-determination

and differentiation pathways (*doublesex, transformer-2* and *sex-lethal*) were expressed at the

hatchling stage, *doublesex* had very low expression, preventing analysis of sex-bias, and

transformer-2 and sex-lethal did not feature sex-biased expression.

Sex-biased genes are enriched for development-related processes

Genes sex-biased at the hatchling stage were enriched for GO-terms related to

developmental processes (e.g., "regulation of cell development", "regulation of cell

proliferation", "cuticle development"), and some to female specific functions such as

oogenesis ("oocyte construction", "oocyte development") (Supplemental figure 8a, and

Supplemental table 10). At the juvenile stage, sex-biased genes were enriched for GO-terms

related to metabolic processes, in particular to catabolism (e.g., "lipid catabolic processes",

"cellular catabolic processes", "regulation of catabolic processes") (Supplemental figure 8b,

and Supplemental table 11). At the adult stage, sex-biased genes were enriched for GO-terms

related to diverse metabolic and physiological processes with no clear association to sexual

differentiation (Supplemental figure 8c, and Supplemental table 12); however, several terms

were related to chemosensory and olfactory behavior, which may play a role in mate

detection. Furthermore, adult stage, sex-biased genes were enriched for pigmentation (e.g.,

"developmental pigmentation", "eye pigmentation"), which is a sexually dimorphic trait in

Timema californicum.

Figure 2 | Gene expression (log$_2$CPM) in *T. californicum* males and females at the hatchling (left), juvenile (middle) and adult stage (right). The number of differentially expressed genes (SBG) is shown at the bottom right corner of each plot, as well as the total number of expressed genes at each stage (T). Genes are classified based on their sex-bias into seven categories: "slight FB"- female bias (<2 log$_2$FC), "strong FB"- female bias (≥2 log$_2$FC), "female limited"- with no expression in males, "slight MB"- male bias (<2 log$_2$FC), "strong MB"- male bias (≥2 log$_2$FC), "male limited"- no expression in females, "Not DE"- not differentially expressed genes.

Figure 3 | Sex-biased genes shared across development. (**a**) Venn-diagram of sex-biased

genes in three developmental stages: hatchling (yellow), juvenile (purple), and adult (green).

The number of genes shared between all three stages was greater than expected by chance

(observed overlap= 9, expected overlap= 0.007, Exact test of multi set interactions: p-value <

0.0001). (**b**) A heatmap showing the 26 sex-biased genes at the hatchling stage with their

annotations and their expression in three developmental stages. Genes in red are female-

biased, blue are male-biased. Saturation of the colors increases with the $\log_2$FC. Genes

without a label have no annotation, annotation in purple is for genes that were sex-biased in

all three stages

Sex-biased genes have more stage-specific expression

Expression levels of sex-biased genes tend to be specific to only one developmental stage

while unbiased genes have a more constant expression level across development (Figure 4).

Male-biased genes have the most stage specific expression in both males and females.

Female-biased genes are only slightly more stage specific than un-biased genes, but the trend

is consistent and the difference is significant notably in both male and female adults.

Figure 4 | Tau index of gene expression at three developmental stages in males (a) and females (b). Tau range from zero (similarly expressed over development) to one (gene expressed in only one stage). Three gene categories are depicted with different colors; female-biased in red, male-biased in blue, and un-biased in grey. Boxplots represent the median, lower and upper quartiles, and whiskers the minimum and maximum values (in the limit of 1.5x interquartile range). Adjusted p values of Wilcoxon rank sum tests are summarized above the box plots (*** = p < 0.001, * = p < 0.05, ˙p=0.06).

Sex-biased genes have faster sequence evolution rates

Male-biased genes have faster rates of sequence evolution (dN/dS) compared to both un-biased and female-biased genes, at juvenile and adult stages (Figure 5). At the hatchling stage, female-biased genes evolve faster than the un-biased genes, but not at the other two developmental stages. Elevated rates of sequence evolution are likely the result of reduced purifying selection, but stronger positive selection may also contribute to fast sequence evolution in some genes. To test this possibility, we repeated our analysis after excluding 19

313 (11 F- biased and 8 M-biased) genes for which a previous study (Jaron et al., 2020) detected

314 signatures of positive selection. Excluding these 19 genes did not change our results

315 (Supplemental Table 7a). More generally, we did not find an enrichment of positively selected

316 genes among those with sex-biased expression (Fisher's exact test; p-value=0.39, odds ratio:

317 1.1) (Supplemental Table 7b).

318  We performed partial correlation analyses to test the effect of sex bias strength on

319 sequence evolution rate, controlling for the average expression level of the gene

320 (Supplemental Table 6). For both male-biased and female-biased genes, stronger sex-bias is

321 associated with faster sequence evolution at the adult stage (Figure 6). Partial correlations

322 are not significant at juvenile or hatchling stages.

Figure 5 | Divergence rates (the ratio of nonsynonymous to synonymous substitutions, d_N/d_S),

at three developmental stages; hatchling, juvenile and adult stage. Boxplots represent the

median, lower and upper quartiles, and whiskers the minimum and maximum values (in the

limit of 1.5x interquartile range). Adjusted p values of Wilcoxon rank sum tests are

summarized above the box plots (*** = p < 0.001, * = p < 0.05).

Figure 6 | Spearman's rank correlations between the divergence rate of genes, d_N/d_S, and

their strength of the sex bias |$\log_2$FC| at three developmental stages. Note that all genes are

displayed on each plot but only male- respectively female-biased ones are used for

correlation tests.

Sex-biased gene expression increases abruptly in adult *D. melanogaster*

In order to compare the dynamics of sex-biased gene expression over development between

hemimetabolous and holometabolous insects, we reanalyzed publicly available data from *D.*

melanogaster from (Ingleby et al., 2016)*, using the same pipeline as for *T. californicum.* This

experiment was chosen for comparison as it has a similar design, i.e. whole-body RNA-seq

data from several developmental stages. In addition, although *D.* melanogaster from this

experiment were lab-reared, they had a similar between sample variance as *T. californicum*

(biological coefficient of variation at the adult stage: *D. melanogaster* = 0.266, *T. californicum*

= 0.348).

In contrast to the gradual increase in the amount of sex-biased expression over development

in *T. californicum,* sex-biased gene expression increased abruptly at the adult stage in *D.*

melanogaster (Figure 8), from 22% to 85% of sex-biased genes (Figure 7). Overall, at each

developmental stage, there were more male than female-biased genes (Supplemental table

2). The majority (85% and 63%) of sex-biased genes at the two preadult stages had strong sex

bias ($\geq$2 $\log_2$FC, P_{adj} <0.05) in contrast to the adult stage where weaker sex-bias was more

common (51% of the sex-biased genes had <2 $\log_2$FC, P_{adj} <0.05) (Supplemental table 2). Out

of 2113 sex-biased genes at the larval stage, 58% remained sex-biased throughout

development, and 93% of the pupal sex-biased genes remained sex-biased in adults (Figure

7b).

Figure 7 | (a) Gene expression (log$_2$CPM) in males and females at the larval (left), pupal

(middle) and adult stage (right). The number of differentially expressed genes (SBG) is shown

at bottom right corner of each plot, as well as the total number of expressed genes at each

stage (T). Genes are classified based on their sex-bias into seven categories: "slight FB"-

female bias (<2 log$_2$FC), "strong FB"- female bias ($\geq$2 log$_2$FC), "female limited"- with no

expression in males, "slight MB"- male bias (<2 log$_2$FC), "strong MB"- male bias ($\geq$2 log$_2$FC),

"male limited"- no expression in females, "Not DE"- not differentially expressed genes. **(b)**

Venn-diagram of sex-biased genes in three developmental stages: larval (yellow), pupal

(purple), and adult (green). The number of genes shared between all three stages was greater

than expected by chance (observed overlap= 1233, expected overlap=195.2), Exact test of

multi set interactions: p-value < 0.0001).

Figure 8 | Percentage of sex-biased genes (SBG) in *T. californicum* (blue) and *D. melanogaster* (red) at three developmental stages.

Discussion

In the stick insect *T. californicum,* gene expression differs between males and females before morphologically detectable sexual dimorphism. Over development, most of the sex-biased genes are added gradually and cumulatively, meaning that once a gene gains sex-biased expression in *T. californicum* it generally remains sex-biased at later developmental stages. In addition, genes were consistent in their sex-bias direction; only a single gene shifted from male- to female-biased expression. Surprisingly, such a gradual and consistent addition of sex-bias does not appear to be a general pattern over animal development. For example, in humans or mosquitos, some stages have bursts of sex-biased gene expression, suggesting that specific developmental stages contribute discretely to sexual differentiation, and that sexual phenotypes are not gradually established (Magnusson et al., 2011; Shi et al., 2016).

Although only 26 genes were sex-biased in *T. californicum* at the hatchling stage, they

all had a strong sex-bias and nine of them stayed sex-biased at juvenile and adult stages. As

such these 26 genes likely represent key genes involved in the sexual differentiation of

Timema. Most of the sex-biased genes at the hatchling stage were female-biased. This might

suggest that the process of building a female phenotype begins earlier in development. Later

developmental stages had more male-biased than female- biased genes. A similar shift from

female- to male-bias in gene expression during development was found in *Nasonia* jewel

wasps, where little male-biased expression was detected until the pupal stage, during the

activation of spermatogenesis. Similar to *Timema*, the shift was interpreted as a different

developmental timing of the two sexes (Rago et al., 2020). In *Timema*, male-biased genes

were also expressed in a relatively stage-specific manner, in contrast to a more constant

expression of female-biased genes. This further supports the idea that *Timema* individuals

first develop along the female trajectory (or the female phenotype being the "default" in

Timema) and may indicate greater pleiotropy of female-biased genes. Additional support for

this interpretation comes from the lower sequence divergence rates of female-biased genes

than male-biased genes, implicating stronger selective constraints on female- than male-

biased genes. In addition, we observed that genes with greater sex-bias have faster sequence

divergence rates. This is largely due to relaxed purifying selection, with positive selection

contributing little to the accelerated evolutionary rate of sex-biased genes. These findings are

consistent with the idea that sex-biased genes evolve from genes with few evolutionary

constraints and that are relatively "dispensable" (Catalan et al., 2018; Mank and Ellegren,

2009).

Sex-biased genes in *Timema* hatchlings were enriched for processes related to

oogenesis. Moreover, one of the most consistently and strongly female-biased genes across

development was annotated as *vitellogenin receptor*. In adult flies, *vitellogenin receptor*

transfers yolk protein precursors into the oocytes and is necessary for the development of

the oocytes and later embryo (Liu et al., 2016). The role of these processes in *Timema*

hatchlings remain to be investigated. Sex-biased genes in juvenile and adult stages were not

enriched for processes obviously related to sex- specific functions. However, some processes

such as pigmentation and chemosensory/olfactory behavior could have a sex-related role.

Pigmentation is a sexually dimorphic trait in *Timema* (Sandoval, 2008), while chemosensory

and olfactory behavior are important for mate recognition and mating (Parker et al., 2019b;

Schwander et al., 2013). We additionally looked for the expression of two key genes involved

in insect sex-determination pathways: *transformer* and *doublesex*. In holometabolous insects,

transformer acts as a splicing regulator of *doublesex*, a sex differentiation master switch gene

(Geuverink and Beukeboom, 2014). The *Doublesex- Transformer* role in sex-differentiation is

conserved among Diptera, Coleoptera and Hymenoptera (Verhulst et al., 2010; Wexler et al.,

2019). Emerging studies suggest that *doublesex* and *transformer* likely play the same role in

sex differentiation in hemimetabolous insects (Wexler et al., 2019; Zhuo et al., 2018). In *T.*

californicum, we found that *doublesex* was expressed in all three developmental stages, but

too lowly for analyzing sex-bias. *Transformer* is not annotated in the available *T. californicum*

genome; a *Transformer-2 homolog* is annotated but did not feature sex-bias at any of the

three developmental stages. As such, whether or not these classic sex-determining genes

influence sexual differentiation in *Timema* is unclear, requiring future studies.

We hypothesized that the dynamics of sex biased gene expression would be strongly

affected by hemimetabolous vs holometabolous development in insects, with a gradual

increase of sex-biased gene expression over development in the former and an abrupt

increase at the adult stage in the latter. Hemimetabolous insects such as *T. californicum* have

multiple nymphal stages that progressively resemble the adult stage, together with a gradual

increase in sexual dimorphism. On the other hand, a holometabolous insect like *D.*

melanogaster has more monomorphic larval stages and pupae, with no resemblance to the

sexually dimorphic adult stage. The change in sexually dimorphic gene expression across

development mirrored these different developmental patterns, with the proportion of sex-

biased genes increasing gradually in *T. californicum* as sexual dimorphism became more

pronounced, whereas sexually dimorphic gene expression increased abruptly at the adult

stage in *D. melanogaster*.

Although the change in sexually dimorphic gene expression across development fit

with our expectations, we were surprised to find a much higher proportion of sex-biased

genes in *D. melanogaster* than in *T. californicum* overall. This difference is clearest at the adult

stage when most of the expressed genes in *D. melanogaster* were sex-biased (85%),

compared to only 20% in *T. californicum*. Similar to *Drosophila*, other holometabolous insects

also featured sex-bias for more than half of the genes expressed at the adult stage (Dean A

Baker et al., 2011; Rago et al., 2020), while all studied hemimetabolous insects had a much

smaller proportion of sex-biased genes (<10%) (Pal and Vicoso, 2015). Why there is such a

difference in the prevalence of sex-biased genes between species with different

developmental strategies is still an open question. One potential explanation is that the

proportion of sex-biased genes at the adult stage may be less constrained under holo- than

hemimetabolous development. For example, the complete reorganization of cells and tissues

during metamorphosis in holometabolous insects may allow for more drastic phenotypic

changes between the sexes, while incomplete metamorphosis constrains the evolution of

distinct sexual phenotypes. Data on additional species are required to confirm that this

difference is not simply due to the species investigated, but is indeed general.

Holometabolous insects studied are often established lab-models, with high fecundity, and

may thus be characterized by comparatively large gonads, which would result in large

proportions of sex-biased genes in adults. By contrast, all hemimetabolous insects

characterized for sex-biased expression are non-model species. However, the choice of model

species is unlikely to be the sole explanation for the different amount of sex bias in holo- and

hemimetabolous species. Indeed, in crickets and *Timema* stick insects, even the most sexually

differentiated tissue (gonads) feature sex-bias of fewer than 30% of genes (Parker et al.,

2019a; Whittle et al., 2020), suggesting that extensive differences are not completely driven

by large gonads.

Overall, our results describe the dynamics of sex- biased gene expression over

development in a hemimetabolous insect, and suggest profound differences of these

dynamics between holometabolous and hemimetabolous species. We showed that the

extent of sex-biased gene expression follows the development of sexual dimorphism.

Generating sexually dimorphic phenotypes is a developmental process, and sex-biased genes

are added gradually over development, with the majority of early sex-biased genes

consistently sex-biased throughout development.

Data availability

Raw reads have been deposited in SRA under PRJNA678950 bioproject, with accession codes

SRR13084978- SRR13085001 (Supplemental table 9). Mapped read counts are provided in

Supplemental table 13 and fold changes in Supplemental table 14.

Acknowledgment

This study was supported by Swiss Natural Science Foundation Grant 31003A_182495.

Bibliography

Alexa, A., Rahnenführer, J., and Lengauer, T. (2006). Improved scoring of functional groups from gene expression data by decorrelating GO graph structure. Bioinformatics *22*, 1600-1607.

Anders, S., Pyl, P.T., and Huber, W. (2014). HTSeq—a Python framework to work with high-throughput sequencing data. Bioinformatics *31*, 166-169.

Andrews, S. (2014). FastQC. v.0.11.2.

Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics *30*, 2114-2120.

Catalan, A., Macias-Munoz, A., and Briscoe, A.D. (2018). Evolution of sex-biased gene expression and dosage compensation in the eye and brain of Heliconius butterflies. Mol Biol Evol *35*, 2120-2134.

Chauhan, P., Wellenreuther, M., and Hansson, B. (2016). Transcriptome profiling in the damselfly Ischnura elegans identifies genes with sex-biased expression. BMC Genomics *17*, 985.

Chen, H., and Boutros, P.C. (2011). VennDiagram: a package for the generation of highly-customizable venn and euler diagrams in R. BMC Bioinformatics *12*.

Chen, S., Yang, P., Jiang, F., Wei, Y., Ma, Z., and Kang, L. (2010). De novo analysis of transcriptome dynamics in the migratory locust during the development of phase traits. PLoS One *5*, e15633.

Chen, Y., Lun, A.T., and Smyth, G.K. (2016). From reads to genes to pathways: differential expression analysis of RNA-Seq experiments using Rsubread and the edgeR quasi-likelihood pipeline. F1000Res *5*, 1438.

Dean A Baker, Tony Nolan, Bettina Fischer, Alex Pinder, and, A.C., and Russell, S. (2011). A

comprehensive gene expression atlas of sex- and tissue- specificity in the malaria

vector, Anopheles gambiae. BMC genomics *12*, 12.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M.,

and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*,

15-21.

Ellegren, H., and Parsch, J. (2007). The evolution of sex-biased genes and sex-biased gene

expression. Nat Rev Genet *8*, 689-698.

Faircloth, B.C. (2008). Msatcommander: detection of microsatellite repeat arrays and

automated, locus-specific primer design. Mol Ecol Resour *8*, 92-94.

Geuverink, E., and Beukeboom, L.W. (2014). Phylogenetic distribution and evolutionary

dynamics of the sex determination genes doublesex and transformer in Insects. Sexual

Development *8*, 38-49.

Gramates, L.S., Marygold, S.J., Santos, G.D., Urbano, J.M., Antonazzo, G., Matthews, B.B., Rey,

A.J., Tabone, C.J., Crosby, M.A., Emmert, D.B.*, et al.* (2017). FlyBase at 25: looking to the

future. Nucleic Acids Res *45*, D663-D671.

Grath, S., and Parsch, J. (2012). Rate of amino acid substitution is influenced by the degree

and conservation of male-biased transcription over 50 myr of Drosophila evolution.

Genome Biol Evol *4*, 346-359.

Haynes, W. (2013). "Benjamini–Hochberg method". In Encyclopedia of Systems Biology, W.

Dubitzky, O. Wolkenhauer, K.-H. Cho, and H. Yokota, eds. (New York, NY: Springer New

York), pp. 78-78.

Ingleby, F., Webster, C., Pennell, T., Flis, I., and Morrow, E. (2016). Sex-biased gene expression

in Drosophila melanogaster is constrained by ontogeny and genetic architecture.

bioRxiv, 034728.

Ingleby, F.C., Ilona, F., and Morrow, E.H. (2015). Sex-biased gene expression and sexual

conflict throughout development. Cold Spring Harbor Laboratory Press *7*, a017632.

Jaron, K.S., Parker, D.J., Anselmetti, Y., Van, P.T., Bast, J., Dumas, Z., Figuet, E., François, C.M.,

Hayward, K., Rossier, V.*, et al.* (2020). Convergent consequences of parthenogenesis on

stick insect genomes. bioRxiv *2020.11.20.391540*

Khila, A., Abouheif, E., and Rowe, L. (2012). Function, developmental genetics, and fitness

consequences of a sexually antagonistic trait. Science *336*, 585-589.

Kim, S. (2015). ppcor: an R package for a fast calculation to semi-partial correlation

coefficients. Commun Stat Appl Methods *22*, 665-674.

Kolde, R. (2018). pheatmap: Pretty Heatmaps, R package version 1.0.12.

Lande, R. (1980). Sexual dimorphism, sexual selection, and adaptation in polygenic

characters. Evolution *34*, 292-305.

Liu, J., and Robinson-Rechavi, M. (2018). Developmental constraints on genome evolution in

four bilaterian model species. Genome Biol Evol *10*, 2266-2277.

Liu, L., Wang, Y., Li, Y., Lin, Y., Hou, Y., Zhang, Y., Wei, S., Zhao, P., Zhao, P., and He, H. (2016).

LBD1 of vitellogenin receptor specifically binds to the female-specific storage protein

SP1 via LBR1 and LBR3. PLOS ONE *11*, e0162317.

Lowe, R., Gemma, C., Rakyan, V.K., and Holland, M.L. (2015). Sexually dimorphic gene

expression emerges with embryonic genome activation and is dynamic throughout

development. BMC Genomics *16*, 295.

Loytynoja, A. (2014). Phylogeny-aware alignment with PRANK. Methods Mol Biol *1079*, 155-

170.

Magnusson, K., Mendes, A.M., Windbichler, N., Papathanos, P.A., Nolan, T., Dottorini, T., Rizzi,

E., Christophides, G.K., and Crisanti, A. (2011). Transcription regulation of sex-biased

genes during ontogeny in the malaria vector Anopheles gambiae. PLoS One *6*, e21572.

Mank, J.E. (2017). The transcriptional architecture of phenotypic dimorphism. Nat Ecol Evol

1, 6.

Mank, J.E., and Ellegren, H. (2009). Are sex-biased genes more dispensable? Biol Lett *5*, 409-

412.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing

reads. EMBnetjournal *17*, 10-12.

McCarthy, D.J., Chen, Y., and Smyth, G.K. (2012). Differential expression analysis of

multifactor RNA-Seq experiments with respect to biological variation. Nucleic Acids Res

40, 4288-4297.

Pal, A., and Vicoso, B. (2015). The X Chromosome of Hemipteran Insects: Conservation,

Dosage Compensation and Sex-Biased Expression. Genome Biol Evol *7*, 3259-3268.

Paris, M., Villalta, J.E., Eisen, M.B., and Lott, S.E. (2015). Sex bias and maternal contribution

to gene expression divergence in Drosophila blastoderm Embryos. PLOS Genetics *11*,

e1005592.

Parker, D.J., Bast, J., Jalvingh, K., Dumas, Z., Robinson-Rechavi, M., and Schwander, T. (2019a).

Sex-biased gene expression is repeatedly masculinized in asexual females. Nat Commun

10, 4638.

Parker, D.J., Djordjevic, J., and Schwander, T. (2019b). Olfactory proteins in Timema stick

insects. Frontiers in Ecology and Evolution *7*.

Perry, J.C., Harrison, P.W., and Mank, J.E. (2014). The ontogeny and evolution of sex-biased

gene expression in Drosophila melanogaster. Mol Biol Evol *31*, 1206-1219.

Pointer, M.A., Harrison, P.W., Wright, A.E., and Mank, J.E. (2013). Masculinization of gene

expression is associated with exaggeration of male sexual dimorphism. PLoS Genet *9*,

e1003697.

Rago, A., Werren, J.H., and Colbourne, J.K. (2020). Sex biased expression and co-expression

networks in development, using the hymenopteran Nasonia vitripennis. PLoS Genet *16*,

e1008518.

Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2009). edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. Bioinformatics *26*, 139-

140.

Sandoval, C.P. (1994). Geographic, ecological and behavioral factors affecting spatial variation

in color morph frequency in the walking-stick, Timema cristinae.

Sandoval, C.P. (2008). Differential visual predation on morphs of Timema cristinae

(Phasmatodeae:Timemidae) and its consequences for host range. Biological Journal of

the Linnean Society *52*, 341-356.

Schwander, T., and Crespi, B.J. (2009). Multiple direct transitions from sexual reproduction to

apomictic parthenogenesis in Timema stick insects. Evolution *63*, 84-103.

Schwander, T., Crespi, B.J., Gries, R., and Gries, G. (2013). Neutral and selection-driven decay

of sexual traits in asexual stick insects. Proceedings of the Royal Society B: Biological

Sciences *280*, 20130823.

Shi, L., Zhang, Z., and Su, B. (2016). Sex biased gene expression profiling of human brains at

major developmental stages. Sci Rep *6*, 21181.

Supek, F., Bosnjak, M., Skunca, N., and Smuc, T. (2011). REVIGO summarizes and visualizes

long lists of gene ontology terms. PLoS One *6*, e21800.

Team, R. (2019). RStudio: Integrated Development Environment for R.

Verhulst, E.C., van de Zande, L., and Beukeboom, L.W. (2010). Insect sex determination: it all

evolves around transformer. Curr Opin Genet Dev *20*, 376-383.

Vickery, V.R., and Sandoval, C.P. (2001). Descriptions of three new species of Timema

(Phasmatoptera: Timematodea: Timematidae) and notes on three other species.

Journal of Orthoptera Research *10*, 53-61.

Wang, M., Zhao, Y., and Zhang, B. (2015). Efficient test and visualization of multi-set

intersections. Sci Rep *5*, 16923.

Weigmann, K., Klapper, R., Strasser, T., Rickert, C., Technau, G., Jäckle, H., Janning, W., and

Klämbt, C. (2003). FlyMove- a new way to look at development of Drosophila. Trends in

Genetics *19*, 310-311.

Wexler, J., Delaney, E.K., Belles, X., Schal, C., Wada-Katsumata, A., Amicucci, M.J., and Kopp,

A. (2019). Hemimetabolous insects elucidate the origin of sexual development via

alternative splicing. eLife *8*, e47490.

Whittle, C.A., Kulkarni, A., and Extavour, C.G. (2020). Sex-biased genes expressed in the

cricket brain evolve rapidly. bioRxiv *2020.07.07.192039*.

Wickham, H. (2016). ggplot2: Elegant Graphics for Data Analysis (Springer-Verlag New York).

Yanai, I., Benjamin, H., Shmoish, M., Chalifa-Caspi, V., Shklar, M., Ophir, R., Bar-Even, A., Horn-

Saban, S., Safran, M., Domany, E.*, et al.* (2004). Genome-wide midrange transcription

profiles reveal expression level relationships in human tissue specification.

Bioinformatics *21*, 650-659.

Yang, Z. (2007). PAML 4: phylogenetic analysis by maximum likelihood. Molecular Biology and

Evolution *24*, 1586-1591.

Zhuo, J.C., Hu, Q.L., Zhang, H.H., Zhang, M.Q., Jo, S.B., and Zhang, C.X. (2018). Identification

and functional analysis of the doublesex gene in the sexual development of a

hemimetabolous insect, the brown planthopper. Insect Biochem Mol Biol *102*, 31-42.