1   P4HA2-induced prolyl hydroxylation suppresses YAP1-mediated prostate cancer cell migration,

2   invasion, and metastasis

Ming Zhu[1]*, Ruiqing Peng[1, 7]*, Xin Liang[1], Zhengdao Lan[2], Ming Tang[3], Pingping Hou[2], Jian H.

Song[1], Celia Sze Ling Mak[1], Jiwon Park[1], Shui-er Zheng[1], Ailing Huang[1], Xingdi Ma[2], Ruidong

Chen[1], Qing Chang[3], Christopher J. Logothetis[1], Abhinav K. Jain[4], Sue-Hwa Lin[1,5], Hiroyuki

Katayama[6], Samir M. Hanash[6], Guocan Wang[1,2].

[1]Department of Genitourinary Medical Oncology and the David H. Koch Center for Applied

Research of Genitourinary Cancers, [2]Department of Cancer Biology, [3]Department of Genomic

Medicine, [4]Department of Epigenetics and Molecular Carcinogenesis & Epigenomics Profiling

Core Facility, [5]Department of Translational Molecular Pathology, [6]Department of Clinical

Cancer Prevention, The University of Texas MD Anderson Cancer Center, Houston, Texas,

USA. [7]Sun Yat-sen University Cancer Center; State Key Laboratory of Oncology in South

China; Collaborative Innovation Center for Cancer Medicine, Guangzhou, China.

*These authors contributed equally to the work.

Correspondence: Guocan Wang (gwang6@mdanderson.org)

Running title: The role of prolyl hydroxylation of YAP1 in prostate cancer

Key words: YAP1, prolyl hydroxylation, P4A2, prostate cancer, cell migration, invasion,

metastasis

Conflict of interest: C. J. Logothetis reports receiving commercial research grants from Bayer,

Sanofi, Janssen, Astellas Pharma, Pfizer; and honoraria from Bayer, Janssen, Sanofi, Astellas

Pharma. No potential conflicts of interest were disclosed by the other authors.

ABSTRACT

Yes-associated protein 1 (YAP1), a key player in the Hippo pathway, has been shown to play a critical role in tumor progression. However, the role of YAP1 in prostate cancer cell invasion, migration, and metastasis is not well defined. Through functional, transcriptomic, epigenomic, and proteomic analyses, we showed that prolyl hydroxylation of YAP1 plays a critical role in the suppression of cell migration, invasion, and metastasis in prostate cancer. Knockdown (KD) or knockout (KO) of *YAP1* led to an increase in cell migration, invasion, and metastasis in prostate cancer cells. Microarray analysis showed that the EMT pathway was activated in *Yap1*-KD cells. ChIP-seq analysis showed that YAP1 target genes are enriched in pathways regulating cell migration. Mass spectrometry analysis identified P4H prolyl hydroxylase in the YAP1 complex and YAP1 was hydroxylated at multiple proline residues. Proline-to-alanine mutations of YAP1 isoform 3 identified proline 174 as a critical residue, and its hydroxylation suppressed cell migration, invasion, and metastasis. KO of *P4ha2* led to an increase in cell migration and invasion, which was reversed upon *Yap1* KD. Our study identified a novel regulatory mechanism of YAP1 by which P4HA2-dependent prolyl hydroxylation of YAP1 determines its transcriptional activities and its function in prostate cancer metastasis.

INTRODUCTION

Yes-associated protein 1 (YAP1), a key transcriptional coactivator in the Hippo pathway, is an important driver in cancer development and progression (1). Although YAP1 plays an oncogenic role in various cancer types, multiple studies also support a tumor-suppressive function for YAP1 in head and neck (2), breast (3-5), hematological (6), and colorectal (7, 8) cancers. Thus, the functions of YAP1 are likely context-dependent (9). YAP1 was shown to be overexpressed in prostate adenocarcinoma (PCa) and associated with cell proliferation and invasiveness in castration-resistant prostate cancer (CRPC) (10-12). However, YAP1 was found to be downregulated in the highly aggressive NEPC subset (13). Importantly, *YAP1* deletion (heterozygous and homozygous) and mutation were observed in ~3.6% of prostate cancers and was strongly associated with metastasis (**Suppl. Fig. 1A-B & Suppl. Table 1-3**). On the contrary, deletion/mutation of TAZ, another transcriptional coactivator in the Hippo pathway, was not significantly different between the primary and metastatic PCa (**Suppl. Fig. 1B**). On the cellular level, YAP1 promotes prostate cancer cell proliferation through cell-autonomous and non-autonomous mechanisms (11, 12, 14, 15), but its role in prostate cancer metastasis is not clearly defined.

Post-translational modification of YAP1, such as phosphorylation, has also been shown to regulate YAP1 cellular localization, stability, and activities (16, 17). Interestingly, we found that YAP1 proteins in prostate cancer cells are modified by proline hydroxylation, an important post-translational modification that modulates protein folding and stability in mammalian cells (18, 19). Proline hydroxylation is induced by prolyl hydroxylases, such as prolyl hydroxylase domain proteins (PHD) and collagen prolyl 4-hydroxylase (P4H). Whether YAP1 is subjected to proline hydroxylation was previously unknown, as were the effects of such modification on

YAP1 function. In this study, we identified a surprising role for YAP1 in the suppression of cell

migration, invasion, and metastasis in prostate, pancreatic, and breast cancers. We found that

YAP1 interacts with the P4H complex and is hydroxylated at multiple proline residues. The

status of proline hydroxylation of YAP1 determines its oncogenic activity in regulating cell

migration, invasion, and metastasis in prostate cancer and possibly in other cancer cell types.

RESULTS

YAP1 suppresses cancer cell migration, invasion, and metastasis

Previously, we showed that YAP1 was highly expressed in primary tumors from the metastatic

Pten/Smad4 prostate conditional knockout (KO) model (12). We first examined the effect of

Yap1 knockdown (KD) on cell migration, invasion, and metastasis in a highly metastatic

Pten/Smad4-deficient CRPC cell line (referred to as PS cells hereafter) (20). Surprisingly, *Yap1*

KD and *Yap1* KO led to a significant increase in cell migration and invasion (**Fig. 1A-C &**

Suppl. Fig. 2A). Of note, *Yap1* KD or KO did not have a significant effect on cell proliferation

as measured by the total number of cells at the end of the assays (**data not shown**). Furthermore,

re-expression of human *YAP1* in *Yap1*-KO cells suppressed cell migration and invasion (**Fig. 1D**

& Suppl. Fig. 2B). *YAP1*-KD in C4-2b, IGR-CaP1, and PC3 cells similarly increased migration

and invasion (**Fig. 1E-G & Suppl. Fig. 2C**). However, *YAP1* KD in DU145 led to a decrease in

cell migration (**Suppl. Fig. 2D**). Also, we examined whether YAP1 also suppressed cell

migration in other metastatic cancers in which YAP1 has been implicated to play an important

role in tumor progression (21-25). We found that *Yap1* KO or KD led to increased cell migration

in iKPC mouse pancreatic cancer cells (26) and MDA-MB-231 human breast cancer cells (**Fig.**

1H-I) but not in SYO-1 synovial sarcoma cells (**Suppl. Fig. 2E**). Moreover, *Yap1* KD in PS cells

promoted lung metastasis (**Fig. 1J**). Taken together, our data suggest that YAP1 suppresses cell

migration, invasion, and metastasis in multiple cancer types.

YAP1 interacts with the prolyl 4-hydroxylase complex, and its proline residues are

hydroxylated

To understand the mechanisms by which YAP1 suppresses cell migration, invasion, and

metastasis, we performed microarray analysis of RNA isolated from *Yap1*-KD and control PS

cells (**Suppl. Table 4**). As expected, gene set enrichment analysis (GSEA) (27) identified

epithelial to mesenchymal transition (EMT) as the top pathway activated in *Yap1*-KD cells (**Fig.**

2A). We also confirmed that the expression of several EMT genes, including *Postn*, *Cdh11*,

Acta2, *Rgs4*, and *Mgp*, were upregulated upon *Yap1* KD (**Suppl. Fig. 3A**).

Since YAP1 acts as a transcriptional coactivator, we sought to determine whether these

upregulated EMT genes are direct target genes of YAP1 using ChIP-seq in PS cells. We found

that YAP1 binds mostly to the distant intergenic region and other introns (**Fig. 2B**), which is

consistent with previous reports (28). Pathway analyses of the top 2000 YAP1 binding sites

using Gene Ontology (GO) and the Kyoto Encyclopedia of Genes and Genomes (KEGG)

identified multiple pathways related to cell migration as top pathways and the expected "Hippo

signaling pathway" (**Fig. 2C & Suppl. Fig. 3B**). Motif analysis using Homer (29) identified

TEAD1 and AP1 motifs (**Suppl. Fig. 3C & data not shown**), which is consistent with the

known physical and functional interaction between YAP1, TEAD, and AP1 (28). Importantly,

we found that 194 upregulated genes and 247 downregulated genes in *Yap1* KD cells were

among the top 6000 YAP1-target genes predicted by Cistrome-GO (30) (**Suppl. Table 5-7**).

Among these genes, YAP1 binds to the distant intergenic region upstream or downstream of

several genes upregulated in *Yap1*-KD cells (e.g., *Postn*, *Cdh11*, *Acta2*, *Rgs4*, and *Mgp*) (**Suppl.**

Fig. 3D & data not shown), suggesting that YAP1 directly represses their expression.

Additionally, we confirmed the binding of YAP1 to its known target genes, including *Cxcl5* and

Ccnd1 (**Suppl. Fig. 3E**).

Since the functions of YAP1 are regulated through its interacting partners, as well as by

post-translational modifications (16, 17), we performed immunoprecipitation (IP) of YAP1 in PS

cells followed by liquid chromatography-tandem mass spectrometry (LC-MS/MS) to identify

YAP1-interacting proteins and novel post-translational modifications (**Fig. 2D**). As expected, we

identified multiple proteins previously shown to interact with YAP1, such as AMOT and the

SWI/SNF complex (**Fig. 2D**). Interestingly, proteins of P4H complex (P4HA1, P4HA2, and

P4HB) were identified among the top YAP1-interacting proteins (**Fig. 2D**). On the contrary,

prolyl 3-hydroxylase was not identified in the IP-MS (**data not shown**). Collagen P4H, an $\alpha 2\beta 2$

tetrameric complex, specifically catalyzes 4-hydroxylation of proline (18, 19) through its

catalytic α subunit (P4HA). Because P4HA2 was the most abundant protein of the P4H complex

pulled down by YAP1, we focused on its interaction with YAP1. We showed that overexpressed

Flag-YAP1 efficiently pulled down overexpressed P4HA2 in 293T cells (**Fig. 2E**). Also,

endogenous YAP1 was found to interact with P4HA2 in PS cells (**Fig. 2F**).

Give the proline hydroxylase activity of the P4H complex (18, 19), we examined whether

proline residues in YAP1 were hydroxylated. We identified nine hydroxylated proline residues

(proline 60, 70, 105, 157, 159, 347, 353, 395, 398) in mouse YAP1, eight of which were

evolutionarily conserved between mouse and human (**Fig. 2G & Suppl. Fig. 4A-B**), suggesting

that they might play a role in regulating YAP1 functions.

Hydroxylation at proline 174 of YAP1 plays a critical role in suppressing cell migration, invasion, and metastasis

Given the critical regulatory roles of proline hydroxylation in proteins (18, 19), we decided to examine whether proline hydroxylation modulates YAP1 functions. We first generated hydroxylation-defective human YAP1 mutants by mutating proline to alanine (Mut1-3: P75/85/120A; Mut4-5: P172/174A; Mut6-9: P348/352/394/397A) (**Fig. 3A & Suppl. Fig. 4B**). These YAP1 mutants were overexpressed in *Yap1*-KO PS cells to avoid the possible interference of the endogenous wild type (WT) YAP1. The expression of all the YAP1 mutants was similar but higher than the WT (**Fig. 3B**). We did not observe any significant difference in cell growth *in vitro* and *in vivo* (**Suppl. Fig. 5A-B**) between YAP1 WT and mutants. However, we found that there is an increase in cell migration and invasion in Mut4-5–overexpressing cells compared to WT-overexpressing cells (**Fig. 3C-D**), suggesting that mutations of proline 172 and 174 to alanine abolished the activity of YAP1 in suppressing cell migration and invasion. Consistent with the *in vitro* findings, PS cells with YAP1 Mut4-5 overexpression increased the colonization of cancer cells in the lung compared to WT, Mut1-3, and Mut6-9 (**Fig. 3E & Suppl. Fig. 5C**).

To determine whether prolyl hydroxylation controls the transcriptional activities of YAP1, we examined the expression of YAP1 target genes *Ccnd1*, *Mcm6*, *Cxcl1*, and *Cxcl5* in *Yap1*-KO cells that overexpressed YAP1 WT, Mut4-5, or the constitutively active S127A mutant (31). Both Mut4-5 and the S127A mutant dramatically increased the expression of these genes compared to the YAP1 WT (**Fig. 3F & Suppl. Fig. 5D**), suggesting that the non-hydroxylated YAP1 is more transcriptionally active than the hydroxylated YAP1.

To further pinpoint which proline residue of YAP1 is critical for its function in suppressing cell migration and invasion, we generated site-specific proline-to-alanine mutants of

human YAP1 (P172A and P174A mutants), which corresponded to proline 157 and 159 in

mouse YAP1. We overexpressed GFP control, YAP1 WT, and YAP1 mutants (Mut4-5:

P172/174A; Mut4: P172A; Mut5: P174A) in *Yap1*-KO cells (**Suppl. Fig. 5E**) and examined their

effects on cell migration and invasion. We found that both Mut4-5 and Mut5, but not Mut4,

dramatically increased cell migration and invasion compared to the GFP control (**Fig. 3G-H**),

suggesting proline 174 is the critical hydroxylation site that regulates YAP1 activity in cell

migration and invasion. Importantly, overexpression of Mut4-5 and Mut5 similarly increased the

cell migration of PC3 and TRAMPC2 cells (**Suppl. Fig. 5F**). We then performed ChIP-qPCR to

determine whether the increased expression of YAP1 target genes in Mut4-5 expressing cells is

due to increased YAP1 binding to chromatin. We found that Mut5 binding to the promoter/distal

enhancers of its target genes was significantly increased compared to WT (**Fig. 3I & Suppl. Fig.**

5G). To examine the transcriptional activation and repression of YAP1 target genes, we

examined H3K9me3, a mark associated with transcriptional repression (32), and H3K4me3, a

hallmark of active chromatin enriched at active promoters and correlates with transcriptional

activity (33), in the promoters/enhancers of YAP1 target genes. We found that H3K9me3 was

significantly decreased and H3K4me4 was significantly increased in the regulatory region of

several YAP1 target genes (e.g., *Col12a1*, *Mgp*, *Postn*, *Cxcl12*) (**Fig. 3I & Suppl. Fig. 5G**).

Taken together, our data suggest that hydroxylation at proline 174 of YAP1 plays a critical role

in regulating cell migration, invasion, and metastasis by repressing the expression of a subset of

its target genes.

Loss of P4HA2 promotes cell migration and invasion through YAP1

Given the observed hydroxylation of YAP1 in PS cells (**Fig. 2G**) and the effect of hydroxylation-

defective YAP1 mutants on migration and invasion (**Fig. 3C-H & Suppl. Fig. 5F**), the ability of

YAP1 to suppress cell migration and invasion appeared to be regulated by the P4H complex. We

examined the effect of *P4ha2* KO on cell migration and invasion and found that *P4ha2* KO

significantly increased cell migration and invasion compared to WT cells (**Fig. 4A-C**), which

was not due to an increase in cell proliferation (**data not shown**). Interestingly, we found that

YAP1 target genes *Postn*, *Col12a1*, *Mgp, Ccl5, and Cxcl12* were significantly upregulated in

P4ha2-KO cells compared to control cells (**Fig. 4D & data not shown**), suggesting YAP1 is

transcriptionally more active upon loss of P4HA2. Importantly, KD of *Yap1* in *P4ha2*-KO cells

abolished the effect of *P4ha2* KO on cell migration and invasion (**Fig. 4E-F & Suppl. Fig. 5H**).

Taken together, our data indicate that P4HA2 suppresses cell migration and invasion through

proline hydroxylation of YAP1.

DISCUSSION

In contrary to previous findings that KD of *YAP1* in LNCaP-C4-2 cells impaired cell migration

and invasion, we demonstrated that KD or KO of *Yap1* in mouse (PS and TRAMPC2) and

human (C4-2b, IGR-CaP1, and PC3) prostate cancer cells led to enhanced cell migration,

invasion, and metastasis. The discrepancy between our study and the previous one is not clear.

Our data also showed that *YAP1* KD in DU145 cells suppressed cell migration whereas YAP1

KD/KO promoted cell migration in iKPC cells and MDA-MB-231 cells. These findings strongly

suggest that YAP1 plays a context-dependent function in cell migration, invasion, and

metastasis. Further studies are necessary to define molecular basis underlying the context-

dependent functions of YAP1 in cell migration, invasion, and metastasis. Importantly, the

clinical significance of YAP1 loss in PCa patients was supported by the strong association of

YAP1 deletion with metastatic PCa and the loss of YAP1 protein in advanced PCa (12, 13). Loss

of YAP1 function via post-translational modification by proline hydroxylation will be also an

important mechanism of clinical significance. Furthermore, our unpublished data showed that

TAZ similarly suppresses cell migration and regulates a common set of genes as YAP1,

suggesting functional redundancy between YAP1 and TAZ.

Mechanistically, our data suggest that prolyl hydroxylation plays an important role in the

regulation of YAP1 activities, which can both activate and repress transcription (**Fig. 4G**). In

P4HA2 WT cells, YAP1 may suppress gene expression through its interaction with SWI/SNF

complex, which was identified as YAP1-interating proteins in our study and has been shown to

regulate both activation and repression of the same promoters (34, 35), in part through

corepressor NCoR1 (36). Also, YAP1 may recruit YY1 and EZH2 (37) or recruit the NuRD

complex to suppress the expression of its target genes (38). Interestingly, proline 174 is within

the first WW domain of YAP1, which is crucial for the transcriptional activities of YAP1

through its interaction with transcription factors that contain PPxY motifs (39). Thus, our

findings suggest that prolyl hydroxylation at P174 of YAP1 may impair its interaction with key

transcription [e.g., c-JUN (28), TEAD], resulting in a decrease in both binding to its target gene

and reduced transcriptional activation. This notion is supported by our findings that YAP1-Mut5

binds more efficiently to its target genes. YAP1 hydroxylation may also be required for the

efficient recruitment of the transcription corepressor complexes (e.g., NCoR1, NuRD), as our

data showed that YAP1 P174A OE increased H3K4me3 and reduced H3K9me3 for a subset of

YAP1 target genes. Thus, our data suggest that P174A YAP1 mutant not only has increased

binding to its target genes, but also induces a switch from repressive transcription to active

transcription for a subset of genes. Further studies are needed to delineate the effect of prolyl

hydroxylation of YAP1 on the dynamics of the epigenomic landscape. YAP1 is known to be

regulated by phosphorylation at multiple serine residues (40). Our studies shed light on a new

aspect of YAP1 regulation, which may also be involved in many YAP1-mediated cellular

activities yet to be identified.

The P4H α subunit (P4HA) has three isoforms (P4HA1-3) in mammalian cells (19).

Since P4HA1 and P4HA2 were both identified as YAP1-interacting proteins in our MS analysis,

P4HA1 may also suppress cell migration, invasion, and metastasis through prolyl hydroxylation

of YAP1. However, P4HA1 was previously shown to promote prostate cancer progression (41),

and further studies are needed to clarify the role of P4HA1 in regulating YAP1 functions and cell

migration, invasion, and metastasis. Since LS-MS mass spectrometry analysis cannot distinguish

3-prolyl hydroxylation from 4-prolyl hydroxylation, we cannot rule out the presence of 3-

hydroxyl proline in YAP1. Given that P3H1 was not identified as YAP1-interacting proteins, the

high specificity of the P4H and P3H towards prolyl hydroxylation strongly indicates that the

hydroxyproline identified in YAP1 is 4-hydroxyproline. Future experiments combining liquid

chromatography retention time differences with mass spectrometry using ETD-HCD

fragmentation, complemented by ab initio calculations are needed to address this issue (42).

Although collagen deposition is generally associated with tumor progression and invasive

behavior (43), it also plays a tumor-suppressive role (44). P4HA1 is the major isoenzyme in most

cells, and *P4ha1−/−* leads to embryonic lethality in mice due to abnormal deposition of collagen

IV (45). In contrast, *P4ha2−/−* mice had no apparent abnormalities (46). Given that we did not

observe any significant difference in the expression of P4HA1 between *P4ha2* WT and KO cells

(**Suppl. Fig. 5I**), collagen deposition in *P4ha2* KO/KD cells may not be significantly impacted.

Given the role of P4HA2 in promoting secretion and deposition of collagen and cell invasion in

breast cancer (47, 48), the seemly contradictory findings on the differential effect of *P4ha2*

KO/KD and collagen deposition on cell migration may warrant further studies. Furthermore, due

to the lack of an antibody that can specifically recognize the hydroxylated proline 174 of YAP1,

we cannot assess the clinical relevance of prolyl hydroxylation of YAP1 in tumor samples from

prostate cancer patients. The development of such antibodies is warranted and would allow us to

examine the association of prolyl hydroxylation of YAP1 and P4HA2 expression in prostate

cancer specimens.

In summary, our findings support a model in which P4HA2-mediated prolyl

hydroxylation serves as a molecular switch that controls the activities of YAP1 in cell migration,

invasion, and metastasis (**Fig. 4G**).

MATERIALS AND METHODS

The reagents and assays as well as bioinformatic/statistical analyses were described in

Supplementary Information.

ACKNOWLEDGEMENTS

We thank Ronald DePinho and Haoqiang Ying for helpful discussions, Sarah E. Townsend for

editing, and Dr. Timothy C. Thompson for providing RM-1 cells. G.W. is supported by the

funding from MDACC (Moon Shot, IRG, and PCRP), UT STARs Award, NIH R00 CA194289

and P50 CA140388 as well as NIH P30CA016672 for the use of Research Animal Support

Facility, Flow Cytometry and Cellular Imaging Core Facility, and Functional Genomics Core.

CONTRIBUTIONS

MZ and GW contributed to the study's conception and design of this study. GW, MZ, RP, XL, ZL, MT, PH, JHS, CSM, JP, SZ, AH, XM, RC, QC, AKJ, and HK performed the experiments and acquired, analysed, and interpreted the data (e.g., statistical analysis, biostatistics, computational analysis). CJL, SMH, and GW contributed to the supervision of the project. SHL and GW contributed to the writing and editing the manuscript.

REFERENCES

1.   Calses PC, Crawford JJ, Lill JR, Dey A. Hippo Pathway in Cancer: Aberrant Regulation and Therapeutic Opportunities. Trends Cancer. 2019;5(5):297-307.

2.   Ehsanian R, Brown M, Lu H, Yang XP, Pattatheyil A, Yan B, et al. YAP dysregulation by phosphorylation or DeltaNp63-mediated gene repression promotes proliferation, survival and migration in head and neck cancer subsets. Oncogene. 2010;29(46):6160-71.

3.   Yuan M, Tomlinson V, Lara R, Holliday D, Chelala C, Harada T, et al. Yes-associated protein (YAP) functions as a tumor suppressor in breast. Cell Death Differ. 2008;15(11):1752-9.

4.   Yu SJ, Hu JY, Kuang XY, Luo JM, Hou YF, Di GH, et al. MicroRNA-200a promotes anoikis resistance and metastasis by targeting YAP1 in human breast cancer. Clinical cancer research : an official journal of the American Association for Cancer Research. 2013;19(6):1389-99.

5.   Matallanas D, Romano D, Yee K, Meissl K, Kucerova L, Piazzolla D, et al. RASSF1A elicits apoptosis through an MST2 pathway directing proapoptotic transcription by the p73 tumor suppressor protein. Mol Cell. 2007;27(6):962-75.

6.   Cottini F, Hideshima T, Xu C, Sattler M, Dori M, Agnelli L, et al. Rescue of Hippo

coactivator YAP1 triggers DNA damage-induced apoptosis in hematological cancers. Nature

medicine. 2014;20(6):599-606.

7.   Levy D, Adamovich Y, Reuven N, Shaul Y. The Yes-associated protein 1 stabilizes p73

by preventing Itch-mediated ubiquitination of p73. Cell Death Differ. 2007;14(4):743-51.

8.   Cheung P, Xiol J, Dill MT, Yuan WC, Panero R, Roper J, et al. Regenerative

Reprogramming of the Intestinal Stem Cell State via Hippo Signaling Suppresses Metastatic

Colorectal Cancer. Cell Stem Cell. 2020;27(4):590-604 e9.

9.   Ou C, Sun Z, Li S, Li G, Li X, Ma J. Dual roles of yes-associated protein (YAP) in

colorectal cancer. Oncotarget. 2017;8(43):75727-41.

10.   Cheng S, Prieto-Dominguez N, Yang S, Connelly ZM, StPierre S, Rushing B, et al. The

expression of YAP1 is increased in high-grade prostatic adenocarcinoma but is reduced in

neuroendocrine prostate cancer. Prostate Cancer Prostatic Dis. 2020.

11.   Nguyen LT, Tretiakova MS, Silvis MR, Lucas J, Klezovitch O, Coleman I, et al. ERG

Activates the YAP1 Transcriptional Program and Induces the Development of Age-Related

Prostate Tumors. Cancer cell. 2015;27(6):797-808.

12.   Wang G, Lu X, Dey P, Deng P, Wu CC, Jiang S, et al. Targeting YAP-Dependent MDSC

Infiltration Impairs Tumor Progression. Cancer Discov. 2016;6(1):80-95.

13.   Cheng S, Prieto-Dominguez N, Yang S, Connelly ZM, StPierre S, Rushing B, et al. The

expression of YAP1 is increased in high-grade prostatic adenocarcinoma but is reduced in

neuroendocrine prostate cancer. Prostate Cancer Prostatic Dis. 2020;23(4):661-9.

14.   Zhang L, Yang S, Chen X, Stauffer S, Yu F, Lele SM, et al. The hippo pathway effector

YAP regulates motility, invasion, and castration-resistant growth of prostate cancer cells.

Molecular and cellular biology. 2015;35(8):1350-62.

15.   Kuser-Abali G, Alptekin A, Lewis M, Garraway IP, Cinar B. YAP1 and AR interactions

contribute to the switch from androgen-dependent to castration-resistant growth in prostate

cancer. Nat Commun. 2015;6:8126.

16.   Chen YA, Lu CY, Cheng TY, Pan SH, Chen HF, Chang NS. WW Domain-Containing

Proteins YAP and TAZ in the Hippo Pathway as Key Regulators in Stemness Maintenance,

Tissue Homeostasis, and Tumorigenesis. Front Oncol. 2019;9:60.

17.   Yan F, Qian M, He Q, Zhu H, Yang B. The posttranslational modifications of Hippo-

YAP pathway in cancer. Biochim Biophys Acta Gen Subj. 2020;1864(1):129397.

18.   Zurlo G, Guo J, Takada M, Wei W, Zhang Q. New Insights into Protein Hydroxylation

and Its Important Role in Human Diseases. Biochim Biophys Acta. 2016;1866(2):208-20.

19.   Gorres KL, Raines RT. Prolyl 4-hydroxylase. Crit Rev Biochem Mol Biol.

2010;45(2):106-24.

20.   Su W, Han HH, Wang Y, Zhang B, Zhou B, Cheng Y, et al. The Polycomb Repressor

Complex 1 Drives Double-Negative Prostate Cancer Metastasis by Coordinating Stemness and

Immune Suppression. Cancer cell. 2019;36(2):139-55 e10.

21.   Kapoor A, Yao W, Ying H, Hua S, Liewen A, Wang Q, et al. Yap1 activation enables

bypass of oncogenic Kras addiction in pancreatic cancer. Cell. 2014;158(1):185-97.

22.   Shao DD, Xue W, Krall EB, Bhutkar A, Piccioni F, Wang X, et al. KRAS and YAP1

converge to regulate EMT and tumor survival. Cell. 2014;158(1):171-84.

23. Hiemer SE, Szymaniak AD, Varelas X. The transcriptional regulators TAZ and YAP

direct transforming growth factor beta-induced tumorigenic phenotypes in breast cancer cells. J

Biol Chem. 2014;289(19):13461-74.

24. Kim T, Yang SJ, Hwang D, Song J, Kim M, Kyum Kim S, et al. A basal-like breast

cancer-specific role for SRF-IL6 in YAP-induced cancer stemness. Nat Commun. 2015;6:10186.

25. Isfort I, Cyra M, Elges S, Kailayangiri S, Altvater B, Rossig C, et al. SS18-SSX-

Dependent YAP/TAZ Signaling in Synovial Sarcoma. Clinical cancer research : an official

journal of the American Association for Cancer Research. 2019;25(12):3718-31.

26. Ying H, Kimmelman AC, Lyssiotis CA, Hua S, Chu GC, Fletcher-Sananikone E, et al.

Oncogenic Kras maintains pancreatic tumors through regulation of anabolic glucose metabolism.

Cell. 2012;149(3):656-70.

27. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al.

Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide

expression profiles. Proceedings of the National Academy of Sciences of the United States of

America. 2005;102(43):15545-50.

28. Zanconato F, Forcato M, Battilana G, Azzolin L, Quaranta E, Bodega B, et al. Genome-

wide association between YAP/TAZ/TEAD and AP-1 at enhancers drives oncogenic growth. Nat

Cell Biol. 2015;17(9):1218-27.

29. Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, et al. Simple combinations of

lineage-determining transcription factors prime cis-regulatory elements required for macrophage

and B cell identities. Mol Cell. 2010;38(4):576-89.

30. Li S, Wan C, Zheng R, Fan J, Dong X, Meyer CA, et al. Cistrome-GO: a web server for

functional enrichment analysis of transcription factor ChIP-seq peaks. Nucleic Acids Res.

2019;47(W1):W206-W11.

31. Zhao B, Wei X, Li W, Udan RS, Yang Q, Kim J, et al. Inactivation of YAP oncoprotein

by the Hippo pathway is involved in cell contact inhibition and tissue growth control. Genes &

development. 2007;21(21):2747-61.

32. Becker JS, Nicetto D, Zaret KS. H3K9me3-Dependent Heterochromatin: Barrier to Cell

Fate Changes. Trends Genet. 2016;32(1):29-41.

33. Pekowska A, Benoukraf T, Zacarias-Cabeza J, Belhocine M, Koch F, Holota H, et al.

H3K4 tri-methylation provides an epigenetic signature of active enhancers. EMBO J.

2011;30(20):4198-210.

34. Zhang B, Chambers KJ, Faller DV, Wang S. Reprogramming of the SWI/SNF complex

for co-activation or co-repression in prohibitin-mediated estrogen receptor regulation. Oncogene.

2007;26(50):7153-7.

35. Nagl NG, Jr., Wang X, Patsialou A, Van Scoy M, Moran E. Distinct mammalian

SWI/SNF chromatin remodeling complexes with opposing roles in cell-cycle control. EMBO J.

2007;26(3):752-63.

36. Underhill C, Qutob MS, Yee SP, Torchia J. A novel nuclear receptor corepressor

complex, N-CoR, contains components of the mammalian SWI/SNF complex and the

corepressor KAP-1. J Biol Chem. 2000;275(51):40463-70.

37. Hoxha S, Shepard A, Troutman S, Diao H, Doherty JR, Janiszewska M, et al. YAP-

Mediated Recruitment of YY1 and EZH2 Represses Transcription of Key Cell-Cycle Regulators.

Cancer research. 2020;80(12):2512-22.

38.   Kim M, Kim T, Johnson RL, Lim DS. Transcriptional co-repressor function of the hippo

pathway transducers YAP and TAZ. Cell Rep. 2015;11(2):270-82.

39.   Zhao B, Kim J, Ye X, Lai ZC, Guan KL. Both TEAD-binding and WW domains are

required for the growth stimulation and oncogenic transformation activity of yes-associated

protein. Cancer research. 2009;69(3):1089-98.

40.   Ma S, Meng Z, Chen R, Guan KL. The Hippo Pathway: Biology and Pathophysiology.

Annu Rev Biochem. 2019;88:577-604.

41.   Chakravarthi BV, Pathi SS, Goswami MT, Cieslik M, Zheng H, Nallasivam S, et al. The

miR-124-prolyl hydroxylase P4HA1-MMP1 axis plays a critical role in prostate cancer

progression. Oncotarget. 2014;5(16):6654-69.

42.   van Huizen NA, Burgers PC, Saintmont F, Brocorens P, Gerbaux P, Stingl C, et al.

Identification of 4-Hydroxyproline at the Xaa Position in Collagen by Mass Spectrometry. J

Proteome Res. 2019;18(5):2045-51.

43.   Discher DE, Smith L, Cho S, Colasurdo M, Garcia AJ, Safran S. Matrix

Mechanosensing: From Scaling Concepts in 'Omics Data to Mechanisms in the Nucleus,

Regeneration, and Cancer. Annu Rev Biophys. 2017;46:295-315.

44.   Ozdemir BC, Pentcheva-Hoang T, Carstens JL, Zheng X, Wu CC, Simpson TR, et al.

Depletion of carcinoma-associated fibroblasts and fibrosis induces immunosuppression and

accelerates pancreas cancer with reduced survival. Cancer cell. 2014;25(6):719-34.

45.   Holster T, Pakkanen O, Soininen R, Sormunen R, Nokelainen M, Kivirikko KI, et al.

Loss of assembly of the main basement membrane collagen, type IV, but not fibril-forming

collagens and embryonic death in collagen prolyl 4-hydroxylase I null mice. J Biol Chem.

2007;282(4):2512-9.

46.   Aro E, Salo AM, Khatri R, Finnila M, Miinalainen I, Sormunen R, et al. Severe

Extracellular Matrix Abnormalities and Chondrodysplasia in Mice Lacking Collagen Prolyl 4-

Hydroxylase Isoenzyme II in Combination with a Reduced Amount of Isoenzyme I. J Biol

Chem. 2015;290(27):16964-78.

47.   Xiong G, Deng L, Zhu J, Rychahou PG, Xu R. Prolyl-4-hydroxylase alpha subunit 2

promotes breast cancer progression and metastasis by regulating collagen deposition. BMC

Cancer. 2014;14:1.

48.   Gilkes DM, Chaturvedi P, Bajpai S, Wong CC, Wei H, Pitcairn S, et al. Collagen prolyl

hydroxylases are essential for breast cancer metastasis. Cancer research. 2013;73(11):3285-96.

FIGURE LEGENDS

Figure 1. YAP1 suppresses cell migration, invasion, and metastasis. (**A-B**) Cell migration and invasion assay using PS cells transduced with control shRNA and *Yap1* shRNAs. The *Yap1* KD efficiency was confirmed by WB analysis. (**C**) Cell migration assay using *Yap1*-WT PS cells and *Yap1*-KO cells. WB analysis confirmed the KO of YAP1 expression. (**D**) Cell migration assay using *Yap1*-KO cells with GFP overexpression and YAP1 overexpression. WB analysis confirmed the overexpression of YAP1 in *Yap1*-KO PS cells. (**E-F**) Cell migration and invasion assay using C4-2b cells transduced with control shRNA and *YAP1* shRNAs. The *YAP1* KD efficiency was confirmed by WB analysis. (**G-I**) Cell migration assay using IGR-CaP1 (G), iKPC (H), and MDA-MB-231 (I) with *YAP1* KD or KO compared to control cells. (**J**) Luciferase imaging in mice injected with PS cells transduced with control shRNA and *Yap1* shRNAs through the tail vein.

Figure 2. Microarray, ChIP-seq, and immunoprecipitation-mass spectrometry analyses. (**A**) GSEA analysis of microarray data from PS cells transduced with doxycycline-inducible *Yap1* shRNA identified EMT as the top pathway activated in *Yap1*-KD cells. (**B-C**) ChIP-seq

analysis identified YAP1 binding sites and YAP1-regulated pathways. (**D**) Immunoprecipitation-mass spectrometry analysis identified known YAP1-interacting proteins and novel YAP1-interacting proteins. (**E**) Exogenous YAP1 interacts with exogenous P4HA2 when overexpressed in 293T cells by transfection of the indicated plasmids for co-immunoprecipitation experiments. (**F**) Endogenous YAP1 interacts with endogenous P4HA2 in PS cells. (**F**) Multiple prolyl hydroxylation sites were identified in peptides of YAP1 isoform 3 from the LC- MS/MS analysis.

Figure 3. Prolyl hydroxylation of YAP1 suppressed cell migration, invasion, and metastases. (**A**) Scheme showing the strategy to generate prolyl hydroxylation–defective YAP1 mutants by mutating proline to alanine (PA): Mut1-3 (P75/85/120A), Mut4-5 (P172/174A), and Mut6-9 (P348/352/394/397). Human YAP1 isoform 3 was used. (**B**) Expression of YAP1 WT and PA mutants in *Yap1*-KO PS cells. (**C-D**) Cell migration and invasion assay in *Yap1*-KO PS cells with overexpression of YAP1 WT and PA mutants. (**E**) Tail vein injection of *Yap1*-KO cells with overexpression of YAP1 WT and PA mutants. (**F**) qPCR analysis of YAP1 target genes in *Yap1*-KO PS cells with overexpression of YAP1 WT, Mut4-5, and the constitutively active S127A mutant. (**G-H**) Cell migration and invasion assay using *Yap1*-KO PS cells with overexpression of GFP, YAP1 WT, Mut4-5, Mut4, and Mut5. (**I**) ChIP-qPCR analysis of YAP1, H3K4me3, and H3K9me4 binding sites in *Col12a1* and *Mgp*.

Figure 4. P4HA2 suppresses cell migration and invasion through Yap1. (**A**) WB analysis of P4HA2 in *P4ha2*-WT and *P4ha2*-KO PS cells. (**B-C**) Cell migration and invasion using *P4ha2*-WT and *P4ha2*-KO PS cells. (**D**) qPCR analysis of YAP1 target genes (*Postn, Col12a1, and Mgp*) in *P4ha2* KO and WT cells. (**E**) WB analysis of YAP1 in *P4ha2*-WT and *P4ha2*-KO PS

cells transduced with Yap1 shRNAs. (**F**) Cell invasion assay in *P4ha2*-WT and *P4ha2*-KO PS

cells transduced with shYap1#434. (**G**) A model for hydroxylation-dependent YAP1 function in

cell migration, invasion, and metastasis (Created with BioRender.com). Left: P4HA2-mediated

hydroxylation of YAP1 may impair its interactions with transcription factors such as JUN or

enhance the recruitment of corepressor, such as SWI/SNF-NCoR1, NuRD, and EZH2/YY1,

which results in a decrease in the expression of genes involved in cell migration, invasion, and

metastasis. Right: In the absence of P4HA2, non-hydroxylated YAP1 may efficiently interact

with transcription factors such as JUN to activate genes involved in cell migration, invasion, and

metastasis.

A

Enrichment plot:
HALLMARK_EPITHELIAL_MESENCHYMAL_TRANSIT
ION

NES=-1.45
p=0.0

shCon   shYap1

B

- Promoter (<=1kb)
- Promoter (1−2kb)
- Promoter (2−3kb) ()
- 5' UTR
- 3' UTR
- 1st Exon
- Other Exon
- 1st Intron
- Other Intron
- Downstream (<=3kb)
- Distal Intergenic

C

positive regulation of locomotion
positive regulation of cellular component movement
angiogenesis
positive regulation of cell motility
positive regulation of cell migration
negative regulation of intracellular signal transduction
reproductive system development
gland development
reproductive structure development
tube morphogenesis

Count
● 60
● 70
● 80

p.adjust
1.0e-10
7.5e-11
5.0e-11
2.5e-11

D

Pulldown peptide #

	IgG	IP: Yap1	
YAP1	0	133	
AMOT	2	75	
AMOTL2	0	58	
AMOTL1	0	27	Known
LATS1	0	11	Yap1
YWHAG	1	24	interactors
YWHAB	0	20	
YWHAH	0	19	
YWHAZ	0	19	
YWHAE	2	14	
P4HA2	1	53	P4H
P4HB	4	31	complex
P4HA1	0	13	
SMARCC1	0	32	SWI/SNF
SMARCA4	1	26	complex
SMARCE1	0	11	

E

293T

Flag-YAP1   +   +   +   +
V5-P4HA2   -   +   -   +

Input   IP: V5

WB:Flag

WB: V5

F

PS cells (endogenous)

IP:
Input   IgG   YAP1

WB: P4HA2

WB: YAP1

G

YAP1 mouse isoform 1 (NP_001164618)

P60 P70   P105   P157 P159   P347 P353   P395 P398

Proline-rich   TEAD binding   WW   WW   Transactivation domain

Peptides identified from LC-MS/MS
Proline 60: GDSETDLEALFNAVMN**p**K
Proline 70: TANV**p**QTVPMRLR
Proline 105: QASTDAGTAGALT**p**QHVR
Proline 157/159: QSSFEIPDDV**p**L**p**AGWEMAK
Proline 347/353: SQLPTLEQDGGT**p**NAVSS**p**GMSQELR
Proline 395: DESTDSGLSMSSYSI**p**R
Proline 398: T**p**DDFLNSVDEMDTGDTISQSTLPSQQSR

Figure 3