NOMIS: Quantifying morphometric deviation from normality over the lifetime in the adult human brain

Olivier Potvin[1], PhD, Louis Dieumegarde[1], BSc, and Simon Duchesne, PhD[12] for the Alzheimer's Disease Neuroimaging Initiative* and CIMA-Q**

[1] CERVO Brain Research Centre, 2601, de la Canardière, Québec, Canada, G1J 2G3

[2] Département de radiologie, Faculté de médecine, Université Laval, 1050, avenue de la Médecine, Québec, Canada, G1V 0A6

Key words: magnetic resonance imaging, atrophy, morphometry, normality, aging, sex.

Correspondence:

Simon Duchesne, PhD
CERVO Brain Research Centre
F-3582, 2601, de la Canardière, Québec, Canada, G1J 2G3
Phone: 418 663-5000 ext.4777
Fax: 418 663-9540
simon.duchesne@fmed.ulaval.ca

*Part of the data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

** Part of the data used in this article were obtained from the Consortium pour l'identification précoce de la maladie Alzheimer - Québec (CIMA-Q; cima-q.ca). As such, the investigators within the CIMA-Q contributed to the design, the implementation, the acquisition of clinical, cognitive, and neuroimaging data and biological samples. A list of the CIMA-Q investigators is available on www.cima-q.ca.

Abstract

We present NOMIS (https://github.com/medicslab/NOMIS), a comprehensive open MRI tool to assess morphometric deviation from normality in the adult human brain. Based on MR anatomical images from 6,909 cognitively healthy individuals aged 18-100 years, we modeled 1,344 measures computed using the open access *FreeSurfer* pipeline, taking into account personal characteristics (age, sex, head size), scanner characteristics (manufacturer and magnetic field strength), and image quality, providing expected values for any new individual. Then, for each measure, the NOMIS tool was built to generate Z-score effect sizes denoting the extent of deviation from the normative sample. Depending on the user need, NOMIS offers four versions of Z-score adjusted on different sets of variables. While all versions take into account head size, image quality and scanner characteristics, they can also incorporate age and/or sex, thereby facilitating multi-site neuromorphometric research across adulthood.

Introduction

Despite the popularity of magnetic resonance imaging (MRI) to examine abnormalities in brain

morphometry, tools quantifying normality are lacking. While age, sex and intracranial volume are

well-known to influence brain volume and shape[1, 2] the determination of whether an

individual's brain region measurements are within normality faces multiple major challenges

such as the lack of normative data across appropriate age groups, the influence of the MRI

processing pipeline, the variety in neuroanatomical atlases used for parcellation and the quality

of the image itself[3, 4]. We made previous attempts[5-8] to produce such normative data in

adulthood based on *FreeSurfer*, an open-access and fully automated segmentation software

(http://freesurfer.net), for two specific brain atlases, namely Desikan-Killiany[9] (DK) and

Desikan-Killiany-Tourville[10] (DKT). This initial foray allowed for the quantification of the extent

of deviation from normality for a given individual, according to personal characteristics such as

age, sex and estimated intracranial volume (eTIV), while controlling for scanner magnetic field

strength (MFS) and scanner manufacturer (OEM). Although this work has already gathered more

than a hundred citations in the last three years, several researchers solicited the expansion of

the norms to include other brain regions, as well as different atlases; as well as the production

of variants, for example, of only adjusting for head size and scanner characteristics. For the latter,

it became clear to us that we also needed to control for image quality. Further, we recognized

the need to increase the size of our normative sample to ensure better representation of middle

age.

Leveraging this prior work, we offer a comprehensive tool called NOMIS (NOrmative

Morphometry Image Statistics; https://github.com/medicslab/NOMIS). NOMIS can be used for

new adult individuals, healthy or otherwise. Using this individual's T1-weighted MRI, processed

via the *FreeSurfer* 6.0 toolkit, one can derive Z-score effect sizes denoting the extent of deviation

from the normative sample according to the individual's characteristics (age, sex, and eTIV), while

taking into account scanner information (MFS, OEM) and now voxel size (resolution) and image

quality (contrast-to-noise ratio (CNR) and holes in surface reconstruction). Our model takes into

account 1,344 brain measures generated by *FreeSurfer* on 6,909 healthy individuals aged 18 to

100 years (mean ± sd: 55.0 ± 20.0; 56.8% female). The normative data includes as before the

DK[9] and DKT[10] atlases, as well as the Destrieux (a2009s)[11] neocortical atlas; neocortical pial

and white surface areas, volumes and thicknesses; *FreeSurfer*'s default subcortical atlas[12],

hippocampal subfields, brainstem subregions; its ex vivo-based labeling protocol atlas[13]; and

the subcortical white matter parcellation according to the adjacent neocortical areas.

Furthermore, to fulfill specific needs from researchers, we propose four versions of Z-score

adjusted on different sets of variables. While all versions are adjusted for head size, image quality

and scanner characteristics, the full version includes both age and sex whereas the three other

versions are without age, without sex and without age and sex. Thus, a research group working

on aging aiming at removing the variance of hippocampal volumes due to head size, sex, scanner,

and image quality could use the version without age, which preserves the variance due to aging.

Materials and methods

Normative sample

The norms are based on a cross-sectional sample of 6,909 (initial sample: 7,399) cognitively

healthy individuals aged 18 to 100 years, (mean ±sd; 55.0 ±20.0; 56.8% female), gathered from

27 different datasets (Table 1; Acknowledgments). Scans were acquired from one of the three

leading OEM (e.g. Siemens Healthcare (Erlangen, Germany); Philips Medical Systems (Best,

Netherlands); or GE Healthcare (Milwaukee, WI)) at MFS of either 1.5 or 3 Tesla. For each dataset,

approval from the local ethics board and informed consent of the participants were obtained.

Table 1. Datasets included in the normative sample

Dataset	n
Autism Brain Imaging Data Exchange (ABIDE)[14]	183
Alzheimer's Disease Neuroimaging Initiative (ADNI)[15]	672
Australian Imaging Biomarkers and Lifestyle flagship study of ageing (AIBL)[16]	157
Berlin Mind and Brain (Margulies, Villringer) CoRR sample (BMB)[17, 18]	50
Cambridge Centre for Ageing and Neuroscience (CamCAN)[19, 20]	630
Center of Biomedical Research Excellence (COBRE)[21]	70
Cleveland CCF[22]	30
Consortium for the Early Identification of Alzheimer's Disease (CIMA-Q)[23]	29
Dallas Lifespan Brain Study (DLBS)[24]	304
FIND lab sample (FIND) Lab[25]	13
Functional Biomedical Informatics Research Network (FBIRN)[26]	33
Lifespan Human Connectome Project in Aging (HCP-Aging)[27]	612
International Consortium for Brain Mapping (ICBM) - MNI[28]	147
Information eXtraction from Images (IXI)[29]	554
F.M. Kirby Research Center neuroimaging reproducibility data (KIRBY-21)[30]	20
Minimal Interval Resonance Imaging in Alzheimer's Disease (MIRIAD)[31]	21
National Alzheimer's Coordinating Center (NACC)[32]	1562
National Database for Autism Research (NDAR)[33]	56
Nathan Kline Institute Rockland (NKI-RS)[34]	138
Nathan Kline Institute Rockland (NKI-RS) Enhanced[34]	436
Open Access Series of Imaging Studies (OASIS)[35]	288
POWER Neuroimage sample (POWER)[36]	26
Parkinson's Progression Markers Initiative (PPMI)[37]	158
Southwest University Adult Lifespan Dataset (SALD)[38]	490
University of Wisconsin (Birn, Prabhakaran, Meyerand) CoRR sample (UWM)[17]	25
Wayne State EF Dataset[39]	108
Yale Low-Resolution Controls Dataset[40]	97
Total	6909

Among the datasets are the Alzheimer's Disease Neuroimaging Initiative (ADNI), the Australian

Imaging, Biomarkers and Lifestyle study of aging (AIBL) and the Consortium pour l'identification

précoce de la maladie Alzheimer - Québec (CIMA-Q) datasets. The ADNI (adni.loni.usc.edu) was

launched in 2003 as a public-private partnership, led by Principal Investigator Michael W. Weiner,

MD. (www.adni-info.org). The AIBL data was collected by the AIBL study group and AIBL study

methodology has been reported previously by Ellis et al. (2009). For each dataset, approval from

the local ethics board and informed consent of the participants were obtained. Founded in 2013

with a \$2,500,000 grant from the Fonds d'Innovation Pfizer - Fond de Recherche Québec – Santé

sur la maladie d'Alzheimer et les maladies apparentées, the main objective of CIMA-Q is to build

a cohort of participants characterized in terms of cognition, neuroimaging and clinical outcomes

in order to acquire biological samples allowing (1) to establish early diagnoses of Alzheimer's

disease, (2) to provide a well characterized cohort and (3) to identify new therapeutic targets.

The principal investigator and director of CIMA-Q is Dr Sylvie Belleville from the Centre de

recherche de l'Institut universitaire de gériatrie de Montréal, CIUSSS Centre-sud-de-l'Île-de-

Montréal. CIMA-Q represent a common effort of several researchers from Québec affiliated to

Université Laval, Université McGill, Université de Montréal, et Université de Sherbrooke. CIMA-

Q recruited 350 cognitively healthy participants, with subjective cognitive impairment, mild

cognitive impairment, or Alzheimer's disease, between 2013–2016. Volunteers were recruited

from memory clinics, through advertisements posted in the community and amongst participants

from the NuAge population study.

From all the samples mentioned, only cognitively healthy (control) participants were

included. For the Nathan Kline Institute samples, which were projects recruiting in the general

population, we excluded participants with history of schizophrenia or other psychotic disorders,

bipolar disorders, major depressive disorders (recurrent), posttraumatic stress disorder,

substance abuse/dependence disorders, neurodegenerative and neurological disorders, head

injury with loss of consciousness/amnesia, and lead poisoning. Moreover, for the Parkinson's

Progression Markers Initiative dataset, we excluded participants with a Geriatric Depression

Scale[41] score of more than 5.

Brain segmentation

Brain segmentation was conducted using *FreeSurfer* version 6.0, a widely used and freely

available automated processing pipeline that quantifies brain anatomy (http://freesurfer.net).

All raw T1-weighted images were processed using the "recon-all -all" *FreeSurfer* command with

the fully-automated directive parameters (no manual intervention or expert flag options) on the

CBRAIN platform[42]. Normative data were computed for volumes, neocortical thicknesses and

white and pial surfaces areas for all atlases comprised in *FreeSurfer* 6.0: the default subcortical

atlas[12] (aseg.stats), the Desikan-Killiany atlas[9] (DK, aparc.stats file), the Desikan-Killiany-

Tourville atlas[10] (DKT, aparc.DKT.stats file), the Destrieux atlas[11] (aparc.a2009s.stats file), the

ex vivo atlas,[43] including entorhinal and perirhinal cortices, the brainstem sub-regions

atlas[44], the Brodmann area maps which includes somatosensory areas, several motor and

visual areas, as well as the hippocampal subfields atlas[45].

The technical details of *FreeSurfer*'s procedures are described in prior publications.

Briefly, this processing includes motion correction, removal of non-brain tissue using a hybrid

watershed/surface deformation procedure, automated Talairach transformation, intensity

normalization, tessellation of the gray matter white matter boundary, automated topology

correction, and surface deformation following intensity gradients to optimally place the

gray/white and gray/cerebrospinal fluid borders at the location where the greatest shift in

intensity defines the transition to the other tissue class. Once the cortical models are complete,

a number of deformable procedures can be performed for further data processing and analysis

including surface inflation, registration to a spherical atlas which is based on individual cortical

folding patterns to match cortical geometry across subjects and parcellation of the cerebral

cortex into units with respect to gyral and sulcal structure. This method uses both intensity and

continuity information from the entire three-dimensional MRI volume in segmentation and

deformation procedures to produce representations of cortical thickness, calculated as the

closest distance from the gray/white boundary to the gray/CSF boundary at each vertex on the

tessellated surface. The maps are created using spatial intensity gradients across tissue classes

and are therefore not simply reliant on absolute signal intensity. Procedures for the

measurement of cortical thickness have been validated against histological analysis [46] and

manual measurements[47, 48]. Estimated intracranial volume[49] was taken from the aseg.stats

FreeSurfer output file. We added the total ventricle volume (labeled as "ventricles") using the

sum of all ventricles and the corpus callosum (labeled as "cc") using the sum of all corpus

callosum segments.

Quality control and sample selection

A flow chart detailing the final analysis sample is shown in Fig 1. From an initial pool of 7,399

MRIs, nine images failed the *FreeSurfer* pipeline. Following processing, each of the remaining

7,390 brain segmentations was visually inspected through at least 20 evenly distributed coronal

sections. After quality control, 445 images (6.0%) were removed from further analyses due to

segmentation problems, the main reason being that parts of the brain were not completely

segmented (e.g. temporal and occipital poles). During visual inspection, 26 brains were found to

have clear significant brain lesions and were excluded. In addition to visual inspection, we

excluded participants if at least one of the 1,344 brain region measures was missing (n=10). In

fine, the analysis sample was composed of 6,909 individual MRIs.

Fig 1. Flow chart of the images.

Training, validation and test sample

We randomly selected 10% of the whole sample (n=691) to test the models in an

independent sample (age: 55.1 ±20.1, range 18-100; 58.5% female). This test sample was not

used to build the models predicting normative values. The remaining 90% was used as training

sample (age: 54.9 ±20.0, range 18-100; 56.7% female) to build and validate the models. Leave-

10%-out cross-validation was used to validate the model in the training sample.

Clinical samples evaluations

We evaluated the usefulness of normative values using clinical samples of individuals with

schizophrenia (n=72; Age: 38.2 ±13.9, range 18-65; 19% female) from the COBRE dataset, as well

as participants with clinically ascertained mild Alzheimer's disease (n=157 Age: 74.8 ±8.1, range

55-90; 43% female) from the baseline ADNI-2 dataset.

Scanner-related predictors

Scanner-related predictors included manufacturer (OEM), magnetic field strength (MFS), and

voxel size (resolution).

Image quality predictors

Image quality can have an effect on brain segmentation quality[50, 51]. We therefore included

in the prediction models contrast-to-noise ratio (CNR) for each area as well as the total number

of holes in surface reconstruction prior to fixing, since this measure is correlated with visual

assessment of brain segmentation[50]. Defect holes – topological errors in the cortical surface

reconstructions – were extracted from the aseg.stats *FreeSurfer* output file and CNR was assessed

after *FreeSurfer* preprocessing using the brain.mgz file with gray matter (GM) and cerebral white

matter (WM) voxel intensities for each area with the following formula:

194
$$CNR = \frac{(GM\ mean\ -\ WM\ mean\ )^2}{(GM\ variance\ +\ WM\ variance)}$$

Regression models and statistical analyses

Linear regression models predicting each measure were built using age, sex, eTIV, MFS, OEM,

voxel size, surface defect holes and the CNR of the region as predictors. To obtain normal

distributions, surface holes and ventricles (except the 4th) were log transformed. For ventricles

and white matter regions, CNR of the total brain gray matter was used while for the brainstem

subregions and hippocampal subfields, CNR from the whole brainstem and whole hippocampus

were used, respectively. Quadratic and cubic terms for age, eTIV, and surface holes were

included, as well as the following interactions: age X sex, age X eTIV, sex X eTIV, eTIV X MFS and

MFS X OEM. Feature selection was conducted with a 10-fold cross-validation[52] backward

elimination procedure, retaining the model with the subset of predictors that produced the

lowest predicted residual sum of squares. For each selected final model, the fit of the data was

assessed using R^2 coefficient of determination:

208
$$R^2 = 1 - \frac{\sum_i (Y_i - f_i)^2}{\sum_i (Y_i - \hat{Y})^2}$$

where the numerator is the residual sum of squares (Y is the value of the variable to be predicted

and f is the predicted value) and the denominator is the total sum of squares ($\hat{Y}$ is the mean). To

assess the unique contribution of each predictor, we used the lmg metric in the *R* package[53]

relaimpo[54]. This metric is a R^2 partitioned by averaging sequential sums of squares over all

orderings of the predictors.

For each brain measure, in order to exclude potential abnormalities, outliers with Z scores higher

than 3.29 ($p < .001$) were removed to compute the statistical model. This was done in proportion

to eTIV for volumes and surfaces and on raw values for cortical thicknesses. The number of

outliers was below 1% for all regions (mean ±sd of all atlases: 0.45% ±0.10%) except the right

long insular gyrus and central sulcus of the insula white surface (1.1%) and pericallosal sulcus

volume (1.1%) of the Destrieux atlas. Detailed results can be found in the supplementary

material. Brain figures were made using the ggseg R package[55].

The models were verified by examining the difference between CV-10 R^2 of the training sample

and R^2 of the independent test sample of healthy controls. We then examined the validity of the

normative values to show expected patterns of normality deviations using the Z score effect sizes

in the validation samples of healthy individuals and of individuals with AD and SZ.

225
$$Z_{OP} = \frac{Y_o - \hat{Y}}{RMSE}$$

226   Z score effect sizes (Z_{OP}) were obtained by subtracting the *Predicted value* ($\hat{Y}$) from the *Observed*

227   *value* (Y_o) divided by the root mean square error (RMSE) of the model predicting the value [56].

Results

Total variance explained by the models

The cross-validation 10-fold (CV-10) R^2 for the overall explanatory variance of the models

ranged between 0.02 to 0.84, with a mean $\pm$ sd of 0.37 $\pm$ 0.10. The highest R^2 were observed in

the largest regions (total brain volume 0.84, neocortex volume 0.82, pial surface 0.80, and white

surface 0.79). The inferior occipital gyrus and sulcus (0.02) and the anterior transverse collateral

sulcus thicknesses in the Destrieux atlas (0.02), as well as the DK and DKT left parahippocampal

thickness (0.02) had the lowest R^2. As examples, figures in this report display subcortical and DK

neocortical atlases results. Full detailed results for all atlases are provided as supplementary

information. Fig 2 illustrates the total R^2 for neocortical volumes and thicknesses of the DK atlas

parcellation, as well as subcortical volumes. As shown, a higher amount of variance was generally

explained for volumes compared to cortical thicknesses.

Independent test

The models predicting normative values were tested in an independent, healthy adults

randomly chosen 10% sample. Nearly all models showed equivalent or higher R^2 on the test set

than on the training set by CV-10 (difference for all atlases: 0.01 ±0.016). The lowest test

differences were in the Destrieux atlas where 37 measures out of 592 were below -0.05, the

worse being the fronto-marginal gyrus (of Wernicke) and sulcus (0.11), the superior occipital

gyrus (0.10) and the superior temporal sulcus (0.10) pial surface areas. Fig 2 displays the R^2

difference between training and test sets for the DK atlas. Few minimal worse test values were

observed for some cortical and subcortical volumes while all cortical thicknesses and most other

volumes had equivalent or better R^2 prediction values.

Fig 2. Top: R^2 from 10-fold cross-validation (10-fold CV) for cortical volumes and thicknesses

from the DK atlas and subcortical volumes. Bottom: Difference between R^2 from the training

set (10-fold CV) and the independent test set. Worse prediction from the test set are shown

in green while better prediction are grayed.

Variance explained by each predictor

Figs 3 and 4 show variances due to biological and MRI factors. As expected, effects differ

highly from one region to another, however, globally, age and eTIV had the largest effects on

volumes, while age was the essential factor on neocortical thickness. Scanner, as well as image

quality had statistically significant, but smaller effects on morphometric measures. Fig 5

illustrates the effect of age and sex on the four different NOMIS Z scores versions on the

independent healthy test set. While the age effect is clearly apparent on the sex and the version

without age and sex, it is null on the age and sex and age Z scores versions. The sex effect is also

affected, but is relatively small.

Fig 3. Effects of age, sex, and estimated total intra-cranial volume (eTIV) on cortical volumes

and thicknesses from the DK atlas and subcortical volumes.

Fig 4. Effects of scanner and image quality on cortical volumes and thicknesses from the DK

atlas and subcortical volumes.

Fig 5. Age and sex effects on the left cortical thickness across the four NOMIS Z scores

alternatives.

Clinical validation

We validated the normative values in individuals with clinically ascertained mild

Alzheimer's disease and schizophrenia, which showed expected patterns of deviations from

otherwise cognitively/behaviorally healthy individuals (Fig 6). In the Alzheimer's disease group,

deviations from normality covered the frontal, temporal and parietal cortices with enlarged

ventricles, but were especially more pronounced in the hippocampus and entorhinal cortex. In

schizophrenia, atrophy was widespread to nearly all of the cortex.

Fig 6. Mean normative Z scores on cortical volumes and thicknesses from the DK atlas and

subcortical volumes of participants with mild Alzheimer's disease and with schizophrenia.

Discussion

NOMIS strengths and limits

Prior normative data[5-7] were relatively limited in terms of atlases and sample size. With

nearly seven thousand participants and 1,344 brain measures, NOMIS offers a comprehensive

neuromorphometric normative tool based on a very large sample. In addition, an innovation of

NOMIS is its flexibility. Depending on the user need, it has four versions of Z-score adjusted on

different sets of variables. All versions include head size, image quality and scanner

characteristics, but can also take into account age and/or sex or without age and sex. Therefore,

research groups looking for traditional norms, as well as others wanting to lower the variance

due to head size and image quality/scanner characteristics while preserving age and/or sex

variances can take advantage of NOMIS. Another strength of NOMIS is that the normative values

were created on a various amalgam of cognitively healthy participants from multiple countries,

with data acquired from a wide variety of MRI scanners and image quality, maximizing its

generalizability. A novelty to prior existing normative data, is the addition of the image quality

impact on the morphometry measures. Figure 4 shows that its effect is not trivial on cortical

volume and thickness. Thus, our new normative data should help to remove some noise due to

image quality.

Despite these strengths, users should keep in mind that before using NOMIS, it is

mandatory to verify *FreeSurfer* segmentations and that while it will remove parts of variance due

to the scanner and image quality, it won't correct for segmentation errors or image artefacts.

Moreover, the normative sample, comprised essentially of research volunteers in academic-led

environments, was recruited using a non-probability sampling method and may not be

representative of the targeted population by the user.

Using NOMIS

The NOMIS tool is a user-friendly automated script in Python, freely accessible

(https://github.com/medicslab/NOMIS). Users only need to pre-process their images with

FreeSurfer 6.0 using automated directive parameters, then specify the individuals' characteristics

to the script, which will automatically compute Z-scores based on the *FreeSurfer* output. One can

choose the version of the Z-score by including in the csv file, only the variables that needs to be

adjusted and the script automatically selects the appropriate version of predictors. The predictive

models and all statistical parameters are provided along with the script. We anticipate that this

tool will be of broad interest to the neuroscientific community.

Financial Disclosure Statement

OP and LD are supported by grants from the Canadian Consortium on Neurodegeneration in Aging (CCNA) and the Canadian Institutes of Health Research (#IC119923) for which SD is a co-investigator and a primary investigator, respectively. CCNA is supported with funding from several partners including the Alzheimer Society of Canada, Sanofi, and Women's Brain Health Initiative.

Acknowledgments

This study comprises multiple samples of healthy individuals. We wish to thank all principal investigators who collected these datasets and agreed to let them accessible.

Autism Brain Imaging Data Exchange (ABIDE): Primary support for the work by Adriana Di Martino was provided by the NIMH (K23MH087770) and the Leon Levy Foundation. Primary support for the work by Michael P. Milham and the INDI team was provided by gifts from Joseph P. Healy and the Stavros Niarchos Foundation to the Child Mind Institute, as well as by an NIMH award to MPM (R03MH096321). http://fcon_1000.projects.nitrc.org/indi/abide/

Alzheimer's Disease Neuroimaging Initiative (ADNI): The investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-

content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf. ADNI was funded by the

Alzheimer's Disease Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01

AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI

is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and

Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer's

Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen;

Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli

Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company

Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.;Janssen Alzheimer Immunotherapy

Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development

LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx Research;

Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging;

Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes

of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector

contributions are facilitated by the Foundation for the National Institutes of Health

(www.fnih.org). The grantee organization is the Northern California Institute for Research and

Education, and the study is coordinated by the Alzheimer's Therapeutic Research Institute at

the University of Southern California. ADNI data are disseminated by the Laboratory for Neuro

Imaging at the University of Southern California.

http://adni.loni.usc.edu/

Australian Imaging Biomarkers and Lifestyle flagship study of ageing (AIBL): Part of the data

used in this study was obtained from the Australian Imaging Biomarkers and Lifestyle flagship

study of ageing (AIBL) funded by the Commonwealth Scientific and Industrial Research

Organisation (CSIRO) which was made available at the ADNI database

(www.loni.usc.edu/ADNI). The AIBL researchers contributed data but did not participate in

analysis or writing of this report. AIBL researchers are listed at www.aibl.csiro.au

Berlin Mind and Brain (Margulies, Villringer) CoRR sample (BMB). Zuo, X.N., et al. (2014). An

open science resource for establishing reliability and reproducibility in functional connectomics.

Scientific data, 1, 140049. doi: 10.1038/sdata.2014.49.

http://fcon_1000.projects.nitrc.org/indi/CoRR/html/bmb_1.html

Cambridge Centre for Ageing and Neuroscience (CamCAN): CamCAN funding was provided by the

UK Biotechnology and Biological Sciences Research Council (grant number BB/H008217/1),

together with support from the UK Medical Research Council and University of Cambridge, UK.

http://www.mrc-cbu.cam.ac.uk/datasets/camcan/

Center of Biomedical Research Excellence (COBRE): The imaging data and phenotypic

information was collected and shared by the Mind Research Network and the University of New

Mexico funded by a National Institute of Health COBRE: 1P20RR021938-01A2.

http://fcon_1000.projects.nitrc.org/indi/retro/cobre.html

Cleveland Clinic (Cleveland CCF): Funded by the National Multiple Sclerosis Society.

http://fcon_1000.projects.nitrc.org/indi/retro/ClevelandCCF.html

Consortium for the Early Identification of Alzheimer's Disease (CIMA-Q): Part of the data used in

this article were obtained from the Consortium pour l'identification précoce de la maladie

Alzheimer - Québec (CIMA-Q). As such, the investigators within the CIMA-Q contributed to the

design, the implementation, the acquisition of clinical, cognitive, and neuroimaging data and

biological samples. A list of the CIMA-Q investigators is available on cima-q.ca.

Dallas Lifespan Brain Study (DLBS): This study is supported by the Center for Vital Longevity, the

University of Texas at Dallas, the University of Texas Southwestern Medical Center, the National

Institutes of Health and Aging, AVID Radiopharmaceuticals, the Aging Mind Foundation and the

Alzheimer's Association. http://fcon_1000.projects.nitrc.org/indi/retro/dlbs.html

FIND lab sample. Funded by the Dana Foundation; John Douglas French Alzheimer's

Foundation; National Institutes of Health (AT005733, HD059205,HD057610, NS073498,

NS058899). http://fcon_1000.projects.nitrc.org/indi/retro/find_stanford.html

Functional Biomedical Informatics Research Network (FBIRN): Provided by the Biomedical

Informatics Research Network under the following support: U24-RR021992, by the National

Center for Research Resources at the National Institutes of Health, U.S.A.

http://www.birncommunity.org/resources/data/

Lifespan Human Connectome Project in Aging (HCP-Aging): HCP-Aging data were obtained

from the National Institute of Mental Health (NIMH) Data Archive (NDA). NDA is a collaborative

informatics system created by the National Institutes of Health to provide a national resource

to support and accelerate research in mental health. Dataset identifier:

http://dx.doi.org/10.15154/1520138. This manuscript reflects the views of the authors and may

not reflect the opinions or views of the NIH or of the Submitters submitting original data to

NDA. http://nda.nih.gov

International Consortium for Brain Mapping (ICBM). The ICBM (Principal Investigator: John

Mazziotta, MD, PhD) was funded was provided by the National Institute of Biomedical Imaging

and BioEngineering. ICBM is the result of efforts of co-investigators from UCLA, Montreal

Neurologic Institute, University of Texas at San Antonio, and the Institute of Medicine,

Juelich/Heinrich Heine University - Germany." https://ida.loni.usc.edu/login.jsp?project=ICBM

Information eXtraction from Images (IXI): Data collected as part of the project

EPSRC GR/S21533/02 - http://brain-development.org/ixi-dataset/

F.M. Kirby Research Center neuroimaging reproducibility data (KIRBY-21). Landman, B.A. et al.

"Multi-Parametric Neuroimaging Reproducibility: A 3T Resource Study", NeuroImage. (2010)

NIHMS/PMC:252138 doi:10.1016/j.neuroimage.2010.11.047

https://www.nitrc.org/projects/multimodal

Minimal Interval Resonance Imaging in Alzheimer's Disease (MIRIAD): The MIRIAD investigators

did not participate in analysis or writing of this report. The MIRIAD dataset is made available

through the support of the UK Alzheimer's Society (RF116). The original data collection was

funded through an unrestricted educational grant from GlaxoSmithKline (6GKC).

http://miriad.drc.ion.ucl.ac.uk

National Alzheimer's Coordinating Center (NACC): The NACC database is funded by NIA/NIH

Grant U01 AG016976. NACC data are contributed by the NIA-funded ADCs: P30 AG019610 (PI

Eric Reiman, MD), P30 AG013846 (PI Neil Kowall, MD), P30 AG062428-01 (PI James Leverenz,

MD) P50 AG008702 (PI Scott Small, MD), P50 AG025688 (PI Allan Levey, MD, PhD), P50

AG047266 (PI Todd Golde, MD, PhD), P30 AG010133 (PI Andrew Saykin, PsyD), P50 AG005146

(PI Marilyn Albert, PhD), P30 AG062421-01 (PI Bradley Hyman, MD, PhD), P30 AG062422-01 (PI

Ronald Petersen, MD, PhD), P50 AG005138 (PI Mary Sano, PhD), P30 AG008051 (PI Thomas

Wisniewski, MD), P30 AG013854 (PI Robert Vassar, PhD), P30 AG008017 (PI Jeffrey Kaye, MD),

P30 AG010161 (PI David Bennett, MD), P50 AG047366 (PI Victor Henderson, MD, MS), P30

AG010129 (PI Charles DeCarli, MD), P50 AG016573 (PI Frank LaFerla, PhD), P30 AG062429-01(PI

James Brewer, MD, PhD), P50 AG023501 (PI Bruce Miller, MD), P30 AG035982 (PI Russell

Swerdlow, MD), P30 AG028383 (PI Linda Van Eldik, PhD), P30 AG053760 (PI Henry Paulson, MD,

PhD), P30 AG010124 (PI John Trojanowski, MD, PhD), P50 AG005133 (PI Oscar Lopez, MD), P50

AG005142 (PI Helena Chui, MD), P30 AG012300 (PI Roger Rosenberg, MD), P30 AG049638 (PI

Suzanne Craft, PhD), P50 AG005136 (PI Thomas Grabowski, MD), P30 AG062715-01 (PI Sanjay

Asthana, MD, FRCP), P50 AG005681 (PI John Morris, MD), P50 AG047270 (PI Stephen

Strittmatter, MD, PhD). https://www.alz.washington.edu/

National Database for Autism Research (NDAR): Data were obtained from the National Institute

of Mental Health (NIMH) Data Archive (NDA). NDA is a collaborative informatics system created

by the National Institutes of Health to provide a national resource to support and accelerate

research in mental health. Dataset identifier: http://dx.doi.org/10.15154/1520138. This

manuscript reflects the views of the authors and may not reflect the opinions or views of the

NIH or of the Submitters submitting original data to NDA. http://nda.nih.gov

Nathan Kline Institute Rockland (NKI-R) sample (NKI-RS) and Enhanced Sample (NKI-RS):

Principal support for the enhanced NKI-RS project is provided by the NIMH BRAINS

R01MH094639-01. Funding for key personnel also provided in part by the New York State Office

of Mental Health and Research Foundation for Mental Hygiene. Funding for the decompression

and augmentation of administrative and phenotypic protocols provided by a grant from the

Child Mind Institute (1FDN2012-1). Additional personnel support provided by the Center for the

Developing Brain at the Child Mind Institute, as well as NIMH R01MH081218, R01MH083246,

and R21MH084126. Project support also provided by the NKI Center for Advanced Brain

Imaging (CABI), the Brain Research Foundation, the Stavros Niarchos Foundation and the NIH

P50 MH086385-S1 (NKI-RS). http://fcon_1000.projects.nitrc.org/indi/pro/nki.html

http://fcon_1000.projects.nitrc.org/indi/enhanced/

Open access series of imaging studies (OASIS): The OASIS project was funded by grants P50

AG05681, P01 AG03991, R01 AG021910, P50 MH071616, U24 RR021382, and R01 MH56584.

http://www.oasis-brains.org/

POWER: This database was supported by NIH R21NS061144 R01NS32979 R01HD057076

U54MH091657 K23DC006638 P50 MH71616 P60 DK020579-31 , McDonnell Foundation

Collaborative Action Award, NSF IGERT DGE-0548890, Simon's Foundation Autism Research

Initiative grant, Burroughs Wellcome Fund, Charles A. Dana Foundation, Brooks Family Fund,

Tourette Syndrome Association, Barnes-Jewish Hospital Foundation, McDonnell Center for

Systems Neuroscience, Alvin J. Siteman Cancer Center, American Hearing Research Foundation

grant, Diabetes Research and Training Center at Washington University grant.

http://fcon_1000.projects.nitrc.org/indi/retro/Power2012.html

Parkinson's Progression Markers Initiative (PPMI): PPMI – a public-private partnership – is

funded by the Michael J. Fox Foundation for Parkinson's Research and funding partners,

including Abbvie, Allergan, Amathus, Avid Radiopharmaceuticals, Biogen Idec, BioLegend,

Bristol-Myers, Celgene, Cenali, Covance, GE Healthcare, Genentech, GlaxoSmithKline, Glolub

Capital, Handl Therapeutics, Insitro, Janssen Neuroscience, Eli Lilly and Company, Lundbeck,

Merck, Meso Scale Discovery, Neurocrine, Pfizer, Piramal, Prevail, Roche, Sanofi Genzyme,

Servier, Takeda, Teva, UCB, Verily, and Voyager Therapeutics. See http://www.ppmi-info.org for

further details.

Southwest University Adult Lifespan Dataset (SALD): SALD was supported by the National

Natural Science Foundation of China (31470981; 31571137; 31500885), National Outstanding

young people plan, the Program for the Top Young Talents by Chongqing, the Fundamental

Research Funds for the Central Universities (SWU1509383,SWU1509451,SWU1609177), Natural

Science Foundation of Chongqing (cstc2015jcyjA10106), Fok Ying Tung Education Foundation

(151023) , General Financial Grant from the China Postdoctoral Science Foundation

(2015M572423, 2015M580767), Special Funds from the Chongqing Postdoctoral Science

Foundation (Xm2015037, Xm2016044), Key research for Humanities and social sciences of

Ministry of Education (14JJD880009). http://fcon_1000.projects.nitrc.org/indi/retro/sald.html

University of Wisconsin, Madison (Birn, Prabhakaran, Meyerand) CoRR sample (UWM): Zuo,

X.N., et al. (2014). An open science resource for establishing reliability and reproducibility in

functional connectomics. *Scientific data, 1*, 140049. doi: 10.1038/sdata.2014.49

http://fcon_1000.projects.nitrc.org/indi/CoRR/html/uwm_1.html

Wayne State EF Dataset: This dataset was supported by National Institute on Aging grants R01-

AG011230, R37-AG011230, R03-AG024630 to Naftali Raz, Ph.D.

http://fcon_1000.projects.nitrc.org/indi/retro/wayne_EF.html

Yale Low-Resolution Controls Dataset: Scheinost D, Tokoglu F, Shen X, Finn ES, Noble S,

Papademetris X, Constable RT. Fluctuations in Global Brain Activity Are Associated With

Changes in Whole-Brain Connectivity of Functional Networks. IEEE Trans Biomed Eng. 2016

Dec;63(12):2540-2549. Epub 2016 Aug 16.

http://fcon_1000.projects.nitrc.org/indi/retro/yale_lowres.html

References

1.   Pfefferbaum A, Rohlfing T, Rosenbloom MJ, Chu W, Colrain IM, Sullivan EV. Variation in longitudinal trajectories of regional brain volumes of healthy men and women (ages 10 to 85 years) measured with atlas-based parcellation of MRI. Neuroimage. 2013;65:176-93. doi: 10.1016/j.neuroimage.2012.10.008. PubMed PMID: 23063452; PubMed Central PMCID: PMC3516371.

2.   Crivello F, Tzourio-Mazoyer N, Tzourio C, Mazoyer B. Longitudinal assessment of global and regional rate of grey matter atrophy in 1,172 healthy older adults: modulation by sex and age. PLoS One. 2014;9(12):e114478. doi: 10.1371/journal.pone.0114478. PubMed PMID: 25469789; PubMed Central PMCID: PMC4255026.

3.   Govindarajan KA, Freeman L, Cai C, Rahbar MH, Narayana PA. Effect of intrinsic and extrinsic factors on global and regional cortical thickness. PLoS One. 2014;9(5):e96429. doi: 10.1371/journal.pone.0096429. PubMed PMID: 24789100; PubMed Central PMCID: PMCPMC4008620.

4.   Kruggel F, Turner J, Muftuler LT. Impact of scanner hardware and imaging protocol on image quality and compartment volume precision in the ADNI cohort. Neuroimage. 2010;49(3):2123-33. doi: 10.1016/j.neuroimage.2009.11.006. PubMed PMID: 19913626; PubMed Central PMCID: PMC2951115.

5.   Potvin O, Dieumegarde L, Duchesne S. Normative morphometric data for cerebral cortical areas over the lifetime of the adult human brain. Neuroimage. 2017;156:315-39. doi: 10.1016/j.neuroimage.2017.05.019. PubMed PMID: 28512057.

6.   Potvin O, Dieumegarde L, Duchesne S. Freesurfer cortical normative data for adults

using Desikan-Killiany-Tourville and ex vivo protocols. Neuroimage. 2017;156:43-64. doi:

10.1016/j.neuroimage.2017.04.035. PubMed PMID: 28479474.

7.   Potvin O, Mouiha A, Dieumegarde L, Duchesne S. Normative data for subcortical

regional volumes over the lifetime of the adult human brain. Neuroimage. 2016;137:9-20. doi:

10.1016/j.neuroimage.2016.05.016. PubMed PMID: 27165761.

8.   Potvin O, Mouiha A, Dieumegarde L, Duchesne S, Alzheimers Disease Neuroimaging I.

FreeSurfer subcortical normative data. Data Brief. 2016;9:732-6. doi:

10.1016/j.dib.2016.10.001. PubMed PMID: 27830169; PubMed Central PMCID:

PMCPMC5094268.

9.   Desikan RS, Segonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, et al. An automated

labeling system for subdividing the human cerebral cortex on MRI scans into gyral based

regions of interest. Neuroimage. 2006;31(3):968-80. doi: 10.1016/j.neuroimage.2006.01.021.

PubMed PMID: 16530430.

10.   Klein A, Tourville J. 101 labeled brain images and a consistent human cortical labeling

protocol. Front Neurosci. 2012;6:171. doi: 10.3389/fnins.2012.00171. PubMed PMID:

23227001; PubMed Central PMCID: PMC3514540.

11.   Destrieux C, Fischl B, Dale A, Halgren E. Automatic parcellation of human cortical gyri

and sulci using standard anatomical nomenclature. Neuroimage. 2010;53(1):1-15. doi:

10.1016/j.neuroimage.2010.06.010. PubMed PMID: 20547229; PubMed Central PMCID:

PMCPMC2937159.

12.   Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C, et al. Whole brain

segmentation: automated labeling of neuroanatomical structures in the human brain. Neuron.

2002;33(3):341-55. PubMed PMID: 11832223.

13.   Augustinack JC, Magnain C, Reuter M, van der Kouwe AJ, Boas D, Fischl B. MRI

parcellation of ex vivo medial temporal lobe. Neuroimage. 2014;93 Pt 2:252-9. doi:

10.1016/j.neuroimage.2013.05.053. PubMed PMID: 23702414; PubMed Central PMCID:

PMCPMC3883990.

14.   Di Martino A, Yan CG, Li Q, Denio E, Castellanos FX, Alaerts K, et al. The autism brain

imaging data exchange: towards a large-scale evaluation of the intrinsic brain architecture in

autism. Mol Psychiatry. 2014;19(6):659-67. Epub 2013/06/19. doi: 10.1038/mp.2013.78.

PubMed PMID: 23774715; PubMed Central PMCID: PMCPMC4162310.

15.   Mueller SG, Weiner MW, Thal LJ, Petersen RC, Jack CR, Jagust W, et al. Ways toward an

early diagnosis in Alzheimer's disease: The Alzheimer's Disease Neuroimaging Initiative (ADNI).

Alzheimers Dement. 2005;1(1):55-66. PubMed PMID: 17476317.

16.   Ellis KA, Bush AI, Darby D, De Fazio D, Foster J, Hudson P, et al. The Australian Imaging,

Biomarkers and Lifestyle (AIBL) study of aging: methodology and baseline characteristics of

1112 individuals recruited for a longitudinal study of Alzheimer's disease. Int Psychogeriatr.

2009;21(4):672-87. doi: 10.1017/S1041610209009405. PubMed PMID: 19470201.

17.   Zuo XN, Anderson JS, Bellec P, Birn RM, Biswal BB, Blautzik J, et al. An open science

resource for establishing reliability and reproducibility in functional connectomics. Scientific

data. 2014;1:140049. doi: 10.1038/sdata.2014.49. PubMed PMID: 25977800; PubMed Central

PMCID: PMC4421932.

18.   Rohr CS, Okon-Singer H, Craddock RC, Villringer A, Margulies DS. Affect and the brain's

functional organization: a resting-state connectivity approach. PLoS One. 2013;8(7):e68015.

Epub 2013/08/13. doi: 10.1371/journal.pone.0068015. PubMed PMID: 23935850; PubMed

Central PMCID: PMCPMC3720669.

19.   Shafto MA, Tyler LK, Dixon M, Taylor JR, Rowe JB, Cusack R, et al. The Cambridge Centre

for Ageing and Neuroscience (Cam-CAN) study protocol: a cross-sectional, lifespan,

multidisciplinary examination of healthy cognitive ageing. BMC Neurol. 2014;14:204. Epub

2014/11/22. doi: 10.1186/s12883-014-0204-1. PubMed PMID: 25412575; PubMed Central

PMCID: PMCPMC4219118.

20.   Taylor JR, Williams N, Cusack R, Auer T, Shafto MA, Dixon M, et al. The Cambridge

Centre for Ageing and Neuroscience (Cam-CAN) data repository: Structural and functional MRI,

MEG, and cognitive data from a cross-sectional adult lifespan sample. Neuroimage. 2017;144(Pt

B):262-9. Epub 2015/09/17. doi: 10.1016/j.neuroimage.2015.09.018. PubMed PMID: 26375206;

PubMed Central PMCID: PMCPMC5182075.

21.   Mayer AR, Ruhl D, Merideth F, Ling J, Hanlon FM, Bustillo J, et al. Functional imaging of

the hemodynamic sensory gating response in schizophrenia. Hum Brain Mapp.

2013;34(9):2302-12. Epub 2012/03/31. doi: 10.1002/hbm.22065. PubMed PMID: 22461278;

PubMed Central PMCID: PMCPMC4020570.

22.   Beall EB, Lowe MJ. Isolating physiologic noise sources with independently determined

spatial measures. Neuroimage. 2007;37(4):1286-300. doi: 10.1016/j.neuroimage.2007.07.004.

PubMed PMID: 17689982.

23.   Belleville S, LeBlanc AC, Kergoat MJ, Calon F, Gaudreau P, Hebert SS, et al. The

Consortium for the early identification of Alzheimer's disease-Quebec (CIMA-Q). Alzheimers

Dement (Amst). 2019;11:787-96. Epub 2019/12/04. doi: 10.1016/j.dadm.2019.07.003. PubMed

PMID: 31788534; PubMed Central PMCID: PMCPMC6880140.

24.   Dallas Lifespan Brain Study (DLBS). Available from:

http://fcon_1000.projects.nitrc.org/indi/retro/dlbs.html.

25.   Shirer WR, Ryali S, Rykhlevskaia E, Menon V, Greicius MD. Decoding subject-driven

cognitive states with whole-brain connectivity patterns. Cereb Cortex. 2012;22(1):158-65. Epub

2011/05/28. doi: 10.1093/cercor/bhr099. PubMed PMID: 21616982; PubMed Central PMCID:

PMCPMC3236795.

26.   Keator DB, van Erp TGM, Turner JA, Glover GH, Mueller BA, Liu TT, et al. The Function

Biomedical Informatics Research Network Data Repository. Neuroimage. 2016;124(Pt B):1074-

9. Epub 2015/09/15. doi: 10.1016/j.neuroimage.2015.09.003. PubMed PMID: 26364863;

PubMed Central PMCID: PMCPMC4651841.

27.   Bookheimer SY, Salat DH, Terpstra M, Ances BM, Barch DM, Buckner RL, et al. The

Lifespan Human Connectome Project in Aging: An overview. Neuroimage. 2019;185:335-48.

Epub 2018/10/18. doi: 10.1016/j.neuroimage.2018.10.009. PubMed PMID: 30332613; PubMed

Central PMCID: PMCPMC6649668.

28.   Mazziotta J, Toga A, Evans A, Fox P, Lancaster J, Zilles K, et al. A probabilistic atlas and

reference system for the human brain: International Consortium for Brain Mapping (ICBM).

Philos Trans R Soc Lond B Biol Sci. 2001;356(1412):1293-322. PubMed PMID: 11545704.

29.   IXI Dataset. Available from: http://brain-development.org/ixi-dataset/.

30.   Landman BA, Huang AJ, Gifford A, Vikram DS, Lim IA, Farrell JA, et al. Multi-parametric

neuroimaging reproducibility: a 3-T resource study. Neuroimage. 2011;54(4):2854-66. doi:

10.1016/j.neuroimage.2010.11.047. PubMed PMID: 21094686; PubMed Central PMCID:

PMC3020263.

31.   Malone IB, Cash D, Ridgway GR, MacManus DG, Ourselin S, Fox NC, et al. MIRIAD--Public

release of a multiple time point Alzheimer's MR imaging dataset. Neuroimage. 2013;70:33-6.

Epub 2013/01/01. doi: 10.1016/j.neuroimage.2012.12.044. PubMed PMID: 23274184; PubMed

Central PMCID: PMCPMC3809512.

32.   Beekly DL, Ramos EM, van Belle G, Deitrich W, Clark AD, Jacka ME, et al. The National

Alzheimer's Coordinating Center (NACC) Database: an Alzheimer disease database. Alzheimer

Dis Assoc Disord. 2004;18(4):270-7. PubMed PMID: 15592144.

33.   Hall D, Huerta MF, McAuliffe MJ, Farber GK. Sharing heterogeneous data: the national

database for autism research. Neuroinformatics. 2012;10(4):331-9. Epub 2012/05/25. doi:

10.1007/s12021-012-9151-4. PubMed PMID: 22622767; PubMed Central PMCID:

PMCPMC4219200.

34.   Nooner KB, Colcombe SJ, Tobe RH, Mennes M, Benedict MM, Moreno AL, et al. The NKI-

Rockland Sample: A Model for Accelerating the Pace of Discovery Science in Psychiatry. Front

Neurosci. 2012;6:152. doi: 10.3389/fnins.2012.00152. PubMed PMID: 23087608; PubMed

Central PMCID: PMC3472598.

35.   Marcus DS, Wang TH, Parker J, Csernansky JG, Morris JC, Buckner RL. Open Access Series

of Imaging Studies (OASIS): Cross-sectional MRI Data in Young, Middle Aged, Nondemented,

and Demented Older Adults. J Cogn Neurosci. 2007;19(9):1498-507. PubMed PMID: 17714011.

36.   Power JD, Barnes KA, Snyder AZ, Schlaggar BL, Petersen SE. Spurious but systematic

correlations in functional connectivity MRI networks arise from subject motion. Neuroimage.

2012;59(3):2142-54. doi: 10.1016/j.neuroimage.2011.10.018. PubMed PMID: 22019881;

PubMed Central PMCID: PMC3254728.

37.   Parkinson Progression Marker I. The Parkinson Progression Marker Initiative (PPMI).

Prog Neurobiol. 2011;95(4):629-35. Epub 2011/09/21. doi: 10.1016/j.pneurobio.2011.09.005.

PubMed PMID: 21930184.

38.   Wei D, Zhuang K, Ai L, Chen Q, Yang W, Liu W, et al. Structural and functional brain scans

from the cross-sectional Southwest University adult lifespan dataset. Scientific data.

2018;5:180134. Epub 2018/07/18. doi: 10.1038/sdata.2018.134. PubMed PMID: 30015807;

PubMed Central PMCID: PMCPMC6049036.

39.   Moffat SD, Kennedy KM, Rodrigue KM, Raz N. Extrahippocampal contributions to age

differences in human spatial navigation. Cereb Cortex. 2007;17(6):1274-82. Epub 2006/07/22.

doi: 10.1093/cercor/bhl036. PubMed PMID: 16857855.

40.   Scheinost D, Tokoglu F, Shen X, Finn ES, Noble S, Papademetris X, et al. Fluctuations in

Global Brain Activity Are Associated With Changes in Whole-Brain Connectivity of Functional

Networks. IEEE Trans Biomed Eng. 2016;63(12):2540-9. Epub 2016/08/20. doi:

10.1109/TBME.2016.2600248. PubMed PMID: 27541328; PubMed Central PMCID:

PMCPMC5180443.

41.   Sheikh JI, Yesavage JA. Geriatric Depression Scale (GDS): Recent evidence and

development of a shorter version. Clinical Gerontology: a Guide to Assessment and

Intervention. New York: The Haworth Press; 1986. p. 165-73.

42.   Sherif T, Rioux P, Rousseau ME, Kassis N, Beck N, Adalat R, et al. CBRAIN: a web-based,

distributed computing platform for collaborative neuroimaging research. Front Neuroinform.

2014;8:54. Epub 2014/06/07. doi: 10.3389/fninf.2014.00054. PubMed PMID: 24904400;

PubMed Central PMCID: PMCPMC4033081.

43.   Augustinack JC, Huber KE, Stevens AA, Roy M, Frosch MP, van der Kouwe AJ, et al.

Predicting the location of human perirhinal cortex, Brodmann's area 35, from MRI. Neuroimage.

2013;64:32-42. doi: 10.1016/j.neuroimage.2012.08.071. PubMed PMID: 22960087; PubMed

Central PMCID: PMCPMC3508349.

44.   Iglesias JE, Van Leemput K, Bhatt P, Casillas C, Dutt S, Schuff N, et al. Bayesian

segmentation of brainstem structures in MRI. Neuroimage. 2015;113:184-95. doi:

10.1016/j.neuroimage.2015.02.065. PubMed PMID: 25776214; PubMed Central PMCID:

PMCPMC4434226.

45.   Iglesias JE, Augustinack JC, Nguyen K, Player CM, Player A, Wright M, et al. A

computational atlas of the hippocampal formation using ex vivo, ultra-high resolution MRI:

Application to adaptive segmentation of in vivo MRI. Neuroimage. 2015;115:117-37. doi:

10.1016/j.neuroimage.2015.04.042. PubMed PMID: 25936807; PubMed Central PMCID:

PMCPMC4461537.

46.   Rosas HD, Liu AK, Hersch S, Glessner M, Ferrante RJ, Salat DH, et al. Regional and

progressive thinning of the cortical ribbon in Huntington's disease. Neurology. 2002;58(5):695-

701. PubMed PMID: 11889230.

47.   Salat DH, Buckner RL, Snyder AZ, Greve DN, Desikan RS, Busa E, et al. Thinning of the

cerebral cortex in aging. Cereb Cortex. 2004;14(7):721-30. PubMed PMID: 15054051.

48.   Kuperberg GR, Broome MR, McGuire PK, David AS, Eddy M, Ozawa F, et al. Regionally

localized thinning of the cerebral cortex in schizophrenia. Arch Gen Psychiatry. 2003;60(9):878-

88. PubMed PMID: 12963669.

49.   Buckner RL, Head D, Parker J, Fotenos AF, Marcus D, Morris JC, et al. A unified approach

for morphometric and functional data analysis in young, old, and demented adults using

automated atlas-based head size normalization: reliability and validation against manual

measurement of total intracranial volume. Neuroimage. 2004;23(2):724-38. doi:

10.1016/j.neuroimage.2004.06.018. PubMed PMID: 15488422.

50.   Klapwijk ET, van de Kamp F, van der Meulen M, Peters S, Wierenga LM. Qoala-T: A

supervised-learning tool for quality control of FreeSurfer segmented MRI data. Neuroimage.

2019;189:116-29. doi: 10.1016/j.neuroimage.2019.01.014. PubMed PMID: 30633965.

51.   Backhausen LL, Herting MM, Buse J, Roessner V, Smolka MN, Vetter NC. Quality Control

of Structural MRI Images Applied Using FreeSurfer-A Hands-On Workflow to Rate Motion

Artifacts. Front Neurosci. 2016;10:558. doi: 10.3389/fnins.2016.00558. PubMed PMID:

27999528; PubMed Central PMCID: PMCPMC5138230.

52.   Hastie T, Tibshirani R, Friedman J. The elements of statistical learning. Data mining,

inference, and prediction. Statistics, editor: Springer; 2008. 745 p.

53.   R Development Core Team. R: A language and environment for statistical computing.

Vienna, Austria: R Foundation for Statistical Computing. http://www.R-project.org2010.

54.   Grömping U. Relative Importance for Linear Regression in R: The Package relaimpo.

Journal of Statistical Software. 2006;17(1):1-27.

55.   Mowinckel AM, Vidal-Pineiro V. Visualisation of Brain Statistics with R-packages ggseg

and ggseg3d. arXiv. 2019. doi: https://arxiv.org/abs/1912.08200.

56.   Crawford JR, Garthwaite PH, Denham AK, Chelune GJ. Using regression equations built

from summary data in the psychological assessment of the individual case: extension to

multiple regression. Psychological assessment. 2012;24(4):801-14. doi: 10.1037/a0027699.

PubMed PMID: 22449035.

Total variance explained
Training 10-fold CV

Volume

Thickness

Subcortical volume

Test validation (test - training values)

Volume

Thickness

Subcortical volume

Biological effects

Age

Sex

eTIV

MRI effects
Scanner

Volume   Thickness

Subcortical volume

Image quality

Volume   Thickness

Subcortical volume

Norms for age and sex

Alzheimer's disease

Volume

Thickness

Subcortical volume

Z score

Schizophrenia

Volume

Thickness

Subcortical volume

Z score