NOMIS: Quantifying morphometric deviations from normality over the lifetime of the adult human brain

Olivier Potvin[1], PhD, Louis Dieumegarde[1], BSc, and Simon Duchesne, PhD[12] for the Alzheimer's Disease Neuroimaging Initiative,* the CIMA-Q** and the CCNA*** groups

[1] CERVO Brain Research Centre, 2601, de la Canardière, Québec, Canada, G1J 2G3

[2] Département de radiologie, Faculté de médecine, Université Laval, 1050, avenue de la Médecine, Québec, Canada, G1V 0A6

Key words: magnetic resonance imaging, atrophy, morphometry, normality, aging, sex.

Correspondence:

Simon Duchesne, PhD
CERVO Brain Research Centre
F-3582, 2601, de la Canardière, Québec, Canada, G1J 2G3
Phone: 418 663-5000 ext.4777
Fax: 418 663-9540
simon.duchesne@fmed.ulaval.ca

*Part of the data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

** Part of the data used in this article were obtained from the Consortium pour l'identification précoce de la maladie Alzheimer - Québec (CIMA-Q; cima-q.ca). As such, the investigators within the CIMA-Q contributed to the design, the implementation, the acquisition of clinical, cognitive, and neuroimaging data and biological samples. A list of the CIMA-Q investigators is available on www.cima-q.ca.

** Part of the data used in this article were obtained from the Canadian Consortium on Neurodegeneration in Aging (CCNA; www.ccna-ccnv.ca) .

Abstract

We present NOMIS (https://github.com/medicslab/NOMIS), a comprehensive open MRI tool to assess morphometric deviation from normality in the adult human brain. Based on MR anatomical images from 6,909 cognitively healthy individuals aged 18-100 years, we modeled 1,344 measures computed using the open access *FreeSurfer* pipeline, considering account personal characteristics (age, sex, intracranial volume) and image quality (resolution, contrast-to-noise ratio and surface reconstruction defect holes), and providing expected values for any new individual. Then, for each measure, the NOMIS tool was built to generate Z-score effect sizes denoting the extent of deviation from the normative sample. Depending on the user need, NOMIS offers four versions of Z-score adjusted on different sets of variables. While all versions consider head size and image quality, they can also incorporate age and/or sex, thereby facilitating multi-site neuromorphometric research across adulthood.

Introduction

Despite the popularity of magnetic resonance imaging (MRI) to examine abnormalities in brain

morphometry, tools quantifying normality are lacking. While age, sex and intracranial volume are

well-known to influence brain volume and shape[1, 2] the determination of whether an

individual's brain region measurements are within normality faces multiple major challenges

such as the lack of normative data across appropriate age groups, the influence of the MRI

processing pipeline, the variety in neuroanatomical atlases used for parcellation and the

uniqueness of the image acquisition itself[3, 4]. We made previous attempts[5-8] to produce such

normative data in adulthood based on *FreeSurfer*, an open-access and fully automated

segmentation software (http://freesurfer.net), for two specific brain atlases, namely Desikan-

Killiany[9] (DK) and Desikan-Killiany-Tourville[10] (DKT). This initial foray allowed for the

quantification of the extent of deviation from normality for a given individual, according to

personal characteristics such as age, sex and estimated intracranial volume (eTIV), while

controlling for scanner magnetic field strength (MFS) and original equipment manufacturer

(OEM).

Leveraging this prior work, we offer a comprehensive tool called NOMIS (NOrmative

Morphometry Image Statistics; https://github.com/medicslab/NOMIS). NOMIS can be used to

produce normative values for any new adult individual, cognitively healthy or otherwise. Using

this individual's T1-weighted MRI, processed via the *FreeSurfer* 6.0 toolkit, one can derive Z-score

effect sizes denoting the extent of deviation from the normative sample according to the

individual's characteristics (age, sex, and eTIV), while taking into account image quality

information (resolution, contrast-to-noise ratio (CNR) and holes in surface reconstruction)[11,

12]. NOMIS contains 1,344 brain measures generated by *FreeSurfer* on 6,909 healthy individuals

aged 18 to 100 years (mean ±sd: 55.0 ±20.0; 56.8% female). The normative data includes as

before the DK[9] and DKT[10] atlases, as well as the Destrieux (a2009s)[13] neocortical atlas;

neocortical pial and white surface areas, volumes and thicknesses; *FreeSurfer*'s default

subcortical atlas[14], hippocampal subfields, brainstem subregions; its ex vivo-based labeling

protocol atlas[15]; and the subcortical white matter parcellation according to the adjacent

neocortical areas. Furthermore, to fulfill specific needs from researchers, we propose four

versions of Z-score adjusted on different sets of variables. While all versions are adjusted for head

size and image quality, the full version includes both age and sex whereas the three other

versions are without age, without sex and without age and sex. Thus, a research group working

on aging aiming at removing the variance of hippocampal volumes due to head size, sex, and

image quality could use the version without age, which preserves the variance due to aging.

When compared to our previous work on normative values, there are important new

contributions in NOMIS:

• The norms were calculated using a newer *FreeSurfer* software version

• New variables were added to remove undesirable variance (CNR, surface holes,

resolution)

• New atlases were processed, such as Destrieux, hippocampal subfields, brainstem

subregions, ex vivo-based labeling protocol atlas, subcortical white matter parcellation

according to the adjacent neocortical areas

• The possibility of calculating normative scores while adjusting only some selected

variables was introduced (intracranial volume with image quality in combination, or not,

with age and/or sex)

• The sample size of the normative sample was doubled, making the age distribution more

uniform than previously

The multiple scanner problem

Different scanners produce different images, even in the same individuals, which produce

in the end different morphometric values. One way of capturing inter-scanner variance is using

information about the scanner (e.g. magnetic field strength and vendor). For the creation of

NOMIS, and contrary to our previous work, we chose not to incorporate such information since

the samples of individuals within each combination of scanner characteristic is likely to be

different and thus, possibly bias-inducing due to known or unknown individuals' characteristics

stemming from recruitment in a particular study included in the training data. Therefore, to

minimize inter-scanner variance, NOMIS strictly uses image information.

Moreover, as a final validation step, we have compared the basic version of NOMIS (i.e.

only adjusting for head size and image quality) with two global scaling harmonization techniques,

namely NeuroCombat[16] and NeuroHarmonize[17] on their ability to reduce the mean effect

and variance induced by different scanners. Such techniques model the differences between

scanners to apply a post-hoc correction on morphometric estimates based on the complete set

of data in the study. In that, they are fundamentally different from our attempt here at a

normative tool to be used in new, separate studies.

Nevertheless, it should be noted that, while they are gaining popularity, harmonization

techniques can potentially induce biases due to the different participants' characteristics at each

scanner[18, 19]. The main challenge to show that harmonization is actually working is that MRI

provides relative measures for which that there is no gold standard; each scanner yields its own

measure, given its hardware software and other factors, even time of the day[20]. In order to

properly test harmonization, we defined our own gold standard by using the Single Individual

Across Networks (SIMON) dataset[21], comprised of images from a single person that was

scanned within a short span at 12 sites for quality control purposes in the context of within two

Canadian studies. By harmonizing these 12 scanners using 547 MRIs from individuals scanned in

these studies, as well as the quality-control 48 SIMON MRIs, we verified whether the variance of

the SIMON measures was lower or not. Unfortunately, we conclude that none of the

harmonization techniques reduce real inter-scanner variance. While neither does NOMIS, such is

not our purpose.

127 Materials and methods

128 Normative sample

The norms are based on a cross-sectional sample of 6,909 (initial sample: 7,399) cognitively

healthy individuals aged 18 to 100 years, (mean $\pm$sd; 55.0 $\pm$20.0; 56.8% female), gathered from

27 different datasets (Table 1). Supplementary Fig 1 shows the age distribution within each

dataset. Scans were acquired from one of the three leading OEM (e.g. Siemens Healthcare

(Erlangen, Germany); Philips Medical Systems (Best, Netherlands); or GE Healthcare (Milwaukee,

WI)) at MFS of either 1.5 or 3 Tesla. This study received the approval of the Institutional review

board of neuroscience and mental health of the CIUSSS de la Capitale- Nationale (#217207).

From all the samples mentioned, only cognitively healthy (control) participants were

included. For the Nathan Kline Institute samples, which were projects recruiting in the general

population, we excluded participants with history of schizophrenia or other psychotic disorders,

bipolar disorders, major depressive disorders (recurrent), posttraumatic stress disorder,

substance abuse/dependence disorders, neurodegenerative and neurological disorders, head

injury with loss of consciousness/amnesia, and lead poisoning. Moreover, for the Parkinson's

Progression Markers Initiative dataset, we excluded participants with a Geriatric Depression

Scale[22] score of more than 5.

Table 1. Datasets included in the normative sample

Dataset	n
Autism Brain Imaging Data Exchange (ABIDE)[23]	183
Alzheimer's Disease Neuroimaging Initiative (ADNI)[24]	672
Australian Imaging Biomarkers and Lifestyle flagship study of ageing (AIBL)[25]	157
Berlin Mind and Brain (Margulies, Villringer) CoRR sample (BMB)[26, 27]	50
Cambridge Centre for Ageing and Neuroscience (CamCAN)[28, 29]	630
Center of Biomedical Research Excellence (COBRE)[30]	70
Cleveland CCF[31]	30
Consortium for the Early Identification of Alzheimer's Disease (CIMA-Q)[32]	29
Dallas Lifespan Brain Study (DLBS)[33]	304
FIND lab sample (FIND) Lab[34]	13
Functional Biomedical Informatics Research Network (FBIRN)[35]	33
Lifespan Human Connectome Project in Aging (HCP-Aging)[36]	612
International Consortium for Brain Mapping (ICBM) - MNI[37]	147
Information eXtraction from Images (IXI)[38]	554
F.M. Kirby Research Center neuroimaging reproducibility data (KIRBY-21)[39]	20
Minimal Interval Resonance Imaging in Alzheimer's Disease (MIRIAD)[40]	21
National Alzheimer's Coordinating Center (NACC)[41]	1562
National Database for Autism Research (NDAR)[42]	56

Nathan Kline Institute Rockland Sample(NKI-RS)[43]	138
Nathan Kline Institute Rockland Enhanced Sample (NKI-RES) [43]	436
Open Access Series of Imaging Studies (OASIS)[44]	288
POWER Neuroimage sample (POWER)[45]	26
Parkinson's Progression Markers Initiative (PPMI)[46]	158
Southwest University Adult Lifespan Dataset (SALD)[47]	490
University of Wisconsin (Birn, Prabhakaran, Meyerand) CoRR sample (UWM)[26]	25
Wayne State EF Dataset (Wayne State)[48]	108
Yale Low-Resolution Controls Dataset (Yale)[49]	97
Total	6909

Among the datasets are the Alzheimer's Disease Neuroimaging Initiative (ADNI), the Australian

Imaging, Biomarkers and Lifestyle study of aging (AIBL) and the Consortium pour l'identification

précoce de la maladie Alzheimer - Québec (CIMA-Q) datasets. The ADNI (adni.loni.usc.edu) was

launched in 2003 as a public-private partnership, led by Principal Investigator Michael W. Weiner,

MD. (www.adni-info.org). The AIBL data was collected by the AIBL study group and AIBL study

methodology has been reported previously by Ellis et al. (2009). For each dataset, approval from

the local ethics board and informed consent of the participants were obtained. Founded in 2013,

the main objective of CIMA-Q is to build a cohort of participants characterized in terms of

cognition, neuroimaging and clinical outcomes in order to acquire biological samples allowing (1)

to establish early diagnoses of Alzheimer's disease, (2) to provide a well characterized cohort and

(3) to identify new therapeutic targets. The principal investigator and director of CIMA-Q is Dr

Sylvie Belleville from the Centre de recherche de l'Institut universitaire de gériatrie de Montréal,

CIUSSS Centre-sud-de-l'Île-de-Montréal. CIMA-Q represent a common effort of several

researchers from Québec affiliated to Université Laval, Université McGill, Université de Montréal,

et Université de Sherbrooke. CIMA-Q recruited cognitively healthy participants, participants with

subjective cognitive impairment, mild cognitive impairment, or Alzheimer's disease, between

2013–2016. Volunteers were recruited from memory clinics, through advertisements posted in

the community and amongst participants from the NuAge population study.

Harmonization test sample

For the harmonization test, we used three datasets: 1) the complete CIMA-Q sample (n=208

participants; 286 MRIs), which was described earlier in the method section, 2) the

Comprehensive Assessment of Neurodegeneration and Dementia (COMPASS-ND; n=393) study

conducted by the Canadian Consortium on Neurodegeneration in Aging (CCNA), and the 3) the

SIMON dataset[19], comprised of images from single healthy volunteer scanned on the same

scanner as those used for CIMA-Q and COMPASS-ND. From COMPASS-ND, we used participants

that were either cognitively unimpaired participants (CU), with mild cognitive impairment (MCI),

and with probable Alzheimer's disease (AD), for a total of 273 participants. While CIMA-Q and

COMPASS-ND were acquired at 18 different sites, we selected only data from scanners that had

at least three participants other than SIMON, which resulted in a total of 547 images (300 CU,

193 MCI, 54 AD) from 12 different scanners; each ranging from 7 to 145 participants. On those

12 scanners, SIMON was scanned 48 times and was aged between 42-46 years old during that

time.

Brain segmentation

Brain segmentation was conducted using *FreeSurfer* version 6.0, a widely used and freely

available automated processing pipeline that quantifies brain anatomy (http://freesurfer.net).

All raw T1-weighted images were processed using the "recon-all -all" *FreeSurfer* command with

the fully-automated directive parameters (no manual intervention or expert flag options) on the

CBRAIN platform[50]. Normative data were computed for volumes, neocortical thicknesses and

white and pial surfaces areas for all atlases comprised in *FreeSurfer* 6.0: the default subcortical

atlas[14] (aseg.stats), the Desikan-Killiany atlas[9] (DK, aparc.stats file), the Desikan-Killiany-

Tourville atlas[10] (DKT, aparc.DKT.stats file), the Destrieux atlas[13] (aparc.a2009s.stats file), the

ex vivo atlas,[51] including entorhinal and perirhinal cortices, the brainstem sub-regions

atlas[52], the Brodmann area maps which includes somatosensory areas, several motor and

visual areas, as well as the hippocampal subfields atlas[53].

The technical details of *FreeSurfer*'s procedures are described in prior publications.

Briefly, this processing includes motion correction, removal of non-brain tissue using a hybrid

watershed/surface deformation procedure, automated Talairach transformation, intensity

normalization, tessellation of the gray matter white matter boundary, automated topology

correction, and surface deformation following intensity gradients to optimally place the

gray/white and gray/cerebrospinal fluid borders at the location where the greatest shift in

intensity defines the transition to the other tissue class. Once the cortical models are complete,

a number of deformable procedures can be performed for further data processing and analysis

including surface inflation, registration to a spherical atlas which is based on individual cortical

folding patterns to match cortical geometry across subjects and parcellation of the cerebral

cortex into units with respect to gyral and sulcal structure. This method uses both intensity and

continuity information from the entire three-dimensional MRI volume in segmentation and

deformation procedures to produce representations of cortical thickness, calculated as the

closest distance from the gray/white boundary to the gray/cerebrospinal fluid boundary at each

vertex on the tessellated surface. The maps are created using spatial intensity gradients across

tissue classes and are therefore not simply reliant on absolute signal intensity. Procedures for the

measurement of cortical thickness have been validated against histological analysis [54] and

manual measurements[55, 56]. Estimated intracranial volume[57] was taken from the aseg.stats

FreeSurfer output file. We added the total ventricle volume (labeled as "ventricles") using the

sum of all ventricles and the corpus callosum (labeled as "cc") using the sum of all corpus

callosum segments.

Quality control and sample selection

A flow chart detailing the final analysis sample is shown in Fig 1. From an initial pool of 7,399

MRIs, nine images failed the *FreeSurfer* pipeline. Following processing, each of the remaining

7,390 brain segmentations was visually inspected through at least 20 evenly distributed coronal

sections by O.P. (see Supplementary materials for quality control examples). After quality control,

445 images (6.0%) were removed from further analyses due to segmentation problems, the main

reason being that parts of the brain were not completely segmented (e.g. temporal and occipital

poles. During visual inspection, 26 brains were found to have signal alterations or clear significant

brain lesions and were excluded. Quality control image examples are displayed as Supplementary

materials (setup, segmentation errors and abnormalities). In addition to visual inspection, we

excluded participants if at least one of the 1,344 brain region measures was missing (n=10). In

fine, the analysis sample was composed of 6,909 individual MRIs.

Fig 1. Flow chart of the images.

Training, validation and test sample

We randomly selected 10% of the whole sample (n=691) to test the models in an

independent sample (age: 55.1 ±20.1, range 18-100; 58.5% female). This test sample was not

used to build the models predicting normative values. The remaining 90% was used as training

sample (age: 54.9 ±20.0, range 18-100; 56.7% female) to build and validate the models. Leave-

10%-out cross-validation was used to validate the model in the training sample.

Clinical samples evaluations

We evaluated the usefulness of normative values using clinical samples of individuals with

schizophrenia (n=72; Age: 38.2 ±13.9, range 18-65; 19% female) from the COBRE dataset, as well

as participants with clinically ascertained Alzheimer's disease (n=157 Age: 74.8 ±8.1, range 55-

90; 43% female) from the baseline ADNI-2 dataset.

Image quality predictors

Image quality predictors included voxel size (resolution) and two measures of image quality,

one global, and the second local. The first was the total number of defect holes over the whole

cortex, i.e. topological errors in the initial cortical surface reconstructions. The total number

correlated well with visual inspection of the whole image by trained raters [11]. This measure

was extracted from the aseg.stats *FreeSurfer* output file. The second measure was contrast-to-

noise ratio (CNR) assessed in each region (R) and therefore used as a regional measure of image

quality. For each region, CNR was calculated after *FreeSurfer* preprocessing using gray matter

(GM) and cerebral white matter (WM) intensities from the brain.mgz file and the following

formula:

246
$$CNR_R = \frac{(GM_R\ mean\ -\ WM\ mean\ )^2}{(GM_R\ variance\ +\ WM\ variance)}$$

Outliers removal

For each brain measure, to exclude potential abnormalities, outliers with Z scores lower than -

3.29 and higher than 3.29 (p < .001) were removed before computing the statistical model. This

procedure allowed the identification of brain regions that were either very small or very large

when compared to the rest of the sample and thus, might not be good representative of

normality. For volumes and surfaces, this procedure was applied in proportion to eTIV (i.e.

regional measure/eTIV). Since cortical thickness is not affected by eTIV, the outliers screening

procedure was applied directly on the raw values. The number of outliers was below 1% for all

regions (mean ±sd of all atlases: 0.45% ±0.10%) except the right long insular gyrus and central

sulcus of the insula white surface (1.1%) and pericallosal sulcus volume (1.1%) of the Destrieux

atlas. Detailed results can be found in the supplementary material as csv files.

Regression models and statistical analyses

For each brain region measure, the normative values were produced following two linear

regression models. First, a Model 1 was conducted with image quality predictors (voxel size,

surface defect holes and CNR) and eTIV. Then, Model 2 with age and sex was applied on the

residuals of Model 1. In order to respect the normality of the residuals, surface holes and all

ventricles variables, except the 4th (3rd, lateral, inferior lateral and the sum of all ventricles), were

log transformed. For ventricles and white matter regions, CNR of the total brain gray matter was

used while for the brainstem subregions and hippocampal subfields, CNR from the whole

brainstem and whole hippocampus were used, respectively. Quadratic and cubic terms for age,

CNR and surface holes were included. Since voxel size has a relatively limited variability (mean:

1.02, std: 0.24, range: 0.18-2.2), we chose not to include quadratic and cubic terms for this

variable. We also included all interactions except for voxel size (Model 1: eTIV X surface holes,

eTIV x CNR, CNR X surface holes ; Model 2: age X sex). Feature selection was conducted with a

10-fold cross-validation[58] backward elimination procedure, retaining the model with the

subset of predictors that produced the lowest predicted residual sum of squares. For each

selected final model, the fit of the data was assessed using R^2 coefficient of determination:

275
$$R^2 = 1 - \frac{\sum_i (Y_i - f_i)^2}{\sum_i (Y_i - \hat{Y})^2}$$

where the numerator is the residual sum of squares (Y is the value of the variable to be predicted

and f is the predicted value), the denominator is the total sum of squares ($\hat{Y}$ is the mean) and i is

the index over subjects. To assess the unique contribution of each predictor, we used the lmg

metric in the R package[59] relaimpo[60]. This metric is a R^2 partitioned by averaging sequential

sums of squares over all orderings of the predictors. Brain figures were made using the ggseg R

package[61]. In order to compare the effects of each predictor, the sum of all relaimpo R^2 terms

related to each variable was computed (i.e. quadratic, cubic, and half of interaction values). For

example, the variance explained by age includes the R^2 sum for age, age^2, age^3, age X sex /2.

When a term was not included within a model, its R^2 value was given 0.

The models were verified by examining the difference between R^2 of the training sample

and R^2 of the independent test sample of healthy controls. It was expected that the test R^2 would

be within 10% from the value of the training R^2. Then, patterns of normality deviations were

examined with the Z score effect sizes using the validation samples of healthy individuals and of

individuals with AD and SZ.

290
$$Z_{OP} = \frac{Y_o - \hat{Y}}{RMSE}$$

Z score effect sizes (Z_{OP}) were obtained by subtracting the *Predicted value* ($\hat{Y}$) from the *Observed*

value (Y_o) divided by the root mean square error (RMSE) of the model predicting the value [62].

Harmonization test

While the goal of NOMIS differs, we compared its results with twoharmonization procedures,

NeuroCombat[16] and NeuroHarmonize[17] on the aseg volume and DKT cortical volume and

thickness measures (matrix of 146 brain measures) from the harmonization dataset (SIMON,

CIMA-Q and COMPASS-ND). We used the scanner identification number as "batch" (i.e. site)

variable. For NeuroCombat, we also specified age and eTIV as covariates to preserve their effects.

To compare harmonization procedures with NOMIS, after harmonization, eTIV was regressed out

from the brain volume measures. Finally, to compare them on the same scale for statistical

analyses on the variance and figure presentations, all measures were transformed into T and Z

scores, respectively (see Supplementary Fig 2 as example).

We had three expectations following harmonization procedures. Compared to raw data, these

procedures should:

• Reduce the variance of the measures from the 48 SIMON MRIs

• Maintain or increase effect sizes for hippocampi volumes and entorhinal thicknesses

between CU, MCI, and AD groups

• Maintain or increase effect sizes for the relationships between hippocampi volumes and

entorhinal thicknesses and episodic memory as measured by the delayed recall performance of

the Rey Auditory Verbal Learning Test (RAVLT)[63] and Logical Memory Test[64].

The change in variance was assess using the quartile coefficient of dispersion (QCD):

312
$$\frac{Q3 - Q1}{Q3 + Q1}$$

313   where Q1 and Q3 are the first and third quartile, respectively, and the Levene's test for

314   homogeneity of variance:

315
$$Z_{ij} = \left|Y_{ij} - \bar{Y}_i\right|$$

where *i* and *j* are the groups and the individuals, respectively. The Levene's test is equivalent to

a one-way between-groups analysis of variance (ANOVA) with the dependent variable being the

absolute value of the difference between a score (Y) and the mean of the group ($\bar{Y}$). For each

harmonization procedure, a one-way ANOVA on the QCD of the 146 measures before (raw) and

after harmonization was conducted and a Levene's test on each of the 146 measures was

conducted.

All statistics were conducted using python's module Scikit-learn[65] and Statsmodel[66].

Results

As examples, figures in this report display results for subcortical volumes and DKT neocortical

atlases volumes and thicknesses. Full detailed results for all atlases are provided as

supplementary information as csv files.

Model 1 – Image quality and eTIV

The R^2 for model 1 ranged between 0.003 to 0.75, with a mean $\pm$sd of 0.23 ± 0.15. The highest

R^2 were observed in brain measures with the largest volumes and surface areas (i.e. left and

right white surface areas 0.82 and 0.81, and left and right pial surface areas 0.79 and 0.78, brain

segmentation volume 0.75, supratentorial volume 0.75). Fig 2 shows the R^2 portion due to

image quality and eTIV for neocortical volumes and thicknesses of the DKT atlas parcellation, as

well as subcortical volumes. Image quality had a substantial impact with a mean ±sd of 0.08

±0.09, 0.14 ±0.05 and 0.13 ±0.05, for subcortical, neocortical volume and thickness values,

respectively. A high amount of variance due to eTIV was observed in subcortical and neocortical

volumes 0.22 ±0.12 and 0.26 ±0.08, respectively while it had nearly impact on cortical thickness

measures 0.01 ±0.01.

Model 2 – Age and sex

The R^2 for model 2 ranged between 0.02 to 0.51, with a mean ±sd of 0.23 ±0.14, 0.08

±0.04 and 0.11 ±0.07, for subcortical volumes, neocortical volumes and thicknesses, respectively.

One should note that the R^2 in model 2 cannot be compared to that of model 1 since the total

variance in model 2 is the remaining variance after model 1 (residuals). The highest R^2 were

observed in the largest regions and ventricles (i.e. all ventricles volume 0.51, brain segmentation

volume 0.49, left and right lateral ventricles 0.49 and 0.49, supratentorial volumes 0.46). The

lowest age and sex effects were generally on pial and white surface areas (0.00 ±0.10 and 0.03

±0.02). Fig 3 illustrates the R^2 for age and sex. As shown, sex did not explain much variance while

age had a very different impact depending on the part of the brain with a higher impact in

subcortical volumes and associative cortices, both in volume and thickness.

Models validation

Model 1 and Model 2 were examined in the independent test sample and nearly all

models showed equivalent or higher R^2 on the test set than on the training set (the difference

test minus training for all atlases: Model 1 -0.026 ±0.027, Model 2 -0.005 ±0.018; Fig. 4). In model

1, the worse test differences were in the Destrieux atlas where 5 measures out of 592 were below

-10%: pial surface areas of the left superior temporal sulcus (-0.13) and right fronto-marginal

gyrus (of Wernicke) and sulcus (-0.12), the left subcallosal area, subcallosal gyrus volume (-0.10),

the white surface area of the left lateral aspect of the superior temporal gyrus (0.10) and the

right fronto-marginal gyrus (of Wernicke) and sulcus volume (-0.10). In Model 2, all measures had

R^2 differences below 10%, the worse being the left and right putamen volumes (-0.09 and -0.08).

One should note that the models for these measures appear to be slightly less generalizable than

the others.

Fig 5 and Fig 6 show the mean and std Z scores adjusted for age and sex when the models

are applied on the independent young and older healthy controls samples. As expected, the

mean Z scores were very close to 0 while the std were very close to 1 (mean ±std, Young,

subcortical regions: -0.06 ±1.04, cortical volume: 0.04 ±1.13, cortical thickness: 0.04 ±0.99; Older,

subcortical regions: 0.02 ±1.04, cortical volume: 0.01 ±0.96, cortical thickness: 0.01 ±1.02).

Using the independent healthy control test sample, Fig 7 and Fig 8 display examples of

how the normative values remove the different effects on the left cortical thickness and volumes.

Fig 2. R^2 from Model 1 for cortical volumes and thicknesses from the DKT atlas and subcortical

volumes. Top: Variance due to image quality predictors. Bottom: Variance due to estimated

intra-cranial volume (eTIV).

Fig 3. R^2 from Model 2 for neocortical volumes and thicknesses from the DKT atlas and

subcortical volumes. Top: Variance due to age. Bottom: Variance due to sex. One should note

that the R^2 in model 2 cannot be compared to that of model 1 since the total variance in

model 2 is the remaining variance after model 1 (residuals).

377 **Fig 4. Difference of R^2 between training and test samples. Top: Model 1 for Image quality and**

378 **eTIV. Bottom: Model 2 for Age and sex.**

379 **Fig 5. Mean normative Z scores on cortical volumes and thicknesses from the DKT atlas and**

380 **subcortical volumes of young (18-34 years old) and older (65-92 years old) healthy**

381 **participants. Note that the scaling chosen was to be comparable to that of Fig 9.**

382 **Fig 6. Variance of the normative Z scores on cortical volumes and thicknesses from the DKT**

383 **atlas and subcortical volumes of young (18-34 years old) and older (65-92 years old) healthy**

384 **participants. Std: standard deviation. The Std is expected to be near 1.**

385 **Fig 7. Example of the four NOMIS Z scores alternatives on the left cortical thickness values of**

386 **the test sample. Note that all four alternative are adjusted for image quality.**

387 **Fig 8. Examples of the impact of contrast-to-noise ratio (CNR) and surface holes on the raw**

388 **and normed data of the test sample. Top: CNR on Left cortical thickness. Bottom: Surface**

389 **holes on left cortical volume. Left: Raw data. Right: Normed data.**

391 **Clinical validation**

392   We validated the normative values in individuals with clinically ascertained Alzheimer's

393 disease and schizophrenia, which showed expected patterns of mean deviations from otherwise

394 cognitively/behaviorally healthy individuals (Fig 9). In the Alzheimer's disease group, the mean

395 deviations from normality covered the frontal, temporal and parietal cortices with enlarged

396 ventricles, but were especially more pronounced in the hippocampus and entorhinal cortex. In

397 schizophrenia, atrophy was widespread to nearly all of the cortex. Supplementary Fig 3 displays

398 the variance of the scores in those two groups.

Fig 9. Mean normative Z scores on cortical volumes and thicknesses from the DKT atlas and subcortical volumes of participants with Alzheimer's disease and with schizophrenia.

Comparison of NOMIS and harmonization procedures

Fig 10 shows the SIMON subcortical and left cortical morphometric values (see Supplementary Fig 4 for right cortical values) before (raw) and after harmonization procedures and NOMIS normalization. Qualitatively, the variance of all measures was high before and after harmonization, as well as after NOMIS normalization. Fig 11 displays the QCD for subcortical and left cortical morphometric measures (see Supplementary Fig 5 for right cortical QCD values). QCD was highly different from a measure to another and globally harmonization procedures did not significantly lower QCD (NeuroCombat F: 1.96, p=.16; NeuroHarmonize F: 2.34, p=.13). On the other hand, NOMIS significantly reduced the QCD (F: 4.14, p = .04). Levene's tests also indicated that the harmonization procedures and NOMIS values had equivalent variance than that of the raw values, even without correction for multiple comparison (p values ranging from .26 to .99). Fig 12 shows two examples of measures (left hippocampal volume and left entorhinal thickness) across the 12 different sites and reveals that, before and after harmonization or NOMIS, there is a high variability between sites, but also within some sites.

Both harmonization procedures systematically diminished all effect sizes between CU and MCI (range: -0.01 to -0.11) and CU and AD (range: -0.09 to -0.20) groups. After NOMIS normalization, the effects sizes increased for hippocampal volumes (CU-MCI left: +0.14, right: +0.14; CU-AD: left: +0.12, right: +0.11) while it decreased for entorhinal volume (CU-MCI left: -

0.01, right: -0.02; CU-AD: left: -0.08, right: -0.09) and thickness (CU-MCI left: -0.12, right: -0.16;

CU-AD: left: -0.21, right: -0.28). These results are shown in Fig 13 (left hemisphere) and

Supplementary Fig 6 (right hemisphere). Finally, we observed (Fig. 14) that both harmonization

procedures systematically lowered the magnitude of the correlations between the six

morphometric values and episodic memory scores. NOMIS on the other hand increased the

correlations with hippocampal volumes, slightly decreased the ones with entorhinal cortices and

performed similarly to harmonization procedures for entorhinal cortices thicknesses.

Fig 10. Boxplots showing the SIMON subcortical volumes and left neocortical volumes and

thicknesses before (Raw) and after harmonization procedures (NeuroCombat and

NeuroHarmonize) and NOMIS. Boxes show the first and third quartiles with the line denoting

the median. Whiskers represent the lowest/highest datum still within 1.5 interquartile range

(IQR) of the lower/higher quartile. Right neocortical volumes and thicknesses are shown in

Supplementary Fig 4.

Fig 11. Quartile coefficient of dispersion of the SIMON subcortical volumes and left neocortical

volumes and thicknesses before (Raw) and after harmonization procedures (NeuroCombat and

NeuroHarmonize) and NOMIS. Values for the right neocortical volumes and thicknesses are

shown in Supplementary Fig 5.

Fig 12. Boxplots showing the SIMON left hippocampal volumes and right entorhinal thicknesses

before (Raw) and after harmonization procedures (NeuroCombat and NeuroHarmonize) and

NOMIS for the 12 different sites. Boxes show the first and third quartiles with the line denoting

the median. Whiskers represent the lowest/highest datum still within 1.5 interquartile range

(IQR) of the lower/higher quartile.

Fig 13. Boxplots showing the effect sizes (Cohen's d) between cognitively unimpaired (CU), mild

cognitive impairment (MCI) and Alzheimer's disease (AD) groups for the left hippocampal

volume and left entorhinal volume and thickness before (Raw) and after harmonization

procedures (NeuroCombat and NeuroHarmonize) and NOMIS. Boxes show the first and third

quartiles with the line denoting the median. Whiskers represent the lowest/highest datum still

within 1.5 interquartile range (IQR) of the lower/higher quartile. Values for the right

hemisphere are shown in Supplementary Fig 6.

Fig 14. Correlations between episodic memory score and morphometric measures before (Raw)

and after harmonization procedures (NeuroCombat and NeuroHarmonize) and NOMIS.

Discussion

Recent initiatives for morphometric normative data includes percentile fitting curves on

subcortical regions [67], deep learning-based segmentation of subcortical regions and cortical

lobes for east Asians [68], and yearly percentage of brain volume changes [69]. To our knowledge,

there is no other automated calculator for normative morphometric values available to

researchers except the one from our previous work using *FreeSurfer* 5.3

(https://github.com/medicslab/mNormsFS53). These prior normative data from our group[5-7]

were relatively limited in terms of atlases and sample size. With nearly seven thousand

participants and 1,344 brain measures, NOMIS offers a comprehensive neuromorphometric

normative tool based on a very large sample. In addition, an innovation of NOMIS is its flexibility.

Depending on the user need, it has four versions of Z-score adjusted on different sets of variables.

All versions include head size and image quality, but can also take into account age and/or sex or

without age and sex. Therefore, research groups looking for traditional norms, as well as others

wanting to lower the variance due to head size and image quality while preserving age and/or

sex variances can take advantage of NOMIS. Another strength of NOMIS is that the normative

values were created on a various amalgam of cognitively healthy participants from multiple

countries, with data acquired from a wide variety of MRI scanners and image quality, maximizing

its generalizability. A novelty to prior existing normative data, is the addition of the image quality

impact on the morphometry measures. Figures 2 shows that its effect is not trivial on cortical

volume and thickness. As shown by our results, our new normative data should help to remove

some undesirable variance due to scanners and image quality. Furthermore, the results from

NOMIS also show that in independent samples, the Z scores behaved as expected, that is with a

mean of 0 and standard deviation of 1 in healthy individuals and with marked mean deviations

targeted to the medial temporal lobes in participants with AD and throughout the cortex in

participants with SZ.

Despite these strengths, users should keep in mind that before using NOMIS, it is

mandatory to verify *FreeSurfer* segmentations and that while it will remove parts of variance due

to head size and image quality, it won't correct for segmentation errors or image artefacts.

Moreover, the normative sample, comprised essentially of research volunteers in academic-led

environments, was recruited using a non-probability sampling method and may not be

representative of the targeted population by the user.

Norms and multi-site data harmonization

The main aim of the normative values is to quantify the deviation from normality of

measurements for a new individual (i.e. one who is not in the sample used to define normality).

Because the norms remove some variance due to image quality, they can also be a useful and

simple way of lowering the noise between scanners in multi-site studies. However, norms should

not be considered an optimal technique to remove variances in multi-site studies; other

strategies are meant to specifically tackle multi-site variance. Studies should use in fact a

combination of approaches, including harmonized procedures for data acquisition, normative

values such as the one proposed herein, and post-hoc correction.

A harmonized scanning procedure, such as the Canadian Dementia Imaging Protocol

(CDIP)[70], addresses variations due to parameter and sequence dissimilarities, including quality

control and assurance, for example through scanning at all sites of an object of known geometric

and contrast properties (i.e. a "phantom") as well as human volunteers. Recent data from the

SIMON dataset [21], including non-harmonized and CDIP-harmonized scans, demonstrated how

using a harmonized protocol reduces variability across sites; however, some notable variance

remained[71, 72], (see also Figs 10-12). The idea of a harmonized protocol is however limited to

specific initiatives due to the high amount of resources it requires to implement.

Post-hoc harmonization procedures on the other hand have been developed to pool data

from different sites in large studies. Various of procedures have been proposed [16-19, 73] and

aim to lower differences in morphometric data between sites by generally applying scaling

corrections based on differences in the morphometric data themselves. The scaling corrections

are applicable for the sites/scanners included in the analysis and not for future sites/scanners or

data. This makes such post-hoc correction analysis-specific and needs to be conducted each time

some data are removed or added to an analysis. Such an approach can be very useful for large

multi-centric studies but is not applicable for generating normative values aiming to be applied

on future data. It is also vulnerable to selection bias since the scaling factors are not based on the

images or scanner characteristics, but on the difference of data between sites/scanners[18, 19].

Thus, distinct characteristics of the participants at a given site can affect the scaling factors and

post-hoc scaling factors should be used when the aim of a study is not vulnerable to sources of

variance between sites that are not related to image acquisition.

We compared NOMIS values to two post-hoc harmonization procedures, namely

NeuroCombat[16] and NeuroHarmonize[17] and while globally NOMIS slightly lowered the

variance of the values from the same individuals originating from 12 different scanners, these

two procedures were worse than NOMIS and did not significantly reduce true variance induced

by different scanners. We also verified effect sizes of well-established effects in MCI and AD

participants and once again the harmonization procedures were either similar or worse than

NOMIS. NeuroCombat and NeuroHarmonize systematically lowered the morphometric

differences between CU, MCI and AD participants while NOMIS lowered the entorhinal volume

and thickness effect sizes and increased the hippocampal volume differences between these

groups. These results suggest that caution should be exercised when using post-hoc

harmonization; the use of a calibration technique (e.g. repeated scans of human volunteers as

part of the study) is strongly encouraged.

Using NOMIS

The NOMIS tool is a user-friendly automated script in Python, freely accessible

(https://github.com/medicslab/NOMIS). Users only need to pre-process their images with

FreeSurfer 6.0 using automated directive parameters, then specify the individuals' characteristics

to the script, which will automatically compute Z-scores based on the *FreeSurfer* output. One can

choose the version of the Z-score by including in the csv file only the variables that need to be

adjusted and the script automatically selects the appropriate version of predictors. The predictive

models and all statistical parameters are provided along with the script. We anticipate that this

tool will be of broad interest to the neuroscientific community.

Financial Disclosure Statement

OP and LD are supported by a grant from the Canadian Institutes of Health Research (#IC119923). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgments

This study comprises multiple samples of healthy individuals. We wish to thank all principal investigators who collected these datasets and agreed to let them accessible.

Autism Brain Imaging Data Exchange (ABIDE): Primary support for the work by Adriana Di Martino was provided by the NIMH (K23MH087770) and the Leon Levy Foundation. Primary support for the work by Michael P. Milham and the INDI team was provided by gifts from Joseph P. Healy and the Stavros Niarchos Foundation to the Child Mind Institute, as well as by an NIMH award to MPM (R03MH096321). http://fcon_1000.projects.nitrc.org/indi/abide/

Alzheimer's Disease Neuroimaging Initiative (ADNI): The investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf. ADNI was funded by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01

AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI

is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and

Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer's

Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen;

Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli

Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company

Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.;Janssen Alzheimer Immunotherapy

Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development

LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx Research;

Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging;

Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes

of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector

contributions are facilitated by the Foundation for the National Institutes of Health

(www.fnih.org). The grantee organization is the Northern California Institute for Research and

Education, and the study is coordinated by the Alzheimer's Therapeutic Research Institute at

the University of Southern California. ADNI data are disseminated by the Laboratory for Neuro

Imaging at the University of Southern California.

http://adni.loni.usc.edu/

Australian Imaging Biomarkers and Lifestyle flagship study of ageing (AIBL): Part of the data

used in this study was obtained from the Australian Imaging Biomarkers and Lifestyle flagship

study of ageing (AIBL) funded by the Commonwealth Scientific and Industrial Research

Organisation (CSIRO) which was made available at the ADNI database

(www.loni.usc.edu/ADNI). The AIBL researchers contributed data but did not participate in

analysis or writing of this report. AIBL researchers are listed at www.aibl.csiro.au

Berlin Mind and Brain (Margulies, Villringer) CoRR sample (BMB). Zuo, X.N., et al. (2014). An

open science resource for establishing reliability and reproducibility in functional connectomics.

Scientific data, 1, 140049. doi: 10.1038/sdata.2014.49.

http://fcon_1000.projects.nitrc.org/indi/CoRR/html/bmb_1.html

Cambridge Centre for Ageing and Neuroscience (CamCAN): CamCAN funding was provided by the

UK Biotechnology and Biological Sciences Research Council (grant number BB/H008217/1),

together with support from the UK Medical Research Council and University of Cambridge, UK.

http://www.mrc-cbu.cam.ac.uk/datasets/camcan/

Center of Biomedical Research Excellence (COBRE): The imaging data and phenotypic

information was collected and shared by the Mind Research Network and the University of New

Mexico funded by a National Institute of Health COBRE: 1P20RR021938-01A2.

http://fcon_1000.projects.nitrc.org/indi/retro/cobre.html

Cleveland Clinic (Cleveland CCF): Funded by the National Multiple Sclerosis Society.

http://fcon_1000.projects.nitrc.org/indi/retro/ClevelandCCF.html

Comprehensive Assessment of Neurodegeneration and Dementia (COMPASS-ND) study: The

COMPASS-ND study is conducted by the Canadian Consortium on Neurodegeneration in Aging

(CCNA; www.ccna-ccnv.ca). The CCNA is supported by a grant from the Canadian Institutes of

Health Research (CIHR) with funding from several partners.

Consortium for the Early Identification of Alzheimer's Disease (CIMA-Q): Part of the data used in

this article were obtained from the Consortium pour l'identification précoce de la maladie

Alzheimer - Québec (CIMA-Q). As such, the investigators within the CIMA-Q contributed to the

design, the implementation, the acquisition of clinical, cognitive, and neuroimaging data and

biological samples. A list of the CIMA-Q investigators is available on cima-q.ca. CIMA-Q was

funded in 2013 with a $2,500,000 grant from the Fonds d'Innovation Pfizer - Fond de Recherche

Québec – Santé sur la maladie d'Alzheimer et les maladies apparentées.

Dallas Lifespan Brain Study (DLBS): This study is supported by the Center for Vital Longevity, the

University of Texas at Dallas, the University of Texas Southwestern Medical Center, the National

Institutes of Health and Aging, AVID Radiopharmaceuticals, the Aging Mind Foundation and the

Alzheimer's Association. http://fcon_1000.projects.nitrc.org/indi/retro/dlbs.html

FIND lab sample. Funded by the Dana Foundation; John Douglas French Alzheimer's

Foundation; National Institutes of Health (AT005733, HD059205,HD057610, NS073498,

NS058899). http://fcon_1000.projects.nitrc.org/indi/retro/find_stanford.html

F.M. Kirby Research Center neuroimaging reproducibility data (KIRBY-21). Landman, B.A. et al.

"Multi-Parametric Neuroimaging Reproducibility: A 3T Resource Study", NeuroImage. (2010)

NIHMS/PMC:252138 doi:10.1016/j.neuroimage.2010.11.047

https://www.nitrc.org/projects/multimodal

Minimal Interval Resonance Imaging in Alzheimer's Disease (MIRIAD): The MIRIAD investigators

did not participate in analysis or writing of this report. The MIRIAD dataset is made available

through the support of the UK Alzheimer's Society (RF116). The original data collection was

funded through an unrestricted educational grant from GlaxoSmithKline (6GKC).

http://miriad.drc.ion.ucl.ac.uk

National Alzheimer's Coordinating Center (NACC): The NACC database is funded by NIA/NIH

Grant U01 AG016976. NACC data are contributed by the NIA-funded ADCs: P30 AG019610 (PI

Eric Reiman, MD), P30 AG013846 (PI Neil Kowall, MD), P30 AG062428-01 (PI James Leverenz,

MD) P50 AG008702 (PI Scott Small, MD), P50 AG025688 (PI Allan Levey, MD, PhD), P50

AG047266 (PI Todd Golde, MD, PhD), P30 AG010133 (PI Andrew Saykin, PsyD), P50 AG005146

(PI Marilyn Albert, PhD), P30 AG062421-01 (PI Bradley Hyman, MD, PhD), P30 AG062422-01 (PI

Ronald Petersen, MD, PhD), P50 AG005138 (PI Mary Sano, PhD), P30 AG008051 (PI Thomas

Wisniewski, MD), P30 AG013854 (PI Robert Vassar, PhD), P30 AG008017 (PI Jeffrey Kaye, MD),

P30 AG010161 (PI David Bennett, MD), P50 AG047366 (PI Victor Henderson, MD, MS), P30

AG010129 (PI Charles DeCarli, MD), P50 AG016573 (PI Frank LaFerla, PhD), P30 AG062429-01(PI

James Brewer, MD, PhD), P50 AG023501 (PI Bruce Miller, MD), P30 AG035982 (PI Russell

Swerdlow, MD), P30 AG028383 (PI Linda Van Eldik, PhD), P30 AG053760 (PI Henry Paulson, MD,

PhD), P30 AG010124 (PI John Trojanowski, MD, PhD), P50 AG005133 (PI Oscar Lopez, MD), P50

AG005142 (PI Helena Chui, MD), P30 AG012300 (PI Roger Rosenberg, MD), P30 AG049638 (PI

Suzanne Craft, PhD), P50 AG005136 (PI Thomas Grabowski, MD), P30 AG062715-01 (PI Sanjay

Asthana, MD, FRCP), P50 AG005681 (PI John Morris, MD), P50 AG047270 (PI Stephen

Strittmatter, MD, PhD). https://www.alz.washington.edu/

National Database for Autism Research (NDAR): Data were obtained from the National Institute

of Mental Health (NIMH) Data Archive (NDA). NDA is a collaborative informatics system created

by the National Institutes of Health to provide a national resource to support and accelerate

research in mental health. Dataset identifier: http://dx.doi.org/10.15154/1520138. This

manuscript reflects the views of the authors and may not reflect the opinions or views of the

NIH or of the Submitters submitting original data to NDA. http://nda.nih.gov

Nathan Kline Institute Rockland (NKI-R) sample (NKI-RS) and Enhanced Sample (NKI-RES):

Principal support for the NKI-RES project is provided by the NIMH BRAINS R01MH094639-01.

Funding for key personnel also provided in part by the New York State Office of Mental Health

and Research Foundation for Mental Hygiene. Funding for the decompression and

augmentation of administrative and phenotypic protocols provided by a grant from the Child

Mind Institute (1FDN2012-1). Additional personnel support provided by the Center for the

Developing Brain at the Child Mind Institute, as well as NIMH R01MH081218, R01MH083246,

and R21MH084126. Project support also provided by the NKI Center for Advanced Brain

Imaging (CABI), the Brain Research Foundation, the Stavros Niarchos Foundation and the NIH

P50 MH086385-S1 (NKI-RS). http://fcon_1000.projects.nitrc.org/indi/pro/nki.html

http://fcon_1000.projects.nitrc.org/indi/enhanced/

Open access series of imaging studies (OASIS): The OASIS project was funded by grants P50

AG05681, P01 AG03991, R01 AG021910, P50 MH071616, U24 RR021382, and R01 MH56584.

http://www.oasis-brains.org/

POWER: This database was supported by NIH R21NS061144 R01NS32979 R01HD057076

U54MH091657 K23DC006638 P50 MH71616 P60 DK020579-31 , McDonnell Foundation

Collaborative Action Award, NSF IGERT DGE-0548890, Simon's Foundation Autism Research

Initiative grant, Burroughs Wellcome Fund, Charles A. Dana Foundation, Brooks Family Fund,

Tourette Syndrome Association, Barnes-Jewish Hospital Foundation, McDonnell Center for

Systems Neuroscience, Alvin J. Siteman Cancer Center, American Hearing Research Foundation

grant, Diabetes Research and Training Center at Washington University grant.

http://fcon_1000.projects.nitrc.org/indi/retro/Power2012.html

Parkinson's Progression Markers Initiative (PPMI): PPMI – a public-private partnership – is

funded by the Michael J. Fox Foundation for Parkinson's Research and funding partners,

including Abbvie, Allergan, Amathus, Avid Radiopharmaceuticals, Biogen Idec, BioLegend,

Bristol-Myers, Celgene, Cenali, Covance, GE Healthcare, Genentech, GlaxoSmithKline, Glolub

Capital, Handl Therapeutics, Insitro, Janssen Neuroscience, Eli Lilly and Company, Lundbeck,

Merck, Meso Scale Discovery, Neurocrine, Pfizer, Piramal, Prevail, Roche, Sanofi Genzyme,

Servier, Takeda, Teva, UCB, Verily, and Voyager Therapeutics. See http://www.ppmi-info.org for

further details.

Southwest University Adult Lifespan Dataset (SALD): SALD was supported by the National

Natural Science Foundation of China (31470981; 31571137; 31500885), National Outstanding

young people plan, the Program for the Top Young Talents by Chongqing, the Fundamental

Research Funds for the Central Universities (SWU1509383,SWU1509451,SWU1609177), Natural

Science Foundation of Chongqing (cstc2015jcyjA10106), Fok Ying Tung Education Foundation

(151023) , General Financial Grant from the China Postdoctoral Science Foundation

(2015M572423, 2015M580767), Special Funds from the Chongqing Postdoctoral Science

Foundation (Xm2015037, Xm2016044), Key research for Humanities and social sciences of

Ministry of Education (14JJD880009). http://fcon_1000.projects.nitrc.org/indi/retro/sald.html

University of Wisconsin, Madison (Birn, Prabhakaran, Meyerand) CoRR sample (UWM): Zuo,

X.N., et al. (2014). An open science resource for establishing reliability and reproducibility in

functional connectomics. *Scientific data, 1*, 140049. doi: 10.1038/sdata.2014.49

http://fcon_1000.projects.nitrc.org/indi/CoRR/html/uwm_1.html

Wayne State EF Dataset: This dataset was supported by National Institute on Aging grants R01-

AG011230, R37-AG011230, R03-AG024630 to Naftali Raz, Ph.D.

http://fcon_1000.projects.nitrc.org/indi/retro/wayne_EF.html

Yale Low-Resolution Controls Dataset: Scheinost D, Tokoglu F, Shen X, Finn ES, Noble S,

Papademetris X, Constable RT. Fluctuations in Global Brain Activity Are Associated With

Changes in Whole-Brain Connectivity of Functional Networks. IEEE Trans Biomed Eng. 2016

Dec;63(12):2540-2549. Epub 2016 Aug 16.

http://fcon_1000.projects.nitrc.org/indi/retro/yale_lowres.html

References

1.   Pfefferbaum A, Rohlfing T, Rosenbloom MJ, Chu W, Colrain IM, Sullivan EV. Variation in longitudinal trajectories of regional brain volumes of healthy men and women (ages 10 to 85 years) measured with atlas-based parcellation of MRI. Neuroimage. 2013;65:176-93. doi: 10.1016/j.neuroimage.2012.10.008. PubMed PMID: 23063452; PubMed Central PMCID: PMC3516371.

2.   Crivello F, Tzourio-Mazoyer N, Tzourio C, Mazoyer B. Longitudinal assessment of global and regional rate of grey matter atrophy in 1,172 healthy older adults: modulation by sex and age. PLoS One. 2014;9(12):e114478. doi: 10.1371/journal.pone.0114478. PubMed PMID: 25469789; PubMed Central PMCID: PMC4255026.

3.   Govindarajan KA, Freeman L, Cai C, Rahbar MH, Narayana PA. Effect of intrinsic and extrinsic factors on global and regional cortical thickness. PLoS One. 2014;9(5):e96429. doi: 10.1371/journal.pone.0096429. PubMed PMID: 24789100; PubMed Central PMCID: PMCPMC4008620.

4.   Kruggel F, Turner J, Muftuler LT. Impact of scanner hardware and imaging protocol on image quality and compartment volume precision in the ADNI cohort. Neuroimage. 2010;49(3):2123-33. doi: 10.1016/j.neuroimage.2009.11.006. PubMed PMID: 19913626; PubMed Central PMCID: PMC2951115.

5.   Potvin O, Dieumegarde L, Duchesne S. Normative morphometric data for cerebral cortical areas over the lifetime of the adult human brain. Neuroimage. 2017;156:315-39. doi: 10.1016/j.neuroimage.2017.05.019. PubMed PMID: 28512057.

6.   Potvin O, Dieumegarde L, Duchesne S. Freesurfer cortical normative data for adults

using Desikan-Killiany-Tourville and ex vivo protocols. Neuroimage. 2017;156:43-64. doi:

10.1016/j.neuroimage.2017.04.035. PubMed PMID: 28479474.

7.   Potvin O, Mouiha A, Dieumegarde L, Duchesne S. Normative data for subcortical

regional volumes over the lifetime of the adult human brain. Neuroimage. 2016;137:9-20. doi:

10.1016/j.neuroimage.2016.05.016. PubMed PMID: 27165761.

8.   Potvin O, Mouiha A, Dieumegarde L, Duchesne S, Alzheimers Disease Neuroimaging I.

FreeSurfer subcortical normative data. Data Brief. 2016;9:732-6. doi:

10.1016/j.dib.2016.10.001. PubMed PMID: 27830169; PubMed Central PMCID:

PMCPMC5094268.

9.   Desikan RS, Segonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, et al. An automated

labeling system for subdividing the human cerebral cortex on MRI scans into gyral based

regions of interest. Neuroimage. 2006;31(3):968-80. doi: 10.1016/j.neuroimage.2006.01.021.

PubMed PMID: 16530430.

10.   Klein A, Tourville J. 101 labeled brain images and a consistent human cortical labeling

protocol. Front Neurosci. 2012;6:171. doi: 10.3389/fnins.2012.00171. PubMed PMID:

23227001; PubMed Central PMCID: PMC3514540.

11.   Klapwijk ET, van de Kamp F, van der Meulen M, Peters S, Wierenga LM. Qoala-T: A

supervised-learning tool for quality control of FreeSurfer segmented MRI data. Neuroimage.

2019;189:116-29. doi: 10.1016/j.neuroimage.2019.01.014. PubMed PMID: 30633965.

12.   Backhausen LL, Herting MM, Buse J, Roessner V, Smolka MN, Vetter NC. Quality Control

of Structural MRI Images Applied Using FreeSurfer-A Hands-On Workflow to Rate Motion

Artifacts. Front Neurosci. 2016;10:558. doi: 10.3389/fnins.2016.00558. PubMed PMID:

27999528; PubMed Central PMCID: PMCPMC5138230.

13.   Destrieux C, Fischl B, Dale A, Halgren E. Automatic parcellation of human cortical gyri

and sulci using standard anatomical nomenclature. Neuroimage. 2010;53(1):1-15. doi:

10.1016/j.neuroimage.2010.06.010. PubMed PMID: 20547229; PubMed Central PMCID:

PMCPMC2937159.

14.   Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C, et al. Whole brain

segmentation: automated labeling of neuroanatomical structures in the human brain. Neuron.

2002;33(3):341-55. PubMed PMID: 11832223.

15.   Augustinack JC, Magnain C, Reuter M, van der Kouwe AJ, Boas D, Fischl B. MRI

parcellation of ex vivo medial temporal lobe. Neuroimage. 2014;93 Pt 2:252-9. doi:

10.1016/j.neuroimage.2013.05.053. PubMed PMID: 23702414; PubMed Central PMCID:

PMCPMC3883990.

16.   Fortin JP, Cullen N, Sheline YI, Taylor WD, Aselcioglu I, Cook PA, et al. Harmonization of

cortical thickness measurements across scanners and sites. Neuroimage. 2018;167:104-20.

Epub 2017/11/21. doi: 10.1016/j.neuroimage.2017.11.024. PubMed PMID: 29155184; PubMed

Central PMCID: PMCPMC5845848.

17.   Pomponio R, Erus G, Habes M, Doshi J, Srinivasan D, Mamourian E, et al. Harmonization

of large MRI datasets for the analysis of brain imaging patterns throughout the lifespan.

Neuroimage. 2020;208:116450. Epub 2019/12/11. doi: 10.1016/j.neuroimage.2019.116450.

PubMed PMID: 31821869; PubMed Central PMCID: PMCPMC6980790.

18.   Pinto MS, Paolella R, Billiet T, Van Dyck P, Guns PJ, Jeurissen B, et al. Harmonization of

Brain Diffusion MRI: Concepts and Methods. Front Neurosci. 2020;14:396. Epub 2020/05/22.

doi: 10.3389/fnins.2020.00396. PubMed PMID: 32435181; PubMed Central PMCID:

PMCPMC7218137.

19.   Dinsdale NK, Jenkinson M, Namburete AIL. Deep learning-based unlearning of dataset

bias for MRI harmonisation and confound removal. Neuroimage. 2021;228:117689. Epub

2021/01/02. doi: 10.1016/j.neuroimage.2020.117689. PubMed PMID: 33385551; PubMed

Central PMCID: PMCPMC7903160.

20.   Trefler A, Sadeghi N, Thomas AG, Pierpaoli C, Baker CI, Thomas C. Impact of time-of-day

on brain morphometric measures derived from T1-weighted magnetic resonance imaging.

Neuroimage. 2016;133:41-52. Epub 2016/02/28. doi: 10.1016/j.neuroimage.2016.02.034.

PubMed PMID: 26921714; PubMed Central PMCID: PMCPMC5602560.

21.   Duchesne S, Dieumegarde L, Chouinard I, Farokhian F, Badhwar A, Bellec P, et al.

Structural and functional multi-platform MRI series of a single human volunteer over more than

fifteen years. Scientific data. 2019;6(1):245. Epub 2019/11/02. doi: 10.1038/s41597-019-0262-

8. PubMed PMID: 31672977; PubMed Central PMCID: PMCPMC6823440.

22.   Sheikh JI, Yesavage JA. Geriatric Depression Scale (GDS): Recent evidence and

development of a shorter version. Clinical Gerontology: a Guide to Assessment and

Intervention. New York: The Haworth Press; 1986. p. 165-73.

23.   Di Martino A, Yan CG, Li Q, Denio E, Castellanos FX, Alaerts K, et al. The autism brain

imaging data exchange: towards a large-scale evaluation of the intrinsic brain architecture in

autism. Mol Psychiatry. 2014;19(6):659-67. Epub 2013/06/19. doi: 10.1038/mp.2013.78.

PubMed PMID: 23774715; PubMed Central PMCID: PMCPMC4162310.

24.   Mueller SG, Weiner MW, Thal LJ, Petersen RC, Jack CR, Jagust W, et al. Ways toward an

early diagnosis in Alzheimer's disease: The Alzheimer's Disease Neuroimaging Initiative (ADNI).

Alzheimers Dement. 2005;1(1):55-66. PubMed PMID: 17476317.

25.   Ellis KA, Bush AI, Darby D, De Fazio D, Foster J, Hudson P, et al. The Australian Imaging,

Biomarkers and Lifestyle (AIBL) study of aging: methodology and baseline characteristics of

1112 individuals recruited for a longitudinal study of Alzheimer's disease. Int Psychogeriatr.

2009;21(4):672-87. doi: 10.1017/S1041610209009405. PubMed PMID: 19470201.

26.   Zuo XN, Anderson JS, Bellec P, Birn RM, Biswal BB, Blautzik J, et al. An open science

resource for establishing reliability and reproducibility in functional connectomics. Scientific

data. 2014;1:140049. doi: 10.1038/sdata.2014.49. PubMed PMID: 25977800; PubMed Central

PMCID: PMC4421932.

27.   Rohr CS, Okon-Singer H, Craddock RC, Villringer A, Margulies DS. Affect and the brain's

functional organization: a resting-state connectivity approach. PLoS One. 2013;8(7):e68015.

Epub 2013/08/13. doi: 10.1371/journal.pone.0068015. PubMed PMID: 23935850; PubMed

Central PMCID: PMCPMC3720669.

28.   Shafto MA, Tyler LK, Dixon M, Taylor JR, Rowe JB, Cusack R, et al. The Cambridge Centre

for Ageing and Neuroscience (Cam-CAN) study protocol: a cross-sectional, lifespan,

multidisciplinary examination of healthy cognitive ageing. BMC Neurol. 2014;14:204. Epub

2014/11/22. doi: 10.1186/s12883-014-0204-1. PubMed PMID: 25412575; PubMed Central

PMCID: PMCPMC4219118.

29.   Taylor JR, Williams N, Cusack R, Auer T, Shafto MA, Dixon M, et al. The Cambridge

Centre for Ageing and Neuroscience (Cam-CAN) data repository: Structural and functional MRI,

MEG, and cognitive data from a cross-sectional adult lifespan sample. Neuroimage. 2017;144(Pt

B):262-9. Epub 2015/09/17. doi: 10.1016/j.neuroimage.2015.09.018. PubMed PMID: 26375206;

PubMed Central PMCID: PMCPMC5182075.

30.   Mayer AR, Ruhl D, Merideth F, Ling J, Hanlon FM, Bustillo J, et al. Functional imaging of

the hemodynamic sensory gating response in schizophrenia. Hum Brain Mapp.

2013;34(9):2302-12. Epub 2012/03/31. doi: 10.1002/hbm.22065. PubMed PMID: 22461278;

PubMed Central PMCID: PMCPMC4020570.

31.   Beall EB, Lowe MJ. Isolating physiologic noise sources with independently determined

spatial measures. Neuroimage. 2007;37(4):1286-300. doi: 10.1016/j.neuroimage.2007.07.004.

PubMed PMID: 17689982.

32.   Belleville S, LeBlanc AC, Kergoat MJ, Calon F, Gaudreau P, Hebert SS, et al. The

Consortium for the early identification of Alzheimer's disease-Quebec (CIMA-Q). Alzheimers

Dement (Amst). 2019;11:787-96. Epub 2019/12/04. doi: 10.1016/j.dadm.2019.07.003. PubMed

PMID: 31788534; PubMed Central PMCID: PMCPMC6880140.

33.   Dallas Lifespan Brain Study (DLBS). Available from:

http://fcon_1000.projects.nitrc.org/indi/retro/dlbs.html.

34.   Shirer WR, Ryali S, Rykhlevskaia E, Menon V, Greicius MD. Decoding subject-driven

cognitive states with whole-brain connectivity patterns. Cereb Cortex. 2012;22(1):158-65. Epub

2011/05/28. doi: 10.1093/cercor/bhr099. PubMed PMID: 21616982; PubMed Central PMCID:

PMCPMC3236795.

35. Keator DB, van Erp TGM, Turner JA, Glover GH, Mueller BA, Liu TT, et al. The Function

Biomedical Informatics Research Network Data Repository. Neuroimage. 2016;124(Pt B):1074-

9. Epub 2015/09/15. doi: 10.1016/j.neuroimage.2015.09.003. PubMed PMID: 26364863;

PubMed Central PMCID: PMCPMC4651841.

36. Bookheimer SY, Salat DH, Terpstra M, Ances BM, Barch DM, Buckner RL, et al. The

Lifespan Human Connectome Project in Aging: An overview. Neuroimage. 2019;185:335-48.

Epub 2018/10/18. doi: 10.1016/j.neuroimage.2018.10.009. PubMed PMID: 30332613; PubMed

Central PMCID: PMCPMC6649668.

37. Mazziotta J, Toga A, Evans A, Fox P, Lancaster J, Zilles K, et al. A probabilistic atlas and

reference system for the human brain: International Consortium for Brain Mapping (ICBM).

Philos Trans R Soc Lond B Biol Sci. 2001;356(1412):1293-322. PubMed PMID: 11545704.

38. IXI Dataset. Available from: http://brain-development.org/ixi-dataset/.

39. Landman BA, Huang AJ, Gifford A, Vikram DS, Lim IA, Farrell JA, et al. Multi-parametric

neuroimaging reproducibility: a 3-T resource study. Neuroimage. 2011;54(4):2854-66. doi:

10.1016/j.neuroimage.2010.11.047. PubMed PMID: 21094686; PubMed Central PMCID:

PMC3020263.

40. Malone IB, Cash D, Ridgway GR, MacManus DG, Ourselin S, Fox NC, et al. MIRIAD--Public

release of a multiple time point Alzheimer's MR imaging dataset. Neuroimage. 2013;70:33-6.

Epub 2013/01/01. doi: 10.1016/j.neuroimage.2012.12.044. PubMed PMID: 23274184; PubMed

Central PMCID: PMCPMC3809512.

41.   Beekly DL, Ramos EM, van Belle G, Deitrich W, Clark AD, Jacka ME, et al. The National

Alzheimer's Coordinating Center (NACC) Database: an Alzheimer disease database. Alzheimer

Dis Assoc Disord. 2004;18(4):270-7. PubMed PMID: 15592144.

42.   Hall D, Huerta MF, McAuliffe MJ, Farber GK. Sharing heterogeneous data: the national

database for autism research. Neuroinformatics. 2012;10(4):331-9. Epub 2012/05/25. doi:

10.1007/s12021-012-9151-4. PubMed PMID: 22622767; PubMed Central PMCID:

PMCPMC4219200.

43.   Nooner KB, Colcombe SJ, Tobe RH, Mennes M, Benedict MM, Moreno AL, et al. The NKI-

Rockland Sample: A Model for Accelerating the Pace of Discovery Science in Psychiatry. Front

Neurosci. 2012;6:152. doi: 10.3389/fnins.2012.00152. PubMed PMID: 23087608; PubMed

Central PMCID: PMC3472598.

44.   Marcus DS, Wang TH, Parker J, Csernansky JG, Morris JC, Buckner RL. Open Access Series

of Imaging Studies (OASIS): Cross-sectional MRI Data in Young, Middle Aged, Nondemented,

and Demented Older Adults. J Cogn Neurosci. 2007;19(9):1498-507. PubMed PMID: 17714011.

45.   Power JD, Barnes KA, Snyder AZ, Schlaggar BL, Petersen SE. Spurious but systematic

correlations in functional connectivity MRI networks arise from subject motion. Neuroimage.

2012;59(3):2142-54. doi: 10.1016/j.neuroimage.2011.10.018. PubMed PMID: 22019881;

PubMed Central PMCID: PMC3254728.

46.   Parkinson Progression Marker I. The Parkinson Progression Marker Initiative (PPMI).

Prog Neurobiol. 2011;95(4):629-35. Epub 2011/09/21. doi: 10.1016/j.pneurobio.2011.09.005.

PubMed PMID: 21930184.

47.   Wei D, Zhuang K, Ai L, Chen Q, Yang W, Liu W, et al. Structural and functional brain scans

from the cross-sectional Southwest University adult lifespan dataset. Scientific data.

2018;5:180134. Epub 2018/07/18. doi: 10.1038/sdata.2018.134. PubMed PMID: 30015807;

PubMed Central PMCID: PMCPMC6049036.

48.   Moffat SD, Kennedy KM, Rodrigue KM, Raz N. Extrahippocampal contributions to age

differences in human spatial navigation. Cereb Cortex. 2007;17(6):1274-82. Epub 2006/07/22.

doi: 10.1093/cercor/bhl036. PubMed PMID: 16857855.

49.   Scheinost D, Tokoglu F, Shen X, Finn ES, Noble S, Papademetris X, et al. Fluctuations in

Global Brain Activity Are Associated With Changes in Whole-Brain Connectivity of Functional

Networks. IEEE Trans Biomed Eng. 2016;63(12):2540-9. Epub 2016/08/20. doi:

10.1109/TBME.2016.2600248. PubMed PMID: 27541328; PubMed Central PMCID:

PMCPMC5180443.

50.   Sherif T, Rioux P, Rousseau ME, Kassis N, Beck N, Adalat R, et al. CBRAIN: a web-based,

distributed computing platform for collaborative neuroimaging research. Front Neuroinform.

2014;8:54. Epub 2014/06/07. doi: 10.3389/fninf.2014.00054. PubMed PMID: 24904400;

PubMed Central PMCID: PMCPMC4033081.

51.   Augustinack JC, Huber KE, Stevens AA, Roy M, Frosch MP, van der Kouwe AJ, et al.

Predicting the location of human perirhinal cortex, Brodmann's area 35, from MRI. Neuroimage.

2013;64:32-42. doi: 10.1016/j.neuroimage.2012.08.071. PubMed PMID: 22960087; PubMed

Central PMCID: PMCPMC3508349.

52.   Iglesias JE, Van Leemput K, Bhatt P, Casillas C, Dutt S, Schuff N, et al. Bayesian

segmentation of brainstem structures in MRI. Neuroimage. 2015;113:184-95. doi:

10.1016/j.neuroimage.2015.02.065. PubMed PMID: 25776214; PubMed Central PMCID:

PMCPMC4434226.

53.   Iglesias JE, Augustinack JC, Nguyen K, Player CM, Player A, Wright M, et al. A

computational atlas of the hippocampal formation using ex vivo, ultra-high resolution MRI:

Application to adaptive segmentation of in vivo MRI. Neuroimage. 2015;115:117-37. doi:

10.1016/j.neuroimage.2015.04.042. PubMed PMID: 25936807; PubMed Central PMCID:

PMCPMC4461537.

54.   Rosas HD, Liu AK, Hersch S, Glessner M, Ferrante RJ, Salat DH, et al. Regional and

progressive thinning of the cortical ribbon in Huntington's disease. Neurology. 2002;58(5):695-

701. PubMed PMID: 11889230.

55.   Salat DH, Buckner RL, Snyder AZ, Greve DN, Desikan RS, Busa E, et al. Thinning of the

cerebral cortex in aging. Cereb Cortex. 2004;14(7):721-30. PubMed PMID: 15054051.

56.   Kuperberg GR, Broome MR, McGuire PK, David AS, Eddy M, Ozawa F, et al. Regionally

localized thinning of the cerebral cortex in schizophrenia. Arch Gen Psychiatry. 2003;60(9):878-

88. PubMed PMID: 12963669.

57.   Buckner RL, Head D, Parker J, Fotenos AF, Marcus D, Morris JC, et al. A unified approach

for morphometric and functional data analysis in young, old, and demented adults using

automated atlas-based head size normalization: reliability and validation against manual

measurement of total intracranial volume. Neuroimage. 2004;23(2):724-38. doi:

10.1016/j.neuroimage.2004.06.018. PubMed PMID: 15488422.

58.   Hastie T, Tibshirani R, Friedman J. The elements of statistical learning. Data mining,

inference, and prediction. Statistics, editor: Springer; 2008. 745 p.

59.   R Development Core Team. R: A language and environment for statistical computing.

Vienna, Austria: R Foundation for Statistical Computing. http://www.R-project.org2010.

60.   Grömping U. Relative Importance for Linear Regression in R: The Package relaimpo.

Journal of Statistical Software. 2006;17(1):1-27.

61.   Mowinckel AM, Vidal-Pineiro V. Visualisation of Brain Statistics with R-packages ggseg

and ggseg3d. arXiv. 2019. doi: https://arxiv.org/abs/1912.08200.

62.   Crawford JR, Garthwaite PH, Denham AK, Chelune GJ. Using regression equations built

from summary data in the psychological assessment of the individual case: extension to

multiple regression. Psychological assessment. 2012;24(4):801-14. doi: 10.1037/a0027699.

PubMed PMID: 22449035.

63.   Rey A. L'examen clinique en psychologie. Paris: Presses Universitaires de France; 1964.

64.   Wechsler D. WMS-R Wechsler Memory Scale - Revised Manual. New York, NY: The

Psychological Corporation, Harcourt Brace Jovanovich, Inc.; 1987.

65.   Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-learn:

Machine learning in Python. Journal of machine learning research. 2011;12(Oct):2825-30.

66.   Seabold S, Perktold J, editors. Statsmodels: Econometric and statistical modeling with

python. Proceedings of the 9th Python in Science Conference; 2010 June 28 - July 3; Austin, TX.

67.   Vinke EJ, Huizinga W, Bergtholdt M, Adams HH, Steketee RME, Papma JM, et al.

Normative brain volumetry derived from different reference populations: impact on single-

subject diagnostic assessment in dementia. Neurobiol Aging. 2019;84:9-16. Epub 2019/09/07.

doi: 10.1016/j.neurobiolaging.2019.07.008. PubMed PMID: 31491596.

68.   Kim REY, Lee M, Kang DW, Wang SM, Kim NY, Lee MK, et al. Deep Learning-Based

Segmentation to Establish East Asian Normative Volumes Using Multisite Structural MRI.

Diagnostics (Basel). 2020;11(1). Epub 2020/12/31. doi: 10.3390/diagnostics11010013. PubMed

PMID: 33374745; PubMed Central PMCID: PMCPMC7824436.

69.   Battaglini M, Gentile G, Luchetti L, Giorgio A, Vrenken H, Barkhof F, et al. Lifespan

normative data on rates of brain volume changes. Neurobiol Aging. 2019;81:30-7. Epub

2019/06/18. doi: 10.1016/j.neurobiolaging.2019.05.010. PubMed PMID: 31207467.

70.   Duchesne S, Chouinard I, Potvin O, Fonov VS, Khademi A, Bartha R, et al. The Canadian

Dementia Imaging Protocol: Harmonizing National Cohorts. J Magn Reson Imaging.

2019;49(2):456-65. Epub 2019/01/13. doi: 10.1002/jmri.26197. PubMed PMID: 30635988.

71.   Potvin O, Chouinard I, Dieumegarde L, Bartha R, Bellec P, Collins DL, et al. The Canadian

Dementia Imaging Protocol: Harmonization validity for morphometry measurements.

NeuroImage Clinical. 2019;24:101943. Epub 2019/07/28. doi: 10.1016/j.nicl.2019.101943.

PubMed PMID: 31351228; PubMed Central PMCID: PMCPMC6661407.

72.   Potvin O, Khademi A, Chouinard I, Farokhian F, Dieumegarde L, Leppert I, et al.

Measurement Variability Following MRI System Upgrade. Frontiers in Neurology. 2019;10. doi:

10.3389/fneur.2019.00726.

73.   Maikusa N, Zhu Y, Uematsu A, Yamashita A, Saotome K, Okada N, et al. Comparison of

traveling-subject and ComBat harmonization methods for assessing structural brain

characteristics. Hum Brain Mapp. 2021;42(16):5278-87. Epub 2021/08/18. doi:

10.1002/hbm.25615. PubMed PMID: 34402132; PubMed Central PMCID: PMCPMC8519865.

Model 2

Age

Volume   Thickness

Subcortical volume

r^2
0.5
0.4
0.3
0.2
0.1
0.0

Sex

Volume   Thickness

Subcortical volume

r^2
0.5
0.4
0.3
0.2
0.1
0.0

Test validation (test - training)

Model 1 - Image quality and eTIV

Volume   Thickness

Subcortical volume   r²
                                              0.4
                                              0.0
                                              −0.4

Model 2 - Age and sex

Volume   Thickness

Subcortical volume   r²
                                              0.50
                                              0.25
                                              0.00
                                              −0.25
                                              −0.50

Z scores adjusted for age and sex

Young healthy controls (n=157)

Variance of Z scores adjusted for age and sex

Young healthy controls (n=157)

Image quality effects examples on the test sample

Z scores adjusted for age and sex

Alzheimer's disease (n=157)

Examples of SIMON values across the 12 sites

Episodic memory and morphometry correlations
Hippocampal volume

Entorhinal cortex volume

Entorhinal cortex thickness