# Switches, stability and reversals:
# the evolutionary history of sexual systems in fish

Susanna Pla[1#], Chiara Benvenuto[2#], Isabella Capellini[3*] & Francesc Piferrer[1*]

[1]Institut de Ciències del Mar, Spanish National Research Council (CSIC), Barcelona, Spain

[2]School of Science, Engineering and Environment, University of Salford, Salford, UK

[3]School of Biological Sciences, Queen's University Belfast, UK

[#]Joint first authors

*Correspondence: Dr. Francesc Piferrer, Institut de Ciències del Mar, Spanish National Research Council (CSIC), Passeig Marítim, 37-49, 08003 Barcelona. Tel. +34-932 309 567; E-mail: piferrer@icm.csic.es

*Correspondence: Dr. Isabella Capellini, School of Biological Sciences, Queen's University Belfast, 19 Chlorine Gardens, Belfast, BT9 5DL, Northern Ireland, United Kingdom; E-mail: I.Capellini@qub.ac.uk

ORCID #
S.P. #  0000-0001-5354-7032
C.B. # 0000-0002-8378-8168
I.C. # 0000-0001-8065-2436
F.P. # 0000-0003-0903-4736

**Abstract**

Sexual systems are highly diverse and have profound consequences for population dynamics

and resilience. Yet, little is known about how they evolved. Using phylogenetic Bayesian

modelling and a sample of 4614 species, we show that gonochorism is the likely ancestral

condition in teleost fish. While all hermaphroditic forms revert quickly to gonochorism,

protogyny and simultaneous hermaphroditism are evolutionarily more stable than protandry.

In line with theoretical expectations, simultaneous hermaphroditism does not evolve directly

from gonochorism but can evolve slowly from sequential hermaphroditism, particularly

protandry. We find support for the predictions from life history theory that protogynous, but

not protandrous, species live longer than gonochoristic species and invest the least in male

gonad mass. The distribution of teleosts' sexual systems on the tree of life does not seem to

reflect just adaptive predictions, suggesting that adaptations alone may not fully explain why

some sexual forms evolve in some taxa but not others (Williams' paradox). We propose that

future studies should incorporate mating systems, spawning behaviours, and the diversity of

sex determining mechanisms. Some of the latter might constrain the evolution of

hermaphroditism, while the non-duality of the embryological origin of teleost gonads might

explain why protogyny predominates over protandry in teleosts.

**Keywords**

Hermaphroditism, phylogeny, Dollo's law, developmental plasticity, sex determination

## Introduction

Sexual reproduction is a unifying feature of eukaryotes[1] and yet it is extremely diverse[2]. Sexual systems (also known as "sexual patterns"), defined as the pattern of distribution of the male and female function among the individuals of a given species, vary from separate fixed sexes (known as gonochorism in animals and dioecy in plants) to simultaneous hermaphroditism (each individual produces both male and female gametes at the same time). These two sexual systems can be viewed as the extremes in a sexually plastic gradient[3] of intermediate systems (sequential hermaphroditism) and mixed systems (coexistence of males and/or females with hermaphrodites)[4,5]. Sexual systems have a profound influence on individuals' mating success and fitness[6], population sex ratios and effective sizes[7], as well as colonization events and habitat use[8]. As a result, sexual systems influence the population dynamics and resilience to natural and anthropogenic stressors of ecologically and commercially important species that are often endangered or overexploited[9].

Hermaphroditism is predominant in flowering plants (angiosperms)[10], where 94% of the species have male and female sex organs in the same individual/flower, and it is widespread in invertebrates and teleost fish (the only vertebrates to exhibit hermaphroditism[11]), totalling 5% of animal species or up to ~30% if insects are excluded[12]. While this diversity suggests multiple evolutionary transitions between sexual systems in response to selection, current evolutionary models on the adaptive advantage of different sexual systems explain little about how and why sexual systems evolve and thus their large-scale distribution across the tree of life. This might indicate that adaptive predictions alone fail to fully explain why some sexual forms evolve in some taxa but not others (Williams' paradox)[4,13]. Therefore, unravelling the evolutionary history of sexual systems and quantifying how frequently and in what direction transitions occur is key to revealing which sexual systems are evolutionarily labile or stable, elucidating how one changes into another over evolutionary time, and identifying the environmental, genetic and developmental drivers favouring or opposing these changes. Yet, our understanding of how sexual systems evolve is still limited, particularly in animals.

Theoretical models, initially developed for plants, suggest that simultaneous hermaphroditism and dioecy are evolutionary stable conditions that are retained over long evolutionary time and unlikely lost once evolved, while mixed sexual systems represent evolutionary intermediate stages[4,5,14] (Fig. 1).

**Fig. 1.** Theoretical framework for the evolution of sexual systems: potential evolutionary transitions between gonochorism and simultaneous hermaphroditism via mixed systems (mixed pathways) as described in plants and some animals; via sequential hermaphroditism (sequential pathways) as recently suggested[4]; or without intermediate states (direct pathways) as proposed for plants[6]. Double-headed arrows indicate theoretical pathways.

Simultaneous hermaphroditism is likely the ancestral state in angiosperms from which dioecy, a rare sexual system in plants[6], has evolved independently several times, possibly to avoid inbreeding[15,16]. Theoretical models predict that separate sexes in plants evolve from hermaphroditism in different ways: 1) primarily through the intermediate state of gynodioecy[17], a common sexual system in plants that occurs when a male-sterile mutant invades an hermaphroditic population resulting in the coexistence of hermaphrodites and females; 2) through androdioecy, a less common system[18,19] in which mutations resulting in female sterility lead to the coexistence of hermaphrodites and males; 3) via trioecy, i.e., the coexistence of hermaphrodites, males and females, which is very rare; and 4) less frequently, via a direct transition[6,10] (Fig. 1). However, in animals no evidence of a direct transition between hermaphroditism and gonochorism exists. Once gained, dioecy was believed to be an irreversible condition[20], a conclusion based on the assumption that returning to a simultaneous expression of male- and female-specific genes would likely produce contrasting effects on sex-specific physiology. Recent studies, however, reject this claim in plants, as phylogenetic

reconstructions of direct transitions from dioecy/gonochorism to simultaneous
hermaphroditism have been documented[10,21].

The same theoretical framework with mixed pathways has been proposed also for
animals where, in contrast to plants, gonochorism is the most common sexual system,
androdioecy is more common than gynodioecy[5,14] and trioecy is very rare[22]. However, several
reproductive characteristics in plants differ substantially from those in animals[23], albeit
similarities can be found in some invertebrates[24]; hence, different theoretical frameworks are
required (Fig. 1). Furthermore, evolutionary transitions between sexual systems in teleost fish
(~34000 species, comprising the overwhelming majority of the ray-finned fishes,
Actinopterygii)[25], might be less likely to occur via a mixed pathway (Fig. 1) given that in this
group only a few killifish species of the genus *Kryptolebias* (formerly *Rivulus*) are truly
androdioecious[5,26,27]. Beyond teleosts, the presence of gynodioecy and trioecy among
vertebrates is still debated in the jawless hagfish *Myxine glutinosa* (Myxini)[14,28]. Recently,
sequential hermaphroditism has been suggested as a possible intermediary state that may
facilitate evolutionary changes between gonochorism and simultaneous hermaphroditism[4]
(Fig. 1). However, phylogenetic studies on the evolution of hermaphroditism at large scale do
not typically discriminate between the different forms of hermaphroditism and treat sexual
system as binary trait[29]. Thus, we currently have no robust large-scale study on the evolution
of sexual systems in animals and we do not know whether sequential hermaphroditism
represents an evolutionary intermediate stage between gonochorism and simultaneous
hermaphroditism, whether protogyny and protandry act equally as transitional forms between
the two, and whether gonochorism and simultaneous hermaphroditism are evolutionary stable
conditions in animals as they are in plants.

The evolution of hermaphroditism in animals has mostly been interpreted in the context
of its adaptive advantages relative to gonochorism, as proposed by the low density and the size
advantage models[30]. The former predicts that simultaneous hermaphroditism evolves under
low population densities and/or low dispersal capacity as, in these conditions, individuals with
this sexual system can maximize their chances of securing a mate compared to sex-changing
or gonochoristic individuals[31]. Note, however, that advantages of self-fertilizing simultaneous
hermaphrodites may be offset by increased risk of inbreeding. The size advantage model
proposes adaptive explanations for the evolution of sequential hermaphroditism based on the
relationship between size and fecundity[30,32–34]. Specifically, since most fish, including
sequential hermaphrodites, have indeterminate growth[35] (i.e., they can keep growing as far as
food resources and environment allow) sex change should be favoured when the reproductive

value of an individual depends on size (and thus age), and this affects in particular one of the

sexes. Thus, individuals change from a smaller first sex to a larger second sex and the direction

of sex change depends on the sex that maximizes its reproductive value with a larger size[36].

The size advantage model has been supported in crustaceans[37], molluscs[38] and teleost fish[39].

The interdependence between size, fecundity and fitness is affected by a species'

mating system, defined as the pattern of sexual interactions given the number of reproducing

males and females (Table 1). Therefore, among sequential hermaphrodites, protandry (male-

to-female sex change) is usually expected in species that reproduce in monogamous or random

pairs and where individuals switch from small males to large, highly fecund females, achieving

higher reproductive potential. Conversely, protogyny (female-to-male sex change) is usually

expected in polygynous/group-mating species, where small females become large dominant

males that monopolize females, often grouped in harems (Table 1). In both systems, cases exist

with a few individuals born directly as the second sex. Specifically, in digynic protandrous

species, primary females directly develop as such and secondary females develop from males

after sex change[40]. Likewise, in diandric protogynous species, primary males develop directly

as such whereas secondary males develop from females after sex change[41].

Crucially, life history traits underpin the formulation and assumptions of the size

advantage model. Life history theory is central to the study of sexual systems evolution since

it allows to derive clear predictions about why and when individuals should allocate energy

among different life history traits, including sexual functions, to optimize fitness[42]. However,

life history traits are surprisingly not explicitly and formally incorporated in the size advantage

model, nor tested in empirical studies[13]. Longevity, maximum size and age/size at maturity are

key life history traits because they determine individual fitness, influence demographic

parameters of populations[43] and impact on populations' genetic diversity[44]. These traits evolve

and are under several selective forces at the population level, but differences in the intensity of

selection among species can lead to large-scale diversity, thus allowing large-scale comparative

studies to inform our understanding of how and why they evolved[45]. Since sequential

hermaphrodites achieve higher fitness when reproducing as the second sex[36] (hence the

advantage of changing sex), they should, on average, benefit more than gonochoristic and

simultaneous hermaphroditic species from increased longevity (overall and/or as the second

sex in particular) or larger size (especially in protandry where females are the larger sex ad size

give fecundity advantage). In general, larger females tend to produce more eggs than smaller

ones both within and across species[46], while larger males do not necessarily increase as much

their sperm production with size. In males, larger size gives an advantage to secure

dominance/change sex (and increase fertilization rates), but not necessarily fecundity.
Alternatively, sequential hermaphrodites could mature, on average, earlier as the first sex
compared to the same sex in gonochoristic species and capitalize on reproduction as second
sex. These predictions, however, remain to be tested.

Although exceptions occur, spawning behaviour, i.e., how the two sexes interact to
release the gametes, can be broadly classified in fish as pair spawning, involving only two
individuals at the time, and group spawning, comprising large breeding groups[47] (Table 1).
Mating system and spawning behaviour together determine the intensity of direct male-male
competition and sperm competition (i.e., the competition between the sperm of two or more
males for fertilization of the same eggs), and thus certainty of paternity. Sperm competition is
a key selective force shaping male reproductive anatomy, physiology and behaviour across
diverse animal groups[48,49]. In general, sperm competition is assumed to be low in haremic
systems[39,50,51] where large dominant males can better monopolize groups of females (pair
spawning) with no or limited competition by other males[52]. Likewise, low sperm competition
is expected under monogamy. Group spawning is commonly found in promiscuous mating,
leading to intense sperm competition[53] as many males try to fertilize the eggs of multiple
females. The intensity of sperm competition has been incorporated in the size advantage
model[54] as it can play a significant role in the advantage of protogyny: changing sex from
female to male should be more advantageous when paternity assurance is high due to reduced
sperm competition[55]. Consistent with these predictions, the gonadosomatic index (GSI),
defined as the percentage of body mass devoted to the gonads[56] and a reliable indicator of the
intensity of sperm competition[57], is significantly lower in protogynous teleost species than in
gonochoristic congeners[51,52,58]. However, protandrous teleost fish do not always conform to
theoretical expectations, exhibiting higher GSI as males than expected[52]. We have recently
proposed that, at least in the family Sparidae, high male GSI in protandrous fish can be
explained not only by group spawning and high sperm competition in some species, but also
because high investment in the gonads can represent a compensatory mechanism that allows
small males to fertilize highly fecund females much larger than themselves[58].
205
206

**Table 1.** Predictions of associations between most common sexual systems (distribution of the male and female function among the individuals of a given species), mating systems (pattern of sexual interactions that take place considering the number of males and females involved in reproduction), adult sex ratio, size of mates and spawning behaviour (how the two sexes interact to release the gametes) in teleosts. This general set of predictions is applicable to most species, but exceptions are found in species with less common sex determination mechanisms and mating or spawning behaviour.

| Sexual system | Mating system | Adult sex ratio | Size of mates | Spawning behaviour |
|---|---|---|---|---|
| ♀♂ **GONOCHORISM (G)**<br>Individuals reproduce as one sex throughout their lifetime (male or female) | ♂ ↔ ♀♀♀♂♂♂ ↔ ♀♀♀<br>♂ ↔ ♀♀♀ | Variable | ♂ = ♀<br>♂ > ♀<br>♀ > ♂ | ♂♀<br>♂♂♂♀♀♀ |
| ♀→♂ **PROTOGYNY (PG)**<br>Female-first sequential hermaphroditism: individuals first reproduce as females, change sex once with increasing size/age and then reproduce as males | ♂ ↔ ♀♀♀<br>♂♂♂ ↔ ♀♀♀ | ♀ biased | ♂ > ♀ | ♂♀<br>♂♂♂♀♀♀ |
| ♂→♀ **PROTANDRY (PA)**<br>Male-first sequential hermaphroditism: individuals first reproduce as males, change sex once with increasing size/age and then reproduce as females | ♂ ↔ ♀ | ♂ biased | ♀ > ♂ | ♂♀<br>♂♂♂♀♀♀ |
| ♀↔♂ **BIDIRECTIONAL (BD)**<br>Individuals can change sex more than once, in either direction, throughout their lifespan, usually starting from PG | ♂ ↔ ♀<br>♂ ↔ ♀♀♀ | ♀ biased | ♂ > ♀ | ♂♀ |
| ♂♀ **SIMULTANEOUS (SH)**<br>Individuals produce gametes of both sexes at the same time or in a short period of time | ♂ ↔ ♀ | 1:1 | ♂ = ♀ | ♂♀ |

Mating system: ♂ ↔ ♀ = monogamy (pair bond) or random pairing; ♂♂♂ ↔ ♀♀♀ = promiscuity; ♂ ↔ ♀♀♀ = harem polygyny or temporary lek-like systems (many females in a territory defended by a male). Spawning behaviour: ♂♀ = pair spawning (pair of individuals); ♂♂♂♀♀♀ = group spawning. Symbols do not reflect bias in sex ratios.

Teleosts account for more than 50% of the extant species of vertebrates and are
characterized not only by their extraordinary diversity in morphology, physiology, ecology and
habitat, but also by different sexual systems, including gonochorism, different forms of
hermaphroditism —the only group among vertebrates— and unisexuality (all-female
populations)[11,59,60]. Hermaphroditism in teleosts is broadly divided into simultaneous
(synchronous) and sequential (consecutive) hermaphroditism, the latter in the form of
protandry, protogyny and bidirectional sex change (Table. 1). Thus, the remarkable diversity
in sexual systems in teleost fish makes them an ideal group in which to study the evolution of
different forms of hermaphroditism[29]. Here, we investigate the evolutionary origin and
transitions among sexual systems across 4614 teleost species belonging to 49 orders and 293
families using a recent time calibrated phylogeny[61] and modern phylogenetic comparative
approaches. Our large-scale approach allows us to fully unravel how sexual patterns evolved
and identify which ones represent evolutionary stable conditions. We focus on gonochorism,
protogyny, protandry and simultaneous hermaphroditism as these are the most common sexual
systems in teleosts. For hermaphrodites, we only included species for which functional
hermaphroditism could be confirmed by primary literature; all remaining species, following
the sexual system obtained from FishBase[62], were classified as gonochoristic, excluding the
species with ambiguous information (see Methods below and Supplementary Figure 1). We do
not distinguish digynic and diandric species (or populations) in this study because the number
of sequentially hermaphroditic species in our dataset is not sufficient for splitting them in
narrower categories. Thus, separating digynic and diandric species would lead to small sample
size per category while increasing the number of parameters to be estimated, ultimately eroding
power for the analysis. Likewise, unisexual species ("biotypes", hybrid in origin)[59,63] are too
few to be incorporated in any formal analyses in our study.

Here, we demonstrate that gonochorism is the likely ancestral condition in teleosts and
it is an evolutionarily stable state from which protogyny and protandry evolve at a moderate
evolutionary rate. Consistent with theoretical predictions, we show for the first time that
simultaneous hermaphroditism cannot evolve directly from gonochorism but rather through the
intermediate stage of sequential hermaphroditism, most likely protandry. Further, we expand
the theoretical framework of the evolution of hermaphroditism investigating how life history
traits and male GSI differ between sexual systems, as predicted by life history theory. In
support of these predictions, we found evidence of longer lifespan in protogynous species
compared to gonochoristic and strong evidence of smaller GSI in protogynous males. However,
contrary to predictions, we found no difference in maximum size and age or size at maturity

across sexual systems. We discuss how our results should be incorporated in a broader
framework with sex determining mechanisms and gonadal plasticity as possible constraining
and facilitating mechanisms respectively, to gain a fuller understanding of the evolution of
sexual systems and possibly resolve Williams' paradox.
255

## Results

### *Evolutionary history of sexual systems*

258 Our dataset includes 4614 extant teleost species, of which 294 are hermaphroditic
259 (protogynous: n = 196; protandrous: n = 36; bidirectional sex changers: n = 16; simultaneous
260 hermaphrodites: n = 46; Fig. 2; Supplementary Information).
261

262
263 **Fig. 2.** Sexual systems of extant species of teleosts, colour coded for gonochorism (n = 4320;
264 grey), protogyny (n = 196; red), protandry (n = 36; blue), bidirectional sex change (n = 16;
265 green) and simultaneous hermaphroditism (n = 46; yellow). Families (n = 32) with
266 hermaphroditic species are labelled. Silhouettes have been obtained from fishualize[64],
267 Phylopics (http://www.phylopic.org/) or drawn by the authors.
268

269 We used Discrete models of evolution to reconstruct the evolutionary history of sexual systems
270 using Reversible Jump (RJ) Markov chain Monte Carlo (MCMC) in *Bayes Traits* (Methods,

Phylogenetic comparative analysis). Treating sexual systems as a two-character state (gonochoristic or hermaphroditic) our analysis reveals that gonochorism is the most likely ancestral character state in teleosts (Fig. 3a-b; Supplementary Table 1) and that hermaphroditism evolves slowly from, and reverts very quickly and multiple times back to, gonochorism (Fig. 3a and 3c; Supplementary Table 1). This indicates that gonochorism is an evolutionarily stable state in teleosts.

**Fig. 3.** (a) The evolutionary history of sexual system as two-character state (gonochorism or hermaphroditism) is visually summarised with a maximum likelihood ancestral state reconstruction that best approximates results of our RJ-MCMC Multistate model. The sexual systems of extant species and their ancestors are colour coded for gonochorism (n = 4320; grey) and hermaphroditism (n = 294; magenta). (b) Density plots from RJ-MCMC Multistate models for the estimated probability of character state at the root of the phylogeny colour coded for gonochorism (mean = 66%; grey) and hermaphroditism (mean: 34%; magenta). (c) RJ-MCMC multistate posterior distributions of the transition rates from gonochorism to hermaphroditism (magenta) and from hermaphroditism to gonochorism (grey).

Treating sexual system as four-character states (gonochoristic, protandric, protogynic, simultaneous hermaphroditic; Fig. 2) reveals that both types of sequential hermaphroditism evolve at a very low rate from gonochorism and revert very rapidly back to it (Fig. 4, Table 2). In contrast, direct transitions between gonochorism and simultaneous hermaphroditism are very slow if they happen at all, given that over 60% and 31% of the models estimate the transition from gonochorism to simultaneous hermaphroditism and the reversal respectively to be equal to zero. Our analysis also shows that protogyny evolves as slowly from gonochorism as it switches to protandry and simultaneous hermaphroditism, although 32% of models estimate the latter transition to be equal to zero. Conversely, protandry is lost quickly to protogyny and simultaneous hermaphroditism, and very rapidly to gonochorism. Therefore, simultaneous hermaphroditism evolves from sequential hermaphroditism, most likely from protandry, and is lost to gonochorism, protogyny and protandry at similar low rates, although approximately half of the models estimate transition rates to sequential hermaphroditism to be equal to zero. Altogether, these results clearly indicate that in teleosts gonochorism is an evolutionarily stable state; protogyny is evolutionarily more stable than protandry, while simultaneous hermaphroditism evolves rarely, most like from protandry, and is evolutionarily stable being lost slowly to gonochorism, and less likely, to protogyny and protandry (Fig. 4, Table 2).

**Fig. 4.** Summary of RJ-MCMC Multistate analysis with density plots of the posterior distributions of the transition rates to gonochorism (grey), protogyny (red), protandry (blue), and simultaneous hermaphroditism (yellow). Gonochorism is the estimated likely ancestral condition. Note, only x-axis, but not y-axis, are the same for each pair of gain and loss between two-character states. The thickness of the arrows is roughly proportional to the mean magnitude of the transition rates from the posterior distribution. Dotted lines indicate transition rates estimated to be equal to 0 in over 40% of the models in the posterior distributions. Sample sizes of extant species for each sexual system category are indicated between parentheses.

**Table 2.** Results of the RJ-MCMC Multistate analysis in *BayesTraits* of sexual systems as a four-state categorical variable: gonochorism (G), protogyny (PG), protandry (PA) or simultaneous hermaphroditism (SH). For each posterior distribution, we report the effective sample size (ESS), the mean and 95% high posterior density intervals (95-HPD), the mode and the percentage of models in which the parameter is estimated as zero. This analysis is based on 4598 extant species (G: n = 4320; PG: n = 196; PA: n = 36; SH: n = 46). Note: 16 species of bidirectional sex change were not included in this analysis due to their low number.

| Transition rates | ESS | Mean | 95-HPD | Mode | % Zero |
|---|---|---|---|---|---|
| G → PG | 1153 | 0.014 | 0.000 – 0.023 | 0.014 | 7.4 |
| PG → G | 1600 | 1.077 | 0.804 – 1.396 | 1.116 | 0.0 |
| G → PA | 1600 | 0.015 | 0.005 – 0.025 | 0.014 | 0.0 |
| PA → G | 1143 | 1.617 | 0.760 – 4.292 | 1.105 | 0.0 |
| G → SH | 1366 | 0.004 | 0.000 – 0.014 | 0.000 | 60.7 |
| SH → G | 1600 | 0.010 | 0.000 – 0.023 | 0.000 | 31.7 |
| PG → PA | 1600 | 0.013 | 0.000 – 0.023 | 0.014 | 18.7 |
| PA → PG | 1600 | 0.976 | 0.000 – 1.321 | 1.107 | 2.9 |
| PG → SH | 1600 | 0.009 | 0.000 – 0.023 | 0.000 | 35.8 |
| SH → PG | 1600 | 0.009 | 0.000 – 0.021 | 0.000 | 51.7 |
| PA → SH | 1155 | 0.602 | 0.000 – 1.236 | 1.105 | 18.1 |
| SH → PA | 1600 | 0.008 | 0.000 – 0.021 | 0.000 | 48.6 |
| **Root probabilities** | | | | | |
| G | 1324 | 46.1 | 36.1 – 55.0 | 49.5 | 0.0 |
| PG | 1258 | 31.4 | 22.7 – 37.8 | 34.9 | 0.0 |
| PA | 1167 | 22.4 | 13.1 – 37.0 | 14.9 | 0.0 |
| SH | 1600 | <0.1 | 0.0 – <0.1 | 0 | 6.8 |

*Life history traits and sexual systems in teleosts*

Using life history theory, we predicted that sequential hermaphrodites live longer and/or reach a larger adult size and/or mature earlier as the first sex. The phylogenetic generalized least square (PGLS) analyses revealed that protogynous, but not protandrous, species live longer than gonochoristic species (Fig. 5a; Table 3). Larger species however might live longer, therefore we repeated the analysis controlling for allometry; even so, adding size (maximum length) as a covariate did not alter this result (Fig. 5b; Supplementary Table 2). Contrary to predictions, we did not find any significant size difference across sexual systems (Fig. 5c; Table 3). Female and male age at maturity does not differ across species with different sexual systems (Fig. 5d-e; Table 3), even when accounting for allometry (length at maturity; Supplementary Table 4), nor does sex-specific length at maturity (Table 3). Finally, the PGLS revealed that protogynous males have lower GSI values than gonochoristic and protandrous ones, but GSI does not differ significantly between gonochoristic and protandric males (Fig. 5f; Table 3) even when considering allometry (Supplementary Table 2).

**Fig. 5.** Life history traits by sexual system. Phylogenetic estimated mean and phylogenetic
standard error from the PGLS results of: **a)** longevity (year, $log_{10}$ transformed; G: n = 758; PG:
n = 69; PA: n = 17); **b)** longevity while controlling for maximum length (G: n = 575; PG: n =
61; PA: n = 8); **c)** maximum length (cm, $log_{10}$ transformed; G: n = 2612; PG: n = 167; PA: n =
20); **d)** male age at first maturity (year, $log_{10}$ transformed; G: n = 259; PG: n = 15; PA: n = 9);
**e)** female age at first maturity (year, $log_{10}$ transformed; G: n = 282; PG: n = 30; PA: n = 5); **f)**
male gonadosomatic index, GSI ($log_{10}$ transformed; G: n = 44; PG: n = 38; PA: n = 15). In all
panels gonochorism (G) is depicted in grey, protogyny (PG) in red and protandry (PA) in blue.
350

**Table 3.** Results of phylogenetic generalized least square (PGLS) model of longevity (year; $\log_{10}$ transformed), maximum length (cm; $\log_{10}$ transformed), age at first maturity (year; $\log_{10}$ transformed), length at first maturity (cm; $\log_{10}$ transformed) per each sex (♂: male; ♀: female), and male gonadosomatic index (GSI; $\log_{10}$ transformed) across sexual systems: gonochorism (G); protogyny (PG); protandry (PA). For each independent variable we report the parameter estimate (Beta), t-statistics (T), *P*-value (P; two-sided test), and the model statistics including the degrees of freedom (df), the maximum likelihood estimation of the phylogenetic signal ($\lambda$) and $R^2$. Significant differences are indicated in bold. Results of analyses controlling for allometry are available in Supplementary Table 2. See Supplementary Table 3 for sexual system and sex-specific sample sizes.

| Dependent | Variable Independent | Beta | T | P | Df | $\lambda$ | $R^2$ |
|---|---|---|---|---|---|---|---|
| Longevity | Sexual system - PA[1] | 0.015 | 0.128 | 0.898 | 2; 841 | 0.914 | 0.007 |
| | Sexual system - PG[1] | 0.161 | 2.340 | **0.019** | | | |
| | Sexual system - PG[2] | 0.146 | 1.205 | 0.229 | | | |
| Max length | Sexual system - PA[1] | 0.068 | 1.036 | 0.300 | 2; 2796 | 0.972 | 0.0004 |
| | Sexual system - PG[1] | -0.036 | -0.098 | 0.922 | | | |
| | Sexual system - PG[2] | -0.071 | -0.971 | 0.332 | | | |
| Age at maturity ♂ | Sexual system - PA[1] | -0.015 | -0.129 | 0.897 | 2; 280 | 0.859 | 0.002 |
| | Sexual system - PG[1] | 0.070 | 0.684 | 0.495 | | | |
| | Sexual system - PG[2] | 0.085 | 0.623 | 0.534 | | | |
| Age at maturity ♀ | Sexual system - PA[1] | 0.165 | 1.217 | 0.225 | 2; 314 | 0.862 | 0.005 |
| | Sexual system - PG[1] | 0.029 | 0.333 | 0.739 | | | |
| | Sexual system - PG[2] | -0.135 | -0.945 | 0.345 | | | |
| Length at maturity ♂ | Sexual system - PA[1] | -0.060 | -0.809 | 0.419 | 2; 359 | 0.974 | 0.002 |
| | Sexual system - PG[1] | -0.020 | -0.337 | 0.736 | | | |
| | Sexual system - PG[2] | 0.040 | 0.448 | 0.654 | | | |
| Length at maturity ♀ | Sexual system - PA[1] | -0.018 | -0.169 | 0.866 | 2; 340 | 0.971 | 0.0009 |
| | Sexual system - PG[1] | -0.041 | -0.565 | 0.572 | | | |
| | Sexual system - PG[2] | -0.023 | -0.200 | 0.842 | | | |
| GSI ♂ | Sexual system - PA[1] | 0.092 | 0.736 | 0.464 | 2; 94 | 0.835 | 0.234 |
| | Sexual system - PG[1] | -0.500 | -4.977 | **<0.001** | | | |
| | Sexual system - PG[2] | -0.592 | -4.209 | **<0.001** | | | |

[1] G as reference level; [2] PA as reference level

## Discussion

Our large-scale phylogenetic study has tested for the first time the theoretical predictions on how sexual systems evolve and has revealed the evolutionary origin of and transitions between different sexual systems in the highly diverse teleosts. We identify gonochorism and simultaneous hermaphroditism as stable conditions over evolutionary time. In support to recent theoretical models[4], our study demonstrates that simultaneous hermaphroditism is unlikely to evolve directly from gonochorism and instead requires the intermediate step of sequential hermaphroditism, most likely protandry. We find support for the predictions derived from life

history theory that protogynous species live longer than gonochoristic species but no evidence

that sequential hermaphrodites attain a larger size or mature earlier than gonochoristic species.

Finally, we find strong evidence that protogynous males invest the least in male gonad tissues

(quantified by the gonadosomatic index) relative to gonochoristic and protandric males.

Combined, these results suggest that the two forms of sequential hermaphroditism must be

treated separately in theoretical and empirical studies as protandry and protogyny are

characterized by distinct life history strategies[36], even though they both entail sex change.

Using the largest dataset ever collected with four sexual systems in teleosts, our study

reveals a complex and dynamic way through which sexual systems evolve and switch between

one another. Sequential hermaphroditism can evolve slowly from gonochorism, the ancestral

state in teleosts, but revert to gonochorism rapidly. Although gonochorism is an evolutionarily

stable condition, gained faster than it is lost, these results refute the assumption that the

transition to gonochorism is irreversible[20] and represent another example[65,66] against Dollo's

law of irreversibility[67], as previously suggested[29]. Conversely, sequential hermaphroditism in

teleosts, particularly protandry, is less evolutionarily stable than gonochorism. Our results

however contradict Pennell et al.'s finding[29] that evolutionary transition from gonochorism to

hermaphroditism occur over twice as fast than the reverse, suggesting rapid evolution of

hermaphroditism from gonochorism, a conclusion that the same authors acknowledge is

counterintuitive. We note that Pennell et al.[29] used a much smaller dataset biased towards a

greater proportion of hermaphroditic than gonochoristic species than what is observed in

teleosts, and did not discriminate between different types of hermaphroditism. In contrast, we

find that the evolutionary gain of hermaphroditism is slower than its loss to gonochorism,

regardless of whether we treat sexual system as binary trait (gonochorism vs hermaphroditism)

or discriminate between forms of hermaphroditism. Heterogeneity in the rate of gain and losses

across large phylogenies can potentially bias the estimates of the faster transition rate for binary

traits[68]. However, our analysis at four states reveals that protandry is lost rapidly to both

protogyny and gonochorism, and to a lesser degree, to simultaneous hermaphroditism.

Altogether, our results at four states indicate that rapid transition rates from hermaphroditism

to gonochorism in our analysis at two states are robust and reveal that protandry and protogyny

—but not simultaneous hermaphroditism— evolve much more slowly from gonochorism than

the reverse.

Importantly, our study demonstrates that simultaneous hermaphroditism does not

originate directly from gonochorism but rather through sequential hermaphroditism, most

likely protandry. However, simultaneous hermaphroditism is lost preferentially to

gonochorism than to either form of sequential hermaphroditism. Thus, our analyses

demonstrate that an intermediate stage is required for the gain of simultaneous

hermaphroditism from gonochorism but not the loss back to it. These results support theoretical

predictions (Fig. 1) that sex-specific gene expression in gonochoristic species may prevent

direct evolutionary transitions between gonochorism and simultaneous hermaphroditism, and

intermediate stages, like sequential hermaphroditism, are required[4]. Overall, our study is

consistent with suggestions that the complexity of sex-specific physiology and behaviour is

likely to constrain some transitions between sexual systems. Androdioecy is considered an

intermediate stage from simultaneous hermaphroditism to dioecy in plants and from

gonochorism to hermaphroditism in some invertebrates[5,14]. However, this sexual system is

extremely rare in fish and cannot explain the evolution of diverse sexual system in this

vertebrate group, where instead sequential hermaphroditism seems to play a similar role. We

suggest that future studies in other taxa may also consider sequential hermaphroditism (if

present) together with other mixed systems as an important stepping stone for evolutionary

changes between gonochorism and simultaneous hermaphroditism as we have found in

teleosts.

According to life history theory, sequential hermaphrodites should, on average, live

longer, grow bigger and/or mature earlier as the first sex than gonochoristic species. We find

that protogynous and protandrous species differ in their life history strategies: protogynous, but

not protandrous species live longer than gonochoristic species. These differences reflect the

fact that protogyny and protandry maximise their fitness as the second sex[36] which differs

between the two systems. Therefore, the longer life in protogynous species favours large

successful males (second sex) that can monopolize females in harems or in spawning grounds.

Conversely, protandrous species benefit primarily by achieving a larger size, as larger females

(second sex) are more fecund than smaller ones. In addition, male investment in gonad tissue

(as quantified by the gonadosomatic index) is lower in protogyny, as expected by theory[52,58],

since large males can better monopolize mating opportunities and face low levels of sperm

competition in harems and group spawning (Table 1). Small-sized protandrous males in group

spawning instead need to boost their investment in the gonads but even in the absence of sperm

competition (monogamy) they require large gonads to fertilize highly fecund females, larger

than themselves[58]. Thus, sexual systems and mating strategies affect life history traits

differentially in protogynous and protandrous species. It is well known that in sequential

hermaphrodites the second sex always matures later and is larger than the first sex, so it is not

surprising that in protandrous species females are significantly larger than males when reaching

maturity, while in protogynous species males are significantly larger than females[55] (excluding

the cases of primary females and primary males, respectively). Yet, no comparison has been

made for size/age at first maturity for males and females across sexual systems. Life history

theory predicts that the first sex of sequential hermaphrodites matures earlier than the same sex

in gonochorism, but, with the data currently available, we find no evidence for this.

Our study is the first to attempt to include explicitly life history traits into a theoretical

framework for the evolution of sexual systems and provide some evidence in support to

theoretical predictions, but records on life history traits for teleosts species in general and

hermaphroditic species in particular, are still currently too scarce. Even less complete and

reliable data are available on mating systems and spawning behaviours, which should be

incorporated in future studies aiming at obtaining a more complete understanding of the role

that life history traits play. Particularly necessary to fully assess theoretical predictions are sex-

specific data for size and time spent as females and as males in sequential hermaphrodites, and

for investment in male *vs* female function in simultaneous hermaphrodites, for which currently

little is known. Future studies should re-evaluate these relationships as more data become

available for a large number of species.

While we have shown that life history theory can provide a major contribution to our

understanding of sexual system evolution, below we present a general model for studying

sexual systems and propose that the highly dynamic picture revealed by this study should be

expanded using a more comprehensive approach that includes not only selection and

adaptation, but also sex determining mechanisms and gonadal plasticity (Fig. 6).

**Fig. 6.** Theoretical framework for the evolution of sexual systems: overview of parameters
(with some examples) considered in the low density and the size advantage models (*), used
in our analyses (**) and proposed in the present study (***).

Sex determination in gonochoristic animals is determined either at fertilization by
different genetic mechanisms, including male (XX/XY) or female (ZW/ZZ) heterogamety with
homomorphic or heteromorphic sex chromosomes, or after conception by environmental
factors, or by a combination of both[69,70]. Fish are characterized by an incredible diversity[71–73]
and plasticity[29,74] of sex determining mechanisms. Many fish do not have sex chromosomes[72].
When they are present, they might not always be clearly differentiated since sex determining
loci might not be easily identifiable[75] and in some cases the sex can be determined by a change
in a single nucleotide[72]. High turnover of sex chromosomes has also been detected in some fish
lineages (e.g., sticklebacks[76,77]), including reversal to autosomes[29]. It has been suggested that
fixed, strongly canalized, genetic sex determination (culminating in the formation of fully
differentiated and stable heteromorphic sex chromosomes) might constrain the evolution of
hermaphroditism, acting as an evolutionary trap[2,78–80]. Even if this is not the case[29,81], sequential
hermaphrodites do not appear to have sexually differentiated chromosomes[82], but data are

currently scarce for formal analyses. Finally, sequential hermaphroditism can be regarded as a
clear example of phenotypic plasticity, and since epigenetics underlies phenotypic plasticity,
epigenetic mechanisms have been proposed to participate in the evolutionary transitions
between different sexual systems and sex-determining mechanisms[83]. Therefore, although
complete genetic control of hermaphroditism is common in plants[84], a better knowledge of the
genetic and epigenetic mechanisms of sex determination could be helpful to explain how
hermaphroditism in teleosts has evolved in some taxonomic groups but not in others under
similar ecological pressures.

Previous attempts to connect the distribution of sexual systems have invoked
morphological[85] and developmental[86] aspects. Developmental plasticity is uniquely
documented in teleosts via the bipotential nature of their gonads and gonoducts[86]. Thus, while
in most vertebrate taxa gonads develop from two distinct germinal layers (medulla, endodermal
in origin, which gives rise to the testes; and cortex, mesodermal in origin, which gives rise to
the ovary), in teleosts the gonads consist entirely of the cortex homolog[87]. Moreover, teleosts
are the only group of vertebrates where the Müllerian duct is absent, and the gonoduct has the
same origin in both sexes, being the reproductive systems completely independent of the
excretory system[86]. Therefore, anatomically all teleosts could, in principle, be
hermaphrodites[86]. Furthermore, the transition to protogyny may be favoured by the
peculiarities of gonadal development in many gonochoristic teleost species, which develop a
female gonad, complete with ovaries containing cysts of oocytes, during the initial stage of
gonadal formation[88,89]. Only later testicular development is triggered and superimposed on this
arrangement so that the individual ultimately matures functionally as a male[90–92]. Thus,
protogyny might be favoured because female gonads are often the first to develop albeit
temporarily, even in protandrous species[92]. Bidirectional sex change, a rarer system in teleosts
(Table 1), further demonstrates the importance of gonadal plasticity. In most cases, the initial
strategy is protogyny[93], but males can revert back to females when triggered by new social
conditions. The retention of some female gonadal tissue in males facilitates a new change of
sex, if and when required[94]. The maintenance of both gonadal tissues could facilitate a
transition to simultaneous hermaphroditism. Thus, the study of the evolution of sexual systems
in fish (and possibly other taxa) could greatly benefit from taking into consideration the
facilitating/constraining aspects linked to gonadal developmental plasticity and the existence
of different sex determining mechanisms.

In conclusion, our study reveals that gonochorism is the most likely ancestral state and
the most evolutionary stable sexual system in teleosts. In support to theoretical predictions, we

demonstrate that simultaneous hermaphroditism cannot evolve directly from gonochorism but

requires an intermediate step, most likely through protandry. However, simultaneous

hermaphroditism is more likely to be lost to gonochorism than to sequential hermaphroditism

in teleosts. Overall, our study reveals that the evolution of sexual systems is evolutionarily

more dynamic and complex than commonly assumed. Our results support theoretical

assumptions that changes between sexual systems are likely constrained by sex-specific gene

expression, physiology and behaviour. In addition, we propose that the adaptive advantage of

different sexual systems is further underpinned in fish by their extraordinary and unique

developmental plasticity[95], including common and fast transitions among different sex

determining mechanisms[29,69]. Our study also reveals that different sexual systems exhibit

different life history strategies that allow species with sequential hermaphroditism to maximise

fitness as the second sex[36], particularly in protogynous species, and highlights the need for

more sex-specific life history data to gain a fuller and deeper understanding of the interplay

between life history strategies and sexual system. Altogether we propose that a comprehensive

framework that incorporates life history traits, sex determining mechanisms and gonadal

plasticity into traditional theoretical models of sexual system adaptive value will be essential

if we are to fully understand the evolution of sexual systems, their phylogenetic distribution

and their implications for conservation and management.

**Methods**

*Data collection and verification*

We compiled the most comprehensive database on sexual systems in teleosts to date.

Information on sexual system were first extracted from FishBase[62]. Next, species were classed

as hermaphroditic only if functional hermaphroditism could be confirmed by primary literature,

as recently compiled elsewhere[96] (see Supplementary Data for details). For the remaining

species, we maintained the gonochoristic classification of FishBase[62], unless recent literature

stated otherwise. Indeed, gonochorism is rarely confirmed in literature even when present, so

including as gonochoristic only species for which this sexual pattern is confirmed would

strongly bias the dataset against gonochorism, ultimately undermining the robustness of the

analyses. Importantly, species for which there is contrasting information in the literature were

discarded (Supplementary Methods). Of this database, 4614 species are included in the most

recent and largest molecular phylogeny for the class[61] (available at https://fishtreeoflife.org)

and used in this study. Altogether our final dataset included 4320 gonochoristic and 294

hermaphrodite species (Supplementary Figure 1), of which there were 196 protogynous, 36

protandrous, 16 bidirectional species and 46 simultaneous hermaphrodites. Unisexual species

were not included in the analyses, due to their extremely low number and hybrid orgin[59,63]; we

also did not have enough data (and power) to consider separately digynic and diandric species.

Life history traits (Supplementary Table 3) were also collected from primary literature,

FishBase[62] and *rFishBase*[97]: longevity (in years), maximum length (in cm); length (in cm) and

age at maturity (in years) of males and females; male gonadosomatic index (GSI; the maximum

value recorded, expected to coincide with the peak of the reproductive season).

### *Phylogenetic comparative analyses*

We investigated the evolutionary history of sexual systems of 4614 teleost species using

Multistate models in *BayesTraits* V.3[98,99] in a Bayesian framework. Multistate estimates

instantaneous transition rates between alternative character states of a single categorical

variable (i.e., the rate of change between states along the branches of a phylogeny), based on a

continuous-time Markov model of evolution for discrete traits[100,101]. A high transition rate from

one state to another indicates that the first state changes rapidly to the second state over

evolutionary time. Therefore, a character state is evolutionarily stable when it is lost more

slowly than it is gained[102]. Multistate also produces posterior distributions of the ancestral

character state at the root of the phylogeny. We scaled the tree by a default constant (mean of

0.1) in all analyses[102] and used an exponential prior whose mean was seeded from a uniform

hyperprior ranging from 0 to 10 to reduce inherent uncertainty and biases of prior choice[99]. We

ran all Multistate analyses with Reversible Jump (RJ) Markov chain Monte Carlo (MCMC)

methods. MCMC samples models in direct proportion to their fit to the data, generating a

posterior distribution of parameter estimates for each transition rate, and RJ sets some

parameters equal to zero or equal to one another, thereby reducing model complexity and over-

parametrization[98,99,101]. As a result, posterior distributions of parameter estimates may not be

normal; we thus summarised results by presenting mean and mode of the posterior distributions

of each parameter estimate, 95% higher posterior density, and percentage of models with

parameters estimated to be 0. We ran all MCMC chains for 320 million iterations in addition

to a burn-in of half a million iterations, sampling every 200000 iterations. All chains converged

and showed good mixing as indicated by their effective sample sizes of 2000 and visual

inspections of their traces in Tracer v1.6[103]. All analyses were run in triplicate and the three

independent chains converged on very similar solutions, leading to qualitatively similar results.

Here we present the results from the first chain. We ran RJ-MCMC Multistate analysis on

sexual system (Supplementary Table 4) as a binary state (gonochoristic or hermaphrodite) and

as a four-state categorical variable (gonochorism, protandry, protogyny, simultaneous
hermaphroditism). Bidirectional sequential hermaphrodites were excluded from the latter
analysis as the sample size of extant species was too low so that the chains failed to converge
and mix properly when sexual system was studied as a five states categorical variable. For the
analyses with two-character state, we graphed the evolutionary history of sexual systems on
the phylogeny using maximum likelihood (ML) in the R package *ape* v.5.3[104], which provided
a reasonably close approximation of the RJ-MCMC Multistate results (this was not the case
with the four-character state analysis).

We used phylogenetic generalized least square (PGLS) models[105-107] to test for the
association of each life history trait, entered as dependent variables, with sexual systems
entered as independent discrete variable with three possible states (gonochorism, protogyny,
protandry), as not enough data were available for simultaneous hermaphroditic species and
bidirectional sex-changers. PGLS models were run with the R package *caper*[108] in Maximum
Likelihood. The parameter $\lambda$ of PGLS models quantifies the strength of the phylogenetic signal
in the model residuals[105]. $\lambda$ ranges between zero (there is no phylogenetic structure in the data)
and one (the species share similarity in trait values directly proportional to their common
evolutionary time, under Brownian motion model of evolution[105,107]). Continuous variables
were $\log_{10}$-transformed to meet assumptions of normality.

**Acknowledgments**
This project was supported by Spanish Ministry of Science and Innovation Grant PID2019-
108888RB-I00 to F.P. and the Santander Universities Travel Award Grant 2016/2017 to C.B.
and S.P. that facilitated the collaboration among the three institutions involved in this study.
I.C. was supported by NERC (grant no. NE/K013777/1). We acknowledge the funding of the
Spanish government through the 'Severo Ochoa Centre of Excellence' accreditation
(CEX2019-000928-S). We acknowledge Professor Stefano Mariani for helpful discussion in
the first stages of the project and Professors Stephen Weeks and Manfred Schartl for valuable
comments on the manuscript.

**Author contributions**
F.P. conceived the study. S.P., C.B., I.C., and F.P. designed the study. S.P. collected data with
assistance from C.B. Data analyses were performed by S.P., C.B., and I.C. and S.P., C.B., I.C.,
and F.P. wrote the manuscript.

## Data availability

All data collected or analysed during this study are included in this published article and its supplementary information files.

## Competing interests

The authors declare no competing interests.

## References

1. Speijer, D., Lukeš, J. & Eliáš, M. Sex is a ubiquitous, ancient, and inherent attribute of eukaryotic life. *Proceedings of the National Academy of Sciences* **112,** 8827–8834 (2015).

2. Bachtrog, D. *et al.* Sex determination: why so many ways of doing it? *PLoS Biology* **12,** e1001899 (2014).

3. Ah-King, M. & Nylin, S. Sex in an evolutionary perspective: just another reaction norm. *Evolutionary Biology* **37,** 234–246 (2010).

4. Leonard, J. L. The evolution of sexual systems in animals. In Leonard, J.L. (ed.). *Transitions between sexual systems: understanding the mechanisms of, and pathways between, dioecy, hermaphroditism and other sexual systems*, 1–58 Springer (2019).

5. Weeks, S. C., Benvenuto, C. & Reed, S. K. When males and hermaphrodites coexist: a review of androdioecy in animals. *Integrative and Comparative Biology* **46,** 449–64 (2006).

6. Goldberg, E. E. *et al.* Macroevolutionary synthesis of flowering plant sexual systems. *Evolution* **71,** 898–912 (2017).

7. Waples, R. S., Mariani, S. & Benvenuto, C. Consequences of sex change for effective population size. *Proceedings of the Royal Society B: Biological Sciences* **285,** 20181702 (2018).

8. Benvenuto, C. & Weeks, S. C. Hermaphroditism and gonochorism. *The Natural History of the Crustacea: Reproductive Biology: Volume VI* 197–241 (2020).

9. Mariani, S., Sala-Bozano, M., Chopelet, J. & Benvenuto, C. Spatial and temporal patterns of size-at-sex-change in two exploited coastal fish. *Environmental Biology of Fishes* **96,** 535–541 (2013).

10. Käfer, J., Marais, G. A. & Pannell, J. R. On the rarity of dioecy in flowering plants. *Molecular Ecology* **26,** 1225–1241 (2017).

662    11.   Atz, J. Intersexuality in Fishes. In C.N. Amstrong and A.J. Marshall (eds).
663          *Intersexuality in vertebrates including man,* 145-232 Academic Press, London (1964).
664

665    12.   Jarne, P. & Auld, J. R. Animals mix it up too: the distribution of self-fertilization among
666          hermaphroditic animals. *Evolution* **60,** 1816–1824 (2006).
667

668    13.   Leonard, J. L. Williams' paradox and the role of phenotypic plasticity in sexual
669          systems. *Integrative and Comparative Biology* **53,** 671–688 (2013).
670

671    14.   Weeks, S. C. The role of androdioecy and gynodioecy in mediating evolutionary
672          transitions between dioecy and hermaphroditism in the animalia. *Evolution* **66,** 3670–
673          3686 (2012).
674

675    15.   Renner, S. S. The relative and absolute frequencies of angiosperm sexual systems:
676          dioecy, monoecy, gynodioecy, and an updated online database. *American Journal of*
677          *botany* **101,** 1588–1596 (2014).
678

679    16.   Bawa, K. S. Evolution of dioecy in flowering plants. *Annual review of ecology and*
680          *systematics* **11,** 15–39 (1980).
681

682    17.   Charlesworth, B. & Charlesworth, D. A model for the evolution of dioecy and
683          gynodioecy. *The American Naturalist* **112,** 975–997 (1978).
684

685    18.   Charlesworth, D. Androdioecy and the evolution of dioecy. *Biological Journal of the*
686          *Linnean Society* **22,** 333–348 (1984).
687

688    19.   Pannell, J. R. The evolution and maintenance of androdioecy. *Annual Review of*
689          *Ecology and Systematics* 397–425 (2002).
690

691    20.   Bull, J. & Charnov, E. On irreversible evolution. *Evolution* **39,** 1149–1155 (1985).
692

693    21.   Barrett, S. C. The evolution of plant reproductive systems: how often are transitions
694          irreversible? *Proceedings of the Royal Society B: Biological Sciences* **280,** 20130913
695          (2013).
696

697    22.   Oyarzún, P. A., Nuñez, J. J., Toro, J. E. & Gardner, J. P. Trioecy in the marine mussel
698          *Semimytilus algosus* (Mollusca, Bivalvia): stable sex ratios across 22 degrees of a
699          latitudinal gradient. *Frontiers in Marine Science* **7,** 348 (2020).
700

701    23.   Dani, K. & Kodandaramaiah, U. Plant and animal reproductive strategies: lessons from
702          offspring size and number tradeoffs. *Frontiers in Ecology and Evolution* **5,** 38 (2017).
703

704    24.   Avise, J. & Mank, J. Evolutionary perspectives on hermaphroditism in fishes. *Sexual*
705          *Development* **3,** 152–163 (2009).
706

707    25.   Dornburg, A. & Near, T. J. The Emerging phylogenetic perspective on the evolution of
708          Actinopterygian fishes. *Annual Review of Ecology, Evolution, and Systematics* **52, 427-**
709          **452** (2021).
710

26.　Costa, W. J., Lima, S. M. & Bartolette, R. Androdioecy in *Kryptolebias* killifish and the
evolution of self-fertilizing hermaphroditism. *Biological Journal of the Linnean Society*
**99,** 344–349 (2010).

27.　Costa, W. Colouration, taxonomy and geographical distribution of mangrove killifishes,
the *Kryptolebias marmoratus* species group, in southern Atlantic coastal plains of Brazil
(Cyprinodontiformes: Rivulidae). *Ichthyological Exploration of Freshwaters* **27,** 183–
192 (2016).

28.　Powell, M. L., Kavanaugh, S. I. & Sower, S. A. Seasonal concentrations of reproductive
steroids in the gonads of the Atlantic hagfish, *Myxine glutinosa*. *Journal of*
*Experimental Zoology Part A Comparative Experimental Biology* **301,** 352–60 (2004).

29.　Pennell, M. W., Mank, J. E. & Peichel, C. L. Transitions in sex determination and sex
chromosomes across vertebrate species. *Molecular Ecology* **27,** 3950–3963 (2018).

30.　Ghiselin, M. T. The evolution of hermaphroditism among animals. *Quarterly Review of*
*Biology* **44,** 189–208 (1969).

31.　Eppley, S. M. & Jesson, L. K. Moving to mate: the evolution of separate and combined
sexes in multicellular organisms. *Journal of Evolutionary Biology* **21,** 727–36 (2008).

32.　Warner, R. R. The adaptive significance of sequential hermaphroditism in animals.
*American Naturalist* **109,** 61–82 (1975).

33.　Warner, R. R., Robertson, D. R. & Leigh, E. G. Sex change and sexual selection.
*Science* **190,** 633–638 (1975).

34.　Charnov, E. L. *The Theory of Sex Allocation*. Princeton University Press, USA (1982).

35.　Policansky, D. Sex change in plants and animals. *Annual Review of Ecology and*
*Systematics* **13,** 471–495 (1982).

36.　Benvenuto, C., Coscia, I., Chopelet, J., Sala-Bozano, M. & Mariani, S. Ecological and
evolutionary consequences of alternative sex-change pathways in fish. *Scientific*
*Reports* **7,** 9084 (2017).

37.　Charnov, E. L. Natural selection and sex change in pandalid shrimp: test of a life-
history theory. *The American Naturalist* **113,** 715–734 (1979).

38.　Broquet, T. *et al.* The size advantage model of sex allocation in the protandrous sex-
changer *Crepidula fornicata*: role of the mating system, sperm storage, and male
mobility. *The American Naturalist* **186,** 404–420 (2015).

39.　Erisman, B. E., Craig, M. T. & Hastings, P. A. A phylogenetic test of the size-advantage
model: evolutionary changes in mating behavior influence the loss of sex change in a
fish lineage. *The American Naturalist* **174,** E83–E99 (2009).
758

759    40.  Buxton, C. D. & Garratt, P. A. Alternative reproductive styles in seabreams (Pisces:
760          Sparidae). *Environmental Biology of Fishes* **28,** 113–124 (1990).
761

762    41.  Shapiro, D. Y. Social behavior, group structure, and the control of sex reversal in
763          hermaphroditic fish. *Advances in the Study of Behavior* **10,** 43–102 (1979).
764

765    42.  Stearns, S. C. Life history evolution: successes, limitations, and prospects.
766          *Naturwissenschaften* **87,** 476–486 (2000).
767

768    43.  Waples, R. S., Luikart, G., Faulkner, J. R. & Tallmon, D. A. Simple life-history traits
769          explain key effective population size ratios across diverse taxa. *Proceedings of the
770          Royal Society of London B: Biological Sciences* **280,** 20131339 (2013).
771

772    44.  Martinez, A. S., Willoughby, J. R. & Christie, M. R. Genetic diversity in fishes is
773          influenced by habitat type and life-history variation. *Ecology and Evolution* **8,** 12022–
774          12031 (2018).
775

776    45.  Harvey, P. H. & Pagel, M. D. *The comparative method in evolutionary biology*. Oxford
777          University Press, USA (1991).
778

779    46.  Barneche, D. R., Robertson, D. R., White, C. R. & Marshall, D. J. Fish reproductive-
780          energy output increases disproportionately with body size. *Science* **360,** 642–645
781          (2018).
782

783    47.  Brandl, S. J. & Bellwood, D. R. Pair-formation in coral reef fishes: an ecological
784          perspective. *Oceanography and Marine Biology: An Annual Review* **52,** 1–80 (2014).
785

786    48.  Fitzpatrick, J. L. Sperm competition and fertilization mode in fishes. *Philosophical
787          Transactions of the Royal Society B: Biological Sciences* **375,** 20200074 (2020).
788

789    49.  Parker, G. A. Conceptual developments in sperm competition: a very brief synopsis.
790          *Philosophical Transactions of the Royal Society B: Biological Sciences* **375,** 20200061
791          (2020).
792

793    50.  Warner, R. R. Sex change in fishes: hypotheses, evidence, and objections.
794          *Environmental Biology of Fishes* **22,** 81–90 (1988).
795

796    51.  Molloy, P. P., Goodwin, N. B., Côté, I. M., Reynolds, J. D. & Gage, M. J. Sperm
797          competition and sex change: a comparative analysis across fishes. *Evolution* **61,** 640–
798          652 (2007).
799

800    52.  Erisman, B. E., Petersen, C. W., Hastings, P. A. & Warner, R. R. Phylogenetic
801          perspectives on the evolution of functional hermaphroditism in teleost fishes.
802          *Integrative and Comparative Biology* **53,** 736–754 (2013).
803

804    53.  Sadovy, Y., Colin, P. & Domeier, M. Aggregation and spawning in the tiger grouper,
805          *Mycteroperca tigris* (Pisces: Serranidae). *Copeia* **1994,** 511–516 (1994).
806

807    54.  Muñoz, R. C. & Warner, R. R. A new version of the size-advantage hypothesis for sex
808          change: incorporating sperm competition and size-fecundity skew. *The American

809         *Naturalist* **161,** 749–761 (2003).
810

811  55.  Horne, C. R., Hirst, A. G. & Atkinson, D. Selection for increased male size predicts
812         variation in sexual size dimorphism among fish species. *Proceedings of the Royal*
813         *Society B: Biological Sciences* **287,** 20192640 (2020).
814

815  56.  Parker, G. The evolution of expenditure on testes. *Journal of Zoology* **298,** 3–19 (2016).
816

817  57.  Stockley, P., Gage, M., Parker, G. & Møller, A. Sperm competition in fishes: the
818         evolution of testis size and ejaculate characteristics. *The American Naturalist* **149,** 933–
819         954 (1997).
820

821  58.  Pla, S., Benvenuto, C., Capellini, I. & Piferrer, F. A phylogenetic comparative analysis
822         on the evolution of sequential hermaphroditism in seabreams (Teleostei: Sparidae).
823         *Scientific Reports* **10,** 3606 (2020).
824

825  59.  Vrijenhoek, R. C. Unisexual fish: model systems for studying ecology and evolution.
826         *Annual Review of Ecology and Systematics* **25,** 71–96 (1994).
827

828  60.  Sadovy de Mitcheson, Y. & Liu, M. Functional hermaphroditism in teleosts. *Fish and*
829         *Fisheries* **9,** 1–43 (2008).
830

831  61.  Rabosky, D. L. *et al.* An inverse latitudinal gradient in speciation rate for marine fishes.
832         *Nature* **559,** 392 (2018).
833

834  62.  Froese, R., Pauly, D. & Editors. FishBase. World Wide Web electronic publication.
835         www.fishbase.org (2018).
836

837  63.  Moore, W. S. Evolutionary ecology of unisexual fishes. In *Evolutionary genetics of*
838         *fishes* 329–398 Springer (1984).
839

840  64.  Schiettekatte, N., Brandl, S. & Casey, J. Fishualize: Color palettes based on fish species.
841         *R package v0.2.2* (2021).
842

843  65.  Collin, R. & Miglietta, M. P. Reversing opinions on Dollo's Law. *Trends in Ecology &*
844         *Evolution* **23,** 602–609 (2008).
845

846  66.  Domes, K., Norton, R. A., Maraun, M. & Scheu, S. Re-evolution of sexuality breaks
847         Dollo's law. *Proceedings of the National Academy of Sciences* **104,** 7139–7144 (2007).
848

849  67.  Dollo, L. Les lois de l'évolution. *Bulletin de la Société belge de géologie, de*
850         *paléontologie et d'hydrologie* **7,** 164–166 (1893).
851

852  68.  King, B. & Lee, M. S. Ancestral state reconstruction, rate heterogeneity, and the
853         evolution of reptile viviparity. *Systematic Biology* **64,** 532–544 (2015).
854

855  69.  Uller, T. & Helanterä, H. From the origin of sex-determining factors to the evolution of
856         sex-determining systems. *The Quarterly Review of Biology* **86,** 163–180 (2011).
857

70. Devlin, R. H. & Nagahama, Y. Sex determination and sex differentiation in fish: an
overview of genetic, physiological, and environmental influences. *Aquaculture* **208,**
191–364 (2002).

71. Volff, J.-N., Nanda, I., Schmid, M. & Schartl, M. Governing sex determination in fish:
regulatory putsches and ephemeral dictators. *Sexual Development* **1,** 85–99 (2007).

72. Nagahama, Y., Chakraborty, T., Paul-Prasanth, B., Ohta, K. & Nakamura, M. Sex
determination, gonadal sex differentiation and plasticity in vertebrate species.
*Physiological Reviews* **101,** 1237-1308 (2020).

73. Penman, D. J. & Piferrer, F. Fish gonadogenesis. Part I: genetic and environmental
mechanisms of sex determination. *Reviews in Fisheries Science* **16(S1),** 16–34 (2008).

74. Mank, J. E., Promislow, D. E. L. & Avise, J. C. Evolution of alternative sex-
determining mechanisms in teleost fishes. *Biological Journal of the Linnean Society* **87,**
83–93 (2006).

75. Galetti, P. M., Aguilar, C. T. & Molina, W. F. An overview of marine fish cytogenetics.
*Hydrobiologia* **420,** 55–62 (2000).

76. Yoshida, K. *et al.* Sex chromosome turnover contributes to genomic divergence
between incipient stickleback species. *PLoS Genetics* **10,** e1004223 (2014).

77. Ross, J. A., Urton, J. R., Boland, J., Shapiro, M. D. & Peichel, C. L. Turnover of sex
chromosomes in the stickleback fishes (Gasterosteidae). *PLoS Genetics* **5,** e1000391
(2009).

78. Vicoso, B. Molecular and evolutionary dynamics of animal sex-chromosome turnover.
*Nature Ecology & Evolution* 1–10 (2019).

79. Gamble, T. *et al.* Restriction site-associated DNA sequencing (RAD-seq) reveals an
extraordinary number of transitions among gecko sex-determining systems. *Molecular*
*Biology and Evolution* **32,** 1296–1309 (2015).

80. Pokorná, M. & Kratochvíl, L. Phylogeny of sex-determining mechanisms in squamate
reptiles: are sex chromosomes an evolutionary trap? *Zoological Journal of the Linnean*
*Society* **156,** 168–183 (2009).

81. Furman, B. L. *et al.* Sex chromosome evolution: So many exceptions to the rules.
*Genome Biology and Evolution* **12,** 750-763 (2020).

82. Carvalho, N. D. M. *et al.* Cytogenetics of Synbranchiformes: a comparative analysis of
two *Synbranchus* Bloch, 1795 species from the Amazon. *Genetica* **140,** 149–58 (2012).

83. Piferrer, F. Epigenetic mechanisms in sex determination and in the evolutionary
transitions between sexual systems. *Philosophical Transactions of the Royal Society B:*
*Biological sciences* **376,** 20200110 (2021).

84. Grant, S. *et al.* Genetics of sex determination in flowering plants. *Developmental*
*Genetics* **15,** 214–230 (1994).
909

910 85. Harrington Jr, R. W. How ecological and genetic factors interact to determine when
911    self-fertilizing hermaphrodites of *Rivulus marmoratus* change into functional secondary
912    males, with a reappraisal of the modes of intersexuality among fishes. *Copeia* 389–432
913    (1971).
914

915 86. Adolfi, M. C., Nakajima, R. T., Nóbrega, R. H. & Schartl, M. Intersex,
916    Hermaphroditism, and gonadal plasticity in vertebrates: Evolution of the Müllerian duct
917    and Amh/Amhr2 signalling. *Annual Review of Animal Biosciences* (2018).
918

919 87. Adkins-Regan, E. Early organizational effects of hormones: an evolutionary
920    perspective. In Adler, N.T. (ed.) *Neuroendocrinology of reproduction: physiology and*
921    *behavior* 159–228 Springer USA (1981).
922

923 88. Navara, K. J. The truth about Nemo's dad: sex-changing behaviors in fishes. In
924    *Choosing Sexes* 183–212 Springer, Cham (2018).
925

926 89. Orban, L., Sreenivasan, R. & Olsson, P.E. Long and winding roads: testis differentiation
927    in zebrafish. *Molecular and Cellular Endocrinology* **312,** 35–41 (2009).
928

929 90. Zohar, Y., Abraham, M. & Gordin, H. The gonadal cycle of the captivity-reared
930    hermaphroditic teleost *Sparus aurata* (L.) during the first two years of life. *Annales de*
931    *Biologie Animale Biochimie Biophysique* **18,** 877–882 (1978).
932

933 91. Chang, C.-F. & Yueh, W.-S. Annual cycle of gonadal histology and steroid profiles in
934    the juvenile males and adult females of the protandrous black porgy, *Acanthopagrus*
935    *schlegelii*. *Aquaculture* **91,** 179–196 (1990).
936

937 92. Miura, S., Nakamura, S., Kobayashi, Y., Piferrer, F. & Nakamura, M. Differentiation of
938    ambisexual gonads and immunohistochemical localization of P450 cholesterol side-
939    chain cleavage enzyme during gonadal sex differentiation in the protandrous
940    anemonefish, *Amphiprion clarkii*. *Comparative Biochemistry and Physiology Part B:*
941    *Biochemistry and Molecular Biology* **149,** 29–37 (2008).
942

943 93. Yamaguchi, S. & Iwasa, Y. Advantage for the sex changer who retains the gonad of the
944    nonfunctional sex. *Behavioral Ecology and Sociobiology* **71,** 39 (2017).
945

946 94. Munday, P. L., Kuwamura, T. & Kroon, F. J. Bi-directional sex change in marine fishes.
947    In: Cole, K.S. (ed.) *Reproduction and sexuality in marine fishes: Patterns and*
948    *processes.* University of California Press, Berkeley, USA 241–271 (2010).
949

950 95. Uller, T., Feiner, N., Radersma, R., Jackson, I. S. & Rago, A. Developmental plasticity
951    and evolutionary explanations. *Evolution & Development* **22,** 47–55 (2020).
952

953 96. Pla, S., Maynou, F. & Piferrer, F. Hermaphroditism in fish: incidence, distribution and
954    associations with abiotic environmental factors. *Reviews in Fish Biology and Fisheries*
955    **31,** 935–955 (2021).
956

957  97. Boettiger, C., Lang, D. T. & Wainwright, P. C. *rfishbase*: exploring, manipulating and
958      visualizing FishBase data from R. *Journal of Fish Biology* **81,** 2030–9 (2012).
959

960  98. Pagel, M., Meade, A. & Barker, D. Bayesian estimation of ancestral character states on
961      phylogenies. *Systematic Biology* **53,** 673–684 (2004).
962

963  99. Pagel, M. & Meade, A. Bayesian analysis of correlated evolution of discrete characters
964      by reversible-jump Markov chain Monte Carlo. *The American Naturalist* **167,** 808–25
965      (2006).
966

967  100. Pagel, M. Detecting correlated evolution on phylogenies: a general method for the
968      comparative analysis of discrete. *Proceedings of the Royal Society B: Biological
969      Sciences* **255,** 37-45 (1994).
970

971  101. Currie, T. E. & Meade, A. in *Modern phylogenetic comparative methods and their
972      application in evolutionary biology* 263–286 Springer (2014).
973

974  102. Furness, A. I. & Capellini, I. The evolution of parental care diversity in amphibians.
975      *Nature Communications* **10,** 1–12 (2019).
976

977  103. Rambaut, A., Drummond, A. J., Xie, D., Baele, G. & Suchard, M. A. Posterior
978      summarization in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology* **67,** 901
979      (2018).
980

981  104. Paradis, E. & Schliep, K. ape 5.0: an environment for modern phylogenetics and
982      evolutionary analyses in R. *Bioinformatics* **35,** 526–528 (2019).
983

984  105. Freckleton, R. P., Harvey, P. H. & Pagel, M. Phylogenetic analysis and comparative
985      data: a test and review of evidence. *The American Naturalist* **160,** 712–726 (2002).
986

987  106. Pagel, M. Inferring evolutionary processes from phylogenies. *Zoologica Scripta* **26,**
988      331–348 (1997).
989

990  107. Pagel, M. Inferring the historical patterns of biological evolution. *Nature* **401,** 877–84
991      (1999).
992

993  108. Orme, D. The *caper* package: Comparative analysis of phylogenetics and evolution in R
994      (2013). *https://cran. r-project. org/web/packages/caper/vignettes/caper. pdf* (2018).
995

# Switches, stability and reversals:
# the evolutionary history of sexual systems in fish

Susanna Pla[1#], Chiara Benvenuto[2#], Isabella Capellini[3*] & Francesc Piferrer[1*]

[1]Institut de Ciències del Mar, Spanish National Research Council (CSIC), Barcelona, Spain

[2]School of Science, Engineering and Environment, University of Salford, Salford, UK

[3]School of Biological Sciences, Queen's University Belfast, UK

#Joint first authors

*Correspondence: Dr. Francesc Piferrer, Institut de Ciències del Mar, Spanish National Research Council (CSIC), Passeig Marítim, 37-49, 08003 Barcelona. Tel. +34-932 309 567; E-mail: piferrer@icm.csic.es

*Correspondence: Dr. Isabella Capellini, School of Biological Sciences, Queen's University Belfast, 19 Chlorine Gardens, Belfast, BT9 5DL, Northern Ireland, United Kingdom; E-mail: I.Capellini@qub.ac.uk

# Supplementary information

- Supplementary tables (pp 2–5)
- Supplementary methods (pp 6-7)
- Supplementary references (pp 8–29)
- Supplementary data are attached as an Excel file

# Supplementary tables

**Supplementary Table 1.** Results of the RJ-MCMC Multistate analysis in *BayesTraits* of sexual systems as a binary trait: gonochorism (G) or hermaphroditism (H). For each posterior distribution, we report the effective sample size (ESS), the mean and 95% high posterior density (95-HPD) intervals, the mode, and the percentage of models in which the parameter is estimated as zero. Analysis based on 4614 extant teleost species (G: n = 4320; H: n = 294)

| Transition rates | ESS | Mean | 95-HPD | Mode | % Zero |
|---|---|---|---|---|---|
| G → H | 1600 | 0.034 | 0.021 – 0.047 | 0.032 | **0** |
| H → G | 1600 | 0.826 | 0.608 – 1.038 | 0.823 | **0** |
| **Root probabilities** | | | | | |
| G | 1600 | 66.0 | 59.1 – 74.1 | 65.1 | **0** |
| H | 1600 | 34.0 | 25.9 – 40.9 | 34.9 | **0** |

**Supplementary Table 2.** Results of phylogenetic generalized least square (PGLS) model of longevity (year; $\log_{10}$ transformed), controlled for allometry (maximum length, in cm; log10 transformed; age at first maturity (year; $\log_{10}$ transformed) per each sex (♂: male; ♀: female), controlling for allometry (length at maturity in cm; $\log_{10}$ transformed); and male gonadosomatic index (GSI; log10 transformed), controlling for allometry (male length at maturity in cm; $\log_{10}$ transformed) across sexual systems: gonochorism (G); protogyny (PG); protandry (PA). For each independent variable we report the parameter estimate (Beta), t-statistics (T), P-value (P; two-sided test), and the model statistics including the degrees of freedom (df), the maximum likelihood estimation of the phylogenetic signal ($\lambda$) and $R^2$. Significant differences are indicated in bold. See Supplementary Table 3 for sexual system and sex-specific data

| | **Variable** | **Beta** | **T** | **P** | **Df** | **Model statistics** | |
| **Dependent** | **Independent** | | | | | $\lambda$ | $R^2$ |
|---|---|---|---|---|---|---|---|
| Longevity | Max length | 0.680 | 19.968 | **<0.0001** | 3; 640 | 0.864 | 0.388 |
| | Sexual system - PA[1] | -0.043 | -0.417 | 0.677 | | | |
| | Sexual system - PG[1] | 0.129 | 2.193 | **0.029** | | | |
| | Sexual system - PG[2] | 0.172 | 1.538 | 0.124 | | | |
| Age at maturity ♂ | Length at maturity ♂ | 0.602 | 8.360 | **<0.001** | 3; 149 | 0.785 | 0.325 |
| | Sexual system - PA[1] | -0.090 | -0.956 | 0.340 | | | |
| | Sexual system - PG[1] | 0.068 | 0.774 | 0.440 | | | |
| | Sexual system - PG[2] | 0.159 | 1.434 | 0.154 | | | |
| Age at maturity ♀ | Length at maturity ♀ | 0.536 | 8.062 | **<0.001** | 3; 166 | 0.829 | 0.282 |
| | Sexual system - PA[1] | 0.016 | 0.117 | 0.907 | | | |
| | Sexual system - PG[1] | -0.014 | -0.167 | 0.867 | | | |
| | Sexual system - PG[2] | -0.029 | -0.217 | 0.828 | | | |
| GSI ♂ | Length at maturity ♂ | 0.041 | 0.236 | 0.814 | 3; 51 | 0.000 | 0.376 |
| | Sexual system - PA[1] | 0.164 | 1.197 | 0.237 | | | |
| | Sexual system - PG[1] | -0.513 | -4.631 | **<0.001** | | | |
| | Sexual system - PG[2] | -0.678 | -4.582 | **<0.001** | | | |

**Supplementary Table 3.** Sample sizes for longevity (year), maximum length (cm), age at maturity (years), length at maturity (cm) and gonadosomatic index (GSI) for each sexual system with sex-specific male (♂) and female (♀) data, when available. In italics data not used in the analyses. G = gonochorism; PG = protogyny; PA = protandry; SH = simultaneous hermaphroditism; BD = bidirectional hermaphroditism

| Life-history traits | Sex | G | PG | PA | Total for analyses | *SH* | *BD* |
|---|---|---|---|---|---|---|---|
| **Longevity** | | 758 | 69 | 17 | 844 | *7* | *3* |
| **Maximum length** | | 2612 | 167 | 20 | 2799 | *28* | *11* |
| **Age at maturity** | ♂ | 259 | 15 | 9 | 283 | *1* | - |
| | ♀ | 282 | 30 | 5 | 317 | *2* | - |
| **Length at maturity** | ♂ | 305 | 42 | 15 | 362 | *9* | - |
| | ♀ | 297 | 36 | 10 | 343 | *2* | - |
| **GSI** | ♂ | 44 | 38 | 15 | 97 | *3* | - |

**Supplementary Table 4.** Number of species used to study the evolutionary transitions among different sexual systems. Note that androdioecious species are not included in the analyses

| Sexual system | 2 state [G, H] | 4 state [G, PG, PA, SH] |
|---|---|---|
| Gonochorism [G] | 4320 | 4320 |
| Hermaphroditism [H] | 294* | |
| Protogyny [PG] | | 196 |
| Protandry [PA] | | 36 |
| Simultaneous hermaphroditism [SH] | | 46 |
| Total | 4614 | |

*Includes 16 species of bidirectional sequential hermaphrodites, which were not included in other analyses due to their small sample size when added as a separate category.

# Supplementary methods

**Data collection and verification**

Information on the sexual system and life history traits was previously collected from FishBase (www.fishbase.org) for a total of 10914 actinopterygian species, of which 10875 were teleosts. Of these, we only retained the 4740 teleost species that were also present in the phylogenetic tree of Rabosky et al. (2018), of which 4320 were gonochoristic and 420 had different forms of hermaphroditism (Supplementary Figure 1). Of the latter, we retained only the species in which functional hermaphroditism has been reported in the primary literature (compiled in Pla et al., 2021) plus some additional species added also from the primary literature (all used references are provided). Regarding gonochoristic species, we only considered those that were regarded as such in FishBase, unless recent literature states otherwise. Importantly, species for which there is contrasting information in the literature were discarded and not used for this study.

**Supplementary Figure 1.** Diagram of the data acquisition and curation followed in this study.

We also extracted data from primary literature, FishBase (www.fishbase.org; CD-ROM version), and rfishbase (https://www.rdocumentation.org/packages/rfishbase/versions/3.0.4; Boettiger et al., 2012), on the following life-history traits:

1) Longevity (in years). When more than one value was present for a given species, we used the maximum value reported in the wild.

2) Maximum length, specifically total length (TL; in cm).

3) Age (in years) and length (in cm) at first maturity for each sex.

4) Male gonadosomatic index (GSI). The maximum value (expected to coincide with the peak of the reproductive season) was used.

We controlled for allometry as follows: longevity was controlled for maximum length (available for both sexes combined); age at maturity was controlled for length at maturity (by sex). GSI was controlled for male length at maturity (male-specific): in this case we could not use maximum length, not sex-specific, which would give an incorrect length of males in protandric species, where the larger individuals are females.

**References**

Boettiger, C., Lang, D.T. & Wainwright, D.C. (2012) rfishbase: exploring, manipulating and visualizing FishBase data from R. *Journal of Fish Biology* 81, 2030–2039.

Pla, S., Maynou, F. & Piferrer, F. (2021) Hermaphroditism in fish: incidence, distribution and associations with abiotic environmental factors. *Reviews in Fish Biology and Fisheries* 31, 935–955.

Rabosky, D. L. et al. (2018) An inverse latitudinal gradient in speciation rate for marine fishes. *Nature* 559, 392-395.

# Supplementary data

Full dataset attached as an Excel file:
- Sheet 1. Sexual system and life history traits for a total of 4614 teleosts.
- Sheet 2. List Sexual system of the 294 hermaphroditic species used in this study and the supporting primary literature (listed below).

## References to Supplementary Data

Abd-el-Aziz, S.H. & Ramadan, A.A. (1990) Sexuality and hermaphroditism in fishes. I. Synchronous functional hermaphroditism in the serranid fish *Serranus scriba* L. *Folia Morphologica, 38*(1), 86-100.

Abou-Seedo, F., Wright, J.M. & Clayton, D. (1990) Aspects of the biology of *Diplodus sargus kotschyi* (Sparidae) from Kuwait Bay. *Cybium, 14*(3), 217-223.

Abou-Seedo, F.S., Dadzie, S. & Al-Kanaan, K.A. (2003) Sexuality, sex change and maturation patterns in the yellowfin seabream, *Acanthopagrus latus* (Teleostei: Sparidae) (Houttuyn, 1782). *Journal of Applied Ichthyology, 19*(2), 65-73. http://doi.org/ 10.1046/j.1439-0426.2003.00355.x

Abu-Hakima, R. (1984) Some aspects of the reproductive biology of Acanthopagrus spp. (Family: Sparidae). *Journal of Fish Biology, 25*(5), 515-526. http://doi.org/10.1111/j.1095-8649.1984.tb04898.x

Adams, S. (2003) Morphological ontogeny of the gonad of three plectropomid species through sex differentiation and transition. *Journal of Fish Biology, 63*(1), 22-36. http://doi.org/**1**0.1046/j.1095-8649.2003.00098.x

Adreani, M.S. & Allen, L.G. (2008) Mating system and reproductive biology of a temperate wrasse, *Halichoeres semicinctus*. *Copeia, 2008*(2), 467-475. http://doi.org/10.1643/CP-06-265

Adreani, M.S., Erisman, B.E. & Warner, R.R. (2004) Courtship and spawning behavior in the California sheephead, *Semicossyphus pulcher* (Pisces: Labridae). *Environmental Biology of Fishes, 71*(1), 13-19. http://doi.org/10.1023/B:EBFI.0000043177.98895.f7

Aldenhoven, J.M. (1984) S*ocial organization and sex change in an angelfish, Centropyge bicolor on the Great Barrier Reef.* Unpublished PhD dissertation, Macquarie University, North Ryde, Australia

Alekseev, F.E. (1982) Hermaphroditism in sparid fishes (Perciformes, Sparidae). I. Protogyny in porgies, *Pagrus, P. orphus, P. ehrenbergi* and *P. auriga*, from West Africa. *Journal of Ichthyology, 22*, 85-94

Alekseev, F.E. (1983) Hermaphroditism in porgies (Perciformes, Sparidae). II. Sexual structure of the populations, mechanism of its formation and evolution in scups, *Pagrus, P. orphus, P. ehrenbergi* and *P. auriga. Journal of Ichthyology, 23*, 61-67

Anderson, W.D. & Baldwin, C.C. (2000) A new species of Anthias (Teleostei: Serranidae: Anthiinae) from the Galápagos Islands, with keys to Anthias and eastern Pacific Anthiinae. *Proceedings of the Biological Society of Washington, 113*(2), 369-385.

Adreani, M.S. & Allen, L.G. (2008) Mating system and reproductive biology of a temperate wrasse, *Halichoeres semicinctus*. *Copeia, 2008*(2), 467-475. http://doi.org/10.1643/CP-06-265

Aoyama, T. (1955) On the hermaphroditism in the yellow sea bream, *Taius tumifrons*. *Japanese Journal of Ichthyology, 4*(4-6), 119-129. http://doi.org/10.11369/jji1950.4.119

Aronov, A. & Goren, M. (2008) Ecology of the mottled grouper (*Mycteroperca rubra*) in the eastern Mediterranean. *Electronic Journal of Ichthyology, 2*, 43-55

Asoh, K. & Yoshikawa, T. (2003) Gonadal development and an indication of functional protogyny in the Indian damselfish (*Dascyllus carneus*). *Journal of Zoology, 260*(1), 23-39. http://doi.org/10.1017/S0952836903003418

Asoh, K. (2004) Gonadal development in the coral reef damselfish *Dascyllus flavicaudus* from Moorea, French Polynesia. *Marine Biology, 146*(1), 167-179. http://doi.org/10.1007/s00227-004-1414-7

Asoh K (2005a) Frequency of functional sex change in two populations of *Dascyllus melanurus* conforms to a prediction from sex allocation theory. *Copeia* 2005:732-744.

Asoh, K. (2005b) Gonadal development and diandric protogyny in two populations of *Dascyllus reticulatus* from Madang, Papua New Guinea. *Journal of Fish Biology, 66*(4), 1127-1148. http://doi.org/10.1111/j.0022-1112.2005.00669.x

Bacci, G. & Razzauti, A. (1958) Protogynous hermaphroditism in *Coris julis* L. *Nature, 181*(4606), 432-433.

Bañón, R., Villegas-Ríos, D., Rodriguez Tamargo, P., Alonso-Fernández, A., Barros-García, D. and De Carlos, A. (2018) First record of *Epinephelus costae* (Actinopterygii: Perciformes: Epinephelidae) from Galician waters (north-western Spain): Exploring the northward range expansion. *Acta Ichthyologica et Piscatoria* , *48*(4), 399-402 http://doi.org/10.3750/AIEP/02490

Barlow, G.W. (1975) On the sociobiology of some hermaphroditic serranid fishes, the hamlets, in Puerto Rico. *Marine Biology, 33*(4), 295-300. http://doi.org/10.1007/BF00390567

Bean, K., Mapstone, B.D., Davies, C.R., Murchie, C.D. & Williams, A.J. (2003) Gonad development and evidence of protogyny in the red-throat emperor on the Great Barrier Reef. *Journal of Fish Biology, 62*(2), 299-310. http://doi.org/10.1046/j.1095-8649.2003.00021.x

Bentivegna, F. & Rasotto, M.B. (1987) Protogynous hermaphroditism in *Xyrichthys novacula* (L. 1758). *Cybium, 11*(1), 75-78.

Bertrand, J. (1986) Données concernant la reproduction de *Lethrinus mahsena* (Forsskal 1775) sur les bancs de Saya de Malha (Océan Indien)." *Cybium (Paris)* p. 15-29.

Besseau, L. & Bruslé-Sicard, S. (1995) Plasticity of gonad development in hermaphroditic sparids: ovotestis ontogeny in a protandric species, *Lithognathus mormyrus*. *Environmental Biology of Fishes, 43*(3), 255-267. http://doi.org/10.1007/BF00005857

Bhandari, R.K., Alam, M.A.l., Higa, M., Soyano, K. & Nakamura, M. (2005) Evidence that estrogen regulates the sex change of honeycomb grouper (*Epinephelus merra*), a protogynous hermaphrodite fish. *Journal of Experimental Zoology Part A: Comparative Experimental Biology, 303*(6), 497-503. http://doi.org/10.1002/jez.a.178

Bhandari, R.K., Komuro, H., Nakamura, S., Higa, M. & Nakamura, M. (2003) Gonadal restructuring and correlative steroid hormone profiles during natural sex change in protogynous honeycomb grouper (*Epinephelus merra*). *Zoological science, 20*(11), 1399-1405. http://doi.org/10.2108/zsj.20.1399

Bortone, S.A. (1971) Studies on the biology of the sand perch, *Diplectrum formosum* (Perciformes: Serranidae). *Florida Department of Natural Resources, Marine Research Laboratory Series 65*, 1-27.

Bortone, S.A. (1977a) Gonad morphology of the hermaphroditic fish *Diplectrum pacificum* (Serranidae). *Copeia 1977*(3), 448-453. http://doi.org/10.2307/1443262

Bortone, S.A. (1977b) Observations on the life history of *Diplectrum pacificum* and *D. macropoma* (Pisces, Serranidae) from the Bay of Panama. *Pacific Science, 31*(1), 49-60.

Bortone, S.A. (1977c) Revision of the sea basses ofthe genus Diplectrum (Pisces: Serranidae). *U.S. Department of Commerce, NOAA Technical Reports, 404*, 1-49.

Brickle, P., Laptikhovsky, V., & Arkhipkin, A. (2005). Reproductive strategy of a primitive temperate notothenioid *Eleginops maclovinus*. *Journal of Fish*

*Biology, 66*(4), 1044-1059.

Bruce, R.W. (1980) Protogynous hermaphroditism in two marine angelfishes. *Copeia, 1980*(2), 353-355. http://doi.org/10.2307/1444017

Brulé, T., Colás-Marrufo, T.E., Tuz-Sulub, A. & Déniel, C. (2000) Evidence for protogynous hermaphroditism in the serranid fish *Epinephelus drummondhayi* (Perciformes: Serranidae) from the Campeche Bank in the southern Gulf of Mexico. *Bulletin of Marine Science, 66*(2), 513-521.

Brulé, T., Déniel, C., Colás-Marrufo, T. & Sánchez-Crespo, M. (1999) Red grouper reproduction in the southern Gulf of Mexico. *Transactions of the American Fisheries Society, 128*(3), 385-402.

Bruslé, J. & Bruslé, S. (1975) Ovarian and testicular intersexuality in two protogynous mediterranean groupers, *Epinephelus aeneus* and *Epinephelus guaza*. In: R. Reinboth (Ed.), *Intersexuality in the Animal Kingdom* (pp. 222-227). Springer, Berlin, Heidelberg. http://doi.org/10.1007/978-3-642-66069-6_21

Bruslé, S. (1983) Contribution to the sexuality of a Hermaphroditic teleost, *Serranus hepatus* L. *Journal of Fish Biology, 22*(3), 283-292. http://doi.org/10.1111/j.1095-8649.1983.tb04752.x

Bruslé, S. (1987) Sex-inversion of the hermaphroditic, protogynous teleost *Coris julis* L. (Labridae). *Journal of Fish Biology, 30*(5), 605-616. http://doi.org/10.1111/j.1095-8649.1987.tb05788.x

Bullock, L.H. & Murphy, M.D. (1994) Aspects of the life history of the yellowmouth grouper, *Mycteroperca interstitialis*, in the eastern Gulf of Mexico. *Bulletin of Marine Science, 55*(1), 30-45.

Bullock, L.H. & Smith, G.B. (1991) Seabasses (Pisces: Serranidae). *Memoirs of the Hourglass Cruises, 8*, 1-243.

Burgos, J.M., Sedberry, G.R., Wyanski, D.M. & Harris, P.J. (2007) Life history of red grouper (*Epinephelus morio*) off the coasts of North Carolina and South Carolina. *Bulletin of Marine Science, 80*(1), 45-65.

Burnett-Herkes, J.N. (1975) *Contributions to the biology of the red hind, Epinephelus guttatus, a commercially important serranid fish from the tropical western Atlantic.* PhD dissertation, University of Miami., Coral Gables, 154 pp.

Burton, M. L., Potts, J. C., Poholek, A. J., Ostrowski, A., & Page, J. (2019). Age, growth, natural mortality, and reproductive seasonality of knobbed porgy from southeastern United States waters. *Marine and Coastal Fisheries*, *11*(2), 231-245.

Bustos, R., Luque-Escalona, A. & Pajuelo, J.G. (2010) Reproductive biology of the island grouper (*Mycteroperca fusca*) in the Canary Islands, northwest coast of Africa. *Scientia Marina*, *74*(3) 613-619. http://doi.org/10.3989/scimar.2010.74n3613

Buxton, C.D. & Clarke, J.R. (1986) Age, growth and feeding of the blue hottentot *Pachymetopon aeneum* (Pisces: Sparidae) with notes on reproductive biology. *African Zoology, 21*(1), 33-38.

Buxton, C.D. & Garratt, P.A. (1990) Alternative reproductive styles in seabrearns (Pisces: Sparidae). In: M.N. Bruton (Ed.), *Alternative life-history styles of fishes* (pp. 113-124). Springer, Dordrecht. http://doi.org/10.1007/978-94-009-2065-1_7

Buxton, C.D. (1989) Protogynous hermaphroditism in *Chrysoblephus laticeps* (Cuvier) and *C. cristiceps* (Cuvier) (Teleostei: Sparidae). *African Zoology, 24*(3), 212-216.

Cabiddu, S., Follesa, M.C., Porcu, C. & Cau, A. (2010) Gonad development and reproduction in the monoecious species *Chlorophthalmus agassizi* (Actinopterygii: Aulopiformes: Chlorophthalmidae) from the Sardinian waters (Central-Western Mediterranean). *Acta Ichthyologica et Piscatoria, 40*, 167-177.

Calvo, J., Morriconi, E., Rae, G.A. & San Roman, N.A. (1992) Evidence of protandry in

a subantarctic notothenid, *Eleginops maclovinus* (Cuv. & Val., 1830) from the Beagle Channel, Argentina. *Journal of Fish Biology, 40*(2), 157-164. http://doi.org/10.1111/j.1095-8649.1992.tb02563.x

Candi, G., Castriota, L., Andaloro, F., Finoia, M.G. & Marino, G. (2004) Reproductive cycle and sex inversion in razor fish, a protogynous labrid in the southern Mediterranean Sea. *Journal of Fish Biology, 64*(6), 1498-1513. http://doi.org/10.1111/j.0022-1112.2004.0404.x

Chang, C.-F. & Yueh, W.-S. (1990) Annual cycle of gonadal histology and steroid profiles in the juvenile males and adult females of the protandrous black porgy, *Acanthopagrus schlegelii*. *Aquaculture,* 91, 179-196. http://doi.org/10.1016/0044-8486(90)90187-R

Chang-Po, C., Hwey-Lian, H. & Kun-Hsiung, C. (1980) Some aspects of the sex change and reproductive biology of the grouper, *Epinephelus diacanthus* (Cuvier et Valenciennes). *Bulletin of the Institute of Zoology, Academia Sinica, 19*(1)**,** 11-17.

Choat, J.H. (1969) Studies on the biology of labroid fishes (Labridae and Scaridae) at Heron Island, Great Barrier Reef. PhD Thesis, University of Queensland

Choat, J.H. & Robertson, D.R. (1975) Protogynous hermaphroditism in fishes of the family Scaridae. In: R. Reinboth (Ed.), *Intersexuality in the Animal Kingdom* (pp. 263-283). Springer, Berlin, Heidelberg. http://doi.org/10.1007/978-3-642-66069-6_26

Clark, E. & Shen, D. (1986) Territoriality of Red Sea sand-diving fishes of the genera *Xyrichtys* and *Trichonotus.* In: T. Uyeno, R. Arai, T. Taniuchi, & K., Matsuura, (Eds.), *Indo-Pacific Fish Biology: Proceedings of the Second International Conference on Indo-Pacific Fishes, Tokyo.*

Coates, D. (1982) Some observations on the sexuality of humbug damselfish, *Dascyllus aruanus* (Pisces, Pomacentridae) in the field. *Ethology, 59*(1), 7-18. http://doi.org/10.1111/j.1439-0310.1982.tb00328.x

Cole, K.S. & Hoese, D.F. (2001) Gonad morphology, colony demography and evidence for hermaphroditism in *Gobiodon okinawae* (Teleostei, Gobiidae). *Environmental Biology of Fishes, 61*(2), 161-173. http://doi.org/10.1023/A:1011032228716

Cole, K.S. & Robertson, R.D. (1988) Protogyny in the Caribbean reef goby, *Coryphopterus personatus*: gonad ontogeny and social influences on sex-change. *Bulletin of Marine Science, 42*(2), 317-333.

Cole, K.S. & Shapiro, D.Y. (1990) Gonad structure and hermaphroditism in the gobiid genus Coryphopterus (Teleostei: Gobiidae). *Copeia***,** *1990*(4), 996-1003. http://doi.org/10.2307/1446482

Cole, K.S. & Shapiro, D.Y. (1992) Gonadal structure and population characteristics of the protogynous goby *Coryphopterus glaucofraenum. Marine Biology, 113*(2), 1-9. http://doi.org/10.1007/BF00367632

Cole, K.S. & Shapiro, D.Y. (1995) Social facilitation and sensory mediation of adult sex change in a cryptic, benthic marine goby. *Journal of Experimental Marine Biology and Ecology, 186*(1), 65-75. http://doi.org/10.1016/0022-0981(94)00152-4

Cole, K.S. (1983) Protogynous hermaphroditism in a temperate zone territorial marine goby, *Coryphopterus nicholsi*. *Copeia, 1983*(3), 809-812. http://doi.org/10.2307/1444350

Cole, K.S. (1990) Patterns of gonad structure in hermaphroditic gobies (Teleostei Gobiidae). *Environmental Biology of Fishes, 28*(1-4), 125-142. http://doi.org/10.1007/BF00751032

Cole, K.S. (2002) Gonad morphology, sexual development, and colony composition in the obligate coral-dwelling damselfish *Dascyllus aruanus*. *Marine Biology,*

*140*(1), 151-163. http://doi.org/10.1007/s002270100681

Cole, K.S. (2003) Hermaphroditic characteristics of gonad morphology and inferences regarding reproductive biology in *Caracanthus* (Teleostei, Scorpaeniformes). *Copeia, 2003*(2), 68-80. http://doi.org/10.1643/0045-8511(2003)003[0068:HCOGMA]2.0.CO;2

Cole, K.S. (2010) Gonad morphology in hermaphroditic gobies. In: K.S. Cole (Ed.), *Reproduction and Sexuality in Marine Fishes: Patterns and Processes* (pp. 117–164). University of California Press, Berkeley, CA.

Coleman, F.C. (1981) Protogynous hermaphroditism in the anthiine serranid fish *Holanthias martinicensis. Copeia, 1981*(4), 893-895.

Coleman, F.C., Koenig, C.C. & Collins, L.A. (1996) Reproductive styles of shallow-water groupers (Pisces: Serranidae) in the eastern Gulf of Mexico and the consequences of fishing spawning aggregations. *Environmental Biology of Fishes, 47*(2), 129-141. http://doi.org/10.1007/BF00005035

Collins, L.A., Johnson, A.G., Koenig, C.C. & M.S., B.J. (1998) Reproductive patterns, sex ratio, and fecundity in gag, *Mycteroperca microlepis* (Serranidae), a protogynous grouper from the northeastern Gulf of Mexico. *Fishery Bulletin, 96*, 415-427.

Collins, M. R., Waltz, C. W., Roumillat, W. A., & Stubbs, D. L. (1987). Contribution to the life history and reproductive biology of gag, *Mycteroperca microlepis* (serrandidae), in the South Atlantic bight. *South Carolina State Documents Depository*.

Cowen, R.K. (1990) Sex change and life history patterns of the labrid, *Semicossyphus pulcher*, across an environmental gradient. *Copeia, 1990*(3), 787-795. http://doi.org/10.2307/1446444

Crabtree, R.E. & Bullock, L.H. (1998) Age, growth, and reproduction of black grouper, *Mycteroperca bonaci. Florida waters. Fishery Bulletin, 96*(4), 735-753.

Craig, M.T. (2007) Preliminary observations on the life history of the white-streaked grouper, *Epinephelus ongus*, from Okinawa, Japan. *Ichthyological Research, 54*(1), 81-84.

Craig, M.T., Hastings, P.A., Pondella, D.J., Ross Robertson, D. & Rosales-Casián, J.A. (2006) Phylogeography of the flag cabrilla *Epinephelus labriformis* (Serranidae): implications for the biogeography of the Tropical Eastern Pacific and the early stages of speciation in a marine shore fish. *Journal of Biogeography, 33*(6), 969-979. http://doi.org/10.1111/j.1365-2699.2006.01467.x

D'Ancona, U. (1949) Ermafroditismo ed intersessualita nei Teleostei. *Experientia, 5*, 381-389.

Dantchakoff, V. (1936) Sur les facteurs de l'histogenese chez des hermaphrodites. *Comptes Rendus de la Société de Biologie, 123*, 856-858.

David, G., Coutinho, R., Quagio-Grassiotto, I. & Verani, J. (2005) The reproductive biology of *Diplodus argenteus* (Sparidae) in the coastal upwelling system of Cabo Frio, Rio de Janeiro, Brazil. *African Journal of Marine Science*, *27*(2), 439-447.

Davis, M.P. & Fielitz, C. (2010) Estimating divergence times of lizardfishes and their allies (Euteleostei: Aulopiformes) and the timing of deep-sea adaptations. *Molecular Phylogenetics and Evolution, 57*, 1194-1208. http://doi.org/10.1016/j.ympev.2010.09.003

DeLoach, N. & Humann, P. (1999) Reef fish behavior: Florida, Caribbean, Bahamas. Jacksonville, FL. New World Publications.

DeMartini, E.E., Friedlander, A.M. & Holzwarth, S.R. (2005) Size at sex change in protogynous labroids, prey body size distributions, and apex predator densities at

NW Hawaiian atolls. *Marine Ecology Progress Series*, *297*, 259–271.

DeMartini, E.E., Everson, A.R. & Nichols, R.S. (2011) Estimates of body sizes at maturation and at sex change, and the spawning seasonality and sex ratio of the endemic Hawaiian grouper (H*yporthodus quernus*, F. Epinephelidae). *Fishery Bulletin, 109*(1)**,** 123-134.

Dipper, F.A. & Pullin, R.S.V. (1979) Gonochorism and sex-inversion in British Labridae (Pisces). *Journal of Zoology, 187*(1)**,** 97-112. http://doi.org/10.1111/j.1469-7998.1979.tb07716.x

Dulčić, J., Kraljević, M., Grbec, B. & Cetinić, P. (2000) Age, growth and mortality of blotched picarel *Spicara maena* L. (Pisces: Centracanthidae) in the eastern central Adriatic. *Fisheries Research, 48*(1), 69-78. http://doi.org/10.1016/S0165-7836(00)00112-0

Dulčić, J., Pallaoro, A., Cetinić, P., Kraljević, M., Soldo, A. & Jardas, I. (2003) Age, growth and mortality of picarel, *Spicara smaris* L. (Pisces: Centracanthidae), from the eastern Adriatic (Croatian coast). *Journal of Applied Ichthyology, 19*(1), 10-14. http://doi.org/10.1046/j.1439-0426.2003.00345.x

Ebisawa, A. (1990) Reproductive biology of *Lethrinus nebulosus* (Pisces; Lethrinidae) around the Okinawan waters [Japan]. *Bulletin of the Japanese Society of Scientific Fisheries (Japan)*, 56*(12),* 1941-1954. http://doi.org/10.2331/suisan.56.1941

Ebisawa, A. (1997) Some aspects of reproduction and sexuality in the spotcheek emperor, *Lethrinus rubrioperculatus*, in waters off the Ryukyu Islands. *Ichthyological Research, 44*(2-3)*,* 201-212.

Ebisawa, A. (1999) Reproductive and sexual characteristics in the Pacific yellowtail emperor, *Lethrinus atkinsoni*, in waters off the Ryukyu Islands. *Ichthyological Research, 46*(4)*,* 341-358.

Ebisawa, A. (2006) Reproductive and sexual characteristics in five Lethrinus species in waters off the Ryukyu Islands. *Ichthyological Research, 53*(3), 269-280. http://doi.org/10.1007/s10228-006-0345-3

Ebisawa, A., Kanashiro, K., Kyan, T. & Motonaga, F. (1995) Aspects of reproduction and sexuality in the black-spot tuskfish, *Choerodon schoenleinii*. *Japanese Journal of Ichthyology, 42*(1), 121-130. http://doi.org/10.11369/jji1950.42.121

El-Etreby, S.G., Roberts, C., Ghobashy, A.A. & Zyadah, M.A. (1993) Coral reef groupers (family: Serranidae) in south Sinai: reproduction and hermaphroditism in the fish *Cephalopholis hemistiktos* (Rüppell, 1830). *Journal of the Egyptian-German Society of Zoology (Anatomy and Embryology), 12(*B), 1-19.

Erisman, B.E. (2008) Reproductive biology and evolution of epinephelid and serranid fishes (Perciformes, Epinephelidae, Serranidae). PhD Thesis. Scripps Institute of Oceanography, University of California, San Diego.

Erisman, B.E., Craig, M.T. & Hastings, P.A. (2010) Reproductive biology of the Panama graysby *Cephalopholis panamensis* (Teleostei: Epinephelidae). *Journal of Fish Biology, 76*(6), 1312-1328.http://doi.org/10.1111/j.1095-8649.2010.02567.x

Etassami, S. (1983) Hermaphroditism in one Sparidae of the Persian Gulf: *Acanthopagrus bifasciatus* (Forssk.). *Cybium, 7*(2)*,* 87-91.

Fairclough, D.V., Edmonds, J.S., Lenanton, R.C., Jackson, G., Keay, I.S., Crisafulli, B.M. & Newman, S.J. (2011) Rapid and cost-effective assessment of connectivity among assemblages of *Choerodon rubescens* (Labridae), using laser ablation ICP-MS of sagittal otoliths. *Journal of Experimental Marine Biology and Ecology*, *403*(1-2), 46-53.

Ferreira, B.P. (1993) Reproduction of the inshore coral trout *Plectropomus maculates* (Perciformes: Serranidae) from the central Great Barrier Reef, Australia. *Journal*

of Fish Biology, *42*(6), 831-844. http://doi.org/10.1111/j.1095-8649.1993.tb00393.x

Ferreira, B.P. (1995) Reproduction of the common coral trout *Plectropomus leopardus* (Serranidae: Epinephelinae) from the central and northern Great Barrier Reef, Australia. *Bulletin of Marine Science, 56*(2), 653-669.

Fujii, T. (1970) Hermaphroditism and sex reversal in the fishes of the Platycephalidae - I. Sex reversal of *Onigocia macrolepis* (Bleeker). *Japanese Journal of Ichthyology*, *17*(1), 14-21.

Fischer, E.A. & Petersen, C.W. (1987) The evolution of sexual patterns in the seabasses. *Bioscience, 37*(7), 482-489. http://doi.org/10.2307/1310420

Fischer, E.A. (1980) The relationship between mating system and simultaneous hermaphroditism in the coral reef fish, *Hypoplectrus nigricans* (Serranidae). *Animal Behaviour, 28*(2), 620-633. http://doi.org/10.1016/S0003-3472(80)80070-4

Fischer, E.A. (1981) Sexual allocation in a simultaneously hermaphroditic coral reef fish. *The American Naturalist, 117*(1), 64-82. http://doi.org/10.1086/283686

Fischer, E.A. (1984) Egg trading in the chalk bass, *Serranus tortugarum*, a simultaneous hermaphrodite. *Zeitschrift für Tierpsychologie, 66*(2), 143-151. http://doi.org/10.1111/j.1439-0310.1984.tb01361.x

Fishelson, L. (1970) Protogynous sex reversal in the fish *Anthias squamipinnis* (Teleostei, Anthiidae) regulated by the presence or absence of a male fish. *Nature, 227*(5253), 90–91.

Fishelson, L. (1975) Ecology and physiology of sex reversal in *Anthias squamipinnis* (Peters),(Teleostei: Anthiidae). In: R. Reinboth (Ed.), *Intersexuality in the animal kingdom* (pp. 284-294). Springer, Berlin, Heidelberg. http://doi.org/10.1007/978-3-642-66069-6_27

Fishelson, L. (1989) Bisexuality and pedogenesis in gobies (Gobiidae: Teleostei) and other fish, or why so many fish in tropical seas? *Senckenbergiana Maritima, 20*, 147-170.

Fishelson, L. (1992) Comparative gonad morphology and sexuality of the Muraenidae (Pisces, Teleostei). *Copeia*, *1992*(1), 197-209. http://doi.org/10.2307/1446552

Francis, M.P., Harasti, D. & Malcolm, H.A. (2016) Surviving under pressure and protection: a review of the biology, ecology and population status of the highly vulnerable grouper *Epinephelus daemelii*. *Marine and Freshwater Research*. *67*(8), 1215-1228. http://dx.doi.org/10.1071/MF15099

Fricke, H. & Fricke, S. (1977) Monogamy and sex change by aggressive dominance in coral reef fish. *Nature, 266*(5606), 830-832. http://doi.org/10.1038/266830a0

Fricke, H.W. (1979) Mating system, resource defence and sex change in the anemonefish *Amphiprion akallopisos*. *Zeitschrift für Tierpsychologie, 50*(3), 313-326. http://doi.org/10.1111/j.1439-0310.1979.tb01034.x

Fricke, H.W. (1983) Social control of sex: field experiments with the anemonefish *Amphiprion bicinctus*. *Zeitschrift für Tierpsychologie, 61*(1), 71-77. http://doi.org/10.1111/j.1439-0310.1983.tb01327.x

García-Cagide, A. & García, T. (1996) Reproducción de Mycteroperca bonaci y M*ycteroperca venenosa* (Pisces: Serranidae) en la plataforma cubana. *Revista de Biología Tropical, 44*, 771-780.

García-Cagide, A.R., Claro, R. & García-Arteaga, J.P. (1999) Biología del bonací gato, *Mycteroperca tigris* (Pisces: Serranidae) en la plataforma SW de Cuba. I. Características generales y reproducción. *Revista de Investigaciones Marinas, 20*, 8-14.

García-Díaz, M.M., González, J.A., Lorente, M.J. & Tuset, V.M. (2006) Spawning season, maturity sizes, and fecundity in blacktail comber (*Serranus atricauda*) (Serranidae) from the eastern-central Atlantic. *Fishery Bulletin, 104*(4), 159-166.

García-Díaz, M.M., Lorente, M.J., González, J.A. & Tuset, V.M. (2002) Morphology of the ovotestis of *Serranus atricauda* (Teleostei, Serranidae). *Aquatic sciences, 64*(1), 87-96.

García-Díaz, M.M., Tuset, V.M., González, J.A. & Socorro, J. (1997) Sex and reproductive aspects in *Serranus cabrilla* (Osteichthyes: Serranidae): macroscopic and histological approaches. *Marine Biology, 127*(3), 379-386. http://doi.org/10.1007/s002270050024

Garratt, P.A. (1986) Protogynous hermaphroditism in the slinger, *Chrysoblephus puniceus* (Gilchrist & Thompson, 1908) (Teleosteir: Sparidae). *Journal of Fish Biology, 28*(3), 297-306. http://doi.org/10.1111/j.1095-8649.1986.tb05167.x

Ghorbel, M. (1996). Le pageot commun *Pagellus erythrinus* (poison, Sparidae) ecobiologie et etat d'exploitation dans le Golfe de Gabes. *PhD Thesis in Marine Biology and Oceanography, University of Sfax*.

Gillanders, B.M. (1995) Reproductive biology of the protogynous hermaphrodite *Achoerodus viridis* (Labridae) from south-eastern Australia. *Marine and Freshwater Research, 46*(7), 999-1008. http://doi.org/10.1071/MF9950999

Gilmore, G.R. & Jones, R.S. (1992) Color variation and associated behavior in the epinepheline groupers, *Mycteroperca microlepis* (Goode and Bean) and *M. phenax* Jordan and Swain. *Bulletin of Marine Science, 51*(1), 83-103.

Godwin, J. (1994a) Behavioural aspects of protandrous sex change in the anemonefish, *Amphiprion melanopus*, and endocrine correlates. *Animal Behaviour, 48*(3), 551-567. http://doi.org/10.1006/anbe.1994.1275

Godwin, J. (1994b) Historical aspects of protandrous sex change in the anemonefish *Amphiprion melanopus* (Pomacentridae, teleostei). *Journal of Zoology, 232*(2), 199-213. http://doi.org/10.1111/j.1469-7998.1994.tb01569.x

Godwin, J. (1995) Phylogenetic and habitat influences on mating system structure in the humbug damselfishes (Dascyllus, Pomacentridae). *Bulletin of Marine Science, 57*(3), 637-652.

Goeden, G.B. (1978) A monograph of the coral trout, *Plectropomus leopardus* (Lacepede). Research Bulletin No. 1, Queensland Fisheries Service, Brisbane.

Grandcourt, E.M. (2002) Demographic characteristics of a selection of exploited reef fish from the Seychelles: preliminary study. *Marine and Freshwater Research, 53*(2), 123-130. http://doi.org/10.1071/MF01123

Grandcourt, E.M., Al Abdessalaam, T.Z., Francis, F. & Al Shamsi, A.T. (2004) Biology and stock assessment of the Sparids, *Acanthopagrus bifasciatus* and *Argyrops spinifer* (Forsskål, 1775), in the Southern Arabian Gulf. *Fisheries Research, 69*(1), 7-20. http://doi.org/10.1016/j.fishres.2004.04.006

Guiguen, Y., Cauty, C.I., Fostier, A., Fuchs, J. & Jalabert, B. (1994) Reproductive cycle and sex inversion of the seabass, *Lates calcarifer,* reared in sea cages in French Polynesia: histological and morphometric description. *Environmental Biology of Fishes, 39*(3), 231-247. http://doi.org/10.1007/BF00005126

Hamzeh, S., Keivany, Y., Mahboobi Soofiani, N. & Jamshid-Aen, K. (2017) Reproductive Biology of Kingsoldier Bream (*Argyrops spinifer*) in the Persian Gulf. *Iranian Scientific Fisheries Journal, 26*(3), 91-104.

Harris, P.J., Wyanski, D.M., White, D.B. & Moore, J.L. (2002) Age, growth, and reproduction of scamp, *Mycteroperca phenax*, in the southwestern North Atlantic, 1979–1997. *Bulletin of Marine Science, 70*(1), 113-132.

Hassin, S., De Monbrison, D., Hanin, Y., Elizur, A., Zohar, Y. & Popper, D.M. (1997) Domestication of the white grouper, *Epinephelus aeneus* 1. Growth and reproduction. *Aquaculture, 156*(3-4), 305-316. http://doi.org/10.1016/S0044-8486(97)00136-1

Hastings, P.A. & Petersen, C.W. (1986) A novel sexual pattern in serranid fishes: simultaneous hermaphrodites and secondary males in *Serranus fasciatus*. *Environmental Biology of Fishes, 15*(1), 59-68. http://doi.org/10.1007/BF00005389

Hastings, P.A. (1981) Gonad morphology and sex succession in the protogynous hermaphrodite *Hemanthias vivanus* (Jordan and Swain). *Journal of Fish Biology, 18*(4), 443-454. http://doi.org/10.1111/j.1095-8649.1981.tb03785.x

Hesp, S.A., Potter, I.C. & Hall, N.G. (2004) Reproductive biology and protandrous hermaphroditism in *Acanthopagrus latus*. *Environmental Biology of Fishes, 70*(3), 257-272. http://doi.org/10.1023/B:EBFI.0000033344.21383.00

Hioki, S. & Suzuku, K. (1996) Sex changing from male to female on the way of protogynous process in three *Centropyge angelfishes* (Pomacanthidae: Teleostei). *Bulletin of institute of Oceanic Research and Development, Tokai University, 17,* 27-34.

Hioki, S., Tanaka, Y. & Suzuki, K. (1995) Reproductive behavior, eggs, larvae, and sexuality of two angelfishes, *Genicanthus watanabei* and *G. bellus,* in an aquarium. *Journal of the School of Marine Science and Technology Tokai University, 40*, 151-171.

Hodge, J.R., Santini, F. & Wainwright, P.C. (2020) Correlated evolution of sex allocation and mating system in wrasses and parrotfishes, *The American Naturalist*, *196*, 57–73.

Hoffman, S.G. (1980) *Sex-related social, mating, and foraging behavior in some sequentially hermaphroditic reef fishes.* PhD dissertation, University of California, Santa Barbara, CA.

Hoffman, S.G. (1983) Sex-related foraging behavior in sequentially hermaphroditic hogfishes (*Bodianus* spp.). *Ecology, 64*(4), 798-808. http://doi.org/10.2307/1937203

Hoffman, S.G. (1985) Effects of size and sex on the social organization of reef-associated hogfishes, *Bodianus* spp. *Environmental Biology of Fishes*, *14*(2), 185-197.

Horvath, M.L., Grimes, C.B. & Huntsman, G.R. (1990) Growth, mortality, reproduction and feeding of knobbed porgy, *Calamus nodosus*, along the southeastern United States coast. *Bulletin of Marine Science, 46*(3), 677-687.

Hourigan, T.F. & Kelley, C.D. (1985) Histology of the gonads and observations on the social behavior of the Caribbean angelfish *Holacanthus tricolor*. *Marine Biology, 88*(3), 311-322. http://doi.org/10.1007/BF00392592

Howard, J. & Randali, E. (1986) A review of the parrotfishes (family Scaridae) of the Great Barrier Reef of Australia with description of a new species. *Records of the Australian Museum, 38*, 175-228.

Ismail, R.F., Mourad, M.M. & Farrag, M.M.S. (2018) Gonadal development and hermaphroditism of bluespotted seabream, *Pagrus caeruleostictus* (Valenciennes, 1830) from the Mediterranean Sea, Egypt. *The Egyptian Journal of Aquatic Research, 44*(2), 163-171. http://doi.org/10.1016/j.ejar.2018.05.003

Kadota, T., Osato, J., Nagata, K. & Sakai, Y. (2012) Reversed sex change in the haremic protogynous hawkfish *Cirrhitichthys falco* in natural conditions. *Ethology, 118*(3), 226-234.

Kawaguchi, K. & Marumo, R. (1967) Biology of *Gonostoma gracile* (Gonostomatidae)

I. Morphology, life history and sex reversal. *Inform. Bull. Planktology Japan, 1967,* 53-69.

Kline, R.J., Khan, I.A. & Holt, G.J. (2011) Behavior, color change and time for sexual inversion in the protogynous grouper (*Epinephelus adscensionis*). *PLoS One, 6*(5), e19576. http://doi.org/10.1371/journal.pone.0019576

Kožul, V., Glavić, N., Tutman, P., Bolotin, J. & Onofri, V. (2011) The spawning, embryonic and early larval development of the green wrasse *Labrus viridis* (Linnaeus, 1758)(Labridae) in controlled conditions. *Animal reproduction science. 125*(1-4), 196-203. http://doi.org/10.1016/j.anireprosci.2011.01.013

Kobayashi, K. & Suzuki, K. (1992) Hermaphroditism and sexual function in *Cirrhitichthys aureus* and the other Japanese hawkfishes (Cirrhitidae: Teleostei). *Japanese Journal of Ichthyology, 38*(4), 397-410. http://doi.org/https://doi.org/10.11369/jji1950.38.397

Kokokiris, L., Bruslé, S., Kentouri, M. & Fostier, A. (1999) Sexual maturity and hermaphroditism of the red porgy *Pagrus* (Teleostei: Sparidae). *Marine Biology, 134*(4), 621-629. http://doi.org/10.1007/s002270050577

Krug, H.M. (1990) The Azorean blackspot seabream, *Pagellus bogaraveo* (Brünnich, 1768) (Teleostei, Sparidae). Reproductive cycle, hermaphroditism, maturity and fecundity. *Cybium, 14*(2), 151-159.

Kusen, J.D. (1991) Protogynous hermaphroditism in the sand diver *Trichonotus filamentosus. Bulletin of the Japanese Society for the Science of Fish, 57*(1), 35-40. http://doi.org/10.2331/suisan.57.35

Kuwamura, T., Nakashimn, Y. & Yogo, Y. (1994) Sex change in either direction by growth-rate advantage in the monogamous coral goby, *Paragobiodon echinocephalus. Behavioral Ecology, 5*(4), 434-438. http://doi.org/10.1093/beheco/5.4.434

Kuwamura, T., Suzuki, S. & Kadota, T. (2016) Male-to-female sex change in widowed males of the protogynous damselfish *Dascyllus aruanus. Journal of ethology, 34*(1), 85-88. http://doi.org/10.1007/s10164-015-0450-8

Kuwamura, T., Tanaka, N., Nakashima, Y., Karino, K. & Sakai, Y. (2002) Reversed sex-change in the protogynous reef fish *Labroides dimidiatus. Ethology, 108*(5), 443-450. http://doi.org/10.1046/j.1439-0310.2002.00791.x

Lamrini, A. (1986) Sexualité de *Pagellus acarne* (Risso, 1826) (Teleosteen Sparidae) de la côte Atlantique méridionale du Maroc (21°-26° N). *Cybium (Paris), 10*(1), 3-14.

Lavenda, N. (1949) Sexual differences and normal protogynous hermaphroditism in the Atlantic sea bass, *Centropristes striatus. Copeia, 1949*(3), 185-194. http://doi.org/10.2307/1438985

Lee, Y.D. (1995) Sexual phenomenon of protogynous serranid fish. *Exploited Marine Resources,* 1, 143-150.

Lee, Y.H., Du, J.L., Yueh, W.S., Lin, B.Y., Huang, J.D., Lee, C.Y., Lee, M.F., Lau, E.L., Lee, F.Y. & Morrey, C. (2001) Sex change in the protandrous black porgy, *Acanthopagrus schlegeli*: a review in gonadal development, estradiol, estrogen receptor, aromatase activity and gonadotropin. *Journal of experimental Zoology, 290*(7), 715-726. http://doi.org/10.1002/jez.1122

Leem, J.B., Sakamoto, K., Tsuruda, Y. & Nakazono, A. (1998) Sexual pattern of the labrid fishes collected from Kuchinoerabu-jima, Kagoshima, Japan. *J Journal of the Faculty of Agriculture Kyushu University, 42*(3-4), 409-420.

Leonard, J.L. (1993) Sexual conflict in simultaneous hermaphrodites: evidence from serranid fishes. *Environmental Biology of Fishes, 36*(2), 135-148.

Liem, K.F. (1963) Sex reversal as a natural process in the synbranchiform fish *Monopterus albus*. *Copeia 1963*(2), 303-312. http://doi.org/10.2307/1441348

Liem, K.F. (1968) Geographical and taxonomic variation in the pattern of natural sex reversal in the teleost fish order Synbranchiformes. *Journal of Zoology, 156*(2), 225-238. http://doi.org/10.1111/j.1469-7998.1968.tb05930.x

Lin, Y.-J., Roa-Ureta, R.H., Rabaoui, L., Grandcourt, E.M., Maneja, R.H., Al-Abdulkader, K. & Qurban, M.A. (2019) Association to vegetated habitats and different vulnerability to habitat degradation for two fish species, *Epinephelus areolatus* (Serranidae) and *Siganus canaliculatus* (Siganidae), from the western Arabian Gulf. *Marine pollution bulletin*, *141*, 482-492. https://doi.org/10.1016/j.marpolbul.2019.03.011

Lissia-Frau, A.M. & Casu, S. (1968) Il differenziamento sessuale di *Lithognathus mormyrus* (L.) e di *Oblada melanura* (L.). *Studi Sassaresi, 46*, 1-19.

Lissia-Frau, A.M., Pala, M. & Casu, S. (1976) Observations and considerations on protandrous hermaphroditism in some species of sparid fishes (Teleostei, Perciformes). *Studi Sassaresi,* 54, 147-167.

Liu, M. & Sadovy, Y. (2004a) Early gonadal development and primary males in the protogynous epinepheline, *Cephalopholis boenak*. *Journal of Fish Biology, 65*(4), 987-1002. http://doi.org/10.1111/j.0022-1112.2004.00503.x

Liu, M. & Sadovy, Y. (2004b) The influence of social factors on adult sex change and juvenile sexual differentiation in a diandric, protogynous epinepheline, *Cephalopholis boenak* (Pisces, Serranidae). *Journal of Zoology*, *264*(3), 239-248. http://doi.org/10.1017/S0952836904005631

Liu, M., Wang, Y.-Y., Shan, X.-J., Kang, B. and Ding, S.-X. (2016) Primary male development of two sequentially hermaphroditic groupers, *Epinephelus akaara* and *Epinephelus awoara* (Perciformes: Epinephelidae). *Journal of Fish Biology*, *88*(4), 1598-1613.

Lloret, J., Muñoz, M. & Casadevall, M. (2012) Threats posed by artisanal fisheries to the reproduction of coastal fish species in a Mediterranean marine protected area. *Estuarine, Coastal and Shelf Science*, *113*, 133-140. http://dx.doi.org/10.1016/j.ecss.2012.07.015

Longley, W.H. & Hildebrand, S.F. (1941) *Systematic catalogue of the fishes of Tortugas, Florida: with observations on color, habits, and local distribution*. Papers Tortugas Laboratory, *34*, 1-331.

Lorenzo, J.M., Pajuelo, J.G., Méndez-Villamil, M., Coca, J. & Ramos, A.G. (2002) Age, growth, reproduction and mortality of the striped seabream, *Lithognathus mormyrus* (Pisces, Sparidae), off the Canary Islands (central-east Atlantic). *Journal of Applied Ichthyology, 18*(3), 204-209. http://doi.org/10.1046/j.1439-0426.2002.00318.x

Lou, D.C. (1992) *Age specific patterns of growth and reproduction in tropical herbivorous fishes.* PhD Thesis. James Cook University, Queensland, Australia.

Lucks, D.K. (1970) *Aspects of the biology of the white steenbras (Lithognathus aureti Smith, 1962) in the Sandwich estuary.* PhD dissertation. University of Stellenbosch, South Africa. 49 pp.

Lutnesky, M.M.F. (1994) Density-dependent protogynous sex change in territorial-haremic fishes: models and evidence. *Behavioral Ecology, 5*(4), 375-383. http://doi.org/10.1093/beheco/5.4.375

Lutnesky, M.M.F. (1996) Size-dependent rate of protogynous sex change in the pomacanthid angelfish, *Centropyge potteri*. *Copeia 1996*(1), 209-212. http://doi.org/10.2307/1446961

Mackie, M. (2000) Reproductive biology of the halfmoon grouper, *Epinephelus rivulatus,* at Ningaloo Reef, Western Australia. *Environmental Biology of Fishes, 57*(4)*,* 363-376. http://doi.org/10.1023/A:1007658027359

Mackie, M.C. (2003) Socially controlled sex-change in the half-moon grouper, *Epinephelus rivulatus*, at Ningaloo Reef, Western Australia. *Coral Reefs, 22*(2)*,* 133-142. http://doi.org/10.1007/s00338-003-0296-3

Mackie, M.C. (2006) Anatomical changes to the gonad during protogynous sex change in the half-moon grouper *Epinephelus rivulatus* (Valenciennes). *Journal of Fish Biology, 69*(1)*,* 176-186. http://doi.org/10.1111/j.1095-8649.2006.01084.x

Mackie, M. (2007) Reproductive behavior of the halfmoon grouper, *Epinephelus rivulatus*, at Ningaloo Reef, Western Australia. *Ichthyological Research, 54*(3), 213-220. http://doi.org/10.1007/s10228-006-0393-8

Madhu, K. & Madhu, R. (2006) Protandrous hermaphroditism in the clown fish *Amphiprion percula* from Andaman and Nicobar islands. *Indian Journal of Fisheries, 53*(4)*,* 373-382.

Manabe, H., Ishimura, M., Shinomiya, A. & Sunobe, T. (2007a) Field evidence for bi-directional sex change in the polygynous gobiid fish *Trimma okinawae*. *Journal of Fish Biology, 70*(2), 600-609. http://doi.org/10.1111/j.1095-8649.2007.01338.x

Manabe, H., Ishimura, M., Shinomiya, A. & Sunobe, T. (2007b) Inter-group movement of females of the polygynous gobiid fish *Trimma okinawae* in relation to timing of protogynous sex change. *Journal of ethology, 25*(2)*,* 133-137. http://doi.org/10.1007/s10164-006-0007-y

Manabe, H., Toyoda, K., Nagamoto, K., Dewa, S.-I., Sakurai, M., Hagiwara, K., Shinomiya, A. & Sunobe, T. (2013) Bidirectional sex change in seven species of *Priolepis* (Actinopterygii: Gobiidae). *Bulletin of Marine Science, 89*(2), 635-642.

Marino, G., Azzurro, E., Massari, A., Finoia, M.G. & Mandich, A. (2001) Reproduction in the dusky grouper from the southern Mediterranean. *Journal of Fish Biology, 58*(4), 909-927. http://doi.org/10.1111/j.1095-8649.2001.tb00544.x

Mathew, G. & Kuruvilla, M. (2010) Anatomical changes during early gonad development in the protogynous greasy grouper *Epinephelus tauvina* (Forsskal). *Indian Journal of Fisheries, 57*(2), 21-24.

Matos-Caraballo, D., Posada, J.M. & Luckhurst, B.E. (2006) Fishery-dependent evaluation of a spawning aggregation of tiger grouper (*Mycteroperca tigris*) at Vieques Island, Puerto Rico. *Bulletin of Marine Science, 79*(1), 1-16.

Maxfield, J.M. & Cole, K.S. (2019) Structural changes in the ovotestis of the bidirectional hermaphrodite, the blue-banded goby (*Lythrypnus dalli*), during transition from ova production to sperm production. *Environmental Biology of Fishes, 102*(11)*,* 1393-1404. http://doi.org/10.1007/s10641-019-00914-2

McBride, R.S. & Johnson, M.R. (2007) Sexual development and reproductive seasonality of hogfish (Labridae: *Lachnolaimus maximus*), an hermaphroditic reef fish. *Journal of Fish Biology, 71*(5)*,* 1270-1292. http://doi.org/10.1111/j.1095-8649.2007.01580.x

McBride, R.S., Sulak, K.J., Thurman, P.E. & Richardson, A.K. (2009) Age, growth, mortality, and reproduction of Roughtongue bass, *Pronotogrammus martinicensis* (Serranidae), in the northeastern Gulf of Mexico. *Gulf of Mexico Science, 27*(1), 4. 10.18785/goms.2701.04

McErlean, A.J. & Smith, C.L. (1964) The age of sexual succession in the protogynous hermaphrodite *Mycteroperca microlepis*. *Transactions of the American Fisheries Society, 93*(3), 301-302. http://doi.org/10.1577/1548-8659(1964)93[301:TAOSSI]2.0.CO;2

McIlwain, J., Hermosa, G.V., Claereboudt, M., Al-Oufi, H.S. & Al-Awi, M. (2006) Spawning and reproductive patterns of six exploited finfish species from the Arabian Sea, Sultanate of Oman. *Journal of Applied Ichthyology, 22*(2)*, 167-176. http://doi.org/10.1111/j.1439-0426.2006.00723.x

Mead, G.W. (1959) Hermaphroditism in archibenthic and pelagic fishes of the order Iniomi. *Deep Sea Research (1953), 6,* 234-IN19.

Merrett, N.R., Badcock, J. & Herring, P.J. (1973) The status of *Benthalbella infans* (Pisces: Myctophoidei), its development, bioluminescence, general biology and distribution in the eastern North Atlantic. *Journal of Zoology, 170*(1), 1-48. http://doi.org/10.1111/j.1469-7998.1973.tb05042.x

Meyer, K.A. (1977) Reproductive behavior and patterns of sexuality in the Japanese labrid fish *Thalassoma cupido*. *Japanese Journal of Ichthyology, 24*(2), 101-112. http://doi.org/10.11369/jji1950.24.101

Micale, V. & Perdichizzi, F. (1994) Further studies on the sexuality of the hermaphroditic teleost *Diplodus sargus*, with particular reference to protandrous sex inversion. *Journal of Fish Biology, 45*(4), 661-670.http://doi.org/10.1111/j.1095-8649.1994.tb00932.x

Micale, V., Maricchiolo, G. & Genovese, L. (2002) The reproductive biology of blackspot sea bream *Pagellus bogaraveo* in captivity. I. Gonadal development, maturation and hermaphroditism. *Journal of Applied Ichthyology, 18*(3), 172-176. http://doi.org/10.1046/j.1439-0426.2002.00328.x

Micale, V., Perdichizzi, F. & Santangelo, G. (1987) The gonadal cycle of captive white bream, *Diplodus sargus* (L.). *Journal of Fish Biology, 31*(3), 435-440. http://doi.org/10.1111/j.1095-8649.1987.tb05247.x

Moore, R. (1979) Natural sex inversion in the giant perch (*Lates calcarifer*). *Marine and Freshwater Research, 30*(6), 803-813. http://doi.org/10.1071/MF9790803

Moss, J.W., Adams, S. & Welch, D.J. (2002) Bommie cod (*Cephalopholis cyanostigma*): a big surprise from a little fish. Bridging the gap: A workshop linking student research with fisheries stakeholde*rs*, pp. 94-107.

Moussac, G.D. (1986) Mise en évidence de l'hermaphrodisme protogyne d'*Epinephelus chlorostigma* (Valenciennes, 1828) aux Seychelles (Pisces, Serranidae). *Cybium, 10*(3), 249-262.

Moyer, J.T. & Nakazono, A. (1978) Protandrous hermaphroditism in six species of the anemonefish genus Amphiprion in Japan. *Japanese Journal of Ichthyology, 25*(2), 101-106.http://doi.org/10.11369/jji1950.25.101

Moyer, J.T. & Zaiser, M.J. (1984) Early sex change: a possible mating strategy of Centropyge angelfishes (Pisces: Pomacanthidae). *Journal of ethology, 2*(1)*, 63-67. http://doi.org/10.1007/BF02348208

Munday, P.L., Caley, M.J. & Jones, G.P. (1998) Bi-directional sex change in a coral-dwelling goby. *Behavioral Ecology and Sociobiology, 43*(6), 371-377.http://doi.org/10.1007/s002650050504

Nagelkerken, W.P. (1979) Biology of the graysby, *Epinephelus cruentatus*, of the coral reef of Curaçao. *Studies on the fauna of Curaçao and other Caribbean Islands, 60*(1), 1-118.

Nakai, T. & Sano, M. (2002) Evidence of protogynous hermaphroditism in the darkfin hind *Cephalopholis urodeta* (Serranidae) at Iriomote Island, southern Japan. *Fisheries Science, 68*(3)*, 697-699. http://doi.org/10.1046/j.1444-2906.2002.00479.x

Nakashima, Y., Kuwamura, T. & Yogo, Y. (1995) Why Be a Both-ways Sex Changer? *Ethology, 101*(4), 301-307. http://doi.org/10.1111/j.1439-0310.1995.tb00367.x

Nakashima, Y., Kuwamura, T. & Yogo, Y. (1996) Both-ways sex change in monogamous coral gobies, Gobiodon spp. *Environmental Biology of Fishes, 46*(3), 281-288. http://doi.org/10.1007/BF00005004

Nakazono, A. & Kusen, J.D. (1991) Protogynous hermaphroditism in the wrasse *Choerodon azurio*. *Nippon Suisan Gakkaishi, 57*(3), 417-420. http://doi.org/10.2331/suisan.57.417

Nakazono, A. (1979) Studies on the sex reversal and spawning behavior of five species of Japanese labrid fishes. *Report of Fishery Research Laboratory, Kyushu University, 4*, 1-64.

Ndiaye, A.M. (2014) Etude du cycle sexuel et l'inversion sexuelle de *Pagellus bellottii* (Téléostéen: Sparidae) dans les eaux sénégalaises. *Afrique Science: Revue Internationale des Sciences et Technologie, 10*(4), 257-266.

Nemtzov, S.C. (1985) Social control of sex change in the Red Sea razorfish *Xyrichtys pentadactylus* (Teleostei, Labridae). *Environmental Biology of Fishes, 14*(2-3), 199-211. http://doi.org/10.1007/BF00000827Nemtzov, S.C. (2007) Intraspecific variation in home range exclusivity by female green razorfish, Xyrichtys splendens (family Labridae), in different habitats. *Environmental Biology of Fishes* 50, 371–381. https://doi.org/10.1023/A:1007319622028

Oh, S.-R., Kang, H.-C., Lee, C.-H., Hur, S.-W. & Lee, Y.-D. (2013) Sex reversal and masculinization according to growth in longtooth grouper *Epinephelus bruneus*. *Development & Reproduction, 17*(2), 79-85. http://doi.org/10.12717/DR.2013.17.2.079

Ohta, I., Akita, Y., Uehara, M. & Ebisawa, A. (2017) Age-based demography and reproductive biology of three *Epinephelus* groupers, *E. polyphekadion*, *E. tauvina*, and *E. howlandi* (Serranidae), inhabiting coral reefs in Okinawa. *Environmental Biology of Fishes*, *100*(11), 1451-1467. http://doi.org/10.1007/s10641-017-0655-5

Ohta, I. & Ebisawa, A. (2015) Reproductive biology and spawning aggregation fishing of the white-streaked grouper, *Epinephelus ongus,* associated with seasonal and lunar cycles. *Environmental Biology of Fishes, 98*(6), 1555-1570. http://doi.org/10.1007/s10641-015-0382-8

Ohta, K., Sundaray, J.K., Okida, T., Sakai, M., Kitano, T., Yamaguchi, A., Takeda, T. & Matsuyama, M. (2003) Bi-directional sex change and its steroidogenesis in the wrasse, *Pseudolabrus sieboldi. Fish physiology and biochemistry, 28*(1-4), 173-174. http://doi.org/10.1023/B:FISH.0000030517.06738.e7

Okada, Y.K. (1965a) Bisexuality in Sparid Fishes I. Origin of bisexual gonads in *Mylio macrocephalus. Proceedings of the Japan Academy, 41*(4), 294-299. http://doi.org/10.2183/pjab1945.41.294

Okada, Y.K. (1965b) Bisexuality in Sparid Fishes II. *Proceedings of the Japan Academy, 41*(4), 300-304. http://doi.org/10.2183/pjab1945.41.300

Okada, Y.K. (1966) Sex reversal in *Inegocia meerdervoorti* with special reference to repetition of hermaphroditic state. *Proceedings of the Japan Academy, 42*(5), 497-502. http://doi.org/10.2183/pjab1945.42.497

Okumura, S. (2001) Evidence of sex reversal towards both directions in reared red spotted grouper *Epinephelus akaara*. *Fisheries Science, 67*(3), 535-537. http://doi.org/10.1046/j.1444-2906.2001.00287.x

Ota, K., Kobayashi, T., Ueno, K. & Gojobori, T. (2000) Evolution of heteromorphic sex chromosomes in the order Aulopiformes. *Gene, 259*(1-2), 25-30. http://doi.org/10.1016/S0378-1119(00)00460-1

Paiva, R.B., Neves, A., Sequeira, V., Vieira, A.R., Costa, M.J. & Gordo, L.S. (2018) Age,

growth and reproduction of the protandrous hermaphrodite fish, *Sarpa salpa*, from the Portuguese continental coast. *Journal of the Marine Biological Association of the United Kingdom, 98*(2), 269-281. http://doi.org/10.1017/S0025315416001405

Pajuelo, J.G. & Lorenzo, J.M. (1995) Biological parameters reflecting the current state of the exploited pink dentex, *Dentex gibbosus* (Pisces: Sparidae) population off the Canary Islands. *South African Journal of Marine Science, 16*(1), 311-319. http://doi.org/10.2989/025776195784156421

Pajuelo, J.G. & Lorenzo, J.M. (1998) Population biology of the common pandora *Pagellus erythrinus* (Pisces: Sparidae) off the Canary Islands. *Fisheries Research, 36*(2-3)**,** 75-86. http://doi.org/10.1016/S0165-7836(98)00110-6

Pajuelo, J.G. & Lorenzo, J.M. (1999) Life history of black seabream, *Spondyliosoma cantharus*, off the Canary Islands, Central-east Atlantic. *Environmental Biology of Fishes, 54*(3), 325-336. http://doi.org/10.1023/A:1007515301745

Pajuelo, J.G. & Lorenzo, J.M. (2001) Biology of the annular seabream, *Diplodus annularis* (Sparidae), in coastal waters of the Canary Islands. *Journal of Applied Ichthyology, 17*(3), 121-125. http://doi.org/10.1046/j.1439-0426.2001.00292.x

Pajuelo, J.G. & Lorenzo, J.M. (2004) Basic characteristics of the population dynamic and state of exploitation of Moroccan white seabream *Diplodus sargus cadenati* (Sparidae) in the Canarian archipelago. *Journal of Applied Ichthyology, 20*(1), 15-21. http://doi.org/10.1046/j.0175-8659.2003.00540.x

Parvez, K. & Al-Marzouk, A. (2000) First observations on natural sex reversal in a protandrous bream (*Sparidentex hasta*: Sparidae) from Kuwait. *Pakistan Journal of Zoology, 32*(3), 229-244.

Pears, R.J., Choat, J.H., Mapstone, B.D. & Begg, G.A. (2006) Demography of a large grouper, *Epinephelus fuscoguttatus*, from Australia's Great Barrier Reef: implications for fishery management. *Marine Ecology Progress Series, 307*, 259-272.

Pears, R.J., Choat, J.H., Mapstone, B.D. & Begg, G.A. (2007) Reproductive biology of a large, aggregation-spawning serranid, *Epinephelus fuscoguttatus* (Forsskål): management implications. *Journal of Fish Biology, 71*(3)*,* 795-817. http://doi.org/10.1111/j.1095-8649.2007.01545.x

Pember, M.B. (2006) *Characteristics of fish communites in coastal waters of north-western Australia, including the biology of the threadfin species Eleutheronema tetradactylum and Polydactylus macrochir.* PhD Thesis, Murdoch University.

Pember, M.B., Newman, S.J., Hesp, S.A., Young, G.C., Skepper, C.L., Hall, N.G. & Potter, I.C. (2005) *Biological parameters for managing the fisheries for blue and king threadfin salmons, estuary rockcod, Malabar grouper and mangrove jack in north-western Australia*. Fisheries Research and Development Corporation Report Murdoch University, Western Australia.

Penrith, M.J. (1972) Sex reversal in the sparid fish *Chrysoblephus laticeps*. *Koedoe, 15*(1), 135-139. http://doi.org/10.4102/koedoe.v15i1.679

Petersen, C.W. & Fischer, E.A. (1986) Mating system of the hermaphroditic coral-reef fish, *Serranus baldwini. Behavioral Ecology and Sociobiology, 19*(3), 171-178. http://doi.org/10.1007/BF00300857

Petersen, C.W. (1991) Sex allocation in hermaphroditic sea basses. *The American Naturalist, 138*(3), 650-667. http://doi.org/10.1086/285240

Petersen, C.W. (1995) Reproductive behavior, egg trading, and correlates of male mating success in the simultaneous hermaphrodite, *Serranus tabacarius. Environmental Biology of Fishes, 43*(4), 351-361. http://doi.org/10.1007/BF00001169

Platten, J., Tibbetts, I. and Sheaves, M. (2002) The influence of increased line-fishing

mortality on the sex ratio and age of sex reversal of the venus tusk fish. *Journal of Fish Biology*, *60*(2), 301–318.

Pollock, B.R. (1985) The reproductive cycle of yellowfin bream, *Acanthopagrus australis* (Günther), with particular reference to protandrous sex inversion. *Journal of Fish Biology, 26*(3), 301-311. http://doi.org/10.1111/j.1095-8649.1985.tb04269.x

Pressley, P.H. (1981) Pair formation and joint territoriality in a simultaneous hermaphrodite: the coral reef fish *Serranus tigrinus*. *Zeitschrift für Tierpsychologie, 56*(1)*, 33-46. http://doi.org/10.1111/j.1439-0310.1981.tb01282.x

Ramirez, J. (1984) *Aspects of the reproductive biology of three morphospecies in the genus Hypoplectrus; H. chlorurus, H. puella, H. unicolor.* Unpublished MS Thesis, University of Puerto Rico, Mayagüez, Puerto Rico.

Randall, J.E. & Bruce, R.W. (1983) The parrotfishes of the subfamily Scarinae of the western Indian Ocean with descriptions of three new species. *Ichthyological Bulletin 47*, 1-19.

Reinboth, R. (1962) Morphologische und funktionelle Zweigeschlechtigkeit bei marinen Teleostiern (Serranidae, Sparidae, Centracanthidae, Labridae). *Zoologische Jahrbücher Physiologie Bd, 69,* 405–480.

Reinboth, R. (1963) Naturlicher geschlechtswechsel bei *Sacura margaritacea* (Hilgendorf) (Serranidae). *Japanese Annotnes Zoology, 36*(4), 173-178.

Reinboth, R. (1964) Inversion du sexe chez *Anthias* (L.) (Serranidae). *Vie Milieu, 17*, 499-503.

Reinboth, R. (1970) Intersexuality in fishes. *Mem. Soc Endocrinology***,** *18*,515-543.

Reinboth, R., Becker, B. & Latz, M. (1986) In vitro studies on steroid metabolism by gonadal tissues from ambisexual teleosts: II. Conversion of [14C] androstenedione by the heterologous gonadal tissues of the protandric sea bream *Pagellus acarne* (Risso). *General and comparative endocrinology, 62*(2), 335-340. http://doi.org/10.1016/0016-6480(86)90124-3

Reñones, O., Grau, A., Mas, X., Riera, F. & Saborido-Rey, F. (2010) Reproductive pattern of an exploited dusky grouper *Epinephelus marginatus* (Lowe 1834) (Pisces: Serranidae) population in the western Mediterranean. *Scientia Marina, 74*(3), 523-537.

Roberts Jr, D.E. & Schlieder, R.A. (1983) Induced sex inversion, maturation, spawning and embryogeny of the protogynous grouper, *Mycteroperca microlepis*. *Journal of the World Mariculture Society, 14*(1-4)*, 637-649. http://doi.org/10.1111/j.1749-7345.1983.tb00116.x

Robertson, D.R. & Choat (1974) Protogynous hermaphroditism and social systems in labrid fish. *Proc. 2nd Inter. Coral Reef Symp., 1974*(1), pp. 217-225.

Robertson, D.R. & Justines, G. (1982) Protogynous hermaphroditism and gonochorism in four Caribbean reef gobies. *Environmental Biology of Fishes, 7*(2), 137-142. http://doi.org/10.1007/BF00001783

Robertson, D.R. & Warner, R.R. (1978) Sexual patterns in the labroid fishes of the Western Caribbean, II. The Parrotfishes (Scaridae). *Smithsonian Contributions to Zoology, 255*, 1-26.

Robertson, D.R. (1972) Social control of sex reversal in a coral-reef fish. *Science, 177*(4053), 1007-1009. http://doi.org/10.1126/science.177.4053.1007

Robertson, R.D., Reinboth, R. & Bruce, R.W. (1982) Gonochorism, protogynous sex-change and spawning in three sparisomatinine parrotfishes from the western Indian Ocean. *Bulletin of Marine Science, 32*(4), 868-879.

Robinson, G.A. (1976) Sex reversal in the dageraad *Chrysoblephus cristiceps* (Pisces:

Sparidae). *Koedoe, 19*(1), 43-48. http://doi.org/10.4102/koedoe.v19i1.1180

Rodgers, E.W., Drane, S. & Grober, M.S. (2005) Sex reversal in pairs of *Lythrypnus dalli*: behavioral and morphological changes. *The Biological Bulletin, 208*(2)*,* 120-126. http://doi.org/10.2307/3593120

Roede, M.J. (1972) Color as related to size, sex and behavior in seven Caribbean Labrid fish species (Genera *Thalassoma*, *Halichoeres* and *Hemipteronotus*). *Studies on the Fauna of Curacao and other Caribbean Islands 32*, 1–266.

Ross, R.M. (1982) *Sex change in the endemic Hawaiian labrid Thalassoma duperrey (Quoy and Gaimard): a behavioral and ecological analysis*. Unpublished PhD dissertation, University of Hawaii.

Ross, R.M. (1984) Anatomical changes associated with sex reversal in the fish *Thalassoma duperrey* (Teleostei: Labridae). *Copeia, 1984*(1), 245-248. http://doi.org/10.2307/1445069

Ryen, C. (2007) Sex-specific growth dynamics in protogynous hermaphrodites. Master's Thesis. James Cook University, Queensland, Australia.

Sadovy de Mitcheson, Y., Liu, M. & Suharti, S. (2010) Gonadal development in a giant threatened reef fish, the humphead wrasse *Cheilinus undulatus*, and its relationship to international trade. *Journal of Fish Biology, 77*(3), 706-718. http://doi.org/10.1111/j.1095-8649.2010.02714.x

Sadovy, Y. & Donaldson, T.J. (1995) Sexual pattern of *Neocirrhites armatus* (Cirrhitidae) with notes on other hawkish species. *Environmental Biology of Fishes, 42*(2)*,* 143-150. http://doi.org/10.1007/BF00001992

Sadovy, Y., Kulbicki, M., Labrosse, P., Letourneur, Y., Lokani, P. & Donaldson, T.J. (2003) The humphead wrasse, *Cheilinus undulatus*: synopsis of a threatened and poorly known giant coral reef fish. *Reviews in Fish Biology and Fisheries, 13*(3), 327-364. http://doi.org/10.1023/B:RFBF.0000033122.90679.97

Sadovy, Y., Rosario, A. & Román, A. (1994) *Reproduction in an aggregating grouper, the red hind*, *Epinephelus guttatus*. In: E.K. Balon, M.N. Bruton & D.L.G. Noakes (Eds.), Women in ichthyology: an anthology in honour of ET, Ro and Genie (pp. 269-286). Springer, Dordrecht. http://doi.org/10.1007/978-94-011-0199-8_21

Sakai, Y. (1997) Alternative spawning tactics of female angelfish according to two different contexts of sex change. *Behavioral Ecology, 8*(4), 372-377. http://doi.org/10.1093/beheco/8.4.372

Sakai, Y., Karino, K., Kuwamura, T., Nakashima, Y. & Maruo, Y. (2003a) Sexually dichromatic protogynous angelfish *Centropyge ferrugata* (Pomacanthidae) males can change back to females. *Zoological science, 20*(5), 627-634. http://doi.org/10.2108/zsj.20.627

Sakai, Y., Tsujimura, C., Nakata, Y., Tanabe, H. & Maejima, G. (2003) Rapid transition in sexual behaviors during protogynous sex change in the haremic angelfish *Centropyge vroliki* (Pomacanthidae). *Ichthyological Research*, *50*(1), 30-35.

Salekhova, L.P. (1961) Hermaphroditism of annular bream *Diplodus annularis* (L.). *Trudy Sevastopol Biological Station, 14*, 257-268.

Schwarz, A.L. & Smith, L.C. (1990) Sex change in the damselfish *Dascyllus reticulatus* (Richardson) (Perciformes: Pomacentridae). *Bulletin of marine science, 46*(3), 790-798.

Sellami, A. & Bruslé, J. (1979) Contribution a l'étude de la pêche, de la sexualité et de la reproduction de la mendole *Maena* (Linnaeus, 1758), Téléostéens, Maenidae, des cotes tunisiennes. *Mem Biol Mar Ocean IX***,** 91-109.

Shapiro, D.Y. & Rasotto, M.B. (1993) Sex differentiation and gonadal development in the diandric, protogynous wrasse, *Thalassoma bifasciatum* (Pisces, Labridae).

*Journal of Zoology, 230*(2), 231-245. http://doi.org/10.1111/j.1469-7998.1993.tb02685.x

Shapiro, D.Y. (1981) Size, maturation and the social control of sex reversal in the coral reef fish *Anthias squamipinnis* (Peters). . *Journal of Zoology London, 193*(1), 105–128. http://doi.org/10.1111/j.1469-7998.1981.tb01494.x

Shapiro, D.Y. (1992) Plasticity of gonadal development and protandry in fishes. *Journal of experimental Zoology, 261*(2), 194-203. http://doi.org/10.1002/jez.1402610210

Shapiro, D.Y., Garcia-Moliner, G. & Sadovy, Y. (1994) Social system of an inshore stock of the red hind grouper, *Epinephelus guttatus* (Pisces: Serranidae). In: E.K. Balon, M.N. Bruton & D.L.G. Noakes (Eds.), *Women in ichthyology: an anthology in honour of ET, Ro and Genie. Developments in environmental biology of fishes* (pp. 415-422). Springer, Dordrecht.

Shapiro, D.Y., Sadovy, Y. & McGehee, M.A. (1993a) Periodicity of sex change and reproduction in the red hind, *Epinephelus guttatus*, a protogynous grouper. *Bulletin of Marine Science, 53*(3), 1151-1162.

Shapiro, D.Y., Sadovy, Y. & McGehee, M.A. (1993b) Size, composition, and spatial structure of the annual spawning aggregation of the red hind, *Epinephelus guttatus* (Pisces: Serranidae). *Copeia, 1993*(2), 399-406. http://doi.org/10.2307/1447138

Sheaves, M. (1995) Large lutjanid and serranid fishes in tropical estuaries: are they adults or juveniles? *Marine Ecology Progress Series, 129*, 31-41. http://doi.org/10.3354/meps129031.

Shibuno, T., Chiba, I., Gushima, K., Kakudal, S. & Hashimotol, H. (1993a) Reproductive behavior of the wrasse, *Halichoeres marginatus*, at Kuchierabu-jima. *Japanese Journal of Ichthyology, 40*(3), 351-359.

Shibuno, T., Chiba, I., Hashimoto, H. & Gushima, K. (1994) Reproductive behavior of the wrasse, *Thalassoma lutescens*, at Kuchierabu-jima [in Japan]. *Journal of the Faculty of Applied Biological Science-Hiroshima University (Japan), 33,* 43-50.

Shibuno, T., Gushima, K. & Kakuda, S. (1993b) Female spawning migrations of the protogynous wrasse, *Halichoeres marginatus*. *Japanese Journal of Ichthyology, 39*(4), 357-362.

Shihab, I., Gopalakrishnan, A., Vineesh, N., Muktha, M., Akhilesh, K.V. & Vijayagopal, P. (2017) Histological profiling of gonads depicting protandrous hermaphroditism in *E leutheronema tetradactylum*. *Journal of Fish Biology, 90*(6), 2402-2411. http://doi.org/10.1111/jfb.13324

Shitamitsu, T. & Sunobe, T. (2018) Protandry of the flathead *Suggrundus meerdervoortii* (Teleostei: Platycephalidae). *Ichthyological Research, 65*(4), 507-509. http://doi.org/10.1007/s10228-018-0651-6

Shpigel, M. & Fishelson, L. (1986) Behavior and physiology of coexistence in two species of Dascyllus (Pomacentridae, Teleostei). *Environmental Biology of Fishes, 17*(4), 253-265. http://doi.org/10.1007/BF00001492

Shpigel, M. & Fishelson, L. (1991) Territoriality and associated behaviour in three species of the genus Cephalopholis (Pisces: Serranidae) in the Gulf of Aqaba, Red Sea. *Journal of Fish Biology, 38*(6), 887-896. http://doi.org/10.1111/j.1095-8649.1991.tb03628.x

Shpigel, M. (1985) *Aspects of the biology and ecology of the Red Sea groupers of the genus Cephalopholis (Serranidae, Teleostei).* PhD dissertation, Tel Aviv University.176 pp.

Siau, Y. & Bouain, A. (1994) Preliminary indications on growth and reproduction in the protogynous grouper *Mycteroperca rubra* (Pisces, Serranidae). *Journal of African*

*Zoology, 108*(4), 353-360.

Smith, C. L., & Atz, E.H. (1969). The sexual mechanism of the reef bass *Pseudogramma bermudensis* and its implications in the classification of the Pseudogrammidae (Pisces: Perciformes). *Zeitschrift für Morphologie der Tiere*, *65*(4), 315-326.

Smith, C.L. & Atz, E.H. (1973) Hermaphroditism in the mesopelagic fishes *Omosudis lowei* and *Alepisaurus ferox*. *Copeia*, *1973*(1), 41-44. http://doi.org/10.2307/1442355

Smith, C.L. (1959) Hermaphroditism in some serranid fishes from Bermuda. *Papers Michigan Academy of Science, Arts and Letters, 44*, 111-119.

Smith, C.L. (1964) Hermaphroditism in Bahama groupers. *Bulletin of the Museum of Natural History, New York, 73*, 42-47.

Smith, C.L. (1965) The patterns of sexuality and the classification of serranid fishes. *American Museum novitates, 2207*, 1-20.

Smith, C.L. (1975) The evolution of hermaphroditism in fishes. In: R. Reinboth (Eds.), *Intersexuality in the Animal Kingdom* (pp. 295-310). Springer, Berlin, Heidelberg. http://doi.org/10.1007/978-3-642-66069-6_28

Sordi, M. (1962). Ermafroditismo proteroginico in *Labrus turdis* L. e in *L. merula* L. *Monitore Zoologico Italiano* 69, 69-89.

St. Mary, C.M. (1996) Sex allocation in a simultaneous hermaphrodite, the zebra goby *Lythrypnus zebra*: insights gained through a comparison with its sympatric congener, *Lythrypnus dalli. Environmental Biology of Fishes, 45*(2), 177-190. http://doi.org/10.1007/BF00005232

Sulak, K.J., Wenner, C.A., Sedberry, G.R. & Guelpen, L.V. (1985) The life history and systematics of deep-sea lizard fishes, genus *Bathysaurus* (Synodontidae). *Canadian Journal of Zoology, 63*(3), 623-642. http://doi.org/10.1139/z85-091

Sumpton, W. & Brown, I. (2004) Reproductive biology of the red throat emperor *Lethrinus miniatus* (Pisces: Lethrinidae) from the southern Great Barrier Reef, Australia. . *Bulletin of marine science, 74*(2), 423-432.

Sunobe, T. & Nakazono, A. (1993) Sex change in both directions by alteration of social dominance in *Trimma okinawae* (Pisces: Gobiidae). *Ethology, 94*(4), 339-345. http://doi.org/10.1111/j.1439-0310.1993.tb00450.x

Sunobe, T., Sado, T., Hagiwara, K., Manabe, H., Suzuki, T., Kobayashi, Y., Sakurai, M., Dewa, S.I., Matsuoka, M., Shinomiya, A., Fukuda, K. & Miya, M. (2017) Evolution of bidirectional sex change and gonochorism in fishes of the gobiid genera Trimma, Priolepis, and Trimmatom. *The Science of Nature, 104*(3-4), 15. http://doi.org/10.1007/s00114-017-1434-z

Suzuki, K. (1979) Spawning behavior, k eggs, larvae, and sex reversal of two pomacanthine fishes, *Genicanthus lamarck* and *G. semifasciatus*, in the aquarium. *Journal of Faculty of Marine Science and Technology, Tokai University, 12*, 149-165.

Suzuki, K., Kobayashi, K., Hioki, S. & Sakamoto, T. (1978) Ecological studies of the anthiine fish, *Franzia squamipinnis*, in Suruga Bay, Japan. *Japanese Journal of Ichthyology, 25*(2), 124-140. http://doi.org/10.11369/jji1950.25.124

Suzuki, S., Toguchi, K., Makino, Y., Kuwamura, T., Nakashima, Y. & Karino, K. (2008) Group spawning results from the streaking of small males into a sneaking pair: male alternative reproductive tactics in the threespot wrasse *Halichoeres trimaculatus. Journal of ethology, 26*(3), 397-404. http://doi.org/10.1007/s10164-008-0102-3

Tanaka, H. (1990) Sexual maturation and sex reversal in red spotted grouper, *Epinephelus akaara. Bulletin of National Research Institute of Aquaculture, 17*, 1-15.

Tan-Fermin, J.D., Garcia, L.M.B. & Castillo, A.R. (1994) Induction of sex inversion in juvenile grouper, *Epinephelus suillus*, (Valenciennes) by injections of 17α-methyltestosterone. *Japanese Journal of Ichthyology, 40*(4), 413-420. http://doi.org/10.11369/jji1950.40.413

Taylor, R.G., Whittington, J.A., Grier, H.J. & Crabtree, R.E. (2000) Age, growth, maturation, and protandric sex reversal in common snook, *Centropomus undecimalis*, from the east and west coasts of South Florida. *Fishery Bulletin, 98*(3), 612-612.

Thompson, R. & Munro, J.L. (1974) The biology, ecology and bionomics of Caribbean reef fishes: Serranidae (hinds and groupers). *Research Reports Zoology Department University of the West Indies, 5*, 1-82.

Thompson, R. & Munro, J.L. (1978) Aspects of the biology and ecology of Caribbean reef fishes: Serranidae (hinds and groupers). *Journal of Fish Biology, 12*(2), 115-146. http://doi.org/10.1111/j.1095-8649.1978.tb04158.x

Tobin, A.J., Sheaves, M.J. & Molony, B.W. (1997) Evidence of protandrous hermaphroditism in the tropical sparid *Acanthopagrus berda. Journal of Fish Biology, 50*(1), 22-33. http://doi.org/10.1111/jfb.1997.50.1.22

Touart, L.W. & Bortone, S.A. (1980) The accessory reproductive structure in the simultaneous hermaphrodite *Diplectrum bivittatum. Journal of Fish Biology, 16*(4), 397-403. http://doi.org/10.1111/j.1095-8649.1980.tb03717.xTuset, V.M., García-Díaz, M.M., Gonzalez, J.A., Lorente, M.J. & Lozano, I.J. (2005) Reproduction and growth of the painted comber *Serranus scriba* (Serranidae) of the Marine Reserve of Lanzarote Island (Central-Eastern Atlantic). *Estuarine, Coastal and Shelf Science, 64*(2-3), 335-346. http://doi.org/10.1016/j.ecss.2005.02.026

Vadiya, V. (1984) Reproductive systems of *Epinephelus aeneus* and *Epinephelus alexandrinus* (Serranidae) from the southeastern Meditteranean. *Journal of Ichthyology, 24*, 77-81.

Van der Walt, B.A. & Mann, B.Q. (1998) Aspects of the reproductive biology of *Sarpa salpa* (Pisces: Sparidae) off the east coast of South Africa. *African Zoology, 33*(4), 241-248.Van Oördt, G.J. (1929) Zur mikroskopischen Anatomie der Ovariotestes von Serranus und Sargus (Teleostei). *Zeitschrift für mikroskopisch-anatomische Forschung, 19*, 1-17.

Van Oordt, G.J. (1933) Zur sexualität der gattung Epinephelus (Serranidae, Teleostei). *Zeitschrift für mikroskopisch-anatomische Forschung, 33*, 525-533.

Victor, B.C. (1987) The mating system of the Caribbean rosy razorfish, *Xyrichtys martinicensis. Bulletin of marine science, 40*(1), 152-160.

Von Brandt, V.I. (1979) *Soziale kontrole des wachstums und geschlechtsdifferenzierung bei jungen anemonenfischen (Amphiprion bicinctus)*. University of Munich, Munich, Germany.

Wakefield, C.B., Williams, A.J., Newman, S.J., Bunel, M., Boddington, D.K., Vourey, E. & Fairclough, D.V. (2015) Variations in growth, longevity and natural mortality for the protogynous hermaphroditic eightbar grouper *Hyporthodus octofasciatus* between the Indian and Pacific Oceans. *Fisheries Research, 172*, 26-33. http://dx.doi.org/10.1016/j.fishres.2015.06.0210165-7836

Warner, R.R. & Lejeune, P. (1985) Sex change limited by paternal care: a test using four Mediterranean labrid fishes, genus *Symphodus. Marine Biology, 87*(1), 89-99. http://doi.org/10.1007/BF00397010

Warner, R.R. & Robertson, D.R. (1978) Sexual Patterns in the Labroid Fishes of the Western Caribbean, I: The Wrasses (Labridae). *Smithsonian Contributions to*

*Zoology, 254*, 38 pp.

Warner, R.R. (1975) The reproductive biology of the protogynous hermaphrodite *Pimelometopon pulchrum* (Pisces: Labridae). *Fishery Bulletin, 73*(2), 262-283.

Warner, R.R. (1982) Mating systems, sex change and sexual demography in the rainbow wrasse, *Thalassoma lucasanum*. *Copeia, 1982*(3), 653-661. http://doi.org/10.2307/1444666

Wenner, C.A., Roumillat, W.A. & Waltz, C.W. (1986) Contributions to the life history of black sea bass, *Centropristis striata*, off the southeastern United States. *Fishery Bulletin, 84*.

Wernerus, F.M. & Tessari, V. (1991) The influence of population density on the mating system of *Thalassoma pavo*, a protogynous Mediterranean labrid fish. *Marine Ecology, 12*(4), 361-368. http://doi.org/10.1111/j.1439-0485.1991.tb00264.x

Whaylen, L., Pattengill-Semmens, C.V., Semmens, B.X., Bush, P.G. & Boardman, M.R. (2004) Observations of a Nassau grouper, *Epinephelus striatus*, spawning aggregation site in Little Cayman, Cayman Islands, including multi-species spawning information. *Environmental Biology of Fishes, 70*(3), 305-313. http://doi.org/10.1023/B:EBFI.0000033341.57920.a8

White, D.B., Wyanski, D.M., Eleby, B.M. & Lilyestrom, C.G. (2002) Tiger grouper (*Mycteroperca tigris*): profile of a spawning aggregation. *Bulletin of marine science, 70*(1), 233-240.Williams, A.J., Davies, C.R., Mapstone, B.D., Currey, L.M., Welch, D.J., Begg, G.A., Ballagh, A.C., Choat, J.H., Murchie, C.D. & Simpfendorfer, C.A. (2009) Age-based demography of humpback grouper *Cromileptes altivelis*: implications for fisheries management and conservation. *Endangered Species Research*, *9*(1), 67-79.

Wittenrich, M.L. & Munday, P.L. (2005). Bi-directional sex change in coral reef fishes from the family Pseudochromidae: an experimental evaluation. *Zoological Science*, *22*(7), 797-803.

Wong, M.Y., Munday, P.L. & Jones, G.P. (2005) Habitat patch size, facultative monogamy and sex change in a coral-dwelling fish, *Caracanthus unipinna*. *Environmental Biology of Fishes, 74*(2), 141-150. http://doi.org/10.1007/s10641-005-6715-2

Wyanski, D.M., White, D.B. & Barans, C.A. (2000) Growth, population age structure, and aspects of the reproductive biology of snowy grouper, *Epinephelus niveatus*, off North Carolina and South Carolina. *Fishery Bulletin, 98*.

Yeh, S.L., Dai, Q.C., Chu, Y.T., Kuo, C.M., Ting, Y.Y. & Chang, C.F. (2003) Induced sex change, spawning and larviculture of potato grouper, *Epinephelus tukula*. *Aquaculture, 228*(1-4), 371-381. http://doi.org/10.1016/S0044-8486(03)00316-8

Yeung, W.S.B. & Chan, S.T.H. (1987) The gonadal anatomy and sexual pattern of the protandrous sex-reversing fish, *Rhabdosargus sarba* (Teleostei: Sparidae). *Journal of Zoology, 212*(3), 521-532. http://doi.org/10.1111/j.1469-7998.1987.tb02922.x

Young, P.C. & Martin, R.B. (1982) Evidence for protogynous hermaphroditism in some lethrinid fishes. *Journal of Fish Biology, 21*(4), 475-484. http://doi.org/10.1111/j.1095-8649.1982.tb02853.x

Young, P.C. & Martin, R.B. (1985) Sex ratios and hermaphroditism in nemipterid fish from northern Australia. *Journal of Fish Biology, 26*(3), 273-287. http://doi.org/10.1111/j.1095-8649.1985.tb04266.x

Zamboni, A. & Relini, G. (1986) Note di biologia di *Spicara flexuosa* (Osteichthyes, Centracanthidae) del Mar Ligure. *Bollettino dei Musei e Degli Istituti Biologici dell'Università di Genova, 52*, 251-265.

Zei, M. (1950) Typical sex-reversal in Teleosts. *Proceedings of the Zoological Society of London*, *119*(4), 917-920. http://doi.org/10.1111/j.1096-3642.1950.tb00917.x

Zohar, Y., Abraham, M. & Gordin, H. (1978) The gonadal cycle of the captivity-reared hermaphroditic teleost *Sparus aurata* (L.) during the first two years of life. *Annales de Biologie Animale Biochimie Biophysique*, *18*(4), 877-882.

Zorica, B., Sinovčić, G. & Čikeš Keč, V. (2005) Reproductive period and histological analysis of the painted comber, *Serranus scriba* (Linnaeus, 1758), in the Trogir Bay area (eastern mid-Adriatic). *Acta Adriatica: International journal of Marine Sciences, 46*(1), 77-82.