Switches, stability and reversals:
the evolutionary history of sexual systems in fish

Susanna Pla[1#], Chiara Benvenuto[2#], Isabella Capellini[3*] & Francesc Piferrer[1*]

[1]Institut de Ciències del Mar, Spanish National Research Council (CSIC), Barcelona, Spain

[2]School of Science, Engineering and Environment, University of Salford, Salford, UK

[3]School of Biological Sciences, Queen's University Belfast, UK

[#]Joint first authors

*Correspondence: Dr. Francesc Piferrer, Institut de Ciències del Mar, Spanish National Research Council (CSIC), Passeig Marítim, 37-49, 08003 Barcelona. Tel. +34-932 309 567; E-mail: piferrer@icm.csic.es

*Correspondence: Dr. Isabella Capellini, School of Biological Sciences, Queen's University Belfast, 19 Chlorine Gardens, Belfast, BT9 5DL, Northern Ireland, United Kingdom; E-mail: I.Capellini@qub.ac.uk

Abstract

Sexual systems are highly diverse and have profound consequences for population dynamics and resilience. Yet, little is known about how they evolved. Using phylogenetic Bayesian modelling on 4740 species, we show that gonochorism is the likely ancestral condition in teleost fish. While all hermaphroditic forms revert quickly to gonochorism, protogyny and simultaneous hermaphroditism are evolutionarily more stable than protandry. Importantly, simultaneous hermaphroditism can evolve directly from gonochorism, in contrast to theoretical expectations. We find support for predictions from life history theory that protogynous species live longer than gonochoristic species, are smaller than protandrous species, have males maturing later than protandrous males, and invest the least in male gonad mass. The large-scale distribution of sexual systems on the tree of life does not seem to reflect just adaptive predictions and thus does not fully explain why some sexual forms evolve in some taxa but not others (William's paradox). We propose that future studies should take into account the diversity of sex determining mechanisms. Some of these might constrain the evolution of hermaphroditism, while the non-duality of the embryological origin of teleost gonads might explain why protogyny predominates over protandry in this extraordinarily diverse group of animals.

Keywords

Hermaphroditism, phylogeny, Williams' paradox, Dollo's law, developmental plasticity, sex determination

Introducction

Sexual reproduction is a unifying feature of Eukaryotes[1] and yet it is extremely diverse[2]. Sexual systems, defined as the distribution of the male and female function among the individuals of a given species, vary from separate fixed sexes (known as gonochorism in animals and dioecy in plants) to simultaneous hermaphroditism (each individual produces both male and female gametes at the same time). These two sexual systems can be viewed as the extremes in a sexually plastic gradient[3] of intermediate (sequential hermaphroditism) and mixed (coexistence of males and/or females with hermaphrodites)[4,5] systems. Sexual systems have a profound influence on individuals' mating success and fitness[6], population sex ratios and effective sizes[7], and colonization events and habitat use[8]. As a result, sexual system influence the population dynamics and resilience to natural and anthropogenic stressors of ecologically and commercially important species that are often endangered or overexploited[9].

Hermaphroditism is predominant in flowering plants, i.e., angiosperms[10], where 94% of the species have male and female sex organs in the same individual/flower, and it is widespread in invertebrates and fish, totalling 5% of animal species or up to ~30% if insects are excluded[11]. While this diversity suggests multiple evolutionary transitions between sexual systems in response to selection, current evolutionary models on the adaptive advantage of different sexual systems explain little about how and why sexual systems evolve and thus their large-scale distribution across the tree of life (Williams' paradox[4,12]). Thus, unravelling the evolutionary history of sexual systems and quantifying how frequently and in what direction transitions occur, is key to revealing which sexual systems are evolutionarily labile or stable, elucidating how one changes into another over evolutionary time, and identifying the environmental, genetic and developmental drivers favouring or opposing these changes. Yet, our understanding of how sexual systems evolve is still limited, particularly in animals.

Theoretical models, initially developed for plants, suggest that simultaneous hermaphroditism and dioecy are evolutionary stable conditions that are retained over long evolutionary time and unlikely lost once evolved, while mixed sexual systems represent evolutionary intermediate stages[4,5,13] (Fig. 1).

Fig. 1. Theoretical framework for the evolution of sexual systems: potential evolutionary transitions between gonochorism and simultaneous hermaphroditism via mixed systems (mixed pathways) as described in plants and some animals; via sequential hermaphroditism (sequential pathways) as recently suggested[4]; or without intermediate states (direct pathways) as proposed for plants[6]. Double-headed arrows indicate theoretical pathways. Large arrows identify the directionality more frequently investigated for plants (green) and animals (blue), from the most common to the rarest state.

Simultaneous hermaphroditism is likely the ancestral state in angiosperms from which dioecy, a rare sexual system in plants[6], has evolved independently several times, possibly to avoid inbreeding[14,15]. Theoretical models predict that separate sexes in plants evolve from hermaphroditism in different ways: 1) primarily through the intermediate state of gynodioecy[16], a sexual system common in plants that occurs when a male-sterile mutant invades an hermaphroditic population resulting in the coexistence of hermaphrodites and females; 2) androdioecy, where the mutation results in female sterility, resulting in the coexistence of hermaphrodites and males, which is less common in plants[17,18]; 3) possibly via trioecy, i.e., the coexistence of hermaphrodites, males and females, which is very rare; and 4) less frequently, via a direct transition[6,10] (Fig. 1). However, in animals no evidence of a direct transition exists.

Once gained, dioecy was believed to be an irreversible condition[19], a conclusion based on the

assumption that returning to a simultaneous expression of male- and female-specific genes

would likely produce contrasting effects on sex-specific physiology. Recent studies, however,

reject this claim in plants, as phylogenetic reconstructions of direct transitions from

dioecy/gonochorism to simultaneous hermaphroditism have been documented[10,20].

The same theoretical framework with mixed pathways has been proposed also for

animals where, in contrast to plants, gonochorism is the most common sexual system,

androdioecy is more common than gynodioecy[5,13] and trioecy is very rare[21]. However, several

reproductive characteristics in plants are quite different from the ones evolved in animals[22],

albeit similarities can be found in some invertebrates[23]; hence, different theoretical frameworks

are required (Fig. 1). Furthermore, evolutionary transitions between sexual systems in fish, the

only vertebrates to exhibit hermaphroditism[24], might be less likely to occur preferentially via a

mixed pathway (Fig. 1) given that in fish (>30000 species) only a few killifish species in the

genus *Krytpolebias* (formerly *Rivulus*) are truly androdioecious[5,25,26] while the presence of

gynodioecy and trioecy is still debated in the hagfish *Myxine glutinosa*[13,27]. Recently,

sequential hermaphroditism has been suggested as a possible intermediary state that may

facilitate evolutionary changes between gonochorism and simultaneous hermaphroditism[4]

(Fig. 1). However, we currently have no robust study on the evolution of sexual systems in

animals and thus we currently do not know whether sequential hermaphroditism represents an

evolutionary intermediate stage between gonochorism and simultaneous hermaphroditism,

whether protogyny and protandry act equally as transitional forms between the two, and

whether gonochorism and simultaneous hermaphroditism are evolutionary stable conditions in

animals as they are in plants.

The evolution of hermaphroditism in animals has mostly been interpreted in the context

of its adaptive advantages relative to gonochorism, as proposed by the low density and the size

advantage models[28]. The former predicts that simultaneous hermaphroditism evolves under

low population densities and/or low dispersal capacity as, in these conditions, individuals with

this sexual system can maximize their chances of securing a mate than sex-changing or

gonochoristic individuals[29]. Of note, the possible advantages of self-fertilizing simultaneous

hermaphrodites would be offset by increased risk of inbreeding. The size advantage model

proposes adaptive explanations for the evolution of sequential hermaphroditism based on the

relationship between size and fecundity[28,30–32]. Specifically, since sequential hermaphrodites

have indeterminate growth[33], sex change should be favoured when the reproductive value of

an individual depends on size (and thus age), and this affects in particular one of the sexes.

Thus, individuals change from a smaller first sex to a larger second sex and the direction of sex
change depends on the sex that maximizes its reproductive value with a larger size[34]. The
interdependence between size, fecundity and fitness is also affected by the mating system of
the species, defined as the pattern of sexual interactions that take place considering the number
of males and females involved in reproduction (Table 1). Therefore, protandry (male-to-female
sex change) is expected in species that reproduce in monogamous or random pairs and where
individuals switch from small males to large, highly fecund females, achieving higher
reproductive potential. Conversely, protogyny (female-to-male sex change) is expected in
polygynous/group-mating species, where small females become large dominant males that
monopolize females, often grouped in harems (Table 1). In both systems exceptions exist with
a few individuals born directly as second sex (primary females in digynic protandrous species[35]
and primary males in diandric protogynous species[36]).

Crucially, life history traits underpin the size advantage model and life history theory
is central to the study of sexual systems evolution since it allows to derive clear predictions
about how and when individuals should allocate energy among different life history traits,
including sexual functions, to optimize fitness[37]. However, life history traits are surprisingly
not explicitly incorporated in the theory, nor tested in empirical studies[12]. Longevity, maximum
size and age/size at maturity are key life history traits because they determine individual fitness,
influence demographic parameters of populations[38] and impact on the population's genetic
diversity[39]. Since sequential hermaphrodites achieve higher fitness when reproducing as the
second sex[34], they should benefit more from increased longevity or larger size than
gonochoristic and simultaneous hermaphroditic species. Alternatively, sequential
hermaphrodites could mature earlier as the first sex compared to the same sex in gonochoristic
species and capitalize on reproduction as second sex. These predictions, however, remain to be
tested.

The size advantage model has been supported in crustaceans[40], molluscs[41] and fish[42].
Spawning behaviour, i.e., how the two sexes interact to release the gametes, can be classified
in fish as pair spawning, involving only two individuals at the time, and spawning aggregations,
comprising large breeding groups[43] (Table 1). Mating system and spawning behaviour together
determine the intensity of direct male-male competition and sperm competition (i.e., the
competition between the sperm of two or more males for fertilization of the same eggs), and
thus certainty of paternity. Sperm competition is a key selective force shaping male
reproductive anatomy, physiology and behaviour across diverse animal groups[44,45] and is
assumed to be low in haremic systems[42,46,47] where large dominant males can better monopolize

groups of females (pair spawning) with no or limited competition by other males[48]. Likewise,

low sperm competition is expected under monogamy. Group spawning is found in promiscuous

mating or aggregations, leading to intense sperm competition[49] as many males try to fertilize

the eggs of multiple females. The intensity of sperm competition has been incorporated in the

size advantage model[50] as it can play a significant role in the advantage of protogyny: changing

sex from female to male should be more advantageous when paternity assurance is high due to

reduced sperm competition[51]. Consistent with these predictions, the gonadosomatic index

(GSI), defined as the percentage of body mass devoted to the gonads[52] and a reliable indicator

of the intensity of sperm competition[53], is significantly lower in protogynous teleost species

than in gonochoristic congeners[47,48,54]. However, protandrous teleost fish do not always

conform to theoretical expectations, exhibiting higher GSI as males than expected[48]. We have

recently proposed that, at least in the family Sparidae, high male GSI in protandrous fish can

be explained not only by spawning aggregations and high sperm competition in some species,

but also because high investment in the gonads represents a compensatory mechanism that

allows small males to fertilize highly fecund females much larger than themselves[54].

Table 1. Predictions of associations between sexual systems, mating systems, sex ratio, size of mates and spawning behaviour in teleosts.

Sexual System	Mating system	Sex ratio	Size of mates	Spawning behaviour
GONOCHORISM [G]	♂ ↔ ♀		♂ = ♀	
Individuals reproduce as one sex throughout their lifetime (male or female)	♂♂♂ ↔ ♀♀♀	1:1	♂ > ♀	♂♀
	♂ ↔ ♀♀♀		♀ > ♂	♂♂♂♀♀♀
PROTOGYNY [PG]				
Female-first sequential hermaphroditism: individuals first reproduce as females, change sex once with increasing size/age and then reproduce as males	♂ ↔ ♀♀♀ ♂♂♂ ↔ ♀♀♀	♀ Biased	♂ > ♀	♂♀ ♂♂♂♀♀♀
PROTANDRY [PA]				
Male-first sequential hermaphroditism: individuals first reproduce as males, change sex once with increasing size/age and then reproduce as females	♂ ↔ ♀	♂ Biased	♀ > ♂	♂♀ ♂♂♂♀♀♀
BIDIRECTIONAL [BD]				
Individuals can change sex more than once, in either directions throughout their lifespan usually from PG	♂ ↔ ♀ ♂ ↔ ♀♀♀	♀ Biased	♂ > ♀	♂♀
SIMULTANEOUS [SH]				
Individuals produce gametes of both sexes at the same time or in a short period of time	♂ ↔ ♀	1:1	♂ = ♀	♂♀

Mating system: ♂ ↔ ♀ = monogamy (pair bond) or random pairing; ♂♂♂ ↔ ♀♀♀ = promiscuity; ♂ ↔ ♀♀♀ = harem polygyny or temporary lek-like systems (many females in a territory defended by a male). Spawning behaviour: ♂♀ = pair spawning (pair of individuals); ♂♂♂♀♀♀ = group spawning/spawning aggregations.

Fish account for more than 50% of the extant species of vertebrates and are characterized not only by their extraordinary diversity in morphology, physiology, ecology and habitat, but also by different sexual systems, including gonochorism, different forms of hermaphroditism —the only group among vertebrates— and unisexuality (all-female populations)[24,55,56]. Hermaphroditism in fish is broadly divided into simultaneous (synchronous) and sequential (consecutive) hermaphroditism, the latter in the form of protandry, protogyny and bidirectional sex change (Table. 1). This remarkable diversity in sexual systems in fish, particularly in the infraclass teleosts accounting for 96% of species, makes them an ideal group in which to study the evolution of different forms of

hermaphroditism. Here, we investigate the evolutionary origin and transitions among sexual

systems across 4740 teleost species belonging to 294 families using a recent time calibrated

phylogeny and modern phylogenetic comparative approaches. Our large-scale approach allows

us to fully unravel how sexual patterns evolved and identify which ones represent evolutionary

stable conditions. We demonstrate that gonochorism is the likely ancestral condition in teleosts

and it is an evolutionarily stable state from which not only protogyny and protandry, but also

simultaneous hermaphroditism evolve at a moderate evolutionary rate. In contrast to theoretical

predictions, we show for the first time that simultaneous hermaphroditism can evolve directly

from (and be directly lost back to) gonochorism, without the need of intermediary evolutionary

steps. Further, we expand the theoretical framework of the evolution of hermaphroditism

investigating how life history traits and male GSI differ between sexual systems, as predicted

by life history theory. In support of predictions, we found longer lifespan in protogynous

species compared to gonochoristic, larger size in protandrous species compared to

protogynous, and smaller GSI in protogynous males. However, contrary to predictions, we

found no differences in age or size at maturity across sexual systems, with the exception of late

maturing protogynous males. Finally, we discuss how our results should be incorporated in an

even broader framework that includes sex determining mechanisms and gonadal plasticity as

possible constraint and facilitation mechanisms, respectively, to gather a better understanding

of the evolution of sexual systems and possibly resolve Williams' paradox.

Results

Evolutionary history of sexual systems

Our dataset includes 4740 extant teleost species, of which 403 hermaphroditic (protogynous: n

= 285; protandrous: n = 57; bidirectional sex change: n = 14; simultaneous hermaphroditism:

n = 47; see Methods, Data collection). We used Discrete models of evolution to reconstruct the

evolutionary history of sexual systems using Reversible Jump (RJ) Markov chain Monte Carlo

(MCMC) in *BayesTraits* (Methods, Phylogenetic comparative analysis). Treating sexual

systems as a two character states (gonochoristic or hermaphroditic) our analysis revealed that

gonochorism is the most likely ancestral character state in teleosts (Fig. 2; Supplementary

Table 1) and that hermaphroditism slowly evolves from, and reverts very quickly and multiple

times back to, gonochorism (Fig. 2). This indicates that gonochorism is an evolutionarily stable

state in teleosts (Supplementary Fig. 1).

Fig. 2. Teleost phylogeny with the evolutionary history of sexual systems in maximum likelihood using the R package Ape v4. In the inner circle and in the internal nodes of the phylogeny, the sexual systems of extant species and their ancestors are colour coded for gonochorism (grey) and hermaphroditism (light blue). In the external circle, the sexual systems of extant species are colour coded for gonochorism (grey), protogyny (red), protandry (blue), bidirectional hermaphroditism (green) and simultaneous hermaphroditism (yellow). The density plots in the inset show the estimated probability of character state at the root of the phylogeny (RJ-MCMC Multistate) and colour coded for gonochorism (grey) and hermaphroditism (light blue).

Treating sexual system as four character states (gonochoristic, protandric, protogynic, simultaneous hermaphroditic) our analysis revealed that all three types of hermaphroditism evolve at a very low rate from gonochorism (Fig. 3, Table 2). Protogyny and protandry revert very rapidly back to gonochorism. Furthermore, protogyny is as likely to evolve from gonochorism as it is to switch to simultaneous hermaphroditism or protandry, although nearly 20% of the models estimate the latter transition rate to be equal to zero. Conversely, protandry is lost as quickly to protogyny as it is to gonochorism and, to a much smaller extent, to simultaneous hermaphroditism given that 38% of the models estimated this latter transition rate to be zero. Finally, simultaneous hermaphroditism evolves directly from and is lost back to

gonochorism at similar low rates, with 15% of the models estimating these transition rates to be equal to zero. Therefore, simultaneous hermaphroditism can evolve at similar rates directly from gonochorism or from protogyny, but less likely from protandry. Conversely, while simultaneous hermaphroditism is also lost to protogyny and protandry at similar low rates as to gonochorism, transitions from simultaneous hermaphroditism to sequential hermaphroditism are much less likely given that over 45% of the models estimated them to be equal to zero. Altogether, these results clearly indicate that in teleosts gonochorism is an evolutionarily stable state; protogyny is evolutionarily more stable than protandry, while simultaneous hermaphroditism evolves rarely but is evolutionarily stable, being lost slowly to gonochorism, and possibly to protogyny and protandry (Fig. 3, Table 2).

Fig. 3. Summary of RJ-MCMC Multistate analysis with density plots of the posterior distributions of the transition rates to gonochorism (G, in grey), protogyny (PG, in red), protandry (PA in blue), and simultaneous hermaphroditism (SH, in yellow). Note, only x-axis, but not y-axis, are the same for each pair of gain and loss between two character states. The thickness of the arrows is roughly proportional to the mean magnitude of the transition rates from the posterior distribution. Dotted lines indicate transition rates estimated to be equal to 0 in over 30% of the models in the posterior distributions. Sample sizes of extant species for each sexual system category are indicated between parentheses.

Table 2. Results of the RJ-MCMC Multistate analysis in BayesTraits of sexual systems as a
four state categorical variable: gonochorism (G), protogyny (PG), protandry (PA) or
simultaneous hermaphroditism (SH). For each posterior distribution, we report the effective
sample size (ESS), the mean and 95% high density intervals (95-HPD), the mode and the
percentage of models in which the parameter is estimated as zero. This analysis is based on
4726 extant species (G: n = 4337; PG: n = 285; PA: n = 57; SH: n = 47). Note: 14 species of
bidirectional sex change were not included due to their low number.
285

Transition rates	ESS	Mean	95-HPD	Mode	% Zero
G → PG	2000	0.0172	0.0097 – 0.0291	0.0156	0.2
PG → G	2000	1.1370	0.8381 – 1.4181	1.1096	0
G → PA	2000	0.0172	0.0097 – 0.0291	0.0156	0
PA → G	2000	1.1770	0.8427 – 1.4866	1.1096	0
G → SH	2000	0.0132	0 – 0.0201	0.0156	15
SH → G	2000	0.0147	0 – 0.0275	0.0162	15
PG → PA	2000	0.0142	0 – 0.0277	0.0161	19
PA → PG	2000	1.1160	0 – 1.4893	1.1096	0.3
PG → SH	2000	0.0170	0.0089 – 0.0309	0.0156	2
SH → PG	2000	0.0102	0 – 0.0252	0	45
PA → SH	2000	0.0611	0 – 0.0589	0	38
SH → PA	2000	0.0090	0 – 0.0247	0	48
Root probabilities					
G	2000	40.7	35.7 – 48.2	39.2	0
PG	2000	29.9	25.5 – 32.8	30.3	0
PA	2000	29.3	23.7 – 37.0	30.4	0
SH	2000	<0.1	0 – 0.9	<0.01	2.3

Life history traits and sexual systems in teleosts

The phylogenetic generalized least square (PGLS) analysis revealed that protogynous, but not

protandrous, species live longer than gonochoristic species (Fig. 4a; Table 3). Protandrous

species reach larger maximum adult size than protogynous species (Fig. 4b; Table 3). Female

and male age at maturity do not differ across species with different sexual systems (Fig. 4c-d;

Table 3) nor does sex-specific length at maturity (Table 3). However, when accounting for

allometry, males in protogynous systems mature later than males in protandrous systems (Fig.

4e). Finally, protogynous males have lower GSI values than gonochoristic and protandrous

ones, but GSI does not differ significantly between gonochoristic and protandric males (Fig.

4f; Table 3).

Fig. 4. Selected life history traits by sexual system. Phylogenetic estimated mean and standard error from the PGLS results of: **a)** longevity (G: n = 759; PG: n = 88; PA: n = 19); **b)** maximum length (G: n = 2624; PG: n = 243; PA: n = 36); **c)** male age at first maturity (G: n = 258; PG: n = 20; PA: n = 12); **d)** female age at first maturity (G: n = 281; PG: n = 33; PA: n = 5); **e)** male age at first maturity corrected for male length at first maturity (G: n = 136; PG: n = 12; PA: n = 9); **f)** male gonadosomatic index, GSI (G: n = 45; PG: n = 45; PA: n = 15). In all panels gonochorism (G) in grey, protogyny (PG) in red and protandry (PA) in blue.

Table 3. Results of phylogenetic generalized least square (PGLS) model of longevity (year; $\log_{10}$ transformed), length at first maturity (cm; $\log_{10}$ transformed) across sexual systems: gonochorism (G); protogyny (PG); protandry (PA). For each independent variable we report the parameter estimate (Beta), t-statistics (T), *P*-value (P), and the model statistics including the degrees of freedom (df), the maximum likelihood estimation of the phylogenetic signal (λ) and R^2. Refer to supplementary table 2 for sexual system and sex-specific data. Significant differences are indicated in bold.

Variable Dependent	Variable Independent	Beta	T	P	Df	Model statistics λ	R^2
Longevity	Sexual system - PA[1]	0.035	0.345	0.730	2; 863	0.909	0.005
	Sexual system - PG[1]	0.127	2.079	**0.038**			
	Sexual system - PG[2]	0.092	0.878	0.380			
Max length	Sexual system - PA[1]	0.094	1.615	0.106	2; 2900	0.973	0.001
	Sexual system - PG[1]	-0.037	-1.111	0.267			
	Sexual system - PG[2]	-0.131	-2.025	**0.043**			
Age at maturity ♂	Sexual system - PA[1]	-0.117	-1.183	0.237	2; 287	0.865	0.009
	Sexual system - PG[1]	0.067	0.763	0.446			
	Sexual system - PG[2]	0.184	1.595	0.112			
Age at maturity ♀	Sexual system - PA[1]	0.172	1.288	0.198	2; 316	0.861	0.005
	Sexual system - PG[1]	0.006	0.070	0.944			
	Sexual system - PG[2]	-0.166	0.138	0.231			
Length at maturity ♂	Sexual system - PA[1]	-0.031	-0.468	0.640	2; 368	0.973	0.001
	Sexual system - PG[1]	-0.033	-0.601	0.548			
	Sexual system - PG[2]	-0.003	-0.034	0.973			
Length at maturity ♀	Sexual system - PA[1]	0.030	0.344	0.731	2; 348	0.970	0.0003
	Sexual system - PG[1]	0.013	0.199	0.842			
	Sexual system - PG[2]	-0.017	-0.174	0.862			
Age at maturity ♂	Length at maturity ♂	0.581	8.186	**<0.001**	3; 153	0.782	0.318
	Sexual system - PA[1]	-0.145	-1.680	0.095			
	Sexual system - PG[1]	0.091	1.184	0.238			
	Sexual system - PG[2]	0.236	2.427	**0.016**			
Age at maturity ♀	Length at maturity ♀	0.535	8.203	**<0.001**	3; 168	0.831	0.287
	Sexual system - PA[1]	0.032	0.243	0.808			
	Sexual system - PG[1]	0.008	0.112	0.911			
	Sexual system - PG[2]	-0.023	-0.181	0.856			
GSI ♂	Sexual system - PA[1]	0.072	0.553	0.581	2; 102	0.904	0.143
	Sexual system - PG[1]	-0.367	-3.722	**<0.001**			
	Sexual system - PG[2]	-0.433	-3.214	**0.002**			

[1] G as reference level; [2] PA as reference level

Discussion

365  Our large-scale phylogenetic study has, for the first time, tested the theoretical

366 predictions for how sexual systems have evolved and has revealed the evolutionary origin of

367 and transitions between different sexual systems in the highly diverse teleosts. We thus identify

368 gonochorism, and to some extent simultaneous hermaphroditism, as stable conditions over

369 evolutionary time, being lost at lower rates than they are gained. Protandry is evolutionarily

370 the most labile condition. In contrast to theoretical predictions, our study clearly demonstrates

371 that, while simultaneous hermaphroditism can evolve from protogyny and possibly from

372 protandry, it is as likely to evolve directly from —and revert directly back to— gonochorism.

373 We find partial support for the predictions derived from life history theory that sequential

374 hermaphrodites live longer and grow bigger than teleosts with other sexual systems.

375 Specifically, protogynous species live longer than gonochoristic species and protandrous

376 species attain a larger size than protogynous ones. Finally, protogynous males invest the least

377 in male gonad tissues (quantified by the gonadosomatic index) relative to males with other

378 sexual systems and mature later for their size than protandrous males. However, female age at

379 maturity does not differ between sexual systems. Combined, these results suggest that the two

380 forms of sequential hermaphroditism must be treated separately in theoretical and empirical

381 studies as protandry and protogyny are characterized by distinct life history strategies[34], even

382 though they both entail sex change.

383  Using the largest dataset ever collected with four sexual systems in teleost fish, our

384 study reveals a complex and dynamic way through which sexual systems evolve and switch

385 between one another. All forms of hermaphroditism can evolve directly from gonochorism (the

386 ancestral state in teleosts) and revert back to gonochorism even faster. Therefore, even though

387 gonochorism is an evolutionarily stable condition, gained faster than it is lost, our results refute

388 the assumption that the transition to gonochorism is irreversible[19] and represent another

389 example[57,58] against Dollo's law of irreversibility[59]. Conversely, in teleosts, all forms of

390 hermaphroditism are less evolutionarily stable than gonochorism. Importantly, our analysis

391 reveals that simultaneous hermaphroditism can originate directly from gonochorism or through

392 sequential hermaphroditism, but simultaneous hermaphroditism is lost preferentially to

393 gonochorism rather than either form of sequential hermaphroditism. Our results therefore

394 refute the conclusion of Pennell et al[60]. Specifically, using a much smaller dataset biased

395 towards a greater proportion of hermaphrodite than gonochoristic species than what is observed

396 in teleosts, and without discriminating between different types of hermaphroditism, Pennell et

397 al[60] found that the evolutionary transition from gonochorism to hermaphroditism is over twice

as fast than the reverse, suggesting rapid evolution of hermaphroditism from gonochorism, a

conclusion that these authors acknowledge is counterintuitive. Importantly, our analyses

demonstrate that no intermediate stages are strictly required to either gain or lose simultaneous

hermaphroditism from gonochorism. These results call into question theoretical predictions

(Fig. 1) that sex-specific gene expression in gonochoristic species prevents direct evolutionary

transitions between gonochorism and simultaneous hermaphroditism, and intermediate stages,

like sequential hermaphroditism, are required[4]. Androdioecy is considered an intermediate

stage from simultaneous hermaphroditism to gonochorism in plants and from gonochorism to

hermaphroditism in some invertebrates[5,13], but this sexual system is extremely rare in fish and

cannot explain the evolution of diverse sexual system in this vertebrate group. Overall, and

contrary to what had previously been surmised, our analyses suggest that the complexity of

sex-specific physiology and behaviour do not constrain transitions between sexual systems,

and that even changes to simultaneous expression of both sexual functions can be

evolutionarily gained and lost easily from gonochorism.

According to life history theory, sequential hermaphrodites should live longer, grow

bigger and mature earlier as the first sex than gonochoristic species. We find that protandrous

and protogynous species differ in their life history strategies: protandrous species attain a

bigger size than protogynous species while protogynous species live longer than gonochoristic

species. These differences reflect the fact that protandry and protogyny maximise their fitness

as the second sex[34] which differs between the two systems. Therefore, protandrous species

benefit primarily by achieving a larger size, as larger females (second sex) are more fecund

than smaller ones. Conversely, the longer life in protogynous species maximises the fitness of

the large successful males (second sex) that can monopolize females in harems or in spawning

grounds. In addition, male investment in gonad tissue (as quantified by the gonadosomatic

index) is lower in protogyny, as expected by theory[48,54], since large males can better

monopolize mating opportunities and face low levels of sperm competition in harems and

spawning aggregations (Table 2). Small-sized protandrous males in spawning aggregations

instead need to boost their investment in the gonads but even in the absence of sperm

competition (monogamy) they require large gonads to fertilize highly fecund females larger

than themselves[54]. Thus, sexual systems and mating strategies affect life history traits

differentially in protandrous and protogynous species. It is well known that in sequential

hermaphrodites the second sex always matures later and is larger than the first sex, so it is not

surprising that in protandrous species females are significantly larger than males when reaching

maturity, while in protogynous species males are significantly larger than females[51] (excluding

the cases primary females and primary males, respectively). Yet, no comparison has been made

for size/age at first maturity for males and females across sexual systems. Life history theory

predicts that the first sex of sequential hermaphrodites matures earlier than the same sex in

gonochorism, but we find no evidence for this. However, future studies should re-evaluate

these relationships as more data become available for a large number of sex-changing species,

as currently sex-specific life history data are scarce for hermaphroditic fish species. Data are

particularly necessary to fully assess theoretical predictions for size and time spent as females

and as males in sequential hermaphrodites. Finally, when accounting for allometry, we find

that protogynous species mature as males (second sex) at a later age than males of protandrous

species. Such difference for males but not females in sequential hermaphrodites might reflect

social regulation associated with sex change in protogyny. Specifically, protogynous females

typically change sex when a dominant male is not present[36]; thus, social regulation might delay

sex change in females until later in life.

445   While we have shown that life history theory can provide a major contribution to our

understanding of sexual system evolution, we propose that the highly dynamic picture revealed

by this study should be expanded using a more comprehensive approach that includes not only

selection and adaptation, but also genetic and developmental mechanisms (Fig. 5). We propose

that more data should be gathered on sex determining mechanisms to gain a better

understanding of the evolution of sexual systems. Fixed, strongly canalized, genetic sex

determination (culminating in the formation of fully differentiated and stable heteromorphic

sex chromosomes) might constrain the evolution of hermaphroditism (acting as an evolutionary

trap[2,61–63], but see Pennell et al[60]). Similarly, the peculiarities of the embryonic origin of the

urogenital system in teleosts, compared to other vertebrates[64], may favour a certain sexual

system (protogyny) over another (protandry) in this group. Sex determination in gonochoristic

animals is determined either at fertilization by different genetic mechanisms, including male

(XX/XY) or female (ZW/ZZ) heterogamety with homomorphic or heteromorphic sex

chromosomes, or after conception by environmental factors, or by a combination of both[65,66].

Fish are characterized by an incredible diversity[67–69] and evolutionary plasticity[60,70] of sex

determining mechanisms. Many fish do not have sex chromosomes[68] and when they are

present, they might not always be clearly differentiated, with the sex determining loci often not

easily identifiable[71]. In some cases, sex determination can be due to a difference in a single

nucleotide[68]. Moreover, high turnover of sex chromosomes has been detected in some fish

lineages (e.g., sticklebacks[72,73]), including reversal to autosomes[60]. Although data are currently

scarce for formal analyses, sequential hermaphrodites do not appear to have sexually

differentiated chromosomes[74], but additional research on the underlying genetic basis of sex determination specifically linked to hermaphroditism in teleosts is clearly needed. Therefore, although complete genetic control of hermaphroditism is common in plants[75], a better knowledge of the genetic mechanisms of sex determination could be helpful to explain how hermaphroditism in teleosts has evolved in some taxonomic groups but not in others under similar ecological pressures. Previous attempts to connect the distribution of sexual systems and sex determining mechanisms have invoked morphological[76] and developmental[64] aspects. Furthermore, sequential hermaphroditism can be regarded as a clear example of phenotypic plasticity, and since epigenetics underlies phenotypic plasticity, common epigenetic mechanisms have been proposed to participate in the evolutionary transitions between different sexual systems and sex-determining mechanisms[77].

Fig. 5. Theoretical framework for the evolution of sexual systems: overview of parameters (with some examples) considered in the low density and the size advantage models (*), used in our analyses (**) and proposed in the present study (***).

Overall, sex determination is achieved through plastic switches that channel an
individual's development towards one sex or the other, or both at different or at the same time,
usually with the concurrence of biotic or abiotic influences[78]. Further developmental plasticity
is uniquely documented in fish via the bipotential nature their gonads and gonoducts[64]. Thus,
while in most vertebrate taxa gonads develop from two distinct germinal layers (medulla,
endodermal in origin, which gives rise to the testes; and cortex, mesodermal in origin, which
gives rises to the ovary), in teleosts the gonads consist entirely of the cortex homolog[79].
Therefore, anatomically all fish could be, in principle, hermaphrodites[64]. The transition to
protogyny may be favoured by the peculiarities of gonadal development in many fish species,
which develop a female gonad, complete with ovaries containing cysts of oocytes, during the
initial stage of gonadal formation[80,81]. Only later testicular development is triggered and
superimposed on this arrangement so that the individual ultimately matures functionally as a
male[82–84]. Thus, protogyny might be favoured because female gonads are often the first to
develop. Bidirectional sex change, a rarer system in fish (Table 1), further demonstrates the
importance of gonadal plasticity. In most cases, the initial strategy is protogyny[85], but males
can revert back to females when triggered by new social conditions. The retention of some
female gonadal tissue in males facilitates a new change of sex, if and when required[86]. Thus, it
has been argued that in bidirectional sex change it should be less expensive for a male to
maintain a non-functional female gonad than for a female to maintain a non-functional male
gonad[85]: females typically use the full gonad to produce eggs, while males could afford to spare
some tissue as ovarian given that their mating success should not be greatly affected by a slight
reduction in sperm production[86] (e.g., in some monogamous gobies of the genus *Trimma*). The
maintenance of both gonadal tissues could facilitate a transition to simultaneous
hermaphroditism. Consistent with this suggestion, our analyses indicate that evolutionary
transition to simultaneous hermaphroditism are more likely from protogyny than from
protandry: protogyny might change to bidirectional sex change and, with maintenance of both
gonadal tissues, to simultaneous hermaphroditism (Fig. 3, Table 2).
In conclusion, our study reveals that gonochorism is the most likely ancestral state and
the most evolutionary stable sexual system in teleosts. In contrast to theoretical predictions, we
demonstrate that simultaneous hermaphroditism can evolve directly to and from gonochorism
and, overall, the evolution of sexual systems is evolutionarily more dynamic than commonly
assumed. Our results call into question theoretical assumptions that changes between sexual
systems are constrained by sex-specific gene expression, physiology and behaviour. Instead
we propose that the adaptive advantage of different sexual systems is underpinned in fish by

their extraordinary and unique developmental plasticity[87], including common and fast transitions among different sex determining mechanisms[60,65]. Our study also reveals that different sexual systems exhibit different life history strategies that allow species with sequential hermaphroditism to maximise fitness as the second sex[34], and highlights the need for more sex-specific life history data to gain a fuller and deeper understanding of the interplay between life history strategies and sexual system. Altogether we propose that a comprehensive framework that incorporates life history traits, sex determining mechanisms and gonadal plasticity into traditional theoretical models of sexual system adaptive value will be essential if we are to fully understand the evolution of sexual systems, their phylogenetic distribution and their implications for conservation and management.

Methods

Data collection and verification

We compiled the most comprehensive database on sexual systems in teleost fish to date (9005 species). Of these, 4740 (403 hermaphroditic) are included in the most recent and largest phylogeny for the class[88] (Supplementary Data). Information on sexual system were first extracted from FishBase[89] and then verified individually for the majority of the species against the primary literature available to date (Supplementary Methods). Life history traits were also collected from the primary literature and FishBase[90]: longevity (in years), maximum length (in cm); length (in cm) and age at maturity (in years) of males and females; male gonadosomatic index (GSI; Supplementary Table 2).

Phylogenetic comparative analyses

We investigated the evolutionary history of sexual systems of 4740 teleost species using Multistate models in BayesTraits V.3[91,92] in a Bayesian framework. Multistate estimates instantaneous transition rates between alternative character states of a single categorical variable (i.e., the rate of change between states along the branches of a phylogeny), based on a continuous-time Markov model of evolution for discrete traits[93,94]. A high transition rate from one state to another indicates that the first state changes rapidly to the second state over evolutionary time. Therefore, a character state is evolutionarily stable when it is lost more slowly than it is gained[95]. Multistate also produces posterior distributions of the ancestral character state at the root of the phylogeny. We scaled the tree by a default constant (mean of 0.1) in all analyses[95] and used an exponential prior whose mean was seeded from a uniform hyperprior ranging from 0 to 10 to reduce inherent uncertainty and biases of prior choice[92]. We

ran all Multistate analyses with Reversible Jump (RJ) Markov chain Monte Carlo (MCMC) methods. MCMC samples models in direct proportion to their fit to the data, generating a posterior distribution of parameter estimates for each transition rate, and RJ sets some parameters equal to zero or equal to one another, thereby reducing model complexity and over-parametrization[91,92,94]. As a result, posterior distributions of parameter estimates may not be normal; we thus summarised results by presenting mean and mode of the posterior distributions of each parameter estimate, 95% higher posterior density, and percentage of models with parameters estimated to be 0. We ran all MCMC chains for 400 million iterations in addition to a burn-in of half a million iterations, sampling every 200000 iterations. All chains converged and showed good mixing as indicated by their effective sample sizes of 2000 and visual inspections of their traces in Tracer v1.6[96]. All analyses were run in triplicate and the three independent chains converged on very similar solutions, leading to qualitatively similar results. Here we present the results from the first chain. We ran RJ-MCMC Multistate analysis on sexual system (Supplementary Table 3) as a binary state (gonochoristic or hermaphrodite) and as a four state categorical variable (gonochorism, protandry, protogyny, simultaneous hermaphroditism). Bidirectional sequential hermaphrodites were excluded from the latter analysis as the sample size of extant species was too low so that the chains failed to converge and mix properly when sexual system was studied as a five states categorical variable. For the analyses with two character state, we graphed the evolutionary history of sexual systems on the phylogeny using maximum likelihood (ML) in the R package Ape v.5.3[97], which provided a reasonably close approximation of the RJ-MCMC Multistate results (this was not the case with the four character state analysis).

We used phylogenetic generalized least square (PGLS) models[98–100] to test for the association of each life history trait, entered as dependent variables, with sexual systems entered as independent discrete variable with three possible states (gonochorism, protogyny, protandry), as not enough data were available for simultaneous hermaphroditic species and bidirectional sex-changers. PGLS models were run with the R package *caper*[101] in Maximum Likelihood. The parameter λ of PGLS models quantifies the strength of the phylogenetic signal in the model residuals[98]. λ ranges between zero (there is no phylogenetic structure in the data) and one (the species share similarity in trait values directly proportional to their common evolutionary time, under Brownian motion model of evolution[98,100]). Continuous variables were $\log_{10}$-transformed to meet assumptions of normality.

Acknowledgments

This project was supported by Spanish Government Grant Epipure (PID2019-

108888RB-I00) to FP and the Santander Universities Travel Award Grant 2016/2017 to CB

and SP that facilitated the collaboration among the three institutions involved in this study. IC

was supported by NERC (grant no. NE/K013777/1). We acknowledge Dr. Stefano Mariani

(Univ. John Moores, UK) for helpful discussion in the first stages of the project and Drs.

Stephen Weeks (Univ. Akron, USA) and Manfred Schartl (Univ. Würzburg) for helpful

comments on the manuscript.

Author contributions

F.P. conceived the project. S.P., C.B., I.C., and F.P. designed the study. S.P. collected data with

assistance from C.B. Data analyses were performed by S.P., C.B., and I.C. and S.P., C.B., I.C.,

and F.P. wrote the manuscript.

References

1.   Speijer, D., Lukeš, J. & Eliáš, M. Sex is a ubiquitous, ancient, and inherent attribute of
eukaryotic life. *Proc. Natl Acad. Sci. USA* **112,** 8827–8834 (2015).
603

604   2.   Bachtrog, D. *et al.* Sex determination: why so many ways of doing it? *PLoS Biol.* **12,**
605   e1001899 (2014).
606

607   3.   Ah-King, M. & Nylin, S. Sex in an evolutionary perspective: just another reaction norm.
608   *Evol. Biol.* **37,** 234–246 (2010).
609

610   4.   Leonard, J. L. The evolution of sexual systems in animals. In: Leonard J.L. (ed)
611   *Transitions between sexual systems: understanding the mechanisms of, and pathways*
612   *between, dioecy, hermaphroditism and other sexual systems*. pp. 1–58 Cham: Springer
613   (2019).
614

615   5.   Weeks, S. C., Benvenuto, C. & Reed, S. K. When males and hermaphrodites coexist: a
616   review of androdioecy in animals. *Integr. Comp. Biol.* **46,** 449–64 (2006).
617

618   6.   Goldberg, E. E. *et al.* Macroevolutionary synthesis of flowering plant sexual systems.
619   *Evolution* **71,** 898–912 (2017).
620

621   7.   Waples, R. S., Mariani, S. & Benvenuto, C. Consequences of sex change for effective
622   population size. *Proc. R. Soc. B* 285: 20181702 (2018).
623

624   8.   Benvenuto, C. & Weeks, S. C. Hermaphroditism and gonochorism. In: Cothran R.D.
625   and Thiel M. (eds) *The Natural History of the Crustacea: Reproductive Biology.*

Volume VI pp. 197-241 New York: Oxford University Press (2020).

9. Mariani, S., Sala-Bozano, M., Chopelet, J. & Benvenuto, C. Spatial and temporal patterns of size-at-sex-change in two exploited coastal fish. *Environ. Biol. Fish.* **96,** 535-541 (2013).

10. Käfer, J., Marais, G. A. & Pannell, J. R. On the rarity of dioecy in flowering plants. *Mol. Ecol.* **26,** 1225–1241 (2017).

11. Jarne, P. & Auld, J. R. Animals mix it up too: the distribution of self-fertilization among hermaphroditic animals. *Evolution* **60,** 1816–1824 (2006).

12. Leonard, J. L. Williams' paradox and the role of phenotypic plasticity in sexual systems. *Integr. Comp. Biol.* **53,** 671–688 (2013).

13. Weeks, S. C. The role of androdioecy and gynodioecy in mediating evolutionary transitions between dioecy and hermaphroditism in the Animalia. *Evolution* **66,** 3670–3686 (2012).

14. Renner, S. S. The relative and absolute frequencies of angiosperm sexual systems: dioecy, monoecy, gynodioecy, and an updated online database. *Am. J. Bot.* **101,** 1588–1596 (2014).

15. Bawa, K. S. Evolution of dioecy in flowering plants. *Ann. Rev. Ecol. Syst.* **11,** 15–39 (1980).

16. Charlesworth, B. & Charlesworth, D. A model for the evolution of dioecy and gynodioecy. *Am. Nat.* **112,** 975–997 (1978).

17. Charlesworth, D. Androdioecy and the evolution of dioecy. *Biol. J. Linn. Soc.* **22,** 333–348 (1984).

18. Pannell, J. R. The evolution and maintenance of androdioecy. *Ann. Rev. Ecol. Syst.* **33,** 397–425 (2002).

19. Bull, J. & Charnov, E. On irreversible evolution. *Evolution* **39,** 1149–1155 (1985).

20. Barrett, S. C. The evolution of plant reproductive systems: how often are transitions irreversible? *Proc. R. Soc. B* **280,** 20130913 (2013).

21. Oyarzún, P. A., Nuñez, J. J., Toro, J. E. & Gardner, J. P. Trioecy in the marine mussel *Semimytilus algosus* (Mollusca, Bivalvia): stable sex ratios across 22 degrees of a latitudinal gradient. *Front. Mar. Sci.* **7,** 348 (2020).

22. Dani, K. & Kodandaramaiah, U. Plant and animal reproductive strategies: lessons from offspring size and number tradeoffs. *Front. Ecol. Evol.* **5,** 38 (2017).

23. Avise, J. & Mank, J. Evolutionary perspectives on hermaphroditism in fishes. *Sexual Development* **3,** 152–163 (2009).

676   24.   Atz, J. Intersexuality in fishes. In: Amstrong C.N. and Marshall A.J. (eds).
677   *Intersexuality in Vertebrates Including Man*, pp. 145-232. London: Academic Press
678   (1964).

679   25.   Costa, W. J., Lima, S. M. & Bartolette, R. Androdioecy in *Kryptolebias* killifish and the
680   evolution of self-fertilizing hermaphroditism. *Biol. J. Linn. Soc.* **99,** 344–349 (2010).
681

682   26.   Costa, W. Colouration, taxonomy and geographical distribution of mangrove killifishes,
683   the *Kryptolebias marmoratus* species group, in southern Atlantic coastal plains of Brazil
684   (Cyprinodontiformes: Rivulidae). *Ichthyol. Explor. Freshw.* **27,** 183–192 (2016).
685

686   27.   Powell, M. L., Kavanaugh, S. I. & Sower, S. A. Seasonal concentrations of reproductive
687   steroids in the gonads of the Atlantic hagfish, *Myxine glutinosa. J. Exp. Zoolog. Part A
688   Comp. Exp. Biol.* **301,** 352–60 (2004).
689

690   28.   Ghiselin, M. T. The evolution of hermaphroditism among animals. *Q. Rev. Biol.* **44,**
691   189–208 (1969).
692

693   29.   Eppley, S. M. & Jesson, L. K. Moving to mate: the evolution of separate and combined
694   sexes in multicellular organisms. *J. Evol. Biol.* **21,** 727–36 (2008).
695

696   30.   Warner, R. R. The adaptive significance of sequential hermaphroditism in animals. *Am.
697   Nat.* **109,** 61–82 (1975).
698

699   31.   Warner, R. R., Robertson, D. R. & Leigh, E. G. Sex change and sexual selection.
700   *Science* **190,** 633–638 (1975).
701

702   32.   Charnov, E. L. *The Theory of Sex Allocation*. New Jersey: Princeton University Press
703   (1982).
704

705   33.   Policansky, D. Sex change in plants and animals. *Ann. Rev. Ecol. Syst.* **13,** 471–495
706   (1982).
707

708   34.   Benvenuto, C., Coscia, I., Chopelet, J., Sala-Bozano, M. & Mariani, S. Ecological and
709   evolutionary consequences of alternative sex-change pathways in fish. *Sci. Rep.* **7,** 9084
710   (2017).
711

712   35.   Buxton, C. D. & Garratt, P. A. Alternative reproductive styles in seabreams (Pisces:
713   Sparidae). *Envir. Biol. Fishes* **28,** 113–124 (1990).
714

715   36.   Shapiro, D. Y. Social behavior, group structure, and the control of sex reversal in
716   hermaphroditic fish. *Advances in the Study of Behavior* **10,** 43–102 (1979).
717

718   37.   Stearns, S. C. Life history evolution: successes, limitations, and prospects.
719   *Naturwissenschaften* **87,** 476–486 (2000).
720

721   38.   Waples, R. S., Luikart, G., Faulkner, J. R. & Tallmon, D. A. Simple life-history traits
722   explain key effective population size ratios across diverse taxa. *Proc. R. Soc. B* **280,**
723   20131339 (2013).
724

725 39. Martinez, A. S., Willoughby, J. R. & Christie, M. R. Genetic diversity in fishes is
726   influenced by habitat type and life-history variation. *Ecol. Evol.* **8,** 12022–12031 (2018).
727

728 40. Charnov, E. L. Natural selection and sex change in pandalid shrimp: test of a life-
729   history theory. *Am. Nat.* **113,** 715–734 (1979).
730

731 41. Broquet, T. *et al.* The size advantage model of sex allocation in the protandrous sex-
732   changer *Crepidula fornicata*: role of the mating system, sperm storage, and male
733   mobility. *Am. Nat.* **186,** 404–420 (2015).
734

735 42. Erisman, B. E., Craig, M. T. & Hastings, P. A. A phylogenetic test of the size-advantage
736   model: evolutionary changes in mating behavior influence the loss of sex change in a
737   fish lineage. *Am. Nat.* **174,** E83–E99 (2009).
738

739 43. Brandl, S. J. & Bellwood, D. R. Pair-formation in coral reef fishes: an ecological
740   perspective. *Oceanogr. Mar. Biol.* **52,** 1–80 (2014).
741

742 44. Fitzpatrick, J. L. Sperm competition and fertilization mode in fishes. *Phil. Trans. R.
743   Soc. B* **375,** 20200074 (2020).
744

745 45. Parker, G. A. Conceptual developments in sperm competition: a very brief synopsis.
746   *Phil. Trans. R. Soc B* **375,** 20200061 (2020).
747

748 46. Warner, R. R. Sex change in fishes: hypotheses, evidence, and objections. *Environ.
749   Biol. Fishes* **22,** 81–90 (1988).
750

751 47. Molloy, P. P., Goodwin, N. B., Côté, I. M., Reynolds, J. D. & Gage, M. J. Sperm
752   competition and sex change: a comparative analysis across fishes. *Evolution* **61,** 640–
753   652 (2007).
754

755 48. Erisman, B. E., Petersen, C. W., Hastings, P. A. & Warner, R. R. Phylogenetic
756   perspectives on the evolution of functional hermaphroditism in teleost fishes. *Integr.
757   Comp. Biol.* **53**, 736-754 (2013).
758

759 49. Sadovy, Y., Colin, P. & Domeier, M. Aggregation and spawning in the tiger grouper,
760   *Mycteroperca tigris* (Pisces: Serranidae). *Copeia* **1994,** 511–516 (1994).
761

762 50. Muñoz, R. C. & Warner, R. R. A new version of the size-advantage hypothesis for sex
763   change: incorporating sperm competition and size-fecundity skew. *Am. Nat.* **161,** 749–
764   761 (2003).
765

766 51. Horne, C. R., Hirst, A. G. & Atkinson, D. Selection for increased male size predicts
767   variation in sexual size dimorphism among fish species. *Proc. R. Soc. B* **287,** 20192640
768   (2020).
769

770 52. Parker, G. The evolution of expenditure on testes. *J. Zool.* **298,** 3–19 (2016).
771

772 53. Stockley, P., Gage, M., Parker, G. & Møller, A. Sperm competition in fishes: the
773   evolution of testis size and ejaculate characteristics. *Am. Nat.* **149,** 933–954 (1997).
774

775 54. Pla, S., Benvenuto, C., Capellini, I. & Piferrer, F. A phylogenetic comparative analysis
776   on the evolution of sequential hermaphroditism in seabreams (Teleostei: Sparidae). *Sci*
777   *Rep* **10,** 3606 (2020).
778
779 55. Vrijenhoek, R. C. Unisexual fish: model systems for studying ecology and evolution.
780   *Ann. Rev. Ecol. Syst.* **25,** 71–96 (1994).
781
782 56. Sadovy de Mitcheson, Y. & Liu, M. Functional hermaphroditism in teleosts. *Fish Fish.*
783   **9,** 1–43 (2008).
784
785 57. Collin, R. & Miglietta, M. P. Reversing opinions on Dollo's law. *Trends Ecol. Evol.* **23,**
786   602–9 (2008).
787
788 58. Domes, K., Norton, R. A., Maraun, M. & Scheu, S. Reevolution of sexuality breaks
789   Dollo's law. *Proc. Natl Acad. Sci. USA* **104,** 7139–7144 (2007).
790
791 59. Dollo, L. Les lois de l'évolution. *Bulletin de la Société belge de géologie, de*
792   *paléontologie et d'hydrologie* **7,** 164–166 (1893).
793
794 60. Pennell, M. W., Mank, J. E. & Peichel, C. L. Transitions in sex determination and sex
795   chromosomes across vertebrate species. *Mol. Ecol.* **27,** 3950–3963 (2018).
796
797 61. Vicoso, B. Molecular and evolutionary dynamics of animal sex-chromosome turnover.
798   *Nat. Ecol. Evol.* 1–10 (2019).
799
800 62. Gamble, T. *et al.* Restriction site-associated DNA sequencing (RAD-seq) reveals an
801   extraordinary number of transitions among gecko sex-determining systems. *Mol. Biol.*
802   *Evol.* **32,** 1296–1309 (2015).
803
804 63. Pokorná, M. & Kratochvíl, L. Phylogeny of sex-determining mechanisms in squamate
805   reptiles: are sex chromosomes an evolutionary trap? *Zool. J. Linn. Soc.* **156,** 168–183
806   (2009).
807
808 64. Adolfi, M. C., Nakajima, R. T., Nóbrega, R. H. & Schartl, M. Intersex,
809   hermaphroditism, and gonadal plasticity in vertebrates: evolution of the Müllerian duct
810   and Amh/Amhr2 signaling. *Annu. Rev. Anim. Biosci.* **7:**7.1–7.24 (2018).
811
812 65. Uller, T. & Helanterä, H. From the origin of sex-determining factors to the evolution of
813   sex-determining systems. *Q. Rev. Biol.* **86,** 163–180 (2011).
814
815 66. Devlin, R. H. & Nagahama, Y. Sex determination and sex differentiation in fish: an
816   overview of genetic, physiological, and environmental influences. *Aquaculture* **208,**
817   191–364 (2002).
818
819 67. Volff, J.-N., Nanda, I., Schmid, M. & Schartl, M. Governing sex determination in fish:
820   regulatory putsches and ephemeral dictators. *Sexual Development* **1,** 85–99 (2007).
821
822 68. Nagahama, Y., Chakraborty, T., Paul-Prasanth, B., Ohta, K. & Nakamura, M. Sex
823   determination, gonadal sex differentiation and plasticity in vertebrate species.

824   *Physiological Reviews* (2020). https://doi.org/10.1152/physrev.00044.2019
825

826   69.   Penman, D. J. & Piferrer, F. Fish gonadogenesis. Part I: genetic and environmental
827   mechanisms of sex determination. *Reviews in Fisheries Science* **16(S1),** 16–34 (2008).
828

829   70.   Mank, J. E., Promislow, D. E. L. & Avise, J. C. Evolution of alternative sex-
830   determining mechanisms in teleost fishes. *Biol. J. Linn. Soc.* **87,** 83–93 (2006).
831

832   71.   Galetti, P. M., Aguilar, C. T. & Molina, W. F. An overview of marine fish cytogenetics.
833   *Hydrobiologia* **420,** 55–62 (2000).
834

835   72.   Yoshida, K. *et al.* Sex chromosome turnover contributes to genomic divergence
836   between incipient stickleback species. *PLoS Genetics* **10,** e1004223 (2014).
837

838   73.   Ross, J. A., Urton, J. R., Boland, J., Shapiro, M. D. & Peichel, C. L. Turnover of sex
839   chromosomes in the stickleback fishes (Gasterosteidae). *PLoS Genetics* **5,** e1000391
840   (2009).
841

842   74.   Carvalho, N. D. M. *et al.* Cytogenetics of Synbranchiformes: a comparative analysis of
843   two *Synbranchus* Bloch, 1795 species from the Amazon. *Genetica* **140,** 149–58 (2012).
844

845   75.   Grant, S. *et al.* Genetics of sex determination in flowering plants. *Dev. Genet.* **15,** 214–
846   230 (1994).
847

848   76.   Harrington Jr, R. W. How ecological and genetic factors interact to determine when
849   self-fertilizing hermaphrodites of *Rivulus marmoratus* change into functional secondary
850   males, with a reappraisal of the modes of intersexuality among fishes. *Copeia* 389–432
851   (1971).
852

853   77.   Piferrer, F. Epigenetic mechanisms in sex determination and in the evolutionary
854   transitions between sexual systems. *Phil. Trans. R. Soc B* in press (2021).
855

856   78.   Capel, B. Vertebrate sex determination: evolutionary plasticity of a fundamental switch.
857   *Nat. Rev. Genet.* **18,** 675–689 (2017).
858

859   79.   Adkins-Regan, E. Early Organizational Effects of Hormones: An Evolutionary
860   Perspective. In: Adler N.T. (ed) *Neuroendocrinology of Reproduction: Physiology and*
861   *Behavior* pp.159–228 Boston: Springer (1981).
862

863   80.   Navara, K. J. Mechanisms of environmental sex determination in fish, amphibians, and
864   reptiles. In: *Choosing Sexes* 213–240 Cham: Springer (2018).
865

866   81.   Orban, L., Sreenivasan, R. & Olsson, P.E. Long and winding roads: testis differentiation
867   in zebrafish. *Mol. Cell. Endocrinol.* **312,** 35–41 (2009).
868

869   82.   Zohar, Y., Abraham, M. & Gordin, H. The gonadal cycle of the captivity-reared
870   hermaphroditic teleost *Sparus aurata* (L.) during the first two years of life. *Ann. Biol.*
871   *Anim. Bioch. Biophys.* **18,** 877–882 (1978).
872

873   83.   Chang, C.-F. & Yueh, W.-S. Annual cycle of gonadal histology and steroid profiles in
874   the juvenile males and adult females of the protandrous black porgy, *Acanthopagrus*
875   *schlegeli. Aquaculture* **91,** 179–196 (1990).
876
877   84.   Miura, S., Nakamura, S., Kobayashi, Y., Piferrer, F. & Nakamura, M. Differentiation of
878   ambisexual gonads and immunohistochemical localization of P450 cholesterol side-
879   chain cleavage enzyme during gonadal sex differentiation in the protandrous
880   anemonefish, *Amphiprion clarkii. Comparative Biochemistry and Physiology Part B:*
881   *Biochemistry and Molecular Biology* **149,** 29–37 (2008).
882
883   85.   Yamaguchi, S. & Iwasa, Y. Advantage for the sex changer who retains the gonad of the
884   nonfunctional sex. *Behav. Ecol. Sociobiol.* **71,** 39 (2017).
885
886   86.   Munday, P. L., Kuwamura, T. & Kroon, F. J. Bidirectional sex change in marine fishes.
887   In: *Reproduction and sexuality in marine fishes: Patterns and processes* 241–271
888   (2010).
889
890   87.   Uller, T., Feiner, N., Radersma, R., Jackson, I. S. & Rago, A. Developmental plasticity
891   and evolutionary explanations. *Evol. Dev.* **22,** 47–55 (2020).
892
893   88.   Rabosky, D.L., Chang, J., Title, P.O., Cowman, P.F., Sallan, L., Friedman, M.,
894   Kaschner, K., Garilao, C., Near, T.J., Coll, M. & Alfaro, M.E. An inverse latitudinal
895   gradient in speciation rate for marine fishes. *Nature*, **559,** 392-395 (2018).
896
897   89.   Froese, R., Pauly, D. (eds) FishBase. World Wide Web electronic publication (2018).
898   www.fishbase.org
899
900   90.   Boettiger, C., Lang, D. T. & Wainwright, P. C. rfishbase: exploring, manipulating and
901   visualizing FishBase data from R. *J. Fish Biol.* **81,** 2030–9 (2012).
902
903   91.   Pagel, M., Meade, A. & Barker, D. Bayesian estimation of ancestral character states on
904   phylogenies. *Syst. Biol.* **53,** 673–684 (2004).
905
906   92.   Pagel, M. & Meade, A. Bayesian analysis of correlated evolution of discrete characters
907   by reversible-jump Markov chain Monte Carlo. *Am. Nat.* **167,** 808–25 (2006).
908
909   93.   Pagel, M. Detecting correlated evolution on phylogenies: a general method for the
910   comparative analysis of discrete. *Proc. R. Soc. B* **255,** 37-45 (1994).
911
912   94.   Currie, T. E. & Meade, A. Keeping yourself updated: Bayesian approaches in
913   phylogenetic comparative methods with a focus on Markov chain models of discrete
914   character evolution. In: *Modern phylogenetic comparative methods and their*
915   *application in evolutionary biology* pp. 263–286 Berlin, Heidelberg: Springer (2014).
916
917   95.   Furness, A. I. & Capellini, I. The evolution of parental care diversity in amphibians.
918   *Nat. Commun.* **10,** 1–12 (2019).
919
920   96.   Rambaut, A., Drummond, A. J., Xie, D., Baele, G. & Suchard, M. A. Posterior
921   summarization in Bayesian phylogenetics using Tracer 1.7. *Syst. Biol.* **67,** 901 (2018).
922

97. Paradis, E. & Schliep, K. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* **35,** 526–528 (2019).

98. Freckleton, R. P., Harvey, P. H. & Pagel, M. Phylogenetic analysis and comparative data: a test and review of evidence. *Am. Nat.* **160,** 712–726 (2002).

99. Pagel, M. Inferring evolutionary processes from phylogenies. *Zool. Scr.* **26,** 331–348 (1997).

100. Pagel, M. Inferring the historical patterns of biological evolution. *Nature* **401,** 877–84 (1999).

101. Orme, D. The caper package: Comparative analysis of phylogenetics and evolution in R (2013). See *https://cran. r-project. org/web/packages/caper/vignettes/caper.pdf* (2018).

Switches, stability and reversals:
the evolutionary history of sexual systems in fish

Susanna Pla[1#], Chiara Benvenuto[2#], Isabella Capellini[3*] & Francesc Piferrer[1*]

[1]Institut de Ciències del Mar, Spanish National Research Council (CSIC), Barcelona, Spain

[2]School of Science, Engineering and Environment, University of Salford, Salford, UK

[3]School of Biological Sciences, Queen's University Belfast, UK

#Joint first authors

*Correspondence: Dr. Francesc Piferrer, Institut de Ciències del Mar, Spanish National Research Council (CSIC), Passeig Marítim, 37-49, 08003 Barcelona. Tel. +34-932 309 567; E-mail: piferrer@icm.csic.es

*Correspondence: Dr. Isabella Capellini, School of Biological Sciences, Queen's University Belfast, 19 Chlorine Gardens, Belfast, BT9 5DL, Northern Ireland, United Kingdom; E-mail: I.Capellini@qub.ac.uk

Supplementary information

Supplementary figure

Supplementary Fig. 1. RJ-MCMC Multistate posterior distributions of the transition rates (two character state) from gonochorism (n = 4337 species) to hermaphroditism (n = 403 species) in light blue and from hermaphroditism to gonochorism in grey.

Supplementary tables

Supplementary Table 1. Results of the RJ-MCMC Multistate analysis in BayesTraits of sexual systems as a binary trait: gonochorism (G) or hermaphroditism (H). For each posterior distribution, we report the effective sample size (ESS), the mean and 95% high posterior density (95-HPD) intervals, the mode, and the percentage of models in which the parameter is estimated as zero. This analysis is based on 4740 extant teleost species (G: n = 4337; H: n = 403)

Transition rates	ESS	Mean	95-HPD		Mode	% Zero
G → H	2000	0.047	0.032 – 0.065		0.045	0
H → G	2000	0.864	0.663 – 1.098		0.831	0
Root probabilities						
G	2000	63.8	57.3 – 70.7		64.5	0
H	2000	36.2	29.3 – 42.7		35.5	0

Supplementary Table 2. Sample sizes for longevity, maximum length, age at maturity and length at maturity for each sexual system with sex-specific male (♂) and female (♀), when data were available. In italics data not used in the analyses (due to the small male size). G = gonochorism; H = hermaphroditism; PG = protogyny; PA = protandry; SH = simultaneous hermaphroditism; BD = Bidirectional hermaphroditism

Life-history traits	Sex	G	PG	PA	Total for analyses	*SH*	*BD*
Longevity		759	88	19	866	*7*	*2*
Max length		2624	243	36	2903	*28*	*10*
Age maturity	♂	258	20	12	290	*1*	-
	♀	281	33	5	319	*2*	-
Length maturity	♂	303	50	18	371	*9*	-
	♀	295	43	13	351	*2*	-
GSI	♂	45	45	15	105	*3*	-

Supplementary Table 3. Number of species used to study the evolutionary transitions among different sexual systems, out of a total sample of 4740 teleosts. Note that androdioecious species are not included in the analyses

Sexual system	2 state [G, H]	4 state [G, PG, PA, SH]
Gonochorism [G]	4337	4337
Hermaphroditism [H]	403*	
Protogyny [PG]		285
Protandry [PA]		57
Simultaneous hermaphroditism [SH]		47
Total	4740	4726

*Includes 14 (13 verified against primary literature) species of bidirectional sequential hermaphrodites, which were not included in other analyses due to their small sample size as a separate category.

Supplementary methods

Data collection and verification The sexual system was extracted from FishBase (www.fishbase.org) and the classification of 403 hermaphroditic species was cross-checked against more recent primary literature: 67% of those species (i.e., more than two out of three) were confirmed to be functional hermaphrodites (see Supplementary Data and full references below).

The following life history traits were obtained from FishBase (www.fishbase.org; CD-ROM version) and updated with primary literature:

- Longevity (in years): also updated with data extracted with rfishbase (Boettiger et al., 2012; https://www.rdocumentation.org/packages/rfishbase/versions/3.0.4). When more than one value was present for species, we used the max value reported (in the wild);
- maximum length (TL: total length; in cm);
- sex-specific age (in years) and length (in cm) at maturity for each sex.

See supplementary Data for full dataset attached as an Excel file.

Supplementary references used to verify functional hermaphroditism in fish

Abd-el-Aziz, S.H. & Ramadan, A.A. (1990) Sexuality and hermaphroditism in fishes. I. Synchronous functional hermaphroditism in the serranid fish *Serranus scriba* L. *Folia morphologica, 38*(1), 86-100.

Abou-Seedo, F., Wright, J.M. & Clayton, D. (1990) Aspects of the biology of *Diplodus sargus kotschyi* (Sparidae) from Kuwait Bay. *Cybium, 14*(3), 217-223.

Abou-Seedo, F.S., Dadzie, S. & Al-Kanaan, K.A. (2003) Sexuality, sex change and maturation patterns in the yellowfin seabream, *Acanthopagrus latus* (Teleostei: Sparidae) (Houttuyn, 1782). *Journal of Applied Ichthyology, 19*(2), 65-73. http://doi.org/ 10.1046/j.1439-0426.2003.00355.x

Abu-Hakima, R. (1984) Some aspects of the reproductive biology of Acanthopagrus spp. (Family: Sparidae). *Journal of Fish Biology, 25*(5), 515-526. http://doi.org/10.1111/j.1095-8649.1984.tb04898.x

Adams, S. (2003) Morphological ontogeny of the gonad of three plectropomid species through sex differentiation and transition. *Journal of Fish Biology, 63*(1), 22-36. http://doi.org/10.1046/j.1095-8649.2003.00098.x

Adreani, M.S. & Allen, L.G. (2008) Mating system and reproductive biology of a temperate wrasse, *Halichoeres semicinctus*. *Copeia, 2008*(2), 467-475. http://doi.org/10.1643/CP-06-265

Adreani, M.S., Erisman, B.E. & Warner, R.R. (2004) Courtship and spawning behavior in the California sheephead, *Semicossyphus pulcher* (Pisces: Labridae). *Environmental Biology of Fishes, 71*(1), 13-19. http://doi.org/10.1023/B:EBFI.0000043177.98895.f7

Aldenhoven, J.M. (1984) S*ocial organization and sex change in an angelfish, Centropyge bicolor on the Great Barrier Reef.* Unpublished PhD dissertation, Macquarie University, North Ryde, Australia

Alekseev, F.E. (1982) Hermaphroditism in sparid fishes (Perciformes, Sparidae). I. Protogyny in porgies, *Pagrus pagrus, P. orphus, P. ehrenbergi* and *P. auriga*, from West Africa. *Journal of Ichthyology, 22*, 85-94.

Alekseev, F.E. (1983) Hermaphroditism in porgies (Perciformes, Sparidae). II. Sexual structure of the populations, mechanism of its formation and evolution in scups, *Pagrus pagrus, P. orphus, P. ehrenbergi* and *P. auriga. Journal of Ichthyology, 23*, 61-67.

Anderson, W.D. & Baldwin, C.C. (2000) A new species of Anthias (Teleostei: Serranidae: Anthiinae) from the Galápagos Islands, with keys to Anthias and eastern Pacific Anthiinae. *Proceedings of the Biological Society of Washington, 113*(2), 369-385.

Aoyama, T. (1955) On the hermaphroditism in the yellow sea bream, *Taius tumifrons*. *Japanese Journal of Ichthyology, 4*(4-6), 119-129. http://doi.org/10.11369/jji1950.4.119

Aronov, A. & Goren, M. (2008) Ecology of the mottled grouper (*Mycteroperca rubra*) in the eastern Mediterranean. *Electronic Journal of Ichthyology, 2*, 43-55.

Asoh, K. (2004) Gonadal development in the coral reef damselfish *Dascyllus flavicaudus* from Moorea, French Polynesia. *Marine Biology, 146*(1), 167-179. http://doi.org/10.1007/s00227-004-1414-7

Asoh, K. (2005) Gonadal development and diandric protogyny in two populations of

Dascyllus reticulatus from Madang, Papua New Guinea. *Journal of Fish Biology, 66*(4), 1127-1148. http://doi.org/10.1111/j.0022-1112.2005.00669.x

Asoh, K. & Yoshikawa, T. (2003) Gonadal development and an indication of functional protogyny in the Indian damselfish (*Dascyllus carneus*). *Journal of Zoology, 260*(1), 23-39. http://doi.org/10.1017/S0952836903003418

Bacci, G. & Razzauti, A. (1958) Protogynous hermaphroditism in *Coris julis* L. *Nature, 181*(4606), 432-433.

Barlow, G.W. (1975) On the sociobiology of some hermaphroditic serranid fishes, the hamlets, in Puerto Rico. *Marine Biology, 33*(4), 295-300. http://doi.org/10.1007/BF00390567

Bean, K., Mapstone, B.D., Davies, C.R., Murchie, C.D. & Williams, A.J. (2003) Gonad development and evidence of protogyny in the red-throat emperor on the Great Barrier Reef. *Journal of Fish Biology, 62*(2), 299-310. http://doi.org/10.1046/j.1095-8649.2003.00021.x

Bentivegna, F. & Rasotto, M.B. (1987) Protogynous hermaphroditism in *Xyrichthys novacula* (L. 1758). *Cybium, 11*(1), 75-78.

Bertrand, J. (1986) Données concernant la reproduction de *Lethrinus mahsena* (Forsskal 1775) sur les bancs de Saya de Malha (Océan Indien)." *Cybium (Paris)* p. 15-29.

Besseau, L. & Bruslé-Sicard, S. (1995) Plasticity of gonad development in hermaphroditic sparids: ovotestis ontogeny in a protandric species, *Lithognathus mormyrus*. *Environmental Biology of Fishes, 43*(3), 255-267. http://doi.org/10.1007/BF00005857

Bhandari, R.K., Alam, M.A.l., Higa, M., Soyano, K. & Nakamura, M. (2005) Evidence that estrogen regulates the sex change of honeycomb grouper (*Epinephelus merra*), a protogynous hermaphrodite fish. *Journal of Experimental Zoology Part A: Comparative Experimental Biology, 303*(6), 497-503. http://doi.org/10.1002/jez.a.178

Bhandari, R.K., Komuro, H., Nakamura, S., Higa, M. & Nakamura, M. (2003) Gonadal restructuring and correlative steroid hormone profiles during natural sex change in protogynous honeycomb grouper (*Epinephelus merra*). *Zoological science, 20*(11), 1399-1405. http://doi.org/10.2108/zsj.20.1399

Bortone, S.A. (1971) Studies on the biology of the sand perch, *Diplectrum formosum* (Perciformes: Serranidae). *Florida Department of Natural Resources, Marine Research Laboratory Series 65*, 1-27.

Bortone, S.A. (1977a) Gonad morphology of the hermaphroditic fish *Diplectrum pacificum* (Serranidae). *Copeia 1977*(3), 448-453. http://doi.org/10.2307/1443262

Bortone, S.A. (1977b) Observations on the life history of *Diplectrum pacificum* and *D. macropoma* (Pisces, Serranidae) from the Bay of Panama. *Pacific Science, 31*(1), 49-60.

Bortone, S.A. (1977c) Revision ofthe sea basses ofthe genus Diplectrum (Pisces: Serranidae). *U.S. Department of Commerce, NOAA Technical Reports, 404*, 1-49.

Brickle, P., Laptikhovsky, V., & Arkhipkin, A. (2005). Reproductive strategy of a primitive temperate notothenioid Eleginops maclovinus. *Journal of Fish Biology*, *66*(4), 1044-1059.

Bruce, R.W. (1980) Protogynous hermaphroditism in two marine angelfishes. *Copeia, 1980*(2), 353-355. http://doi.org/10.2307/1444017

Brulé, T., Colás-Marrufo, T.E., Tuz-Sulub, A. & Déniel, C. (2000) Evidence for protogynous

hermaphroditism in the serranid fish *Epinephelus drummondhayi* (Perciformes: Serranidae) from the Campeche Bank in the southern Gulf of Mexico. *Bulletin of Marine Science, 66*(2), 513-521.

Brulé, T., Déniel, C., Colás-Marrufo, T. & Sánchez-Crespo, M. (1999) Red grouper reproduction in the southern Gulf of Mexico. *Transactions of the American Fisheries Society, 128*(3), 385-402.

Bruslé, J. & Bruslé, S. (1975) Ovarian and testicular intersexuality in two protogynous mediterranean groupers, *Epinephelus aeneus* and *Epinephelus guaza*. In: R. Reinboth (Ed.), *Intersexuality in the Animal Kingdom* (pp. 222-227). Springer, Berlin, Heidelberg. http://doi.org/10.1007/978-3-642-66069-6_21

Bruslé, S. (1983) Contribution to the sexuality of a Hermaphroditic teleost, *Serranus hepatus* L. *Journal of Fish Biology, 22*(3), 283-292. http://doi.org/10.1111/j.1095-8649.1983.tb04752.x

Bruslé, S., Debas, L. & Cauty, C. (1989) Morphological and cytological aspects of sex inversion in a protogynous hermaphrodite, *Epinephelus microdon* (Teleostei, Serranidae). *Advances in Tropical Aquaculture*, nº9, chap.53, 559 –562.

Brusléa, S. (1987) Sex-inversion of the hermaphroditic, protogynous teleost *Coris julis* L. (Labridae). *Journal of Fish Biology, 30*(5), 605-616. http://doi.org/10.1111/j.1095-8649.1987.tb05788.x

Bullock, L.H. & Murphy, M.D. (1994) Aspects of the life history of the yellowmouth grouper, *Mycteroperca interstitialis*, in the eastern Gulf of Mexico. *Bulletin of Marine Science, 55*(1), 30-45.

Bullock, L.H. & Smith, G.B. (1991) Seabasses (Pisces: Serranidae). *Memoirs of the Hourglass Cruises, 8*, 1-243.

Burgos, J.M., Sedberry, G.R., Wyanski, D.M. & Harris, P.J. (2007) Life history of red grouper (*Epinephelus morio*) off the coasts of North Carolina and South Carolina. *Bulletin of Marine Science, 80*(1), 45-65.

Burnett-Herkes, J.N. (1975) *Contributions to the biology of the red hind, Epinephelus guttatus, a commercially important serranid fish from the tropical western Atlantic.* PhD dissertation, University of Miami., Coral Gables, 154 pp.

Burton, M. L., Potts, J. C., Poholek, A. J., Ostrowski, A., & Page, J. (2019). Age, growth, natural mortality, and reproductive seasonality of knobbed porgy from southeastern United States waters. *Marine and Coastal Fisheries, 11*(2), 231-245.

Buxton, C.D. (1989) Protogynous hermaphroditism in *Chrysoblephus laticeps* (Cuvier) and *C. cristiceps* (Cuvier) (Teleostei: Sparidae). *African Zoology, 24*(3), 212-216.

Buxton, C.D. & Clarke, J.R. (1986) Age, growth and feeding of the blue hottentot *Pachymetopon aeneum* (Pisces: Sparidae) with notes on reproductive biology. *African Zoology, 21*(1), 33-38.

Buxton, C.D. & Garratt, P.A. (1990) Alternative reproductive styles in seabrearns (Pisces: Sparidae). In: M.N. Bruton (Ed.), *Alternative life-history styles of fishes* (pp. 113-124). Springer, Dordrecht. http://doi.org/10.1007/978-94-009-2065-1_7

Cabiddu, S., Follesa, M.C., Porcu, C. & Cau, A. (2010) Gonad development and reproduction in the monoecious species *Chlorophthalmus agassizi* (Actinopterygii: Aulopiformes: Chlorophthalmidae) from the sardinian waters (Central-Western Mediterranean). *Acta Ichthyologica et Piscatoria, 40*, 167-177.

Calvo, J., Morriconi, E., Rae, G.A. & San Roman, N.A. (1992) Evidence of protandry in a subantarctic nototheniid, *Eleginops maclovinus* (Cuv. & Val., 1830) from the Beagle

Channel, Argentina. *Journal of Fish Biology, 40*(2), 157-164. http://doi.org/10.1111/j.1095-8649.1992.tb02563.x

Candi, G., Castriota, L., Andaloro, F., Finoia, M.G. & Marino, G. (2004) Reproductive cycle and sex inversion in razor fish, a protogynous labrid in the southern Mediterranean Sea. *Journal of Fish Biology, 64*(6), 1498-1513. http://doi.org/10.1111/j.0022-1112.2004.0404.x

Chang-Po, C., Hwey-Lian, H. & Kun-Hsiung, C. (1980) Some aspects of the sex change and reproductive biology of the grouper, *Epinephelus diacanthus* (Cuvier et valenciensis). *Bulletin of the Institute of Zoology, Academia Sinica, 19*(1), 11-17.

Chang, C.-F. & Yueh, W.-S. (1990) Annual cycle of gonadal histology and steroid profiles in the juvenile males and adult females of the protandrous black porgy, *Acanthopagrus schlegeli. Aquaculture,* **91,** 179-196. http://doi.org/10.1016/0044-8486(90)90187-R

Chen, F.Y., Chow, M., Chao, T.M. & Lim, R. (1977) Artificial spawning and larval rearing of the grouper, *Epinephelus tauvina* (Forskal) in Singapore. *Singapore Journal of Primary Industries,* **5,** 1-21.

Choat, J.H. & Robertson, D.R. (1975) Protogynous hermaphroditism in fishes of the family Scaridae. In: R. Reinboth (Ed.), *Intersexuality in the Animal Kingdom* (pp. 263-283). Springer, Berlin, Heidelberg. http://doi.org/10.1007/978-3-642-66069-6_26

Clark, E. & Shen, D. (1986) Territoriality of Red Sea sand-diving fishes of the genera Xyrichtys and Trichonotus. In: T. Uyeno, R. Arai, T. Taniuchi, & K., Matsuura, (Eds.), *Indo-Pacific Fish Biology: Proceedings of the Second International Conference on Indo-Pacific Fishes, Tokyo.*

Coates, D. (1982) Some observations on the sexuality of humbug damselfish, *Dascyllus aruanus* (Pisces, Pomacentridae) in the field. *Ethology, 59*(1), 7-18. http://doi.org/10.1111/j.1439-0310.1982.tb00328.x

Coetzee, P.S. (1986) Diet composition and breeding cycle of blacktail, *Diplodus sargus capensis* (Pisces: Sparidae), caught off St Croix Island, Algoa Bay, South Africa. *African Zoology, 21*(3), 237-243.

Cole, K.S. (1983) Protogynous hermaphroditism in a temperate zone territorial marine goby, *Coryphopterus nicholsi. Copeia, 1983*(3), 809-812. http://doi.org/10.2307/1444350

Cole, K.S. (1990) Patterns of gonad structure in hermaphroditic gobies (Teleostei Gobiidae). *Environmental Biology of Fishes, 28*(1-4), 125-142. http://doi.org/10.1007/BF00751032

Cole, K.S. (2002) Gonad morphology, sexual development, and colony composition in the obligate coral-dwelling damselfish *Dascyllus aruanus. Marine Biology, 140*(1), 151-163. http://doi.org/10.1007/s002270100681

Cole, K.S. (2003) Hermaphroditic characteristics of gonad morphology and inferences regarding reproductive biology in Caracanthus (Teleostei, Scorpaeniformes). *Copeia, 2003*(2), 68-80. http://doi.org/10.1643/0045-8511(2003)003[0068:HCOGMA]2.0.CO;2

Cole, K.S. (2010) Gonad morphology in hermaphroditic gobies. In: K.S. Cole (Ed.), *Reproduction and Sexuality in Marine Fishes: Patterns and Processes* (pp. 117–164). University of California Press, Berkeley, CA.

Cole, K.S. & Hoese, D.F. (2001) Gonad morphology, colony demography and evidence for hermaphroditism in *Gobiodon okinawae* (Teleostei, Gobiidae). *Environmental*

Biology of Fishes, 61(2), 161-173. http://doi.org/10.1023/A:1011032228716

Cole, K.S. & Robertson, R.D. (1988) Protogyny in the Caribbean reef goby, *Coryphopterus personatus*: gonad ontogeny and social influences on sex-change. *Bulletin of Marine Science, 42*(2), 317-333.

Cole, K.S. & Shapiro, D.Y. (1990) Gonad structure and hermaphroditism in the gobiid genus Coryphopterus (Teleostei: Gobiidae). *Copeia*, *1990*(4), 996-1003. http://doi.org/10.2307/1446482

Cole, K.S. & Shapiro, D.Y. (1992) Gonadal structure and population characteristics of the protogynous goby *Coryphopterus glaucofraenum*. *Marine Biology, 113*(2), 1-9. http://doi.org/10.1007/BF00367632

Cole, K.S. & Shapiro, D.Y. (1995) Social faciliation and sensory mediation of adult sex change in a cryptic, benthic marine goby. *Journal of Experimental Marine Biology and Ecology, 186*(1), 65-75. http://doi.org/10.1016/0022-0981(94)00152-4

Coleman, F.C. (1981) Protogynous hermaphroditism in the anthiine serranid fish *Holanthias martinicensis*. *Copeia, 1981*(4), 893-895.

Coleman, F.C. (1983) *Hemanthias peruanus*—another hermaphroditic anthiine serranid. *Copeia, 1983*(1), 252-253.

Coleman, F.C., Koenig, C.C. & Collins, L.A. (1996) Reproductive styles of shallow-water groupers (Pisces: Serranidae) in the eastern Gulf of Mexico and the consequences of fishing spawning aggregations. *Environmental Biology of Fishes, 47*(2), 129-141. http://doi.org/10.1007/BF00005035

Collins, M. R., Waltz, C. W., Roumillat, W. A., & Stubbs, D. L. (1987). Contribution to the life history and reproductive biology of gag, *Mycteroperca microlepis* (serrandidae), in the South Atlantic bight. *South Carolina State Documents Depository*.

Collins, L.A., Johnson, A.G., Koenig, C.C. & M.S., B.J. (1998) Reproductive patterns, sex ratio, and fecundity in gag, *Mycteroperca microlepis* (Serranidae), a protogynous grouper from the northeastern Gulf of Mexico. *Fishery Bulletin, 96*, 415-427.

Cowen, R.K. (1990) Sex change and life history patterns of the labrid, *Semicossyphus pulcher*, across an environmental gradient. *Copeia, 1990*(3), 787-795. http://doi.org/10.2307/1446444

Crabtree, R.E. & Bullock, L.H. (1998) Age, growth, and reproduction of black grouper, *Mycteroperca bonaci*. *Florida waters. Fishery Bulletin, 96*(4), 735-753.

Craig, M.T. (2007) Preliminary observations on the life history of the white-streaked grouper, *Epinephelus ongus*, from Okinawa, Japan. *Ichthyological Research, 54*(1), 81-84.

Craig, M.T., Hastings, P.A., Pondella, D.J., Ross Robertson, D. & Rosales-Casián, J.A. (2006) Phylogeography of the flag cabrilla *Epinephelus labriformis* (Serranidae): implications for the biogeography of the Tropical Eastern Pacific and the early stages of speciation in a marine shore fish. *Journal of Biogeography, 33*(6), 969-979. http://doi.org/10.1111/j.1365-2699.2006.01467.x

D'Ancona, U. (1949) Ermafroditismo ed intersessualita nei Teleostei. *Experientia, 5*, 381-389.

Dantchakoff, V. (1936) Sur les facteurs de l'histogenes chez des hermaphrodites. *Comptes Renduesociété de Biologie, 123*, 856-858.

Davis, M.P. & Fielitz, C. (2010) Estimating divergence times of lizardfishes and their allies (Euteleostei: Aulopiformes) and the timing of deep-sea adaptations. *Molecular*

Phylogenetics and Evolution, 57, 1194-1208. http://doi.org/10.1016/j.ympev.2010.09.003

Debas, L. (1989) Etude biométrique, histologique et endocrinologique de la sexualité du mérou Epinephelus microdon dans le milieu naturel et en élevage: caractérisation de l'hermaphrodisme protérogyne, description du phénomène d'inversion et mise en évidence du phénomène de reversion. *Thèse de doctorat de l'universite d'Aix-Marseille, France.* 108 pp.

DeLoach, N. & Humann, P. (1999) Reef fish behavior: Florida, Caribbean, Bahamas. Jacksonville, FL. New World Publications.

DeMartini, E.E., Everson, A.R. & Nichols, R.S. (2011) Estimates of body sizes at maturation and at sex change, and the spawning seasonality and sex ratio of the endemic Hawaiian grouper (H*yporthodus quernus,* F. Epinephelidae). *Fishery Bulletin, 109*(1)*,* 123-134.

Dipper, F.A. & Pullin, R.S.V. (1979) Gonochorism and sex-inversion in British Labridae (Pisces). *Journal of Zoology, 187*(1)*,* 97-112. http://doi.org/10.1111/j.1469-7998.1979.tb07716.x

Dulčić, J., Kraljević, M., Grbec, B. & Cetinić, P. (2000) Age, growth and mortality of blotched picarel *Spicara maena* L. (Pisces: Centracanthidae) in the eastern central Adriatic. *Fisheries Research, 48*(1), 69-78. http://doi.org/10.1016/S0165-7836(00)00112-0

Dulčić, J., Pallaoro, A., Cetinić, P., Kraljević, M., Soldo, A. & Jardas, I. (2003) Age, growth and mortality of picarel, *Spicara smaris* L. (Pisces: Centracanthidae), from the eastern Adriatic (Croatian coast). *Journal of Applied Ichthyology, 19*(1), 10-14. http://doi.org/10.1046/j.1439-0426.2003.00345.x

Ebisawa, A. (1990) Reproductive biology of *Lethrinus nebulosus* (Pisces; Lethrinidae) around the Okinawan waters [Japan]. *Bulletin of the Japanese Society of Scientific Fisheries (Japan)*, 56*(12),* 1941-1954. http://doi.org/10.2331/suisan.56.1941

Ebisawa, A. (1997) Some aspects of reproduction and sexuality in the spotcheek emperor, *Lethrinus rubrioperculatus*, in waters off the Ryukyu Islands. *Ichthyological Research, 44*(2-3)*,* 201-212.

Ebisawa, A. (1999) Reproductive and sexual characteristics in the Pacific yellowtail emperor, *Lethrinus atkinsoni*, in waters off the Ryukyu Islands. *Ichthyological Research, 46*(4)*,* 341-358.

Ebisawa, A. (2006) Reproductive and sexual characteristics in five Lethrinus species in waters off the Ryukyu Islands. *Ichthyological Research, 53*(3), 269-280. http://doi.org/10.1007/s10228-006-0345-3

Ebisawa, A., Kanashiro, K., Kyan, T. & Motonaga, F. (1995) Aspects of reproduction and sexuality in the black-spot tuskfish, *Choerodon schoenleinii. Japanese Journal of Ichthyology, 42*(1), 121-130. http://doi.org/10.11369/jji1950.42.121

El-Etreby, S.G., Roberts, C., Ghobashy, A.A. & Zyadah, M.A. (1993) Coral reef groupers (family: Serranidae) in south Sinai: reproduction and hermaphroditism in the fish *Cephalopholis hemistiktos* (Rüppell, 1830). *Journal of the Egyptian German Society of Zoology (Anatomy and Embryology), 12(*B), 1-19.

Erisman, B.E. (2008) Reproductive biology and evolution of epinephelid and serranid fishes (Perciformes, Epinephelidae, Serranidae). Doctoral thesis. Scripps Institute of Oceanography, University of California, San Diego.

Erisman, B.E., Craig, M.T. & Hastings, P.A. (2010) Reproductive biology of the Panama graysby *Cephalopholis panamensis* (Teleostei: Epinephelidae). *Journal of Fish*

Biology, 76(6), 1312-1328.http://doi.org/10.1111/j.1095-8649.2010.02567.x

Etassami, S. (1983) Hermaphroditism in one Sparidae of the Persian Gulf: *Acanthopagrus bifasciatus* (Forssk.). *Cybium, 7*(2)*,* 87-91.

Ferreira, B.P. (1993) Reproduction of the inshore coral trout *Plectropomus maculates* (Perciformes: Serranidae) from the central Great Barrier Reef, Australia. *Journal of Fish Biology, 42*(6), 831-844. http://doi.org/10.1111/j.1095-8649.1993.tb00393.x

Ferreira, B.P. (1995) Reproduction of the common coral trout *Plectropomus leopardus* (Serranidae: Epinephelinae) from the central and northern Great Barrier Reef, Australia. *Bulletin of Marine Science, 56*(2), 653-669.

Fischer, E.A. (1980) The relationship between mating system and simultaneous hermaphroditism in the coral reef fish, *Hypoplectrus nigricans* (Serranidae). *Animal Behaviour, 28*(2), 620-633. http://doi.org/10.1016/S0003-3472(80)80070-4

Fischer, E.A. (1981) Sexual allocation in a simultaneously hermaphroditic coral reef fish. *The American Naturalist, 117*(1), 64-82. http://doi.org/10.1086/283686

Fischer, E.A. (1984) Egg trading in the chalk bass, *Serranus tortugarum*, a simultaneous hermaphrodite. *Zeitschrift für Tierpsychologie, 66*(2), 143-151. http://doi.org/10.1111/j.1439-0310.1984.tb01361.x

Fischer, E.A. & Petersen, C.W. (1987) The evolution of sexual patterns in the seabasses. *Bioscience, 37*(7), 482-489. http://doi.org/10.2307/1310420

Fishelson, L. (1970) Protogynous sex reversal in the fish *Anthias squamipinnis* (Teleostei, Anthiidae) regulated by the presence or absence of a male fish. *Nature, 227*(5253), 90–91.

Fishelson, L. (1975) Ecology and physiology of sex reversal in *Anthias squamipinnis* (Peters),(Teleostei: Anthiidae). In: R. Reinboth (Ed.), *Intersexuality in the animal kingdom* (pp. 284-294). Springer, Berlin, Heidelberg. http://doi.org/10.1007/978-3-642-66069-6_27

Fishelson, L. (1989) Bisexuality and pedogenesis in gobies (Gobiidae: Teleostei) and other fish, or why so many fish in tropical seas? *Senckenbergiana Maritima, 20*, 147-170.

Fishelson, L. (1992) Comparative gonad morphology and sexuality of the Muraenidae (Pisces, Teleostei). *Copeia*, *1992*(1), 197-209. http://doi.org/10.2307/1446552

Fricke, H. & Fricke, S. (1977) Monogamy and sex change by aggressive dominance in coral reef fish. *Nature, 266*(5606), 830-832. http://doi.org/10.1038/266830a0

Fricke, H.W. (1979) Mating system, resource defence and sex change in the anemonefish *Amphiprion akallopisos*. *Zeitschrift für Tierpsychologie, 50*(3), 313-326. http://doi.org/10.1111/j.1439-0310.1979.tb01034.x

Fricke, H.W. (1983) Social control of sex: field experiments with the anemonefish *Amphiprion bicinctus*. *Zeitschrift für Tierpsychologie, 61*(1), 71-77. http://doi.org/ 10.1111/j.1439-0310.1983.tb01327.x

García-Cagide, A. & García, T. (1996) Reproducción de Mycteroperca bonaci y Mycteroperca venenosa (Pisces: Serranidae) en la plataforma cubana. *Revista de Biología Tropical, 44*, 771-780.

García-Cagide, A.R., Claro, R. & García-Arteaga, J.P. (1999) Biología del bonací gato, *Mycteroperca tigris* (Pisces: Serranidae) en la plataforma SW de Cuba. I. Características generales y reproducción. *Revista de Investigaciones Marinas, 20*, 8-14.

García-Díaz, M.M., González, J.A., Lorente, M.J. & Tuset, V.M. (2006) Spawning season,

maturity sizes, and fecundity in blacktail comber (*Serranus atricauda*) (Serranidae) from the eastern-central Atlantic. *Fishery Bulletin, 104*(4), 159-166.

García-Díaz, M.M., Lorente, M.J., González, J.A. & Tuset, V.M. (2002) Morphology of the ovotestis of *Serranus atricauda* (Teleostei, Serranidae). *Aquatic sciences*, *64*(1), 87-96.

García-Díaz, M.M., Tuset, V.M., González, J.A. & Socorro, J. (1997) Sex and reproductive aspects in *Serranus cabrilla* (Osteichthyes: Serranidae): macroscopic and histological approaches. *Marine Biology, 127*(3), 379-386. http://doi.org/10.1007/s002270050024

Garratt, P.A. (1986) Protogynous hermaphroditism in the slinger, *Chrysoblephus puniceus* (Gilchrist & Thompson, 1908) (Teleosteir: Sparidae). *Journal of Fish Biology, 28*(3), 297-306. http://doi.org/10.1111/j.1095-8649.1986.tb05167.x

Ghorbel, M. (1996). Le pageot commun *Pagellus erythrinus* (poison, Sparidae) ecobiologie et etat d'exploitation dans le Golfe de Gabes. *Ph. D. Thesis in Marine Biology and Oceanography, University of Sfax.*

Gilmore, G.R. & Jones, R.S. (1992) Color variation and associated behavior in the epinepheline groupers, *Mycteroperca microlepis* (Goode and Bean) and *M. phenax* Jordan and Swain. *Bulletin of Marine Science, 51*(1), 83-103.

Gillanders, B.M. (1995) Reproductive biology of the protogynous hermaphrodite *Achoerodus viridis* (Labridae) from south-eastern Australia. *Marine and Freshwater Research, 46*(7), 999-1008. http://doi.org/10.1071/MF9950999

Godwin, J. (1994a) Behavioural aspects of protandrous sex change in the anemonefish, *Amphiprion melanopus*, and endocrine correlates. *Animal Behaviour, 48*(3), 551-567. http://doi.org/10.1006/anbe.1994.1275

Godwin, J. (1994b) Historical aspects of protandrous sex change in the anemonefish *Amphiprion melanopus* (Pomacentridae, teleostei). *Journal of Zoology, 232*(2), 199-213. http://doi.org/10.1111/j.1469-7998.1994.tb01569.x

Godwin, J. (1995) Phylogenetic and habitat influences on mating system structure in the humbug damselfishes (Dascyllus, Pomacentridae). *Bulletin of Marine Science, 57*(3), 637-652.

Goeden, G.B. (1978) A monograph of the coral trout, *Plectropomus leopardus* (Lacepede). Research Bulletin No. 1, Queensland Fisheries Service, Brisbane.

Grandcourt, E.M. (2002) Demographic characteristics of a selection of exploited reef fish from the Seychelles: preliminary study. *Marine and Freshwater Research, 53*(2), 123-130. http://doi.org/10.1071/MF01123

Grandcourt, E.M., Al Abdessalaam, T.Z., Francis, F. & Al Shamsi, A.T. (2004) Biology and stock assessment of the Sparids, *Acanthopagrus bifasciatus* and *Argyrops spinifer* (Forsskål, 1775), in the Southern Arabian Gulf. *Fisheries Research, 69*(1), 7-20. http://doi.org/10.1016/j.fishres.2004.04.006

Guiguen, Y., Cauty, C.I., Fostier, A., Fuchs, J. & Jalabert, B. (1994) Reproductive cycle and sex inversion of the seabass, *Lates calcarifer,* reared in sea cages in French Polynesia: histological and morphometric description. *Environmental Biology of Fishes, 39*(3), 231-247. http://doi.org/10.1007/BF00005126

Hamzeh, S., Keivany, Y., Mahboobi Soofiani, N. & Jamshid-Aeen, K. (2017) Reproductive Biology of Kingsoldier Bream (*Argyrops spinifer*) in the Persian Gulf. *Iranian Scientific Fisheries Journal, 26*(3), 91-104.

Harris, P.J., Wyanski, D.M., White, D.B. & Moore, J.L. (2002) Age, growth, and reproduction

of scamp, *Mycteroperca phenax*, in the southwestern North Atlantic, 1979–1997. *Bulletin of Marine Science, 70*(1), 113-132.

Hassin, S., De Monbrison, D., Hanin, Y., Elizur, A., Zohar, Y. & Popper, D.M. (1997) Domestication of the white grouper, *Epinephelus aeneus* 1. Growth and reproduction. *Aquaculture, 156*(3-4), 305-316. http://doi.org/10.1016/S0044-8486(97)00136-1

Hastings, P.A. (1981) Gonad morphology and sex succession in the protogynous hermaphrodite *Hemanthias vivanus* (Jordan and Swain). *Journal of Fish Biology, 18*(4), 443-454. http://doi.org/10.1111/j.1095-8649.1981.tb03785.x

Hastings, P.A. & Petersen, C.W. (1986) A novel sexual pattern in serranid fishes: simultaneous hermaphrodites and secondary males in *Serranus fasciatus*. *Environmental Biology of Fishes, 15*(1), 59-68. http://doi.org/10.1007/BF00005389

Hesp, S.A., Potter, I.C. & Hall, N.G. (2004) Reproductive biology and protandrous hermaphroditism in *Acanthopagrus latus*. *Environmental Biology of Fishes, 70*(3), 257-272. http://doi.org/10.1023/B:EBFI.0000033344.21383.00

Hioki, S. & Suzuku, K. (1996) Sex changing from male to female on the way of protogynous process in three *Centropyge angelfishes* (Pomacanthidae: Teleostei). *Bulletin of institute of Oceanic Research and Development, Tokai University, 17,* 27-34.

Hioki, S., Tanaka, Y. & Suzuki, K. (1995) Reproductive behavior, eggs, larvae, and sexuality of two angelfishes, *Genicanthus watanabei* and *G. bellus,* in an aquarium. *Journal of the School of Marine Science and Technology Tokai University, 40*, 151-171.

Hoffman, S.G. (1980) *Sex-related social, mating, and foraging behavior in some sequentially hermaphroditic reef fishes.* PhD dissertation, University of California, Santa Barbara, CA.

Hoffman, S.G. (1983) Sex-related foraging behavior in sequentially hermaphroditic hogfishes (Bodianus spp.). *Ecology, 64*(4), 798-808. http://doi.org/10.2307/1937203

Horvath, M.L., Grimes, C.B. & Huntsman, G.R. (1990) Growth, mortality, reproduction and feeding of knobbed porgy, *Calamus nodosus*, along the southeastern United States coast. *Bulletin of Marine Science, 46*(3), 677-687.

Hourigan, T.F. & Kelley, C.D. (1985) Histology of the gonads and observations on the social behavior of the Caribbean angelfish *Holacanthus tricolor*. *Marine Biology, 88*(3), 311-322. http://doi.org/10.1007/BF00392592

Howard, J. & Randali, E. (1986) A review of the parrotfishes (family Scaridae) of the Great Barrier Reef of Australia with description of a new species. *Records of the Australian Museum, 38*, 175-228.

Ismail, R.F., Mourad, M.M. & Farrag, M.M.S. (2018) Gonadal development and hermaphroditism of bluespotted seabream, *Pagrus caeruleostictus* (Valenciennes, 1830) from the Mediterranean Sea, Egypt. *The Egyptian Journal of Aquatic Research, 44*(2), 163-171. http://doi.org/10.1016/j.ejar.2018.05.003

Kadota, T., Osato, J., Nagata, K. & Sakai, Y. (2012) Reversed sex change in the haremic protogynous hawkfish *Cirrhitichthys falco* in natural conditions. *Ethology, 118*(3), 226-234.

Kawaguchi, K. & Marumo, R. (1967) Biology of *Gonostoma gracile* (Gonostomatidae) I. Morphology, life history and sex reversal. *Inform. Bull. Planktology Japan, 1967,* 53-69.

Kinoshita, Y. (1939) Studies on the sexuality of genus Sparus (Teleostei). *Journal of science of the Hiroshima University. Series B, Division 1, Zoology, 17*, 25-37.

Kline, R.J., Khan, I.A. & Holt, G.J. (2011) Behavior, color change and time for sexual inversion in the protogynous grouper (*Epinephelus adscensionis*). *PLoS One, 6*(5), e19576. http://doi.org/10.1371/journal.pone.0019576

Kobayashi, K. & Suzuki, K. (1992) Hermaphroditism and sexual function in *Cirrhitichthys aureus* and the other Japanese hawkfishes (Cirrhitidae: Teleostei). *Japanese Journal of Ichthyology, 38*(4), 397-410. http://doi.org/https://doi.org/10.11369/jji1950.38.397

Kokokiris, L., Bruslé, S., Kentouri, M. & Fostier, A. (1999) Sexual maturity and hermaphroditism of the red porgy *Pagrus pagrus* (Teleostei: Sparidae). *Marine Biology, 134*(4), 621-629. http://doi.org/10.1007/s002270050577

Krug, H.M. (1990) The azorean blackspot seabream, *Pagellus bogaraveo* (Brünnich, 1768) (Teleostei, Sparidae). Reproductive cycle, hermaphrodism, maturity and fecundity. *Cybium, 14*(2), 151-159.

Kusen, J.D. (1991) Protogynous hermaphroditism in the sand diver *Trichonotus filamentosus. Bulletin of the Japanese Society for the Science of Fish, 57*(1), 35-40. http://doi.org/10.2331/suisan.57.35

Kusen, J.D. & Nakazono, A. (1991) Protogynous hermaphroditism in the parrotfish, *Calotomus japonicus. Japanese Journal of Ichthyology, 38*(1), 41-45.

Kuwamura, T., Nakashimn, Y. & Yogo, Y. (1994) Sex change in either direction by growth-rate advantage in the monogamous coral goby, *Paragobiodon echinocephalus. Behavioral Ecology, 5*(4), 434-438. http://doi.org/10.1093/beheco/5.4.434

Kuwamura, T., Suzuki, S. & Kadota, T. (2016) Male-to-female sex change in widowed males of the protogynous damselfish *Dascyllus aruanus. Journal of ethology, 34*(1), 85-88. http://doi.org/10.1007/s10164-015-0450-8

Kuwamura, T., Suzuki, S., Tanaka, N., Ouchi, E., Karino, K. & Nakashima, Y. (2007) Sex change of primary males in a diandric labrid *Halichoeres trimaculatus*: coexistence of protandry and protogyny within a species. *Journal of Fish Biology, 70*(6), 1898-1906. http://doi.org/10.1111/j.1095-8649.2007.01464.x

Kuwamura, T., Tanaka, N., Nakashima, Y., Karino, K. & Sakai, Y. (2002) Reversed sex-change in the protogynous reef fish *Labroides dimidiatus. Ethology, 108*(5), 443-450. http://doi.org/10.1046/j.1439-0310.2002.00791.x

Lamrini, A. (1986) Sexualité de *Pagellus acarne* (Risso, 1826) (Teleosteen Sparidae) de la côte Atlantique méridionale du Maroc (21°-26° N). *Cybium (Paris), 10*(1), 3-14.

Lavenda, N. (1949) Sexual differences and normal protogynous hermaphroditism in the Atlantic sea bass, *Centropristes striatus. Copeia, 1949*(3), 185-194. http://doi.org/10.2307/1438985

Lee, S., Kime, D., Chao, T., Lim, H., Chou, R., Lam, T. & Tan, C. (1995) In vitro metabolism of testosterone by gonads of the grouper (*Epinephelus tauvina*) before and after sex inversion with 17α-methyltestosterone. *General and comparative endocrinology, 99*(1), 41-49. http://doi.org/10.1006/gcen.1995.1083

Lee, Y.D. (1995) Sexual phenomenon of protogynous serranid fish. *Exploited Marine Resources,* 1, 143-150.

Lee, Y.H., Du, J.L., Yueh, W.S., Lin, B.Y., Huang, J.D., Lee, C.Y., Lee, M.F., Lau, E.L., Lee, F.Y. & Morrey, C. (2001) Sex change in the protandrous black porgy, *Acanthopagrus schlegeli*: a review in gonadal development, estradiol, estrogen receptor,

aromatase activity and gonadotropin. *Journal of experimental Zoology, 290*(7)*,* 715-726. http://doi.org/10.1002/jez.1122

Leem, J.B., Sakamoto, K., Tsuruda, Y. & Nakazono, A. (1998) Sexual pattern of the labrid fishes collected from Kuchinoerabu-jima, Kagoshima, Japan. *J Journal of the Faculty of Agriculture Kyushu University, 42*(3-4), 409-420.

Leonard, J.L. (1993) Sexual conflict in simultaneous hermaphrodites: evidence from serranid fishes. *Environmental Biology of Fishes, 36*(2), 135-148.

Liem, K.F. (1963) Sex reversal as a natural process in the synbranchiform fish *Monopterus albus. Copeia 1963*(2), 303-312. http://doi.org/10.2307/1441348

Liem, K.F. (1968) Geographical and taxonomic variation in the pattern of natural sex reversal in the teleost fish order Synbranchiformes. *Journal of Zoology, 156*(2), 225-238. http://doi.org/10.1111/j.1469-7998.1968.tb05930.x

Lissia-Frau, A.M. (1968) Research on the sexual differentiation of *Boops boops* (L.) (Teleostei, Sparidae). *Boll. Pesca. Piscic. Idrobiol,* 21, 9-21.

Lissia-Frau, A.M. & Casu, S. (1968) Il differenziamento sessuale di *Lithognathus mormyrus* (L.) e di *Oblada melanura* (L.). *Studi Sassaresi, 46,* 1-19.

Lissia-Frau, A.M., Pala, M. & Casu, S. (1976) Observations and considerations on protandrous hermaphroditism in some species of sparid fishes (Teleostei, Perciformes). *Studi Sassaresi,* 54, 147-167.

Liu, M. & Sadovy, Y. (2004a) Early gonadal development and primary males in the protogynous epinepheline, *Cephalopholis boenak. Journal of Fish Biology, 65*(4), 987-1002. http://doi.org/10.1111/j.0022-1112.2004.00503.x

Liu, M. & Sadovy, Y. (2004b) The influence of social factors on adult sex change and juvenile sexual differentiation in a diandric, protogynous epinepheline, *Cephalopholis boenak* (Pisces, Serranidae). *Journal of Zoolog*y, *264*(3), 239-248. http://doi.org/10.1017/S0952836904005631

Longley, W.H. & Hildebrand, S.F. (1941) *Systematic catalogue of the fishes of Tortugas, Florida: with observations on color, habits, and local distribution.* Papers Tortugas Laboratory, *34,* 1-331.

Lorenzo, J.M., Pajuelo, J.G., Méndez-Villamil, M., Coca, J. & Ramos, A.G. (2002) Age, growth, reproduction and mortality of the striped seabream, *Lithognathus mormyrus* (Pisces, Sparidae), off the Canary Islands (central-east Atlantic). *Journal of Applied Ichthyology, 18*(3), 204-209. http://doi.org/10.1046/j.1439-0426.2002.00318.x

Lou, D.C. (1992) *Age specific patterns of growth and reproduction in tropical herbivorous fishes.* PhD dissertation. James Cook University.

Lucks, D.K. (1970) *Aspects of the biology of the white steenbras (Lithognathus aureti Smith, 1962) in the Sandwich estuary.* PhD dissertation. University of Stellenbosch, South Africa.49 pp.

Lutnesky, M.M.F. (1994) Density-dependent protogynous sex change in territorial-haremic fishes: models and evidence. *Behavioral Ecology, 5*(4), 375-383. http://doi.org/10.1093/beheco/5.4.375

Lutnesky, M.M.F. (1996) Size-dependent rate of protogynous sex change in the pomacanthid angelfish, *Centropyge potteri. Copeia 1996*(1), 209-212. http://doi.org/10.2307/1446961

Mackie, M. (2000) Reproductive biology of the halfmoon grouper, *Epinephelus rivulatus,* at Ningaloo Reef, Western Australia. *Environmental Biology of Fishes, 57*(4)*,* 363-

376. http://doi.org/10.1023/A:1007658027359

Mackie, M.C. (2003) Socially controlled sex-change in the half-moon grouper, *Epinephelus rivulatus*, at Ningaloo Reef, Western Australia. *Coral Reefs, 22*(2)*, 133-142. http://doi.org/10.1007/s00338-003-0296-3

Mackie, M.C. (2006) Anatomical changes to the gonad during protogynous sex change in the half-moon grouper *Epinephelus rivulatus* (Valenciennes). *Journal of Fish Biology, 69*(1)*, 176-186. http://doi.org/10.1111/j.1095-8649.2006.01084.x

Mackie, M. (2007) Reproductive behavior of the halfmoon grouper, *Epinephelus rivulatus*, at Ningaloo Reef, Western Australia. *Ichthyological Research, 54*(3), 213-220. http://doi.org/10.1007/s10228-006-0393-8

Madhu, K. & Madhu, R. (2006) Protandrous hermaphroditism in the clown fish *Amphiprion percula* from Andaman and Nicobar islands. *Indian Journal of Fisheries, 53*(4)*, 373-382.

Manabe, H., Ishimura, M., Shinomiya, A. & Sunobe, T. (2007a) Field evidence for bi-directional sex change in the polygynous gobiid fish *Trimma okinawae*. *Journal of Fish Biology, 70*(2), 600-609. http://doi.org/10.1111/j.1095-8649.2007.01338.x

Manabe, H., Ishimura, M., Shinomiya, A. & Sunobe, T. (2007b) Inter-group movement of females of the polygynous gobiid fish *Trimma okinawae* in relation to timing of protogynous sex change. *Journal of ethology, 25*(2)*, 133-137. http://doi.org/10.1007/s10164-006-0007-y

Manabe, H., Toyoda, K., Nagamoto, K., Dewa, S.-i., Sakurai, M., Hagiwara, K., Shinomiya, A. & Sunobe, T. (2013) Bidirectional sex change in seven species of Priolepis (Actinopterygii: Gobiidae). *Bulletin of Marine Science, 89*(2), 635-642.

Marino, G., Azzurro, E., Massari, A., Finoia, M.G. & Mandich, A. (2001) Reproduction in the dusky grouper from the southern Mediterranean. *Journal of Fish Biology, 58*(4), 909-927. http://doi.org/10.1111/j.1095-8649.2001.tb00544.x

Mathew, G. & Kuruvilla, M. (2010) Anatomical changes during early gonad development in the protogynous greasy grouper *Epinephelus tauvina* (Forsskal). *Indian Journal of Fisheries, 57*(2), 21-24.

Matos-Caraballo, D., Posada, J.M. & Luckhurst, B.E. (2006) Fishery-dependent evaluation of a spawning aggregation of tiger grouper (*Mycteroperca tigris*) at Vieques Island, Puerto Rico. *Bulletin of Marine Science, 79*(1), 1-16.

Maxfield, J.M. & Cole, K.S. (2019) Structural changes in the ovotestis of the bidirectional hermaphrodite, the blue-banded goby (*Lythrypnus dalli*), during transition from ova production to sperm production. *Environmental Biology of Fishes, 102*(11)*, 1393-1404. http://doi.org/10.1007/s10641-019-00914-2

McBride, R.S. & Johnson, M.R. (2007) Sexual development and reproductive seasonality of hogfish (Labridae: *Lachnolaimus maximus*), an hermaphroditic reef fish. *Journal of Fish Biology, 71*(5)*, 1270-1292. http://doi.org/10.1111/j.1095-8649.2007.01580.x

McBride, R.S., Sulak, K.J., Thurman, P.E. & Richardson, A.K. (2009) Age, growth, mortality, and reproduction of Roughtongue Bass, *Pronotogrammus martinicensis* (Serranidae), in the northeastern Gulf of Mexico. *Gulf of Mexico Science, 27*(1), 4. 10.18785/goms.2701.04

McErlean, A.J. & Smith, C.L. (1964) The age of sexual succession in the protogynous hermaphrodite *Mycteroperca microlepis*. *Transactions of the American Fisheries Society, 93*(3), 301-302. http://doi.org/10.1577/1548-

8659(1964)93[301:TAOSSI]2.0.CO;2

McIlwain, J., Hermosa, G.V., Claereboudt, M., Al-Oufi, H.S. & Al-Awi, M. (2006) Spawning and reproductive patterns of six exploited finfish species from the Arabian Sea, Sultanate of Oman. *Journal of Applied Ichthyology, 22*(2)*,* 167-176. http://doi.org/10.1111/j.1439-0426.2006.00723.x

Mead, G.W. (1959) Hermaphroditism in archibenthic and pelagic fishes of the order Iniomi. *Deep Sea Research (1953),* **6,** 234-IN19.

Merrett, N.R., Badcock, J. & Herring, P.J. (1973) The status of *Benthalbella infans* (Pisces: Myctophoidei), its development, bioluminescence, general biology and distribution in the eastern North Atlantic. *Journal of Zoology, 170*(1), 1-48. http://doi.org/10.1111/j.1469-7998.1973.tb05042.x

Meyer, K.A. (1977) Reproductive behavior and patterns of sexuality in the Japanese labrid fish *Thalassoma cupido*. *Japanese Journal of Ichthyology, 24*(2), 101-112. http://doi.org/10.11369/jji1950.24.101

Micale, V., Maricchiolo, G. & Genovese, L. (2002) The reproductive biology of blackspot sea bream *Pagellus bogaraveo* in captivity. I. Gonadal development, maturation and hermaphroditism. *Journal of Applied Ichthyology, 18*(3), 172-176. http://doi.org/10.1046/j.1439-0426.2002.00328.x

Micale, V. & Perdichizzi, F. (1994) Further studies on the sexuality of the hermaphroditic teleost *Diplodus sargus*, with particular reference to protandrous sex inversion. *Journal of Fish Biology, 45*(4), 661-670.http://doi.org/10.1111/j.1095-8649.1994.tb00932.x

Micale, V., Perdichizzi, F. & Santangelo, G. (1987) The gonadal cycle of captive white bream, *Diplodus sargus* (L.). *Journal of Fish Biology, 31*(3), 435-440. http://doi.org/10.1111/j.1095-8649.1987.tb05247.x

Moore, R. (1979) Natural sex inversion in the giant perch (*Lates calcarifer*). *Marine and Freshwater Research, 30*(6), 803-813. http://doi.org/10.1071/MF9790803

Morton, J.K. (2007) *The ecology of three species of wrasse (Pisces: Labridae) on temperate rocky reefs of New South Wales, Australia*. PhD Thesis, University of Newcastle, Callaghan, NSW, Australia.

Moss, J.W., Adams, S. & Welch, D.J. (2002) Bommie cod (*Cephalopholis cyanostigma*): a big surprise from a little fish. Bridging the gap: A workshop linking student research with fisheries stakeholder*s*, pp. 94-107.

Moussac, G.d. (1986) Mise en évidence de l'hermaphrodisme protogyne d'*Epinephelus chlorostigma* (Valenciennes, 1828) aux Seychelles (Pisces, Serranidae). *Cybium, 10*(3), 249-262.

Moyer, J.T. & Nakazono, A. (1978) Protandrous hermaphroditism in six species of the anemonefish genus Amphiprion in Japan. *Japanese Journal of Ichthyology, 25*(2), 101-106.http://doi.org/10.11369/jji1950.25.101

Moyer, J.T. & Zaiser, M.J. (1984) Early sex change: a possible mating strategy of Centropyge angelfishes (Pisces: Pomacanthidae). *Journal of ethology, 2*(1)*,* 63-67. http://doi.org/10.1007/BF02348208

Munday, P.L., Caley, M.J. & Jones, G.P. (1998) Bi-directional sex change in a coral-dwelling goby. *Behavioral Ecology and Sociobiology, 43*(6), 371-377.http://doi.org/10.1007/s002650050504

Munday, P.L., Pierce, S.J., Jones, G.P. & Larson, H.K. (2002) Habitat use, social organization and reproductive biology of the seawhip goby, *Bryaninops yongei*. *Marine and*

Freshwater Research, 53(4), 769-775. http://doi.org/10.1071/MF01205

Nagelkerken, W.P. (1979) Biology of the graysby, *Epinephelus cruentatus*, of the coral reef of Curaçao. *Studies on the fauna of Curaçao and other Caribbean Islands, 60*(1), 1-118.

Nakai, T. & Sano, M. (2002) Evidence of protogynous hermaphroditism in the darkfin hind *Cephalopholis urodeta* (Serranidae) at Iriomote Island, southern Japan. *Fisheries Science, 68*(3)*,* 697-699. http://doi.org/10.1046/j.1444-2906.2002.00479.x

Nakashima, Y., Kuwamura, T. & Yogo, Y. (1995) Why Be a Both-ways Sex Changer? *Ethology, 101*(4), 301-307. http://doi.org/10.1111/j.1439-0310.1995.tb00367.x

Nakashima, Y., Kuwamura, T. & Yogo, Y. (1996) Both-ways sex change in monogamous coral gobies, Gobiodon spp. *Environmental Biology of Fishes, 46*(3), 281-288. http://doi.org/10.1007/BF00005004

Nakazono, A. (1979) Studies on the sex reversal and spawning behavior of five species of Japanese labrid fishes. *Report of Fishery Research Laboratory, Kyushu University, 4*, 1-64.

Nakazono, A. & Kusen, J.D. (1991) Protogynous hermaphroditism in the wrasse *Choerodon azurio*. *Nippon suisan gakkaishi, 57*(3)*,* 417-420. http://doi.org/10.2331/suisan.57.417

Nakazono, A., Nakatani, H. & Tsukahara, H. (1985) Reproductive ecology of the Japanese reef fish, *Parapercis snyderi*. *Proceedings of the Fifth International Coral Reef Congress, Tahiti*, pp. 355-360.

Ndiaye, A.M. (2014) Etude du cycle sexuel et l'inversion sexuelle de *Pagellus bellottii* (Téléostéen: Sparidae) dans les eaux sénégalaises. *Afrique Science: Revue Internationale des Sciences et Technologie, 10*(4), 257-266.

Nemtzov, S.C. (1985) Social control of sex change in the Red Sea razorfish *Xyrichtys pentadactylus* (Teleostei, Labridae). *Environmental Biology of Fishes, 14*(2-3)*,* 199-211. http://doi.org/10.1007/BF00000827

Oh, S.-R., Kang, H.-C., Lee, C.-H., Hur, S.-W. & Lee, Y.-D. (2013) Sex reversal and masculinization according to growth in longtooth grouper *Epinephelus bruneus*. *Development & Reproduction, 17*(2)*,* 79-85. http://doi.org/10.12717/DR.2013.17.2.079

Ohta, I. & Ebisawa, A. (2015) Reproductive biology and spawning aggregation fishing of the white-streaked grouper, *Epinephelus ongus,* associated with seasonal and lunar cycles. *Environmental Biology of Fishes, 98*(6)*,* 1555-1570. http://doi.org/10.1007/s10641-015-0382-8

Ohta, K., Sundaray, J.K., Okida, T., Sakai, M., Kitano, T., Yamaguchi, A., Takeda, T. & Matsuyama, M. (2003) Bi-directional sex change and its steroidogenesis in the wrasse, *Pseudolabrus sieboldi. Fish physiology and biochemistry, 28*(1-4), 173-174. http://doi.org/10.1023/B:FISH.0000030517.06738.e7

Okada, Y.K. (1965a) Bisexuality in Sparid Fishes I. Origin of bisexual gonads in *Mylio macrocephalus.* . *Proceedings of the Japan Academy, 41*(4)*,* 294-299. http://doi.org/10.2183/pjab1945.41.294

Okada, Y.K. (1965b) Bisexuality in Sparid Fishes II. *Proceedings of the Japan Academy, 41*(4), 300-304. http://doi.org/10.2183/pjab1945.41.300

Okada, Y.K. (1966) Sex reversal in *Inegocia meerdervoorti* with special reference to repetition of hermaphroditic state. *Proceedings of the Japan Academy, 42*(5)*,* 497-502. http://doi.org/10.2183/pjab1945.42.497

Okumura, S. (2001) Evidence of sex reversal towards both directions in reared red spotted grouper *Epinephelus akaara*. *Fisheries Science, 67*(3)*,* 535-537. http://doi.org/10.1046/j.1444-2906.2001.00287.x

Ota, K., Kobayashi, T., Ueno, K. & Gojobori, T. (2000) Evolution of heteromorphic sex chromosomes in the order Aulopiformes. *Gene, 259*(1-2), 25-30. http://doi.org/10.1016/S0378-1119(00)00460-1

Paiva, R.B., Neves, A., Sequeira, V., Vieira, A.R., Costa, M.J. & Gordo, L.S. (2018) Age, growth and reproduction of the protandrous hermaphrodite fish, *Sarpa salpa*, from the Portuguese continental coast. *Journal of the Marine Biological Association of the United Kingdom, 98*(2), 269-281. http://doi.org/10.1017/S0025315416001405

Pajuelo, J.G. & Lorenzo, J.M. (2001) Biology of the annular seabream, *Diplodus annularis* (Sparidae), in coastal waters of the Canary Islands. *Journal of Applied Ichthyology, 17*(3), 121-125. http://doi.org/10.1046/j.1439-0426.2001.00292.x

Pajuelo, J.G. & Lorenzo, J.M. (1995) Biological parameters reflecting the current state of the exploited pink dentex *Dentex gibbosus* (Pisces: Sparidae) population off the Canary Islands. *South African Journal of Marine Science, 16*(1), 311-319. http://doi.org/10.2989/025776195784156421

Pajuelo, J.G. & Lorenzo, J.M. (1998) Population biology of the common pandora *Pagellus erythrinus* (Pisces: Sparidae) off the Canary Islands. *Fisheries Research, 36*(2-3)**,** 75-86. http://doi.org/10.1016/S0165-7836(98)00110-6

Pajuelo, J.G. & Lorenzo, J.M. (1999) Life history of black seabream, *Spondyliosoma cantharus*, off the Canary Islands, Central-east Atlantic. *Environmental Biology of Fishes, 54*(3), 325-336. http://doi.org/10.1023/A:1007515301745

Pajuelo, J.G. & Lorenzo, J.M. (2004) Basic characteristics of the population dynamic and state of exploitation of Moroccan white seabream *Diplodus sargus cadenati* (Sparidae) in the Canarian archipelago. *Journal of Applied Ichthyology, 20*(1), 15-21. http://doi.org/10.1046/j.0175-8659.2003.00540.x

Parvez, K. & Al-Marzouk, A. (2000) First observations on natural sex reversal in a protandrous bream (*Sparidentex hasta*: Sparidae) from Kuwait. *Pakistan Journal of Zoology, 32*(3), 229-244.

Pears, R.J., Choat, J.H., Mapstone, B.D. & Begg, G.A. (2006) Demography of a large grouper, *Epinephelus fuscoguttatus*, from Australia's Great Barrier Reef: implications for fishery management. *Marine Ecology Progress Series, 307*, 259-272.

Pears, R.J., Choat, J.H., Mapstone, B.D. & Begg, G.A. (2007) Reproductive biology of a large, aggregation-spawning serranid, *Epinephelus fuscoguttatus* (Forsskål): management implications. *Journal of Fish Biology, 71*(3)*,* 795-817. http://doi.org/10.1111/j.1095-8649.2007.01545.x

Pember, M.B. (2006) *Characteristics of fish communites in coastal waters of north-western Australia, including the biology of the threadfin species Eleutheronema tetradactylum and Polydactylus macrochir.* PhD. Thesis, Murdoch University.

Pember, M.B., Newman, S.J., Hesp, S.A., Young, G.C., Skepper, C.L., Hall, N.G. & Potter, I.C. (2005) *Biological parameters for managing the fisheries for blue and king threadfin salmons, estuary rockcod, Malabar grouper and mangrove jack in north-western Australia.* Fisheries Research and Development Corporation Report Murdoch University, Western Australia.

Penrith, M.J. (1972) Sex reversal in the sparid fish *Chrysoblephus laticeps*. *Koedoe, 15*(1), 135-139. http://doi.org/10.4102/koedoe.v15i1.679

Petersen, C.W. (1991) Sex allocation in hermaphroditic sea basses. *The American Naturalist, 138*(3), 650-667. http://doi.org/10.1086/285240

Petersen, C.W. (1995) Reproductive behavior, egg trading, and correlates of male mating success in the simultaneous hermaphrodite, *Serranus tabacarius*. *Environmental Biology of Fishes, 43*(4), 351-361. http://doi.org/10.1007/BF00001169

Petersen, C.W. & Fischer, E.A. (1986) Mating system of the hermaphroditic coral-reef fish, *Serranus baldwini*. *Behavioral Ecology and Sociobiology, 19*(3), 171-178. http://doi.org/10.1007/BF00300857

Pollock, B.R. (1985) The reproductive cycle of yellowfin bream, *Acanthopagms australis* (Günther), with particular reference to protandrous sex inversion. *Journal of Fish Biology, 26*(3), 301-311. http://doi.org/10.1111/j.1095-8649.1985.tb04269.x

Pressley, P.H. (1981) Pair formation and joint territoriality in a simultaneous hermaphrodite: the coral reef fish *Serranus tigrinus*. *Zeitschrift für Tierpsychologie, 56*(1)*, 33-46. http://doi.org/10.1111/j.1439-0310.1981.tb01282.x

Ramirez, J. (1984) *Aspects of the reproductive biology of three morphospecies in the genus Hypoplectrus; H. chlorurus, H. puella, H. unicolor.* Unpublished M.S. thesis, University of Puerto Rico, Mayaguez, Puerto Rico.

Randall, J.E. & Bruce, R.W. (1983) The parrotfishes of the subfamily Scarinae of the western Indian Ocean with descriptions of three new species. *Ichthyological Bulletin 47*, 1-19.

Reinboth, R. (1962) Morphologische und funktionelle Zweigeschlechtigkeit bei marinen Teleostiern (Serranidae, Sparidae, Centracanthidae, Labridae). *Zoologische Jahrbücher Physiologie Bd, 69,* 405–480.

Reinboth, R. (1963) Naturlicher Geschlechtswechsel bei *Sacura margaritacea* (Hilgendorf) (Serranidae). *Japanese Annotnes Zoology, 36*(4), 173-178.

Reinboth, R. (1964) Inversion du sexe chez *Anthias anthias* (L.) (Serranidae). *Vie Milieu, 17*, 499-503.

Reinboth, R. (1970) Intersexuality in fishes. *Mem. Soc Endocrinology*, *18*,515-543.

Reinboth, R., Becker, B. & Latz, M. (1986) In vitro studies on steroid metabolism by gonadal tissues from ambisexual teleosts: II. Conversion of [14C] androstenedione by the heterologous gonadal tissues of the protandric sea bream *Pagellus acarne* (Risso). *General and comparative endocrinology, 62*(2), 335-340. http://doi.org/10.1016/0016-6480(86)90124-3

Reñones, O., Grau, A., Mas, X., Riera, F. & Saborido-Rey, F. (2010) Reproductive pattern of an exploited dusky grouper *Epinephelus marginatus* (Lowe 1834) (Pisces: Serranidae) population in the western Mediterranean. *Scientia Marina, 74*(3), 523-537.

Roberts Jr, D.E. & Schlieder, R.A. (1983) Induced sex inversion, maturation, spawning and embryogeny of the protogynous grouper, *Mycteroperca microlepis. Journal of the World Mariculture Society, 14*(1-4)*, 637-649. http://doi.org/10.1111/j.1749-7345.1983.tb00116.x

Robertson, D.R. (1972) Social control of sex reversal in a coral-reef fish. *Science, 177*(4053), 1007-1009. http://doi.org/10.1126/science.177.4053.1007

Robertson, D.R. & Choat (1974) Protogynous hermaphroditism and social systems in

labrid fish. *Proc. 2nd Inter. Coral Reef Symp., 1974*(1), pp. 217-225.

Robertson, D.R. & Justines, G. (1982) Protogynous hermaphroditism and gonochorism in four Caribbean reef gobies. *Environmental Biology of Fishes, 7*(2), 137-142. http://doi.org/10.1007/BF00001783

Robertson, R.D., Reinboth, R. & Bruce, R.W. (1982) Gonochorism, protogynous sex-change and spawning in three sparisomatinine parrotfishes from the western Indian Ocean. *Bulletin of Marine Science, 32*(4), 868-879.

Robertson, D.R. & Warner, R.R. (1978) Sexual patterns in the labroid fishes of the Western Caribbean, II. The Parrotfishes (Scaridae). *Smithsonian Contributions to Zoology,* 255, 1-26.

Robinson, G.A. (1976) Sex reversal in the dageraad *Chrysoblephus cristiceps* (Pisces: Sparidae). *Koedoe, 19*(1), 43-48. http://doi.org/10.4102/koedoe.v19i1.1180

Rodgers, E.W., Drane, S. & Grober, M.S. (2005) Sex reversal in pairs of *Lythrypnus dalli*: behavioral and morphological changes. *The Biological Bulletin, 208*(2)*,* 120-126. http://doi.org/10.2307/3593120

Roede, M.J. (1972) Color as related to size, sex and behavior in seven Caribbean Labrid fish species (Genera *Thalassoma*, *Halichoeres* and *Hemipteronotus*). *Studies on the Fauna of Curacao and other Caribbean Islands 32*, 1–266.

Ross, R.M. (1982) *Sex change in the endemic Hawaiian labrid Thalassoma duperrey (Quoy and Gaimard): a behavioral and ecological analysis*. Unpublished PhD dissertation, University of Hawaii.

Ross, R.M. (1984) Anatomical changes associated with sex reversal in the fish *Thalassoma duperrey* (Teleostei: Labridae). *Copeia, 1984*(1), 245-248. http://doi.org/10.2307/1445069

Sadovy, Y. & Donaldson, T.J. (1995) Sexual pattern of *Neocirrhites armatus* (Cirrhitidae) with notes on other hawkish species. *Environmental Biology of Fishes, 42*(2)*,* 143-150. http://doi.org/10.1007/BF00001992

Sadovy de Mitcheson, Y., Liu, M. & Suharti, S. (2010) Gonadal development in a giant threatened reef fish, the humphead wrasse *Cheilinus undulatus*, and its relationship to international trade. *Journal of Fish Biology, 77*(3), 706-718. http://doi.org/10.1111/j.1095-8649.2010.02714.x

Sadovy, Y., Kulbicki, M., Labrosse, P., Letourneur, Y., Lokani, P. & Donaldson, T.J. (2003) The humphead wrasse, *Cheilinus undulatus*: synopsis of a threatened and poorly known giant coral reef fish. *Reviews in Fish Biology and Fisheries, 13*(3), 327-364. http://doi.org/10.1023/B:RFBF.0000033122.90679.97

Sadovy, Y., Rosario, A. & Román, A. (1994) *Reproduction in an aggregating grouper, the red hind*, *Epinephelus guttatus*. In: E.K. Balon, M.N. Bruton & D.L.G. Noakes (Eds.), Women in ichthyology: an anthology in honour of ET, Ro and Genie (pp. 269-286). Springer, Dordrecht. http://doi.org/10.1007/978-94-011-0199-8_21

Sakai, Y. (1997) Alternative spawning tactics of female angelfish according to two different contexts of sex change. *Behavioral Ecology, 8*(4), 372-377. http://doi.org/10.1093/beheco/8.4.372

Sakai, Y., Karino, K., Kuwamura, T., Nakashima, Y. & Maruo, Y. (2003a) Sexually dichromatic protogynous angelfish *Centropyge ferrugata* (Pomacanthidae) males can change back to females. *Zoological science, 20*(5), 627-634. http://doi.org/10.2108/zsj.20.627

Sakai, Y., Karino, K., Nakashima, Y. & Kuwamura, T. (2002) Status-dependent behavioural sex change in a polygynous coral-reef fish, *Halichoeres melanurus*. *Journal of ethology, 20*(2)*,* 101-105. http://doi.org/10.1007/s10164-002-0060-0

Salekhova, L.P. (1961) Hermaphroditism of annular bream *Diplodus annularis* (L.). *Trudy Sevastopol Biological Station, 14*, 257-268.

Schwarz, A.L. & Smith, L.C. (1990) Sex change in the damselfish *Dascyllus reticulatus* (Richardson) (Perciformes: Pomacentridae). *Bulletin of marine science, 46*(3), 790-798.

Sellami, A. & Brusle, J. (1979) Contribution a l'étude de la pêche, de la sexualité et de la reproduction de la mendole *Maena maena* (Linnaeus, 1758), Téléostéens, Maenidae, des cotes tunisiennes. *Mem Biol Mar Ocean IX***,** 91-109.

Shapiro, D.Y. (1981) Size, maturation and the social control of sex reversal in the coral reef fish *Anthias squamipinnis* (Peters). . *Journal of Zoology London, 193*(1), 105–128. http://doi.org/10.1111/j.1469-7998.1981.tb01494.x

Shapiro, D.Y. (1992) Plasticity of gonadal development and protandry in fishes. *Journal of experimental Zoology, 261*(2)*,* 194-203. http://doi.org/10.1002/jez.1402610210

Shapiro, D.Y., Garcia-Moliner, G. & Sadovy, Y. (1994) Social system of an inshore stock of the red hind grouper, *Epinephelus guttatus* (Pisces: Serranidae). In: E.K. Balon, M.N. Bruton & D.L.G. Noakes (Eds.), *Women in ichthyology: an anthology in honour of ET, Ro and Genie. Developments in environmental biology of fishes* (pp. 415-422). Springer, Dordrecht.

Shapiro, D.Y. & Rasotto, M.B. (1993) Sex differentiation and gonadal development in the diandric, protogynous wrasse, *Thalassoma bifasciatum* (Pisces, Labridae). *Journal of Zoology, 230*(2), 231-245. http://doi.org/10.1111/j.1469-7998.1993.tb02685.x

Shapiro, D.Y., Sadovy, Y. & McGehee, M.A. (1993a) Periodicity of sex change and reproduction in the red hind, *Epinephelus guttatus*, a protogynous grouper. *Bulletin of Marine Science, 53*(3), 1151-1162.

Shapiro, D.Y., Sadovy, Y. & McGehee, M.A. (1993b) Size, composition, and spatial structure of the annual spawning aggregation of the red hind, *Epinephelus guttatus* (Pisces: Serranidae). *Copeia, 1993*(2), 399-406. http://doi.org/10.2307/1447138

Sheaves, M. (1995) Large lutjanid and serranid fishes in tropical estuaries: are they adults or juveniles? *Marine Ecology Progress Series, 129*, 31-41. http://doi.org/10.3354/meps129031

Shibuno, T., Chiba, I., Gushima, K., Kakudal, S. & Hashimotol, H. (1993a) Reproductive behavior of the wrasse, *Halichoeres marginatus*, at Kuchierabu-jima. *Japanese Journal of Ichthyology, 40*(3), 351-359.

Shibuno, T., Chiba, I., Hashimoto, H. & Gushima, K. (1994) Reproductive behavior of the wrasse, *Thalassoma lutescens*, at Kuchierabu-jima [in Japan]. *Journal of the Faculty of Applied Biological Science-Hiroshima University (Japan), 33,* 43-50.

Shibuno, T., Gushima, K. & Kakuda, S. (1993b) Female spawning migrations of the protogynous wrasse, *Halichoeres marginatus*. *Japanese Journal of Ichthyology, 39*(4), 357-362.

Shihab, I., Gopalakrishnan, A., Vineesh, N., Muktha, M., Akhilesh, K.V. & Vijayagopal, P. (2017) Histological profiling of gonads depicting protandrous hermaphroditism in *E leutheronema tetradactylum*. *Journal of Fish Biology, 90*(6), 2402-2411. http://doi.org/10.1111/jfb.13324

Shitamitsu, T. & Sunobe, T. (2018) Protandry of the flathead *Suggrundus meerdervoortii* (Teleostei: Platycephalidae). *Ichthyological Research,* 65(4), 507-509. http://doi.org/10.1007/s10228-018-0651-6

Shpigel, M. (1985) *Aspects of the biology and ecology of the Red Sea groupers of the genus Cephalopholis (Serranidae, Teleostei).* PhD dissertation, Tel Aviv University.176 pp.

Shpigel, M. & Fishelson, L. (1986) Behavior and physiology of coexistence in two species of Dascyllus (Pomacentridae, Teleostei). *Environmental Biology of Fishes, 17*(4)*,* 253-265. http://doi.org/10.1007/BF00001492

Shpigel, M. & Fishelson, L. (1991) Territoriality and associated behaviour in three species of the genus Cephalopholis (Pisces: Serranidae) in the Gulf of Aqaba, Red Sea. *Journal of Fish Biology, 38*(6)*,* 887-896. http://doi.org/10.1111/j.1095-8649.1991.tb03628.x

Siau, Y. & Bouain, A. (1994) Preliminary indications on growth and reproduction in the protogynous grouper *Mycteroperca rubra* (Pisces, Serranidae). *Journal of African Zoology, 108*(4), 353-360.

Smith, C.L. (1959) Hermaphroditism in some serranid fishes from Bermuda. *Papers Michigan Academy of Science, Arts and Letters, 44*, 111-119.

Smith, C.L. (1964) Hermaphroditism in Bahama groupers. *Bulletin of the Museum of Natural History, New York, 73*, 42-47.

Smith, C.L. (1965) The patterns of sexuality and the classification of serranid fishes. *American Museum novitates, 2207*, 1-20.

Smith, C.L. (1975) The evolution of hermaphroditism in fishes. In: R. Reinboth (Eds.), *Intersexuality in the Animal Kingdom* (pp. 295-310). Springer, Berlin, Heidelberg. http://doi.org/10.1007/978-3-642-66069-6_28

Smith, C. L., & Atz, E.H. (1969). The sexual mechanism of the reef bass *Pseudogramma bermudensis* and its implications in the classification of the Pseudogrammidae (Pisces: Perciformes). *Zeitschrift für Morphologie der Tiere*, 65(4), 315-326.

Smith, C.L. & Atz, E.H. (1973) Hermaphroditism in the mesopelagic fishes *Omosudis lowei* and *Alepisaurus ferox. Copeia*, *1973*(1), 41-44. http://doi.org/10.2307/1442355

St. Mary, C.M. (1996) Sex allocation in a simultaneous hermaphrodite, the zebra goby *Lythrypnus zebra*: insights gained through a comparison with its sympatric congener, *Lythrypnus dalli. Environmental Biology of Fishes, 45*(2)*,* 177-190. http://doi.org/10.1007/BF00005232

Sulak, K.J., Wenner, C.A., Sedberry, G.R. & Guelpen, L.V. (1985) The life history and systematics of deep-sea lizard fishes, genus Bathysaurus (Synodontidae). *Canadian Journal of Zoology, 63*(3), 623-642. http://doi.org/10.1139/z85-091

Sumpton, W. & Brown, I. (2004) Reproductive biology of the red throat emperor Lethrinus miniatus (Pisces: Lethrinidae) from the southern Great Barrier Reef, Australia. . *Bulletin of marine science, 74*(2), 423-432.

Sunobe, T. & Nakazono, A. (1993) Sex change in both directions by alteration of social dominance in *Trimma okinawae* (Pisces: Gobiidae). *Ethology, 94*(4), 339-345. http://doi.org/10.1111/j.1439-0310.1993.tb00450.x

Sunobe, T., Sado, T., Hagiwara, K., Manabe, H., Suzuki, T., Kobayashi, Y., Sakurai, M., Dewa, S.I., Matsuoka, M., Shinomiya, A., Fukuda, K. & Miya, M. (2017) Evolution of bidirectional sex change and gonochorism in fishes of the gobiid genera Trimma, Priolepis, and Trimmatom. *The Science of Nature, 104*(3-4), 15. http://doi.org/10.1007/s00114-017-1434-z

Suzuki, K. (1979) Spawning behavior, k eggs, larvae, and sex reversal of two pomacanthine fishes, *Genicanthus lamarck* and *G. semifasciatus*, in the aquarium. *Journal of Faculty of Marine Science and Technology, Tokai University, 12*, 149-165.

Suzuki, K., Kobayashi, K., Hioki, S. & Sakamoto, T. (1978) Ecological studies of the anthiine fish, *Franzia squamipinnis*, in Suruga Bay, Japan. *Japanese Journal of Ichthyology, 25*(2)*, 124-140. http://doi.org/10.11369/jji1950.25.124

Suzuki, S., Toguchi, K., Makino, Y., Kuwamura, T., Nakashima, Y. & Karino, K. (2008) Group spawning results from the streaking of small males into a sneaking pair: male alternative reproductive tactics in the threespot wrasse *Halichoeres trimaculatus. Journal of ethology, 26*(3)*, 397-404. http://doi.org/10.1007/s10164-008-0102-3

Tan-Fermin, J.D., Garcia, L.M.B. & Castillo, A.R. (1994) Induction of sex inversion in juvenile grouper, *Epinephelus suillus*, (Valenciennes) by injections of 17α-methyltestosterone. *Japanese Journal of Ichthyology, 40*(4), 413-420. http://doi.org/10.11369/jji1950.40.413

Tan, S.M. & Tan, K.S. (1974) Biology of the tropical grouper. *Epinephelus tauvina* (Forskal) I. A preliminary study on hermaphroditism in *E. tauvina. Singapore Journal of Primary Industries, 2*, 123-133.

Tanaka, H. (1990) Sexual maturation and sex reversal in red spotted grouper, *Epinephelus akaara. Bulletin of National Research Institute of Aquaculture, 17*, 1-15.

Taylor, R.G., Whittington, J.A., Grier, H.J. & Crabtree, R.E. (2000) Age, growth, maturation, and protandric sex reversal in common snook, *Centropomus undecimalis*, from the east and west coasts of South Florida. *Fishery Bulletin, 98*(3), 612-612.

Thompson, R. & Munro, J.L. (1974) The biology, ecology and bionomics of Caribbean reef fishes: Serranidae (hinds and groupers). *Research Reports Zoology Department University of the West Indies, 5*, 1-82.

Thompson, R. & Munro, J.L. (1978) Aspects of the biology and ecology of Caribbean reef fishes: Serranidae (hinds and groupers). *Journal of Fish Biology, 12*(2)*, 115-146. http://doi.org/10.1111/j.1095-8649.1978.tb04158.x

Tobin, A.J., Sheaves, M.J. & Molony, B.W. (1997) Evidence of protandrous hermaphroditism in the tropical sparid *Acanthopagrus berda. Journal of Fish Biology, 50*(1)*, 22-33. http://doi.org/10.1111/jfb.1997.50.1.22

Touart, L.W. & Bortone, S.A. (1980) The accessory reproductive structure in the simultaneous hermaphrodite *Diplectrum bivittatum. Journal of Fish Biology, 16*(4), 397-403. http://doi.org/10.1111/j.1095-8649.1980.tb03717.x

Vadiya, V. (1984) Reproductive systems of *Epinephelus aeneus* and *Epinephelus alexandrinus* (Serranidae) from the southeastern Meditteranean. *Journal of Ichthyology, 24*, 77-81.

Van Oördt, G.J. (1929) Zur mikroskopischen Anatomie der Ovariotestes von Serranus und Sargus (Teleostei). *Zeitschrift für mikroskopisch-anatomische Forschung, 19*, 1-17.

Van Oordt, G.J. (1933) Zur sexualität der gattung Epinephelus (Serranidae, Teleostei). *Zeitschrift für mikroskopisch-anatomische Forschung, 33*, 525-533.

Victor, B.C. (1987) The mating system of the Caribbean rosy razorfish, *Xyrichtys martinicensis. Bulletin of marine science, 40*(1), 152-160.

Von Brandt, V.I. (1979) *Soziale kontrolle des wachstums und geschlechtsdifferenzierung bei jungen anemonenfischen (Amphiprion bicinctus)*. University of Munich, Munich, Germany.

Warner, R.R. (1975) The reproductive biology of the protogynous hermaphrodite

Pimelometopon pulchrum (Pisces: Labridae). *Fishery Bulletin, 73*(2), 262-283.

Warner, R.R. (1982) Mating systems, sex change and sexual demography in the rainbow wrasse, *Thalassoma lucasanum. Copeia, 1982*(3), 653-661. http://doi.org/10.2307/1444666

Warner, R.R. & Lejeune, P. (1985) Sex change limited by paternal care: a test using four Mediterranean labrid fishes, genus Symphodus. *Marine Biology, 87*(1), 89-99. http://doi.org/10.1007/BF00397010

Warner, R.R. & Robertson, D.R. (1978) Sexual Patterns in the Labroid Fishes of the Western Caribbean, I: The Wrasses (Labridae). *Smithsonian Contributions to Zoology, 254*, 38 pp.

Wernerus, F.M. & Tessari, V. (1991) The influence of population density on the mating system of *Thalassoma pavo*, a protogynous Mediterranean labrid fish. *Marine Ecology, 12*(4), 361-368. http://doi.org/10.1111/j.1439-0485.1991.tb00264.x

Yeh, S.L., Dai, Q.C., Chu, Y.T., Kuo, C.M., Ting, Y.Y. & Chang, C.F. (2003) Induced sex change, spawning and larviculture of potato grouper, *Epinephelus tukula. Aquaculture, 228*(1-4), 371-381. http://doi.org/10.1016/S0044-8486(03)00316-8

Yeung, W.S.B. & Chan, S.T.H. (1987) The gonadal anatomy and sexual pattern of the protandrous sex-reversing fish, *Rhabdosargus sarba* (Teleostei: Sparidae). *Journal of Zoology, 212*(3), 521-532. http://doi.org/10.1111/j.1469-7998.1987.tb02922.x

Young, P.C. & Martin, R.B. (1982) Evidence for protogynous hermaphroditism in some lethrinid fishes. *Journal of Fish Biology, 21*(4), 475-484. http://doi.org/10.1111/j.1095-8649.1982.tb02853.x

Young, P.C. & Martin, R.B. (1985) Sex ratios and hermaphroditism in nemipterid fish from northern Australia. *Journal of Fish Biology, 26*(3), 273-287. http://doi.org/10.1111/j.1095-8649.1985.tb04266.x

Zamboni, A. & Relini, G. (1986) Note di biologia di *Spicara flexuosa* (Osteichthyes, Centracanthidae) del Mar Ligure. *Bollettino dei Musei e Degli Istituti Biologici dell'Università di Genova, 52*, 251-265.

Zei, M. (1950) Typical sex-reversal in Teleosts. *Proceedings of the Zoological Society of London*, *119*(4), 917-920. http://doi.org/10.1111/j.1096-3642.1950.tb00917.x

Zohar, Y., Abraham, M. & Gordin, H. (1978) The gonadal cycle of the captivity-reared hermaphroditic teleost *Sparus aurata* (L.) during the first two years of life. *Annales de Biologie Animale Biochimie Biophysique*, *18*(4), 877-882.